



Introduction

Petroleum lease sales of the outer continental shelf (OCS) in the northern Gulf of Alaska appear imminent (Dec. 1975). Because of the impending offshore exploration and subsurface construction activities which will follow the sale, information is needed about the shelf sediments. Knowledge of thickness of Holocene sediments facilitates evaluation of environmental problems involving instability of the seafloor and areas of excessive erosion and deposition.

The data used in the construction of this isopach map of the northern Gulf of Alaska OCS are high resolution seismic profiles collected on the cruise of the R/V THORNDIKE G. THOMPSON, September-October, 1974 (sheet A). These records are available in profile form from there and others, 1975. Navigation instrumentation used to locate the seismic lines shown on sheet A include Decca HF/IF, satellite, Loran A, and radar. To aid in the interpretation of the seismic data, supplemental information was obtained from shipboard sediment descriptions recorded on the cruise of the R/V THORNDIKE G. THOMPSON, September-October, 1975.

The four different sedimentary units in the OCS area (Molnia and Carlson, 1975a) have characteristic seismic signatures. The two youngest sedimentary units, both of which are Holocene, resemble much of the OCS region. One consists largely of clayey silt and is characterized by seismic profiles by relatively horizontal and parallel reflectors. Exceptions are in areas where slides or slumps have developed and the resulting reflectors are very disrupted (Carlson, Bruun, and Molnia, 1975) and mounded of the barrier islands of the Copper River where the principal sediment is fine sand and the reflectors are highly irregular. The second Holocene unit is present off the Bering and Malaspina glaciers where the most recent outcrops show a complex mass of irregular reflectors. The Holocene unit is underlain in some parts of the OCS by a glacial marine unit of poorly sorted mud which is characterized by the stratigraphic position between Holocene and older rocks which range from Tertiary to Pleistocene in age, we assign a Quaternary age to this unit. Its unique seismic signature separates it from the overlying Holocene sediments and therefore it is not included within the isopach unit. In other areas of the OCS, the glacial marine unit is absent and the Holocene unit is underlain directly by folded, faulted and in many places truncated, stratified sedimentary rocks which are probably Tertiary or Pleistocene in age.

Distribution of Holocene Sediments

Holocene sediment is present throughout much of the OCS area in thicknesses varying from less than 5 metres to greater than 300 metres. The widest of Holocene fine sand to clayey silt which makes up the Copper River profile is the thickest of all the modern sediments examined, reaching a thickness of about 300 metres just southeast of the main channel of the Copper River.

Other thick sequences of sediment are: (1) seaward of Icy Bay near Malaspina Glacier (140 m), (2) south of the Bering Glacier (200 m), (3) between Hinchinbrook and Montague Islands (220 m), and (4) at the southeast end of Kayak Island (155 m).

Areas Devoid of Holocene Sediment

The largest area free of Holocene sediment in the western half of the map is an irregularly shaped topographic high that includes Tarr Bank and the Middleton Island platform. Truncated, folded, and faulted sedimentary strata of probable Tertiary age appear to crop out at the surface on these bank areas and they are flanked by a thin band of Quaternary glacial marine poorly sorted mud along the west and north sides (Molnia and Carlson, 1975a). Within this area of Tertiary outcrop are small depressions filled with Holocene silty clay from two to 20 metres in thickness. Our sediment sampling suggests that much of Tarr Bank (and similar areas of the OCS) is covered by a thin veneer (< 2 m) of Holocene mud. This Holocene cover is not detectable on the seismic profiles because of the thickness of the sediment (cm), which is less than the resolution of the seismic system, and/or because of the transparent nature of the sediment.

Smaller areas in the western one-half of the map that are devoid of Holocene sediment, and apparently consist of Tertiary bedrock at the surface, are present east of Montague Island and southwest of Hinchinbrook Island. One of these areas, Seal Rocks, a small group of islands 5 km southeast of Hinchinbrook Island, is composed of well-indurated flysch-like sandstone and argillite that is identical to the Orca formation found on both Montague and Hinchinbrook Islands (Winkler, 1973). Wessels Reef, an intertidal shoal on Tarr Bank about 15 km northeast of Middleton Island, exposes friable sandstone and quartz conglomerate that is similar lithologically to rocks of the Katalla formation on Kayak Island (Parker, 1974; Winkler, oral comm., 1975).

The largest area devoid of Holocene sediments in the eastern one-half of the map extends along the continental slope from south of Kayak Island to the asternmost seismic profile line off the Malaspina Glacier. Much of this slope area consists of glacial marine poorly sorted mud. In addition, four areas of older folded and faulted sedimentary rocks crop out along the outer shelf and upper slope (Molnia and Carlson, 1975b). In a few places along the slope, upon this patches of Holocene sediment (2-20 m thick) cover the older materials. These other areas on the shelf where older folded and faulted sedimentary rocks crop out at the surface are the Kayak Island platform and two structural highs near Cape Yakutat.

Sources of Sediment

The two main sources of Holocene sediment in this OCS region are the Copper River, which, according to Molnia (1964), annually supplies  $10^7 \times 10^6$  metric tons of detritus, and the two large adjacent glaciers (Bering and Malaspina). The sediment being supplied from the two glaciers is at present primarily suspended matter, the sedimentary rocks crop out along the outer shelf and upper slope (Molnia and Carlson, 1975b). In a few places along the slope, upon this patches of Holocene sediment (2-20 m thick) cover the older materials. These other areas on the shelf where older folded and faulted sedimentary rocks crop out at the surface are the Kayak Island platform and two structural highs near Cape Yakutat.

The sediment, whether supplied by river, glacial runoff, or wind, is subject to the rigors of the nearshore currents which, with the exception of local eddies, move in a counter clockwise direction similar to the offshore Alaska Current (Belmont and Carlson, 1974). This counterclockwise movement transports the suspended sediment in a westerly direction. Much of the Copper River sediment is being carried into Prince William Sound through channels north and south of Hinchinbrook Island. Sediments which are part of the Bering glacial runoff plume are carried around Kayak Island. Complex gyres of turbid water have been seen on both sides of Kayak Island on satellite imagery (Belmont and Carlson, 1974). It is likely that some of the suspended sediment settles out over Kayak trough. However, the high resolution seismic profiles indicate that very little of the suspended matter from either the Copper River or from sources east of Kayak Island accumulates on Tarr Bank or the Middleton Island platform. This lack of sediment on these topographic highs may perhaps be due to the scouring action of the frequent storm waves that are particularly large and forceful during the winter season of intense low pressure activity in the Gulf of Alaska.

References Cited

Carlson, Paul R., Bruun, Terry R., and Molnia, Bruce F., 1975, Submarine slides and nearshore faults northern Gulf of Alaska. U.S. Geol. Survey open-file report 75-505.

Molnia, Bruce F., and Carlson, Paul R., 1975a, Shelf sediment distribution: northern Gulf of Alaska. ARMO-SEPM Pacific Section, Long Beach, California, April, 1975, Abstracts Program, p. 15.

Molnia, Bruce F., and Carlson, Paul R., 1975b, Surface sediment distribution northern Gulf of Alaska. U.S. Geol. Survey open-file report 75-506.

Plafker, George, 1974, Preliminary geologic map of Kayak and Hinchinbrook Islands, Alaska. U.S. Geol. Survey open-file map 74-62.

Seimmitz, Earl, 1966, Late Quaternary history and sedimentation of the Copper River delta and vicinity, Alaska (unpublished Ph.D. dissertation, La Jolla: California Univ., San Diego, 160 p).



Seimmitz, Earl, and Carlson, Paul R., 1975, Circulation of nearshore surface water in the Gulf of Alaska. In Carlson, P. R., Gosses, T. J., Janda, R. J., and Peterson, D. R., Principal sources and dispersal patterns of suspended particulate matter in nearshore surface waters of the northeast Pacific Ocean. ERM final report, Natl. Tech. Info. Service, 126 p.

von Neube, Roland, Molnia, Bruce F., Bruun, Terry R., and Carlson, Paul R., 1975, Seismic profiles of a portion of the offshore Gulf of Alaska Tertiary province, R/V THORNDIKE Sept.-Oct., 1974. U.S. Geol. Survey open-file report 75-507.

Winkler, G. R., 1973, Geologic map of the Cordova A-7, A-8, B-6, B-7, and B-8 quadrangles (Hinchinbrook Island) Alaska. U.S. Geol. Survey Misc. Field Studies Map 46-531.

EXPLANATION

- DEVOID OF HOLOCENE SEDIMENTS
- BOUNDARY UNCERTAIN, LIMIT OF SEISMIC PROFILES
- THICKNESS CONTOURS, 25 METRE INTERVAL
- 200 METRE ISOBATH
- TICKS SHOW DIRECTION TOWARD THICKER SEDIMENT



A. Map of seismic reflection lines.

# PRELIMINARY ISOPACH MAP OF HOLOCENE SEDIMENTS, NORTHERN GULF OF ALASKA

By  
Paul R. Carlson and Bruce F. Molnia  
1975

U.S. GEOLOGICAL SURVEY  
OPEN FILE MAP 75-507

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey standards and nomenclature.