(200)
R290

76-743

PRELIMINARY GEOLOGIC MAP OF THE TA'IZ REGION

YEMEN ARAB REPUBLIC

LANDSAT-1 IMAGE NO. 1189-06564

By

M. J. Grolier and W. C. Overstreet

U. S. Geological Survey

This report is preliminary and
has not been edited or reviewed for
conformity with Geological Survey
standards or nomenclature

The project report series presents information resulting from
various kinds of scientific, technical, or administrative studies.
Reports may be preliminary in scope, provide interim results in
advance of publication, or may be final documents.

PRELIMINARY GEOLOGIC MAPS OF THE

YEMEN ARAB REPUBLIC

By

M. J. Grolier, and W. C. Overstreet
U. S. Geological Survey

INTRODUCTION

Objectives of the project and scope of the mapping

This is one of nine separate preliminary geologic maps at 1:500,000 scale covering the entire Yemen Arab Republic (YAR). Each of the nine maps corresponds to an area of the Yemen Arab Republic covered by a Landsat (formerly ERTS) image. The respective Landsat images were used as the bases on which each of the maps was compiled, after stratigraphic and structural analysis in the office, and subsequent checking in the field. This work, in 1975, is one of the services called for under PASA ASIA (IC) YEM-925-22-74 between the U. S. Agency for International Development (USAID) and the U. S. Geological Survey (USGS) for a water and mineral survey in the Yemen Arab Republic, to be performed in cooperation with the Central Planning Organization, the Ministries of Agriculture and Economy, and the Mineral and Petroleum Authority of that country. Part of the program called for a Landsat survey of the country, and one of the goals of the sub-project was to produce a Landsat mosaic of the Yemen Arab Republic.

A substantial part of the imagery used in this study was provided by the National Aeronautics and Space Administration (NASA), Goddard Space Flight Center, under a Memorandum of Understanding between the USGS and NASA.

The intent in compiling these geologic maps was to bring together, at a convenient working scale, previously known and recently acquired geologic data. It is hoped that this set of maps can be used as a tool in hydrologic investigations, minerals exploration, in regional planning, economic and industrial development, highway engineering, and, also, as an aid in mapping the regional geology of the YAR at a larger scale, such as at the 1:100,000 scale.

Permission to release these geologic maps to the open file of the U. S. Geological Survey was given on March 2, 1976, by Dr. A. A. El-Eryani, Minister of Development, and Chairman of the Central Planning Organization, Yemen Arab Republic. The maps are being released in this limited preliminary version, prior to formal publication on a Landsat base. A geologic explanation on a separate sheet accompanies each of the nine geologic maps.

Reconnaissance field checking

Two field trips were made to the YAR in connection with this program. The first field trip, which included reconnaissance on the ground and from the air, was made between June 16 and July 13, 1975, jointly by Maurice J. Grolier and William C. Overstreet (USGS). During the trip it was possible to check an early version of the geologic map, to visit several mineral prospects, and to collect samples of rocks, ores, and slags. The samples were analysed in November 1975 at the laboratories of the USGS in Denver, Colorado. Description of the samples and results of the analyses were presented in a previous report (Overstreet and others, 1976). The second field trip for further checking on the ground was made by Grolier alone between February 1 and 29, 1976, in connection with a hydrologic reconnaissance of the country.

Acknowledgments

The authors are pleased to acknowledge the courtesies and interest of the officials who made possible the work that has led to the preparation and release of this geologic map. They were His Excellency, Dr. A. A. El-Eryani, Minister of Development; Hamoud Ahmid Daif Allah, President, Mineral and Petroleum Authority, Ministry of Economy, YAR, and Aldelmo Ruiz, Director, USAID Mission to the YAR. Without their aid, this work could not have been done.

The authors also wish to acknowledge help received from G. C. Tibbitts, Jr., USGS Project Chief, Water and Minerals Survey in the YAR, who made arrangements for both field trips in June-July 1975 and February 1976, and from their associate, Mohammad Mukred Ibrahim, Assistant Chief Minerals Geologist, Mineral and Petroleum Authority, who cleared all the trips through local authorities and who was a constant source of information on the geology, ore deposits, and geography of the YAR. The writers were also fortunate to be accompanied on the 1975 field trips by Mohammad Luft El-Eryani, a third-year undergraduate student in geology at the College of Science, Kuwait University. His careful observations and refreshing questions added greatly to the discussions at the outcrops.

James W. Aubel, a United States Peace Corps Volunteer and geologist working with G. C. Tibbitts, Jr., on the USAID water supply project in Yemen, had discovered several fossil localities in the Amran Series. He kindly led the writers to these localities and helped in making collections of fossils. In many other ways he contributed to the field work. Help was most generously given by Roy O. Jackson, USGS, in planning the work and in interpreting the results.

Discussions in Sana'a with Dr. Joachim Thiele, Party Chief, Mission to Yemen of the Bundesanstalt für Geowissenschaften und Rohstoffe of the Federal Republic of Germany, and Dr. Karl-Heinz Schultze, Chief (in replacement of Dr. Thiele), and members of their staff, particularly Dr. Norbert W. Roland and Dr. von Prosch, were enlightening. Michael Glase, hydrologist, Tipton and Kalmbach, Inc., Denver, Colorado, and Peter S. Walczak, Resident-Oceanographer at Al Hydaydah, U.N. Food and Agricultural Organization, also supplied valuable geologic information.

REFERENCE

Overstreet, W. C., Domenico, J. A., Grolier, M. J., Tibbitts, G. C., Jr., Ibrahim, M. M., 1976, Trace elements in some rocks, saprolite, gossan, and slag from the Yemen Arab Republic, and their bearing on the iron ore near Sa'dah: U. S. Geol. Survey open-file rept. 76-264 ((IR)Y-8), 70 p., 10 figs., 13 tables.

GPO 910-170

(200)
R 290
no. 76-743

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

Prepared in cooperation with the Yemen Arab Republic Ministry of Economic Development, Minerals and Petroleum Authority and under the sponsorship of the Agency for International Development, U.S. Department of State.

PROJECT REPORT (IR)Y-7
OPEN FILE REPORT
(SHEET 2 of 2)

Base: Landsat-1, 1189-06564 (Jan. 1973)
LINEAMENT SHEET

5 0 5 10 15 20 25 KILOMETRES

Scale 1:500,000

Note: Country boundaries indefinite

PRELIMINARY GEOLOGIC MAP OF THE TAIZ REGION, YEMEN ARAB REPUBLIC

By
Maurice J. Grolier and William C. Overstreet
1975

PLEASE REPLACE IN POCKET
IN BACK OF BOUND VOLUME

(200)
R290
no. 76-743

Compiled by Maurice J. Grolier and William C. Overstreet, U. S. Geological Survey, and based on:

A. Geologic interpretation of LANDSAT-1 images, supplemented by reconnaissance airborne and field surveys in June and July 1975.

B. References, as follows:

Brown, G. F., 1970, Eastern margin of the Red Sea and coastal structures in Saudi Arabia: Roy. Soc. London Phil. Trans., v. 267, p. 75-87

and Jackson, R. O., 1959, Geology of the Asir quadrangle, Kingdom of Saudi Arabia: U.S. Geol. Survey Misc. Geol. Inv. Map I-217A, scale 1:500,000

Coleman, R. G., Fleck, R. J., Hedge, C. E., and Ghent, E. D., in press, The volcanic rocks of southwest Saudi Arabia and the opening of the Red Sea: Saudi Arabia Dir. Gen. Mineral Resources, Red Sea Vol.

Cox, L. R., 1931, The geology of the Farsan Islands, Cizan, and Kamaran Island, Red Sea, Part 2, Molluacan Paleontology: Geol. Magazine, v. 68, p. 1-13.

Dodge, F. C. W., and Rossman, D. L., 1975, Mineralization of the Wadi Qatan area, Kingdom of Saudi Arabia: U.S. Geol. Survey open file rept. 75-309, 71 p.

Geukens, F., 1966, Geology of the Arabian Peninsula, Yemen: U. S. Geol. Survey Prof. Paper 560-B, 23 p., incl. geol. map, scale 1:1,000,000

Gillmann, M., Letullier, A., and Renouard, G., 1966, La Mer Rouge: Geologie et probleme petrolier: Revue de l'Inst. Franc. Petrole, v. 21, no. 10, p. 1467-1487, illus.

Goerlich, F. Von, 1956, Mikropaläontologische und stratigraphische Untersuchungen im Königreich Yemen: Paläont. Zeit., v. 30, p. 213-214.

Greenwood, J. E. G. W., Bleackley, D., and Beydoun, Z. R., 1967, Photogeological Map of Western Aden Protectorate: Overseas Geol. Surveys, London, 2 sheets, scale 1:250,000

Greenwood, W. R., Hadley, D. G., Anderson, R. E., Fleck, R. J., and Schmidt, D. W., in press, Late Proterozoic cratonization in southwestern Saudi Arabia: Royal Soc. London

Heybroek, F., 1965, The Red Sea Miocene evaporite basin, in Salt Basins around Africa: The Institute of Petroleum, London, p. 17-40.

Hotchkiss, Henry, 1963, Petroleum developments in Middle East and adjacent countries in 1962: Am. Assoc. Petroleum Geol. Bull., v. 47, no. 7, p. 1420-1421, fig. 11.

Karrenberg, R., 1959, Yünger Magmatismus in Südwest Arabien (Jemen): Techn. Mitt. Krupp, v. 17, no. 1, p. 33-36

MacFadyen, W. A., 1930, The geology of the Farsan Islands, Cizan, and Kamaran Island, Red Sea, Part 1, General Geology: Geol. Magazine, v. 67, p. 310-315.

Overstreet, W. C., and Rossman, D. C., 1970, Reconnaissance geology of the Wadi Wassa quadrangle, Kingdom of Saudi Arabia: U. S. Geol. Survey open file rept. (IR) SA-117, 68 p.

Rathjens, Carl, and Wissman, H. V., 1934, Süd-arabien Reise, Landeskundliche Ergebnisse: Friedericksen, de Gruyter and Co., M. B. H., Hamburg, v. 2 and 3

Beobachtungen in Yemen: Geologische Rundschau, v. 33, p. 248-279

Shukri, N. M., and Baata, E. Z., 1955, Petrography of the alkaline rocks of Yemen: L'Institut d'Egypte Bull., v. 36, p. 129-163

U. S. Geol. Survey, and Arabian American Oil Company, 1963, Geologic Map of the Arabian Peninsula: U. S. Geol. Survey Misc. Geol. Inv. Map I-270A, scale: 2,000,000

NOTES

Copies of this map are available at the Ministry of Economic Development, Minerals and Petroleum Authority, San'a, Yemen Arab Republic, and at the U. S. Geological Survey, Washington, D. C., or U. S. A. The base for this map is a two-, or three-band (5,7) or (4,5,7) false-color composite of the LANDSAT-1 image indexed hereby, and is available in a black and white positive print at the same places.

Indicated positions of boundary lines not demarcated on the ground are not necessarily definitive. Abbreviations: YAR - Yemen Arab Republic; PDYR - Peoples' Democratic Republic of Yemen.

GEOLOGIC EXPLANATION

Double or fractional symbols indicate grouped formations: Symbols queried where identification doubtful.

CORRELATION OF MAP UNITS

DESCRIPTION OF MAP UNITS

Geologic names and symbols given below apply to the whole area of the Yemen Arab Republic; some names and symbols may not appear on the geologic map of an area covered by an individual LANDSAT-1 image. Names and descriptions of geologic units, unless otherwise noted, are adopted from U.S. Geological Survey and Arabian American Oil Company, 1963, Geologic map of the Arabian Peninsula; U.S. Geol. Survey Misc. Geol. Inv. Map 270-A, and Brown, G. F., and Jackson, R. O., 1959, Geology of the Asir quadrangle, Kingdom of Saudi Arabia: U.S. Geol. Survey Misc. Geol. Inv. Map 217-A.

Qsb Silt, clay, and muddy sand; commonly saturated with brine and salt encrusted; in mud flats (sabkhas) along the Red Sea coast

Qu River terrace deposits, alluvial fans, gravel, sand, and silt including unmapped alluvium which overlies rock salt at Jabal Kushah, near Guma; numerous loess deposits particularly in the central plains. Wherever possible, alluvial deposits have been divided regionally on a basis of reflectance, natural vegetation and crops, altitude, and location into six sub-units, as follows:

Qa16, alluvial gravel, sand, and silt restricted to channels and flood plains of present-day ephemeral streams

Qa15, alluvial gravel, sand, and silt on river terraces and fans, adjacent to and higher than the flood plains of present-day streams; generally darker than Qa16; may include colluvium at base of foothills

Qa14, same as above, but darker, and possibly older

Qa13, same as Qa14, but higher and older

Qa12, same as Qa13, but higher and farther inland from the Red Sea Coast

Qa11, alluvial gravel heavily coated with desert varnish, restricted to dissected river terraces on the south valley slope of Wadi Jawf, north of Jabal Bahra and west of Wadi Raghwan

Qk Yellow and green marly limestone, white limestone, and reef limestone, undifferentiated, exposed on Kamaran Island. Fossiliferous, and of probable Pleistocene age (MacFadyen, 1930; Cox, 1931). Probably correlative with unmapped marine terrace deposits which disconformably overlie Pliocene-Miocene tuffaceous sandstone at the Al Luhayyah diapirs

Ql Loess deposits, with calcareous concretions and caliche layers; fossil mollusks abundant locally; may include alluvial silt alternating with alluvial or colluvial gravel

Qe Folian sand, commonly mobile

Qa Basalt flows and dikes; numerous scattered cones and craters; at places covered with tuff and volcanic bombs. May be rock and time equivalent of the Aden Volcanic Series in the People's Democratic Republic of Yemen; in the San'a region, lava flows have been divided regionally on a basis of reflectance into four sub-units, as follows:

Qa1, very dark basaltic lobate flows, extruded in historical times, possibly in 3rd century A. D. (Rathjens, G., and Wissman, H. V., 1934, v. 2, p. 13; v. 3, p. 105, fig. 51; p. 162-163; Rathjens, C., and Wissman, 1942, v. 33, p. 276)

Qa2, dark basaltic flows

Qa3, thin basalt flows, discontinuous over older rocks; appear lighter gray than units Qa2 and Qa4 on LANDSAT-1 images

Qa4, basalt flows forming a continuous mantle over older rocks; Qa1 and Qa2 possible are part of only one eruption phase

Tba BAID FORMATION--Gray, red, and green siliceous and tuffaceous shale and sandstone; also limestone and evaporite layers. Includes rock salt of salt domes at Salif and Jabal Qimnah, and at Jabal Kushah near Guma. Generally unfossiliferous, but middle to late Miocene microfossils reported by Klaus (in Heybroek, 1965, p. 34-35) from rock salt at Jabal Kushah, and at Salif, and late Pliocene microfauna reported from marine sediments overlying salt (Goerlich, 1956, p. 213-214). Correlated with rocks of the Baid Formation exposed in Wadi Baid, Saudi Arabia, because of similar lithology (Gillmann, Letullier, and Renouard, 1966, p. 1479-1480, pl. 1, fig. 4).

Ta Hypabyssal andesite and diabase intrusives, commonly glomeroporphyritic, and in dike swarms

Tgr Alkali granite and diorite in subvolcanic plugs, stocks, and plutons (Karrenberg, 1959, v. 17, no. 1, p. 33-36); leucocratic granite locally has primary flow banding. Crests of unbreached plugs may be overlain by hydrothermally altered rocks of the Yemen Volcanics, locally in northwestern part of the Yemen Arab Republic mapped as Tertiary laccoliths (U.S. Geol. Survey and Arabian American Oil Co., 1963). Some granitic plutons as at Jibal Sabir, south of Taiz, have syenite margins. A K-Ar age of

22.7 ± 0.9 m.y. is reported for a granite sample from Jibal Sabir collected by R. O. Jackson (Field No. ROJ-1), and analysed by R. F. Marvin, H. H. Mehnert, and Violet Merritt (Marvin, 1974, written commun. to G. F. Brown). A similar K-Ar age (22.0 ± 0.7 m.y.) is reported by Marvin (1974, written commun. to Brown) for a syenite sample which had been collected from a plug cutting a laterite deposit in the Sirat Plateau, Saudi Arabia by Brown (Field No. 519B).

Tl Laterite and saprolite, mainly white, may be yellow or red, developed on upper surface of Precambrian crystalline rocks by prolonged weathering during Eocene (?) time, to 50 meters in thickness; probably equivalent to laterite in As Sirat Mountains, Saudi Arabia (Brown and others, 1959)

TKy YEMEN VOLCANICS, undivided--Bedded alkaline flows and pyroclastic rocks including but not restricted to rhyolite, comendite, pantellerite, trachyte, andesite, basalt, and ankaramite (Shukri and Baata, 1955, v. 36, p. 129-163), with interbedded lentils of fluviatile and lacustrine sand, clay, and shale; locally contains fresh-water Oligocene-Miocene fossils; upper surfaces of many volcanic beds weather to reddish paleosols a few centimeters to a few meters thick, particularly in middle and upper parts of the sequence; whole sequence of Yemen Volcanics at least 2,000 meters thick. Term Yemen Volcanics introduced here to replace former name Trap Series (Geukens, 1966), to emphasize presence of thick sequence of highly fractionated felsic volcanic rocks. Wherever possible, the Yemen Volcanics have been divided regionally on basis of reflectivity and stratigraphic succession into six sub-units, as follows:

TKy6, dark basaltic flows;

TKy5, generally leucocratic felsic tuffs with some dark basaltic flows, associated with the formation and collapse of a circular volcanic structure, 8.5 km in diameter, in the north-central part of the area covered by LANDSAT-1 image 1189-06561;

TKy4, predominantly felsic and tuffaceous, with some basaltic flows, underlies TKy6 and TKy5;

TKy3, predominantly felsic and tuffaceous; older than TKy4;

TKy2, predominantly felsic and tuffaceous; older than TKy3;

TKy1, predominantly basaltic, but includes green felsic conglomerate, porphyritic trachyte, and pink tuffs; overlies the Tawilah Group.

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

610

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

774

NOTE

The gossans in the Kingdom of Saudi Arabia at Wadi Wassa (Overstreet and Rossman, 1970), and at Wadi Qatan (Dodge and Rossman, 1975) were formed over extensive deposits of stratabound massive and disseminated pyrite and pyrrhotite in Precambrian volcanogenic rocks. Should the iron deposits near Sa'dah, which are known to extend tens of kilometers northward, and similar deposits gossans over massive sulfide, then the region mined for iron northward from the vicinity of Sa'dah and Majadh to the border between the Yemen Arab Republic and the Kingdom of Saudi Arabia merit geologic, geophysical, and geochemical exploration for base metals, nickel, silver, gold and molybdenum.

INDEX MAP OF YEMEN ARAB REPUBLIC - Showing location of Landsat-1 images used as bases for the geologic investigation maps published by the U.S. Geological Survey. Scale 1:500,000

PLEASE REPLY IN BLOCK
BACK OF BOOK

(200)
R290
no. 76-743

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

Prepared in cooperation with the Yemen Arab Republic Ministry of Economic Development, Minerals and Petroleum Authority and under the sponsorship of the Agency for International Development, U.S. Department of State.

PROJECT REPORT (IR)Y-7
OPEN FILE REPORT
(SHEET 1 of 2)

Base: Landsat-1, 1189-06564 (Jan. 1973)
Geographic coordinates developed by NASA
This geologic map is preliminary and has not been edited for review for conformity with U.S. Geological Survey standards and nomenclature.

5 0 5 10 15 20 25 KILOMETRES

44°00' E

Note: Country boundaries indefinite

Scale 1:500,000

PRELIMINARY GEOLOGIC MAP OF THE TAIZ REGION, YEMEN ARAB REPUBLIC

By
Maurice J. Grolier and William C. Overstreet
1975

PLEASE REPLACE IN POCKET
IN BACK OF BOUND VOLUME