

EXPLANATION

- Holocene**
- QUATERNARY**
- Qc Sand and gravel deposits, undifferentiated
- JURATRIAS**
- ala, qm Intrusive rocks
ala, alaskite
qm, quartz monzonite
- PRECAMBRIAN AND JURATRIAS**
- mig Migmatite
- PRECAMBRIAN**
- gn, bs, qfs Bucaramanga Gneiss
gn, gneiss
bs, biotite schist
qfs, quartzo-feldspathic schist
- Contact—Dashed where approximately located; dotted where concealed
- - - Fault—Dotted where concealed
- 20 Fault, inclined
- ↑ Fault, vertical
- 50 Minor fold
- 65 Inclined foliation
- ↑ Vertical foliation
- ↑ Joint, vertical
- 35 Joint, inclined
- 50 Quartz vein, inclined
- ↑ Quartz vein, vertical
- Tunnel
- Tunnel(closed)

FRESH AND ALTERED ROCK SAMPLES

- N 0 ppm
- L <5 ppm
- ≥5 <10
- ≥10 <50
- ≥50 <100
- ≥100 <500
- ≥500

STREAM SEDIMENT SAMPLES

- N 0 ppm
- L <5 ppm
- △ ≥5 <10
- ▲ ≥10 <50
- △ ≥50 <100
- ▲ ≥100 <500
- ▲ ≥500
- X Sample insufficient for analysis

PAN CONCENTRATE SAMPLES

- N 0 ppm
- L <5 ppm
- ≥5 <10
- ≥10 <50
- ≥50 <100
- ≥100 <500
- ≥500
- X Sample insufficient for analysis




FIGURE 14. MAP SHOWING DISTRIBUTION OF LEAD IN THE VETAS DISTRICT, COLOMBIA

G-26-71