



EXPLANATION

- Holocene**
- Qc Sand and gravel deposits, undifferentiated
- JURASSIC-TRIASSIC QUATERNARY**
- ala qm Intrusive rocks
ala, alaskite
qm, quartz monzonite
- PRECAMBRIAN AND JURA-TRIAS**
- mig Migmatite
- gn bs qfs Bucaramanga Gneiss
gn, gneiss
bs, biotite schist
qfs, quartzo-feldspathic schist
- Contact—Dashed where approximately located, dotted where concealed
- - - Fault—Dotted where concealed
- 70 Fault, inclined
- ↑ Fault, vertical
- 50 Minor fold
- 65 Inclined foliation
- ↑ Vertical foliation
- Joint, vertical
- Joint, inclined
- Quartz vein, inclined
- Quartz vein, vertical
- Tunnel
- Tunnel(closed)
- FRESH AND ALTERED ROCK SAMPLES**
- N 0 ppm
- L <5 ppm
- ≥5 <10
- ≥10 <50
- ≥50 <100
- ≥100 <500
- ≥500
- STREAM SEDIMENT SAMPLES**
- N 0 ppm
- L <5 ppm
- △ ≥5 <10
- ▲ ≥10 <50
- △ ≥50 <100
- ▲ ≥100 <500
- ▲ ≥500
- PAN CONCENTRATE SAMPLES**
- N 0 ppm
- L <5 ppm
- ≥5 <10
- ≥10 <50
- ≥50 <100
- ≥100 <500
- ≥500
- X Sample insufficient for analysis

SCALE 1:5000
 0 100 200 300 400 500 METRES
 0 500 1000 FEET

G-26071

FIGURE 11. MAP SHOWING DISTRIBUTION OF COPPER IN THE VETAS DISTRICT, COLOMBIA