

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

GEOLOGIC MAP OF THE DESCANSO QUADRANGLE,
SAN DIEGO COUNTY, CALIFORNIA

By

Wendy C. Hoggatt and Victoria R. Todd

Open-file report

77-406

This report is preliminary
and has not been edited or
reviewed for conformity with
Geological Survey standards
and nomenclature

Geologic map of the Descanso quadrangle, San Diego County, California

Purpose of project:

Published maps indicate that rocks of the Peninsular Ranges batholith in southern San Diego County comprise a relatively unfaulted block but topographic maps and imagery at all scales reveal numerous prominent lineaments that cross the block in many directions. The purpose of this project, which began in 1974, is to evaluate the structural stability of the block, and in particular, to determine whether lineaments are related to faulting. Figure 1 shows the project area. The age, magnitude and direction of displacement of faults are being evaluated. The primary product of the project will be a number of geologic maps at a scale of 1:24,000. Mapping of the bedrock geology of the batholith became an important part of the project because the geologic maps that were available when the study began were not detailed enough for determination of fault displacements. The Descanso quadrangle is the second of this series.

Introduction:

The Descanso quadrangle lies within the mid-Cretaceous Peninsular Ranges batholith of southern California and Baja California (Fig. 1). Nine plutonic units have been differentiated within the project area and informal names have been assigned to avoid confusion with earlier nomenclature. Five of the units were not recognized previously and published names for rocks in the study area included one or more of these five units. The name Cuyamaca gabbro (Everhart, 1951) has been retained. A tenth bedrock unit consists of meta-sedimentary rocks of uncertain age which occur as roof pendants and septa in the batholith.

Previous usage has been followed in referring to the nine rock units as plutonic, even though all of the rocks in this part of the batholith have undergone dynamothermal metamorphism. The plutonic rocks are gneissic and their foliation consists chiefly of the planar orientation of recrystallized mineral grains and aggregates (Fig. 2). Gneissic textures include augen gneiss and mylonite gneiss. Petrographic study reveals igneous textures modified by strain at temperatures high enough for recrystallization to occur, indicating that the plutonic rocks have undergone solid-state flowage at elevated temperatures. Foliation is locally concordant with plutonic contacts, but commonly crosses contacts and, therefore, is in part younger than the intrusion of the igneous rocks. Where noted by earlier workers, foliation was considered to be a primary igneous structure or protoclastic structure. The foliations form a regional tectonite fabric which is concordant with that of the metasedimentary wall rocks. The metamorphism of the wall rocks was thought to pre-date emplacement of the batholith in this area (Everhart, 1951). Preliminary results of the present study indicate that in this part of the batholith plutons were emplaced during regional metamorphism and deformation and that deformation and recrystallization continued long after emplacement.

Nature of plutonic contacts:

A striking feature of the plutonic rocks in southern San Diego County is the prevalence of synplutonic contacts between plutons of different ages. Along some contacts between two plutons the true age relationship locally appears reversed, so that the older pluton has a chilled margin against the younger pluton, sends dikes into the younger pluton, and carries inclusions of it (Fig. 3). These contacts were deformed during the regional deformation,

with the result that both "normal" dikes (younger pluton of an intrusive pair into older) and synplutonic dikes (older pluton into younger) have been stretched, intricately folded, and pulled apart so that they resemble inclusions. Whether they originated as stopped inclusions or dismembered dikes, blocks of one pluton in the other are increasingly rounded and assimilated away from the contact between the two plutons. The resulting contact relations may be extremely ambiguous, so that many outcrops had to be examined before consistent plutonic age relationships were worked out. The prevalence of synplutonic contacts indicates that the entire batholith remained mobile throughout emplacement. The presence of mutual chilled margins and quenched dikes suggests a continuing magmatic component of mobility, but rock textures and the relation between foliation and plutonic contacts imply solid-state flowage. Some combination of the two probably occurred.

Rock units:

Ten bedrock units are discussed below and their intrusive sequence is depicted in the map legend. Modal data for the plutonic units is given in Figure 4. The following discussion is based in part upon observations of rock units and contact relations in the surrounding quadrangles, which are shown in Figure 5. The bedrock units are overlain locally by unconsolidated Quaternary deposits.

Metasedimentary rocks:--Metasedimentary rocks (Rm) consist predominantly of fine-grained quartzo-feldspathic semischistose rock and micaceous, feldspathic quartzite. Andalusite-bearing schists are interbedded locally with these rocks, but pelitic schist is a minor rock in the Descanso quadrangle. The rocks are medium to dark gray and weather dark reddish-brown. They tend to form topographic highs. The quartzo-feldspathic semischistose rocks and feldspathic quartzites are thin-bedded (<1 foot) and probably were fine-grained, argillaceous sandstones and siltstones with calcareous cement. Fine-grained calc-silicate rock variably composed of epidote, garnet, plagioclase, pyroxene, quartz and hornblende is common in the vicinity of, and as inclusions in Kc and KlB plutons. Fine-grained black amphibolite (meta-basalt or meta-basaltic tuff?) is a minor component of the metasedimentary section, and metamorphosed calcareous pebble and small cobble conglomerate occurs locally. Cross-bedding and graded beds were observed locally but deformation has obliterated sedimentary structures in many places. Most mineral assemblages suggest amphibolite facies metamorphism but proximity to gabbros and pyroxene tonalite plutons locally has increased the grade of metamorphism, which indicates that the sedimentary rocks have undergone a complex combination of dynamothermal and thermal metamorphism. Foliation in the rocks is parallel to steep and near-vertical relict bedding, and axial planes of metamorphic folds are generally concordant with this foliation.

A stipple pattern superimposed on Rm means that the metasedimentary rock is interlayered intimately with Khc.

Cuyamaca Gabbro:--The name Cuyamaca Gabbro was applied by Everhart (1951) to the large mafic body in the Cuyamaca Mountains. He assumed that all of the large mafic bodies (Guatay Mountain, Viejas Mountain, Poser Mountain) in south-central San Diego County were coeval, and the present study agrees with this interpretation. These bodies, consisting of peridotite, olivine gabbros,

hornblende gabbro, and norite were considered by Everhart to post-date the granitic plutons in the northern part of the Cuyamaca Peak quadrangle, but this study has found gabbro to be the oldest plutonic rock. Because the granitic intrusive bodies form sheets in gabbro, it is difficult to determine which rock is older from map relations in any one local area. However, relations over a large area show that the Kc bodies are screens surrounded by sheeted complexes of younger granitic rocks. Thus, the present outlines of gabbro bodies and their distribution probably do not accurately reflect either the original extent of gabbro or the original shapes of gabbro plutons.

The present study has found the Cuyamaca Gabbro unit to be deformed and recrystallized as are the younger granitic units. All gabbro observed in the Cuyamaca Peak and surrounding quadrangles is strongly foliated and in most cases apparent primary compositional layering is parallel or near parallel to the foliation of recrystallized mineral aggregates. In marginal parts of gabbro, foliation is concordant with regional foliation. It is more complex in the interior parts of the bodies where it may parallel regional foliation, but also shows swirling patterns. Foliation within the gabbro, and by inference, the so-called primary structures, appears to have formed by solid-state flowage.

One reason for ambiguity over the age of Cuyamaca Gabbro is that locally the unit has broad, fine-grained and porphyritic margins against younger granitic plutons. Thin sections of gabbro from these areas show relict chilled igneous textures modified by recrystallization. These rocks are commonly associated with zones of intrusion breccia between gabbro and granitic plutons consisting of variably rounded blocks of fine-grained and porphyritic gabbro in a matrix of chilled, contaminated granitic rock. In other places, contacts between gabbro and granitic plutons are sharp and the granitic plutons send dikes into gabbro.

Both types of contact may occur together in a given area. The intrusion breccia zones are deformed strongly parallel to the regional foliation. The intrusion breccia grades into highly contaminated granitic plutons which contain abundant, large, fine-grained gabbro inclusions. Although the intrusion breccia is younger than Kc, it is shown on the map by a block pattern superimposed on Kc because 1) fine-grained gabbro makes up most of the rock, 2) the zones of intrusion breccia grade into gabbro complexly and commonly cannot be mapped as separate zones, and 3) the granitic matrix may have as its source Kcp, Kpv, Kcm or Klb.

The origin of the chilled-appearing gabbro in this zone is still under investigation. Locally, gabbro intruded metasedimentary rocks to form a zone of intrusion breccia consisting of a matrix of fine-grained gabbro surrounding variably assimilated metasedimentary inclusions. Inclusions of metasedimentary rock occur in some intrusion breccia zones between Kc and granitic plutons, but there are places where no metasedimentary rocks appear to be involved and where gabbro grain size decreases systematically as contacts with granitic plutons are approached. These contacts may be synplutonic, i.e., both older and younger rocks were able to chill against one another. Fine-grained gabbro dikes emanating from Kc plutons cut the intrusion breccia zones surrounding several of the large gabbro bodies in the project area. Such dikes also cut units as young as Kem. This suggests that Kc contacts are in part synplutonic with younger plutons.

Tonalite of Las Bancas:--The tonalite of Las Bancas (Klb) is a widespread unit in the project area. The unit was formerly called Green Valley tonalite (Everhart, 1951). It is dark gray on fresh surfaces and weathers to a reddish or buff-gray color. Typically, the tonalite forms extensive, low, bouldery outcrops. The unit has few mafic inclusions in the Descanso quadrangle where it occupies a broad plateau at about 4,000 feet named Las Bancas. The unit is homogeneous and has sharp contacts with granitic units. Klb has complex contact relations with TRm locally, marked by intrusion breccia, abundant partly assimilated metasedimentary inclusions in Klb, and contamination-assimilation reactions between the two units. Locally, injection migmatite occurs between Klb and TRm. Klb has a fine-grained (chilled) margin against Cuyamaca Gabbro in a few places and it is intruded by Kcp, Kcr, Kcm and Kmd.

The tonalite of Las Bancas is a medium-grained gneissic rock, approximately equigranular, with lenticular recrystallized mafic aggregates. Locally it carries 1 cm poikilitic biotite grains. The rock has 15-20 percent quartz, traces of K-feldspar, relict zoned phenocrysts of calcic andesine to labradorite, with a color index ranging from 25-30 percent. The reaction sequence of the mafic assemblage is opx -> cpx -> olive-green hornblende -> yellowish-brown biotite. Most samples contain chiefly pyroxene and biotite, with the hornblende occurring as sparse, narrow rims on corroded pyroxene cores and both pyroxene and hornblende poikilitically enclosed by biotite. These igneous reaction textures have been modified by recrystallization.

Hybrid gneiss of Harper Creek.--The hybrid gneiss of Harper Creek (Khc) is a gray and yellow-weathering, cordierite- and sillimanite-bearing, quartz biotite-plagioclase-K-feldspar/muscovite gneiss with a dynamothermal metamorphic texture. The unit, which is remarkably homogeneous over large areas, includes rocks that closely resemble Kcr in the field as well as rocks that are clearly metasedimentary in origin. Khc contains abundant metasedimentary inclusions, up to several meters in length, as well as evenly and closely spaced micaceous lenses, several cm long, which grade into partly assimilated metasedimentary inclusions. These inclusions, plus the high proportion of quartz and mica attest to the rock's partial sedimentary origin. Local relict igneous textures and gradation into Kcr indicate that the Khc unit originated by thorough mixing of Kcr and TRm enhanced by deformation and metamorphic temperatures that existed for a considerable length of time both before and after emplacement of Kcr. In a few places, Khc has intruded younger plutons synplutonically which indicates that the rock was locally as mobile as the plutonic units.

The hybrid gneiss is not migmatite, although minor migmatite does occur at contacts with TRm. These contacts may be sharp, or may be marked by large areas consisting of alternating layers of Khc and TRm that are too small to be depicted at the map scale. A stipple pattern has been used in these areas to indicate that one of the two units contains up to 50 percent of the other. Contacts between Khc and Kcr may be gradational or sharp. Where they are gradational, a zone of fine-grained Kcr containing abundant partly assimilated inclusions of TRm may occur between Khc and Kcr. These are interpreted as marginal parts of Kcr plutons.

Granodiorite and tonalite of Cuyamaca Reservoir:--The area of outcrop of the granodiorite and tonalite of Cuyamaca Reservoir (Kcr) is small in the Descanso quadrangle, but the unit underlies large areas elsewhere within the project area. Because of this and the fact that Kcr is intruded by younger plutons, it is difficult to determine the original shapes of Kcr plutons solely

from exposures in this quadrangle. Kcr is interlayered with younger units in sheeted complexes and may itself have been emplaced in a sheeted fashion into older plutons. The rock weathers to reddish and yellowish-tan and is light to dark gray on fresh surfaces depending upon its mafic content, which ranges from 14-29 percent. Locally, Kcr contains abundant fine-grained mafic inclusions that are flattened parallel to foliation. Although varying compositionally from tonalite to granodiorite, the unit is texturally homogeneous and the two compositional varieties seem to grade into one another. The rock's texture is fine to medium-grained, very strongly gneissic and, on the average, more strongly deformed than the other plutonic units. In thin section, Kcr shows some of the most strained and recrystallized igneous textures of all units.

All samples of Kcr contain some K-feldspar, the plagioclase is andesine, greatly modified by recrystallization, and pale reddish-brown biotite is the chief mafic mineral. Some samples contain no hornblende, others show a few hornblende relicts within biotite aggregates, but in most rocks intergrowths of actinolite and epidote have replaced hornblende. The chief accessory minerals are allanite and sphene. The mineralogical and textural differences between Kcr and other granitic units may be related to intimate mixing of the Kcr unit with metasedimentary rocks. The granodiorite and tonalite are especially gneissic, locally porphyroclastic and mylonitic, next to large bodies of metasedimentary rock. Here the unit contains abundant partly assimilated inclusions of $\mathbb{R}m$ and is rich in mica. Fine-grained granodiorite and tonalite may grade into Khc which locally occurs as sheets between Kcr and $\mathbb{R}m$.

A noticeable decrease in Kcr grain size occurs 1 to 2 meters from contacts with Cuyamaca Gabbro (Kc) and tonalite of Las Bancas (K1b) suggesting that Kcr is younger. Many of these contacts are poorly exposed so these age relations must still be considered tentative and subject to re-interpretation with further mapping.

Quartz monzonite and granodiorite of Chiquito Peak:--The quartz monzonite and granodiorite of Chiquito Peak (Kcp) is a medium-grained, strongly foliated, white-weathering rock with color index ranging from 5 - 12 percent. The plagioclase feldspar is oligoclase with relict euhedral zoning and the mafic minerals are chiefly dark greenish-brown biotite which appears to be derived from reaction of dark green to brown hornblende. Both biotite and hornblende have recrystallized but igneous relicts are present. Prominent accessory minerals are sphene and allanite. Although the unit forms at least one fairly large pluton (north of the town of Descanso), it typically was emplaced in a series of steeply-dipping interconnected sheets and lenses.

The quartz monzonite and granodiorite of Chiquito Peak intruded older plutons intimately, chilled against them, and locally show a high degree of contamination through assimilation of stopped inclusions. This is particularly true where quartz monzonite is in contact with metasedimentary rock, Kc, and Kem. The contamination and post-intrusive deformation have given rise to complex hybrid zones between these plutons. A contaminated quartz monzonite category has been used (dashed pattern) to designate parts of Kcp plutons which are particularly inclusion-rich and contaminated. These rocks are intimate mixtures of quartz monzonite, granodiorite, and tonalite representing both contaminated quartz monzonite and mafic inclusions which have been almost completely assimilated. Because of the close association of these zones with mafic plutons, there is little doubt that stopping of mafic plutons (both older and younger than Kcp) has given rise to the heterogeneous zones.

A textural variety of Kcp is a fine- to medium-grained, sub-porphyritic (1 cm relict euhedral white K-feldspar phenocrysts) rock locally contaminated by abundant mafic inclusions. This rock appears to be a chilled facies of average Kcp.

A quartz monzonite with color index ranging from 2 to 7 percent, locally no hornblende and slightly more quartz than the average Kcp grades into and intrudes the more mafic quartz monzonite and granodiorite. For example, the rock which underlies Stonewall Peak is leucoadamellite with abundant 1 to 2 cm relict euhedral K-feldspar grains. Some of the leucocratic Kcp is alaskite and aplite that resembles the leucocratic facies of the younger quartz monzonite and granodiorite of Pine Valley (Kpv).

The thin-sheeted style of intrusion, extensive stoping and chemical reactions with mafic rocks, finer grain size and mafic mineral suite of Kcp help to distinguish it from Kpv.

Quartz monzonite and granodiorite of Pine Valley:--In plan view the quartz monzonite and granodiorite of Pine Valley (Kpv) forms large (several km across), discrete plutons which are relatively inclusion-free and uncontaminated compared to Kcp. Mapping indicates that the unit has been emplaced in steeply-dipping sheets.

The rock is white-weathering and underlies highlands. It is chiefly coarse-grained quartz monzonite with color index ranging from 5 to 10 percent. Mafic minerals are dark yellowish-brown biotite and small, skeletal relicts of dark bluish-green hornblende. Many samples contain no hornblende. The plagioclase feldspar is oligoclase that occurs as relict, euhedrally zoned grains. Prominent accessory minerals are sphene, allanite, and epidote. In the Arroyo Seco area (Cuyamaca Peak quadrangle) and in the town of Pine Valley (Descanso quadrangle) the rock contains white, relict euhedral K-feldspar 2 cm long and approaches porphyritic texture. Everywhere, elongate, gray, 2 to 3 cm recrystallized quartz lenticles, probably relicts of large igneous grains, are characteristic of Kpv. The unit locally has chilled margins against, and sends dikes into, all units except Kem, which it locally intrudes synplutonically. The unit is strongly foliated.

Quartz monzonite and granodiorite of Corte Madera:--The quartz monzonite and granodiorite of Corte Madera (Kcm) appears to be a leucocratic variant of Kpv which crops out in the Descanso and Viejas Mountain quadrangles. Although Kpv and Kcm are not shown in contact on these two maps, small amounts of Kpv occur locally in Kcm plutons in a gradational relationship to Kcm. Both Kcm and Kpv are coarse-grained and have elongate gray quartz aggregates up to several cm long. Kcm has the same age relationship to other units as Kpv and typically occurs as sheets in older plutons. The chief difference between Kcm and Kpv are that Kcm has a lower average color index than Kpv. Most parts of the Kcm bodies are leucoadamellite, leucogranodiorite or leucogranite. Kcm and Kpv are probably closely related to one another because of the gradational relationship, textural and petrographic similarities and similar age relationship to other plutonic units. Kcm grades into pegmatite, alaskite and aplite which locally occur as dikes in the unit.

Pegmatite, alaskite and aplite:--Leucocratic dikes (Kl) of pegmatite, alaskite and aplite occur in all units. In some areas they can be traced into plutons of Kpv, Kcm or Kcp. Where no association with larger bodies was established, the dikes have been mapped separately. These dikes share the metamorphic fabric of the other plutonic rocks.

Tonalite, quartz diorite and gabbro of East Mesa:--The tonalite, quartz diorite and gabbro of East Mesa (Kem) is the most texturally and compositionally heterogeneous plutonic unit. In plan view the sizes and shapes of Kem bodies vary because the unit has intruded older plutons in multiple sheets which have moderate to steep dips. The sheets are interconnected and commonly are localized along older contacts.

Tonalite and quartz diorite form the major part of the unit. The rocks are typically dark gray, fine- to medium-grained, and locally sub-porphyritic with relict subhedral phenocrysts of plagioclase and hornblende. A common textural variety has a spotted appearance due to poikilocrysts of biotite in a fine-grained groundmass. Pale tan to green hornblende is either the dominant mafic mineral or is about equal in abundance to pale reddish-brown biotite. Color index decreases in a regular manner as modal quartz increases. Color indices of quartz diorite samples range from 35 to 50 percent, while those of tonalites range from 25 to 35 percent. In some samples, hornblende has been altered to actinolite, and biotite to chlorite. Relict phenocrysts of andesine show strong oscillatory zoning with calcic cores. Medium-grained Kem carries abundant, fine-grained mafic inclusions, less than one foot long, some of which are elongate parallel to foliation but many of which are angular or irregularly shaped blocks.

Dark gray quartz diorite locally grades into fine-grained black dikes, some with scattered plagioclase relict phenocrysts and others choked with relict euhedral plagioclase grains.

A more leucocratic, sub-porphyritic variety of the Kem unit in the southern part of the Descanso 7½' quadrangle intrudes dark gray to black Kem. Distinctive pale greenish-white aplite dikes cut this leucocratic variant and were seen only in this area.

Kem whose hornblendes enclose pyroxene cores (opx -> cpx) grades into the less abundant gabbro of the unit. The gabbro is generally fine- to medium-grained. The distribution of tonalite, quartz diorite and gabbroic rocks within the Kem plutons shows no regular pattern. Although these plutons were not studied in detail, internal contacts were seen locally and undoubtedly the history of the unit is complex.

The Kem unit sends dikes into and has chilled margins against all plutonic units except Klb and Kmd. Kem is locally continuous with, and also cut by, dikes of Kmd. Synplutonic dikes of some host rocks into Kem give rise locally to apparent reversals of age relations. This is especially true where Kem intrudes Kcr and Kcp. In the southeastern part of the Descanso quadrangle, these three units are intimately interlayered, their synplutonic contacts have been deformed, and granitic rocks and Kem have been contaminated by one another. A stippled map pattern for Kem indicates that it contains up to about 50 percent of granitic inclusions, chiefly quartz monzonite.

The Kem bodies are strongly foliated, especially near their margins. The Deer Park body in the eastern part of the Cuyamaca Peak quadrangle, for example, consists almost wholly of mylonite gneiss.

Mafic dikes:--The youngest plutonic unit is an ubiquitous system of mafic dikes (Kmd) which cut all other plutonic rocks. Few are large enough to be shown at 1:24,000 map scale. The dikes are dark gray to black, mostly fine-grained to very fine-grained, but some have fine- to medium-grained centers with chilled margins, while others are variably porphyritic. The dikes consist of plagioclase, hornblende, biotite, sphene and traces of quartz, and overlap Kem in modal composition. Their textures are dynamothermal metamorphic, and only the plagioclase phenocrysts in porphyritic dikes show relict euhedral outlines and oscillatory zoning. The dikes are abundant in the vicinity of bodies of Kem, and locally grade into these bodies. They also cut Kem, and locally crosscut one another.

Although in some places the mafic dikes crosscut regional foliation, for the most part the largest dikes were emplaced concordant to foliation. With few exceptions, foliation of the dikes' mineral grains is parallel to the surrounding regional foliation, regardless of the dikes' orientation.

The mafic dikes have a synplutonic relationship with their host rocks. Most of the dikes are highly deformed; many are pulled apart and resemble inclusions. They typically contain folded bodies of aplite that in some cases are continuous with the granitic host rock but mostly are not.

Surficial deposits:--Surficial deposits have been divided into alluvium, colluvium and landslide materials. Although most of the deposits have not been dated, they are probably Quaternary in age. Alluvium (Qal) consists of sand, silt, clay and gravel in stream valleys and meadows. The alluvium in stream valleys consists of older deposits which cover the valley floors and thin modern deposits in the beds of narrow channels that cut the older alluvium to depths of up to 6 meters. A preliminary C¹⁴ age of 920 ± 60 years has been obtained on charcoal from one of the lowest exposed beds in older alluvium in Pine Valley (Stephen W. Robinson, U.S. Geological Survey, Menlo Park). This data indicates only that the older alluvium is no older than 920 ± 60 years, because the charcoal may have been reworked from midden deposits of Indians living in Pine Valley. Considerable erosion of older alluvium has probably occurred in historic times, as suggested by the headward cutting of gullies 1 to 2 meters deep along wagon ruts in a few places. Since bedrock is exposed in most of the modern streambeds, the total thickness of older alluvium is probably about 6 m or less. Fine-grained sediments in the high meadows probably formed in situ by chemical weathering in addition to some stream deposition. These sediments are being stripped from the meadows by headward-cutting gullies and most meadows contain bedrock exposures.

Colluvial deposits, Qc, consist of poorly sorted sand, silt and gravel mixtures and formed by slope wash; they grade into bouldery deposits that fill small, steep valleys. The latter are probably debris flow deposits and locally, they form topographic ridges which are being eroded by modern gullies. Colluvial deposits have been mapped only where they are thick enough to completely obscure bedrock outcroppings over a large area. Thick colluvium deposits commonly are associated with fault-steepened topography. In general, their thickness is no more than a few meters, indeed, most slopes are bedrock which is essentially bare of colluvium or soil.

Landslide deposits (Qls) are of the slump and rockslide type and occur in terrain which has been oversteepened by faulting. The landslides are small (50-250 m wide) and are characterized by steep scarps, which are free of colluvium and vegetation, above the landslide deposits.

Structure of batholithic rocks:

The plutonic units occur as steeply dipping sheets and lenticular bodies which are separated locally by screens of metasedimentary rock. The sheets, lenticular bodies and screens range from a few meters to several kms in thickness and the larger ones continue for tens of kms along strike. In plan view, the preferred orientation of the long dimensions of plutonic sheets and lenticular bodies, of Tm screens, and of foliation within plutonic and metamorphic rocks, imparts a structural grain to this part of the batholith. Only a small part of this structural grain can be seen in any one 7½' quadrangle. Successive intrusions parallel to this structural grain have resulted in stratiform complexes of 3 to 4 units. The structural grain varies greatly over the project area.

In the Cuyamaca Peak quadrangle, it is north-northwest to northwest and the dip is predominantly eastward, with the exception of the eastern margin of the Cuyamaca Mountains gabbro body whose walls dip inward. In the Descanso quadrangle, the structural grain is north-northwest in the eastern half of the quadrangle and northwest to east-west in the western half; the regional dip is strongly to the northeast. Locally, plutonic contacts and foliation describe large fold forms about steeply plunging axes. Several of these fold forms include metasedimentary screens which are folded concordantly with plutonic contacts and foliation. One such structure, involving Kc and Kcp, occurs just west of the center of the Cuyamaca Peak quadrangle. In the Descanso quadrangle, one involving TRm and Kcr occurs in the center of the map; a second, involving Kc, Kcp and TRm, lies south of it. It is not clear at this time whether these structures are tectonic, i.e., systematic folds, or relicts of local deformation due to plutonic intrusion. Possibly, they are the result of a combination of the two processes.

Locally, the dips of plutonic contacts appear to be gentle to moderate, e.g., the contacts of the Rattlesnake Mountain pluton in the northeastern part of the Cuyamaca Peak quadrangle, and the Kc-Kcp contacts in the Buckman Mountain area, southeastern corner of the Descanso quadrangle. Here, foliation dips more steeply than plutonic contacts. The dips of plutonic contacts may appear less steep than they truly are because of a combination of the sheeted style of intrusion and poor exposure (dense vegetation). Apparently, magma moved upward utilizing steep planes of entry such as bedding in TRm and contacts and foliation in older plutonic rocks. Thus, TRm and older plutonic rocks tend to occur as screens which are underlain by and separated by younger plutonic rock. Where the present level of erosion is close to the highest level to which a younger magma intruded (and therefore to the keel-shaped bottoms of screens of TRm and older plutonic rocks), valleys expose chiefly the underlying, younger plutonic rocks. If exposure is limited, an approximate contact drawn between the younger plutonic rock and the host rocks may cut across many thin screens and thus appear flatter than it truly is.

If, as seems likely, the Cuyamaca Gabbro is the oldest plutonic unit, then the present Kc bodies are not individual plutons but are remnants of original plutons. The distribution of TRm remnants (probable wallrocks) in relation to Kc over the project area suggests that while some present Kc bodies coincide approximately with original plutons, others may not. The Cuyamaca Mountains body may be one of the former in which case it had a north-northwest-elongate shape originally. Younger granitic plutons apparently have divided the original gabbro plutons into a number of lobes--in the case of the Cuyamaca Mountains body, a northern lobe consisting of Middle Peak and North Peak, and a southern, Cuyamaca Peak lobe. The gabbro body north of Corte Madera Mountain in the Descanso quadrangle also shows partial division into lobes by granitic plutons. In the eastern part of this quadrangle there is a bead-like arrangement of Kc bodies in a linear array extending north-northwest from Buckman Mountain through Bear Valley to Guatay Mountain. This arrangement suggests boudinaging of gabbro in a granitic matrix which may have been more mobile under metamorphic conditions because of the presence of the empirically "weaker" minerals quartz and mica.

Faults of the Descanso 7½' quadrangle:

Introduction:--Figure 5 shows the locations of fault zones and lineaments to be discussed. Two zones of faulting occur in the Descanso quadrangle. One is a diffuse zone in the eastern third of the quadrangle which is the probable southern extension of the north-south-trending, en echelon Chariot Canyon-Oriflamme Canyon fault zone (Julian, Monument Peak and Cuyamaca Peak 7½' quadrangles).

The second fault zone is exposed in Pine Creek and Secret Canyons, in the southwestern part of the Descanso quadrangle and southeastern Viejas Mountain quadrangles, respectively, and is part of a complex zone that includes faults in and near Horsethief Canyon. Several faults extend into the northwestern corner of the map in the valley of the Sweetwater River. They represent the southern end of the Cuyamaca Mountains fault zone (Cuyamaca Peak and Tule Springs 7½' quadrangles) and are discussed in the map explanation for the Cuyamaca Peak quadrangle.

Faults in both zones within the Descanso quadrangle have youthful topographic expression--alined linear valleys and ridges, benches, notches, scarps, hillside valleys, vegetation stripes and gullying--and are therefore tentatively considered as Quaternary in age. Some faults in the Pine Creek-Secret Canyon zone appear to crosscut colluvial deposits.

Faults of the Pine Valley - Cottonwood Valley Zone:--The faults occur in groups of several short, steeply-dipping breaks which strike from north-northwest to N40E, but typically north-northeast to north-northwest. Many faults, the longest a few km in length, parallel the older gneissic foliation for at least part of their length. The zone passes through the town of Pine Valley and continues southeastward into the valley of Cottonwood Creek, the site of recently completed Interstate Highway 8 (Mount Laguna 7½' quadrangle). The largest faults found to date occur along the walls of Cottonwood Valley.

The town of Pine Valley and the area to the north is underlain predominantly by strongly foliated and jointed quartz monzonite which presents a bewildering array of linear features on intermediate scale aerial photographs. Features found to be associated with faults are alined linear valleys, benches, notches, scarples, gullies and vegetation stripes. Where gabbro (Kc) is faulted, the rock weathers to light-colored greenish-white hues which may be highly visible from a distance. The number and array of faults shown on the map depend to a considerable extent upon accessibility for there are few through-going trails or roads and the steep quartz monzonite terrain is extremely brushy. Undoubtedly there are more breaks in the area than were located during mapping. Width and nature of crush zones are variable, most in the Pine Valley area consisting of several meters of brownish, gougey gneiss (the gneissic texture is pre-faulting), locally with discrete, thin (<10 cm), anastomosing bands of mylonite and/or gouge. The thickest mylonite and gouge observed is about one meter. Thus, none of these faults appear to be large breaks, unless the linear valley of Pine Creek itself was eroded along a major fault.

The faults bounding the linear two thousand feet-deep valley of Cottonwood Creek (south of the Sunrise highway-Interstate 8 junction) may be more important. They are marked by prominent alined hillside valleys, benches, notches, vegetation stripes and gullying. Although not all of the lineaments of this zone expose gouge, one in the southeastern corner of the Descanso quadrangle is marked by at least 3 to 4 m of continuous gouge.

The faults in the Buckman Mountain area (southeastern part of quadrangle) appear to have chiefly vertical displacements. However, because the faults are roughly parallel to the structural grain and because of the complex intrusive pattern in this area, small lateral displacements would be difficult to prove. There is no single through-going fault in the eastern third of the Descanso quadrangle, and the fault pattern is suggestive of normal faulting.

Faults of the Secret Canyon zone:-- The faults which define the southwest-trending reaches of Pine Creek Canyon in the southwestern part of the Descanso quadrangle, and of Secret Canyon in the Viejas Mountain quadrangle trend about N30E. They are very strongly marked by alined linear valleys, ridges, scarps, benches, notches and vegetation stripes. Vegetation stripes continue from bedrock through Qc deposits. The north wall of Pine Creek Canyon about 120 m east of its junction with Nelson Canyon (where Pine Creek turns southwest) exposes approximately 8-9 m of vertical gouge which separates Kc on the east from K1b on the west. Other less accessible lineaments in the zone are marked by mylonite and gouge. Although there is evidence for as much as 200 m of apparent right-lateral offset along the fault zone, some vertical displacement is required to explain the absence of Kcm east of the zone, as well as to explain the distribution of K1b across the faults. The pattern of multiple short breaks of slightly different strikes also suggests vertical slip.

Many short east-west and west-northwest-striking lineaments are present in this general area; some are marked by mylonite and gouge. The eastern end of a fairly large east-west fault extends into Espinosa Creek Canyon in the extreme southwestern corner of the Descanso quadrangle. Some of the east-west mylonite zones appear gradational to the older gneissic structure. Others appear to curve and join north-south faults. A better understanding of the complex faults of the area awaits further study.

Relationship of major lineaments to faulting:--The Sweetwater River-Horsethief Canyon lineament has been described by Merifield and Lamar (1976) as the second most prominent linear feature on spacecraft photography of the Peninsular Ranges. The lineament extends northeastward from the international border approximately to the Elsinore fault zone, a distance of about 50 km. It consists of closely en echelon, approximately linear stream valleys. The faults in the Sweetwater River valley (northwestern Descanso quadrangle) do not appear to cross Descanso Valley since several northwest-trending lithologic contacts are not offset. No crushed rock was seen in Descanso Valley.

The Pine Valley-Oak Valley lineament is highly visible on 1:62,500 aerial photographs and consists of several alined, linear stream valleys. It extends north-northeast from Corte Madera Valley about 15 km to the Deer Park mining district. In Oak Valley the lineament is crossed by a number of geologic contacts which it does not offset.

REFERENCES CITED

- Everhart, D. L., 1951, Geology of the Cuyamaca Peak quadrangle, San Diego County, California: California Div. Mines Bull. 159, p. 51-115.
- Jahns, R. H., 1954, Geology of the Peninsular Range province, southern California and Baja California, in Geology of Southern California, California Div. Mines Bull. 170, Sept. 1954, p. 29-52.
- Merifield, P. M., and Lamar, D. L., 1976, Fault tectonics and earthquake hazards in parts of southern California: Technical report 76-1, NASA Lyndon B. Johnson Space Center, Houston, Texas.
- Streckeisen, A. L., 1973, Plutonic rocks, classification and nomenclature recommended by the I.U.G.S. Subcommittee on the Systematics of Igneous Rocks: Geotimes, v. 18, no. 10, p. 26-30.

Williams H., Turner, F. J., and Gilbert, C. M., 1954, Petrography, an introduction to the study of rocks in thin sections, W. H. Freeman and Co., San Francisco, 406 pp.

ACKNOWLEDGMENTS

The field study would not have been possible without the cooperation of rangers of the Cleveland National Forest (U.S. Forest Service) and the Cuyamaca Rancho State Park (California Department of Parks and Recreation) who gave advice and practical aid whenever asked. Landowners who have generously allowed access to their property are too numerous to name--their cooperation made the study possible, and their warmth and interest helped to make it a rich personal experience as well. In particular, the owners and foreman of the Corte Madera Ranch generously provided access to a very large and important area.




Figure 1.--Peninsular Ranges batholith in southern California and Baja California and project area (after Jahns, 1954).




Figure 2a.--Kpv outcrop, trace of foliation parallel to pencil. Color index appears higher than 5-10 percent because mafic minerals have broken down and recrystallized into fine-grained aggregates.




Figure 2b.--Slabs cut at right angles to foliation. Left, Kcp granodiorite; right, Kpv quartz monzonite. Stained for K-feldspar and plagioclase; 6-inch scale.




Figure 3.--Sketch diagram of synplutonic relation between Kem dike and Kep host rock.




Figure 4a.--Modal data for granitic rocks from Guyamaca Peak 15' quadrangle (classification according to Streckeisen, 1973).





Figure 4b.--Modal data for granitic rocks from Cuyamaca Peak 15' quadrangle (classification according to Williams and others, 1954).



FAULT ZONE



LINEAMENTS

- GV - S - H : Green Valley -
Sweetwater River -
Horsethief Canyon
- PV - OV : Pine Valley - Oak Valley
- CV : Cottonwood Valley

Figure 5.--Index map showing lineaments, fault zones, and geographic features in and around Cuyamaca Peak and Descanso quadrangles. $7\frac{1}{2}'$ quadrangles: TS - Tule Springs, CP - Cuyamaca Peak, VM - Viejas Mountain, D - Descanso, MP - Monument Peak, ML - Mount Laguna, J - Julian)