

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

MAPS AND TABLES DESCRIBING METALLIFEROUS
MINERAL RESOURCE POTENTIAL OF SOUTHERN ALASKA

By E. M. MacKevett, Jr., D. A. Singer, and C. D. Holloway

TO ACCOMPANY

Geological Survey Open-File Report 78-1E

This report is preliminary
and has not been edited or
reviewed for conformity with
Geological Survey standards
and nomenclature

Menlo Park, California
1978

MAPS AND TABLES DESCRIBING METALLIFEROUS MINERAL RESOURCE
POTENTIAL OF SOUTHERN ALASKA

Explanatory text to accompany U.S. Geological Survey open-file report 78-1-E

INTRODUCTION

This report is the culmination of a regional mineral resource appraisal of southern Alaska by the U.S. Geological Survey. It consists of two maps, designated sheets 1 and 2 of open-file 78-1-E, descriptive and documentary tables that supplement the maps, and this explanatory text. Sheet 1 pertains to that part of southern Alaska east of the 153° meridian and north of the 59° parallel and sheet 2 to the western part of southern Alaska. South of the 59° parallel the eastern boundary of sheet 2 is the 152° meridian. Elsewhere the eastern boundary is the 153° meridian. As used in this study, southern Alaska includes a large area that extends northward from the Pacific Ocean to an irregular boundary that roughly parallels the northernmost forelands of the convex northward, arcuate Alaska Range (see supplementary index maps on the accompanying maps). From its apical region in the Fairbanks quadrangle, the northern boundary extends southwestward to Bristol Bay and southeastward to near latitude 63° 30' at the Canadian border. The easternmost and westernmost extremities of southern Alaska (in our usage) are, respectively, the 138° meridian and Unimak Pass. The purpose of this report is to provide a current and thorough appraisal of the known and potential metallic mineral resources of southern Alaska that utilizes the best geologic and mineral resource data available.

The maps (sheets 1 and 2) show outlines of favorable areas for

metalliferous mineral resources that are mainly based on known deposits and favorable geology for specific deposit types. Forty-three favorable areas are outlined on sheet 1 and thirteen on sheet 2. Supplementary tables that are keyed numerically to outlined areas on the maps describe the known and speculative deposit types in each outlined area, summarize available data on geology, production, reserves, and status of geologic knowledge, and provide the resource estimates, which are the basic objectives of this study. These tables are designated tables 1 (p.33) and 2 (p.34) and, respectively, refer to sheets 1 and 2. Another table (table 4 (p.45)) summarizes the probabilistic grade and tonnage models for specific deposit types.

Background data for this report have been published separately as a folio of open-file reports (table 3 (p. 2)). Those reports, which include pertinent references and other relevant information, are components of a folio of basic data that constitutes the foundations for this report.

Table 3. Component maps of the regional mineral resource appraisal of southern Alaska

Eastern southern Alaska

<u>U.S. Geological Survey open-file map</u>	<u>Subject</u>
OF-77-169-A (MacKevett and Holloway, 1977)	Metalliferous and selected nonmetalliferous mineral deposits
-B (Beikman, Holloway, and MacKevett, 1977)	Generalized geology
-C (Barnes, 1977)	Gravity data
-D (Holloway, 1977)	Coal
-E (Decker and Karl, 1977)	Aeromagnetic data

Western southern Alaska

U.S. Geological Survey open-file map

OF-77-169-F (MacKevett and Holloway, 1977)	Metalliferous mineral deposits
-G (Beikman, Holloway, and MacKevett, 1977)	Generalized geology
-H (Barnes, 1977)	Gravity data
-I (Holloway, 1977)	Coal
-J (Decker and Karl, 1977)	Aeromagnetic data

Fossil fuels, geothermal energy sources, and nonmetallic mineral commodities are not within the purview of this report. However, the folio of basic data includes descriptions of a few deposits of nonmetallic minerals in eastern southern Alaska, plus maps and tables that summarize coal deposits in southern Alaska.

RESPONSIBILITY AND ACKNOWLEDGMENTS

This report represents the combined and cooperative product of the authors. MacKevett and Holloway were largely responsible for geologic descriptions of deposit types and related data such as production, reserves, and status of geologic knowledge for a given area; MacKevett determined extents and configurations of the favorable areas; and Singer was mainly responsible for the resource estimates and appraisals.

The authors are indebted to many people, mainly U.S. Geological Survey colleagues, who facilitated the preparation of this report and the companion reports that provide the fundamental background materials. We are especially grateful to E. H. Cobb for his useful inventories of Alaskan mineral deposits; to B. L. Reed for sharing his extensive knowledge

of the geology and mineral deposits of the western Alaska Range; and to W. D. Menzie for his contributions in developing models for specific deposit types.

PHILOSOPHY AND LIMITATIONS

Our investigation represents a thorough attempt to use the best available and most current relevant information to derive objective mineral resource estimates for southern Alaska. Even so, some disparities exist in our basic data and, correspondingly, in the derivative resource estimates. For example, some areas are geologically poorly known and have been scantily prospected, whereas a few others are geologically well known and locally well prospected. Documentation for individual deposits ranges from a few sentences in old reports that cursorily allude to a deposit to a few modern scientific reports that provide thorough descriptions. Nevertheless, the basic geologic framework of southern Alaska and the types and geologic settings of the region's mineral deposits are reasonably well known.

In a broad sense, just about every area on earth has some resource potential, regardless of how remote or insignificant such a potential may be. In this study only the potentially significant resource areas are identified, delineated, and described; the other areas being excluded after carefully evaluating the basic data. Many of the excluded areas are mantled by thick covers of younger unfavorable rocks, glaciers, or unconsolidated surficial deposits, and even though they may contain concealed deposits at depth, the chances for discovering and exploiting such deposits are minimal.

Speculative or suspected deposit types, one of the criteria used in determining the favorable areas, are inferred from their occurrences

in similar geologic settings elsewhere. A more comprehensive use of this category might be desirable, but to be meaningful, it should be founded on more detailed geologic information than is generally available for southern Alaska. Such deposit types include some that have been known for many years in some other parts of the world and a few others, such as volcanic-type nickel deposits and various types of uranium deposits that have been recognized only recently.

Among the factors worth considering in estimating the mineral resource potential of southern Alaska are:

- (1) Southern Alaska is well endowed with a variety of mineral deposits commensurate with its diverse geology
- (2) With a few exceptions, notably for placer gold, southern Alaska is scantily prospected by modern standards, and the vast majority of known deposits are too poorly explored to permit precise evaluations
- (3) Potentially significant new discoveries have been made in the region during the past decade, notably the extensive belt of submarine volcanogenic base metal-silver deposits along the north flank of the Alaska Range and the copper-molybdenum porphyry province of the Alaska Peninsula and nearby islands; such discoveries augur the continued success of thorough modern exploration
- (4) Southern Alaska contains known deposits of several metals of current national interest, for example, chromium and tin, and it may contain significant resources of these commodities
- (5) Some of the large covered tracts, both within and beyond areas designated as favorable, may contain concealed deposits at shallow depths that are amenable to discovery and exploitation

- (6) Possibly some of the region's diverse known or undiscovered metals may be of future importance in supplying metals for new uses brought about by technologic advances
- (7) Although no assuredly significant uranium deposits are known in southern Alaska, the region contains many geologic settings that are favorable for a variety of uranium deposits, and systematic prospecting for uranium is warranted in some areas .
- (8) Extensive tracts of southern Alaska are geologically poorly known. Some contain geologic settings favorable for significant mineral deposits, and more thorough geologic knowledge of these areas would substantially increase the validity of future mineral resource estimates.

In order for this report to be useful, the purpose of the analysis had to be considered in the design of the resource appraisal (Singer, 1975). The purpose in this case is primarily to provide mineral resource information that can be used in the land classification decisions of Alaska. To achieve this, it is desirable to delineate individual tracts of land and to differentiate them on the basis of their potential for containing mineral resources. For each tract it is also desirable to indicate the quality and quantity of mineral resources with respect to the factors that affect possible economics and technologies of exploitation. Ideally, these factors include grade and tonnage estimates, the physical, chemical, and mineralogical features of the mineralized rock that could affect its treatment and recovery, and whether all of the mineralized rock has been found.

Information concerning many of these factors is probably best conveyed by using mineral deposit types as a basis for the estimates, as we have done. In many cases, deposit types have distinct physical, chemical, and mineralogical features, and some can be characterized as having restricted ranges of grades and tonnages. In addition, because deposit types tend to have certain geologic associations, the resource appraisal can be made relatively straightforward and readily explainable. Estimates of grades and tonnages of similar well explored deposits can be used as models of the incompletely explored and, in many cases, undiscovered deposits of Alaska (table 4).

METHODOLOGY

This report augments the fundamental mineral resource, geologic, and related information in the folio of basic data (table 3) by utilizing various mineral resource appraisal methods in order to fulfill its objectives. In essence, the favorable mineral resource areas are outlined on the basis of their known deposits, including principal occurrences, and their favorability for undiscovered or speculative deposits. No attempt is made to rank the outlined areas relative to their degrees of favorability, but the general potential and rank of a given area can be ascertained from descriptions in the tables. The potential for undiscovered deposits is regarded as a function of favorable geology and, in some cases, supplementary favorable geochemical or geophysical data. The outlined favorable areas and the metals for which they are noteworthy are shown on the accompanying maps. Symbols for the less significant metallic constituents that generally constitute byproducts or potential byproducts are

enclosed in parentheses. Succinct descriptions of the deposit types in the outlined areas are given in the accompanying tables (tables 1 and 2); these tables describe the contained metals, geologic settings, and other information relevant to the deposits. The tabulated descriptions are keyed numerically to the maps. Generally used nomenclature for deposit types, for example porphyry, vein, submarine volcanogenic, and contact metamorphic, are used in this report. Many of these have genetic connotations.

The mineral resource estimates, which are the crux of this report, are derived by integrating and objectively evaluating all available germane data. Mineral resource data for each favorable area outlined on the map are shown in tables 1 and 2. The mineral resource estimates supplement what is known by incorporating a variety of pertinent considerations, such as degrees of geologic, geochemical, and geophysical favorability, extent and adequacy of exploration and geologic knowledge, and, for some deposits, indications of sizes and grades extrapolated from models of better-known deposits of a specific type (table 4 (p.45)). In most cases the basic data are insufficient to justify more than qualitative resource estimates. However, in some instances the data are adequate to permit more quantitative estimates of the number of deposits of a specific type that may be present in a given area and their probable grades and sizes.

The general procedure followed in deriving the resource estimates consisted of: (1) using geology to delineate areas that either have known deposits of a particular type or areas that are favorable for containing them, (2) where possible, providing information on grades and tonnages of similar deposits based upon careful study of the geology and grades and tonnages of well explored deposits, and (3) where possible, subjectively

estimating the number of deposits of each type in each delineated area using the number of known deposits, the amount of favorable geology, the extent of exploration, and in some cases supplementary geochemical and geophysical data.

Estimates of grades and tonnages and of the number of deposits are presented in a range of probabilities. Probabilistic estimates of grades and tonnages (table 4) demonstrate the range of values observed for each deposit type; correlations among grades and tonnages are presented in order to show the degree of linear association between grades and tonnages. Significant correlations mean that probabilities of different grade and tonnage combinations must be calculated based on consideration of both variables, while non-significant correlations mean that the probability of a grade-tonnage combination can be calculated as the product of the two probabilities. Probabilistic estimates of the number of deposits show the degree of certainty that we have concerning the number of deposits that might occur in an area. Typically, estimates of the number of deposits are made only for deposits with tonnages and grades comparable to those used in the grade-tonnage model listed in table 4. Also, estimates are made for a few deposits that lack associated grade-tonnage models.

CONCLUSIONS

Southern Alaska is well endowed with a large variety of mineral deposits. Favorable areas for these deposits are outlined on the accompanying maps and individually described in the accompanying tables. Tables 1 and 2 contain the basic resource estimates and some of the supporting data used in deriving the estimates. Additional documentary data are in map components of a folio of basic data (table 3) that should be used in conjunction with this report.

The outlined areas include potentially significant deposits of many types that contain an array of metal commodities. Discrete deposit types are described in the tables. In current economic context, probably the most significant deposits in southern Alaska are the porphyry-type deposits for copper and(or) molybdenum and the submarine volcanogenic deposits mainly for copper, silver, and zinc. However, the region contains numerous examples of many different deposit types that cumulatively contain a large variety of metals. Many of the known deposits, their undiscovered counterparts, and possibly some deposit types not presently known in the region, are of potentially important economic significance.

REFERENCES

- Barnes, D. F., 1977a, Gravity map of the eastern part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-C, 1 sheet, scale 1:1,000,000.
- 1977b, Gravity map of the western part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-H, 1 sheet, scale 1:1,000,000.
- Beikman, Helen M., Holloway, C. D., and MacKevett, E. M., Jr., compilers, 1977a, Generalized geologic map of the eastern part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-B, 1 sheet, scale 1:1,000,000.
- 1977b, Generalized geologic map of the western part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-G, 1 sheet, scale 1:1,000,000.
- Decker, John, and Karl, Susan, compilers, 1977a, Preliminary aeromagnetic map of the eastern part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-E, 1 sheet, scale 1:1,000,000.
- 1977b, Preliminary aeromagnetic map of the western part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-J, 1 sheet, scale 1:1,000,000.

Holloway, C. D., 1977a, Map showing coal fields and distribution of coal-bearing rocks in the eastern part of southern Alaska: U.S. Geol.

Survey open-file map OF 77-169-D, 1 sheet, scale 1:1,000,000.

-----1977b, Map showing coal fields and distribution of coal-bearing rocks in the western part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-I, 1 sheet, scale 1:1,000,000.

MacKevett, E. M., Jr., and Holloway, C. D., 1977a, Metalliferous and selected nonmetalliferous mineral deposits in the eastern part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-A, 1 sheet, scale 1:1,000,000.

-----1977b, Metalliferous mineral deposits in the western part of southern Alaska: U.S. Geol. Survey open-file map OF 77-169-F, 1 sheet scale 1:1,000,000.

Singer, D. A., 1975, Mineral resource models and the Alaskan mineral resource assessment program, in Vogely, W. A., ed., Mineral material modeling: Washington, D. C., Johns Hopkins Univ. Press, p. 370-382.

(a) The combination of several known minor occurrences in the few areas not covered by glaciers suggest many of these small deposits might be covered by physical anomalies in altered granitic rocks suggest the possibility of porphyry molybdenum deposits.

(b) porphyry molybdenum model

A number of gold veins probably occur in this area

Large tonnage, low grade iron and titanium placers are known, low tonnage gold placers are in part ephemeral are also present.

(e) Gold-bearing beach placers that vary in quality yearly due to winter storms are known (f) Stream and bench gold placers that have been mined; resources remaining are unknown

mafic volcanogenic model

One known occurrence plus other possible glacially covered unfound mafic volcanogenic copper deposits.

Typically estimates of the number of deposits are made only for deposits with tonnage and grades comparable to those used in the grade-tonnage models. Also estimates are made for a few deposits that lack associated grade-tonnage models.

ADDITIONAL COMMENTS

Area 1 is in a remote and rugged part of the St. Elias Mountains that is largely covered by glaciers; no significant mineral deposits are known in the area

The area consists of a partly glacier-covered mountainous region between higher terrain of the St. Elias Mountains and the Yakutat Foreland; the less metamorphosed rocks southwest of the Boundary Fault are regarded as more favorable for gold lodes than the dominantly amphibolite terrane between the Boundary and Boundary Faults

The placers that are mainly for gold are small and in part ephemeral; the iron-titanium placers are large and extend intermittently for more than 20 km along beaches fronting the Gulf of Alaska; they consist of black sands that contain titanium (about 10 percent) and iron (about 10 percent) and are between 1 and 3 m in thickness, and although they contain local small higher-grade zones, their overall grade proximates that given in the "Production and resource information" column

The boundaries of areas 4 are inaccurately known and the area may include recent unfossiliferous placer operations on the perimeter of Icy Bay and westward to include some beaches near Cape Suckling

Area 5 is delineated on the basis of favorable geology—mainly submarine basalts of the Orca Group—and one known occurrence

STATUS OF GEOLOGIC INFORMATION

Broad reconnaissance mapping and widely spaced geochemical sampling by U.S. Geological Survey; essentially unprospected

Reconnaissance mapping and geochemical sampling by U.S. Geological Survey; scant prospecting

Reconnaissance mapping, some geochemical sampling, and local aeromagnetic coverage by U.S. Geological Survey; investigations involving auger-hole drilling and sampling by U.S. Bureau of Mines; scant recent interest by industry

Old and modern, largely reconnaissance, mapping by U.S. Geological Survey; scant aeromagnetic potential of radioactive heavy minerals in the beach sands; sampling of beach sands by U.S. Bureau of Mines

Broad reconnaissance mapping and scant geochemical sampling by U.S. Geological Survey; essentially unprospected

PRODUCTION AND RE-SOURCE INFORMATION

No data

No data

Minor gold production, probably about 6 kg (several hundred ounces), during early 1900's, from small deposits; large, low-grade iron and titanium resources having a general tonnage of 30,000 kg (66,000 lbs) of iron (35 lbs/yr) and 12.2 kg of titanium dioxide per cubic meter (20.5 lbs/yr)

(a) Worked intermittently since 1890 for a total production of about 470 kg and 500 kg (15 and 16 thousand ounces) of gold (b) Mined for a few years during early 1900's; production not accurately known; probably between 30 kg and 60 kg (1 and 2 thousand ounces) of gold

No data

GEOLOGIC CONTROL(S) OF MINERAL RESOURCES

(a) Interpreted as metamorphic deposits whose metals were redistributed and weakly concentrated during metamorphism (b) Probable late-stage differentials of shallow plutons

Typically thin gold-bearing quartz veins that are localized in greenschist or lower grade metamorphosed rocks; sparsely and genetically related to Tertiary plutons

Modern beach and older marine terrace placers; the gold placers are best developed in the vicinity of Yakutat; the iron-titanium placers, which generally contain traces of gold, are best developed on beaches and forelands southeast of Yakutat

(a) Gold-bearing black sands that are interbedded with siltstone beaches fronting the Gulf of Alaska; largely ephemeral deposits concentrated during winter storms (b) Stream and bench placers localized by fluvial processes

Polymetallic copper-rich deposits genetically related to submarine basalts of the Orca Group (Tertiary) and less commonly Valdez Group (Cretaceous)

SUSPECTED OR STREAM-LIVE TYPES OF MINERAL DEPOSITS (UNLESS MINOR OCCURRENCES)

(a) (Cu, Zn, Au)-minor occurrences associated with disseminated pyrite in greenstone and (b) (Cu, Zn, Au)-minor occurrences associated with altered zones in granitic rocks; may represent porphyry type deposits

Au(Ag)—mafic quartz veins in Cretaceous metamorphosed gneiss

Au, Fe, Ti—beach and older marine terrace placers

(a) Au—beach placers and (b) Au—stream and bench placers

Cu(Ag, Pb, Zn)—submarine volcanogenic deposits related to mafic basalts

AREA OUTLINED ON MAP

1.

2.

3.

4.

5.

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
6.	--- Au--thin quartz veins in slate and graywacke	(a) Cu(Ag, Au, Zn)--submarine volcanogenic deposits related to (b) Au--placer	(a) Same as 5 (above) (b) Beach placers and possibly stream or bench placers	No data	Reconnaissance mapping by U. S. Geological Survey; little prospecting	Area 6 contains rocks favorable for submarine volcanogenic deposits but no known deposits of this type; it contains one or more placer and some permissive terrane for other placer gold deposits; the area may extend southward beneath the Gulf of Alaska to include Orca Group volcanic rocks on the north-west tip of Kayak Island	(a) Undiscovered mafic volcanogenic copper deposits may occur (b) One gold prospect on a beach placer and other placer gold deposits	90% 50% 10% 1 2 4 chance that there are or more	(a) mafic volcanogenic model
7.	Au--thin quartz veins in slate and graywacke	---	Gold-bearing quartz veins and veinlets in Orca Group (Tertiary) slate and graywacke; near Tertiary granitic pluton	Minor production, probably about 6 kg (several hundred ounces) of gold, from one property during early 1900's	Old Federal Government reports based on brief examination; reconnaissance mapping by U. S. Geological Survey; little recent interest by industry	Contains one inactive mine and one prospect; parts of the surrounding area may contain similar deposits, but they are largely covered by glaciers or unconsolidated surficial deposits	The known gold veins; similar undiscovered covered deposits possible	90% 50% 10% 1 2 4 chance that there are or more	(a) mafic volcanogenic model
8.	(a) Cu(Ag, Au, Zn)--submarine volcanogenic quartz veins in Valdez Group (b) Au--mainly quartz veins in Valdez Group (c) Au--placer	Cu--magmatic deposits with weakly disseminated pyrrhotite and chalcopyrite in Tertiary diorite	The area is largely underlain by the Cretaceous Valdez Group, including abundant mafic submarine volcanic rocks (a) Typically localized in shear zones in or near the volcanic rocks (b) Quartz stringers and veins, generally less than 1 m thick, genetically related to Tertiary plutons (c) Stream placers	Only production was from the Midas mine, which produced more than 450 tons (a million pounds) of copper; the main ore zone at the Midas is about 1 m wide and 300 m long and contains some reserves	Reconnaissance geologic mapping by U. S. Geological Survey; brief studies of a few deposits; recent exploration interest by industry at the Midas mine and probably nearby areas	Area 8 is outlined mainly on the basis of its potential for submarine volcanogenic deposits; in addition to the Midas mine the area has 7 prospects on mafic copper deposits; its potential for mafic copper deposits is much less than for submarine volcanogenic copper deposits; about half of the area is glacier covered	(a) At least eight mafic volcanogenic deposits are known to be present; they remain to be found in the exposed bedrock and under ice. Estimated number of deposits is for deposits comparable in tonnage to those used in the grade-tonnage model. (b) A few low tonnage gold-quartz veins might occur in this area (c) Two small gold-bearing stream placers are known	90% 50% 10% 1 2 4 chance that there are or more	(a) mafic volcanogenic model
9.	(a) Cu(Ag, Au, Zn)--submarine volcanogenic (b) Au--quartz veins in Orca Group (c) Au--placer	---	A near-coastal area that is underlain by the Tertiary Orca Group and by Tertiary anatectic granitic plutons (a) The submarine volcanogenic deposits are located in near mafic lavas of the Orca Group (b) Veins and veinlets in Orca Group flysch (c) Stream placers	No production or resource data	Reconnaissance geologic mapping by U. S. Geological Survey; little recent interest by industry	The area appears to be sparsely mineralized; its known deposits include four for copper and two for gold. It is geologically favorable for additional similar deposits	(a) At least four known mafic volcanogenic deposits; others possible (b) One gold-quartz vein deposit; other small tonnage veins possible (c) Possibility of small stream gold placers; one deposit known	90% 50% 10% 1 2 4 chance that there are or more	(a) mafic volcanogenic model

AREA OUTLINED ON MAP

MAJOR TYPES OF KNOWN DEPOSITS

SUSPECTED OR SPECULATED TYPES OF MINERAL DEPOSITS (INCLUDING MINOR OCCURRENCES)

GEOLOGIC CONTROL (S) OF MINERAL RESOURCES

PRODUCTION AND RESERVE SOURCE INFORMATION

STATUS OF GEOLOGIC INFORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL RESOURCE POTENTIAL

ESTIMATED NUMBER OF DEPOSITS THAT WOULD BE THE NUMBER PRESENTED OR MORE DEPOSITS

GRANDES AND TONNAGES FOR THIS REPORT (TABLE 2)

10. (a) Cu(Ag,Au,Zn)--submarine volcanogenic (b) Au--quartz lodes in Orca Group (c) Zn(Au,Ag,Cu)--breccia cemented by sulfides

Contains the most important submarine volcanogenic deposits of the Prince William Sound area; area underlain by Orca Group mafic volcanic rocks and scattered Tertiary felsic plutons

Between 1900 and 1930 (4 mines on Orca Group) produced about 97,000 tons (214 million pounds) of copper and subordinate amounts of gold, silver, and zinc; two mines, the Ellener, accounted for more than 96 percent of the production; the few gold mines in the area probably produced total of not more than 31 to 1,000 ounces of gold; resource data are sketchy but the submarine volcanogenic deposits probably represent substantial copper resources; Orca has estimated reserves of at least 1,020,000 tons (1,125,000 st) containing 1.25 percent copper

Production from the gold lodes commenced shortly after 1900, and continued on an intermittent small scale until early 1940's; it consisted of about 1,000 g (45,000 ounces) of gold and minor by-product silver; the gold is localized in high-grade shoots that may contain 340 g/ton (10 oz/st) or more gold; the overall grade of the lodes is probably less than 32 g/ton (1 oz/st); probably small reserves of base and ferrous metals; production from the placers is negligible

Modern reconnaissance mapping accompanied by geophysical studies by U. S. Geological Survey for that part of area within Seaward quadrangle; U. S. Geological Survey sponsored mapping and some sampling for remainder of area; topographic studies of some volcanogenic deposits by Government agencies and industry; recent exploration of some volcanogenic deposits by industry

Sketchy reconnaissance mapping and some old studies of the deposits with recorded production from 20 of them, and two known small gold placers; the Valdez district is partly bounded by glacial-covered mountains

The area contains 43 known lode deposits with recorded production from 20 of them, and two known small gold placers; the Valdez district is partly bounded by glacial-covered mountains

(a) Many small tonnage gold-quartz veins are known; others possible but area is too covered

(b) Several small concentrations of gold in stream placers are known

(a) Over 50 mafic volcanogenic deposits are known; many have been incompletely explored and others probably remain to be found. Estimated number of deposits is only for those suitable to base tonnage model.

(b) Several small tonnage gold-quartz veins are known; others possible.

90% 10% chance that these are 2 4 8 deposits or more

(a) mafic volcanogenic model

(a) One small breccia cemented by zinc and copper sulfides is known.

(a) Many small tonnage gold-quartz veins are known; others possible but area is too covered.

(b) Several small concentrations of gold in stream placers are known.

(a) Many small tonnage gold-quartz veins are known; others possible but area is too covered.

(b) Several small concentrations of gold in stream placers are known.

(a) Many small tonnage gold-quartz veins are known; others possible but area is too covered.

(b) Several small concentrations of gold in stream placers are known.

(a) Many small tonnage gold-quartz veins are known; others possible but area is too covered.

(b) Several small concentrations of gold in stream placers are known.

AREA COVERED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	ESTIMATED NUMBER OF DEPOSITS (SERIAL CHANGE THAT THEY ARE THE NUMBER PRESENT OR MORE DEPOSITS)	SUMMARY OF MINERAL RESOURCE POTENTIAL	GRADES AND TONNAGES FOR THIS REPORT TYPE (IN TABLE 2)
12.	(a) Au(Ag, Sb)--mainly thin quartz veins in Valdez Group	---	Port Wells gold district and nearby area: underlain by metaflysch of the Valdez Group (Cretaceous) and subordinately by Tertiary granitic plutons and felsic dikes; the deposits are mainly in the Valdez Group; they consist of quartz veins, rarely more than 1 m wide, and locally disseminated small lenses, and tabular high-grade shales; breccia fillings; the lodes generally are less than a few hundred meters in strike length; besides gold and quartz they generally contain calcite, pyrite, arsenopyrite, minor uneconomic amounts of base metal sulfides, and a little silver; a few deposits contain stibnite, which might be a by-product of a principal by-product	Production was mainly prior to 1920 and consisted of 657 kg (21,125 ounces) of gold, including 640 kg (20,600 ounces) from the granite mine, and a little by-product silver; most ores in area are highly disseminated, and are dusty	Excellent modern reconnaissance mapping and accompanying geophysical data for that part of area in Seward quadrangle; older U. S. Geological Survey mapping for other parts of area; topographic studies and some mapping of the area by private parties and floods	45 deposits, including 15 mines that, at least, have had minor production, are known in area; 15; the potentially favorable areas are partly delimited by glaciers and floods	Numerous small tonnage gold-quartz veins are known; others possible, particularly under ice	(a) mafic volcanic model	
13.	(a) Cu(Ag, Zn, Au)--submarine volcanic (b) Ni, Cr--Magmatic	Cu--occurrence of weakly disseminated copper and iron sulfides in gabbro	Resurrection Peninsula; Underlain by Valdez Group (Cretaceous), mainly mafic metavolcanic rocks; minor gabbro and serpentinitized dunite (a) mainly as disseminations and breccia cement in sheared Valdez Group mafic volcanic rocks; (b) massive sulfides and thin veins; mainly pyrite with subordinate chalcopyrite, sphalerite, pyrrhotite, and secondary copper minerals (b) minor anomalous amounts of nickel and chromium in serpentinitized dunite	No production or known reserves	Modern reconnaissance geologic, geochemical, and geophysical coverage by U. S. Geological Survey; little industry interest	Area 13 contains 11 scantly explored, essentially inactive prospects, and 2 known occurrences	(a) At least 11 completely explored mafic volcanic copper prospects are known; a few more are possible. The grade-tonnage model may apply to some of these. (b) One small body of serpentinitized dunite containing anomalous values of nickel and chromium is known. A few small tonnage nickel or chromium deposits are possible.	(a) mafic volcanic model	
14.	(a) Au(Ag, Sb)--lodes, typically thin quartz veins (b) Ni(Ag)--placer	Cu--occurrence, vein in sheer zone in Valdez Group	Grovewood, Hope-Gilpatrick, and House Pass mining districts; Underlain by Valdez Group (Cretaceous) metaflysch that locally is cut by Tertiary felsic dikes and granitic plutons (a) gold lodes genetically and spatially related to Tertiary anatectic plutons; veins that are discontinuous, generally less than 1.5 m thick, and less than 2,000 m long; mainly in the flysch; typically with minor amounts of silver and uneconomic scattered base and ferrous metal sulfides; a few deposits have potentials for minor by-or coproduct antimony (b) Stream end bench placers and one beach placer	Total estimated production from the lode about 435 kg (14,000 ounces) of gold, a small amount of silver, and about 90 kg (a few hundred pounds) of antimony; grade data unknown but probably the gold was erratically distributed; antimony mined in small veins; small reserves at a few properties; mostly mined during early 1900's; most placer mines were operated during the early production isn't accurately known, but probably is slightly greater than the lode production	Reconnaissance geologic and geophysical surveys and geochemical sampling by U. S. Geological Survey; local detailed mapping of some mines and prospects; small-scale recent activity at a few placer and lode properties	The area includes moderately accessible parts of the Kensil Mountains, and it has been fairly well explored for gold as attested by numerous mines and prospects	(a) Numerous small tonnage quartz veins are known; a few more deposits might be expected to be found. (b) Many small stream and bench gold placers and one beach placer are known; chances for more are slight.	(a) mafic volcanic model	

HAZARD TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATED TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL (S) OF MINERAL RESOURCES	PRODUCTION AND RESERVE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)
15.	(a) Au(Ag)--mainly thin quartz veins in Valdez Group (b) Au(Ag)--placer	Nuka Bay area; underlain by Valdez Group (Cretaceous) and by local felsic dikes and plutons (Pleistocene). Thin gold-bearing quartz veins that cut Valdez Group metafelsch; local rich shoots. (b) Stream and bench placers	Lode production between 1924 and 1942 about 171 kg (3,500 ounces) of gold and a little bit of silver; average grade a little more than 103 g/t (3 oz/st) some reserves; no known placer production	Local geologic mapping and some sampling near known deposits, but regional geologic, geophysical, and geochronological data are scant; interest by industry	Area delineated mainly on basis of its favorable geologic setting for small gold-bearing shoots; scattered isolated known deposits and local clusters of small deposits; large parts of the area are remote and scantily prospected	90% 50% 10% chance that there are 2 2 3 deposits or more
16. a, b	(a) Au(Ag)--lodes, mainly thin quartz veins in Valdez Group or, less commonly, in area of early felsic plutons (b) Au(Ag, PE)--placer	Area includes most of Chugach and Kenai mountains and contains extensive glacial cover; largely underlain by Valdez Group (Cretaceous) metafelsch, and Tertiary felsic plutons and dikes; (a) Thin gold-bearing quartz veins and a few small lenses mainly localized in Valdez Group; genetically affiliated with Tertiary anatectic plutons. (b) Stream and bench placers; a few small placer sites in the Valdez Group (c) Heavily mineralized vein occurrence in Valdez Group (d) Minor anomalous radioactivity detected at a few sites in the Valdez Group	Estimated total gold production from lodes about 46 kg (1,000 ounces); that from 120 placers (4,000 ounces) silver recovered; the known gold lodes and placers are small but locally rich; they have scant reserves.	Large disparity in geologic data base; most of area mapped by reconnaissance methods, but extensive tracts of the mountainous interior are unmapped; scant local geophysical and geochemical coverage; small-scale recent activity at a few placer and lode gold deposits	Possibly undiscovered deposits of this type exist along the northwest flank of the Kenai Mountains; tectonic settings that are similar to the environs of Red Mountain and Claim Point; however large parts of the inferred favorable areas are covered	Wide scattered gold-ore veins that locally rich grades; remoteness and large amount of glacial cover suggest that most of probably large number of undiscovered; most of those that are found will probably be uneconomic to mine due to their low tonnage (b) Stream gold placers, one of which contains a few bench placers are known; relatively small production and few probably remain to be found (c) One small tonnage copper-silver vein with low grades is known; others possible (d) Minor anomalous radioactivity detected at a few localities; slight chance of large tonnage deposits
17.	Cr--magnetic deposits in layered ultramafic rocks	Disseminated and locally massive chromite in layered dunite and, to a small extent, in pyroxenite and serpentinite; small bodies of the ultramafic masses; Red Mountain, about 6.4 by 3.2 km in outcrop plan, and a smaller near tide-water mass at Claim Point; on basis of recent studies both ultramafic bodies are interpreted as klippen that have been thrust over the Michigh complex (Cretaceous)	Production: 1917-18, about 2,000 tons containing 45 percent Cr ₂ O ₃ ; 1922-24, 6,500 tons; 1925-27, 42 percent Cr ₂ O ₃ ; 1954-57, about 21,000 tons. grade not known but probably about 40 percent Cr ₂ O ₃ ; Cr:Fe ratio 2:3; 1942 estimated reserves of about 30,000 tons of chrome; the total 72,000 tons derived from concentrating lower-grade material	The deposits have been studied in some detail and the surrounding areas mapped in reconnaissance; scant geophysical and geochronological data; tectonic settings that are similar to the environs of Red Mountain and Claim Point; however large parts of the inferred favorable areas are covered	One of two ultramafic masses in this area, Red Mountain reportedly contains a total of about 50,000 tons of shipping ore at 41 percent Cr ₂ O ₃ . The other mass, Claim Point has about 260,000 tons at 17.8 percent Cr ₂ O ₃ that could be concentrated to about 75,000 tons at 45 percent; additional deposits under younger rocks and water are likely at Claim Point; ultramafic deposits possibly exist under covered areas along the northwest flank of the Kenai and Chugach Mountains nickel anomalies are known in both areas; grades and tonnages of podiform deposits are appropriate for unroofed deposits in alpine masses	podiform chromite model
18.	(a) Au(Ag)--placer	Placer gold deposits on beaches fronting lower Cook Inlet and possibly in nearby alluviated valleys; typically small, in part ephemeral deposits; only a few deposits known	Worked intermittently during early 1900's; production not known, probably about 30 kg (1,000 ounces) of gold and a little silver	Tertiary non-marine sedimentary rocks that underlie the region and large parts of the nearby Kenai Lowland are regarded as favorable hosts for placer deposits; stream exploration, no uranium exploration; no uranium deposits are known in the region	(a) A few small, in part ephemeral, gold placers are known; a few others possible (b) Tertiary rocks that underlie this area and large parts of the nearby Kenai Lowland are favorable for uranium; however none has been found despite some exploration	

19. (a) Some small tonnage gold-quartz deposits in the McHugh Complex; a few others possible.

(b) Small tonnage placer gold in beaches, streams, and benches are known; no known reserves.

(c) One possible mafic volcanogenic deposit and a few minor occurrences in the McHugh Complex; others possible.

(d) No known deposits but area is favorable for chromite similar to deposits in area 17 or in small alpine-type ultramafic rocks.

(d) podiform chromite model

20. (a) Cu--migmatic deposits in ultramafic rocks.

(b) Cu,Zn,(Ag)--submarine volcanogenic deposits associated with mafic lavas.

(c) Au(Ag)--iodides, typically thin quartz veins.

(d) Au(Ag)--placer

(e) Ni--local minor anomalous amounts of nickel in ophiolite.

(f) U--one prospect allegedly for uranium, in McHugh Complex.

(a) Two chromite prospects and four occurrences are known in this area; the large area that is favorable for chromite deposits combined with the scanty exploration suggests that a large number of deposits might exist here.

(b) Small low-grade mafic volcanogenic deposits containing copper, zinc, and some silver are known; favorable terrain and lack of systematic exploration, especially in the eastern part of the area, suggest that some possibly larger deposits could exist.

(c) Two thin gold-bearing quartz veins are known; others possible.

(d) Gold-bearing stream placers are known in the western part of area 20; a few similar placers might exist in the eastern part.

21. (a) Two chromite prospects and four occurrences are known in this area; the large area that is favorable for chromite deposits combined with the scanty exploration suggests that a large number of deposits might exist here.

(b) Small low-grade mafic volcanogenic deposits containing copper, zinc, and some silver are known; favorable terrain and lack of systematic exploration, especially in the eastern part of the area, suggest that some possibly larger deposits could exist.

(c) Two thin gold-bearing quartz veins are known; others possible.

(d) Gold-bearing stream placers are known in the western part of area 20; a few similar placers might exist in the eastern part.

AREA OUTLINED ON MAP

MAJOR TYPES OF KNOWN DEPOSITS

SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)

GEOLOGIC CONTROL(S) OF MINERAL RESOURCES

PRODUCTION AND RE-SOURCE INFORMATION

STATUS OF GEOLOGIC INFORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL RESOURCE POTENTIAL

ESTIMATED NUMBER OF DEPOSITS PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)

GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)

ESTIMATED NUMBER OF DEPOSITS PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
90% 50% 10% chance that there are 4 deposits or more	(a) nodiform chromite model
90% 50% 10% chance that there are 1 3 deposits or more	(b) mafic volcanogenic model
90% 50% 10% chance that there are 1 2 5 deposits or more	(c) porphyry copper model
90% 50% 10% chance that there are 1 2 4 deposits or more	(f) nickel sulfide model

Area 21 contains a variety of potential deposits. The most favorable terrain has been scantly prospected; all known deposits are unexplored or scantly explored. This area is known for its ultramafic rocks, which are locally abundant. The area is also known for its mafic-ultramafic rocks, which are locally abundant. The area is also known for its porphyry copper and submarine volcanic deposits.

Some disparity in coverage that part of the area is unexplored or scantly explored. This area is known for its ultramafic rocks, which are locally abundant. The area is also known for its mafic-ultramafic rocks, which are locally abundant. The area is also known for its porphyry copper and submarine volcanic deposits.

Recent gold production, probably on a small scale, is reported from the area. This production is primarily from the mafic-ultramafic rocks, which are locally abundant. The area is also known for its porphyry copper and submarine volcanic deposits.

(1) (Mn)--hydrothermal
 (a) Cr(Ni, Cu, Pt)--magmatic deposits
 (b) Cu(Ag, Zn)--submarine volcanic
 (c) Cu(Mo, Ag, Au)--porphyry
 (d) Zn(Pb, Ag)--replacement deposits
 (e) Au(Ag)--lode deposits, typically thin quartz
 (f) Ni, Cu(Pt)--magmatic
 (g) Au(Ag)--placer
 (h) Cu(Ag)--lode deposits, generally thin veins

Area includes northern flank of upper Paleozoic Chitina Valley; largely underlain by upper Paleozoic metamorphosed sedimentary and volcanic rocks and less extensively by ultramafic-mafic complexes and Jurassic granitic rocks. (a) Thin layers, lenses, and disseminations of chromite in partly sericitized granite that includes ultramafic complexes that are as much as 16 km long and several kms wide but commonly much smaller; local minor anomalous amounts of nickel, copper, and platinum-group elements from parts of the complexes. (b) Mainly disseminated sulfides including chalcopyrite and sphalerite in metamorphosed upper Paleozoic mafic volcanic rocks. (c) Altered zones in Jurassic granitic rocks that contain local sulfide-bearing disseminations and veinlets. (d) Massive and disseminated sulfides, mainly sphalerite and galena, in Permian marble. (e) Thin gold-bearing quartz veins in upper Paleozoic metamorphic rocks, generally near granitic plutons. (f) Small deposits of massive and disseminated sulfides, including pentlandite, bravoite, and chalcopyrite in peridotite dikes or gabbro. (g) Stream placers. (h) Small copper-bearing veins in upper Paleozoic metamorphic rocks; some may be of submarine volcanogenic derivation. (i) Disseminated or massive quartz veinlets in lenticular zones within upper Paleozoic metamorphic rocks.

(a) Scantly explored chromite deposits are known; lack of detailed geologic information and paleogeology suggest that many more deposits exist.

(b) Some disseminated mafic volcanogenic deposits containing copper, silver, and zinc are known; large amount of unexplored favorable terrain; probably undiscovered deposits in area 21.

(c) A few undrilled sulfide bearing altered zones in granitic rocks are known; other deposits probably exist.

(d) Massive and disseminated sulfides containing lead, zinc, and silver are known in Permian marble; other deposits possible.

(e) Small tonnage gold-bearing quartz veins with some production; others possible.

(f) Low tonnage massive and disseminated deposits containing nickel and copper with minor concentrations of platinum-group elements in the ultramafic complexes possible.

(g) Gold placers are known; some production; other small placers possible.

(h) Small tonnage copper veins are known; others possible.

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSIT TYPES (INCLUDES MINOR OCCURRENCES)	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSIT TYPES (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RESOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE IS AT LEAST ONE PRE-SENT OR MORE DEPOSIT)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
22.	(a) Cu, Mo (Ag) -- porphyry type deposits (b) Mo -- veins and stockworks (c) Pb, Zn (Ag, Cu) -- replacement deposits	(d) Ni -- disseminated deposits (e) Fe -- skarn (f) Cu (Ag) -- vein deposits (g) U, Th -- some favorable host rock for U-Th deposits	Extreme southeastern parts of Wrangell Mountains; underlain by metamorphosed mid-Paleozoic, mainly carbonate rocks; weakly metamorphosed upper Paleozoic sedimentary and volcanic rocks; an upper Paleozoic syenite-monzonitic plutonic complex, gabbro, and secondary granitic plutons (a) Associated with altered zones in Tertiary granodiorite; two known deposits; one mainly for copper and the other mainly for molybdenum (b) Molybdenite-bearing quartz veins and small stockworks in Tertiary granitic plutons (c) Small sulfide-bearing pods and disseminations in Permian marble (d) Sparsely disseminated gold in upper Paleozoic volcaniclastic rocks; mineralized zone low in grade and probably local in extent (e) Small magnetite- and hematite-bearing concretions (skarn) deposited in contact to upper Paleozoic monzonite; contains minor amounts of copper (f) Small copper-bearing veins in fault zones (g) Some phases of the syenite-monzonite complex are geologically favorable for uranium-thorium deposits	No production; the main inferred resource of the area are in the copper and molybdenum deposits that are associated with Tertiary plutons	Reconnaissance geologic mapping and supplemental reconnaissance geochemical and geophysical investigations; very little exploration	The area is remote and rugged and, at best, has been cursorily prospected; its potential most significant deposits are copper and molybdenum porphyries associated with Tertiary plutons; these deposits, like others in the area, are geologically favorable for Tertiary plutons and their environments are regarded as highly favorable for exploration; other parts of the area, including extensive tracts underlain by mid-Paleozoic rocks that lack known deposits, have diverse degrees of geologic favorability	(a) One porphyry copper and one porphyry molybdenum deposit are known but undrilled; other deposits likely (b) Molybdenite-bearing quartz veins and small stockworks exist; these deposits may be undrilled or underexplored, such as the porphyries (c) Small tonnage lead-zinc replacement deposits have been found in area 22; other small deposits of this type probably occur here (d) A low grade disseminated gold deposit in volcaniclastic rocks has been found, probably local in extent	90% 50% 10% chance that there are 4 or more deposits 2 3	(a) porphyry copper model and porphyry molybdenum model

ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANGE THAT THERE ARE THE NUMBER PRE-SERIED OR MORE DEPOSITS)

SUMMARY OF MINERAL RESOURCES POTENTIAL

ADDITIONAL COMMENTS

STATUS OF GEOLOGIC INFORMATION

PRODUCTION AND RE-SOURCE INFORMATION

GEOLOGIC CONTROL (S) OF MINERAL RESOURCES

SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)

MAJOR TYPES OF MINERAL DEPOSITS

AREA OUTLINED ON MAP

23.	(a) Cu(Ag)--Kennecott type (b) Cu(Ag)--vein (c) Au(Ag, Sb, Mo)--vein (d) Au(Ag, Cu)--placer (e) Au(Mo)--placer (f) Cu(Ag)--skarn (g) Cu(Ag)--sub-aerial volcano-geologic (h) Ag(Cu, Zn, Pb)--vein (i) Sb(Au, W)--vein	(j) Zn(Ag, Pb)--replacement or vein (k) Mo--vein	South-central flank of Wrangell Mountains, a well mineralized area that contains diverse deposits; underlain by upper Paleozoic and abundant Mesozoic sedimentary and volcanic rocks, local Jurassic and Tertiary plutons represent subvolcanic basaltic rocks--and local Cenozoic lavas with minor sedimentary facies (a) Mainly massive copper sulfide-rich lodes localized in lower, chiefly dolomitic, parts of Upper Triassic Chitstone Limestone (b) Typically quartz-calcite replacement veins, which thin out and eventually are confined to the Greenstone; chief ore minerals, chalcopryrite, bornite, and chalcocite (c) Thin gold-bearing quartz veins genetically related to Tertiary plutons or, rarely, to Jurassic plutons (d) Stream and bench placers contain some reserves of porphyry-type deposits associated with Jurassic granitic plutons (e) Small magnetite-rich contact-metamorphic (skarn) deposits in Triassic carbonate rocks adjacent to Jurassic granitic plutons (f) Native copper-bearing mainly anhydrotite deposits in Triassic basaltic and siliceous rocks (142,500 ounces) of gold and some byproduct silver; since 1959 they produced about 60 kg (a few thousand ounces) of gold and small amounts of native copper nuggets that are utilized in Alaska (g) Native copper-bearing veins are active, but their reserves and resources are probably small; except for the sub-aerial volcanogenic deposits (g), which have yielded a little copper and silver, none of the other deposits have been mined; most of the remaining reserves have not been adequately explored, and they all can be regarded as having some resource potential	Production dominated by Kennecott mines, Alaska's premier producer of copper and silver; during their major operations, between 1913 and 1938, they produced 540,000 tons (1.2 billion pounds) of copper and 280 tons (9 million ounces) of silver; minor post-1938 production from small-scale, largely surficial, operations; production data for other deposits less accurately known; probably about 2,700 tons (5 million pounds) of copper and 6,220 kg (200,000 ounces) of silver; Kennecott type deposits contain some reserves of porphyry-type deposits associated with Jurassic granitic plutons (f) Small magnetite-rich contact-metamorphic (skarn) deposits in Triassic carbonate rocks adjacent to Jurassic granitic plutons (g) Native copper-bearing veins are active, but their reserves and resources are probably small; except for the sub-aerial volcanogenic deposits (g), which have yielded a little copper and silver, none of the other deposits have been mined; most of the remaining reserves have not been adequately explored, and they all can be regarded as having some resource potential	Geologic mapping, ranging from detailed to reconnaissance, and reconnaissance geophysical coverage for engineering studies; local topographic maps, mainly related to the mineral deposits; moderate localized current exploration interest	STATUS OF GEOLOGIC INFORMATION	PRODUCTION AND RE-SOURCE INFORMATION	GEOLOGIC CONTROL (S) OF MINERAL RESOURCES	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	MAJOR TYPES OF MINERAL DEPOSITS	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANGE THAT THERE ARE THE NUMBER PRE-SERIED OR MORE DEPOSITS)
			<p>The area has been well prospected by old, traditional prospecting methods, but only scantily explored by modern, sophisticated techniques; because of its diverse deposit types, the area is regarded as having a strong potential for significant mineral resources; an approximate summary of the number of known deposits in the area follows:</p> <p>Kennecott type, 11 mines and 3 prospects; gold-bearing veins, chiefly in Mineral Greenstone, 3 mines, 33 prospects, 9 occurrences; gold placers, 4 mines and 4 prospects; mainly native copper in basalt, 1 mine, 7 prospects; all other deposit types, 1 mine, 22 prospects, 9 occurrences; of the mines and prospects have been active during recent years; the occurrences were discovered during recent U. S. Geological Survey investigations, and some of them are worthy of exploration</p>							<p>(a) Massive copper sulfide deposits containing silver were the largest producers of copper and silver; reserves about 11 percent copper and 66 g/t silver; all known deposits are exposed, at least in part on surface, and about 130 km² of favorable geologic terrain is covered; approximately 7 of the large tonnage-high grade and numerous smaller deposits are estimated to be unmined</p> <p>(b) Generally small tonnage quartz-calcite veins containing copper and silver; other deposits possible</p> <p>(c) Small tonnage gold-bearing quartz veins are known; others possible</p> <p>(d) Gold-bearing stream and bench placers that contain some silver and copper have been mined; a few are still active; reserves and resources are probably small</p> <p>(e) Three undrilled but apparently weakly mineralized porphyry copper deposits are known; others possible</p> <p>(f) Small tonnage skarn deposits containing iron and copper have been found; unmined deposits probably remain</p> <p>(g) Low grade native copper-bearing deposits are known; mainly amygdaloidal deposits in Triassic basalt; other deposits likely</p> <p>(h) Silver-bearing veins that have small tonnages are known</p> <p>(i) Small high-grade antimony (stibnite)-bearing veins</p>	<p>90% 50% 10% that there are 2 3 5 or more deposits</p> <p>(e) porphyry copper</p>

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RESERVE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	ESTIMATED NUMBER OF DEPOSITS OF PRESENT TYPE (MINIMUM OF ONE OR MORE DEPOSITS)	GRADES AND THICKNESSES FOR THIS DEPOSIT TYPE (IN TABLE 2)
24.	(a) Cu(Ag)--porphyry (b) Au--vein	(a) Cu(Ag)--subaerial volcanic (b) Cu(Ag)--vein (c) Cu--magmatic	Western Wrangell Mountains; small windows that expose upper Paleozoic metamorphic rocks, Mesozoic granitic plutons, and Cretaceous sedimentary rocks; surrounded by Cenozoic Wrangell Lava; (a) The granitic rocks and their environs are favorable for porphyry-type deposits. (b) One granitic vein that cuts the metamorphic rocks	No data	Sketchily mapped; no known geochemical or geophysical investigations; no known recent prospecting	Area outlined on basis of favorable geology	90% 50% 10% chance that there are 0 1 2 deposits or more	(a) porphyry copper model
25.	(a) Cu(Ag)--subaerial volcanic (b) Cu(Ag)--vein (c) Cu--magmatic	(d) Cu(Mo)--porphyry (e) Cu--placer	Northeastern flank of Wrangell Mountains; underlain by upper Paleozoic and Mesozoic volcanic and sedimentary rocks, Cretaceous(?) and Tertiary plutons, and Cenozoic Wrangell Lava (a) Native copper in Tertiary basaltic and boninite or chalcocite and their oxidation products in veins less than 1 m thick or in swarms of veinlets or surface coatings; generally in Triassic basalt (c) Occurrence of disseminated sulfides, including chalcopyrite, in a thick mafic dike (d) Altered zones suggestive of porphyry-type mineralization in granitic rocks (e) Native copper nuggets in stream and bench placers	No production or reserve data but possibly significant copper resources	Reconnaissance geologic, geochemical, and geophysical coverage; scant exploration interest	Area 25 has been only scantily prospected; 7 of its 14 known deposits are occurrences that were discovered during recent U.S. Geological Survey investigations. Significant resource significance of the area's diverse copper deposits cannot be accurately determined without adequate exploration; apparently large but very low grade copper resources in Triassic basalts (Mikolai Greenstone) in this and other areas, notably (23), may constitute resource of the future	90% 50% 10% chance that there are 0 1 2 deposits or more	(d) porphyry copper model
26.	(a) Cu--porphyry (b) Au(Ag, Pt)--placer (c) Cu(Au)--iodine deposits, mainly veins (d) Au(Mg)--vein	---	Upper Metanuska Valley and nearby terrain; in part bounded by major faults; underlain by Mesozoic sedimentary rocks, and Tertiary sedimentary rocks, and Mesozoic and Tertiary intrusive rocks (a) The few known porphyry type deposits are associated with strongly altered zones in Tertiary felsic plutons and nearby rocks (b) Numerous small placer gold deposits along streams and alluvial benches (c) Poorly known; probably mostly veins related to intrusive rocks but may include submarine volcanogenic deposits (d) Thin veins probably genetically related to Mesozoic and Tertiary plutonism	Small, but in-accurately known production, probably about 30 kg of gold from the placers; no lode production; inadequate exploration for valid resource estimate, but resources probably are small to moderate	Reconnaissance and local semidetalled geologic mapping; scant geophysical and geologic coverage; recent interest in placer and lode deposits	The area contains about 39 known placer deposits and extensive deposits north-east of area 26 that are underlain by Cretaceous sedimentary rocks or surficial deposits of the Copper River Basin contain scattered gold placers; parts of the Copper River Basin are geologically permissive for sedimentary placer deposits, but this area is not outlined as favorable because prospecting results have been negative; area 26 contains zeolite deposits, which although a non-metallic commodity, are of possible economic importance	90% 50% 10% chance that there are 0 1 3 deposits or more	(a) porphyry copper model

AREA OUTLINED ON MAP
 MAJOR TYPES OF KNOWN DEPOSITS
 SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)
 GEOLOGIC CONTROL(S) OF MINERAL RESOURCES
 PRODUCTION AND RE-SOURCE INFORMATION
 STATUS OF GEOLOGIC INVESTIGATION
 ADDITIONAL COMMENTS
 SUMMARY OF MINERAL RESOURCE POTENTIAL
 ESTIMATED NUMBER OF DEPOSITS PRESENT OR CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS
 GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)

27.	(a) Cu(Ag)--Iodes, mainly veins (b) Cu(Mo)--porphyry (c) Au--placer	Southern Talcotna Monocline, underlain by granitic mass and by small areas of upper Paleozoic metamorphic rocks (a) Poorly known copper deposits generally represented by thin veins, fracture coatings and local disseminations, typically in amphibolites or granitic rocks; may include some porphyry type and magmatic deposits (b) Geologically favorable for porphyry type deposits, but none definitely known (c) One known stream placer	No data	Reconnaissance geologic mapping and local geochemical and geophysical coverage, little recent interest by industry	Area delineated mainly on basis of its geologic favorability for porphyry copper deposits and lack of systematic modern prospecting	(a) Generally thin copper-bearing veins; some known occurrences may be related to porphyry type and magmatic deposits; lack of systematic prospecting (b) Area favorable for porphyry copper deposits; none known	90% 50% 10% chance that there are 0 0 2 deposits or more	(b) porphyry copper model
-----	---	---	---------	--	---	--	--	---------------------------

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATED TYPES OF MINERAL RESOURCES WITHIN OCCURRENCES	GEOLOGIC CONTROLS OF MINERAL RESOURCES	PRODUCTION AND RECOVERY INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (SEE TABLE 2)
28.	(a) Au(Ag,Te)--veins (b) Au(Ag)--placer	(c) Cu(Mo,Ag)--porphyry	Willow Creek district and southwest part of Talkeetna Mountains; largely underlain by an Upper Cretaceous-tertiary tonalitic batholith; small tracts of upper Paleozoic metamorphic rocks to the batholith included in area. (a) Mainly quartz veins may extend for a few kilometers along strike; contain gold, minor amounts of sulfides, and rarely, tellurides and scheelite (b) Chiefly stream placers (c) One poorly known deposit that may be a porphyry type is reported in the tonalite; others probable	Total production from the Willow Creek gold lodes about 17,800 kg (574,000 ounces) of gold, small amount of silver. Recoverable amount probably a little more than 15 million tellurium; main mining between 1932 and 1942; placer gold production probably 60 kg (several thousand ounces), but not accurately known; lode gold was produced from 15 mines, a few of which contain scheelite. Grade probably about 70 g/t (a few ounces per ton) with local higher-grade shoots	The Willow Creek district has been mapped and studied in some detail; the rest of the area in reconnaissance; reconnaissance geochemical and geophysical mapping is in progress; although no recent activity at a few of the areas; small-scale deposits.	For some unknown reason the productive known gold lodes are concentrated in or near southern batholith; although no gold lodes are known in them, similar gold lodes may occur in other parts of the batholith; the Willow Creek district has been well prospected, and its inferred lode gold resources are largely in deeper parts of known deposits; in inferred deposits, undiscovered deposits in the northern and central parts of the batholith have some potential for porphyry copper deposits as well as gold lodes; one small soapstone deposit in the area is mined intermittently	(a) Numerous gold-bearing quartz veins in southern part of batholith; tonnage small; contain small reserves; majority of resource probably in deeper parts of known deposits or in concealed, undiscovered deposits that may be near the known deposits or in other parts of area 28 that appear geologically similar (b) Small gold-bearing stream placers are known; some others possible (c) One deposit may be a porphyry copper-type; geology is favorable for more	90% 50% 10% chance that there are 0 1 3 deposits or more	(c) porphyry copper model
29.	(a) Fe--contact metamorphic	(b) Au--placer	Area includes marginal facies of Jurassic granitic batholith; nearby Lower Cretaceous sandstone and volcanic rocks (a) Apparently small contact-metamorphic (stern) type deposits that are rich in magnetite (b) Stream placers	No data	Local brief study of known contact metamorphic deposits and reconnaissance geologic mapping; scant known geochemical and geophysical investigations; little recent interest	The known deposits appear to be too small to constitute a significant iron resource, but the area has been only cursorily prospected and may contain larger deposits	(a) Several iron-rich (magnetite) stern deposits; tonnage appears small but other larger deposits may exist; this largely unprospected area		

AREA
DATE
BY

MAJOR TYPES OF KNOWN
DEPOSITS

SUSPECTED OR SPECULA-
TIVE TYPES OF MINERAL
DEPOSITS (INCLUDES
RHYTHM OCCURRENCES)

GEOLOGIC CONTROL(S) OF
MINERAL RESOURCES

PRODUCTION AND RE-
SOURCE INFORMATION

STATUS OF GEOLOGIC IN-
FORMATION

ADDITIONAL COMMENTS

ESTIMATED NUMBER OF
DEPOSITS PERCENT
CHANCE THAT THERE
ARE THE NUMBER PRE-
SENTED OR MORE
DEPOSITS

GRADES AND TONNAGES
FOR THIS DEPOSIT
TYPE (IN TABLE 2)

31. (a) Cu(Mo, Ag, Au)--
porphyry
(b) Au(Ag)--placer
(c) Cu(Ag)--vein
(d) Cu(Ag)--sub-
aerial volcano-
genic
(e) Au(Ag, Cu)--
vein

Extreme northeastern
part of McCarthy quad-
rangle and southeastern
part of Habesna quad-
rangle; contains local
upper Paleozoic and Tri-
assic sedimentary and vol-
canic rocks, abun-
dant upper Mesozoic
flysch, and abundant
Miocene volcanic rocks,
Cretaceous granitic
plutons. Tertiary fel-
sic hypabyssal plutons,
and Cenozoic andesitic
lavas

(f) Au(Cu)--contact
metamorphic?
(g) Cu--magmatic
one occurrence
(h) Ag(Au, Cu, Zn, Pb)--
vein

Placer gold produc-
tion estimated between
1,400 kg and 1,500 kg
(45,000 and 50,000
ounces) of gold and
minor byproduct sil-
ver constitutes the
only production from
the area; the area's
six porphyry copper
resources are estimated
to contain 280 million
tons of 0.2 percent cop-
per and very low molyb-
denum and gold con-
tents; the area may
contain some placer
gold resources of
interest, but re-
sources in other de-
posit types, except
porphyries, are in-
ferred to be minor

The resource po-
tential of the
porphyry copper
deposits over-
laps that of the
other deposit types;
a strong aeromagnetic
anomaly near the
southeastern extreme
of the area is
probably indicative
of a concealed plu-
ton of the Klein Creek
type,
which may be an at-
tractive exploration
target for porphyry-
type deposits

Covered by modern recon-
naissance geologic, geo-
chemical, and geophysical
investigations; recent ex-
ploration interest in the
porphyry deposits, a few
gold placers, and one
gold lode

(a) Nine porphyry cop-
per type deposits have
been found but are in-
completely explored;
six deposits have been
partially drilled; sty-
les are possible in
this area; between areas
31 and 32 lies a covered
region favorable for por-
phyry copper deposits

(b) A number of placer
gold deposits are known;
past production about
1500 kg gold; some gold
probably remains in
known placers; a few
placers might be left
to be found

(c) Small tonnage sul-
fide-bearing veins occur;
others possible

(d) High tonnage of very
low grade copper in ba-
salt; smaller tonnages
with locally high grades
possible

(e) Small tonnage gold-
bearing quartz veins

AREA OUTLINED ON MAP

MAJOR TYPES OF KNOWN DEPOSITS

SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)

GEOLOGIC CONTROL(S) OF MINERAL RESOURCES

PRODUCTION AND RE-SOURCE INFORMATION

STATUS OF GEOLOGIC INFORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL RESOURCE POTENTIAL

ESTIMATED NUMBER OF MINERAL RESOURCES THAT HAVE BEEN DISCOVERED OR ARE THE NUMBER OF DEPOSITS STARTED OR MORE

GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)

90%	50%	10%	chance that there are 9 or more deposits probably porphyry copper; other concealed deposits probably remain to be found	(a) porphyry copper model
4	6	9		
90%	50%	10% <td>chance that there are 4 or more deposits in this area</td> <td>(b) porphyry molybdenum model</td>	chance that there are 4 or more deposits in this area	(b) porphyry molybdenum model
2	3	4		
90%	50%	10% <td>chance that there are 14 deposits or more</td> <td>(c) stannic copper model</td>	chance that there are 14 deposits or more	(c) stannic copper model
8	10	14		

The area is highly and diversely mineralized; it is believed to contain significant resources particularly in porphyry-type deposits; despite a moderate amount of prospecting, the area probably contains some undiscovered, concealed deposits that may be important

Covered by modern reconnaissance geologic, geochemical and geophysical studies by U.S. Geological Survey; local topographic studies, mainly of porphyry type deposits, by government and other geologists; several active porphyry type deposits

The only production from the area consists of a little less than 1,800 kg (57,000 ounces) of gold almost entirely from one contact metamorphic deposit, the Nabesna area. The area is rich in porphyry type deposits, and gold resources are inferred in the porphyry-type deposits; the two largest known and best explored porphyry copper deposits have indicated and inferred resources of about 800 metric tons that average between 0.30 and 0.35 percent copper, 0.02 percent molybdenum, and about .017 g/t and very low amounts of silver; the other porphyry copper and porphyry molybdenum deposits are much smaller; one contact metamorphic deposit contains 4,000 tons that averages 34 g/t (1 oz s/t) gold, and similar resources are inferred in nearby deposits; the other deposit types are inferred to have small resources, but, in general, they haven't been adequately explored

South-central and west-central parts of Nabesna quadrangle; underlain by upper Paleozoic and Mesozoic volcanic and sedimentary rocks, Mesozoic and Tertiary plutons, and local Cenozoic volcanic rocks. (a) Porphyry-type deposits associated with Mesozoic granitic plutons. (b) Porphyry-type deposits generally associated with Tertiary plutons. (c) Contact-metamorphic deposits adjacent to Mesozoic granitic plutons; chiefly for gold or copper. (d) Stockworks of quartz veins in or near Tertiary plutons. (e) Breccia pipes associated with dynamic intrusive activity. (f) Amygdaloidal and weakly disseminated copper deposits in Triassic basalt. (g) Thin copper-bearing veins in various host rocks, mainly Triassic basalt. (h) Small placer gold deposits in streams. (i) Thin gold-bearing veins in diverse geologic settings. (j, k) Minor occurrences with little economic potential.

(j) Cu--magnetite?
(k) Mo--pepmatite
(l) (Ag, Pb, Zn)--vein
(m) (Ag, Pb, Zn)

(a) Cu--Ag, Au--porphyry
(b) Mo--porphyry
(c) Au, Cu(Ag, Fe)--contact metamorphic
(d) Cu(Ag, Au)--stockwork
(e) Cu(Ag, Pb, Zn)--breccia pipe
(f) Cu(Ag)--subaerial volcanic
(g) Cu(Ag)--vein
(h) Au(Ag)--placer
(i) Au(Ag)

32.

33.	<p>(e) Au(Ag, Pb)-- (b) Ag, Au(Cu, Pb)-- vein (c) Cu(Ag)--sub- marine volcano- genic (d) Cu--contact metamorphic (e) Mo, Cu(Ag)-- porphyry and stockwork</p>	<p>Southern flank of east- west-trending range and transsected by Denali fault but mainly south of the fault; under- lain by Paleozoic meta- morphitic rocks north of Denali fault; elsewhere by upper Paleozoic vol- canic, sedimentary, and plutonic rocks and by Mesozoic volcanic and plutonic rocks</p>	<p>The gold placers account for the major production from the area their production is not accurately known but probably on the order of 1,900 kg (60,000 ounces) of gold with a little by- product silver and platinum; the placers are be- lieved to contain silver, lead, and sources for con- tinued small- scale mining; the silver-rich veins associated with the Ahtell pluton appear to be too small to constitute more than a modest re- source; the large- ly untested submarine vol- canogenic porphy- ry deposits may contain significant other resources; types known in the area are re- garded as having minor resource potentials</p>	<p>The geology of the area is mainly new from lo- cal studies; some state geological geochronological but no geophysical studies, accompanied these investigations; parts of the area are well prospected and others scantily pros- pected; recent activ- ity has centered on several placer oper- ations, exploring a new silver-bearing vein and searching for submarine volcano- genic or porphyry-type deposits</p>	<p>The area is well mineralized; it contains a variety of deposit types and possibly significant resources</p>	<p>(a) Gold-bearing stream, barite, and other min- erals have had past pro- duction on the order of 1900 kg gold with some silver and platinum; continued small pro- duction possible (b) Small tonnage quartz or barite-carbonate veins containing silver, gold, and some copper and lead (c) Copper-bearing dis- seminated sulfides that may represent volcano- genic deposits (d) Three contact meta- morphitic deposits con- taining copper known; others possible (e) One weakly miner- alized porphyry molyb- denum deposit whose value is unknown but may be porphyry cop- per are known; area is favorable for porphyry copper or molybdenum deposits and is only partially explored (h) Occurrence of spar- sely disseminated gold in chlorite; may be related to porphyry copper mi- neralization</p>	<p>90% 50% 10% chance that there are 3 5 9 deposits or more 90% 50% 10% chance that there are 0 1 3 deposits or more</p>	<p>(e) porphyry copper model porphyry molybdenum model</p>
-----	--	--	---	---	--	---	--	--

34. AREA COVERED BY MAP

MAJOR TYPES OF KNOWN DEPOSITS

SUSPECTED OR SPECULATED TYPES OF MINERAL DEPOSITS (MINOR OCCURRENCES)

GEOLOGIC CONTROLS OF MINERAL RESOURCES

PRODUCTION AND RESERVE DATA

STATUS OF GEOLOGIC INFORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL RESOURCE POTENTIAL

ESTIMATED NUMBER OF DEPOSITS BY TYPE AND GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)

CHANCE THAT THERE ARE THE NUMBER PERCENT LISTED OR MORE DEPOSITS

<p>(a) Au(Ag)--placer (b) Cu(Mo, Au, Ag)--porphyry (c) Cu(Ag)--submarine volcanic (d) Cu(Fe, Au)--concentrated (e) Ni, Cu(Ag)--porphyry (f) Cu(Ag)--subaerial volcanic (g) Au(Ag)--vein (h) Cu(Ag)--vein (i) Mo(Cu)--porphyry (j) Cu(Ag)--sedimentary?</p>	<p>(k) Au(Cu)--fossiliferous placer? (l) Zn(Cu)--metamorphic replacement (m) Cr--magmatic (n) U--type unknown</p>	<p>Southern flank of east central Alaska Range south of McKinley strand of Denali fault and proximal area to south; underlain by upper Paleozoic and Mesozoic volcanic and sedimentary rocks. Tertiary granitic plutons, and local ultramafic and mafic masses.</p> <p>(a) Stream, bench, and channel placers. (b) Porphyry-type deposits in or near granitic plutons. (c) Disseminated and locally massive sulfides, including chalcopyrite, bornite, and associated with upper Paleozoic volcanic rocks. (d) Skarn-type deposits generally with lean disseminations of chalcopyrite and local concentrations of magnetite. (e) Disseminated and locally massive sulfides in ultramafic dikes or gabbro; contain some chalcopyrite and pentlandite. (f) Low-grade mineralized zones in Triassic mafic lavas. (g) Thin gold-bearing quartz veins in diverse host rocks. (h) Thin veins commonly localized in shear zones; most abundant in Triassic volcanic rocks. (i) Local, apparently low-grade zones of magnetite-bearing quartz veins in Cretaceous? granite. (j) Finely laminated pyrite and chalcopyrite in Triassic sedimentary rocks that interfinger with Triassic basalt. (k) Heavily mineralized gold-bearing upper Paleozoic conglomerate. (l) Occurrence of disseminated and copperiferous rocks in the contact zone. (m) Sparsely disseminated chromite in small masses of serpentized dunitite. (n) Reported claims; geologic setting not known.</p>	<p>The Valdez Creek district has produced about 1,700 kg (54,000 ounces) of placer gold and some byproduct silver; its placer gold resources have been estimated at more than 15,000 kg (405,000 ounces) in buried channels and bench gravels; the other gold placers in the area have yielded minor production; they probably have small resources; no known gold production; potential is high in several types of volcanic rocks.</p> <p>(d) Skarn-type deposits generally with lean disseminations of chalcopyrite and local concentrations of magnetite. (e) Disseminated and locally massive sulfides in ultramafic dikes or gabbro; contain some chalcopyrite and pentlandite. (f) Low-grade mineralized zones in Triassic mafic lavas. (g) Thin gold-bearing quartz veins in diverse host rocks. (h) Thin veins commonly localized in shear zones; most abundant in Triassic volcanic rocks. (i) Local, apparently low-grade zones of magnetite-bearing quartz veins in Cretaceous? granite. (j) Finely laminated pyrite and chalcopyrite in Triassic sedimentary rocks that interfinger with Triassic basalt. (k) Heavily mineralized gold-bearing upper Paleozoic conglomerate. (l) Occurrence of disseminated and copperiferous rocks in the contact zone. (m) Sparsely disseminated chromite in small masses of serpentized dunitite. (n) Reported claims; geologic setting not known.</p>	<p>The area is regarded as being well mineralized and as having a good resource potential; besides its placer gold resources, it has scantily explored deposits of several types that represent potentially significant resources, and it is favorable for additional discoveries.</p>	<p>90% 50% 10% chance that there are 4 6 10 deposits or more</p> <p>90% 50% 10% chance that there are 2 4 9 deposits or more</p>	<p>(b) porphyry copper model (c) felsic and intermediate volcanogenic massive sulfide model (d) copper skarn model (e) nickel sulfide model (f) nickel sulfide model (g) numerous low tonnage quartz veins containing gold and some silver (h) small tonnage copper veins, usually in mafic volcanic rocks (i) one apparently low grade porphyry molybdenum deposit is known; others possible (j) copper and minor silver in sedimentary rocks associated with basalt; deposit type not clear (k) low grade gold-bearing conglomerate (l) apparently low grades of chromite in small masses of dunitite; other deposits possible (m) porphyry molybdenum model (n) podiform chromite model</p>
--	---	---	--	--	--	--

36. (a) Mo-Cu-Au--porphyry and vein (b) Ag-Au-Sb--vein (c) Cu-Ag--porphyry and vein (d) Au-Ag--placer

(e) Cu(Ag)--sub-aerial volcanogenic (f) U--one occurrence reported, adjacent to granitic pluton

Includes broad, mainly mountainous regions in the upper Susitna and Chulitna River drainage systems, largely Mesozoic flysch and Tertiary and Cretaceous granitic plutons; local Tertiary and Cenozoic subaerial volcanic rocks. (a) A few known porphyry-type deposits and thin veins associated with Tertiary volcanic rocks. (b) Locally stibnite-rich veins in diverse host rocks; typically associated with Tertiary intrusive rocks. (c) Poorly known deposits mainly related to granitic rocks; includes at least one copper-bearing porphyry. (d) Stream and bench placers. (e) Tertiary volcanic rocks are favorable hosts for subaerial volcanogenic deposits.

Minor placer gold production; a little silver recovered from the placers. Tertiary deposits apparently have small resources.

Parts of the Talkeetna and Talkeetna Mountains quadrangles that are covered by modern U.S. Geological Survey studies including reconnaissance geology, geochemistry, and geophysics; the remainder of the area has scant geologic coverage and no known geochemistry or geophysics; a little recent interest by industry.

On the basis of its known deposits, the potential of the area is low; however, the area has been scantily prospected, and it is geologically favorable, particularly for porphyry type deposits.

(a) Several porphyry molybdenum deposits have been found; favorable geology and scanty exploration suggest that more deposits may occur. (b) A few small tonnage veins containing gold, silver, and some antimony are known; others likely. (c) Several poorly known deposits that may be the porphyry copper type; others may be present. (d) Gold-bearing stream and bench placers; minor gold and a little silver produced; apparently small amounts remain. (e) Mafic volcanic rocks may contain local concentrations of copper. (f) One uranium-bearing occurrence adjacent to a small granitic pluton.

90% 50% 10% chance that there are 0 1 3 deposits or more

(f) porphyry molybdenum

(a) porphyry molybdenum

(c) porphyry copper

AREA: 40. MAJOR TYPES OF KNOWN DEPOSIT: (a) Cu(Ag,Zn,Pb)--submarine volcanogenic; (b) Sb(Au)--vein; (c) Au(Ag)--placer; (d) Au(Ag)--vein

SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES): (e) Cu(Mo)--porphyry

GEOLOGIC CONTROL(S) OF MINERAL RESOURCES: Northeast flank of eastern Alaska Range; contains Paleozoic and probably Precambrian metamorphosed volcanic and sedimentary rocks, Mesozoic and Tertiary granitic rocks, and Cenozoic surficial deposits. (a) Massive and disseminated sulfide mineralization associated with Paleozoic metamorphic rocks, chiefly felsic metavolcanic rocks; local veins as much as 6 m thick, that mainly cut Paleozoic metamorphic rocks. (c) Stream and bench placers in precious metal-bearing veins, probably related to Tertiary igneous activity. (e) Speculative porphyry-type deposits on the basis of favorable host rocks.

PRODUCTION AND RESOURCE INFORMATION: The only known lode production from the area consists of small tonnages of antimony ore that have been mined intermittently during recent years; small quantities of gold with subordinate byproduct silver have been recovered from the placer deposits; the predominant resource potential of the area is in recently discovered submarine volcanogenic deposits, which, although not as yet thoroughly explored, portend significant resources; the type of the deposit with the association of speculative porphyry-type deposits, are believed to be small

STATUS OF GEOLOGIC INFORMATION: The Hlabosna and Tanacross quadrangle parts of the area are covered by modern reconnaissance geologic, geochemical, and geophysical studies; knowledge of that part of the area in the Mount Hayes quadrangle is based on older U.S. Geological Survey work supported by State geologists; active recent prospecting for submarine volcanogenic deposits and intermittent small-scale activity at an antimony mine and a few gold deposits, chiefly placers

ADDITIONAL COMMENTS: Recent prospecting interest focusing on the submarine volcanogenic deposits has been high and has resulted in a rash of claim staking; these deposits constitute a highly significant potential resource, but although several are currently lack sufficient data for mineral resource estimates; most of the area is considered geologically favorable for the submarine volcanogenic deposits, yet large tracts remain that lack systematic exploration

SUMMARY OF MINERAL RESOURCE POTENTIAL: (a) Large area favorable for submarine volcanogenic deposits that contain copper, lead, zinc, and silver; recent claim staking, yet large tracts remain that lack systematic exploration; possibly large number of deposits some of which could contain large tonnages. (b) Antimony-bearing veins; two deposits are as much as 6 m thick; other deposits possible. (c) Small tonnage placer gold deposits; placers are both stream and bench; small amounts of gold and silver produced. (d) Gold-bearing veins having small tonnages; no recorded production. (e) Favorable host rocks for porphyry copper and molybdenum deposits; no deposits known.

ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE AT LEAST THE NUMBER PREVIOUSLY LISTED):

GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (UN TABLE 2):

41.	(a) Cu(Au, Ag, Zn, Pb)--submarine volcano- genic (b) Au(Ag)--placer (c) Cu(Mo, Ag)--por- phyry (d) Cu(Fe)--con- tact metamorphic (e) Zn(Ag, Pb, Cu)-- replacement (f) Au, Ag--vein (g) Sb(Ag)--vein (h) Pb--sedimen- tary	(1) Mo(Au)--vein and porphyry (2) Cu--magmatic (3) Au, Ag(Cu, Pb)-- breccia pipes	The main production area consists of 1,400 kg and 1,600 kg (45,000 and 50,000 ounces) of gold and some by-product silver re- covered from pla- cer operations pri- or to 1960; since 1960 about 10 of the placer mines have been worked on intermittent small amounts of gold and silver have been recov- ered from a few lode deposits; the mineral re- source potential of the area is large; in partic- ular, the sub- marine volcano- genic deposits, which are the fo- cus of current exploration, probably contain significant re- sources of sev- eral metals; re- sources in zinc- rich replacement deposits in the Mount Eielson district prob- ably are fairly large; porphyry resources may be in several other deposit types, such as the por- phyries and con- tact-metamorphic deposits, both of which are scan- dally explored; precambrian bearing veins are probably small, but some of these depos- its may support small-scale min- ing; identified placer gold re- sources are suf- ficient to sus- tain continued mining op- erations; por- phyry and con- tact-metamorphic resources may be large; the uran- ium resource po- tential of the area is unknown and is based on favorable geol- ogy and small- scale explora- tion	Geologic knowledge of area based on local studies by State Dept. of Geology detailed to reconnaissance; some geochemical and geo- physical studies; no sys- tematic investigations of exploration inter- est; several of these deposits have been drilled or are stated for drilling re- sults are not yet available; there is a strong likelihood that these deposits contain large re- sources of several metals including copper, zinc, gold, and silver; Terti- ary and Cretaceous subaerial sedimentary rocks with as- sociated volcanic rocks mantle large parts of the area; are locally exposed throughout this cover as windows and under- lie the Cretaceous and Tertiary cover at shallow depths at many places; the Cre- taceous and Tertiary sedimentary rocks are geologically favorable for uranium deposits, but to date, the prospecting in them has not disclosed significant deposits	Submarine volcano- genic deposits with metapelite of the mid-paleo- zoic to early Tertiary Schist are the fo- cus of much recent exploration inter- est; several of these deposits have been drilled or are stated for drilling re- sults are not yet available; there is a strong likelihood that these deposits contain large re- sources of several metals including copper, zinc, gold, and silver; Terti- ary and Cretaceous subaerial sedimentary rocks with as- sociated volcanic rocks mantle large parts of the area; are locally exposed throughout this cover as windows and under- lie the Cretaceous and Tertiary cover at shallow depths at many places; the Cre- taceous and Tertiary sedimentary rocks are geologically favorable for uranium deposits, but to date, the prospecting in them has not disclosed significant deposits	Several submarine volcanogenic deposits are known in the area; these deposits contain copper, lead, and zinc and lo- cally economic grades of gold and silver; large favorable area and re- placement exploration inter- est suggest that many deposits could be in area 41 (a) Stream, bench, and fluvial placer con- taining gold and some silver are numerous; past production of about 1500 kg gold; future production from known deposits likely and un- discovered deposits possible. (b) Three possible por- phyry copper deposits; area might contain a few porphyry deposits (c) Contact metamorphic deposits containing copper, iron, and possibly gold (d) Several replacement deposits containing zinc and lead with some silver and copper (e) Small tonnage veins that have gold and minor silver contents (f) Many small tonnage veins that contain anti- mony and gold (g) Favorable geology for sandstone uranium deposits; recent exploration activity has been encouraging; possi- bly a number of unfound deposits (h) Molybdenum-bearing quartz veins at one pros- pect; small tonnage; may be indicative of porphyry molybdenum deposits in area	90% 50% 10% chance that there are 4 9 20 deposits or more	(a) felsic and inter- mediate volcanogenic model
-----	---	---	---	---	--	--	---	--

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROLS OF MINERAL RESOURCES	PRODUCTION AND RESOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT THAT ARE PROBABLY MORE EXTENSIVE OR MORE DEPOSITIVE)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
43.	<p>(a) Cu, Zn (Ag, Pb) -- submarine volcanogenic</p> <p>(b) Sn, Ag (Pb) -- vein and disseminated</p> <p>(c) Ag (Pb) -- replacement, metamorphic?</p> <p>(d) Ag, Pb, Zn (Sb, U, Sn) -- vein</p> <p>(e) Cu -- sedimentary?</p> <p>(f) U (Zn) -- contact metamorphic</p> <p>(g) Sb (Ag) -- vein</p>	<p>(h) Cu -- magmatic</p> <p>(i) Au -- vein</p> <p>(j) Be, Sn -- greisen</p>	<p>Northwest flank of part of west-central Alaska Range; adjoins a favorable area on sheet 2 to the west; underlain by locally metamorphosed Paleozoic sedimentary rocks and Tertiary granitic plutons; small areas of Mesozoic sedimentary and volcanic rocks near southern margin of area disseminated sulfides associated with submarine basalt of Mesozoic age</p> <p>(b) Cassiterite-bearing veins and disseminations in hornfels near Tertiary granite</p> <p>(c) Disseminated silver-rich sulfides in dolomite and pyrrhotite</p> <p>(d) Silver-bearing base metal veins that contain small amounts of tin, antimony, and uranium; typically thin veins that are associated with Tertiary granite</p> <p>(e) Recently discovered thin stratiform lens of chalcocite in Paleozoic phyllite</p> <p>(f) Two old prospects on contact-metamorphic (skarn)-type deposits marginal to Tertiary magmatic pluton</p> <p>(g) Thin quartz veins that contain stibnite and, rarely, a little cinnabar</p> <p>(h) Occurrence of disseminated sulfides in a mafic dike</p> <p>(i) Prospect on thin gold-bearing quartz vein that cuts Paleozoic rocks</p> <p>(j) Beryl in greisen; speculative tin-bearing greisens</p>	<p>No production; small identified resources, mainly in the submarine volcanogenic, silver, and tin lodes; the area has been scantily prospected, and its known deposits are unexplored; possibly, the area contains important resources</p>	<p>Most of the area is in the Talkeetna quadrangle, which is covered by modern reconnaissance geologic, geochemical, and geophysical studies; that part of the area in the Mount McKinley quadrangle has scant coverage; recent exploration at a few of the tin, silver, or submarine volcanogenic deposits</p>	<p>This remote and inadequately explored area is favorable for additional discoveries; at least two of its potentially important deposits were discovered during recent U.S. Geological Survey investigations; at least one, other was found during recent private exploration</p>	<p>(a) Submarine volcanogenic deposits bearing copper, zinc, silver, and lead with gold possible; known deposits incompletely explored and the area is generally not well explored; probably other as yet undiscovered deposits</p> <p>(b) Tin and silver-bearing disseminations and veins; extent not fully known; other deposits possible</p> <p>(c) One known deposit containing silver and lead minerals disseminated in carbonate rocks; other deposits possible</p> <p>(d) Silver, lead, zinc, and minor amounts of tin, antimony, and uranium; contained in veins that are probably small tonnage</p> <p>(e) Thin (5 cm thick) stratiform lens containing copper; extent not known; possibly related to large tonnage sedimentary deposit</p> <p>(f) Two skarn deposits that contain copper and some zinc; a few others possible</p> <p>(g) Several antimony-bearing veins are known in the northern part of area 43; tonnages probably small</p> <p>(h) Several greisen deposits that contain beryl or minor tin; tonnages probably small</p>	(a) mafic volcanogenic mode)	(f) copper skarn mode)

TABLE 2. METALLIFEROUS MINERAL RESOURCE DATA FOR WESTERN SOUTHERN ALASKA
(Refers to sheet 2)

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS PRESENT AND THE NUMBER PROSPECTED OR MORE DEPOSITS	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
1.	(a) Au(Ag)--vein placer (b) Au(Ag,Pt)--placer	(c) Mn--disseminated?	Central parts of Kodiak and Afognak Islands; underlain by upper Mesozoic flysch and local Tertiary granitic plutons (a) Gold-bearing quartz veins, mainly less than 1 m thick, that generally are localized in upper Mesozoic flysch (b) Beach placers and a few stream and dune placers (c) Tungsten prospects in upper Mesozoic graywacke	Production data sketchy; the lone deposits probably produced less than 2000 oz of gold and little byproduct silver; placer gold production probably a little larger and includes minor amounts of byproduct silver and platinum; some resources in placers, but they probably are small	Recent reconnaissance mapping by U.S. Geological Survey and University geologists; scant recent exploration; information from industry and known deposits; little geochemical or geophysical data	Much of area 1 is covered by dense vegetation and is difficult to prospect; on the basis of geologic, mineralogical, and known deposits, the area's main resource potential is for gold, but this is regarded as only of moderate significance	(a) Gold-bearing quartz veins; generally small tonnage; total production probably less than 32 kg gold, probably were difficult to prospect area (b) Beach, a few stream, and dune placers that contain gold and some silver and platinum; some past production; probably small amounts remain	90% 50% 10% chance that there are 0 1 3 deposits or more	(a) mafic volcanogenic model
2.	(a) Cu(Au,Zn,Pb)--submarine volcanogenic (b) Au(Ag)--placer	(c) Cu--magmatic; one occurrence (d) Au(Ag)--vein	Southeastern part of Kodiak Island and some nearby islands; area contains Tertiary, mainly flyschoid, rocks and scattered Tertiary plutons (a) Disseminated sulfides, including chalcopryrite, in Tertiary sedimentary and volcanic rocks (b) Mainly beach placers (c) Mostly disseminated sulfides in a gabbro sill (d) Areas near Tertiary plutons are favorable for thin, gold-bearing quartz veins	Possibly a little placer gold produced; otherwise no production; known resources are scant; resource potential rests on significant new discoveries, particularly of submarine volcanogenic deposits	Recent reconnaissance mapping by Government geologists, but very little geochemical or geophysical data; scant recent industry	The resource potential of area 2 is regarded as low; however, the area is geologically favorable for submarine volcanogenic deposits associated with mafic lavas, and possibly, it contains significant, undisturbed deposits of this type	(a) Several submarine volcanogenic deposits associated with mafic lavas are known; deposits usually contain copper, zinc, and gold; several other deposits possibly exist (b) Gold-bearing beach placers; generally small tonnage	90% 50% 10% chance that there are 0 1 3 deposits or more	(a) mafic volcanogenic model
3.	(a) Au(Ag)--placer (b) Au(Ag)--vein	(c) Cu(Ag)--vein (d) Cu(Ag,Zn)--submarine volcanogenic (e) Cu(Mo)--porphyry (f) Zn--magmatic	Northern parts of Kodiak and Afognak Islands, northwest of Border Ranges fault; underlain by upper Mesozoic flysch and Tertiary volcanic rocks, Cretaceous and Tertiary granitic rocks, and local ultramafic rocks and gabbro (a) Mainly beach placers (b) Thin quartz veins in or near granitic rocks (c) Some prospect on a thin copper-bearing vein in a fault zone (d) Suspected deposits associated with mafic volcanic rocks (e) Suspected deposits associated with granitic rocks (f) Some of the ultramafic rocks are favorable for chromite deposits	Small, but unknown amounts of gold recovered from the beach placers; otherwise no known production; area small but area may contain important undiscovered resources	Covered by recent reconnaissance geologic mapping but little available geochemical or geophysical data; scant recent industry	The area is geologically favorable for several types of mineral deposits, and it has been prospectively investigated for these reasons: it is inferred to have at least a moderate resource potential; the Barran Islands, to the north, are geologically similar to parts of area 3 and are regarded as having some favorability	(a) Mainly beach placers that contain gold and some silver; probably small tonnage (b) Small tonnage gold-bearing quartz veins; several are known, others possible (d) Submarine volcanogenic deposits with copper, zinc and gold are possible; area is not well prospected and is geologically favorable (e) Possible copper-bearing porphyry type deposits; part of area is favorable and completely explored (f) Portion of area is favorable for podiform chromite deposits; none known but area has not been thoroughly prospected	90% 50% 10% chance that there are 0 1 3 deposits or more 90% 50% 10% chance that there are 0 1 2 deposits or more	(d) mafic volcanogenic model (e) porphyry copper model (f) podiform chromite model

MAJOR TYPES OF KNOWN DEPOSITS

SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)

GEOLOGIC CONTROL(S) OF MINERAL RESOURCES

PRODUCTION AND RE-SOURCE INFORMATION

STATUS OF GEOLOGIC INFORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL RESOURCE POTENTIAL

ESTIMATED NUMBER OF DEPOSITS PLACED AT RISK THAT THERE ARE THE NUMBER PRE-SERIALIZED OR MORE DEPOSITS

GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)

4.

(a) Ag (Pb, Zn, Cu)--vein, replacement, breccia pipe
 (b) Cu (Pb, Zn)--vein, replacement
 (c) Pb, Zn--vein, replacement
 (d) Ni, Cu (Zn)--magmatic
 (e) Au (Ag)--placer
 (f) Au--vein

Southeastern part of McGrath quadrangle and north-central part of Lime Hills quadrangle, in northwestern Yukon Territory, Alaska. The area is largely underlain by Paleozoic sedimentary rocks, including abundant carbonate rocks; local Tertiary and Cretaceous(?) plutons. (a) Sulfide-rich veins, as much as 3 m thick, and pods and lenses; all with high silver contents; generally barren. (b) Tertiary plutons; less commonly in the plutons. (c) Chalcopyrite and sphalerite-bearing tuffite zones as much as 3 m wide. (d) Disseminated and locally massive sulfides in diatremes. (e) One stream placer known in area. (f) Two prospects on thin, gold-bearing veins in or near Tertiary granitic plutons.

No production; some identified re-sources in two of the silver-rich lenses. The area has been explored by a few diamond drill holes; probably significant potentials for silver and less important and less important potential resources of the other commodities known in the area.

Broad reconnaissance mapping accompanied by some geochemical sampling and a detailed local study of the area is in progress. The area is remote and during the early days attracted very little prospecting; recent exploration interest in the area is largely attributable to discoveries made during the course of U.S. Geological Survey investigations in the past decade.

The area adjoins area 43 on sheet 1 and locally extends into the McGrath study region; the area is remote and during the early days attracted very little prospecting; recent exploration interest in the area is largely attributable to discoveries made during the course of U.S. Geological Survey investigations in the past decade.

(a) Veins, replacement, and possible breccia pipe deposits containing silver, lead, zinc, and copper; all known deposits have high silver contents; as much as 3 m thick; probably other undiscovered deposits; possibly other lead and zinc-bearing deposit types in the carbonate rocks. (b) Veins containing copper and some lead and zinc; generally small tonnage; may, in some cases, be in the form of disseminated mineralization, such as porphyry copper type. (c) Several contact metamorphic deposits that contain copper, silver and zinc are known; others possible. (d) Disseminated and locally massive sulfides containing nickel and copper exist in at least one locality. (e) One gold-bearing stream placer is known in this area; others possible. (f) Two small tonnage veins bearing gold are known; others possible.

(c) copper stann model

(d) nickel sulfide mine

AREA COVERED BY MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL RESOURCES (INCLUDING MINOR OCCURRENCES)	GEOLOGIC CONTROL (S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)	
5.	(a) Cu-Mo, Ag, Zn--porphyry (b) Mo, Cu, Au, Ag--porphyry (c) Ag, Cu, Au, Pb, Zn--breccia pipe (d) Au, Ag--vein (e) Cu, Zn, Ag, Pb--vein (f) Au, Ag--placer	(g) Cu--contact metamorphic (h) Mo, Cu--vein (i) Sb--vein (j) Cu, Ag, Zn--submarine volcanic-genic	Northwest flank of west-central Alaska Range; joins area 30 of sheet 1; underlain by locally metamorphosed Mesozoic sedimentary rocks and volcanic rocks. Small volcanic rocks, and Mesozoic and Tertiary granitic and Tertiary granitic rocks (a) Disseminated sulfides and sulfide-bearing veins in altered Tertiary plutons that probably mainly represent subvolcanic phases of Tertiary igneous activity (b) Similar to (a) except that the chief ore mineral is molybdenite (c) Precious and base metal lodes associated with Tertiary eruptive centers (d) Thin veins genetically related to Tertiary plutonism; a few are silver-rich (e) Typically thin polymetallic veins in diverse host rocks (f) Stream placers (g) Occurrence of sulfide blebs and disseminations in meta-sedimentary rocks adjacent to Tertiary granite (h) Thin molybdenite-bearing quartz veins in Tertiary igneous rocks (i) Occurrence of a thin stibnite-bearing vein in Mesozoic sedimentary rocks (j) Occurrences of submarine volcanic-genic deposits, mainly associated with felsic volcanic rocks	No production; the main potential re-sources of the area appear to be in porphyry-type deposits; both for copper and molybdenum. The area has been poorly explored and it contains several deposit types that may contain significant resources	Reconnaissance mapping and geochemical sampling for most of area; reconnaissance gravity survey; aeromagnetic survey for that part of area in Lake Clark region. Little interest by industry	The area includes large and poorly explored remote tracts; current exploration interest in the area is largely a result of recent U.S. Geological Survey and U.S. Bureau of Mines sponsored investigations	(a) A number of recently discovered porphyry copper deposits that contain minor contents of molybdenum, silver, and zinc are known; large and poorly explored area and incompletely explored deposits probably discovered (b) About five porphyry-type molybdenum deposits that contain some copper, gold, and silver are now known; others possible (c) Several deposits that are probably breccia pipes have been found; they contain silver, copper, and some gold, lead, and zinc; geology is favorable for other deposits of this type (d) Small tonnage veins containing gold and locally silver; three deposits known; others possible (e) Copper, zinc, lead, and silver-bearing veins; two deposits known; probably low tonnage (f) Several gold and silver-bearing stream placers have been found (g) Two suspected submarine volcanic-genic deposits are known; area is poorly explored and may contain other deposits of this type	(a) porphyry copper model 90% 50% 10% chance there are 5 9 15 deposits or more (b) porphyry molybdenum model 90% 50% 10% chance there are 3 6 10 deposits or more	(j) felsic and intermediate volcanic-genic model

AREA
OUT-
LINED
ON
MAP

MAJOR TYPES OF KNOWN
DEPOSITS

SUSPECTED OR SPECULA-
TIVE TYPES OF MINERAL
DEPOSITS (INCLUDES
MINOR OCCURRENCES)

GEOLOGIC CONTROL(S) OF
MINERAL RESOURCES

PRODUCTION AND RE-
SOURCE INFORMATION

STATUS OF GEOLOGIC IN-
FORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL
RESOURCE POTENTIAL

ESTIMATED NUMBER OF
DEPOSITS (PERCENT
CHANCE THAT THERE
ARE THE NUMBER PRE-
SERVED OR MORE
DEPOSITS)

GRADES AND TONNAGES
FOR THIS DEPOSIT
TYPE (IN TABLE 2)

6.	(a) Cu(Fe,Zn)--contact metamorphic (Zn, Pb)--breccia pipe and vein (c) Cu(Mo, Ag)--porphyry (d) Au(Ag)--placer (e) Mo--vein and porphyry? (f) Au--vein (g) Cu(Mo, Ag)--replacement (h) Fe(Ti)--magmatic	(1) Cu(Zn)--submarine volcanogenic	Southern part of Alaska Range north of Inanna Lake, contains sedimentary volcanic, and intrusive rocks (a) Massive and disseminated deposits typically in Triassic carbonate rocks adjacent to Tertiary granite; generally contain abundant magnetite or hematite, but chert potential are minimal (b) Pyrometallitic preoxides and base metal deposits mainly associated with young eruptive or intrusive centers; chief potential values generally in silver or copper (c) Apparently weakly mineralized porphyry-type deposits associated with Tertiary granitic plutons (d) Streamlined, documented probable porphyry molybdenum deposit and a few deposits represented by molybdenite-bearing quartz veins (f) Thin gold-bearing quartz veins that cut upper Mesozoic flysch (g) Massive and disseminated sulfides mineralized zone 6 to 12 mi wide within Triassic marble (h) Breccia fragments of magnetite-rich pyroxenite in granite (i) Occurrences of copper and zinc sulfides in both felsic and mafic metavolcanic rocks; others suspected	Most of the area is included in U.S. Geological Survey Mine Investigations and is being oriented studies by the State Survey and the U.S. Bureau of Mines; when completed, and integrated with previous work, these studies will provide modern geologic, geochemical, and geophysical data; except for activity at the Kasna Creek property, recent exploration interest in the area is low	Area 6 is diversely mineralized and probably contains significant resources in identified resources on the order of several million tons with slightly less than 1 percent copper and about 27 percent iron; area is favorable for additional discoveries (b) Probably breccia pipe and vein deposits that contain silver and copper in places concentrations of gold, silver and lead; other deposits possible (c) Several apparently weakly mineralized porphyry-type copper deposits are known; other possibly richer deposits may occur in area 6 (d) About 6 stream placers that contain gold have been found; small production (e) One incompletely explored deposit that is probably a porphyry molybdenum type; other molybdenum-bearing veins are known; some may be related to porphyry type deposits (f) Small tonnage gold-bearing veins; a few known, others possible (g) Probable replacement deposit containing copper and minor concentrations of gold and silver; mineralized zone is 6 to 12 mi wide (h) Breccia fragments of iron-bearing magnetite proximate in granite at one locality (i) Several probable submarine volcanogenic deposits containing copper and silver; other deposits of this type	(a) copper starn model (c) porphyry copper model (e) porphyry molybdenum model (f) felsic and intermediate volcano-gulf massive sulfide model
6.	(a) Cu(Fe,Zn)--contact metamorphic (Zn, Pb)--breccia pipe and vein (c) Cu(Mo, Ag)--porphyry (d) Au(Ag)--placer (e) Mo--vein and porphyry? (f) Au--vein (g) Cu(Mo, Ag)--replacement (h) Fe(Ti)--magmatic	(1) Cu(Zn)--submarine volcanogenic	Southern part of Alaska Range north of Inanna Lake, contains sedimentary volcanic, and intrusive rocks (a) Massive and disseminated deposits typically in Triassic carbonate rocks adjacent to Tertiary granite; generally contain abundant magnetite or hematite, but chert potential are minimal (b) Pyrometallitic preoxides and base metal deposits mainly associated with young eruptive or intrusive centers; chief potential values generally in silver or copper (c) Apparently weakly mineralized porphyry-type deposits associated with Tertiary granitic plutons (d) Streamlined, documented probable porphyry molybdenum deposit and a few deposits represented by molybdenite-bearing quartz veins (f) Thin gold-bearing quartz veins that cut upper Mesozoic flysch (g) Massive and disseminated sulfides mineralized zone 6 to 12 mi wide within Triassic marble (h) Breccia fragments of magnetite-rich pyroxenite in granite (i) Occurrences of copper and zinc sulfides in both felsic and mafic metavolcanic rocks; others suspected	Most of the area is included in U.S. Geological Survey Mine Investigations and is being oriented studies by the State Survey and the U.S. Bureau of Mines; when completed, and integrated with previous work, these studies will provide modern geologic, geochemical, and geophysical data; except for activity at the Kasna Creek property, recent exploration interest in the area is low	Area 6 is diversely mineralized and probably contains significant resources in identified resources on the order of several million tons with slightly less than 1 percent copper and about 27 percent iron; area is favorable for additional discoveries (b) Probably breccia pipe and vein deposits that contain silver and copper in places concentrations of gold, silver and lead; other deposits possible (c) Several apparently weakly mineralized porphyry-type copper deposits are known; other possibly richer deposits may occur in area 6 (d) About 6 stream placers that contain gold have been found; small production (e) One incompletely explored deposit that is probably a porphyry molybdenum type; other molybdenum-bearing veins are known; some may be related to porphyry type deposits (f) Small tonnage gold-bearing veins; a few known, others possible (g) Probable replacement deposit containing copper and minor concentrations of gold and silver; mineralized zone is 6 to 12 mi wide (h) Breccia fragments of iron-bearing magnetite proximate in granite at one locality (i) Several probable submarine volcanogenic deposits containing copper and silver; other deposits of this type	(a) copper starn model (c) porphyry copper model (e) porphyry molybdenum model (f) felsic and intermediate volcano-gulf massive sulfide model

AREA
OUT-
LINED
ON
MAP

MAJOR TYPES OF KNOWN
DEPOSITS

SUSPECTED OR SPECULA-
TIVE TYPES OF MINERAL
DEPOSITS (INCLUDES
MINOR OCCURRENCES)

GEOLOGIC CONTROL (S) OF
MINERAL RESOURCES

PRODUCTION AND RE-
SOURCE INFORMATION

STATUS OF GEOLOGIC IN-
FORMATION

ADDITIONAL COMMENTS

SUMMARY OF MINERAL
RESOURCE POTENTIAL

ESTIMATED NUMBER OF
DEPOSITS (PERCENT
CHANCE THAT THERE
ARE THE NUMBER PRE-
SENTED OR MORE
DEPOSITS)

GRADES AND TONNAGES
FOR THIS DEPOSIT
TYPE (SEE TABLE 2)

7.	(a) Cu, Fe (Au, Ag)-- contact metamorphic (b) Fe (Ti)--magnetic (c) Cu (Au, Ag, Mo)-- porphyry (d) Cu (Ag)--vein (e) Cu (Au, Ag)--in- trusive breccia (f) Ag, Au (Pb, Zn)-- vein (g) Au (Ag)--placer (h) Au--vein	(1) Mo (Cu, Ag)--porphyry (2) Ag (Au, Cu)--volcanic breccia	Part of Alaska Peninsula; dated by radiometric methods. Intrusive rocks of Jurassic age that form the core of the Ala- ska-Alutian Ranges batho- olith; local upper Paleo- zoic and Mesozoic sedi- mentary and volcanic rocks that in places are metamorphosed, upper Me- sozoic and Tertiary gran- itic plutons, and fairly extensive Tertiary vol- canic rocks. (a) Massive and dissemi- nated contact-metamorphic, mainly skarn, deposits chiefly localized in car- bonate rocks adjacent to granitic plutons; poten- tial chief values mainly for copper contained in chalcopyrite; a few have principal potential values in iron (from magnetite) (b) Massive and dissemi- nated iron-rich (mainly silicified) magnetite segregations typically localized in mafic phases of Jurassic granitic plu- tons (c) Copper-bearing porphyry- type deposits in Jurassic or Tertiary granitic plu- tons (d) Thin copper-bearing veins in diverse geologic settings (e) Disminuated sulfides in an intrusive breccia quartz dykes and shibbo- rocks may be akin to porphyry- type mineralization (f) Thin polymetallic preci- pitated metal-bearing veins in limestone or volcanic rocks (g) Stream placers (h) Thin gold-bearing quartz veins in Tertiary volcanic rocks (i) Occurrences of weakly mineralized molybdenum porphyry deposits in Tertiary volcanic rocks (j) Suspected precious metal lodes associated with Ter- tiary eruptive centers	Most of the area is in the Iliamna quadrangle, which has good con- ditions for iron, magne- tite, and some super- genetic iron-ore and metamorphic de- posits and probably a little placer gold; the area contains large, but low-grade iron resources incor- porated in mafic in- trusive rocks; these have been estimated to be about 100 million tons, containing between 12 and 15 percent ferrous ox- ide (FeO); several other deposit types, particularly the por- phyry-type and con- tact-metamorphic de- posits, probably contain significant resources	Production from the area consists of the export of iron ore from one of the con- tact-metamorphic de- posits and probably a little placer gold; the area contains large, but low-grade iron resources incor- porated in mafic in- trusive rocks; these have been estimated to be about 100 million tons, containing between 12 and 15 percent ferrous ox- ide (FeO); several other deposit types, particularly the por- phyry-type and con- tact-metamorphic de- posits, probably contain significant resources	Although the area's magnetic iron-titanium resources are concentrated in low grade and in view of the world's vast iron resources, they probably will retain their sub- economic status for a long time; estimat- ing the potential re- sources in the other deposit types is ham- pered by the dearth of data on these types of deposits and the lack of thorough search of new deposits	90% 50% 10% chance that there are 5 7 10 deposits or more	(a) Copper and iron-bearing contact metamorphic deposits; about 6 deposits known; a few un- found deposits may remain (b) At least 11 iron-rich (mainly titaniferous magnetite) magmatic deposits have been found; total estimated resources in these deposits of several million tons of iron (15 percent ferrous oxide) (c) Two copper-bearing porphyry-type deposits that contain gold, silver, and molybdenum are known; favorable geology and scant exploration for this type suggest that un- found deposits remain (d) Several low tonnage copper and silver-bearing veins are known (e) Two probable intrusive breccia deposits containing copper, gold, and silver; may be akin to porphyry-type mineralization (f) Small tonnage veins containing silver, gold, lead and zinc (g) Several stream placer deposits, probably with small gold production (h) Small tonnage gold-bearing quartz veins (i) Several occurrences of weakly mineralized porphyry deposits that contain molybdenum; others possible	(a) copper skarn model (c) porphyry copper model (i) porphyry molybdenum model
----	--	---	--	--	---	---	--	--	---

AREA OUTLINED ON MAP	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDE MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANGE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
8. (Ad- joint area 5 on central Alaska map)	(a) Fe(tl)--magnetic (b) Au--vein (c) Au--placer (d) Hg--vein	Mainly lowlands in central part of Dillingham quadrangle; largely mantled by Quaternary surficial deposits; Mesozoic sedimentary rocks and Cretaceous and Tertiary granitic rocks crop out on isolated low hills throughout the area buried titaniferous magnetite deposit in pyroclastics; discovered by diamond drilling in magnetic area detected during an industry-sponsored aeromagnetic survey in 1968	The one known gold lode in the area reportedly yielded a small amount of gold; the dominant and probably only significant resources in the area are the magmatic iron-titanium deposits; these deposits, which have been reported by diamond drill holes, are believed to contain about 2.4 billion tons of hypothetical resources averaging 15 to 17 percent total iron and 10.5 to 12 percent magnetic iron	The area is poorly known geologically and geochemically; industry-sponsored aeromagnetic survey, but otherwise, scant geophysical coverage; with the exception of diamond drilling the pyroclastic body during the 1960's, there has been scant industry interest in the region	The titaniferous magnetite deposits are the predominant potential resource of the area; the area may contain significant concealed deposits of other types, but, in general, these would be extremely difficult to discover	(a) One buried iron-rich (titaniferous magnetite) magmatic deposit has been discovered; it is believed to contain about 2.4 billion tons averaging 15 to 10 percent total iron and 10 percent magnetic iron; other concealed deposits of other types are possible (b) Several low tonnage gold-bearing veins have been found (c) Gold-bearing stream placers; one prospect and one occurrence known (d) Mercury vein deposits are suspected	(d) mercury mode	42	

AREA OUTLINED BY USGS	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECULATIVE TYPES OF MINERAL DEPOSITS (INCLUDES MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRE-SENTED ON MORE DEPOSITS)	GRADES AND TENNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
9.	(a) Cu(Au, Mo, Ag) -- porphyry (b) Au(Ag) -- vein and lens (c) Au(Ag) -- placer (d) Mo(Co) -- porphyry (e) Cu(Ag) -- vein (f) Cu(Au, Ag) -- replacement (g) Fe -- magmatic	(h) U -- placer	Mainly northeastern part of Alaska Peninsula; geology dominated by Cenozoic volcanic rocks related to Aleutian arc volcanism; also contains Mesozoic and rare upper Paleozoic sedimentary and volcanic rocks; Cenozoic subvolcanic plutons and possibly some Mesozoic plutons (a) Porphyry-type deposits genetically related to subvolcanic plutons associated with Aleutian arc igneous activity; localized in the plutons or in various nearby host rocks (b) Locally rich gold-bearing quartz veins and lenses thick; typically called in tertiary volcanic or intrusive rocks (c) Beach and stream placers (d) Molybdenum-bearing porphyry-type deposits genetically related to tertiary subvolcanic plutons (e) Generally thin copper-bearing veins, mainly in host rocks (f) Chalcopyrite-bearing lenses in Jurassic sedimentary rocks (g) Local concentrations of magnetite in mafic phases of granitic plutons (h) Reported placer claim for uranium; probably contains some radioactive refractory heavy minerals	Lode gold production of approximately 3,200 kg (100,000 ounces) and some by-product silver, mainly from the Apollo mine on Unga Island; placer gold produced in the Apollo area, on 19 kg (600 ounces) chiefly from beach placers; some identified gold resources in the Apollo mine; the principal potential resources of the area are associated with the Mesozoic volcanic rocks, mainly low-grade porphyry copper deposits; about 20 of these deposits have attracted recent exploration activity, including exploratory drilling on several of them; in addition there are numerous unexamined zones in the region that may indicate porphyry-type deposits; the one known porphyry molybdenum deposit in the area is currently under exploration; definitive information on the porphyry deposits is not available; unconfirmed information suggests that the porphyry deposits are large, contain less than 0.4 per cent copper, and are slightly richer in gold than the better-known porphyry deposits in the Mesozoic; resource potential of the other deposits in the area are probably low, but some deposits, particularly the copper-bearing replacements, may be worthy of exploration	Diverse degrees of geologic mapping, but, at least, sketchy reconnaissance mapping for the entire area; the Chituk and Sutwik Island quadrangles are currently being investigated under the U.S. Geological Survey AMHP program in which reconnaissance geologic, geochemical, and geophysical studies in the area by industry, but the results of these investigations are mainly privileged; active industry exploration, mainly for porphyry-type deposits, during the past decade	The area includes a large part of a potentially major porphyry copper province associated with Aleutian arc tectonics and igneous activity; most of these deposits represent recent discoveries and have probably already been explored, probably many deposits in this area (b) Locally rich gold-bearing veins and lenses as much as several meters thick; past production about 3200 kg gold (c) Numerous gold-bearing stream and beach placers; the richest is about 19 kg gold; beaches northwest of area 9 contain local iron and titanium-bearing placers that carry minor amounts of gold and, rarely, platinum (d) Molybdenum-bearing porphyry-type deposits; only a few known or suspected; others possible (e) Copper and gold-bearing veins; probably small tonnage; some may be related to other types of mineralization such as porphyry (f) Possible replacement deposits containing copper, gold and silver (g) Several local concentrations of iron (magnetite) in magmatic deposits	90% 50% 10% chance that there are 30 50 75 deposits or more	(a) Island arc porphyry copper mode (d) porphyry molybdenum mode	

AREA NAME AND LOCATION	MAJOR TYPES OF KNOWN DEPOSITS	SUSPECTED OR SPECIALLY-TYPED MINERAL DEPOSITS (UNUSUAL MINOR OCCURRENCES)	GEOLOGIC CONTROL(S) OF MINERAL RESOURCES	PRODUCTION AND RE-SOURCE INFORMATION	STATUS OF GEOLOGIC INFORMATION	ADDITIONAL COMMENTS	SUMMARY OF MINERAL RESOURCE POTENTIAL	ESTIMATED NUMBER OF DEPOSITS (PERCENT CHANCE THAT THERE ARE THE NUMBER PRESENTED OR MORE DEPOSITS)	GRADES AND TONNAGES FOR THIS DEPOSIT TYPE (IN TABLE 2)
10. 11. 12.	-----	(a) Au(Ag)---ve in	<p>Area 10 includes the Aleutian Islands; area 11 includes the Shumagin Islands; and area 12, the Sanak Islands; all three areas are underlain by Tertiary granitic rocks and upper Mesozoic sedimentary rocks.</p> <p>(a) Suspected thin gold-bearing quartz veins genetically related to the Tertiary granitic rocks.</p>	<p>No known mineral deposits in any of the three areas. Gold-bearing vein deposits are analogous to those found in upper Mesozoic flysch terranes in southern Alaska and are genetically related to Tertiary plutons; the mineral resource potential of areas 10, 11, and 12 is regarded as low.</p>	<p>Some reconnaissance geologic mapping but no available geochemical or geophysical information. Little or no exploration interest for metalliferous deposits.</p>	<p>Chirikof Island, southeast of area 10, contains one deposit known from beach placer, but the resource potential of the island is regarded as minimal.</p>	<p>(a) Gold-bearing quartz veins are suspected; tonnage and grade of deposits known but areas 10, 11, and 12 have favorable geology.</p>		
13.	(a) Cu(Mo, Au, Ag)--- porphyry	-----	<p>Southwestern part of Alaska Peninsula; geology dominated by Cenozoic volcanic rocks related to the Aleutian volcanic arc; local Tertiary subvolcanic plutons, Tertiary subaerial sedimentary rocks, and possibly some Cretaceous sedimentary rocks.</p> <p>(a) Porphyry copper deposits are known in the Aleutian volcanic arc; localized in Tertiary volcanic, intrusive, or sedimentary rocks; more than 20 known occurrences.</p>	<p>No mines or prospects; this scantily explored area contains numerous porphyry copper occurrences and is favorable for significant copper resources and lesser resources of byproduct gold, molybdenum, and silver.</p>	<p>Broad reconnaissance geologic mapping; no available geochemical or geophysical data, although some geochemical exploration was conducted by mining companies; moderate recent interest by industry.</p>	<p>Area 13 is part of a potentially major porphyry copper province that mainly is in area 9; it is poorly explored and is regarded as having a good resource potential.</p>	<p>(a) Porphyry copper deposits associated with subvolcanic plutons, possible dykes, and sills; more than 20 altered areas that may be porphyry copper deposits are known; area is poorly explored and may contain many deposits.</p>	<p>90% 50% 10% chance that there are 10 20 35 deposits or more</p>	(a) Island arc porphyry copper model

TABLE 4. GRADE AND TONNAGE MODELS

(metric units)

NS, not significant; *, significant at 5-percent level; **, significant at 1 percent level

Deposit Type	Variable (units)	Number of deposits used	Correlation Coefficients	90 percent of deposits have at least	50 percent of deposits have at least	10 percent of deposits have at least
Porphyry Copper	Tonnage (millions of tons)	41	with tonnage = -0.07 NS	20	100	430
	Average copper grade (percent)	41		0.1	0.3	0.55
	Average molybdenum grade (percent Mo)	41		0.0	0.008	0.031
Island Arc Porphyry Copper	Tonnage (millions of tons)	41	with tonnage = -0.07 NS	20	100	430
	Average copper grade (percent)	41		0.1	0.3	0.55
	Average molybdenum grade (percent Mo)	41		0.0	0.008	0.031
	Average gold grade - locally significant but not determined					
Porphyry Molybdenum	Tonnage (millions of tons)	31	with tonnage = -0.05 NS	1.6	24	340
	Average molybdenum grade (percent Mo)	31		0.065	0.13	0.25
Podiform Chromite	Tonnage of Cr ₂ O ₃ (tons)	268		15	200	2,700
Copper Skarn	Tonnage (millions of tons)	38	with tonnage = -0.44**	0.08	1.4	24
	Average copper grade (percent)	38		0.86	1.7	3.5
	Average gold grade, locally significant, but not determined					
45 Mafic Volcanogenic	Tonnage (millions of tons)	37	with tonnage = -0.13 NS	0.24	2.3	22.0
	Average copper grade (percent)	37		1.1	2.2	4.1
	Average zinc grade excluding deposits without reported grades (percent)	19	with tonnage = 0.03 NS	0.3	1.3	5.5
	Average gold grade - locally significant but not determined					
Felsic and Intermediate Volcanogenic Massive Sulfide	Tonnage (millions of tons)	89	with tonnage = -0.41**	0.19	1.9	18.0
	Average copper grade (percent)	89		0.54	1.70	5.40
	Average zinc grade excluding deposits without reported grades (percent)	41	with tonnage = 0.25 NS	1.40	3.80	10.00
	Average lead grade excluding deposits without reported grades (percent)	14	with tonnage = -0.02 NS	0.20	0.95	4.80
	Tonnage contained gold excluding deposits without reported gold (tons)	38	with tonnage = 0.78**	0.27	2.90	32.00
	Tonnage contained silver excluding deposits without reported silver (tons)	46	with tonnage = 0.82**	5.00	80.00	1300.00
Nickel Sulfide	Tonnage (millions of tons)	48	with tonnage = -0.03 NS	0.23	1.20	5.90
	Average nickel grade (percent)	48		0.32	0.61	1.20
	Average copper grade (percent)	48	with tonnage = 0.03 NS with nickel grade = 0.04 NS	0.18	0.47	1.20
Mercury	Tonnage of contained mercury (tons)	165		0.09	3.10	120.00
Vein Gold	Tonnage of contained gold (tons)	43		0.29	3.30	38.00
Skarn/Tactite Tungsten	Tonnage (millions of tons)	31	with tonnage = -0.34 NS	0.024	0.63	17
	Average tungsten grade (percent W)	31		0.24	0.51	1.10