


CORRELATION OF MAP UNITS




Geology generalized after Csejtey and others, 1978



DESCRIPTION OF MAP UNITS

- Qs SURFICIAL DEPOSITS, UNDIFFERENTIATED (Quaternary).
- Tv VOLCANIC ROCKS, UNDIFFERENTIATED (Pliocene to Pleistocene(?)--Felsic and mafic subaerial volcanic rocks and related shallow intrusions.
- Tsgd TERTIARY SEDIMENTARY ROCKS, UNDIFFERENTIATED (Pliocene to Miocene)--Terrestrial, mostly fluvialite strata with a few lignite interbeds.
- Tsmg GRANODIORITE (Eocene).
- Tkg BIOTITE AND HORNBLENDE GRANODIORITE (Paleocene, in part early Eocene).
- Pzv SCHIST, MICAHITE, AND GRANITE (Paleocene Intrusive and metamorphic ages)--Migmatitic border zone of biotite and hornblende granodiorite.
- Jpm TONALITE (Upper Cretaceous and lower Paleocene).
- Jtr ADAMELITE (Upper Cretaceous and lower Paleocene).
- Jnc GRANITIC ROCKS, UNDIFFERENTIATED (Cretaceous and/or Tertiary).
- Jta ARKOSE RIDGE FORMATION (Lower and/or Upper Cretaceous).
- Jsn MATANUSKA FORMATION (Lower and Upper Cretaceous).
- Jsk SEDIMENTARY ROCKS, UNDIFFERENTIATED (Lower Cretaceous)--Shallow marine sequence of calcareous sandstone, claystone, and massive chertic limestone.
- Jsv ARGILLITE AND LITHIC GRANWACKE (Lower Cretaceous)--Intercalated, marine, flyschlike sequence.
- Jka SEEDIMENTARY AND VOLCANIC ROCKS, UNDIFFERENTIATED (Upper Jurassic)--Marine sequence of argillite, graptolite, conglomerate, and andesitic to latitic felsopar porphyry dikes and intercalated flows.
- Jtr TRONDHJEMITE (Upper Jurassic).
- Jnc JURASSIC SEDIMENTARY ROCKS, UNDIFFERENTIATED (Middle and Upper Jurassic)--Includes Haknek and Christina Formations, and Tuxedii Group.
- Jta CRYSTAL TUFF, ARGILLITE, CHERT, GRAPTOLITE, AND LIMESTONE (Lower to Upper Jurassic)--Shallow to moderately deep marine, intercalated sequence.
- Jsn PLUTONIC AND METAMORPHIC ROCKS, UNDIFFERENTIATED (Lower to Upper Jurassic)--Mainly quartz diorite, granodiorite, amphibolite, and gneisschist.
- Jsk TALKEETNA FORMATION (Lower Jurassic).
- Jsv METABASALT AND SLATE (Upper Triassic)--Intercalated, shallow-water marine sequence.
- Jka BASALTIC METAVOLCANIC ROCKS (Upper Triassic)--Mainly shallow water marine metabasalt flows.
- Pzv BASALTIC AND ANDESITIC METAVOLCANIC ROCKS (Pennsylvanian(?) and Early Permian)--Metamorphosed marine sequence of inter-layered basaltic to andesitic flows, tuffs, coarse volcanoclastic rocks, and subordinate rhyolite and limestone.
- Jsv ARGILLITE AND LITHIC GRANWACKE (Silurian?) to Middle Devonian--Intercalated marine sequence, probably continental margin deposits.

Base map from U.S. Geological Survey, 1:250,000 Talkeetna Mountains Quadrangle, Alaska, 1955



- EXPLANATION OF GEOCHEMICAL MAP SYMBOLS
- Location of stream sediment sample
 - Location of heavy mineral concentrate sample
 - Location of both stream sediment and heavy mineral concentrate sample
 - Heavy mineral concentrate sample with possibly significant barium value. Increase in symbol size indicates higher analytical value as shown on histogram.

EXPLANATORY STATEMENT

In the course of U.S. Geological Survey investigations of the Talkeetna Mountains quadrangle, 1118 stream sediment, 852 heavy mineral concentrate, and 501 rock samples were collected. All of these samples were analyzed for up to 30 elements by a six-step semi-quantitative spectrographic method (Grimes and Marranzino, 1968). Most of the stream sediment and rock samples were also analyzed for up to 4 elements by atomic absorption spectrophotometry, as described by Ward and others (1969). Although the present map shows all the sites where stream sediments were collected in the quadrangle only a portion of those samples were analyzed for barium; therefore analytical data for barium in stream sediments are not included in this report. The present map also shows the collection sites of all heavy mineral concentrates in the quadrangle. All of these samples have been analyzed for barium by the spectrographic method and the results are tabulated in the histogram and shown on the map. Complete analytical data plus location maps, station coordinates, and discussion of sampling and analytical procedures for samples from sites shown on the present map are published in a report by Miller and others (1978).

Concentration of metals in geochemical samples varies for different lithologies and in different areas. Because of this, as well as variability introduced from other sources such as sampling practice, analytical variance, and degree of chemical weathering, it is impossible to select a specific analytical level above which values might indicate the presence of barium deposits. For this reason, the analytical values have been grouped into ranges (see histogram), each range being represented by a different symbol on the map. Higher values may indicate a greater likelihood of barium deposits, but confidence levels are low for "single-element" anomalies and for results which are not supported by neighboring values.

EXPLANATION OF GEOLOGIC MAP SYMBOLS

- Contact, approximately located
- Approximate contact of surficial deposits
- Fault
- Long dashed where approximately located, short dashed where inferred; dotted where concealed. U indicates upthrust side where direction of displacement is known. Arrow indicates relative lateral movement
- Thrust fault
- Long dashed where approximately located, dotted where concealed. Teeth indicate upthrust side.
- Approximate axis of intense shear zone of variable width, possibly marking a thrust fault
- Dotted where concealed; teeth indicate possible upthrust side of postulated thrust

REFERENCES CITED

Csejtey, Bela, Jr., Nelson, W. H., Jones, D. L., Silberling, N. J., Dean, R. M., Morris, M. S., Lanphere, M. A., Smith, J. C., and Silberman, M. L., 1978, Reconnaissance geologic map and geochronology, Talkeetna Mountains quadrangle, northern part of Anchorage quadrangle, and southwest corner of Healy quadrangle, Alaska: U.S. Geol. Survey open-file rept. 78-558-A, 62p.

Grimes, D. J., and Marranzino, A. P., 1968, Direct-current arc and alternating-current spark emission spectrographic field methods for the semi-quantitative analysis of geologic materials: U.S. Geol. Survey Circ. 591, 6p.

Miller, R. J., Cooley, E. F., O'Leary, R. M., Garmezy, Larry, Csejtey, Bela, Jr., Smith, T. E., and Cleveland M. N., 1978, Analyses of geochemical samples from the Talkeetna Mountains quadrangle, Alaska: U.S. Geol. Survey open-file rept. 78-1052, 279 p.

Ward, F. N., Nakagawa, H. M., Harms, T. F., and Van Sickle, C. H., 1969, Atomic-absorption methods of analysis useful in geochemical exploration: U.S. Geol. Survey Bull. 1289, 45 p.

MAP SHOWING GEOCHEMICAL DISTRIBUTION AND ABUNDANCE OF BARIUM IN HEAVY MINERAL CONCENTRATES, TALKEETNA MOUNTAINS QUADRANGLE, ALASKA

by
R. J. Miller, G. C. Curtin, and Bela Csejtey, Jr.
1978

This report is preliminary and has not been edited or revised for conformity with Geological Survey standards and nomenclature.