

COMPREHENSIVE TABLES GIVING PHYSICAL DATA
AND THERMAL ENERGY ESTIMATES FOR YOUNG IGNEOUS
SYSTEMS OF THE UNITED STATES

by
R. L. Smith, H. R. Shaw, R. G. Luedke,
and S. L. Russell

U. S. Geological Survey
OPEN-FILE REPORT 78-925

This report is preliminary and has not
been edited or reviewed for conformity
with Geological Survey Standards and
nomenclature

INTRODUCTION

This report presents two tables. The first is a comprehensive table of 157 young igneous systems in the western United States, giving locations, physical data, and thermal energy estimates, where appropriate, for each system. The second table is a list of basaltic fields probably less than 10,000 years old in the western United States. These tables are updated and reformatted from Smith and Shaw's article "Igneous-related geothermal systems" in *Assessment of geothermal resources of the United States--1975* (USGS Circular 726, White and Williams, eds., 1975). This Open-File Report is a companion to Smith and Shaw's article "Igneous-related geothermal systems" in *Assessment of geothermal resources in the United States--1978* (USGS Circular 790, Muffler, ed., 1979). The article in Circular 790 contains an abridged table showing only those igneous systems for which thermal estimates were made. The article also gives an extensive discussion of hydrothermal cooling effects and an explanation of the model upon which the thermal energy estimates are based.

Thermal energy is calculated for those systems listed in table 1 that are thought to contribute significant thermal energy to the upper crust. As discussed by Smith and Shaw (1975), silicic volcanic systems are believed to be associated nearly always with high-level (<10 km) magma chambers. The thermal calculations in table 1 are made primarily for volcanic systems which show evidence from the presence of young silicic extrusions that a high-level magma chamber is being formed or has formed in the recent past.

Table 2 lists young basaltic lava fields that are probably less than 10,000 years old. The purpose of this listing is simply to call attention to these areas of very young basaltic eruptions where there may be very small thermal anomalies residual from the last magma injections in the upper crust. These areas and many older ones are shown by Luedke and Smith (1978a, 1978b, and in preparation) in a series of maps showing distribution, composition, and age of all late Cenozoic volcanic centers in the United States. Smith and Shaw (1975, p. 78) discussed young basic lava fields briefly and suggested that they should not be totally ignored for geothermal exploration. They do represent viable mantle sources for new magma and some fields may have small but significant thermal anomalies associated with hidden high-level silicic bodies. However, in general, exclusively basic volcanic systems rarely form thermal anomalies of economic interest because they rarely form high-level magma chambers that remain exclusively basic.

THERMAL ENERGY ESTIMATES

Three categories of thermal energy are given in table 1. ΔQ_{total} (column 10) is the thermal energy liberated if the entire magma chamber cools from an initial temperature of 850°C (the appropriate liquidus temperature for most silicic magmas) to a final temperature of 300°C (assumed ambient temperature) starting from a fixed time (in most cases, the age of the youngest silicic eruption). ΔQ_{now} (column 11) is the thermal energy which remains in the system at the present time within and around the original magma chamber. This energy constitutes the identified accessible resource base for igneous-related geothermal systems as defined in USGS Circular 790 and is summarized in table 3 of Smith and Shaw (1979). ΔQ_{out} (column 12) is the thermal energy transferred from the magma chamber to the roof rocks between the assumed time of emplacement of the intrusive body and the present. Three assumptions were used when making these thermal calculations: 1) a single pulse of magma is instantaneously emplaced and cools conductively from that time, 2) for most systems, the time of emplacement is taken as the age of the youngest silicic extrusion, and 3) no additional thermal energy is contributed by magmatic preheating or resupply.

Calculations of heat contents are approximate. The number of significant figures retained is determined from requirements of internal consistency among columns 10, 11, and 12 for systems so old that much of their heat has been lost at the surface on the basis of the model. For example, the possibility of a slight residual heat content is indicated in Column 11 for OR19 (Wart Peak caldera, Oregon) and is roughly proportioned equally between roof rocks and igneous pluton. Thus, roughly 8×10^{18} J are residual so that about 4×10^{18} J are left, respectively in the pluton and in the roof rocks and 356×10^{18} J have been given up by the pluton to the roof rocks and losses at the surface; that is 352×10^{18} J have been totally lost from the system to the surface.

In very large, older systems like ID6 (Rexburg Caldera), calculations for columns 11 and 12 are very crude, particularly because of the limitations of closed-system models. In this case, the entries in columns 11 and 12 simply represent stabs at the orders of magnitudes of the possible heat balances. In such cases, the estimates are probably conservative.

ENTRY CHANGES

Table 1 of this report has been revised from table 7 of Smith and Shaw (1975; USGS Circular 726). The location number designations have been changed to correspond with the zip code abbreviations for each state (for example, Alaska location A-1 has been changed to AK-1), except Hawaii which has remained as H (not HI). Seven systems have been deleted: Double (A-81), Black (A-82), Odell Butte (O-9), Black Butte (O-10), Cougar Mountain (O-15), Tushar Mountains (U-4), and Topaz Mountain (U-5). Thirteen systems have been added: Ukinrek Maars (AK-89), Hayes Volcano (AK-90), Inyo-Mammoth Fissure System (CA-18), Templeton Domes (CA-19), East Butte (ID-5), Rexburg Caldera (ID-6), Bearwallow Buttes (OR-18), Wart Peak Caldera (OR-19), Frederick Butte (OR-20), Thomas Range (UT-4), Wildcat Hills (UT-7), Clear Fork Dacite (WA-6), and Mann Butte (WA-7). It should be noted that Idaho and Wyoming have been tabulated separately. Yellowstone Caldera is now WY-1. Island Park-Huckleberry Ridge System (IW-1 in 1975) has been changed to Island Park System (ID-1). In table 1, Island Park has two chamber area figures: 3900 Ac - the area of the original caldera system (2 m.y. old) and 2100 Ao - the area of the western part of the system which is not overlapped by the younger (0.6 m.y. old) Yellowstone Caldera (WY-1).

The thermal estimates in this report are given in joules (instead of in calories as in Circular 726). A number of systems have significantly different thermal energies because of recalculations made with new age and size data. These systems include: Adagdak (AK-14), Kendrick Peak (AZ-3), Bill Williams Mountain (AZ-4), Melvin-Three Creeks Buttes (OR-7), Cappy-Burn Butte Area (OR-8), Mineral Mountains (UT-1), and Cove Creek Domes (UT-2).

The coordinates of the igneous-related systems have been revised to best approximate the center of the caldera or the vent distribution. In some cases other physical criteria had to be used.

CIRCULAR 790 MAPS

The young igneous systems in table 1 are plotted on maps 1 and 2 of U. S. Geological Survey Circular 790. Those systems for which an estimate of the thermal energy still remaining in the ground (ΔQ_{now} ; Column 11) can be made are shown on the maps by nested green triangles. For each system the number of triangles indicates the range of values in which the thermal estimate falls. Igneous systems in table 1 for which no estimates of thermal energy are made are symbolized on the maps by

green snowflakes. The young basaltic fields of table 2 are also plotted on maps 1 and 2 of Circular 790, as brown shaded areas. The identifying numbers and letters in column 1 of tables 1 and 2 refer to the individual systems plotted on the maps. Longitude in column 3 is west unless otherwise noted (AK1 to AK7). Volcanic systems marked by asterisks in column 1 are known to have some associated hydrothermal activity (see Brook and others, 1979).

INPUT DATA

The input data from which all thermal estimates are made are shown in columns 4-9 of table 1. The physical basis of specific numbers is indicated by symbols which are explained below. The composition of the last eruption (for example, silicic or basic) and age data are listed in columns 4 and 5, respectively. Area of the magma chamber (column 6) is based on various surface manifestations of volcanism, geologic structure, or geophysics. The volume range of the chamber (column 7) is calculated by assuming the thickness of the magma chamber ranges from 2.5 to 10 km. This range is reduced to a single "best estimate" (column 8) to simplify the thermal calculations. Some volumes are derived by ten-fold extrapolation of ejecta volumes (Smith and Shaw, 1973, and unpublished). Column 9 indicates our best estimate (based on Smith and Shaw, 1979, fig. 3) as to the present thermal state of a magma chamber--that is, whether or not magma exists in the system now. Many entries are shown as greater or less than 650°C, which is the approximate minimum temperature of solidification of granitic melts. For those systems whose age and size data are incomplete, no thermal energy estimates are made.

EXPLANATION OF SYMBOLS IN TABLE 1

AGE = T

Ty - Last eruption
 Tys - Youngest silicic eruption
 Tyb - Youngest basic eruption
 Ts - Age (silicic)
 Tc - Age caldera eruption
 Tg - Greatest known age (composition unspecified)
 Tgs - Greatest age (silicic)
 Tgb - Greatest age (basic)
 Tb - Age (basic)

AREA = A

- Ac - From caldera
- Av - From vent distribution
- As - From shadow
- Af - From fractures
- Au - From uplift
- Ag - From geophysical anomaly (unspecified)
 - Agg - Gravity
 - Agm - Magnetic
 - Ags - Seismic
 - Ago - Other, see remarks
- Ao - Other, see remarks

VOLUME = V

- Vc - From caldera
- Vv - From vent distribution
- Vs - From shadow
- Vf - From fractures
- Vu - From uplift
- Vg - From geophysical anomaly
 - Vgg - Gravity
 - Vgm - Magnetic
 - Vgs - Seismic
 - Vgo - Other, see remarks
- Vo - Other, see remarks
- Vee - From extrapolation of silicic ejecta volume
- Vb - Best estimate

METHODS OF CALCULATION OF THERMAL ENERGY

COLUMN 10: ΔQ_{total} , in units of 10^{18} joules

Assumptions:

- Initial temperature = 850°C
- Latent heat of crystallization = 272 J/g
- Heat capacity = $1.3 \text{ J/g/}^{\circ}\text{C}$
- Mean density of magma = 2.5 g/cm^3

The above values are approximate averages for the composition and temperature ranges of table 1. From these values the heat liberated by crystallization and conductive cooling between 850°C and 650°C is $(850^{\circ}\text{C} - 650^{\circ}\text{C})(1.3 \text{ J/g/}^{\circ}\text{C}) + 271 \text{ J/g} = 523 \text{ J/g}$. The total heat liberated in the same manner between 850°C and 300°C is 963 J/g .

One cubic kilometer of magma represents 2.5×10^{15} g. The total heat liberated (between 850°C and 300°C) per cubic kilometer is $(2.5 \times 10^{15} \text{ g/km}^3)(963 \text{ J/g}) = 2.41 \times 10^{18}$ joules. This number multiplied by the volume V_b in column 8 gives the ΔQ_{total} of column 10.

COLUMN 11, ΔQ_{now} , in units of 10^{18} joules

The time required for a change of the original gradient at the Earth's surface to a steady-state gradient between the surface temperature and the magma chamber temperature is given approximately by relations discussed by Jaeger (1964). For the assumed depth of cover of 4 km and a thermal diffusivity of $0.007 \text{ cm}^2/\text{sec}$, this time is about 360,000 years. Where T_y (age of last eruption) is much younger than this time, the total heat remaining in the system now (column 11) is assumed to be about the same as the total value in column 10. Estimates of losses for older systems require detailed calculations of the disturbance of the geothermal gradient.

The value of thermal diffusivity used is an average estimate for crustal rocks. Roof rocks above large caldera systems such as Yellowstone, Wyoming (WY-1), Valles, New Mexico (NM-1) and Long Valley, California (CA-3) may have smaller values of conductive thermal diffusivity. Hydrothermal convection systems, however, can increase the effective value of thermal diffusivity by a significant amount, depending on average permeabilities of roof rocks.

COLUMN 12, ΔQ_{out} , in units of 10^{18} joules

The total amount of heat transfer per square centimeter from a magma chamber into roof rocks is given by Carslaw and Jaeger (1959, p. 61) and also is discussed by Shaw (1974). Using these relations the total heat transfer (ΔQ_{out}) in column 12 is given by:

$$\Delta Q_{\text{out}} = 216At^{\frac{1}{2}}$$

where A is contact area (from column 6 converted to square centimeters) and t is the time in seconds since T_y . Calculations in column 12 are approximately valid only if the time of solidification is greater than T_y in column 5. The time of solidification is approximated by lines 3 and 4 in figure 3 of Smith and Shaw (1979).

If T_y is much greater than 360,000 years and the time for solidification, the calculation of heat content is ambiguous because of the increasing importance of hydrothermal losses. On the basis of conduction models, however, the total time for decay of igneous-related thermal anomalies may be very long. As an example, the time required for the central temperature in a magma chamber of horizontal slab-like geometry to decay from

the initial magma temperature to nearly ambient temperature is about 2 m.y. for a magma chamber 5 km thick and about 10 m.y. for a chamber 10 km thick. Even a liberal allowance for hydrothermal losses means that the igneous-related thermal anomalies for the largest systems of table 1 probably are preserved for times of the order 10 m.y. or longer.

Queries in columns 10-12 mean that even though data exist, we are not confident that they pertain even approximately to the assumptions of the calculations. Blank spaces in the table mean that more geological and geochronological study is needed before we are willing to make estimates.

REFERENCES CITED

- Brook, C. A., Mariner, R. H., Mabey, D. R., Swanson, J. R., Guffanti, Marianne, and Muffler, L. J. P., 1979, Hydrothermal convection systems with reservoir temperatures $>90^{\circ}\text{C}$, in Muffler, L. J. P., ed., Assessment of geothermal resources of the United States--1978: U. S. Geological Survey Circular 790, p. 18-85.
- Carslaw, H. S., and Jaeger, J. C., 1959, Conduction of heat in solids: Oxford, Clarendon Press, 510 p.
- Jaeger, J. C., 1964, Thermal effects of intrusions: Review of Geophysics v. 2, p. 443-466.
- Luedke, R. G., and Smith, R. L., 1978, Map showing distribution, composition, and age of late Cenozoic volcanic centers in Arizona and New Mexico: U. S. Geological Survey Miscellaneous Investigations Map I-1091A.
- _____, 1978, Map showing distribution, composition, and age of late Cenozoic volcanic centers in Colorado and Utah: U. S. Geol. Survey Miscellaneous Investigations Map I-1091B.
- _____, U. S. Geological Survey Miscellaneous Investigations Maps I-1091C, D, E, and F. (in preparation)
- Muffler, L. J. P., 1979, ed., Assessment of geothermal resources of the United States--1978: U. S. Geological Survey Circular 790.

- Shaw, H. R., 1974, Diffusion of H₂O in granitic liquids: Part I Experimental data; Part II Mass transfer in magma chambers, in Hofmann, A. W., Giletti, B. J., Yoder, H. S., and Yu, R. A., eds., Geochemical transport and kinetics: Carnegie Inst. Washington Pub. 634, p. 139-170.
- Smith, R. L., and Shaw, H. R., 1973, Volcanic rocks as geologic guides to geothermal exploration and evaluation: EOS, American Geophysical Union Transactions, v. 54, p. 1213.
- _____, 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.
- _____, Igneous-related geothermal systems, in Muffler, L. J. P., Assessment of geothermal resources of the United States--1978: U. S. Geological Survey Circular 790, p.12-17.
- White, D. E., and Williams, D. L., eds., 1975, Assessment of geothermal resources of the United States--1975: U. S. Geological Survey Circular 726, 155 p.

REFERENCES FOR TABLE 1

ALASKA:

- 1AK Brew, D. A., Muffler, L. J. P., and Loney, R. A., 1969, Reconnaissance geology of the Mount Edgecumbe volcanic field, Kruzof Island, southeastern Alaska: U. S. Geological Survey Professional Paper 650-D, p. D1-D18.
- 2AK Byers, F. M., Jr., 1959, Geology of Umnak and Bogoslof Islands, Aleutian Islands: U. S. Geological Survey Bulletin 1028-L, p. 297-361.
- 3AK Byers, F. M., Jr., and Barth, T. F., 1953, Volcanic activity on Akun and Akutan Islands, Alaska: Pacific Science Congress, 7th, New Zealand, 1949, Proceedings, v. 2, p. 382-397.

- 4AK Coats, R. R., 1950, Volcanic activity in the Aleutian arc: U. S. Geological Survey Bulletin 974-B, p. 35-49.
- 5AK Drewes, H., Fraser, G., Snyder, G., and Barnett, H., Jr., 1961, Geology of Unalaska and the adjacent insular shelf, Aleutian Islands, Alaska: U. S. Geological Survey Bulletin 1028-S, p. 583-676.
- 6AK Kennedy, G. C., and Waldron, H. H., 1955, Geology of Pavlof volcano and vicinity, Alaska: U. S. Geological Survey Bulletin 1028-A, p. 11-13.
- 7AK Lerbekmo, J. F. and Campbell, F. A., 1969, Distribution, composition, and source of the White River Ash, Yukon Territory: Canadian Journal of Geologic Science, v. 6, p. 109-116.
- 8AK Marsh, B. D., 1976, Some Aleutian andesites: their nature and source: Journal of Geology, v. 84, p. 27-45.
- 9AK Miller, T. P. and Smith, R. L., 1978, personal communication.
- 10AK Miller, T. P., Smith, R. L., Richter, D. H., Lanphere, M. A., and Dalrymple, G. B., (in press), Volcanic history and geothermal potential of Mount Drum volcano, Alaska.
- 11AK Richter, D. H., Smith, R. L., Yehle, L. A., and Miller, T. P., (in press), Geologic map of the Gulkana A-2 quadrangle, Alaska: U. S. Geological Survey Quadrangle Map GQ 1520.
- 12AK Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. H., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.
- 13AK Smith, R. L. and Soule, C. E., 1973, Post-Miocene volcanoes of the world-sheet N II, region: western Alaska and Bering Sea islands, in I.A.V.C.E.I., ed., Data sheets of the post-Miocene volcanoes of the world with index maps: I.A.V.C.E.I. Publication Office, Rome, Italy, unpaginated.

14AK Snyder, G. L., 1954, Geology of Little Sitkin Island, Alaska: U. S. Geological Survey Bulletin 1028-H, p. 177-184.

15AK Waldron, H., 1961, Geologic reconnaissance of Frosty Peak and vicinity, Alaska: U. S. Geological Survey Bulletin 1028-T, p. 677-708.

ARIZONA:

1AZ McKee, E. H., Wolf, E., and Ulrich, G. E., 1978, oral communication.

2AZ Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

CALIFORNIA:

1CA Bacon, C. R., 1978, oral communication.

2CA Bailey, R.A., 1978, personal communication.

3CA Bailey, R. A., Dalrymple, G. B., and Lanphere, M. A., 1976, Volcanism, structure, and geochronology of Long Valley Caldera, Mono County, California: Journal of Geophysical Research, v. 81, no. 5, p. 725-744.

4CA Bateman, P. C., 1965, Geology and tungsten mineralization of the Bishop district, California: U. S. Geological Survey Professional Paper 470, 58 p.

5CA Christiansen, R. L., 1978, oral communication.

6CA Hearn, B. C., Jr., Donnelly, J. M., and Goff, F. E., 1975, Duffield, W. A. and Bacon, C. R., 1977, Preliminary geologic map of the Coso volcanic field and adjacent areas, Inyo County, California, with a table of new K/Ar dates by G. B. Dalrymple: U. S. Geological Survey Open-File Report 77-311.

- 8CA Griscom, A. and Muffler, L. J. P., 1971, Aeromagnetic map and interpretation of the Salton Sea Geothermal Area, California: U. S. Geological Survey Geophysical Investigation Map GP-754.
- 9CA Hearn, B. C., Jr., Donnelly, J. M., and Goff, F. E., (in press) The Clear Lake Volcanics, California: tectonic setting and magmatic sources: U. S. Geological Survey Professional Paper.
- 10CA Isherwood, W. F., 1975, Gravity and magnetic studies of The Geysers-Clear Lake Geothermal Region, California, USA: in Proceedings, Second United Nations Symposium on the development and use of geothermal resources, v. 2, p. 1065-1073.
- 11CA Lachenbruch, A. H., Sass, J. K., Monroe, R. J., and Moses, T. H., Jr., 1975, Geothermal setting and simple heat conduction models for the Long Valley Caldera: Journal of Geophysical Research, v. 81, no. 5, p. 769-784
- 12CA Lanphere, M. A., Dalrymple, G. B., and Smith, R. L., 1975, K-Ar ages of Pleistocene rhyolitic volcanism in the Coso Range, California: Geology, v.3, no. 6, p. 339-341.
- 13CA McKee, E. H., 1974, oral communication.
- 14CA Robinson, P. T., Elders, W. A., and Muffler, L. J. P., 1976, Quaternary volcanism in the Salton Sea Geothermal Field, Imperial Valley, California: Geological Society of America Bulletin, v. 87, no. 3, p. 347-360.
- 15CA Silberman, M. L., Chesterman, C. W., Kleinhampl, F. J., and Gray, C. H., Jr., 1972, K-Ar ages of volcanic rocks and gold-bearing quartz-adularia veins in the Bodie Mining District, Mono County, California: Economic Geology, v. 67, p. 597-604.
- 16CA Smith, G. I., 1964, Geology and volcanic petrology of the Lava Mountains, San Bernardino County, California: U. S. Geological Survey Professional Paper 457, 97p.

17CA Smith, R. L., and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

18CA Williams, Howel and Curtis, G. H., 1977, The Sutter Buttes of California: a study of Plio-Pleistocene volcanism: University of California Publications in Geological Sciences, v. 116, 56 p.

HAWAII:

1HI Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

IDAHO:

1ID Armstrong, R. L., Leeman, W. P., and Malde, H. E., 1975, K-Ar dating, Quaternary and Neogene volcanic rocks of the Snake River Plain, Idaho: American Journal of Science, v. 275, p. 225-251.

2ID Christiansen, R. L., 1975, and 1978, oral communication.

3ID Kuntz, M. A., 1978, oral communication.

4ID Mabey, D. R., 1978, Gravity and aeromagnetic anomalies in the Rexburg area of eastern Idaho: U. S. Geological Survey Open-File Report 78-382.

5ID Prostka, H. and Embree, G., 1978, oral communication.

6ID Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

NEVADA:

- 1NV Robinson, P. T., McKee, E. H., and Moiola, R. J., 1968, Cenozoic volcanism and sedimentation, Silver Peak region western Nevada and adjacent California, in Coats, R. R., Hay, R. L., and Anderson, C. A., Studies in volcanology-a memoir in honor of Howel Williams: Geological Society of America Memoir 116, p. 577-612.
- 2NV Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources in the United States-1975, U. S. Geological Survey Circular 726, p. 58-83.
- 3NV White, D. E., 1975, oral communication.

NEW MEXICO:

- 1NM Doell, Richard, R., Dalrymple, G. B., Smith, R. L., and Bailey, R. A., 1969, Paleomagnetism, potassium-argon ages, and geology of rhyolites and associated rocks of the Valles Caldera, New Mexico, in Coats, R. R., Hay, R. L., and Anderson, C. A., eds., Studies in volcanology-a memoir in honor of Howel Williams: Geological Society of America Memoir 116, p. 211-248.
- 2NM Lipman, P. W., 1978, oral communication.
- 3NM Smith, R. L. and Bailey, R. A., 1968, Resurgent cauldrons, in Coats, R. R., Hay, R. L., and Anderson, C. A., eds., Studies in volcanology-a memoir in honor of Howel Williams: Geological Society of America Memoir 116, p. 613-662.
- 4NM Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

OREGON:

- 1OR Armstrong, R. L., Taylor, E. M., Hales, P. O., and Parker, D. J., 1975, K-Ar dates for volcanic rocks, central Cascade Range of Oregon: Isochron West, no. 13, p. 5-10.
- 2OR MacLeod, N. S., 1978, oral communication.
- 3OR MacLeod, N. S., Walker, G. W., and McKee, E. H., 1967, Geothermal significance of eastward increase in age of Upper Cenozoic rhyolitic domes in southeastern Oregon, in Proceedings 2nd U.N. Symposium on the development and use of geothermal resources, v. 1, p. 465-474.
- 4OR Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.
- 5OR Walker, G. W., Dalrymple, G. B., and Lanphere, M. A., 1974, Index to potassium-argon ages of Cenozoic volcanic rocks of Oregon: U. S. Geological Survey Miscellaneous Field Studies Map MF-569.
- 6OR Walker, G. W., Greene, R. C., and Pattee, E. C., 1966, Mineral resources of the Mount Jefferson primitive areas, Oregon: U. S. Geological Survey Bulletin 1230-D, p. D1-D32.
- 7OR Wise, W. S., 1969, Geology and petrology of the Mt. Hood area--a study of High Cascade volcanism: Geological Society of America Bulletin, v. 80, no. 6, p. 969-1006.

UTAH:

- 1UT Armstrong, R. L., 1970, Geochronology of Tertiary igneous rocks, eastern Basin and Range Province, western Utah, eastern Nevada, and vicinity, U.S.A.: Geochimica et Cosmochimica Acta, v. 34, no. 2, p. 203-232.

2UT Lipman, P. W., Rowley, P. D., Mehnert, H. H., Evans, S. H., Jr., Nash, W. P., and Brown, F. H., 1978, Pleistocene rhyolite of the Mineral Mountains, Utah-geothermal and archeological significance: Journal of Research of the U. S. Geological Survey, v. 6, p. 133-147.

3UT McKee, E. H., 1977, oral communication.

4UT Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

WASHINGTON:

1WA Ellingson, J. A., 1972, The rocks and structure of the White Pass area, Washington: Northwestern Science, v. 46, no. 1, p. 9-24.

2WA Hammond, P. E., Bentley, R. D., Brown, J. C., Ellingson, J. A., and Swanson, D. A., 1977, Field trip 4: Volcanic stratigraphy and structure of the southern Cascade Range, Washington, in Geological excursions in the Pacific Northwest: Geological Society of America 1977 Annual Meeting, Seattle, p. 127-169.

3WA Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

WYOMING

1WY Christiansen, R. L., 1975, Oral communication.

6WY Smith, R. L. and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E. and Williams, D. L., eds., Assessment of geothermal resources of the United States-1975: U. S. Geological Survey Circular 726, p. 58-83.

Table 2.--Basic volcanic fields probably less than 10,000 years old

	<u>Longitude</u>	<u>Latitude</u>
ALASKA		
AK1 Devil Mountain Field -----	66°18'	164°31'
AK2 Imuruk Lake field -----	65°29'	163°17'
AK3 St. Lawrence Island -----	63°37'	170°17'
AK4 St. Michaels field -----	63°36'	162°37'
AK5 Ingrichuak Hill area -----	62°11'	164°06'
AK6 Ingakslugwat Hills area -----	61°22'	163°58'
AK7 Nunivak Island (?) -----	60°01'	166°20'
AK8 Pribilof (St. Paul) Islands(?) -----	57°10'	170°23'
ARIZONA		
AZ1 Unikaret flow -----	36°22'	113°09'
AZ2 Sunset Crater flow -----	35°22'	111°30'
CALIFORNIA		
CA1 Copco Lake area -----	41°59'	122°20'
CA2 Goosenest area -----	41°43'	122°13'
Mt. Lassen - Mt. Shasta area		
CA3a Callahan flows -----	41°41'	121°36'
CA3b Burnt Lava flows -----	41°31'	121°32'
CA3c Paint Pot Crater flow -----	41°33'	121°42'
CA3d Six Shooter Butte flows -----	41°31'	121°37'
CA3e Fall River Mills flow -----	40°57'	121°22'
CA3f Hat Creek flow -----	40°39'	121°26'
CA3g Cinder Cone flow "1851" -----	40°33'	121°19'
CA4 Ubehebe Craters area -----	37°01'	117°27'
CA5 Cima Lava field -----	35°11'	115°49'
CA6 Pisgah field -----	34°45'	116°23'
CA7 Amboy field -----	34°33'	115°47'
COLORADO		
CO1 Dotsero -----	39°40'	107°02'

Table 2.--Basic volcanic fields probably less than 10,000 years old--continued

		<u>Longitude</u>	<u>Latitude</u>
HAWAII			
Hawaii Island			
H 1	Hualalei -----	19°42'	155°52'
H 2	Mauna Kea -----	19°50'	155°29'
H 3	Mauna Loa -----	19°29'	155°36'
H 4	Kilauea -----	19°25'	155°17'
Maui Island			
H 5	Haleakala -----	20°43'	156°13'
IDAHO			
ID1	Craters of the Moon area -----	43°25'	113°32'
ID2	North Robbers flow -----	43°23'	112°59'
ID3	Cerro Grande -----	43°22'	112°53'
ID4	Hells Half Acre -----	43°30'	112°27'
ID5	Wapi -----	42°53'	113°13'
ID6	Kings Bowl -----	42°57'	113°13'
NEVADA			
NV1	Lunar Crater lava field -----	38°29'	115°59'
NEW MEXICO			
NM1	Capulin flow -----	36°47'	103°58'
NM2	McCartys flow -----	34°49'	108°00'
NM3	Carrizozo flow -----	33°47'	105°56'

Table 2.--Basic volcanic fields probably less than 10,000 years old--continued

	<u>Longitude</u>	<u>Latitude</u>
OREGON		
OR1 North Cinder Peak flow -----	44°36'	121°47'
OR2 Nash Crater flow -----	44°25'	121°57'
OR3 Sand Mountain flow -----	44°23'	121°56'
OR4 Belknap lava field -----	44°17'	121°51'
OR5 North Sister lava field -----	44°12'	121°47'
OR6 Le Conte Crater flow -----	44°03'	121°48'
OR7 Cayuse Crater flow -----	44°04'	121°42'
OR8 Bachelor Butte lava field -----	43°59'	121°41'
OR9 Lava Butte flow -----	43°55'	121°21'
OR10 Newberry Crater area (lower flanks)--	43°50'	121°17'
OR11 Wukxi Butte area -----	43°46'	121°45'
OR12 Pine Butte area -----	43°40'	121°51'
OR13 Black Rock Butte area -----	43°29'	121°48'
OR14 Devils Garden area -----	43°30'	120°54'
OR15 Squaw Ridge field -----	43°28'	120°45'
OR16 Four Craters lava field -----	43°22'	120°40'
OR17 Brown Mountain area -----	42°22'	122°17'
OR18 Diamond Craters area -----	43°06'	118°45'
OR19 Jordan Craters field -----	43°02'	117°25'
OR20 Jackies Butte field -----	42°36'	117°35'
UTAH		
UT1 Ice Springs field -----	38°58'	112°30'
UT5 Cove Fort flow -----	38°34'	112°39'
UT2 Markagunt field -----	37°34'	112°43'
UT4 Santa Clara flow -----	37°15'	113°38'
UT3 Crater Hill Flow -----	37°13'	113°06'
WASHINGTON		
WA1 Red Mountain-Big Lava Bed -----	45°55'	121°45'

Table 1.--Magnitudes and heat contents of identified volcanic systems

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume V ₀ (km ³)	Solidification state (°C)	ΔQ total (10 ¹⁸ J)	ΔQ now (10 ¹⁸ J)	ΔQ out (10 ¹⁸ J)	Remarks	References
AK1	Buldir	52 21 175 55 N	Basio	<10 ⁴ ?								>10 km depth	4AX
AK2	Kiska	52 06 177 36 N	Basio	Active								>10 km depth	
AK3	Segula	52 01 178 08 N	Basio Silicic ¹⁰	<10 ⁴ ?								Need data on composition and age	
AK4	Davidof	51 58 178 20 E	No data	<10 ⁴ ?	5 Ao	12.5-50 Vc	12.5	>650?	20	20		Need data on composition and age	
AK5	Little Sittin	51 57 178 32 N	Basio	Active	17.3 Ao	45-180 Vc	75	>850	180	180			10AX
AK6	Semlappochnoi (Cerberus)	51 56 179 36 E	Basio	Active	42.4 Ao	106-424 Vc	150	>950	360	360			4AX
AK7	Sugarloaf	51 53 179 38 N	Basio	<10 ⁴ ?								>10 km depth	4AX
AK8	Garelol	51 48 178 48	Basio	Active								>10 km depth	4AX
AK9	Tanaga	51 53 178 89	Basio?	Active	85.9 Ao	215-660 Vc	480	>950	960	960		Need more data	4AX
AK10	Takawengha	51 52 178 00	Basio?	<10 ⁴ ?	0.9 Ao	22.5-90 Vc	>22.5	>650	54	54		Need data on composition and age	4AX
AK11	Sorof	51 54 177 26	Basio?	<10 ⁴ ?								>10 km depth	4AX
AK12	Kanaga	51 56 177 09	Basio?	Active	23.0 Ao	57.5-230 Vc	75	>850	180	180			4AX
AK13	Noroff	51 56 176 44	Basio	0.14 Tys								>10 km depth?	4AX, 9AX
AK14	Adagdak	51 59 176 35	Basio? Silicic ¹⁰	3.4 x 10 ⁵ Tys 1.4 x 10 ⁵ Tys		25 Vcc	25	<650	59	50	9	Probable low temperature hydrothermal system	4AX, 9AX
AK15	Great Sittin	52 04 176 00	Basio	Active	1.8 Ao	4.5-18 Vc	>5	>850	>13	>13			4AX
AK16	Kasatochi	52 11 175 30	Basio?	Active?								>10 km depth?	4AX
AK17	Koniuj	52 13 175 00	Basio?	Active?								>10 km depth?	4AX

Table 1.--Magnitudes and heat contents of identified volcanic systems--Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume Vo (km ³)	Solidifi- cation state (%)	ΔQ total (10 ¹⁸ J)	ΔQ now (10 ¹⁸ J)	ΔQ out (10 ¹⁸ J)	Remarks	References
A L A S K A													
AK18	Bergief	52 19 174 23	No data	No data								Need data on composition and age	6AX
AK19	Korovin	52 23 174 09	Basic	Active								>10 km depth	6AX
AK20	Ellichaf	52 20 174 09	No data	No data	28.6 Ao	70-280 Vo	>100		240			Need data on composition and age. A18 to A21 may be one system	6AX
AK21	Satichaf	52 19 174 01	Basic?	Active?								>10 km depth	6AX
AK22	Segum	52 19 172 29	No data	Active	20 Ao	100-400 Vo	200	>550	480	480		Appears to be a double caldera. Not reported. Needs investigation.	
AK23	Aukta	52 30 171 15	Basic?	Active								>10 km depth? Need data.	
AK24	Chagulat	52 34 171 08	Basic?	<10 ⁴ ?								>10 km depth? Need data.	
AK25	Yunaska	52 36 170 42	No data Basic?	Active	12.1 Ao	30-120 Vo	40	>550	96	96		Probably two volcanoes. Need data.	
AK26	Berbert	52 45 170 07	Basic?	<10 ⁴ ?								>10 km depth? Need data	
AK27	Carlisle	52 54 170 03	Basic?	Active								>10 km depth? Need data.	
AK28	Cleveland	52 49 169 57	Basic?	Active								>10 km depth? Need data.	
AK29	Diliaga	53 04 169 46	Basic?	<10 ⁴ ?								>10 km depth? Need data.	
AK30	Tuna	52 50 169 46	Basic?	No data								>10 km depth? Need data.	
AK31	Kagamil	52 59 169 43	Basic?	Active								>10 km depth? Need data	
AK32	Vmavidof	53 08 168 41	Sillicic	Active								Need more data. Probable high level chamber.	2AX
AK33	Rechaschnoi	53 09 168 33	Basic? Sillicic?	<10 ⁴								Need more data. Probable high level chamber.	2AX
AK34	Olmok	53 25 168 08	Basic	Active 8 x 10 ³ Vo	62.9 Ao	155-428 Vo	250	>450	603	603			2AX

Table 1.—Magnitudes and heat contents of identified volcanic systems—Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition, last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume Vo (km ³)	Solidifi- cation state (°C)	ΔQ total (10 ¹⁸ J)	ΔQ now (10 ¹⁸ J)	ΔQ out (10 ¹⁸ J)	Remarks	References
AK35	Tulik	53 23 168 03	Basic	No data								>10 km depth	2AK
AK36	Bogosluf	53 56 168 02	Basic	Active								>10 km depth	2AK
AK37	Makushin	53 54 168 56	Silicic?	Active	3.6 Mo	9-36 Vo	>10	>850	25	25			5AK, 9AK
AK38	Table Top	53 50 168 40	Basic	No data								>10 km depth	5AK, 9AK
AK39	Antaa	54 06 168 00	Basic?	Active	3.5 Mo	9-36 Vo	>10	>850	25	25			3AK, 9AK
AK40	Mt. Gilbert (Akum)	54 15 165 39	Basic?	No data								>10 km depth?	3AK, 9AK
AK41	Pogromi	54 34 164 42	Basic	Active?								>10 km depth. Satellite to Westdahl. Possibly active.	4AK, 9AK
AK42	Westdahl	54 31 164 39	No data Basic?	Active								Need more data. >10 km?	4AK, 9AK
AK43	Fisher	54 40 164 21	Basic	Active?	122.6 Mo	300-1200 Vo	600	>850	1440	1440			4AK, 9AK
AK44	Shishaldin	54 45 163 50	Basic	Active								>10 km depth	4AK, 9AK
AK45	Tanotaki	54 45 163 44	Basic?	Active								>10 km depth?	4AK, 9AK
AK46	Roundtop	54 48 163 36	No data	Active?									4AK, 9AK
AK47	Aash	55 25 163 09	Basic?	<10 ⁴								>10 km depth?	4AK, 9AK
AK48	Frosty	55 04 162 49	Silicic? Basic?	No data								Need data on composition and age	9AK, 15AK
AK49	Valrus (Morzhovoi)	55 01 162 50	Basic	No data >10 ⁴								>10 km depth	9AK, 15AK
AK50	Dutton	55 11 162 16	Basic	No data								>10 km depth	9AK
AK51	Ramona	55 20 162 04	Basic	Active	117.3 Mo	300-1200 Vo	600	>850	1440	1440		In Ramona caldera	6AK, 9AK

Table 1.--Magnitudes and heat contents of identified volcanic systems--Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume Vo (km ³)	Solidifi- cation state (°C)	AQ total (10 ¹⁸ J)	AQ now (10 ¹⁸ J)	AQ out (10 ¹⁸ J)	Remarks	References
AK52	Bague	55 22 161 58	Basic	Active								In Emons caldera	9AK
AK53	Double Crater	55 23 161 57	Basic	Active?								In Emons caldera	9AK
AK54	Pavlof	55 25 161 56	Basic	Active								>10 km depth	9AK
AK55	Pavlof Sister	55 27 161 52	Basic	Active								>10 km depth	9AK
AK56	Dana	55 37 161 13	Silicic	No data <10 ⁴	1.6 Mo	4-16 Vo	>5		13			Need age data	9AK
AK57	Kupreanof	56 01 159 48	No data Basic?	Active								Need more data. Probably two volcanoes	9AK
AK58	Ventaminof	56 10 159 23	Basic	Active 3.7 x 10 ³ Yo	50.4 Mo	125-500 Vo	200	>850	481	481			9AK
AK59	Black (Purple)	56 34 158 48	Silicic	<10 ⁴	6.9 Mo	17.5-70 Vo	>20	>650	50	50		Need age data	9AK
AK60	Aniakchak	56 53 158 09	Silicic	Active 3.6 x 10 ³ Yo	55.6 Mo	140-560 Vo	225	>650	540	540			9AK
AK61	Chignik	57 08 157 00	Basic?	Active								Need more data	9AK
AK62	Kialagvik	57 12 156 42	Silicic	No Data								Need age data	9AK
AK63	Peulik (Ugashik caldera)	57 45 156 21	Basic? Silicic	Active? >1.7 x 10 ⁵ Yo	10.5 Mo	25-180 Vo	>30	>650	71	71		Silicic in focus of both caldera and Mt. Peulik; perhaps young basic eruption on north flank of Peulik. Need age data.	9AK
AK64	Martin	58 09 155 23	Basic?	Active								>10 km depth?	
AK65	Magick	58 12 155 15	?	Active								Silicic domes? Need more data.	
AK66	Movarupea	58 16 155 10	Silicic	Active	8.1 Mo	20-80 Vo	50	>650	128	128			
AK67	Mt. Griggs (Knife Peak)	58 21 155 07	Basic?	Active								>10 km depth?	
AK68	Trident	58 14 155 07	Basic	Active								>10 km depth?	

Table 1.--Magnitudes and heat contents of identified volcanic systems--Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition, last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume (km ³)	Solidification state (°C)	AQ total (10 ¹⁰ J)	AQ (10 ¹⁰ J)	AQ (10 ¹⁰ J)	Remarks	References
AK69	Atlatzi	58 16 154 59	Basalt	Active	0.1 Mo	20-40 Vv	>28	>850	50	50			
AK70	Snory	58 20 154 44	No data	No data								No data	
AK71	Denison	58 25 154 27	No data	No data								No data	
AK72	Stoller	58 26 154 24	No data	No data								No data	
AK73	Eukak	58 20 154 21	No data	Active?								No data	
AK74	Devila Desk	58 29 154 18	No data	No data								No data	
AK75	Eguyak	58 37 154 05	Silicic	<10 ⁴	4.1 Mo	10-40 Vv	>15	>850	30	30	13	Need age data	9AK
AK76	Fourpeaked	58 46 153 41	No data	<10 ⁴								No data	
AK77	Douglas	58 52 153 33	No data	Active?								No data	
AK78	Augustine	59 22 153 25	Silicic-Basalt	Active								Need geophysical data?	9AK
AK79	Iliamna	60 02 153 05	Basalt	Active								>10 km depth	
AK80	Redoubt	60 29 152 45	Basalt	Active								>10 km depth	
AK83	Spurr	61 10 152 15	Basalt	Active								Caldera? Need more data.	9AK
AK84	Drum	62 07 144 38	Silicic	2.4 x 10 ⁵ Yr 2.4 x 10 ⁵ Yr 8 x 10 ⁵ Yr	148 Av	350-1400 Vv	400	650	960	>840	>420	A more liberal volume and thermal estimate is given by Miller and others, see reference.	10AK, 11AK
AK85	Sanford	62 13 144 07	No data	<3.2 x 10 ⁵ Yr								Need data on parasitic vents high on flanks of Sanford (inaccessible?)	11AK
AK86	Wrangell	62 00 144 01	Basalt?	Active >1.7 x 10 ⁵ Yr	15 Mo	37.5-158 Vv	50	>850	120	120			11AK
AK87	White River	61 27 141 28	Silicic	1.5 x 10 ³		00 Vv	80	>650	190	190			11AK

Table 1.—Magnitudes and heat contents of identified volcanic systems—Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition last eruption	Age data (Yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume (km ³)	Solidification rate (°C)	AO total (10 ¹⁸ J)	AO low cut (10 ¹⁸ J)	AO cut (10 ¹⁸ J)	Remarks	References
AE88	Edgecumbe	57 01 135 46	Basalt Silicic	Active? <8 x 10 ³ 9 x 10 ³ Yr	74 Av	105-740 Vv	250	>650	602	602	250	Silicic in focus. Basic on flanks.	1A2
AE89	Ukinrek Maars	57 50 156 29	Basic	Active								Two <u>new</u> volcano vents 1977. >10 km depth?	9A2
AE90	Mayes volcano	61 37 132 27	Silicic	<3.7 x 10 ³								Probably potentially active vent area, probably ice capped.	9A2
AE1	San Francisco Mountains	35 20 111 40	Basalt Rhyolite	9 x 10 ² Yrb 2 x 10 ⁵ Yrb 2.7 x 10 ⁶ Yrb	250 Av	625-2500 Vv	1250	>650	3010	3010	1320	Shadow area and volume so derived highly speculative. Migrating system.	1A2, 2A2
AE2	Kendrick Peak	35 24 111 52	Silicic	1.5 x 10 ⁶ Yrb 1.9 x 10 ⁶ Yrb <1.9 x 10 ⁶ Yrb	50 Av	125-500 Vv	250	>650	603	150	519	Tys on Slate Mountain	1A2, 2A2
AE3	Sitgreaves Peak	35 21 112 00	Rhyolite	1.9 x 10 ⁶ Yrb 2.0 x 10 ⁶ Yrb >1.9 x 10 ⁶ Yrb	50 Av	125-500 Vv	200	>650	401	46	456		1A2, 2A2
AE4	Bill Williams Mountain	35 12 112 12	Silicic	3.5 x 10 ⁶ Yrb 4.1 x 10 ⁶ Yrb	20 Av	50-200 Vv	100	>650	240	4	240		1A2, 2A2
CA1	Lassen Peak	40 29 121 30	Rhyodacite	61 Yrs Yrb	80 Av	200-800 Vv	400	>650	960	960	0		17CA
CA2	Clear Lake	38 55 122 45	Basalt?	<10 ⁴ Yrb 2 x 10 ⁴ Yrb 9 x 10 ⁴ Yrb 2 x 10 ⁶ Yrb	256 Av	640-3540 Vv	1300	>650	3610	3610	2050	Chamber volume estimated here is virtually unchanged by new geophysical data (1600 Yrb).	6CA, 5CA, 10CA, 17CA
CA3	Long Valley	37 42 118 52	Rhyolite	10 ⁵ Yrb 7 x 10 ⁵ Yrb 9 x 10 ⁵ Yrb	480 Av	1200-4000 Vv	2400	>650	5700	5700	1800	Considered here as a system independent from Mono-Inyo Domes, but may be influenced by heat from Inyo fissure system.	3CA, 11CA, 17CA
CA4	Salton Sea	33 12 115 37	Rhyolite	1.6 x 10 ⁴ Yrb <3.5 x 10 ⁴ Yrb	50 Av	125-500 Vv	200	>650	480	480	75	More and better age data needed	8CA, 7CA, 17CA
CA5	Cono Mts.	36 02 117 49	Basalt	3 x 10 ⁴ Yrb 4 x 10 ⁴ Yrb 9 x 10 ⁵ Yrb	110 Av	275-1100 Vv	650	>650	1370	1370	270	Circular 726 estimate still considered valid. Data from new studies lead to both more liberal and more conservative estimates of chamber.	1CA, 7CA, 12CA, 17CA
CA6	Mono Domes	37 53 119 00	Rhyolite	<10 ³ Yrb <7 x 10 ⁴ Yrb	130 Avf	325-1300 Vv?	650	>650	1370	1370	50	Mono-Inyo valley systems complicated by Inyo Domes chain which intersects both and is best evidence for viable deep heat source.	2CA, 17CA

Table 1.—Magnitudes and heat contents of identified volcanic systems—Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition, last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume V ₀ (km ³)	Solidifi- cation state (°C)	AQ total (10 ¹⁰ J)	AQ new (10 ¹⁰ J)	AQ out (10 ¹⁰ J)	Remarks	References
CA7	Medicine Lake	41 35 121 37	Rhyolite	<10 ³ Yrs	64 AC 74 AV	160-640 Vv 185-740 Vv	300	>650	724	724	29	Should be studied in greater detail	17CA
CA8	Shasta	41 24 122 12	Andesite	52 ± 10 ² Yrs (5-9.5) × 10 ³ Yrs	50 Acop	125-500 Vvgo	300	>650	724	724	42	By analogy to Crater Lake and large gravity low.	9CA, 17CA
CA9	Sutter Buttes	39 13 121 49	Basalt?	1.4 × 10 ⁵ Yrs 1.5 × 10 ⁶ Yrs 2.4 × 10 ⁶ Yrs	40 AV	100-400 Vv	100	<650	240	<42	2107	Ring of silicic vents	17CA, 18CA
CA10	Morgan Mtn Domes	40 23 121 32	Dacite?	No data Plastic?								Needs investigation, age data.	17CA
CA11	Warner Mtns (Boyd Domes)—(Surprise Valley)	41 43 120 12	Rhyolite	7.1 × 10 ⁵ Yrs 7.7 × 10 ⁶ Yrs								Needs investigation. May be part of large subjacent pluton. See Cougar Peak, Oregon (same age).	13CA, 17CA
CA12	Bridgeport-Rodde Volcanic complex Calif.—Nevada	38 14;20 117 31	Basalt	2.5 × 10 ⁵ Yrs 1.1 × 10 ⁷ Yrs 2.5 × 10 ⁶ Yrs								Needs investigation. May be large low grade system.	15CA, 17CA
CA13	Lava Mountains	35 26 117 31	Silicic	No data Plastic? Pilo.		120 Vvgo	>120	<650	>280			Needs investigation as low grade resources. Volume probably greater by 5-10 times.	16CA, 17CA
CA14	Big Pine	37 03 118 19	Rhyolite	9.0 × 10 ⁵ Yrs 6 × 10 ⁴ Yrs	43 AC	105-420 Vv	100	<650	240	<45	1257	Slightly speculative (Vv).	4CA, 17CA
CA15	Olancha Domes	36 20 117 31	Rhyolite?	No data								Need age data.	1CA, 17CA
CA16	Jackson Buttes	36 16 120 44	Dacite?	No data Pilo?								Need age data.	17CA
CA17	Peoba Island Mono Lake	36 01 119 02	Silicic	85 yrs? Data not definitive								Needs more data. Active fumaroles; considered as separate system from Mono Domes.	2CA, 17CA
CA18	Inyo-Mammoth fiisuee system	37 46 119 01	Silicic?	7.25 × 10 ² Yrs 25 × 10 ² Yrs								May be thermal source and root for Long Valley system.	2CA
CA19	Templeton Domes	36 17 118 12	Silicic	1.9 × 10 ⁵ Yrs 2.4 × 10 ⁶ Yrs	50 AV	125-500 Vv	250	<650	603	603	310		1CA
N 1	Kilauea	19 26 155 18	Basalt	Active Yrs	12.5 Au	37.5-50 Vvgo	40	>650	96	96		Chamber probably a plume of sills and dikes with <5% of hot rock volume; molten at any one time.	18I, 28I
N 2	Mauna Loa	19 29 155 35	Olivine Basalt	Active Yrs								>8 km depth	18I

Table 1.—Magnitudes and heat contents of identified volcano systems—Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume (km ³)	Solidification state (°C)	AQ total (10 ¹⁸ J)	AQ now (10 ¹⁸ J)	AQ out (10 ¹⁸ J)	Remarks	References
H 3	Hualalai	19 42 155 50	Olivine Basalt	1.74 × 10 ² Yr								Potentially active. >5 km depth.	LHI
H 4	Mauna Kea	19 49 155 28	Keselite	Post-glacial Yr								Potentially active? No historic eruptions. >5 km depth.	LHI
H 5	Haleakala	20 43 156 15	Mafic Olivine Basalt	2.25 × 10 ² Yr								Potentially active. >5 km depth.	LHI
ID1	Island Park system	44 17 111 24	Basalt	2 × 10 ⁵ Yr 1.2 × 10 ⁶ Yr 1.2 × 10 ⁶ Yr 2.0 × 10 ⁶ Yr	3900 Ao 2100 Ao	5,250-21,000 Vc	16,200	5650	29,000	16,850	27,600	Compound system with Yellowstone caldera (WT-1). Ao and 16,200 km ² (area and volume of western part of Island Park system, not overlapped by Yellowstone Caldera) are used to calculate thermal energies.	21D, 61D
ID3	Blackfoot Dome	42 49 111 36	Silicofe?	4 × 10 ⁴ Yr 0 × 10 ⁴ Yr	25 AV	60-240 Vc	160	4550	240	240	59		11D, 31D, 61D
ID4	Big Southern Butte	43 24 113 01	Rhyolite	3 × 10 ⁵ Yr	No data	100 Vc	180	4550	240	<240		Some evidence that ID4 & ID5 could be part of single system. If so, huge thermal anomaly is possible.	11D, 61D
ID5	East Butte	43 20 112 40	Rhyolite	6 × 10 ⁵ Yr	No data	2 Vc	2	4550				See remarks ID4	17D
ID6	Fedburg Caldera	43 49 111 47	Rhyolite	4.2 × 10 ⁶ Yr	1800 Ao	4,500-18,000 Vc	9,000	4550	21,660	9,400	16,000		41D, 51D
NV1	Steamboat Springs	39 22 119 46	Rhyolite	1.2 × 10 ⁶ Yr	18 AV	45-180 Vc	90	4550	220	?	?	Fault-controlled system possibly related to chamber much larger than indicated.	28V, 34V
NV2	Silver Peak	37 45 117 52	Basalt	4.0 × 10 ⁶ Yr 6.1 × 10 ⁶ Yr	40 Ao	100-600 Vc	200	4550	480	?		Recent basalt cinder cone on lower flanks.	18V, 24V
NV1	Valles Caldera	35 52 106 34	Rhyolite	<10 ⁵ Yr 1.1 × 10 ⁶ Yr 1.4 × 10 ⁶ Yr	400 Ao	1,000-4,000 Vc	3,500	>650	8,425	9,435	<7,700	>20 rhyolitic eruptions <1.1 × 10 ⁶ Yr. See subject 18M, 38M, 48M to revision. Probably close to solidus temperature.	28M, 48M
NV2	Mt. Taylor	35 14 107 35	Basalt?	2.73 × 10 ⁶ Yr								Need more data.	28M, 48M
NV3	Mt. Agua Dome	36 45 105 57	Rhyolite	3.0 × 10 ⁶ Yr									28M, 48M
OR1	Crater Lake	42 56 122 07	Silicofe?	>7 × 10 ² Yr 6.6 × 10 ³ Yr	50 Ao	125-500 Vc	>120	>650	>770	>770	17		40R

Table 1.—Magnitudes and heat contents of identified volcanic systems—Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition, last eruption	Age date (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume V ₀ (km ³)	Solidifi- cation rate (°C)	ΔQ total (10 ¹⁰ J)	ΔQ new (10 ¹⁰ J)	ΔQ old (10 ¹⁰ J)	Remarks	Reference
OR2	Newberry	43 43 121 15	Rhyolite	1.2 x 10 ³ yrs >6.6 x 10 ³ TC	32 AC	80-320 Vc	100	>450	240	240	13		20R, 40R
OR3	South Sister	44 06 121 46	Rhyolite	<2 x 10 ³ yrs	20 AV 40 AC	75-300 VV	100	>450	240	240	21		40R
OR4	Mt. Hood	45 22 121 42	Andesite	Active								>10 km depth?	40R, 70R
OR5	Mt. McLoughlin	42 47 122 19	Andesite	Active?								>10 km depth	40R
OR6	Mt. Jefferson Domes (Steinbush?)	44 41 121 49	Silicic	No data Plinian.								Need age data.	40R, 60R
OR7	Malvin-Threes Creeks Buttes	44 10 121 36	Rhyolite	4 x 10 ⁵ yrs	10 AV	25-100 VV	40	<450	96	76	50	Need better age data. Three domes.	10R, 20R, 40R
OR8	Cappy-Burn Butte Area	43 19 121 56	Silicic	2.5 x 10 ⁶ yrs	0 AV	20-40 VV	40	<450	96	26	83	Need more data.	20R
OR11	Rustler Peak	42 37 122 21	Dacite?	No data Plinian.								Need age data.	40R
OR12	China Hat and East Butte	43 40 121 01	Rhyolite	7.8 x 10 ⁵ yrs 8.4 x 10 ⁵ yrs		85 Vcc	85	<450	203			These two domes may be close enough in time & space to be part of a single thermal anomaly.	20R, 40R
OR13	Quartz Mountain	43 36 120 53	Rhyolite	1.1 x 10 ⁶ yrs		36 Vcc	36	<450	80				30R, 40R
OR14	Glass Buttes	43 33 120 04	Rhyolite	54.9 x 10 ⁶ yrs		330 Vcc	330	<450	800	<407	750?	Needs investigation as low temperature system.	20R, 40R
OR15	Cougat Peak Area	42 18±30 120 36±20	Rhyolite?	7.1-7.7 x 10 ⁶ yrs				<450				Needs more detailed investigation including gravity data. May be underlain by pluton. See C11—same age.	20R, 40R
OR17	Harney-Malheur	43 15? 119 08?	Rhyolite?	56.6 x 10 ⁶ yrs 9 x 10 ⁶ yrs		2,500 Vcc	2,500	<450	6020	7		Needs more detailed investigation especially more accurate location of chamber area. Probably is large low grade system.	40R, 50R
OR18	Barvalow Buttes	44 05 121 33	Silicic	<10 ⁶ ?	8 AV	20-80 VV	30	<450	71	41	51	Need age data—four domes.	20R, 40R
OR19	Mart Peak Caldera	43 19 121 23	Rhyolite	4.5 x 10 ⁶ yrs TC	40 AC	180-400 Vc	150	<450	340	0	356		20R
OR20	Frederick Butte	43 37 120 28	Rhyolite	3.9 x 10 ⁶ yrs	32 AV AC?	80-320 VV Vc?	125	<450	300	4	300		20R

Table 1.--Magnitudes and heat contents of identified volcanic systems--Continued

1	2	3	4	5	6	7	8	9	10	11	12	13	14
No.	Name of area	Latitude, longitude (deg min)	Composition last eruption	Age data (yr)	Chamber area (km ²)	Chamber volume range (km ³)	Chamber volume Vo (km ³)	Solidifi- cation state (°C)	AQ total (10 ¹⁸ J)	AQ new (10 ¹⁸ J)	AQ out (10 ¹⁸ J)	Remarks	References
UT1	Mineral Mts.	38 26 112 48	Rhyolite	5 x 10 ⁵ Tys 0 x 10 ⁵ Tys	66 Av	165-660 Vv	300	<450	724	718	59		207, 407
UT2	Cove Creek Domes	38 45 112 44	Basalt	2.3 x 10 ⁶ Tys	94 Av	235-940 Vv	400	<450	960	84	920	Need geophysical data. May be low-grade system.	207, 407
UT3	White Mtn. Rhyolite	38 55 112 30	Basalt	<10 ⁴ Tys 4 x 10 ⁵ Tys	No data Small							Need geophysical data.	207, 407
UT4	Thomas Range	39 42 113 07	Rhyolite	6 x 10 ⁶ Tys	>1007 Av	>250-1,000 Vv	>500?	<450	>1,210	42	1,190	Need more data. Possible low-grade system.	107, 407
UT6	Smelter Knoll	39 26 112 50	Rhyolite	3.4 x 10 ⁶ Tys	No data			<450				Need more data.	107, 407
UT7	Wildcat Hills	41 51 113 01	Rhyolite	3.1 x 10 ⁶ Tys	Small			<450				Need more data.	207
NA1	Mt. Baker	40 47 121 49	Andesite	Active			W A O H I N G T O M					>10 km depth	3NA
NA2	Glacier Peak	48 07 121 07	Rhyodacite	1.2 x 10 ⁴ Tys <7 x 10 ⁵ Tys	5 ± 5 Av	12.5-50 Vv	12.5 ± 12.5	>450?	35?	35?	177	Volume by analogy to Mount St. Helens and other cascade volcanoes.	3NA
NA3	Mt. Rainier	46 51 121 45	Andesite	Active								>10 km depth	3NA
NA4	Mt. St. Helens	46 12 122 11	Andesite	Active 1.19 x 10 ² Tys 1.75 x 10 ³ Tys	5 Av	12.5-50 Vv	>12.5	>450	>35	>35			3NA
NA5	Mt. Adams	46 12 121 29	Andesite?	Active?								Need more data. >10 km depth?	3NA
NA6	Clear Fork Dacite	46 37 121 30	Dacite	>2 x 10 ⁴ Tys <3 x 10 ⁴ Tys <4 x 10 ⁴ Tys								Need more data.	3NA
NA7	Mann Butte	45 56 121 39	Rhyolite	(11-80) x 10 ⁴ Tys								Need more data.	1NA
W1	Yellowstone Caldera system	44 31 110 35	Rhyolite	6.9 x 10 ⁴ Tys 6 x 10 ⁵ Tys	2,500 Av	6,250-25,000 Vv	15,000	>450	36,100	36,100	7,950	Yellowstone caldera treated as independent from older system.	1W, 6W