0)
90
78-943

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

Office of Earthquake Studies

PROCEEDINGS OF
CONFERENCE VI
METHODOLOGY FOR IDENTIFYING SEISMIC GAPS
AND SOON-TO-BREAK GAPS
Convened Under Auspices of
NATIONAL EARTHQUAKE HAZARDS REDUCTION PROGRAM
25-27 May, 1978

OPEN-FILE REPORT 78-943

This report is preliminary and has not been
edited or reviewed for conformity with
Geological Survey standards and nomencla-
ture

Menlo Park, California

1978

CONFERENCES TO DATE

Conference I	Abnormal Animal Behavior Prior to Earthquakes, I
Conference II	Experimental Studies of Rock Friction with
Application to Earthquake Prediction
Conference III	Fault Mechanics and Its Relation to Earthquake
Prediction
Conference IV	Use of Volunteers in the Earthquake Hazards
Reduction Program
Conference V	Communicating Earthquake Hazard Reduction Information
Conference VI	Methodology for Identifying Seismic Gaps and
Soon-To-Break Gaps

CONFERENCES IN PROCESS

Conference VII	Measurement of Ground Strain Phenomena Related
to Earthquake Prediction |
|----------------|--|

90
78-943

UNITED STATES
DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY
Office of Earthquake Studies

[Reports - Open file Series]

PROCEEDINGS OF
CONFERENCE VI
METHODOLOGY FOR IDENTIFYING SEISMIC GAPS
AND SOON-TO-BREAK GAPS
Convened Under Auspices of
NATIONAL EARTHQUAKE HAZARDS REDUCTION PROGRAM
25 - 27 May, 1978

TM
cm
T. W. Amato

Co-Organizers

Bryan L. Isacks, 1936-
Department of Geological Sciences
Cornell University
Ithaca, New York 14850

George Plafker, 1929-
United States Geological Survey
Office of Earthquake Studies
Menlo Park, California 94025

Convener

Jack F. Evernden
United States Geological Survey
Office of Earthquake Studies
Menlo Park, California 94025

OPEN-FILE REPORT 78-943

This report is preliminary and has
not been edited or reviewed for con-
formity with Geological Survey standards
and nomenclature

296134

The views and conclusions contained in this document are those of the authors
and should not be interpreted as necessarily representing the official policies,
either expressed or implied, of the U.S. Government.

Menlo Park, California
1978

CONTENTS

Introduction and Summary.....	1
Origin of the Seismic Gap: What Initiates and Stops a Rupture Propagation Along a Plate Boundary? Keiiti Aki.....	3
Estimation of Non-Hydrostatic Stress in the Earth by Seismic Methods: Lithospheric Stress Levels Along Pacific and Nazca Plate Subduction Zones Charles Archambeau.....	47
Seismic Gaps in Coastal Peru James W. Dewey and William Spence.....	139
Seismicity Patterns Precursory to Larger Earthquakes in the Adak Region E.R. Engdahl.....	163
The Oaxaca Gap: A Case History Tomas Garza and Cinna Lomnitz.....	173
Deviatoric Stresses and Earthquake Occurrence at the Outer Rise Thomas C. Hanks.....	189
Some Characteristics of Foreshocks Lucile M. Jones and Peter Molnar.....	211
Statistical Study of the Occurrence of Shallow Earthquakes Yan Kagan and Leon Knopoff.....	267
Nature of Seismic Gaps and Foreshocks Hiroo Kanamori.....	319
Statistical Discrimination of Migrating Seismic Activity Keichi Kasahara.....	335
Bursts of Seismicity as Long-Term Precursors of Strong Earthquakes V.I. Keilis-Borok, L. Knopoff, I.M. Rotvain and T.M. Sidorenko..	351
On Soon-to-Break Seismic Gaps V. Keilis-Borok and J. Rotvain.....	387

A Point Process Analysis of Chinese Earthquake History	
W.H.K. Lee.....	395
Seismic Gaps and Plate Tectonics: Seismic Potential for Major Plate Boundaries	
William R. McCann, Stuart P. Nishenko, Lynn R. Sykes, and Janet Krause.....	441
Patterns of Seismicity Preceding Earthquakes in Central America, Mexico, and California	
Masakazu Ohtake, Tosimatu Matumoto and Gary V. Latham.....	585
The Makran Coast of Iran, A Possible Seismic Gap	
W.D. Page, G. Antonnen, and W.U. Savage.....	611
A Semi-Markov Model for Characterizing Recurrence of Great Earthquakes	
Ashok S. Patwardhan, Ram B. Kulkarni and Don Tocher.....	635
Uplift History and Earthquake Recurrence as Deduced from Marine Terraces on Middleton Island, Alaska	
George Plafker and Meyer Rubin.....	687
Analysis of Seismic Migration	
Albert T. Smith and Jacques Delsemme.....	723
A Model for the Earthquake Cycle in Underthrust Zones	
Wayne Thatcher and John B. Rundle.....	753
Space-Time Migration of Earthquakes Along the North Anatolian Fault Zone and Seismic Gaps	
M.N. Toksüz, A.F. Shakal and A.J. Michael.....	829
Patterns of Faulting and Seismic Gaps in the Great Basin Province	
Robert E. Wallace.....	857
Long Term Precursory Seismicity Fluctuations	
M.Wyss, R.E. Habermann and A.C. Johnston.....	869
Closing Comments	
Keiiti Aki.....	896
Michael Chinnery.....	897
James W. Dewey.....	898

E.R. Engdahl.....	899
Thomas Fitch.....	900
Thomas Hanks.....	901
Lucile Jones.....	902
Yan Kagan.....	903
Hiroo Kanamori.....	904
Keichi Kasahara.....	907
V. Keilis-Borok.....	909
Gary Latham.....	912
Willie Lee.....	913
Cinna Lomnitz.....	914
William McCann.....	915
W.D. Page.....	916
Ashok Patwardhen.....	917
George Plafker.....	918
C. Alfonso Reyes Z.....	919
Albert Smith.....	921
Wayne Thatcher.....	922
M. Nafi Toksöz.....	923
Max Wyss.....	924

INTRODUCTION AND SUMMARY

The fifth conference in the continuing series under the Earthquake Hazard Reduction Program was held at Massachusetts Institute of Technology on May 25-27, 1978. Its theme was "Methodology for Defining Seismic Gaps and Soon-to-Break Seismic Gaps." One of the proposed strategies for acquiring critical data on earthquake prediction is to identify soon-to-break gaps and to deploy a variety of instruments in the expected epicentral region. The Conference had as one of its major purposes the evaluation of our present capability to define gaps in general and soon-to-break gaps in particular, and thus to determine if the first step in the strategy proposed above could be fulfilled today.

This was a conference in which conclusions reached as to state of knowledge and state of capability depended grossly upon the individual attendee. Attitudes ran the gamut from exhilaration at how close we are to being able to define all sites of expected great earthquakes in continental areas (no clear time-scale to prediction) through enthusiasm about specific predictions, to a feeling of near-total lack of progress on the entire matter, the latter view being concerned with reasonably short term predictions (a year or so).

The only presentation that sought to consider the full scope of worldwide seismic activity by any mode of analysis with the hope of elucidating soon-to-break gaps was that by McCann et. al. This was a truly strenuous and important exercise and was the basis for the most extensive discussion at the Conference. Unfortunately, the paper by Archambeau was not presented at the Conference. It is included in the Proceedings because of its possibly direct bearing on the identification of soon-to-break gaps. As there was no discussion of the paper at the Conference, each reader must determine for himself the extent to which the case is made for high-stress drop foreshocks. This approach has not been diligently pursued against current data for any seismic arcs, so we have no predictions based on this concept.

As might be expected, the papers by Kagan and Keilis-Borok solicited much spirited comment. The model of prediction proposed by Keilis-Borok makes little or no pretense of being related to a physical model, it only makes the pretense of working. Until it actually does in a truly predictive mode, I expect the "I need a physical model" types will continue to find the approach of Keilis-Borok and his collaborators to earthquake prediction less than satisfying.

An issue that came into clear focus was the contrast in return times of great earthquakes as reported by seismologists and geologists, there being something like an order of magnitude discrepancy with geologists reporting longer return times. There was not even a serious guess as to the explanation of this apparent anomaly between the two data sets.

For what it is worth, the opinion of this writer is that no good case was made for quiet times before great quakes, or for demonstrable fore-shock patterns. If those techniques must be the basis for identifying soon-to-break gaps, we're in real trouble. One of the potentially interesting results was the suggestion by Smith that one can define a rate of progression of seismic activity along individual arcs. Much further analysis needs to be done, however.

A variety of other excellent papers, relevant to studies of island arc dynamics if not to definition of gaps, were presented and will be read with interest. As is obvious to all students of island arc dynamics, the questions of energy budget, state of stress in the subduction zone and modes of stressing the zone are open questions.

As always, the post-Conference impressions of all participants were invited and are included after the papers.

The poor reproductions of some figures and text result from the adamant refusal of a few participants to cooperate and to send originals of text and figures.

ORIGIN OF THE SEISMIC GAP:
WHAT INITIATES AND STOPS A RUPTURE PROPAGATION
ALONG A PLATE BOUNDARY?

Keiiti Aki

Department of Earth and Planetary Sciences

Massachusetts Institute of Technology

Cambridge, Massachusetts 02139

June, 1978

Abstract

From a reinterpretation of near-source and far-field seismic data on the Parkfield earthquake of 1966, we conclude that the rupture was stopped by a barrier with the specific fracture energy $8 \times 10^8 \text{ erg cm}^{-2}$. Using the formulas of Ida (1973b), we estimated parameters of the barrier as follows; the breaking slip of $15 \sim 23 \text{ cm}$ was required to break bonds with cohesive stress $70 \sim 105 \text{ bars}$ distributed over a non-elastic zone of linear dimension $340 \sim 770 \text{ meters}$. The barrier parameters for the 1857 California earthquake were also obtained from the description of surface fault breaks by Wallace (1968). The result led to the estimation of maximum acceleration of 1.5 g near the fault of the 1857 earthquake.

The barriers along the plate boundaries are classified into (1) fault bend, (2) junction, (3) ductile inhomogeneous barrier, and (4) brittle inhomogeneous barrier. The barriers act not only as a stopper of rupture; but also as an initiator of rupture, as well as a stress concentrator, causing twin earthquakes and migration or progression of major earthquakes along the plate boundary.

Using the functional form proposed by Mogi (1962), we find that the stress sensitivity coefficient β for the probability of fracture occurrence varies by 4 orders of magnitude depending on the kind of data used for estimation. The value of β estimated from migration data is $0.02\text{--}0.05 \text{ bar}^{-1}$, while the geodetic data on ultimate strain, the stress drop and recurrence of major earthquakes, and the correlation with

atmospheric disturbance gave the estimate of 0.3, 1.1, and 40 bar⁻¹, respectively. There is some suggestion of systematic increase of β with increasing wave-length of applied stress.

Introduction

As pointed out by Fedotov (1965) and Mogi (1968), the aftershock areas of large earthquakes in the past tens to hundreds of years tend to fill up a seismic zone without significant overlap. In the context of plate tectonics, this implies that the relative motion of plates is accomplished by a series of ruptures along the plate margin, each associated with a major earthquake. Then, it is natural to anticipate the next earthquake in a gap of aftershock zones unfilled for more than the expected recurrence time of major earthquakes for the seismic zone. Thus, the gap is important for a long range prediction of major earthquakes in the plate margin, as demonstrated, for example, by Sykes (1971) for the Sitka earthquake of 1972 and by Utsu (1972) for the Nemuro-Oki earthquake of 1973.

There have been several attempts to relate the size and location of aftershock zones with structural features such as submarine topography, width of the contact zone between two plates, and distribution of seamounts (Kelleher and McCann, 1976). In this paper, we shall make an inquiry into the physical nature of a seismic gap. First of all, why a plate margin is divided into a series of rupture zones. What are the boundaries between the neighboring rupture zones? What are the conditions of stress and strength in and near a gap?

We shall approach these questions from the point of the physics of rupture propagation by looking at the initiation and stopping of rupture processes along a plate margin

for a few earthquakes for which various pertinent data are available. Let us start with the Parkfield, California, earthquake of 1966.

Stopping of Rupture During the Parkfield Earthquake of 1966

This earthquake, after a dozen years since its occurrence, still offers the best instrumented case for near-source observations. The data obtained for this earthquake and their interpretation in terms of a propagating fault model were summarized in a review paper by Aki (1972). Here, we shall briefly describe a revised model based on a reinterpretation of the strong motion record by Bouchon (1978a,b) using a new seismogram synthesis method and on a re-examination of the observed discrepancy between seismic moments determined from near-source and far-field data.

The fault breaks of this earthquake were observed on the surface over a distance of 35 km along the San Andreas fault. From the first P times observed at local seismic stations, the starting point of the rupture was located near the northern end of the fault breaks. The chronograph at Gold Hill station gave a direct determination of rupture velocity. The rupture propagated southward with a velocity of 2.2 km/sec. This velocity and the fault length inferred from surface breaks were consistent with the observed features of spectra of long-period surface waves recorded around the world.

The accelerograph record at station No. 2, located at 80 meters from the fault trace near the southern end of the

fault break, was reinterpreted by Bouchon (1978b) using a new method applicable to a propagating slip along a finite fault in a layered half space. An excellent agreement was obtained between observed and synthetic records calculated for a simple uniform dislocation model with rupture velocity 2.2 km/sec, rise time 0.3 sec, and final slip 35 cm.

The fault breaks were terminated within a couple of kilometers from station No. 2, suggesting a sudden stop of a full-fledged rupture propagation with a large slip velocity (120 cm/sec). This sudden stop is supported by a sharp stopping phase observed at the Temblor station located at about 10 km in the direction of rupture propagation from the end of the fault breaks.

The seismic moment calculated from the product of the aftershock area (35 km x 15 km) and the slip determined from accelerogram (35 cm) is about 3 times greater than the value (1.4×10^{25} dyne cm) accurately estimated using both Love and Rayleigh waves observed at many WWSSN stations. The discrepancy was attributed earlier (Aki, 1968) to the narrow fault width of the mainshock fault which later extended to deeper parts by aftershocks and fault creep. We now feel that a better explanation may be sought in a fault model with barriers which may be left unbroken after a rupture propagation (Das and Aki, 1977). If some of the barriers on the fault plane of the Parkfield earthquake were unbroken during the mainshock rupture, the average slip may be significantly smaller than the value at station No. 2.

This barrier model may be tested with observed distribution of aftershock hypocenters. The unbroken barriers would cause stress concentration and become sources of aftershocks. On the other hand, the foci of aftershocks may be absent on a smoothly slipped segment of fault.

The aftershock distribution of the Parkfield earthquake studied by Eaton et al. (1970) shows clear evidence for the existence of such an unbroken barrier. (The standard error of epicenter location was less than 0.5 km for 59% and less than 1 km for 80% of all the earthquakes. The error of focal depth was only slightly larger than that of epicenter.) The fault trace of the Parkfield earthquake jumped from one side of Cholame Valley to the other, as shown in Figure 1. The initial plane of the aftershock zone is clearly terminated at this jumping point and a new plane, separated by about 1 km from the initial one, is created there and continues to the south. Figure 2 shows the projection of aftershock hypocenters on the fault plane. They are concentrated in the vicinity of this jumping point. There are several other clusters of aftershocks suggesting other unbroken barriers.

There are, on the other hand, areas with little aftershock activity. We hypothesize, as sketched in Figure 2, that the clusters of aftershocks define unbroken barriers and the areas of few aftershocks define slipped segments of fault. We recognize such a slipped segment near station No. 2 which may be approximated by a circular crack with radius of about 3 km.

Since the slip of 35 cm estimated from the accelerogram

probably refers to this crack, we get an estimate of stress drop as

$$\Delta\sigma = \frac{7\pi}{16}\mu \overline{\Delta u}/a = 50 \text{ bars} \quad (1)$$

where μ is the rigidity,

$\overline{\Delta u}$ is the average slip, and a is the radius of the crack.

As mentioned earlier, the rupture apparently stopped suddenly at the southern end of the zone of fault breaks. In general, a rupture may be stopped either by the presence of a fracture energy barrier or by a disappearance of stress (Husseini et al., 1975). In the latter case, however, we expect an overshoot of rupture propagation into the unstressed region as shown by Husseini et al. and the stopping phase would be insignificant (Dahlen, 1974). Our observation of the stopping phase at Temblor, then, supports the presence of a barrier at the stopping point. The barrier was strong enough to resist the extension of the crack with radius 3 km and stress drop 50 bars. Let us now estimate the fracture energy needed to break the barrier that stopped the Parkfield earthquake.

Fracture Energy Barrier That Stopped the Parkfield Earthquake Rupture

Consider an in-plane shear crack whose tip is extending with a velocity v . For v less than the Rayleigh wave velocity, the stress concentration ahead of the crack tip ensures an energy flow into the tip which will be spent for creating a new rupture surface. The fracture energy G per unit area of new crack surface may be written in terms of instantaneous stress intensity factor K (Freund, 1972) as

$$G = \frac{\pi K^2}{2\mu C} \quad (2)$$

where $C = 4(\beta/v)^2(1-v^2/\beta^2)^{-1/2}[(1-v^2/\alpha^2)^{1/2}(1-v^2/\beta^2)^{1/2} - (1-v^2/2\beta^2)^2]$.

α and β are compressional and shear velocity, respectively.

The instantaneous stress ahead of the crack tip is inversely proportional to the square root of the distance from the tip, and K is the proportionality constant.

To simplify the problem, we shall be more specific about the process around the crack tip. We shall look at the process from two different angles. First, we shall follow Barenblat (1959) and introduce cohesive force distribution in the non-elastic region between intact elastic media and completely broken region of crack surface. The cohesive force σ_c acting over the small region inside the crack just behind the tip can eliminate the stress singularity ahead of the tip if K is related to σ_c by the following equation.

$$K = \frac{1}{\pi} \int_{-d}^0 \sigma_c(\xi) (-\xi)^{-1/2} d\xi \quad (3)$$

where d is the length of the region where the cohesive force exists. For simplicity, we shall assume that $\sigma_c(\xi)$ is constant and equal to σ_c . Then, we have

$$K = \frac{2\sigma_c}{\pi} \sqrt{d} \quad (4)$$

Let us call d as "the barrier length". From (2) and (4), the fracture energy can be written as

$$G = \frac{2\sigma_c^2 d}{\mu\pi C} \quad (5)$$

The fracture energy G can be related to the cohesive force more directly, if we define the cohesive force as a function of relative displacement u between the two faces of the fault plane, as done by Ida (1973a).

$$G = \frac{1}{2} \int_0^D \sigma_c(u) du \quad (6)$$

Again, for the sake of simplicity, we shall assume that $\sigma_c(u)$ is constant independent of u . Then, we have

$$G = \frac{\sigma_c D}{2} \quad (7)$$

where D is the slip across the fault plane required to break the bond completely. Let us call D the "breaking slip".

Ida (1973b) calculated the slip motion for various choices of functional forms of $\sigma_c(u)$. Among other observables, he obtained the time t_M at which the slip velocity becomes maximum for an in-plane shear crack propagating with a uniform velocity v .

$$t_M = \text{const.} \cdot \frac{\pi}{32v} \left(\frac{2\mu G}{\sigma_c^2 C} \right) \quad (8)$$

where C is given in (2) and the constant on the right hand depends on the form of $\sigma_c(u)$, varying from 3 to 16, with the average about 7.7 for six cases studied by Ida.

He also derived the maximum particle velocity in the following

form

$$\dot{u}_M = \text{const.} \cdot \frac{16}{\pi} (\sigma_c v / \mu C) \quad (9)$$

where the constant averages .225 for the cases studied by Ida except for the singular case in which $\sigma_c(u)$ is the box-car function.

In order to find these fracture barrier parameters for actual earthquakes, we shall take a quasi-static view of rupture stopping and derive another relation between the parameters and observables. Consider the state of stress near the crack tip at the time of stopping of the Parkfield earthquake. We inferred from Figure 2 that a crack with radius $a = 3$ km and stress drop $\Delta\sigma = 50$ bars is resisted by a barrier. Such a crack will generate stress ahead of the crack tip according to a well known formula in fracture mechanics (e.g., Knott, 1973),

$$\tau = \Delta\sigma \sqrt{a/2x} \quad (10)$$

where x is the distance from the crack tip along the extension of the crack. A rough estimate of the cohesive force σ_c may be obtained, if we put $x = d$, that is, the barrier length introduced earlier.

Then, we have

$$\sigma_c^2 d = \frac{(\Delta\sigma)^2 a}{2} \quad (11)$$

Putting (11) into (5) we find that the specific fracture energy G can be determined only from $\Delta\sigma$, a , and v , as

$$G = \frac{(\Delta\sigma)^2 a}{\mu\pi C} \quad (12)$$

For the barrier that stopped the Parkfield earthquake, we have the following estimates

$$\Delta\sigma = 50 \text{ bars.}$$

$$a = 3 \text{ km,}$$

$$C = 1.0$$

The above value of C was obtained for $\alpha = 6.0 \text{ km/sec}$, $\beta = 3.5 \text{ km/sec}$, and $v = 2.2 \text{ km/sec}$.

Putting these values into (12), we obtain $G = 8 \times 10^8 \text{ erg cm}^{-2}$.

To find the other parameters, we shall apply Ida's formulas (8) and (9) to slip velocity and rise time determined by Bouchon (1978b) from the accelerogram records. First, we shall assume that the time of maximum velocity occurred in the middle of the ramp of Bouchon's slip function, that is, $t_M = \frac{0.3}{2} \text{ sec}$. From (8), (5), (7), and the parameters obtained above, we get

$$\sigma_c = 105 \text{ bars}$$

$$d = 340 \text{ meters}$$

$$D = 15 \text{ cm}$$

(13)

Another set of parameter estimates can be obtained by putting maximum particle velocity $\dot{u}_M = \frac{120 \text{ cm/sec}}{2}$ (half slip velocity) into (9). We, then, get using (5) and (7)

$$\begin{aligned}\sigma_c &= 70 \text{ bars} \\ d &= 770 \text{ meters} \\ D &= 23 \text{ cm}\end{aligned}\tag{14}$$

For another order of magnitude estimate of these parameters, we shall assume that the stress drops instantaneously over the whole fault plane by σ_c (Brune, 1970). Then, the particle velocity is simply σ_c divided only by impedance, $\rho\beta = 9.5 \times 10^5 \text{ dyne dm}^{-3} \text{ sec}$. For a particle velocity of 60 cm/sec, we obtain $\sigma_c = 60 \text{ bars}$, very close to the value given in (14). The other parameters are also similar to (14).

Considering the simplicity of our model, the agreement between the above three sets of estimates is encouraging. The outcome is somewhat surprising. The stress σ_c required to break the bond is very low, only one-and-a-half or twice higher than the stress drop 50 bars. This implies the weakness of bond involved in the rupture of the Parkfield earthquake. The large estimate of d suggests a large linear dimension over which the stress had to work to break the bond. This length (340~770 meters) may be compared with the diameter ($2a = 6 \text{ km}$) of crack whose extension was stopped by the barrier at the stopping point. We shall call $2a$ the "barrier interval".

An important parameter controlling the stress concentration around a barrier is the ratio of barrier length d to barrier

interval, which is 6-12% in this case. Finally, the breaking slip D was also very large. A large portion (40~65%) of the total slip (35 cm) is expended just to break the bond.

Fracture Energy Barrier for the Great Earthquakes of the San Andreas Fault

As shown by the numerical experiment of Das and Aki (1977), a rupture front can propagate over a fault plane leaving behind unbroken barriers. In a previous section of the present paper, we concluded that the fault plane of the Parkfield earthquake had unbroken barriers on the basis of the observed discrepancy between near-field and far-field estimates of seismic moment and the observed aftershock distribution as shown in Figure 2.

For a great earthquake that would break all the barriers on the fault plane, more work must be done per unit area of fault plane than for a smaller earthquake which breaks barriers only partially. In other words, for the same section of a fault the fracture energy may not be a constant but varies depending on how much energy a rupture front can supply.

Let us now estimate the barrier parameters for the Fort Tejon, California, earthquake of 1857. First we need some estimates of stress drop $\Delta\sigma$ and barrier interval $2a$ for calculating the specific fracture energy G . These estimates were taken from a geologist's description. We quote the following paragraphs from Wallace (1968)

...In attempting to locate the smallest clearcut offset that is related to the 1857 earthquake, 15 streams with no

offset were found. Some of these, no doubt, represent channels developed or straightened by flash floods since 1857, but some are of such large size that they must have existed before 1857, and yet there is no clear evidence of a shift to a new channel cut after 1857. One such large channel displaying no apparent offset is in NW1/4 Sec. 29, T.31S., R.21E., in the Painted Rock Quadrangle.

Because streams with no apparent offset lie along fault segments on which tens of feet of offset occur, one is led to the conclusion that the amount of offset varies along a given fault strand. An example is a particularly well-defined linear fault segment extending from about Highway 58 near Simmler, south for about 9 miles. This fault segment displays over 30 feet of offset near its midsection but has at least one stream channel showing no displacement near its southern end, which bows slightly to join an en echelon segment that overlaps it on the southwest side. A comparable pattern was noted in the surface fractures developed during the earthquakes at Parkfield and Cholame in June-August 1966 (Brown et al., 1967). Here en echelon segments of fracture zones, each several miles in length, had slips that were larger near their midpoints and slips that decreased to zero at the ends. Thus even in great earthquakes, such as that of 1857, which accompany displacements of many feet, the pattern may consist of composite segments, each several miles long, along each of which displacement reaches a maximum near the midsection and dies out at the ends. A total fracture length of 180 miles or more, as in the 1857 and 1906 earthquakes, might be a composite of 20 or more such individual segments.

From the above description, we envision that the fault breaks are a series of cracks with 15 km diameter and maximum slip 10 meters, connected by barriers. In this case, we use the following formula to estimate the stress drop.

$$\Delta\sigma = \frac{7\pi}{24}\mu \Delta u_{\max}/a = 370 \text{ bars} \quad (15)$$

Then, from (10), assuming $C = 1,0$ as in the case of the Parkfield earthquake, we have

$$G = 1 \times 10^{11} \text{ erg/cm}^2$$

which is about 100 times greater than the value obtained for the

Parkfield earthquake.

To obtain other parameters of the barrier, we shall assume that the barrier length is proportional to the barrier interval. In the case of the Parkfield earthquake, we estimated that the barrier length is about 6 to 12% of the barrier interval. Applying the same ratio to the 1857 earthquake, we have

$$d = 1 \sim 2 \text{ km.}$$

Then, putting the above values of G and d into (5), we find

$$\sigma_c = 500 \sim 700 \text{ bars}$$

and from (7),

$$D = 3 \sim 4 \text{ meters.}$$

Thus, again, we find the breaking slip D to be a significant fraction of total slip, and the stress needed to break bonds is also low, about 1-1/2 to twice the stress drop.

Let us make a short digression from the main theme of the present paper and estimate the peak acceleration expected for the 1857 earthquake.

The time t_M of the velocity peak according to (7) and (4) is proportional to the barrier length d . Then, t_M for the 1857 earthquake would be about 0.25 sec. Since the slip velocity is proportional to the stress drop $\Delta\sigma$ (Brune, 1970), the acceleration is proportional to the stress drop divided by t_M . Then, we expect

that the acceleration for the 1857 earthquake to be $\frac{370}{50} \times \frac{0.1}{0.25}$
 = 3 times that of the Parkfield earthquake. Since the peak
 acceleration observed at station No. 2 during the Parkfield
 earthquake was 0.5 g, we expect that the peak acceleration
 observed near the fault during the 1857 earthquake to be about
 1.5 g.

Discussion On the Specific Fracture Energy of G

The magnitude of the fracture energy ($10^9 \sim 10^{11}$ erg cm⁻²)
 obtained for the two California earthquakes is much larger than
 those estimated in the laboratory (10^3 erg cm⁻²) for small
 rock samples (Brace and Walsh, 1962), but is consistent with
 various estimates made for natural earthquakes by several
 investigators.

Kikuchi and Takeuchi (1970) studied the spontaneous
 propagation of longitudinal shear cracks and found that once a
 crack-tip starts propagating, for any initial length, it's speed
 will reach 0.9 β within a time roughly given by $2.5\pi\mu G(\Delta\sigma)^{-2}\beta^{-1}$.
 Assuming that the above time is less than about 10% of the
 observed total rupture propagation time for actual earthquakes,
 they obtained the upper bound for G to be on the order of 10^{10}
 erg cm⁻².

Ida (1973b), using equations (8) and (9) given in this paper
 and another formula for estimating acceleration, showed that
 typical strong motion data (maximum acceleration 1 g, maximum
 particle velocity 1 m/sec and the time scale of 0.1 sec) are
 consistent with $G \approx 10^{10}$ erg cm⁻², $\sigma_c \approx 1$ kbar, and $D \approx 10$ cm.

Husseini et al. (1975) considered two arrest mechanisms for a propagating longitudinal-shear crack, one by the presence of fracture energy barriers and the other by the vanishing of stress. In either case, a rough estimate of fracture energy was obtained by the following equation.

$$G = \frac{R(\Delta\sigma)^2}{2\mu\pi} \quad (16)$$

where $\Delta\sigma$ is the average stress drop and R is the characteristic radius of a circular fault equivalent to an earthquake. Equation (16) is similar to our equation (12), but the stress drop and crack radius used in (12) are associated with a single barrier interval, rather than the average over the entire fault plane. Husseini et al. (1975, 1976) estimated fracture energy for many earthquakes using (16). The fracture energy was found to vary from 10^3 to 10^9 erg cm^{-2} . Their results are probably biased toward underestimates because they use the average stress drop and total fault dimension. For example, they use the stress drop 1.3 bars and radius 13 km for the Parkfield earthquake, and their estimate of G is 1.1×10^5 erg cm^{-2} , that is, 1/7,000 of our estimate. We believe that the local stress drop within a barrier interval should reflect the fracture energy of the barrier more correctly than the average stress drop over the entire fault plane.

Das (1976) made a series of numerical experiments on a spontaneous propagation of an in-plane shear crack and found that if the rupture velocity is below Rayleigh waves in actual

earthquakes, G must be 10^9 ergs cm^{-2} for a fault length of 10 km and must be 10^{11} ergs cm^{-2} for a fault length 1000 km. Her result is in harmony with our estimates of G for the Parkfield and 1857 California earthquakes.

Finally, let us consider how this large amount of energy is consumed in the rupture zone. Take the example of the 1857 earthquake. The energy is absorbed through a slip as much as 3~4 meters at the stress of 500~700 bars. The absorption is taking place over a non-elastic zone with linear dimension of 1~2 km. Interlocked blocks of rock of this dimension are involved in the process of breaking bonds between the two faces of fault rather than interatomic or intergranular bonds. It is, then, likely that the energy is absorbed not into a thin film covering the fault surface, but into plastic deformation and micro-cracking in a zone with thickness tens of cm or meters.

Suppose that the energy is absorbed into the thin film covering the fault surface. In this case, the temperature rise T may be estimated (McKenzie and Brune, 1972) roughly by

$$T \approx \frac{\sigma_c D}{\rho C_p \sqrt{\pi k t}} \quad (17)$$

where k is the thermal diffusivity, t is the rise time of fault slip, ρ is the density and C_p is the specific heat of rock. For $\sigma_c D = 2G = 2 \times 10^{11}$ erg cm^{-2} , $\rho = 3 \text{ gm}^{-3}$, $C_p = 10^7$ erg $\text{g}^{-1} \text{ } ^\circ\text{C}^{-1}$, $k = 7 \times 10^{-3} \text{ cm}^2 \text{sec}^{-1}$, and $t = 1 \text{ sec}$, we find $\sqrt{\pi k t} = 0.15 \text{ cm}$, and $T = 4000^\circ\text{C}$, which will certainly produce a film of melt. On the other hand, if the energy is absorbed in the zone of width, say 15 cm, than the temperature rise should

be only 40°C, and no melting will occur.

Since the high energy barriers may occupy only a small fraction of fault plane, their contribution to the total terrestrial heat flow would be negligible and would not violate the constraint from heat flow data (Brune et al., 1969).

How the Rupture Stopped During the 1857 and 1906 California Earthquakes

According to our estimate, the barriers involved in the 1857 earthquake were as high as 10^{11} erg cm⁻². On the other hand, the Parkfield earthquake rupture was stopped by a barrier with fracture energy around 10^9 erg cm⁻². This stopping point is usually considered also the northern end of the 1857 earthquake rupture. To the north of this point, the fault creep occurs at a rate of about 20 mm/year. But, the rate as measured by the in-line geodimeter (Savage, 1975) drops to 0~10 mm/yr to the south of this point, usually referred to as the "blocked zone" as opposed to the "creeping zone" to the north. It is difficult to imagine that the 1857 rupture which went through the high energy barriers can be stopped by a weak barrier with 2 orders of magnitude lower fracture energy. If the stopping is due to the vanishing of stress, on the other hand, we expect the dynamic stress field concentrated around the crack-tip would result in a considerable overshoot of rupture into unstressed regions (Hussieni et al., 1975). In fact, according to Sieh (1977), the 1857 slip might have propagated into the creeping zone 60 km or more.

Let us next turn to the California earthquake of 1906; a great earthquake which occurred on another blocked zone of the San Andreas fault north of the creeping zone. The rupture propagation of this earthquake was studied by Bolt (1968) and Boore (1977). Boore gives a further support to Bolt's conclusion that the main shock began closer to San Francisco than Olema, as originally proposed by Reid (1910). Since the fault breaks extend from San Juan Bautista (about 130 km south of the revised epicenter) to Cape Mendocina (about 300 km north of the revised epicenter) the rupture propagation must have been bilateral.

We found additional evidence supporting the bilateral propagation for this earthquake. The new evidence comes from the sense of ground motion near the rupture front in the direction perpendicular to the fault trace. As shown by Bouchon (1978b) from the accelerogram records of the Parkfield earthquake obtained at station No. 2, both synthetic and actual displacement seismograms near the fault indicate a small initial perpendicular motion directed to the southwest. A field survey of earthquake damage (Brown et al., 1967) revealed that a tall stack of rolls of heavy wire collapsed in a building near the fault break. They were thrown to the northeast in agreement with the sense of initial ground motion toward the southwest. This suggests the possibility of using unstable objects near a fault trace to determine the direction of rupture propagation. If the direction of rupture propagation of the Parkfield earthquake was northward, instead of southward, the rolls of wire should have been thrown to the southwest.

The report of the 1906 earthquake (Lawson, 1908) contains several descriptions of seismic motion in the vicinity of the fault trace. One of the most reliable descriptions is due to G.K. Gilbert, who made a careful observation of large water tanks in the village of Inverness. Since a large water mass is supported by a structure relatively weak to horizontal force, the direction to which the tank was thrown may correspond to the initial motion due to rupture passage. Out of five tanks, two were thrown in a direction almost due west and demolished. One was shifted to the west without collapse. One fell to the southwest and tank and pedestal were both crushed. The last one was thrown to the west-northwest, being overturned as it fell. This suggests an initial ground motion in an easterly direction. Since this is nearly opposite to the direction observed during the Parkfield earthquake, we suggest that the direction of rupture propagation was also opposite, namely northward.

Another example of the movement of unstable objects was found at the Morrell ranch located 1 mile south of Wright's station (about 20 km north of the end of the fault break). An especially strongly constructed wine cellar located close to the fault trace had the upper portion thrown 3 feet northeast, directly away from the fault line. After the shock, this upper portion of the house was left resting upon the wine tuns, and not upon its original upright support which was very weak to the horizontal force, as shown in Figure 3. This direction of movement of the wine cellar agrees with the one observed for

the rolls of heavy wire thrown during the Parkfield earthquake, suggesting the same direction of rupture propagation, namely, southward. Thus, the two observations of particularly unstable structures located close to the fault support that rupture propagation in the 1906 earthquake was bilateral.

To the north of the epicenter, the final slip averaged around 4 meters and the rupture propagation reached nearly to Cape Mendocino, where the Mendocino fracture zone intersects the San Andreas fault. In other words, the rupture propagated as far as the end of a linear plate boundary. It was stopped because it could not penetrate into a rigid plate. If we assume the same barrier interval (15 km) for this earthquake as for the 1857 earthquake, the fracture energy for the northern section may be estimated roughly as $2 \times 10^{10} \text{ erg cm}^{-2}$.

On the other hand, the final slip near the southern end of the fault is around 1 meter (Boore, 1977). Again, assuming the same barrier interval as the 1857 earthquake, the fracture energy for this part of the fault is estimated roughly as 10^9 erg cm^{-2} . This is rather low and comparable to the value obtained for the Parkfield earthquake. The stopping of rupture at San Juan Bautista without any overshoot into the creeping zone, a fact well established by field observations and geodetic measurements (Thatcher, 1978), may be due to the existence of a barrier at San Juan Bautista.

It is interesting to note here the existence of a prominent seismic velocity anomaly lying over the San Andreas fault near San Juan Bautista discovered by the three-dimensional inversion of teleseismic P-wave data by Zandt (1978). The east-northeast

trending anomaly with about 4% high velocity crosses the San Andreas fault obliquely in the depth range from 10 to 30 km. This velocity anomaly may be related to the barrier that stopped the 1906 earthquake rupture.

An alternative explanation for the stopping of the 1906 rupture may be the absence of stress in the creeping zone south of San Juan Bautista. Thatcher (1978), however, points out a puzzle about this explanation. As shown by Spence and Turcotte (1976), if the plate boundary is stress free except for two long shallow locked strips (corresponding to 1857 and 1906 fault sections) separated by an intervening central crack (corresponding to the creeping zone), the creep rate on the stress free central crack is only a small fraction, about one-tenth, of the relative plate motion rate. This large deficit must be made up by increased creep on the central crack during the relaxation process following a great earthquake on one of the locked segments. Thatcher finds, however, no such evidence. The 1906 earthquake did not generate significant increase of creep rate in the central segment, suggesting a major barrier near San Juan Bautista which decoupled the slip motion of the blocked segment from that of the creeping segment.

Classification of Barriers

The barriers that stop an earthquake rupture extension may be classified broadly into two groups: geometric barriers and inhomogeneous barriers. The geometric barriers refer to the corner at which the plate margin changes direction such as

the junction of the Mendocino fracture zone and the San Andreas fault which stopped the northern extension of the 1906 earthquake. Another well known example is the southern end of the 1960 Chilean earthquake which stopped at the junction of the Chile Rise.

The inhomogeneous barrier refers to the stopping point of earthquake rupture where no obvious geometrical discontinuity exists. Examples are the southern ends of the Parkfield and 1906 earthquake faults. In the latter case, the barrier appears to be associated with an anomaly of seismic velocity in the lower crust. Another example may be the Montagua fault during the Guatemala earthquake of 1976. The western end of this fault appears to have stopped near a line of volcanoes which intersects the extension of the fault obliquely.

Kelleher and McCann (1977) pointed out an interesting correlation between the distribution of sea mounts seaward of the Japan trench and the size of rupture zones of major earthquakes (estimated by the aftershock zone and tsunami source area) occurring on the other side of the trench. As the interval between sea mounts decreases, the size of the rupture zone also decreases. This may be explained, if the sea mounts with their deep roots are acting as inhomogeneous barriers when the lithosphere including them underthrust the Eurasia plate by subduction.

Some barriers are sharp and others are dull stress concentrators. The sharpest stress concentrators may be the barriers between so-called twin earthquakes. For example, Ando (1975) describes a region in southwestern Japan where a major earthquake is repeatedly followed by another in the same neighbor region.

The time lag between the pair was two years for the Tokai (1944) and Nankaido (1946) earthquakes, 32 hours for the Ansei I and II earthquakes (1854), and a few minutes in the case of the Hoei earthquake (1707), which was suspected as a twin because of unusually long duration of strong oscillation.

When the barriers act as stress concentrators one by one to the next neighbor, a migration or progression of major earthquakes can occur along a plate boundary. There are several examples such as the westward migration along the Anatolian fault from 1939 to 1944, the westward migration in the Aleutians from 1938 to 1965, the northward migration in southeastern Alaska from 1949 to 1964, and the southward migration in Chile from 1822 to 1837 and 1906 to 1960.

Let us make an estimate of the stress concentration around a barrier using the model discussed earlier in this paper.

The increase of stress σ around a barrier with length d by a crack with radius a and stress drop $\Delta\sigma$ can be approximated by equation (10),

$$\sigma = \Delta\sigma \sqrt{a/2d} \quad (18)$$

Thus, a barrier with small length d and long interval $2a$ is a sharp stress concentrator.

For the Parkfield earthquake, we estimated $d/(2a)$ to be 6-12%, therefore the stress increase is 1.4-2 times the stress drop. Since, $\Delta\sigma$ was estimated as 50 bars for the Parkfield case, the stress increase around the barrier would be about 70 to 100 bars.

What happens when the stress increase of this magnitude occurs? Apparently various things can happen. In the case of the Parkfield earthquake, there were no aftershocks in the vicinity of the stopping point. The stress concentration may still be there or released through plastic deformation. In many cases, however, aftershocks cluster near the ends of the rupture zone, and migrate away gradually extending the aftershock area. In these cases, the stress concentration is relieved at least partially by brittle fracture.

An example of strong brittle inhomogeneous barriers may be found at the stopping point of the great Alaskan earthquake of 1964. The rupture of this earthquake was started from the northeastern end of the aftershock area and propagated toward the southwest (Kanamori, 1970). The starting point was near a corner of the Pacific plate where a north-south trending transform fault and an east-west running subduction zone meet. It is a junction of different modes of fault slip and a nucleation of rupture may be expected. Interestingly, however, the foreshock activity was concentrated near the stopping point rather than the starting point according to Kelleher and Savino (1975). These foreshocks lie near the northeastern end of the aftershock zone of the 1938 earthquake ($M = 8.7$). The rupture of the 1938 earthquake might be stopped by a strong brittle barrier with the stress concentration causing continuous seismicity over two decades until 1964. The same barrier then stopped the 1964 rupture propagating from the other side.

Thus, some barriers may be ductile, relaxing stress by

plastic deformation, and others may be brittle, relaxing stress by aftershocks.

Let us now turn to the nucleation of rupture, and the possible role of a barrier in starting an earthquake rupture.

The starting point of the Parkfield earthquake was located near Middle Mountain, which separates two different modes of fault slip along the San Andreas fault (Brown, 1970). To the north of Middle Mountain, the trace of recently active faults is segmented, with each segment 2 to 6 miles long. The fault creep rate there is 2 to 3 cm per year. To the south of Middle Mountain, the fault is not broken into short segments, and has repeatedly been a site of moderate magnitude earthquakes. The creep here is not continuous but spasmodic, and the rate of movement is about half of that to the north of Middle Mountain. Such a discontinuous change in the mode of slip may have caused a stress concentration and resulted in the nucleation of the Parkfield earthquake.

There is, however, no obvious reason why the mode of slip should be discontinuous at Middle Mountain, except that a slight change in the fault strike direction occurs there. Such a small change in fault strike direction or a fault bend may be important as a site of earthquake nucleation. For example, the strike direction of the San Andreas fault in the San Francisco peninsula is not parallel with that in Marin County. Therefore, there must be a bend offshore near San Francisco. This bend satisfies the constraint posed by Boore (1977) for the starting point of the 1906 rupture. Another example is the

Izu-Hanto-Oki earthquake of 1974 studied by Matsuda and Yamashina (1974). The epicenter is located near a slight bend of the fault trace, suggesting that the rupture might have started from the bend. These examples suggest the importance of very subtle geometrical barriers in the earthquake rupture process, in addition to inhomogeneous barriers.

Simple-minded mechanical models with homogeneous properties and planar boundaries may be missing some of the key elements of the earthquake phenomena.

Now, we shall attempt to be more quantitative about the effect of stress concentration on the occurrence of earthquakes.

Stress Dependence of the Probability of Earthquake Occurrence

How the probability of earthquake occurrence depends on the applied stress is an extremely complex problem. According to Mogi (1962), the probability of occurrence of fracture in a rock sample increases exponentially with the applied stress. When a constant stress σ is applied at $t=0$, he found that the probability of occurrence of fracture between t and $t+dt$ is independent of t and given by

$$\mu(t)dt = \mu_0 \exp(\beta\sigma) \quad (19)$$

where β is determined as 0.37 bar^{-1} for the tensile fracture of granite samples in the atmosphere. The experiment by Scholtz (1972) on static fatigue of quartz under uniaxial compression showed the same functional dependence on stress, but with β about

one hundred times smaller than Mogi's value.

The coefficient β can be estimated from the recurrence time of major earthquakes in a fault zone and the associated stress drop. For the Hokkaido-Kurile region, Utsu (1972) constructed a statistical model based on (19) and determined the model parameter by fitting the data on recurrence time. Combining Utsu's result with the average stress accumulation rate ($0.3 \text{ bar year}^{-1}$) inferred from the stress drops in major earthquakes in this zone (Fukao and Furumoto, 1978), we find β is 1.1 bar^{-1} . Earlier, Hagiwara (1974) obtained β of about 0.3 bar^{-1} by applying equation (19) to the statistical distribution of ultimate strain obtained by geodetic measurements.

An apparent sensitivity of earthquake occurrence to small stress changes such as due to earth tide and atmospheric pressure gives a large estimate of β . Conrad (1932) shows an impressive evidence for the increase of local earthquake frequency by 30% due to the atmospheric pressure gradient across the Alps by 5 mm Hg (6.7×10^{-3} bars) or greater. This gives β to be about 40 bar^{-1} .

With this great variability of the value of β (0.004 to 40 bar^{-1}) estimated by different methods, it is no use to attempt to estimate the probability of migration of earthquakes on the basis of our barrier model. Rather, we shall estimate the value of β applicable to the migration process.

The migration or progression of major earthquakes typically shows the time interval between the neighbor events to be $1/5$ to $1/100$ of the recurrence time. In other words, the probability

of occurrence may be increased by a factor of 5 to 100. If this increase is caused by a stress concentration of 100 bars, as estimated for the case of the Parkfield barrier, we have

$$\exp(100\beta) = 5 \sim 100$$

which gives the estimate of β in the range 0.02 to 0.05 bar⁻¹.

The value of β estimated by various methods, thus ranges from 0.004 to 40 bar⁻¹ over 4 decades. There is some suggestion of increasing β with increasing wave length of the stress field as we compare the values from migration (0.02-0.05 bar⁻¹), geodetic data on ultimate strain (0.3 bar⁻¹), stress drop and recurrence of major earthquakes (1.1 bar⁻¹) and atmospheric disturbance (40 bar⁻¹). Nevertheless, since the probability depends on the exponential of β times stress, the variability of probability estimate is enormous. To find factors determining β will be the most important step towards a quantitative prediction of earthquake occurrence.

Summary

In order to understand why the plate boundary is divided into seismic gaps, we examined what initiates and stops a rupture propagation along the boundary during major earthquakes.

We studied in detail the stopping mechanism of the Parkfield earthquake of 1966 for which both near-source and far-field data are available. We concluded that the rupture of the

Parkfield earthquake was stopped by a fracture energy barrier and described the barrier with the following physical parameters. The specific fracture energy G was $8 \times 10^8 \text{ erg cm}^{-2}$, cohesive stress σ_c was $70 \sim 105$ bars, the slip D required to break bonds was $15 \sim 23$ cm. We introduced a new concept of barrier length d which specifies the linear dimension of non-elastic zone undergoing the transient state from intact elastic body to completely broken fault zone. This is the linear dimension of the zone over which the cohesive stress must act to break the bond. For the Parkfield earthquake, the barrier length d was estimated as $340 \sim 770$ meters. Thus, the barrier involved in the stopping of the Parkfield earthquake rupture implies interlocked blocks of rock with a linear dimension of a few hundred meters which are forced to slip under the stress of about 100 bars. A slip of around 20 cm is required for breaking the bonds between blocks.

Next we considered the barrier encountered by the 1857 California earthquake rupture based on the geological description of the fault breaks by Wallace (1968). Assuming the same ratio of barrier interval to barrier length as in the Parkfield case, we arrived at the following parameters: $G = 10^{11} \text{ erg cm}^{-2}$; cohesive stress $\sigma_c = 500 \sim 700$ bars; barrier length $d = 1 \sim 2$ km; and breaking slip $D = 3 \sim 4$ meters. Although the fracture energy of $10^{11} \text{ erg cm}^{-2}$ appears to be enormous, it will raise the temperature only by 40°C if the energy is spent in heating the zone with thickness 10 cm. The contribution to the total terrestrial heat flow would be negligible

because these high energy barriers are not distributed uniformly over the entire fault surface, but only over a small fraction of it. The maximum acceleration expected near the fault of the 1857 earthquake is estimated at 1.5 g.

We then attempted to classify the barriers along plate boundaries around the world. They are divided into four major categories: (1) fault bend; (2) junction; (3) ductile inhomogeneous barrier; and (4) brittle inhomogeneous barrier. The corresponding examples are (1) the starting point of the Parkfield earthquake, (2) the stopping point of northward rupture of the 1906 California earthquake, (3) the stopping point of the Parkfield earthquake, and (4) the stopping point of the Alaskan earthquake of 1964.

The stress concentration around a barrier will be high if the ratio of barrier interval to barrier length is large. Sharp stress concentrators may be causing twin earthquakes and migration or progression of major earthquakes along plate boundaries.

Finally, we discussed the stress dependence of the probability of earthquake occurrence using the data on migration. Using the form of equation (19) proposed by Mogi (1962), we find that constant β which describes the stress sensitivity of earthquake occurrence probability varies by four orders of magnitude depending on the kind of data used. The value of β estimated from migration data is $0.02-0.05 \text{ bar}^{-1}$; while the geodetic data on ultimate strain, the stress drop and recurrence of major earthquakes and the correlation with atmospheric

disturbance gave the estimate of 0.3, 1.1, and 40 bar⁻¹, respectively. Although there is some suggestion of systematic increase of β with increasing wavelength of applied stress, the uncertainty in β is enormous. To find factors determining β will be the most important step towards a quantitative prediction of earthquake occurrence.

Acknowledgements

This work was partially supported by National Science Foundation grants 77-13632EAR and 77-23336PFR.

References

- Aki, K., Seismic displacement near a fault, J. Geophys. Res., 73, 5359-5376, 1968.
- Aki, K., Earthquake mechanism, in A.R. Ritsema (editor), The Upper Mantle, Tectonophysics, 13, 423-446, 1972.
- Ando, M., Source mechanism and tectonic significance of historical earthquakes along the Nankai trough, Japan, Tectonophysics, 27, 119-140, 1975.
- Barenblat, G.I., The formation of equilibrium cracks during brittle fracture; general ideas and hypotheses, axially symmetric cracks, J. Appl. Math. Mech., 23, 434-444, 1959.
- Bolt, B.A., The focus of the 1906 California earthquake, Bull. Seis. Soc. Am., 58, 457-471, 1968.
- Boore, D.M., Strong-motion recordings of the California earthquake of April 18, 1906, Bull. Seis. Soc. Am., 67, 561-578, 1977.
- Bouchon, M., Discretization of seismic wave fields in three space dimensions, in preparation, 1978a.
- Bouchon, M., Study of strong motion records obtained during the Parkfield earthquake of 1966, in preparation, 1978b.
- Brown, R.D., Jr., J.G. Vedder, R.E. Wallace, E.F. Roth, R.F. Yerkes, R.O. Castle, A.O. Waananen, R.W. Page, and J.P. Eaton, The Parkfield-Cholame, California, earthquakes of June-August, 1966 - Surface geologic effects, water resources effects, and preliminary seismic data, U.S. Geol. Survey Prof. Paper, 579, 66 p., 1967.

- Brown, J.R., Jr., Map showing recent active breaks along the San Andreas and related faults between the northern Gabilan Range and Cholame Valley, California, Miscellaneous Geologic Investigations, Map I-575, U.S. Geol. Surv., 1970.
- Brune, J.N., T.L. Henyey, and R.F. Roy, Heat flow, stress, and the rate of slip along the San Andreas fault, California, J. Geophys. Res., 74, 3821-3827, 1969.
- Brune, J.N., Tectonic stress and the spectra of seismic shear waves from earthquakes, J. Geophys. Res., 75, 4997-5009, 1970.
- Conrad, V., Die Zeitlich Folge der Erdbeben und Bebenaulösende Ursachen, Handbuch der Geophysik, Bd. IV, 1007-1185, 1932.
- Dahlen, F.A., On the ratio of P-wave to S-wave corner frequencies for shallow earthquake sources, Bull. Seis. Soc. Am., 64, 1159-1180,
- Das, S., A numerical study of rupture propagation and earthquake source mechanism, Ph.D. thesis, Mass. Inst. of Technol., Cambridge, 1976.
- Das, S. and K. Aki, Fault planes with barriers: a versatile earthquake model, J. Geophys. Res., 82, 5658-5670, 1977.
- Eaton, J.P., M.E. O'Neill, and J.N. Murdock, Aftershocks of the 1966 Parkfield-Cholame, California, earthquake: a detailed study, Bull. Seis. Soc. Am., 60, 1151-1197, 1970.
- Fedotov, S.A., Regularities of the distribution of strong earthquakes in Kamchatka, the Kurile Islands, and northeastern Japan, Tr. Inst. Zemli Akad Nauk SSSR, 36, 66, 1965.
- Freund, L.B., Energy flux into the tip of an extending crack in an elastic solid, J. Elasticity, 2, 341-348, 1972.

- Fukao, Y., and M. Furumoto, Stress drop, wave spectrum and recurrence interval of great earthquakes - implication from the Etorofu earthquake of November 6, 1958, Geophys. J.R. Astr. Soc., in press, 1978.
- Hagiwara, Y., Probability of earthquake occurrence estimated from results of rock fracture experiments, Tectonophysics, 23, 99-103, 1974.
- Husseini, M.I., D.B. Jovanovich, M.J. Randall, and L.B. Freund, The fracture energy of earthquakes, Geophys. J. R. Astr. Soc., 43, 367-385, 1975.
- Husseini, M.I., D.B. Jovanovich, and M.J. Randall, Fracture energy and aftershocks, Geophys. J.R. Astr. Soc., 45, 393-406, 1976.
- Ida, Y., Cohesive force across the tip of a longitudinal-shear crack and Griffith's specific surface energy, J. Geophys. Res., 77, 3796-3805, 1973a.
- Ida, U., the maximum acceleration of seismic ground motion, Bull. Seis. Soc. Am., 63, 959-968, 1973b.
- Kanamori, H., The Alaska earthquake of 1964: radiation of long-period surface waves and source mechanisms, J. Geophys. Res., 75, 5029-5040, 1970.
- Kelleher, J. and W. McCann, Buoyant zones, great earthquakes, and unstable boundaries of subduction, J. Geophys. Res., 81, 4885-4896, 1976.
- Kelleher, J.A. and W. McCann, Bathymetric highs and the development of convergent plate boundaries, in M. Talwani and W.C. Pitmann, III (editors), Maurice Ewing Series I: Island Arcs,

- Deep Sea Trenches, and Back Arc Basins, Am. Geophys. Un., 115-122, 1977.
- Kelleher, J.A. and J. Savino, Distribution of seismicity before large strike-slip and thrust-type earthquakes, J. Geophys. Res., 80, 260-271, 1975.
- Kikuchi, M. and H. Takeuchi, Unsteady propagation of longitudinal-shear cracks, Zisin, 23, 304-312,
- Knott, J.F., Fundamentals of Fracture Mechanics, Butterworths, London-Boston, 273 pp, 1973.
- Lawson, A.C., The California earthquake of April 18, 1906, Report of the State Earthquake Investigation Commission, Carnegie Institution of Washington, Vol. 1, 1908.
- Matsuda, T. and K. Yamashina, Surface faults associated with the Izu-Hanto-Oki earthquake of 1974, Japan, Sokuho (in Japanese) Earthq. Res. Inst., Univ. Tokyo, 14, 135-158, 1974.
- McKenzie, D. and J.N. Brune, Melting on fault planes during large earthquakes, Geophys. J.R. Astr. Soc., 29, 65-78, 1972.
- Mogi, K., Study of elastic shocks caused by the fracture of heterogeneous material and its relation to earthquake phenomena, Bull. Earthq. Res. Inst., Univ. Tokyo, 40, 125-173, 1962.
- Mogi, K., Sequential occurrences of recent great earthquakes, J. Phys. Earth, 16, 30-36, 1968.
- Mogi, K., Relationship between the occurrence of great earthquakes and tectonic structures, Bull. Earthq. Res. Inst., Univ. Tokyo, 47, 429-452, 1969.

- Reid, H.F., The California earthquake of April 18, 1906, Vol. 2, The mechanics of the earthquake, Report of the State Earthquake Investigation Commission, Carnegie Institution of Washington, 1910.
- Savage, J., A possible bias in the California state geodimeter data, J. Geophys. Res., 80, 4078-4088, 1975.
- Scholtz, C.H., Static fatigue of quartz, J. Geophys. Res., 77, 2104-2114, 1972.
- Sieh, K., Study of Holocene displacement history along the southcentral reach of the San Andreas fault, Ph.D. thesis, Stanford University, Stanford, 219 p., 1977.
- Spence, D.A. and D.L. Turcotte, An elastostatic model of stress accumulation on the San Andreas fault, Proc. Roy. Soc. London, A349, 319-341, 1976.
- Sykes, L.R., Aftershock zones of great earthquakes, seismicity gaps, and earthquake prediction for Alaska and the Aleutians, J. Geophys. Res., 79, 8021- , 1971.
- Thatcher, T., Systematic inversion of geodetic data in central California, J. Geophys. Res., submitted, 1978.
- Utsu, T., Large earthquakes near Hokkaido and the expectancy of the occurrence of a large earthquake off Nemuro, Report of the Coordinating Committee for Earthquake Prediction, Geographical Survey Inst., Ministry of Construction, Japan, 7, 7-13, 1972.
- Wallace, R.E., Notes on stream channels offset by the San Andreas fault, southern Coast Ranges, California, Proceedings of Conference on Geologic Problems of the San Andreas Fault

System, Stanford University Publications in the Geological Sciences, 11, 6-21, 1968.

Zandt, G., Study of three-dimensional heterogeneity beneath seismic arrays in central California and Yellowstone, Wyoming, Ph.D. Thesis, Massachusetts Institute of Technology, 1978.

Figure Captions

Figure 1. Fault trace and aftershock epicenters of the Parkfield earthquake of 1966, reproduced from Eaton et al. (1970). Not only the surface trace of the fault, but the fault plane at depth, identified by the aftershock zone, jumps from one side of Cholame Valley to the other. The initial fault plane is clearly terminated at the jumping point. Two lines fitting the two zones of aftershock epicenters are drawn by the author.

Figure 2. Projection of hypocenters of the Parkfield aftershocks on the vertical fault plane, reproduced from Eaton et al. (1970). The hypothetical boundary between the slipped section (region with no aftershocks) and the unbroken barrier (region with aftershocks) is sketched by the author. The ends of the zone of surface fracturing are indicated by arrows and the approximate position of the main shock epicenter is shown by the vertical dotted line.

Figure 3. The wine cellar at the Morrell ranch located close to the fault trace. The upper portion is thrown to the northeast, directly away from the fault line. Reproduced from Lawson (1908).

Figure 1

Figure 2

Figure 3

Estimation of Non-Hydrostatic Stress in the Earth
by Seismic Methods: Lithospheric Stress Levels
Along Pacific and Nazca Plate Subduction Zones

by

Charles Archambeau*

*University of Colorado
CIRES
Boulder, Colorado 80309

I. Introduction

Reliable estimates of the earth's non-hydrostatic stress field as a function of position would provide, in the broadest sense, important constraints on dynamical processes taking place within the planet. Smith and Toksoz (1972) for example, have provided illustrations of stress field orientations, magnitudes and distributions within subducting plates as functions of the dynamics of plate motion and the rheology of the surrounding mantle, these results showing that an independent determination of the stress field could be used to infer mantle rheology and perhaps the dominant forces acting to cause subduction of lithospheric plates.

In addition, of course, knowledge of the stress field spatially within the earth would be useful for prediction of earthquakes, certainly in the sense that extensive regions of high stress indicate zones in which a large earthquake is possible. Further, it seems reasonable to expect that the magnitude and distribution of the stress field could provide a basis for both the estimation of the size and type of the largest event to be expected and the ground motion to be associated with this "maximum probable" event. Coupling stress data with seismicity time and spatial patterns might also be a key to the understanding of the entire sequence of phenomena leading to a large earthquake.

The principle and presently the only methods that can be applied to estimate stresses at depth are indirect seismic methods. The most practical

and widely applicable is the estimation of stress drops⁺ by "inversion" of the observed seismic radiation field from earthquakes for source properties, and this approach has been considered in a number of investigations in the past, for example, by Wyss and Brune (1968, 1971), Wyss (1970a,b), Hanks and Wyss (1972), Molnar and Wyss (1972), Hanks and Thatcher (1972), Thatcher and Hanks (1973) and Wyss and Molnar (1972a,b). This series of investigations have utilized a "stress pulse" model of an earthquake, the particular model applied being that devised by Brune (1970, 1971), wherein the source is described in terms of an application of stress at the failure zone boundary equal to the stress drop occurring as a result of failure. This model was then used to provide a fit to observed far field seismic spectra from earthquakes, after correction for propagation effects, by variation of the magnitude of the stress drop and rupture dimensions.

In the present study relaxation source theory models are used, the theory having been described most recently and fully by Archambeau and Minster (1978), while the particular models to be used have been described by Archambeau (1968, 1972) and in most detail by Minster (1973). A somewhat different approach

⁺The term "stress drop" used here and generally in the geophysical literature refers to the change in the stress field due to failure, measured on the surface bounding the failure region. Thus the term refers only to the change in stress level along the failure surface rather than to the volumetric stress changes that, of course, occur. In general this stress change may be expected to be a complicated function of position and so the term "stress drop" generally refers to the average value of the stress change over the failure surface. Further, this stress change could well be a rather complex function of time, probably with the (dynamic) stress change being complex in time and possibly large immediately after failure, with subsequent stress build-up with time corresponding to a recovery of strength of the failed material. Equally possible is a low stress drop immediately upon failure at a point followed by a slow viscous-like relaxation of stress with time. In this study, relatively high frequency seismic radiation is used to estimate stress changes, so the stress drop values refer to the short time changes in stress.

will be adopted for the "inversion" of observational data than has been the usual practice. In particular, synthetic seismograms are generated incorporating appropriate propagational effects directly and then these theoretical results are processed in the same manner as the observed data. For example, both theoretical and observed seismograms are narrow band filtered to obtain spectral amplitude levels for particular energy arrivals, using the same filters for both, so that effects of interfering signals are represented in both results. Similarly, m_b (body wave magnitude) and M_S (surface wave magnitude) data are jointly used to infer source parameters and the theoretical M_S versus m_b magnitude curves used as the interpretational basis for the data are generated by the same procedures used to obtain observational magnitude data.

The procedure to be used in this study is to consider a very large number of small to moderate sized earthquakes and to obtain an average stress drop and rupture length estimate for each event. For a spatially dense distribution of events, then these stress estimates can be used to estimate the spatial variation of tectonic stress and to delineate high stress zones in particular. The only reasonable means of obtaining such estimates for many events from a tectonically active region is to use the easily measured, and therefore abundant, magnitude data for worldwide earthquakes. This data set will therefore be used here and once high stress zones are identified, then more detailed (and time consuming) studies of the events within these regions can be made.

There are a number of fundamental uncertainties connected with the estimation of tectonic stress by seismic methods in general and some additional

problems associated with estimations based on event magnitude data in particular. The most basic uncertainties and ambiguities are related to tradeoffs between source variables in any inversion procedure and with resolution of spatial variations in stress drop along the failure zone boundary. In addition, there are the ever present restrictions and possible errors that can be introduced by propagational effects in the earth and by the earthquake model used to provide a theoretical basis for the inversion.

In the next section the problems associated with tradeoffs between intrinsic source variables are more fully defined and discussed along with the stress resolution problems. In the section following that, the source theory and models derived from it are briefly summarized, along with some results from applications and tests of the theory. In all of this the objective is to establish, in a clearly defined fashion, a systematic and easily implemented basis for stress estimates and, as well, to define the limitations and uncertainties inherent in the results to be obtained.

Once this basic framework is established, applications to a number of active seismic regions are considered and inferences based on the stress results obtained are then made in the context of the basic limitations inherent in seismic inversion methods for source variables.

II. Tradeoff and Resolution Limitations for Stress Estimation

Use of earthquakes themselves as a means of inferring existing tectonic stress involves a number of problems that must be addressed before we can use stress drop results obtained for other inferences, such as those involving earthquake prediction.

First of all it is clear that the occurrence of an earthquake can rather drastically alter the initially prevailing stress field, so that the phenomena

that allows us to "sample" the stress field also produces a (complicated) change in the quantity we wish to estimate. Further, it is only the stress field changes that are reflected in the radiation from an earthquake and relating these changes to the ambient stress field is, at best, an uncertain and somewhat arbitrary procedure. For example, we could argue on physical grounds that shear melting should occur during any failure process at depth in the earth, either as a primary or secondary physical phenomena. In this case a relaxation source theory representation of the radiation from an expanding zone of zero shear strength can be used and would allow inversion of the field for the ambient initial stress. However, shear melting, either as a consequence or cause of failure, is presently only a plausible hypothesis in spite of the arguments we might make involving high temperatures and confining pressures tending to favor it. Further, whether shear melting occurs depends on how large the tectonic shear stresses are in the earth, as was pointed out by McKenzie and Brune (1972), and it is therefore only logical to attempt to determine tectonic stress changes without assuming shear melting and to then use this information to deduce whether shear melting is possible. If melting is then shown to be quite certainly involved, then the stress changes obtained using a relaxation theory earthquake model can be used to deduce the ambient tectonic stress, using the same theory. Of course, work in experimental rock mechanics could provide independent confirmation of the reality of shear melting during failure. In this regard recent experimental work by Friedman, Logan and Rigert (1974) and by Teufel (1976) indicates very high temperatures and melting at asperities for prefactured rocks over a wide range of confining pressures and temperatures during failure tests. Further, evidence from field observations of fault zones - for example by

Scott and Drever (1954) - also suggest similar local melting occurrences. However, there are a number of uncertainties inherent in the very limited experimental results and in the field observations as well, so that a conclusion favoring melting, while very attractive, does not appear warranted yet.

In order to properly account for the limitations in the use of earthquake radiation fields for estimation of stress - these being chiefly connected with the fact that earthquakes change the ambient stress and it is only the change that can, currently, be estimated with reasonable certainty - it is necessary and appropriate, at least initially, to: (1) use only relatively small earthquakes, that do not cause a widespread change in the stress field, as sampling events for the inference of values for the existing nonhydrostatic stress, (2) regard the stress drop estimates obtained from the radiation field inversion for source properties as the stress which can be released by an earthquake and to correspond to the recoverable part of the ambient stress ("recoverable ambient stress") with some unknown level of residual or non-recoverable stress.

Use of small events occurring in a region should give us a reasonably accurate spatial picture of the existing recoverable stress field, provided we have a large and well distributed event set. Large earthquakes can be expected to extensively alter the recoverable prestress over a considerable spatial region, and by separating the time sequence of small events into subsets before and after large events, we should be able to resolve the stress changes due to the large event in terms of a change in the prestress level indicated by the two time segmented event series.

The use of the concept of recoverable ambient stress, as previously defined, seems clearly necessary if we are to have a useful and accurate appreciation of what is being measured. Thus, whatever the mechanism of failure at a given depth (i.e., at a particular pressure and temperature)

the recoverable stress measured is that part of the ambient stress that can change in the process and the rest of the stored strain energy is unavailable to this particular process. Since the process of failure and material resistance to failure could well be different at different depths, and for different materials at the same depth in the earth, then we can expect changes in the recoverable ambient stress because of failure mechanism changes as well as because of stress and/or material strength changes. However, in any case the meaning of what has been called the recoverable ambient stress is still clear and more important, it is the only stress quantity that we can confidently talk about and use. In particular, if the perturbations of the stress field due to the occurrence of small events are neglected, then the recoverable ambient stress levels estimated from the small event data can in fact be used to predict the energy release from a large event. In this case, because of the spatial sampling obtained when a large set of small events is used, any variations of mechanism, strength, etc. are automatically taken into account and for a large event, possibly involving different failure mechanisms at different locations and certainly different strength and prestress levels over the extent of the large final failure zone, the use of recoverable stress in determining radiation field characteristics would account for all these variations if, as seems likely, the same process and property variations have also operated on the well distributed set of small events occurring shortly before the large event. Furthermore, the recoverable stress levels, when high, should be indicative of impending failure and if the zone of high (average) recoverable stress is extensive, then a large event could be

predicted in terms of its location and probable size.

This approach, utilizing small events and an interpretation of the resulting stress estimates as recoverable ambient stress, should be a reasonable framework for a start in the process of determining the tectonic stress field and in assessing the implications of the results. The focus of this study will therefore be on obtaining stress estimates from seismic radiation field observations. The results will be used to identify areas of high (recoverable) tectonic stress and hence areas of special interest for event prediction work. But, in addition, an attempt will be made to understand the space and time patterns of the stress drops obtained in an effort to determine whether any conclusions or reasonable hypotheses can be advanced regarding failure mechanisms operative, the relationship of the recoverable tectonic stress to the ambient stress and the variability of the material strength and/or the recoverable stress field in space and time. In addition, of course, we will seek to verify the validity of the assumptions that small earthquakes do not drastically alter the recoverable stress and can be used to "sample" the existing stress field, and that the spatial "stress map" so obtained is that appropriate for the prediction of a future large event.

In addition to the care that must be used in sampling the stress field and in interpreting the results in terms of ambient tectonic stress, etc., it is essential to note that average stress drops are obtained by seismic methods and that the implicit averaging process is dependent on both the method and data used.

Thus, whereas the previously discussed limitations and ambiguities in determining absolute stresses were due to tradeoff problems between various physical quantities, the averaging process inherent in seismic data inversion

for source properties is associated, in part, with the resolution of the data. In addition, however, the source models usually assume, either explicitly or implicitly, a uniform initial prestress in the medium and so any fit to observations, using a single one of these models to represent the entire source, is achieved by choosing the best single value for the magnitude of the prestress (i.e., the recoverable ambient stress). Thus a model limitation of homogeneous prestress forces selection of what amounts to an average value of the inferred recoverable stress. What sort of average is obtained then depends on what part of the data is fit and how well it is fit. For example, a fit to low frequency free oscillation and surface wave data will give a value for the stress drop that is an average over either the maximum failure zone dimension or the wavelength of the components of the seismic field, whichever is smaller. In fact of course, a spectral range of data is used, so the average will always turn out to be some (complicated) weighted average. Thus, if higher frequency body wave data is also used, the model will force a selection of yet a different value for the stress estimate, due to the weighting of the average by the high frequency information which is related to stress changes near the hypocenter, rather than to averages over the entire failure zone. Since it cannot be expected that the recoverable ambient stress is uniform, these different averages must be interpreted with very considerable caution, particularly in regard to inferences involving stresses required for failure in the earth.

The difficulty with the homogeneous initial stress model can obviously be overcome, at least in principal, by use of a less constrained source model involving variable stress drop and rupture rate. This can be done in the context of relaxation source theory in a very general manner (e.g., Stevens and Archambeau, 1977), and subsequent studies can use such models. One can,

however, use multiple constant initial stress model results to simulate, in an approximate way, the variable initial stress and rupture rates that are usually involved. In this case the total event is, in effect, viewed as a multiple event, each subevent having constant initial stress and a rupture rate that is also constant, but of different magnitude from each elemental event to the next. The radiated field is then simply obtained by superposing the radiation field contributions from each subevent. Since interaction between the subevents is neglected (in that changes in initial recoverable stress are not coupled from one subevent to the next), this approach is approximate. Nevertheless, it is undoubtedly superior to the use of a single constant initial stress model for the entire event.

In the next section we will discuss some results obtained by Bache, Barker and Archambeau (1978), using a multiple event model as a basis for the determination of source properties appropriate to the San Fernando earthquake. The purpose of the discussion of these results here is to illustrate the nature of the averaging process, by contrasting the stress estimates obtained using a single constant initial stress model with a "multiple event model". Further, the resolution of source properties, especially the stress changes are well illustrated by comparison of the spatial variation of stress changes along the failure zone, obtained using far field data (and the multiple event model), with the stress changes that are required when high frequency near field data is also fit.

The resolution of spatial variations of stress, in particular, is a problem that can be reduced to acceptable proportions for most purposes. This is accomplished, assuming a suitably general source model is used, by use of enough observational data, distributed not only spatially around a source but, most importantly, widely distributed in frequency (or time) as well. The cost of achieving high resolution, to some specified degree, is of course, represented by greater time and effort and this price can be, in fact, very high. Indeed high resolution of stress variations for each event is essentially impractical if large numbers of events are to be studied. Therefore, in this study the approach is to use a constant initial stress model and to obtain average stress estimates using published event magnitude data (m_b and M_S) for a large number of small to moderate-sized earthquakes, with the event set providing a reasonably dense sampling of a tectonically active region. When enough small event stress estimates are obtained and give good spatial coverage, then spatial variations of tectonic stress can be reliably resolved to the degree that large zones of particularly high stress can be identified. Once high stress zones are delineated then events from these regions can be studied in greater detail employing variable stress drop models or multiple event models to fit the observed seismic radiation and to more precisely define the nature of the stress distribution within these critical areas. Application of this overall approach, in the present study at least, will be confined to identifying high stress zones in regions where enough small to moderate event magnitude data is available to give reasonably good resolution of spatial variations of (recoverable) tectonic stress. In addition, however, a high resolution method that is practical for application to reasonably large data sets will be described

and applied to a limited set of events by way of illustration. In later work this method will be used to obtain a more detailed picture of the stress state of the critical zones identified here.

III. Theoretical Basis for Stress Estimation

The theoretical basis to be used for the inversion of the observed seismic radiation field for stress estimates is given in most complete detail by Archambeau and Minster (1978). This theoretical exposition constitutes an extension of previous relaxation source theory formulations, such as that given by Archambeau (1968), in that it incorporates the dynamical conditions for failure growth in the representation theory.

A short summary of the basic governing equations for dynamic failure growth, along with the radiation field representation equations, will be given here in order to provide a basis for discussion of important features for tectonic stress estimation.

In particular, the essential theoretical results describing failure in an initially stressed medium, and the resulting relaxation of the initial stress by radiation processes, are given by conservation of mass, momentum and energy relations within the medium and on its boundaries. In the case of a failure process in a stressed medium, the failure region may be viewed as an expanding zone of material transition with its boundary constituting a moving internal boundary to the surrounding stressed medium, across which material properties change discontinuously. This in effect amounts to a growing internal inclusion and requires readjustment

of the stress field in the surrounding stressed medium since it is no longer in equilibrium. The stress changes are accomplished dynamically, and by elastic radiation if the surrounding material is elastic. The relation governing this radiation process is simply the equation of motion expressing conservation of momentum in the material around the failure zone. This may be expressed as⁺:

$$\mathcal{L}_{\alpha\gamma} u_{\gamma} = \rho f_{\alpha} \quad ; \alpha, \gamma = 1, 2, 3, 4 \quad (1)$$

where

$$u_{\gamma} \equiv (u, u_2, u_3, 0)$$

is a four vector representation of the displacement field in the elastic medium. In this representation the fourth component is the time-like component. Similarly,

$$f_{\alpha} \equiv (f_1, f_2, f_3, 0)$$

$$x_{\gamma} \equiv (x_1, x_2, x_3, t)$$

are four vector representations of the body force density and the space-time coordinates. Clearly, neither u_{γ} or f_{α} have time-like components for elastic continua. Here the "elastic operator" $\mathcal{L}_{\gamma\alpha}$ is:

$$\mathcal{L}_{\alpha\gamma} \equiv \frac{\partial}{\partial x_{\beta}} \left(c_{\alpha\beta\gamma\delta} \frac{\partial}{\partial x_{\delta}} \right) \quad (2)$$

⁺A four dimensional cartesian tensor representation of the elastic equations is used here in order to present results in a compact form. In this representation Greek letters will always run over the range 1 to 4, while latin letters will run from 1 to 3.

with the elastic-inertial tensor $C_{\alpha\beta\gamma\delta}$ expressed in terms of the ordinary elastic tensor C_{ijkl} and the density ρ , as:

$$C_{\alpha\beta\gamma\delta} : \begin{cases} C_{\alpha\beta\gamma\delta} = -C_{ijkl} & ; (\alpha, \beta, \gamma, \delta) = (i, j, k, l) \\ C_{i4k4} = C_{4i4k} = \rho \delta_{ik} & ; i, k = 1, 2, 3 \\ C_{\alpha\beta\gamma\delta} = 0 & ; \text{all other } \alpha, \beta, \gamma, \delta \end{cases}$$

The equations (1) and (2) apply to the medium external to the failure zone.

The motion of the elastic medium, including the motion of the internal failure zone boundary, is determined by the boundary conditions that apply to it, these being also expressions of conservation of mass, momentum and energy. If we include moving boundaries as possibilities, so that the failure boundary internal to the elastic medium is included as one of its "generalized" boundaries, then we find that conservation laws on any boundary, whether it is a surface of discontinuity expanding at some velocity $\underline{U}$ or a material boundary moving with the particles of the medium, satisfy the conditions (Archambeau and Minster, 1978):

$$\left[\left[\rho v_{\ell}^* \eta_{\ell} \right] \right]_{\partial V} = 0 \quad (3)$$

$$\left[\left[\tau_{\alpha\beta} \eta_{\beta} \right] \right]_{\partial V} = 0 \quad (4)$$

$$\left[\left(\rho \mathcal{E} v_{\ell}^* - v_k T_{\ell k} + q_{\ell} \right) n_{\ell} \right]_{\partial V} = 0 \quad (5)$$

These relations express conservation of mass, momentum and energy, respectively, on the boundaries ∂V of the elastic medium within the volume V . The double bracket notation $[[\]]_{\partial V}$ applied to any variable denotes the change, or "jump" in the variable across the boundary ∂V . Thus $[[f]]_{\partial V} \equiv f_1 - f_2$; where f_1 and f_2 are the limit values of the variable f when the boundary ∂V is approached from opposite sides.

The variables appearing in (3)-(5) are:

$$\tau_{\alpha\beta} \equiv C_{\alpha\beta\gamma\delta} \frac{\partial u_\gamma}{\partial x_\delta}; \text{ the generalized stress tensor}$$

$$n_\ell = (n_1, n_2, n_3); \text{ the normal to the boundary } \partial V$$

$$\eta_\beta \equiv (n_1, n_2, n_3, -U_\ell^* n_\ell); \text{ the four vector "normal"}$$

$$U_\ell^* \equiv U_\ell - v_\ell, \text{ the boundary velocity relative to the particle velocity } v_\ell$$

$$v_\ell^* \equiv v_\ell - U_\ell = -U_\ell^*; \text{ the particle velocity relative to the boundary velocity } U_\ell.$$

$$\rho \mathcal{E} \equiv \rho U + \rho/2 v_\ell v_\ell + \rho \phi; \text{ the total energy, with } \rho U \text{ the internal energy, } \rho \phi \text{ the potential energy and } \rho/2 v_\ell v_\ell \text{ the kinetic energy.}$$

$$T_{\ell k} \equiv C_{\ell k m n} \frac{\partial u_n}{\partial x_m}; \text{ the cauchy stress tensor}$$

$$q_\ell \equiv \text{the heat flux vector}$$

When a boundary of V is an ordinary material boundary, so that the boundary moves with the material particles in the direction normal to the surface ∂V , then $U_\ell n_\ell = v_\ell n_\ell$ and $U_\ell^* n_\ell = -v_\ell^* n_\ell = 0$. In this case

$\eta_\beta = (n_1, n_2, n_3, 0)$ and η_β corresponds to the ordinary spatial normal to the boundary surface ∂V . The equations (3)-(5) then reduce to the regular boundary conditions for an elastic continuum.

If some part of the boundary of the elastic medium is not, however, a material boundary but is, say, an external boundary of a failure zone, then U_ℓ^* does not vanish and then the equations (3)-(5) apply in their general form. In particular they are the dynamical conditions that apply to the boundary growth during failure.

The Greens function integral equation corresponding to (1) for a stressed elastic medium with both material boundaries and one (or more) moving failure boundaries internal to it is given by Archambeau and Minster (1978) as⁺:

$$\begin{aligned}
 4\pi \tilde{u}_m(\underline{r}, \omega) = & \int_{-\infty}^{+\infty} e^{-i\omega t_0} dt_0 \int_{V(t_0)} \rho f_\ell(\underline{r}_0, t_0) \tilde{H}_\ell^m(\underline{r}, \underline{r}_0; \omega) d^3 x_0 \\
 & - \int_{-\infty}^{+\infty} e^{-i\omega t_0} dt_0 \int_{\partial V(t_0)} \left[\tilde{H}_\ell^m(\underline{r}, \underline{r}_0; \omega) \left\{ T_{\ell k}(\underline{r}_0, t_0) - \rho \frac{\partial u_\ell}{\partial t_0} U_k \right\} \right. \\
 & \left. - u_\ell(\underline{r}_0, t_0) \left\{ \tilde{q}_{\ell k}^m(\underline{r}, \underline{r}_0; \omega) + i\omega \rho \tilde{H}_\ell^m(\underline{r}, \underline{r}_0; \omega) U_k \right\} \right] n_k da_0 \\
 & + i\omega \int_{-\infty}^{+\infty} e^{-i\omega t_0} dt_0 \int_{V(t_0)} \left[\rho \frac{\partial u_k^*(\underline{r}_0, t_0)}{\partial t_0} \right] \tilde{H}_k^m(\underline{r}, \underline{r}_0; \omega) d^3 x_0
 \end{aligned} \tag{6}$$

⁺This equation (6) is actually the Fourier transform with respect to t , the "observers time", of their results. This integral equation is the linearized equivalent of the equation set (1) plus (3) through (5). The linearization involves replacing U_ℓ^* by U_ℓ in the boundary conditions on the failure surface and amounts to neglecting the particle velocity relative to the growth rate of the failure surface. For earthquakes this is certainly a reasonable approximation.

Here $\tilde{u}_m(\underline{r}, \omega)$ represents the Fourier transform of the displacement field defined by

$$\tilde{u}_m(\underline{r}, \omega) \equiv \mathcal{F}_t \{u_m(\underline{r}, t)\} = \int_{-\infty}^{+\infty} u_m(\underline{r}, t) e^{-i\omega t} dt$$

The tensors $\tilde{H}_\ell^m$ and $\tilde{g}_{\ell k}^m$ are the Fourier transforms of Greens tensors corresponding to "propagators" or transfer functions for the medium, with

$$\tilde{g}_{\ell k}^m = c_{\ell k i j} \frac{\partial}{\partial x_i} \tilde{H}_i^m$$

representing the stress associated with the Greens displacement $\tilde{H}_i^m$.

The displacement function $u_k^*(\underline{r}_o, t_o)$ appearing in the final integral term in (6) corresponds to the changes in the equilibrium field due to the creation and growth of an inclusion or failure zone within the initially stressed medium. This integral term is zero if the initial stress is zero, but in the present problem it represents the stress relaxation contributions to the radiation field. For an expanding failure zone with a boundary denoted by Σ ,

$$u_k^*(\underline{r}_o, t_o) = u_m(\underline{r}_o, t_o) - u_m^{(o)}(\underline{r}_o)$$

where $u_m^{(o)}$ is the initial equilibrium displacement field in the medium without a failure zone and $u_m(\underline{r}_o, t_o)$ is the equilibrium field appropriate to a failure zone (inclusion) with a surface $\Sigma(t_o)$ at a (source) time t_o . As the failure region expands with time the equilibrium field appropriate to any instantaneous configuration of the inclusion must be

different than it was previously and hence $u_m^{(1)}$ will depend on the time parameter t_o describing the inclusion or failure surface shape.

The analytical expression for $\underline{u}^*$, for any type or form of growing failure zone and medium, can be shown to be (Archambeau and Stevens, 1978):

$$\underline{u}^*(\underline{r}_o, t_o) = \sum_{\ell, m} [\alpha_{\ell m}(t_o) \underline{M}_{\ell m}^-(\underline{r}) + \beta_{\ell m}(t_o) \underline{N}_{\ell m}^-(\underline{r}) + \gamma_{\ell m}(t_o) \underline{F}_{\ell m}^-(\underline{r})]$$

where $\underline{M}_{\ell m}^-$, $\underline{N}_{\ell m}^-$ and $\underline{F}_{\ell m}^-$ are vector solid harmonics. The coefficients $\alpha_{\ell m}$, $\beta_{\ell m}$ and $\gamma_{\ell m}$ are dependent on the inclusion (failure zone) and surrounding elastic zone material properties, the initial stress and the geometry of the failure zone - and therefore also parametrically on t_o , the time parameter for the failure surface Σ determining the inclusion geometry. These coefficients are fixed by solution of the static boundary value problem involving the failure inclusion with surface $\Sigma(t_o)$, along with prescribed stress at infinity, corresponding to the initial stress field (see for example Landau and Lifshitz, 1959). For complex geometries and initial stress they can be obtained numerically (Stevens and Archambeau, 1977).

It is worth pointing out that while these coefficients depend on the initial stress, they also depend on the properties of the material within the inclusion or failure, among their parameters. It is, primarily, this dependence on two unknowns that gives rise to the difficulty of "inverting" the observed dynamic field $\tilde{u}_m$ to obtain the true (ambient) initial stress, for in fact there is a one to one tradeoff between the initial stress value and the parameters representing the failure zone material properties. As noted earlier however, if we knew what rheology was appropriate to the failure zone material then it would be possible to obtain initial stress estimates and conversely. In general of course, we

know neither and so are forced to define what has been called the "recoverable initial stress" which in effect takes into account this tradeoff.

We note that the final or relaxation integral term in (6) depends on the time derivative of u_k^* . This is as would be expected since we cannot specify displacements absolutely, which means that we should be able to add arbitrary functions of r_0 to u_k^* without affecting the dynamical results, and this is clearly the case.

The second integral term in (6) represents the medium boundary interaction effects, which includes interactions - such as wave reflections and diffraction - from the growing failure boundary as well as the fixed boundaries. However, on the fixed (i.e., material) boundaries, the boundary velocity, with components U_k , is zero to the linear approximation used in obtaining (6). Therefore the factor involving U_k in the surface integrals over fixed boundaries is not present. It is, however, present for the growing failure zone boundary and represents the effects of energy absorption by the material transition process (failure process) occurring at the failure surface. This becomes apparent when one considers that the value of boundary velocity U_k appearing in the integrand is controlled by the energy conservation relation (5). Thus the energy absorption at the boundary will, in part at least, serve to determine the value of the boundary velocity U_k appearing in the equation. This also means that U_k is determined by the nonlinear processes at the boundary, and the properties of the material after failure, as well as before failure, via the energy boundary relation (5).

The first integral term in the integral equation for $\tilde{u}_m$ is due to the action of the time dependent variations of external forces (e.g. gravity)

acting on the medium. Since we are considering a spontaneous process here, then to first order the external force effects are time independent and negligible, so this term is usually not included in calculations for the direct seismic radiation from the failure process. (There are, however, second order effects due to the small time dependent changes of density and gravity, due to changes in the strain field in the region around an earthquake, which are the neglected effects when this integral term is ignored.)

The Greens function $H_{\ell}^m(\underline{r}, t; \underline{r}_o, t)$ appearing in the integral equation is generated from the field equations with a generalized delta function force term and with homogeneous boundary conditions in the medium without a failure zone and without prestress. That is, H_{ℓ}^m is obtained from:

$$\left. \begin{aligned} \mathcal{L}_{\ell k} H_k^m &= 4\pi \Delta_{\ell}^m \\ \Delta_{\ell}^m &\equiv \delta_{\ell}^m \delta(\underline{r}-\underline{r}_o) \delta(t-t_o) \end{aligned} \right\} \quad (7)$$

with the boundary conditions

$$\left. \begin{aligned} \left[\mathcal{H}_{\ell k}^m n_k \right]_{\partial V_o} &= 0 \\ \left[H_{\ell}^m \right]_{\partial V_o} &= 0 \quad \text{or} \quad \left[\frac{\partial}{\partial t} H_{\ell}^m n_{\ell} \right]_{\partial V_o} = 0 \end{aligned} \right\} \quad (8)$$

applying on boundaries ∂V_0 , where V_0 does not contain moving (failure) boundaries; that is it represents the whole medium before failure and in its unstressed state. The first of the conditions in (8) expresses continuity of tractions on material boundaries that are "welded" solid-solid interfaces, or continuity of the normal component of velocity if the boundaries are not welded or fluid-solid interfaces.

The Greens displacement tensor $H_{\ell}^m(\underline{r}, t; \underline{r}_0, t_0)$ is, by inspection of (7), the (ℓ) component of displacement in the medium at the position $\underline{r}$ and time t due to the (m) force component of the vector delta function source acting at the time t_0 and at the point $\underline{r}_0$. Thus H_{ℓ}^m is a two point displacement tensor that depends on the "receiver" position and time $(\underline{r}, t)$ and the "source" position and time $(\underline{r}_0, t_0)$. It simply represents the impulse response of the medium and includes reflections etc., from all fixed medium boundaries.

It is not difficult to show that $\tilde{H}_j^k$ can be expressed in terms of an eigenfunction expansion for a spherically layered earth in the form:

$$\tilde{H}_j^k(\underline{r}, \underline{r}_0; \omega) = \sum_{\ell, m, n} \left[\frac{\psi_k(\underline{r}; n \omega_{\ell}) \bar{\psi}_j(\underline{r}_0; n \omega_{\ell})}{\omega^2 - n^2 \omega_{\ell}^2} \right]$$

where the ψ_k are components of eigenvectors for spheroidal and toroidal oscillations. The eigenfunctions have the analytical form:

$$\psi_k^S(\underline{r}; n \omega_{\ell}^S) = \left[D_{\ell m}(r; n \omega_{\ell}^S) \underline{P}_{\ell m}(\theta, \phi) + E_{\ell m}(r; n \omega_{\ell}^S) \underline{B}_{\ell m}(\theta, \phi) \right] \cdot \hat{e}_k$$

$$\psi_k^T(\underline{r}; n \omega_{\ell}^T) = \left[F_{\ell m}(r; n \omega_{\ell}^T) \underline{C}_{\ell m}(\theta, \phi) \right] \cdot \hat{e}_k$$

where ψ_k^S denotes spheroidal and ψ_k^T toroidal eigenvector components, with $\hat{e}_k$ a unit vector in the (k) coordinate direction. Here ω_ℓ^S and ω_ℓ^T are the eigenfrequencies for the two types of oscillations and $P_{\ell m}$, $B_{\ell m}$ and $C_{\ell m}$ are vector spherical harmonics. The radial functions $D_{\ell m}$, $E_{\ell m}$ and $F_{\ell m}$ can be expressed in closed analytical form in terms of spherical Bessel functions of r , with coefficients that are explicit products of transmission matrices (i.e., Haskell-Thompson matrices). This means then, that the Greens function and its associated stress tensor appearing in (6) can be obtained for a specified layered earth model in closed analytic form, so that the integrals in (6) can be evaluated analytically as well.

In this regard, the direct source radiation due to relaxation of stress is given by the final integral in (6) and the interaction of the failure boundary and the relaxation derived radiation is given by the second (surface) integral taken over the growing failure zone boundary. Neglecting the first integral as small, then the total displacement field may be written as:

$$\tilde{u}_m(\underline{r}, \omega) = \tilde{u}_m^I(\underline{r}, \omega) + \tilde{u}_m^{II}(\underline{r}, \omega) \quad (9)$$

where $\tilde{u}_m^I$ represents the direct relaxation field contribution given by the last integral and $\tilde{u}_m^{II}$ the boundary interaction integral. Using the definition and notation for an inner product over a space (e.g., a two dimensional surface or a three dimensional volume) and also noting that the time integral over t_0 is in the form of a Fourier transform with respect to t_0 , then we can express these fields compactly as:

$$4\pi \tilde{u}_m^I = i\omega \oint_{t_0} \{ \langle \rho \partial_{t_0} u_\ell^*, \tilde{H}_\ell^m \rangle_V \} \quad (10)$$

and

$$4\pi \tilde{u}_\mu^{II} = - \oint_{t_0} \{ \langle \tau_{\alpha\beta} \eta_\beta, \tilde{H}_\alpha^\mu \rangle_\Sigma - \langle u_\alpha, \tilde{\mathcal{H}}_{\alpha\beta}^\mu \eta_\beta \rangle_\Sigma \} \quad (11)$$

where the terms in (11) have been written in four vector notation. Here η_β is the four vector normal defined previously and

$$\tilde{\mathcal{H}}_{\alpha\beta}^\mu \eta_\beta \equiv \tilde{\mathcal{H}}_{\ell k}^m + i\omega\rho \tilde{H}_\ell^m U_k$$

$$\tau_{\alpha\beta} \eta_\beta \equiv T_{\ell k} - \rho \frac{\partial u_\ell}{\partial t_0} U_k$$

Here the volume of the elastic medium surrounding the failure zone, V , depends on the (source) time t_0 , as does the boundary of the failure zone Σ , so that the transform $\oint_{t_0}$ operates on the spatial integral limits $V(t_0)$ and $\Sigma(t_0)$ as well as on the integrands of the spatial inner products. This means, that unless we wish to consider special cases or make approximations of the exact representations in (10) and (11), this is as far as it is possible to go in this formal manner - except for substitution of the known analytic forms for u_ℓ^* , $\tilde{H}_\ell^m$ and $\tilde{\mathcal{H}}_{\ell k}^m$ in (10) and (11). These latter substitutions do in fact give (10) and (11) more explicit form for computational purposes, but do not basically reduce either (10) or (11) any further.

It is therefore at this point where approximations are introduced in order to obtain representations for the field $\tilde{u}_m$ in (9). Ordinarily we term the whole set of approximations, special assumptions and simplifications an

earthquake model. There are, of course, many models but just one theory, which is expressed by (9) through (11). Those that are used for the earthquake models applied in the present study can be enumerated as follows:

(1) "The transparent boundary approximation".

In this case the boundary interaction term is neglected, so that in effect the relaxation induced radiation does not "see" the boundary of the failure zone at all, hence the terminology "transparent boundary approximation". In this case $\tilde{u}_m$ is approximated by $\tilde{u}_m^I$ to first order, so:

$$\tilde{u}_m \approx \tilde{u}_m^I$$

A correction, of second order, to this would involve using $\tilde{u}_m^I$ in equation (11), in the integrals on the right, to obtain an approximate value for $\tilde{u}_m^{II}$, and then adding it to the previously determined $\tilde{u}_m^I$ value for an approximation of $\tilde{u}_m$ to second order. This latter iteration procedure is not, however, applied in the theory to be used in this study. It has, however, been demonstrated by Burridge (1975), Kayama et al. (1973) and Minster and Suteau (1976) that the boundary interaction effects are usually quite small over the entire frequency range of the radiation field.

(2) Special geometric approximations: "Expanding and translating spherical inclusion geometries, with uniform initial stress".

In order to simultaneously be able to handle the time dependence of the volume limit V in (10) and to be able to analytically evaluate the vector harmonic coefficients in the expression for $\underline{u}^*$ in the integrand, it is a practical necessity to consider some form of spherical inclusion geometry and uniform initial stress. Thus for an entirely analytical representation for $\tilde{u}_m^I$, it has been usual to consider

expanding or translating spherical zones as geometric models of the growing failure zone. In addition initial stress is used (i.e., the nonhydrostatic stress components are constants, but not necessarily equal to each other). Approximations designed to obtain representations of nonuniform stresses have also been attempted (e.g., Archambeau, 1968), but are difficult to justify (Archambeau, 1975).

In this study both expanding and translating spherical inclusion zones are used, with the translating model used for stress estimates from the magnitude data. In the latter application, a uniform initial stress is also used. The principal effect of adopting a spherical inclusion zone model for an earthquake is to obtain a larger energy release than would be the case for a thin (ellipsoidal) inclusion zone model, which is more suitable for earthquake failure geometries. However, this more efficient energy release can be simply adjusted to give an accurate representation for the energy release induced by a thin failure zone. In particular, Minster and Suteau (1976) have shown that the difference in the radiated spectrum between an expanding spherical inclusion zone and a disk shaped zone is a factor of approximately 3.4, over the entire frequency range. Thus the spherical model spectra can be simply scaled by this constant factor to give spectra representing ellipsoidal shaped inclusion model spectra. Finally it has been found, by direct calculation of the spectral radiation from the expanding spherical model and from a (specially constructed) translating spherical model, that the translating model gives a result which is essentially the same as the spectrum to be expected for a thin ellipsoid over a spectral range that exceeds the range of interest in the present study (i.e., 10 Hz. to .05 Hz.).

That is, the translating sphere model used in the application to the magnitude data in this study yielded a radiation spectrum that was approximately a factor of 3.4 below that for the expanding spherical model.

The assumption of uniform prestress in the model is presently more difficult to deal with in a quantitative fashion. Thus in the use of a uniform prestress for the interpretation of the event magnitude data in this study, it will merely be assumed that the stress is reasonably uniform over the wave length of the lowest frequency involved (.05 Hz). If this is not the case, we obtain a weighted average stress value for the recoverable stress as was previously noted.

A major difference between relaxation models and other earthquake source models, such as dislocation and stress pulse models, is that the former is a dynamical model while the latter group are all kinematical models, in which the time and space variation of the displacement or stress on the failure zone boundary must be specified. In this case, with either the displacement or stress specified on the failure boundary, then a Greens function surface integral term (like $\tilde{u}_m^{II}$) can be used to propagate this specified field throughout the medium. In the dynamical relaxation theory models however, once the initial equilibrium stress field is specified and the material properties before and after failure are specified (including a failure condition, which simply states what value of strain energy density is required to initiate failure), then the time and space variations of the dynamic fields (displacement, stress, etc.) are determined everywhere in the medium surrounding the growing failure zone, including along the failure boundary itself. This is accomplished by the integral equation (6) or (9), with $\tilde{u}_m^I$ being the true source term and $\tilde{u}_m^{II}$, as noted, a scattering intergral term.

To illustrate the dynamical nature of the representation given by (9)-(11) or (6), and to summarize the important conditions for dynamical failure growth contained in the relaxation theory, consider the boundary conditions (3)-(5) on the failure surface Σ . These conditions show up explicitly in the integral equation (6), for the dynamic field, in terms of the presence of the rupture velocity components U_k in the integrals, as well as implicitly in terms of the time dependence of the failure surface Σ and the elastic volume V , over which the integrations are taken. These latter, of course, depend on time (t_0) because of the failure growth, which is in turn controlled by the dynamic failure conditions contained in (3)-(5). In particular, it has been shown by Archambeau and Minster (1978) that these three conservation relations can be combined to give the following conditions for dynamic failure growth:

$$\rho [v_k] U_R = - [t_k] ; \quad k = 1, 2, 3 \quad (12)$$

$$U_R^2 - \{ [q_k n_k] / (\rho L) \} U_R - [t_k t_k] / (2\rho^2 L) = 0 \quad (13)$$

The first of these is obtained by combining the conservation of mass (3) and momentum (4) conditions and the final relation is obtained from the energy conservation relation (5), using (12). The bracket notation is used to represent jumps across the boundary Σ , but with the boundary designation suppressed, so it is to be understood. Here also:

$$[[v_k]] = v_k^{(1)} - v_k^{(2)} ; \quad \binom{1}{2} = \text{Medium } \binom{\text{exterior}}{\text{interior}} \text{ to the failure boundary } \Sigma$$

$$t_k \equiv T_{\ell k} n_{\ell} = \text{traction component}$$

$$U_R \equiv U_k n_k = \text{Rupture velocity (failure velocity in the normal direction to } \Sigma)$$

$$\rho \equiv \rho^{(1)} = \text{density in the elastic medium surrounding the failure zone}$$

$$[[q_k n_k]] = \text{change in the heat flux normal to the failure boundary}$$

$$L \equiv [[U]] = \text{change in the internal energy of the material upon failure}$$

(difference in the internal energy of the material across the failure boundary).

In this context we treat L as a material variable, much like a phase transition energy or latent heat, so that it is a characteristic of the material, independent of the dynamical variables.

From the energy equation (13) we see that the roots of the quadratic in U_R are real if:

$$[[q_k n_k]]^2 + 2L[[t_k t_k]] \geq 0 \quad (14)$$

This then is the condition that must be met if changes in the failure boundary are to occur - that is, it is the condition for new boundary creation. Since the traction changes would be expected to be much larger than the changes in the heat flux, especially for a fast, high stress process like an earthquake, then to a good approximation this reduces to the condition:

$$L \geq 0 \quad (15)$$

or,

$$L = \llbracket u \rrbracket = T \llbracket S \rrbracket + \llbracket W_{el} \rrbracket \geq 0$$

where $\llbracket S \rrbracket$ is the entropy change and $\llbracket W_{el} \rrbracket$ is the change in elastic energy, that is the reversible work available in the process. Here T is the temperature. Thus $\llbracket S \rrbracket$ is the change in entropy in going from the material (1), the unfailed state, to state (2), the failed state. Thus $\llbracket S \rrbracket = S^{(1)} - S^{(2)}$ and since state (2) is less ordered than state (1), then $S^{(2)} > S^{(1)}$ and $\rho T(S^{(2)} - S^{(1)})$ is the magnitude of the energy required to disorder the state (1) into the failed state (2). Thus we see that the energy condition for failure is

$$\llbracket W_{el} \rrbracket \geq -T \llbracket S \rrbracket ; \llbracket S \rrbracket < 0 \quad (16)$$

so that there must be excess strain energy in the elastic medium at the failure point in the amount $-T \llbracket S \rrbracket$, where $\llbracket S \rrbracket$ corresponds to the "disordering entropy".

A second important consequence of the energy condition (13) is that when the heat flux changes are neglected compared to traction (strain energy) changes, then (13) gives:

$$U_R = \left\{ \llbracket t_k t_k \rrbracket / (2\rho^2 L) \right\}^{1/2} \quad (17a)$$

If we define the traction change magnitude by

$$\left| \llbracket \underline{t} \rrbracket \right| \equiv \sqrt{\llbracket t_k t_k \rrbracket}$$

then we have

$$U_R = \left| \left[\frac{t}{\rho \sqrt{2L}} \right] \right| \quad (17b)$$

Therefore the dynamical conditions for failure growth are that:

$\left[W_{el} \right] \geq -T \left[S \right]$ is required, and when this condition is met, then the local rate of rupture growth is given by (17a) or (17b). The latter relations state that the rupture velocity is proportional to the magnitude of the dynamic traction change across the boundary.

In the elastodynamic source theory the condition (17) has an important role in that when the rupture rate is so specified, then the time dependent failure surface configuration can be calculated and the volume of the surrounding elastic zone V as well. Thus all of the failure dependent variables in the integral equation (6), namely $U_k n_k \equiv U_R$, $\dot{\gamma}(t_0)$ and $V(t_0)$ can be specified by (17) in terms of the local dynamic traction magnitude and the material properties, in particular the transition energy L .

In this regard, solutions of (6) incorporating (17) can be obtained using (9)-(11) and the iterative approach discussed under the transparent boundary approximation. That is, the known equilibrium displacement field u_m^* could be used to calculate an equilibrium value of the traction change across the inclusion boundary and this used in (17), in place of the dynamic traction change, to obtain an estimate of U_R for an assumed transition energy L ; where assumption of L implies an assumption of some particular physical process of failure in the material (e.g., shear melting).

With this estimate for U_R the time dependent elastic volume $V(t_0)$ (and Σ) could be calculated and this would allow the relaxation field integral term for $\tilde{u}_m^I$ to be computed. Then U_R could be recomputed using the field $\tilde{u}_m^I$ instead of the equilibrium field, followed by recomputation of Σ and V and these then used to compute the boundary interaction integral for $\tilde{u}_m^{II}$, using $\tilde{u}_m^I$ in the integral as before. This process could be continued to obtain higher accuracy of course. In any case, it can be seen that the dynamical problem can be solved in its entirety.

In this study, source models incorporating the iteration procedures described were not used, but instead (17) was employed to constrain the rupture velocity. In the modeling by Bache et al. (1978), to be described briefly, this was accomplished by requiring that U_R be proportional to the initial equilibrium stress. This is clearly an approximation since U_R is proportional to the dynamic stress change rather than to the equilibrium stress. However, just as for the iteration procedure outlined earlier, this must be a reasonable first approximation. In the translating sphere model used to generate theoretical m_b and M_S results for the data interpretations of this study, the condition (17) was used in the form of a result obtained by Archambeau and Minster (1978). It was shown that if the shear tractions were negligible within the failure zone (near total loss of shear strength after failure) then (17) predicts that the maximum magnitude of U_R is near the local (elastic zone) shear velocity, V_S . Consequently in the models, in which homogeneous initial shear stress was also assumed, the rupture rate was taken to be $.8 V_S$.

In terms of the implications of (17) for the mechanisms of failure in the earth, that is the value of L , the "latent" internal energy change characteristic of the process and the material, it is interesting to note (and perhaps important to keep in mind) that $U_R \sim 3 \times 10^5$ cm/sec and $\rho \sim .3$ gm/cm³ for earthquakes and for shear melting $L \gtrsim 10^7$ ergs/gm (for most materials $L > 10^8$ ergs/gm in fact, but for weakened silicates this could be considerably reduced), then (17) gives for the dynamic stress changes at a rupture front:

$$\left| \left[\underline{t} \right] \right| = \rho U_R \sqrt{2L} \geq 4.5 \times 10^9 \text{ dynes/cm}^2$$

Thus shear melting can occur if the dynamic traction changes are of the order of 5 kb at the rupture front. In the present context this means that if we observe static stress changes (i.e. recoverable ambient stress changes) that are of this order, or even of the order of 1 kb, then shear melting becomes a very plausible process since the dynamic stress changes in the vicinity of the high curvature rupture front could easily be a factor of 5 to 10 times the initial stress level, which must itself be equal to or larger than the stress drops that are obtained seismically.

In this regard it is also worth noting that under laboratory conditions, rocks fail when the applied tractions are of the order of 5 kb. The dynamic tractions at the rupture front would be higher than this, so that L would again be greater than 10^7 ergs/gm and most likely in the range 10^8 - 10^9 ergs/gm. These transition energies are probably close to those required to cause local melting of asperties and the like.

IV. Theoretical Relaxation Model Predictions with Comparisons to Observations and Detailed Numerical Models

The comparison of results from relaxation theory models with detailed numerical models, wherein it is possible to model thin ellipsoidal or planar failure zones and to include a variety of nonlinear effects that are likely to be involved in failure, can be used to check that the various approximations used in obtaining the analytical theory results do not result in major deviations from "reality". In particular, it is important to gain additional verification regarding the constant scaling (by the 3.4 factor) of the spherical model spectra to the thin ellipsoidal geometry. In addition, there is always some uncertainty concerning the importance of nonlinear effects, within the high stress zone near the front of the advancing failure region, on the seismically radiated energy. In the analytical relaxation models this high stress zone is treated as being elastic up until the instant of failure, whereas it is nearly certain that some plastic deformation occurs at some stage. Thus, it is important to see if a numerical model, with plastic behavior possible at high stress just prior to failure, gives results comparable with the analytical model predictions.

Such a comparison was made by Cherry, Bache and Masso (1976) and the results are shown in Figure 1. The numerical model was a three dimensional model with plastic behavior allowed. The criteria for failure was an energy threshold condition for the amount of plastic work. Specifically when the plastic work at the failure front reached a threshold level the material was allowed to fail, so that a failure zone with static and dynamic frictional surface characteristics was created continuously. In this manner the growth of the rupture was allowed to proceed at its own rate in uniformly prestress

Figure 1. Far-field displacement spectra at $R = 1.5$ km, $\phi = 15^\circ$, $\tau = 30^\circ$, for the finite difference model compared to the analytical relaxation source model. The diameter of the growing spherical volume is indicated at five equally spaced time steps. The unilateral rupture indicated was modified to represent the bilateral rupture geometry used in the finite difference model to give the spectral comparison shown.

medium. The comparison of the numerical model results was with a "tangentially expanding spherical relaxation source" model, which is indicated schematically. Since the relaxation model was a unilaterally expanding model (one point fixed), two such models expanding in opposite directions were used to simulate the bilateral numerical model. All the failure parameters (prestress, rupture rate, etc.) in the two models were the same, except that the numerical model was planar rather than having a thickness like that of the relaxation model. After adjustment for the constant scaling value of 3.4 obtained by Minster and Suteau (1976), the spectra from the two models are quite close to each other, as shown in Figure 1. In particular, the relative levels of P, SV and SH are essentially identical for the two models, and the absolute levels are very close over most of the frequency range. The deviation near the corner frequency, where the numerical model spectrum is somewhat lower than the analytic spectrum, may be due to the effects of plasticity in the numerical model. This however cannot be concluded with any degree of certainty until more numerical models with variable plastic effects have been investigated. Further, the numerical calculations are affected by high frequency numerical noise, due to the finite grid used, and this may account for the deviation at high frequencies.

Clearly however, considering the approximations inherent to the analytical model and the problems with numerical noise, the agreement between the two models is remarkable. One can in fact conclude that the scaling from spherical to thin ellipsoidal or planar failure zone geometry is verified, and that the effects of nonlinear behavior are not great.

The application of the source theory to observed data is perhaps best illustrated by the study of the San Fernando earthquake by Bache et al.

(1978). This work is a good example of the systematic inversion of seismic data for source properties and, in particular, for spatially variable stress drop associated with earthquakes. It illustrates in particular, the necessity of multiple sub-events and spatially variable stress drop and rupture velocity if the detailed properties of earthquakes are to be resolved.

Figure 2 shows, schematically, the results (stress drop, rupture "plane" orientation, etc.) of fitting the observed far field radiation from this event. The event was modeled as a double event, with failure initiating at 11.4 km from the surface, rupturing bilaterally at first, for 3 km in both directions from the point labeled (1) in the figure and then unilaterally upward to the surface, along a zone defined by a plane at 50° to the surface. The stress drop inferred was high (near 750 bars) local to the initiation point and then dropped off, with the rupture velocity also dropping off (see the left hand scale in the lower figure). The rupture rate is taken as being proportional to the stress drop, this being based on the theoretical relations for dynamic failure growth previously discussed. The failure zone was inferred to extend to a region of another high stress concentration, labeled 2 in the figure, and what amounts to the initiation of a second high stress drop "event" occurred with ensuing rupture propagation to the free surface. Figures 3 and 4 show the fits to teleseismic short period observations at numerous azimuths and distances for the model. These records are primarily controlled by the high stress drop segments along the failure plane, and comparison of the fits obtained by using the "double event", rather than only the

San Fernando Earthquake:
Geometry of the failure zone and location, extent and sense of growth for the two "events" used to model the complete earthquake.

Figure 2. Rupture velocity and stress drop plotted as functions of distance along the fault from the hypocenter. Event 2 was started from a finite volume for the failure zone while event 1 started from a point.

Figure 3. San Fernando Earthquake: Theoretical (single event) and observed teleseismic short period vertical ground motion recordings. The gains have been chosen so the first trough is the same size in all traces. The gain ratio (observed/synthetic) is indicated on each record.

Figure 4.. San Fernando Earthquake: Theoretical (double event) and observed displacement at teleseismic distances.

single initial "event", shows that a double event, or highly variable stress drop, is required to fully explain the observations. However, we observe that the first few cycles of ground motion are determined by the high stress drop at the hypocenter.

The low frequency body and surface waves are also fit by this model, but these observations are controlled by the average stress drop over the entire failure plane. Thus, because of the extension of the failure zone due to the presence of the second high stress concentration and the ability of the failure zone to dynamically grow into and through zones of relatively low stress ~~between~~ the high stress concentrations, then the average stress drop for the process was only about 100 bars. If, however, the second stress concentration was not present, then the failure would have stopped and the final failure zone dimension would have been much smaller, resulting in a much lower moment and surface wave magnitude. Since the high frequency body wave magnitude, m_b , is determined more by the hypocentral stress drop, it would remain essentially fixed even if the failure had not continued to grow. The net effect would have been a reduction in M_s without change in m_b .

There is additional evidence, however, that the stress drop very local to the two hypocenters at points (1) and (2) in Figure (2) was even higher than that inferred from the teleseismic "short period" data. This evidence comes from comparison of the high frequency near field results, predicted by the model in Figure (2), with the near field recordings. Figure (5) shows the fit of the model predictions to the observed recording at the Pacoima dam site, located as indicated in Figure (2). While the low

Figure 5. Comparisons of the observed and predicted (dashed lines) velocity and displacement at the Pacoima dam site. Final fault model for the San Fernando earthquake.

frequency character of the observations is quite well fit by the model predictions, the very high frequencies, in the range from 3 cps to around 20 cps are not represented. In view of all the constraints imposed on the model by the far field low and high frequency data, the only way in which this near field "very high frequency" ground motion can be fit is to introduce, spatially confined, very high stress concentrations in the immediate vicinity of the hypocentral points (1) and (2). Preliminary estimates imply, in fact, that these stress levels are near 2 to 3 kilobars and confined to a region of a few hundred meters. This means that earthquake stress drops are, locally at least, much higher than we had hitherto estimated. We will show, in the later sections of this paper, that the study of stress drops based on event magnitude data strongly implies the same conclusion.

As previously discussed, the analysis of observed seismic data for stress estimates will be confined to the use of event magnitude data for large numbers of events. The appropriate way to do this is to generate synthetic seismograms in the time domain and to directly measure magnitudes from the predicted signals at a large number of azimuths and distances and to obtain averages in the same manner as is done in generating observed magnitude data. Since the earthquake data to be analyzed will represent, in general, events occurring in both oceanic and continental tectonic environments, it is necessary to obtain theoretical results for earth structures representing both tectonic continental and oceanic upper mantle characteristics. We have used the continental structures for tectonic areas obtained by Archambeau, Flinn, and Lambert (1969) as well as a modified version of the anelastic Q structure obtained in this same study for the body waves. The surface

wave anelastic structure was adopted from the MM8 model of Anderson and Archambeau (1964). The oceanic velocity model used was based on that given by Toksoz and Anderson (1962), with the Q model modified from the continental model in a manner consistent with the differences between oceanic and tectonic continental mantle velocity distributions. That is, the low Q zone was made to correspond to the low velocity zone in the model.

In addition to the necessity of generating synthetic time series for different mantle structures, it is also clearly necessary to vary the source depth and the failure zone orientation with respect to the free surface, inasmuch as these variations will have important effects on the magnitudes observed. Thus three "types" of events were considered: 45° thrust earthquakes, normal dip slip earthquakes and (90°) strike slip earthquakes. Theoretical seismograms were then generated for these basic event types at five separate depths (10 km, 15 km, 25 km, 35 km, and 45 km) in either an oceanic or a tectonic continental environment. In addition the rupture velocity was varied with depth since all the events used a rupture rate equal to .8 of the local shear velocity, which was different in the two structures used and different at different depths. Finally, of course since we wish to generate theoretical results relating body and surface wave magnitudes to stress drop and rupture zone dimensions, both of these source variables were varied for the different events. In particular, event length dimensions of 1 km, 2.5 km, 5 km, and 10 km were used with each event type, as well as event stress drops of .001 kb, .01 kb, .1 kb and 1 kb. All these variations of structure, event type, depth and failure parameters resulted in the generation of a very large number of synthetic seismograms of course.

Figures (6) through (8) show some examples of these, where variations of the compressional body waves for events of different type at various depths, as well as at different distances and azimuths, are illustrated in Figures (6) and (7). Figure (8) shows typical Rayleigh surface wave seismograms for events at different depths.

The theoretical event magnitudes m_b and M_s were obtained from these synthetic seismograms using the conventional, and rather imprecisely defined, procedure for the measurement and computation of magnitudes. However, only the .05 Hz surface wave magnitude was obtained, in view of the variability of M_s with frequency, and so comparison of the theoretical results to observations requires the use of M_s data observed at .05 Hz. Since this has now become essentially standard observational procedure, this presents no particular difficulties. Similarly, only the vertical component M_s was generated, since this is now the standard measurement, but the radial component M_s value can be easily obtained by multiplication of the vertical amplitudes by the appropriate ellipticity factor.

The theoretical magnitude data generated by this rather massive computational effort is illustrated, in one particular form, in Figure (9) through (11).

These figures show the theoretical body and surface magnitudes for the three basic event types as functions of depth, with event rupture dimension as a parameter, and for the two basic mantle structures used. The stress drop for the events was at a fixed 100 bar level. These three figures together constitute the entire theoretical m_b and M_s set for the events having a fixed 100 bar stress drop. For events with larger or small stress drops the

Figure 6. Theoretical seismograms for five depths and two epicentral distances for a relaxation source model. The faulting is strike-slip and the azimuth is 30° from the strike.

Figure 7. Theoretical seismograms for several azimuths for an earthquake source at two orientations, strike-slip and dip-slip. The source depth is 25 km and the epicentral distance is 4000 km.

Figure 8. Theoretical surface waves from a 45° reversed thrust earthquake at various source depths in a tectonic continental structure (Basin and Range type). Fault length $L = 10$ km, $W/L = 0.83$, $V_R = 0.8V_S$. Distance $\Delta = 4000$ km, $A_z = 30^\circ$. Stress ≈ 700 bars.

Figure 9. Theoretically computed m_b and M_s values for thrust type earthquakes of four different rupture dimensions ($L = 1$ to 10 km), located in different source structures and at variable focal depths.

Figure 10. Theoretically computed m_b and M_s values for normal faulting earthquakes as a function of rupture dimension, source structure and focal depth.

Figure 11. Theoretically computed m_b and M_s values for strike-slip earthquakes as a function of rupture dimension, source structure and focal depth.

magnitude scale is simply shifted linearly. That is, since source spectral amplitude scales linearly with stress drop then for 1000 bar stress drop events the magnitude scale would change by the (base ten) log of the ratio of the 1000 to 100 bar stress drop magnitudes, or by one magnitude unit. Hence for 1 kb stress drop events, the magnitude scale can simply be increased by one unit and the proper predicted m_b and M_s values would be obtained. Hence these figures contain essentially all the magnitude information generated.

The important characteristics of these results, especially for the applications intended, are first that the body wave magnitude shows little variation with event hypocentral depth and essentially no variation between continental and oceanic structures. This is, of course, highly desirable for purposes of source property inversion. Second, as expected, the M_s results shown systematic differences in continental versus oceanic structure and a rather strong variation with depth in the hypocentral depth range 0 to 20 km, in both structures. However the M_s variation with event depth in the range from, somewhat less than, 20 to 50 km is mild and the curves are, in fact, very near flat at a constant M_s level. This is very important for source parameter inversion using M_s data, since it is highly desirable that the data be strongly dependent on only the source parameters of interest (stress and rupture dimension) and essentially independent of depth in particular, since hypocentral locations are usually uncertain to at least 20 to 30%. Thus, we see that because of the rather fortunate circumstance that neither M_s nor m_b values are very sensitive to hypocentral depth in the range from about 15 to 50 km, we can expect to largely avoid uncertainties in stress estimates arising from source depth uncertainties for events in

this depth range. Thus only for events in the 0-20 km depth range is it likely that depth is a critically important factor.

The appropriate form of the theoretical magnitude results is as a set of theoretical curves in a plot of m_b versus M_s value. A set of continuous curves of this sort can be easily generated by interpolating the (m_b, M_s) results obtained, using fault dimension and stress drop as parameters for the curves. The set of Figures (12) through (17) give examples of such curves for 45° thrust and normal dip slip events. The lines labeled 1 km, 5 km, etc., are the loci of events in this M_s vs m_b plane having constant rupture dimension, but variable stress drop. The curves labeled .001 kb, .01 kb, etc., are the loci of events in the magnitude plane with constant stress drop but with variable fault dimension. An important aspect of all the constant stress drop curves is their convergence in the region of low magnitudes. Thus the separation of the "grid" defined by these curves is not constant and, particularly for small magnitude, high stress drop events, the convergence is such that estimates of stress will have large uncertainty even for rather small uncertainties in the observed magnitude values. This is particularly the case for uncertainties in m_b values for events in this magnitude range. For events with depths between 20 and 50 km and with M_s values above about 4.5 however, the stress estimates should be quite good average values with only modest uncertainty due to magnitude data and depth errors and bias (e.g., typically 20 to 30% uncertainties).

In addition to illustrating the nature of the theoretically predicted grid of m_b versus M_s curves, from which estimates of stress and failure dimensions can be obtained, Figures (12) through (17) show magnitude data for the

Figure 12. Surface wave magnitude M_s versus body wave magnitude m_b , showing event data in the depth range 0-10 km from the Aleutian-Alaskan region. The curves are theoretically predicted for thrust type events in this depth range and are used to estimate stress drop and rupture dimensions for the observed events. The solid circles are explosions.

Figure 13. Surface wave magnitude M_s versus body wave magnitude m_b , showing event data in the Alaskan-Aleutian region for the depth range 10-20 km. The theoretical curves are for a 45° thrust event at 15 km with variable stress and dimension.

Figure 14. Surface wave magnitude versus body wave magnitude for events in the depth range 20-33 km. The theoretical curves are for events at 25 km in a continental tectonic type structure.

Figure 15. Data and theoretical event loci (curves) in the $m_b - M_s$ plane, for events in the 33-40 km depth range. The event No. 212 is probably grossly mislocated with respect to depth or a data error.

Figure 16. Data and theoretical event loci (curves) in the $m_b - M_s$ plane, for events in the 40-55 km depth range. Numerous small events appear to have very high stress drops (~ 1 kb) but stress value uncertainties for these small, high stress events is large in general in this depth range.

Figure 17. $m_b - M_s$ data and theoretical curves for dip slip type events in the Alaskan-Aleutian region. Some of the smaller events indicate high average stress drops near 1 kilobar and small failure zone dimensions. The depths of these events are unknown but because of their location on the oceanic side of the trench they are interpreted as being shallow.

Alaskan-Aleutian region (Seismic Region 1). The data has been separated into groupings with respect to event type and depth in order to remove biasing of the source parameter estimates due to differences in the magnitudes for events of different type and at different depths. The distribution of the data (for about 250 events) in the m_b - M_s plane is typical of data world-wide in that in the small to moderate event magnitude range, the average stress drops range from about 1 bar to 2 to 3 kbar with the mean at about 60 bars. The range of event maximum dimension is typically from slightly less than 1 km up to around 30 km, with only the large surface wave magnitude events having dimensions larger than this. The mean event size in this data is about 10 km. As can be seen, nearly all of the events are in the surface wave magnitude range from 4.25 to 6.5 while the m_b values are essentially all below 6.0.

Of all this data, there seems to be only one clearly anomalous event, namely event number 212 in Figure (15).

An explanation of the stress drop range is based on the fact that the values obtained by this method, as previously noted, are average stress drops corresponding to the small of spatial averages of stress changes over the entire failure surface or the average taken over a characteristic dimension of the order of the mean of the longest and shortest wave lengths of the seismic data used. In this case, for m_b and M_s data, this averaging dimension would be of the order of 50 km for events with fault lengths larger than 50 km and of the order of the fault length itself for the small events. We observe from the data distribution in Figures (12) through (17), that most of the very high stress drop events are of small

dimension and that most of the very low stress drop events are associated with large faults. This strongly suggests that stress drop averages over a small fault will be much more strongly influenced by a high stress at the hypocenter and reflect this high stress in the average stress drop inferred from the m_b - M_s approach, while for the large faults the average is over essentially the whole failure region and the high static stress drop occurring at the hypocenter does not influence the average proportionately as much, so a low average is obtained. Figure (18), showing a representative group of event stress drops versus fault dimension shows the trend to high stress drop for smaller faults quite clearly. Note also that the range of stress drops associated with events of fixed dimension is only about one order of magnitude and this range may be due to strength variations or to the fact that the size of local stress concentrations can vary considerably.

These inferences are also consistent with the results of the San Fernando earthquake discussed earlier. In particular, the San Fernando earthquake has an inferred average stress drop of about 80 bars and a fault length of about 15 km based on its observed m_b and M_s values. We saw, however, that the hypocentral stress drop was of the order of 2 to 3 kb. Thus if we were to consider only the first, hypocentral event (1) having a fault dimension of about 3 to 4 km, we would obtain a stress drop of about 600 bars for this sub-event using its m_b and M_s values. In this case, as was noted in the discussion of this event earlier, the m_b value would be the same as that observed for the entire event, but the M_s value would be greatly reduced, so that in the m_b - M_s plane the "event point" would be

Figure 18. Inferred average shear stress drops for a variety of events in the Alaskan-Aleutian region versus fault zone maximum dimension. The results imply a roughly linear trend that indicates that events with larger failure zones have decidedly lower average stress drops than smaller dimension events.

within the high stress drop small fault dimension portion of the theoretical grid. Presumably if one were to use higher frequency magnitude data for the average stress drop inferences, along with appropriate high frequency theoretical predictions, we could begin to resolve the very local and very high stress drop at the hypocenter itself.

Use of "variable frequency magnitude" data is one approach that does in fact use high frequency data to obtain greater resolution of spatial stress variations and also avoids uncertainties associated with the depth dependence of M_s data. Figure (19) illustrates the data generation method used, which is simple (Gaussian) narrow band filtering of the compressional wave train. Body wave magnitudes at different frequencies, $m_b(f)$ values, can be easily computed from the maxima in the envelopes of the filter outputs. In addition the time of the envelope peak can be obtained (group arrival time for energy at the filter center frequency) and so the magnitude can be associated with a particular body wave phase. In applications to stress estimates, hypocentral average stress drop estimates would be associated with first arrival P wave, while average stress estimates for other portions of the failure plane would be obtained from later group arrivals. In principle this approach can be used to obtain spatial estimates of (locally averaged) stress drops all along the failure plane.

Figure (20) illustrates the nature of the theoretically predicted event loci in the " $m_b(f)$ plane", where m_b at .45 Hz and 2.25 Hz have been used. The averaging is over a dimension of about 4 km (the wave length mean for .45 and 2.25 Hz P waves). Clearly the resolution in both stress drop and fault zone maximum dimension is very good and considerably better than was the usual case for the m_b - M_s grid.

Figure 19. Examples of narrow band filtering to generate variable frequency body wave magnitudes $m_b(f)$ for events. An explosion and a shallow earthquake are shown as examples. The $m_b(f)$ values are computed from the peaks in the envelope of the filter output at the first arrival P wave.

Figure 20. Eurasian event data in the $m_b(f)$ plane. The events are located in the areas shown in Figure 20. The theoretical curves are based on relaxation source theory models and provide a basis for the estimation of hypocentral stress drop, that is the average stress change near the hypocenter averaged over a region of dimension given by the location of the observed point in the $m_b(f)$ plane relative to the lines marked: $L = 10 \text{ km}$, $L = 5 \text{ km}$, etc. Discrimination of explosions and earthquakes is also obtained.

The data shown in the figure are associated with Eurasian events, with locations shown in Figure (21). Both trench and intraplate events are represented. It is worth noting that the highest stress drop events are from the interiors of the plates although the data shows wide stress drop variations for both classes of events.

The distribution of ordinary m_b magnitude is rather wide, and events with considerably lower m_b values are represented than was the case for the Alaskan-Aleutian data in the previous figures. Thus, noting the stress drop distribution for the data, it is probably that many of the events had fairly small fault dimensions and corresponding very small zones in which the stress was concentrated in the hypocentral region. Thus the averages over a roughly 4 km area could give relatively low average stress drops of from 10 to 100 bars for the small events in the lower left of the plot.

In any case however, this approach gives typically much higher stress drops than does the $m_b - M_s$ method, in agreement with the interpretation that this is the result of averaging over a smaller region near the high stress hypocenter.

In total then, all these comparisons with observations lead to the conclusion that earthquakes are initiated at quite high stress levels (2-3 kb or somewhat higher) which are comparable to laboratory determined stresses required for failure. Once initiated however the failure zone can continue to dynamically expand through zones of low initial (ambient) tectonic stress because of the large dynamic stresses at the failure front. Thus seismic estimates based on finite wave length radiation will obtain average static stress drops and these will in general be relatively low, of the

Figure 21. Earthquake locations for some of the events studied in Eurasia (solid points). The X denotes the location of the explosions studied. The two earthquakes denoted by the symbols ▲ are very high hypocentral stress drop events and appear near the 10 kb line in Figure 19.

order of 100 bars for methods implying m_b - M_s data for example, but higher if shorter wave length higher resolution methods are used. The results given in this section are quite consistent with this view.

V. Stress Estimates and Stress Maps for the Alaskan-Aleutian Region (1)

Figure (22) through (26) show the results of applying the $m_b - M_s$ method to the fairly large total population of events from the Alaskan-Aleutian region. Here the events were separated into groups based on depth and type so that the stress estimates are plotted in particular depth ranges. These figures, taken together, present a picture of the stress level as a function of depth for the epicentral region covered by the zone outlined as "Seismic Region 1". In this discussion and through the remaining sections, the term "stress" is to mean recoverable ambient stress in the sense previously defined unless stated otherwise.

In this regard the spatial consistency of the stress drop estimates suggests but certainly does not necessarily require, that all the stress drops reduce the stress to a fixed level at the failure boundary or to a fixed fraction of the ambient level. In view of the wide variations in material properties to be expected both in depth and laterally along the plate boundary however, such consistency would suggest a total stress drop for the events.

Certainly one of the striking features of the results is the consistent variation of the stress with depth and along the plate boundary. In addition, the data give very high stresses in the region from Kodiak Island to the end of the Alaskan peninsula and this picture is consistent with the presence of a seismic gap in this region (e.g., Sykes, 1971; Kelleher et al, 1973, 1974) and the fact that the recurrence data for large earthquakes in this region implies another large earthquake zone in the near future (e.g., Shakal and Willis, 1972). Further we find that this stress estimation methods gives rather low stress levels for the zone of the 1964 Alaskan event, which would be expected

Figure 22. Stress map from events in the Alaskan-Aleutian Arc region having focal depths in the range from 0 to 20 km. Each event is denoted by a time sequenced event number and average stress drop. The stress drop values are contoured to show the spatial stress pattern and provide a useful indication of the consistency of the results. The event numbers provide an indication of the time sequence of the sampling. If shear melting occurs in the failure zone (on the failure "plane"), then the stress levels shown are equal to the ambient prestress levels in the material.

Figure 24. Stress map from events in the Alaskan-Aleutian Arc region having focal depths in the 33 to 40 km range. Events listed with hypocenters at 33 km are omitted. The high stress zone along the Alaska Peninsula is seen to persist through this depth interval.

Figure 25. Stress map from events in the Alaskan-Aleutian Arc region having focal depths in the 40 to 55 km range. The high stress zone along the peninsula is larger and more extensive in this depth range than at the shallower depths. It is likely to be the depth range for nucleation of a very large earthquake as a result of failure along the plate boundary from just west of Kodiak Island to the end of the Alaskan Peninsula.

Figure 26. Stress map obtained for events listed with depths of 33 km. The depths were interpreted as shallow due in part of the epicentral locations of the events. The events were interpreted as normal dip slip on the ocean side of the trench zone and strike slip in the region to the East. Several stress zones are indicated.

VI. Stress Estimates and Stress Maps for the Japan-Kuriles-Kamchatka Region (19)

Figures (27) through (30) show the inferred stress levels for the Japan region in the depth range from 0 to 50 km. Somewhat more data is available for this region than for the Alaskan zone and the stress estimates are correspondingly somewhat more detailed--and reliable. An outstanding feature of these results are delineated zones of very high stress, in the depth range 20 - 50 km, off the coasts of Kamchatka, Hokkaido and one off Honshu, near Tokyo. These patterns show some irregularity with depth but in general the high stress patterns are found to persist over a considerable range in depth. As with Alaska, the high stress zones are generally at depths greater than 20 km, with the stress levels near the surface more moderate. The one exception appears to be the very high stresses indicated in the Kuriles region in Figure (27). This stress zone appears to be located in the North American plate, northwest of the trench marking the plate boundary, rather than entirely in the narrow zone along the boundary itself.

A particularly critical test of the entire approach is available using the data set for this region. Specifically, we should obtain a high stress level in a concentrated zone before a large earthquake, while after the event we should obtain a stress pattern which is considerably changed. One would expect a general reduction in the stress with perhaps some high stress concentration levels at the ends of the failure zone. The Japan region data contained two large events with reasonably large numbers of small events occurring both before and after them, and consequently we can test the seismic stress estimates obtained against the criteria that the local stress level be generally lower after such a large event.

Figure 27. Stress map from events in the Japan-Kuriles-Kamchatka region for the depth range from 0 to 20 km. The high stress zone in the Kuriles region corresponds to a shallow stress concentration of apparently large dimension within the Asian plate, which is being underthrust by the Pacific plate.

Figure 29. Stress map from events in the Japan-Kuriles-Kamchatka region for the depth range from 33 to 40 km. The high stress zone off the Honshu coast appears more regular and extensive in this depth range and constitutes a high risk zone.

Figure 30. Stress map for the Japan-Kuriles-Kamchatka seismic region from events having focal depths in the depth range 40 to 50 km, with events occurring before the large event, No. 193, represented. This event and its aftershocks occurred within the moderately high stress zone off the northeast point of Hokkaido and results in a considerable change in the stress level and spatial pattern.

Figure (31) shows the stress magnitude distribution for the 40 to 50 km depth range after one of the large events (No. 193) that would be expected to strongly perturb the stress field in a large zone. We would expect to observe a considerable change in the stress field as obtained from the sampling provided by the subsequent small events. Comparison of Figures (30) and (31) shows a major shift in the stress pattern due to the large event, with the high stress zone shifting landward after the event and a considerable reduction in stress in the immediate vicinity of the large earthquake. Hence, our spatial resolution of the stress appears just sufficient in this case to verify that the stress change is as would be expected.

Figures (32) and (33) show the stress field contours and the event data before and after another large event (Event No. 144). This event occurred just at the point where the Aleutian arc intersects the Kamchatka Peninsula. Figure (32) shows that prior to the event the zone was rather highly stressed, with the stress level reaching at least 500 bars. After the event, Figure (33) shows that the later small events provide a sampling of a stress field that is generally very much lower in this region, with however, some very small zones--presumably near the ends of the newly failed zone associated with the large earthquake--where the stress is quite high. Thus the time-space picture presented by this event sequence again entirely conforms to what we think should occur.

Finally we also observe, as for the Alaska region, that the stress levels obtained by contouring are spatially and temporarily consistent. That is, when a high stress zone is delineated by a

Figure 31. Stress map for the Japan-Kuriles-Kamchatka seismic region from events having focal depths in the depth range 40 to 50 km, with the events used including event No. 193 and its aftershocks, but excluding events before 193 in the Northern Hokkaido region. Comparison with Figure 30 shows a strong shift in the stress pattern with an increase in the stress level shoreward, to a level near 1 kbar, and a decrease of stress in the zone of faulting for this large event.

Figure 32. Stress map for the Japan-Kuriles-Kamchatka seismic region from events having focal depths given as 33 km, with only the events before the large event No. 144 represented. Event No. 144, which occurred off the coast of Kamchatka, and its aftershocks are not included so that the stress field prior to this event is shown. It corresponds to the northern-most high stress zone along the Kamchatka coast.

Figure 33. Epicentral stress map for the Japan-Kuriles-Kamchatka from events with focal depths at 33 km, with event No. 144 and its aftershocks included. The stress field in the northern high stress zone is now much reduced, with some high stress indicating events occurring near what appears to be the edges of the new failure zone. The stress level, as evidenced by the event sampling, therefore appears to be much lower, as would be expected, but with higher stress near the edges of the recent failure "plane" formed by the large event, as also would be expected.

sampling of events for a region, then we find that subsequent (small) events **in** the zone will all be high stress drop events until a large event occurs, at which time the stress field is greatly changed, usually over the entire zone.

We conclude that the general space-time picture obtained from this stress estimation procedure is quite consistent with the expected stress behavior within the active seismic zones studied.

VII. Stress Estimates and Stress Maps for the Andean South America Region (8)

Preliminary results for the Andean South American (Nazca Plate subduction zone) region are shown in Figures 34 and 35. These results suggest at least two high stress zones that persist in depth. Study of this region to delineate these zones more precisely will clearly require much more data. This situation is characteristic of many of the active seismic regions globally, where sufficient m_b - M_s data is not yet available.

Figure 34. Stress map for the Andean South America Region for the depth range 0 to 20 km. Only the event sequence numbers are indicated. The data is rather sparse and a more reliable stress map requires additional data. Several moderately high stress zones are indicated however.

Figure 35. Stress map for the Andean South America Region for the depth range 40 to 50 km. Event sequence numbers are indicated, with stress contours providing the stress drop values. The data is quite sparse but three high stress zones are indicated.

VIII. Results and Conclusions

The interpretations of the observed seismic event data in this study, coupled with theoretical considerations, indicate that:

(1) The non-hydrostatic stress field within the lithosphere is highly variable spatially and that consequently, the stress changes or recoverable ambient stress from an earthquake is also highly variable. This leads to highly variable observed average stress drops for earthquakes which are dependent on the resolution of the method used to obtain the stress estimate.

(2) High non-hydrostatic tectonic stresses appear to be involved in failure initiation, with values in the range 2-5 kbars probable. Failure growth is accomplished dynamically and large failure zones appear to occur only when localized stress concentrations are reasonably close spaced.

(3) Stress drops could well be to an essentially fixed level of non-recoverable ambient stress and roughly independent of location and depth, so that recoverable ambient stress varies directly with total ambient stress. It also appears quite possible that total shear stress drop occurs, in the shear melting and total loss of shear strength associated with high dynamic stress at the failure front may quite likely occur.

(4) Several regions of high average stress, which are correlated with seismicity gaps, were delineated. The most notable of these was one along the Alaskan Peninsula from the east end of Kodiak Island to the beginning of the Aleutian Island chain.

(5) The m_b - M_s magnitude method serves as a good reconnaissance approach to the delineation of high stress zones. Other methods, notably the variable frequency magnitude method are more suitable for high resolution studies using small events and can be used to "map" the stress in detail.

(6) Prediction of location and large event size, in terms of maximum probable energy release, appears reasonable using stress estimation based on the seismic methods discussed. The occurrence time of such a large event could possibly be based on the time sequence of the stress pattern and/or the seismicity pattern. In any case stress estimates on a global scale provides a basis for systematic work focussed on selected areas of high stress.

REFERENCES

- Anderson, D. L., and C. B. Archambeau, 1964. The Anelasticity of the Earth, J. Geophys. Res., 69, 2071-2084.
- Andrews, D. J., 1976. Rupture Propagation with Finite Stress in Anti-plane Strain, J. Geophys. Res., 81, 3575.
- Archambeau, C. B., 1972. The Theory of Stress Wave Radiation from Explosions in Prestressed Media, Geophys. J. Roy. astr. Soc., 29, 329-366.
- Archambeau, C. B., 1968. General Theory of Elastodynamic Source Fields, Rev. of Geophys., 16, 241-288.
- Archambeau, C. B., 1964. Elastodynamic Source Theory; Ph.D. Thesis, Calif. Inst. of Tech., Pasadena, CA.
- Archambeau, C. B., E. A. Flinn and D. G. Lambert, 1969. Fine Structure of the Upper Mantle, J. Geophys. Res., 74, 5825-5865.
- Archambeau, C. B. and J. B. Minster, 1978. Dynamics in prestressed media with moving phase boundaries: A continuum theory of failure in solids, Geophys. J. Roy. astr. Soc., 52, 65-96.
- Bache, T. C., T. G. Barker and C. B. Archambeau, 1978. The San Fernando Earthquake - A model consistent with near-field and far-field observations at long and short periods. Publication in preparation. From: Final Report, USGS Earthquake Hazards Program, for the period Oct. 1, 1976 to Sept. 30, 1977. Systems, Science and Software, La Jolla, California.

- Bache, T. C., and D. G. Harkrider, 1976. The Body Waves Due to a General Seismic Source in a Layered Earth Model: I Formulation of the Theory, Bull. Seism. Soc. Am., (in press.)
- Brune, J. N., 1970. Tectonic Stress and the Spectra of Seismic Shear Waves from Earthquakes, J. Geophys. Res., 75, 4997. Correction: 1971, J. Geophys. Res., 76, 5002.
- Cherry, J. T., T. C. Bache, and J. F. Masso, 1976. A Three Dimensional Finite Difference Simulation of Earthquake Faulting, Abstract, AGU National Meeting, Washington, D. C.
- Friedman, M., J. M. Logan and J. A. Rigert, 1974. Glass-Indurated Quartz Gonge in Sliding-Friction Experiments on Sandstone, Geol. Soc. Am. Bull., 85, 937-942.
- Hanks, T. C., and W. Thatcher, 1972. A Graphical Representation of Seismic Sources, J. Geophys. Res., 77, 4393-4405.
- Helmberger, D. V., and R. A. Wiggins, 1971. Upper Mantle Structure of the Middlewestern United States, J. Geophys. Res., 76, 3229.
- Jungels, P. H., 1973. Models of Tectonic Processes Associated with Earthquakes, Ph.D. Thesis, California Institute of Technology, Pasadena, CA.
- Jungels, P. H., and G. A. Frazier, 1973. Finite Element Analysis of the Residual Displacements for an Earthquake Rupture: Source Parameters for the San Fernando Earthquake, J. Geophys. Res., 78, 5062.

- Kelleher, J. Savino, H. Rawlett, and W. McCann, 1974. Why and where Great Thrust Earthquakes Occur Along Island Arcs, J. Geophys. Res., 79, 4889-4899.
- Kelleher, J., L. Sykes, and J. Oliver, 1973. Possible Criteria for Predicting Earthquake Locations and Their Application to Major Plate Boundaries of the Pacific and the Caribbean, J. Geophys. Res., 78, 2547-2585.
- McKenzie, D. and J. N. Brune, 1972. Melting on Fault Planes During Large Earthquakes, Geophys. J. Roy. astr. Soc., 29, 65-78.
- Minster, J. B., 1973. Elastodynamics of Failure in a Continuum, Ph.D. Thesis, California Institute of Technology, Pasadena, CA.
- Minster, J. B. and A. M. Suteau, 1976. Far-field waveforms from an arbitrarily expanding, transparent spherical cavity in a prestressed medium, Geophys. J. Roy. astr. Soc., 50, 215-233.
- Molnar, P. and M. Wyss, 1972. Moments, magnitudes, source dimensions, and stresses of shallow-focus earthquakes in the Tonga-Kermadec arc, Phys. Earth Planet. Inter., 6, 263.
- Smith, A. T. and M. N. Toksoz, 1972. Stress distribution beneath island arcs, Geophys. J. Roy. astr. Soc., 29, 289-318.
- Teufel, L. W., 1976. The Measurement of Contact Areas and Temperature During Frictional Sliding of Tennessee Sandstone, M.A. Thesis, Texas A & M University.
- Thatcher, W. and T. C. Hanks, 1973. Source Parameters of Southern California Earthquakes, J. Geophys. Res., 78, 8547-8576.

- Trifunac, M. D., 1974. A Three-dimensional Dislocation Model for the San Fernando, California Earthquake of February 9, 1971, Bull. Seism. Soc. Am., 164, 149-172.
- Trifunac, M. D. and F. E. Udevadia, 1974. Parkfield, California Earthquake of June 27, 1966; A Three-dimensional Moving Dislocation, Bull. Seism. Soc. Am., 64, 511-533.
- Wiggins, R. A., and D. V. Helmberger, 1974. Synthetic Seismogram Computation by Expansion in Generalized Rays, Geophys. J. Roy. astr. Soc., 37, 73-90.
- Wyss, M., 1970a. Stress estimates of South American shallow and deep earthquakes, J. Geophys. Res., 75, 1529.
- Wyss, M., 1970b. A comparison of apparent stresses of earthquakes on ridges with earthquakes in trenches, Geophys. J. Roy. astr. Soc., 19(5), 479.
- Wyss, M. and P. Molnar, 1972a. Source parameters of intermediate and deep focus earthquakes in the Tonga arc, Phys. Earth Planet. Inter., 6, 263.
- Wyss, M. and P. Molnar, 1972b. Efficiency, stress-drop, apparent stress, effective stress, and frictional stress of Denver, Colorado, earthquakes, J. Geophys. Res., 77, 1433.
- Wyss, M. and J. N. Brune, 1971. Regional variations of source parameters in Southern California estimated from the ratio of short to long period amplitudes, Bull. Seism. Soc. Amer., 61, 1153.
- Wyss, M., and J. N. Brune, 1968. Seismic moment, stress and source dimensions for earthquakes in the California-Nevada region, J. Geophys. Res., 73, 4681.

SEISMIC GAPS IN COASTAL PERU

by James W. Dewey* and William Spence*

ABSTRACT

Teleseismic relocation of earthquakes in a 650-km long segment of the Peruvian coastal region between 8.5° S. and 14° S. for the interval 1964 to 1974 resolves two parallel zones of shallow ($h < 70$ km) earthquake activity that trend parallel to the axis of the Peru trench. One zone is defined principally by aftershocks of the two major thrust fault earthquakes of 17 October 1966 and 03 October 1974. Following the work of Abe (1972), Stauder (1975), and Isacks and Barazangi (1977), we call this zone the (plate) interface thrust zone. Because the mechanism producing the second zone of activity is still obscure, we call the second zone the W-zone. The second zone is made particularly prominent by aftershocks of the earthquake of 31 May 1970, but is much in evidence in sections of the subduction zone far removed from the source region of that earthquake. The second zone is centered about 50 km inland and 30 km deeper than the interface thrust zone; the above-cited authors argue that the larger earthquakes from this zone occur within a plate, rather than at the interface of the colliding Nazca and South American plates. Within any aseismic segment of the Peruvian subduction zone, there may therefore be effectively two "seismic gaps," one corresponding to each of the prominent zones of shallow earthquake activity. It is not clear that the occurrence of a major earthquake in one of the two gaps lessens the danger of a major earthquake from the other gap. The presence of two nearby seismic zones, each perhaps caused by very different stress regimes, also complicates the job of detecting unusual seismic activity precursory to a great earthquake in one of the two zones, since such precursory activity could be masked by activity occurring in the other zone. In the case of the interface thrust zone earthquake of 03 October 1974, we find that removal of W-zone earthquakes from seismicity plots dramatically emphasizes the high seismic activity in the epicentral region of that earthquake in the five years preceding the main shock. Although the general region of the 1974 earthquake had been identified by Kelleher (1972) as a potentially dangerous seismic gap on the basis of the record of major historical and instrumentally-recorded earthquakes, we find nothing in the spatial pattern of teleseismically-determined areas of high and low seismicity since 1964 that would have enabled an estimate to within a factor of two of the length of the region that actually ruptured in the 1974 earthquake.

INTRODUCTION

The primary goal of this research is to improve teleseismically-determined epicenters to define source regions of future destructive earthquakes and, ultimately, to help predict when the earthquakes will occur. We are initially giving heavy emphasis to methods of obtaining more accurate hypocenters, using previously acquired travel-time data supplemented by small numbers of travel-time data read from archived seismograms, with a view toward more sharply defining seismic gaps in generally active subduction zones and with the hope of finding unusual, moderate magnitude, seismic activity precursory to destructive earthquakes in these gaps.

*U.S. Geological Survey, MS 967, Box 25046, Denver Federal Center
Denver, CO 80225

Here we make a detailed teleseismic study of the seismicity of coastal Peru between 8.5° S. and 14° S. (fig. 1). This region has experienced three major earthquakes since 1964, those of 661017 ($M_S = 7.5$), 700531 ($M_S = 7.8$), and 741003 ($M_S = 7.8$) (dates of earthquakes are hereafter given in year, month, and day.) The earthquakes of 1966 and 1974 were thrust-fault earthquakes (Abe, 1972; Spence, *et al.*, 1975); probably on the interface between the subducting Nazca plate and the overriding South American plate. The earthquake of 1970 was a normal faulting earthquake (Abe, 1972). Abe, Stauder (1975), and Isacks and Barazangi (1977) have explained the 1970 earthquake in terms of stresses within the Nazca plate, since the focal mechanism of that earthquake and its aftershocks are apparently not explicable in terms of thrust faulting on the interface between the Nazca and South American plates.

The shock of 741003 occurred in a region that had been identified by Kelleher (1972) as a potentially-dangerous seismic gap, on the basis of the instrumental and historical record of major earthquakes in the Peruvian coastal region. We chose 1964 as the starting point for this study because the earthquake data catalogues of the International Seismological Center (ISC) are readily available and complete from this date for Peruvian shocks down to $m_b \approx 4.7$ (Langer, 1977). We thus had nearly a decade of moderate earthquake activity to use to see whether Kelleher's "earthquake gap watch" could have been made more precise with the help of readily available teleseismic data.

Routinely determined hypocenters from the Peruvian region, used indiscriminately, reveal little about the structure of the subduction zone (fig. 2). It is possible to greatly extend the usefulness of the routinely-determined hypocenters by culling out earthquakes that are likely to be mislocated because they were poorly recorded, had no depth-phases read for them, etc., (Barazangi and Isacks, 1976). However, our approach is to relocate all hypocenters and then to cull out the less reliably-determined shocks on the basis of their lower estimated precision as indicated by the size of the confidence ellipse on the epicentral coordinates.

METHOD OF RELOCATING EARTHQUAKES

Hypocenters in figures 3 through 10 were determined by the method of Joint Hypocenter Determination (JHD) (the program used is modified from Dewey, 1971) or by the single event method with station weights and station adjustments to the travel time tables determined by JHD. Separate station weights and adjustments were computed for earthquakes lying south of 11° S. and those lying north of 11° S. The calibration event for the JHD for the southern group of earthquakes was a shock of 741010 ($m_b = 5.3$, USGS), restrained to the origin time and location (195300.7 GMT, 12.525° S., 77.623° W., $h = 23.4$ km) determined by Langer and Spence (1978) with data from a temporary regional network of seismographs installed to record aftershocks of the major earthquake of 741003. The calibration event of the group of hypocenters lying north of 11° S. was the shock of 700604, 040927.9 GMT, ($m_b = 5.8$, USGS), whose hypocenter had been first located with respect to the calibration event of the southern group of hypocenters.

Earthquakes relocated were those originally assigned by the ISC to the region shown in figure 1. Arrival time data consisted for the most part of P, pP, and PKP readings listed in the ISC. We read seismograms of selected WWSSN stations to obtain readings of pP to supplement readings of depth phases reported in bulletins. ISC arrival time data for aftershocks of the earthquake of 741003 were supplemented by numerous P- and S-readings from the temporary network of Langer and Spence (1978). We also attempted to use S-readings at regional stations to help improve the location of smaller shocks for which P-wave readings were obtained through only a limited range of azimuths.

EFFECTIVENESS OF LOCATION METHODS

We tried to relocate 478 teleseismically-recorded earthquakes. For 37% of these shocks (hereafter designated b-quality), 90% confidence ellipses on the epicentral coordinates have semiaxes less than 20 km in length. For 36% of the earthquakes (hereafter designated d-quality), 90% confidence ellipses on the epicentral coordinates have semi-major axes between 20 km and 50 km in length. The remainder (27%) of the shocks could not be located to an estimated precision of 50 km or better; these shocks are not plotted in figures 3 through 10. The precision of focal depth determination is, for most shocks, roughly comparable to the precision of the determination of epicentral coordinates. Two-thirds of the hypocenters were located with the use of depth phases and many shocks were deep enough and near enough to the local stations NNA and HUA that P-wave times from these stations enabled a reliable depth computation. Among the tendencies shown in depth sections in figures 3-10, we are most suspicious of the apparent deepening of the hypocenters toward the Peru trench in the region from 100 km to 0 km from the trench. Two factors would tend to cause shocks nearest the trench to be mislocated too deep: (1) a shallower M-discontinuity near the trench would cause earlier Pn-arrivals at the local stations and a consequent overestimation of focal depth, and (2), the interpretation of the pwP-phase as pP would cause deeper focal depths to be estimated for shocks occurring beneath the deeper water of the trench (Mendiguren, 1971).

P-wave readings at the nearest stations ($\Delta < 5^\circ$, variance typically 3 sec^2) and at teleseismic stations of $\Delta > 20^\circ$ (variance typically $1 \frac{1}{2} \text{ sec}^2$) were most consistent. P-wave arrival times at stations in the distance range of 8° to 15° were generally less consistent (variance typically 8 sec^2). The high variance regional stations include such normally reliable stations as ARE, PNS, and LPB. Inspection of seismograms at ARE and LBP shows that the initial P-waves from shocks in the region under study are very emergent and of low amplitude, probably the effect of a P-wave shadow zone (Gutenberg and Richter, 1939), and it is likely that later arriving P-waves are often reported as the initial P-wave. Computer location programs will tend to interpret such late arrivals as indicating greater focal depth than is actually the case. The variance of the set of pP readings was about 6 sec^2 , corresponding to an uncertainty of somewhat more than 10 km in focal depth determinations based on a single pP datum. The somewhat high variances of pP times relative to those of teleseismic P-wave times may arise in part from occasional pwP- or sP-phases being read as pP. The regional S-waves reported in the ISC bulletins had a disappointingly high variance (typically 50 sec^2) and correspondingly

were assigned low weights in the computation of hypocenters. The S-wave data provided by the aftershock network of Langer and Spence did prove highly consistent (variance approximately 3 sec^2) indicating that by considering earthquakes only from one aftershock zone at a time and by devoting special care to the identification of S-waves, valuable use may still be made of the S-waves.

In general, we have had difficulty obtaining precise locations of smaller teleseismically-recorded earthquakes. These shocks tend to be most severely mislocated by the ISC and the USGS. We attribute our difficulty with smaller shocks to the inconsistent times reported as P-wave arrival times from stations in the range 8° to 15° ; for the small shocks, these stations often exert a strong (and undesirable) influence on the determination of the hypocenter.

DIFFERENTIATION OF EARTHQUAKE ZONES PERPENDICULAR TO THE TRENCH AXIS

Projecting a 650 km-long segment of the seismic zone (fig. 1) onto a plane perpendicular to the trench axis (fig. 3), one sees two prominent zones of activity between 60 km and 200 kilometers east of the trench. The seaward and shallower of the two zones is defined principally by aftershocks of the earthquakes of 661017 and 741003, and by a series of moderate shocks that preceded the earthquake of 741003. Since the two major earthquakes, 661017 and 741003, probably occurred as the result of slippage on the interface between the subducting Nazca plate and the overriding South American plate (Abe, 1972; Spence, et al., 1975), we call this zone the interface thrust zone.

We call the second zone the W-zone, because the origin of earthquake-generating stresses in the zone is controversial, and we distinguish it from the interface thrust zone for two reasons. First, the zone of hypocenters we call the W-zone appears to be a cluster of high seismicity spatially separated from the interface thrust zone when the 650 km long seismic region is viewed in the transverse section (fig. 3). It is true that the W-zone receives heavy emphasis from aftershocks of the earthquake of 700531. However, many other earthquakes occurred at the same depth and distance from the trench, north and south of the aftershock zone of 700531 (these are most evident in figures 7 and 8, before 700531). Our second reason for defining the W-zone separately from the interface thrust zone is that focal mechanisms thus far determined for earthquakes in this zone are inconsistent with the earthquakes occurring on an eastward shallowly-dipping interface between the plates; the focal mechanisms imply that the large earthquakes are the result instead of internal plate deformation (Abe, 1972; Stauder, 1975; Isacks and Barazangi, 1977). However, since relatively few focal mechanisms have been well-determined for W-zone earthquakes, and these do not seem to have occurred at the interface between the plates, we do not know where the interface lies with respect to the W-zone, nor can we be sure that some of the W-zone hypocenters plotted in figure 3-10 are not occurring on a downdip extension of the principal interface thrust zone. It is necessary to further define the nature of the W-zone with additional focal mechanisms of the moderately large earthquakes occurring more than 125 km from the trench.

were assigned low weights in the computation of hypocenters. The S-wave data provided by the aftershock network of Langer and Spence did prove highly consistent (variance approximately 3 sec^2) indicating that by considering earthquakes only from one aftershock zone at a time and by devoting special care to the identification of S-waves, valuable use may still be made of the S-waves.

In general, we have had difficulty obtaining precise locations of smaller teleseismically-recorded earthquakes. These shocks tend to be most severely mislocated by the ISC and the USGS. We attribute our difficulty with smaller shocks to the inconsistent times reported as P-wave arrival times from stations in the range 8° to 15° ; for the small shocks, these stations often exert a strong (and undesirable) influence on the determination of the hypocenter.

DIFFERENTIATION OF EARTHQUAKE ZONES PERPENDICULAR TO THE TRENCH AXIS

Projecting a 650 km-long segment of the seismic zone (fig. 1) onto a plane perpendicular to the trench axis (fig. 3), one sees two prominent zones of activity between 60 km and 200 kilometers east of the trench. The seaward and shallower of the two zones is defined principally by aftershocks of the earthquakes of 661017 and 741003, and by a series of moderate shocks that preceded the earthquake of 741003. Since the two major earthquakes, 661017 and 741003, probably occurred as the result of slippage on the interface between the subducting Nazca plate and the overriding South American plate (Abe, 1972; Spence, *et al.*, 1975), we call this zone the interface thrust zone.

We call the second zone the W-zone, because the origin of earthquake-generating stresses in the zone is controversial, and we distinguish it from the interface thrust zone for two reasons. First, the zone of hypocenters we call the W-zone appears to be a cluster of high seismicity spatially separated from the interface thrust zone when the 650 km long seismic region is viewed in the transverse section (fig. 3). It is true that the W-zone receives heavy emphasis from aftershocks of the earthquake of 700531. However, many other earthquakes occurred at the same depth and distance from the trench, north and south of the aftershock zone of 700531 (these are most evident in figures 7 and 8, before 700531). Our second reason for defining the W-zone separately from the interface thrust zone is that focal mechanisms thus far determined for earthquakes in this zone are inconsistent with the earthquakes occurring on an eastward shallowly-dipping interface between the plates; the focal mechanisms imply that the large earthquakes are the result instead of internal plate deformation (Abe, 1972; Stauder, 1975; Isacks and Barazangi, 1977). However, since relatively few focal mechanisms have been well-determined for W-zone earthquakes, and these do not seem to have occurred at the interface between the plates, we do not know where the interface lies with respect to the W-zone, nor can we be sure that some of the W-zone hypocenters plotted in figure 3-10 are not occurring on a downdip extension of the principle interface thrust zone. It is necessary to further define the nature of the W-zone with additional focal mechanisms of the moderately large earthquakes occurring more than 125 km from the trench.

If we accept the hypothesis that there are two extensive shallow ($h < 70$ km) seismic zones, each capable of producing destructive earthquakes, then we must consider the possibility that a conventional "seismic gap" inland from the Peru Trench has the potential of producing two major (rupture length greater than 100 km) earthquakes, or, as a corollary, that the occurrence of one major earthquake does not necessarily preclude the occurrence of another major earthquake in the same section of arc. Although the earthquake of 700531 was the only disastrous "W-zone" earthquake in this region from the period 1964-1974, it is possible that the major earthquake, $M_S = 8.0$, of 24 May 1940 may have been a similar subcrustal earthquake, originating within the W-zone south of the focal region of 700531 and inland from the southern portion of the focal region of the thrust fault earthquake of 661017 (J. Kelleher, personal commun., 1978; Silgado, 1973).

The presence of two nearby seismic zones complicates the job of detecting unusual seismic activity precursory to a great earthquake in one of the two zones. The mechanisms proposed to account for the largest of the W-zone earthquakes (Abe, 1972; Stauder, 1975; Barazangi and Isacks, 1976) involve stresses generated by large scale deformation of the subducting Nazca plate and it is not clear how these stresses will be affected by stress conditions at the interface between the Nazca and South American plate. Therefore, it is quite possible, for example, that the W-zone will be seismically active when the interface zone is inactive, and that a resumption of distinctive activity in the interface zone prior to a major interface earthquake would go unnoticed amidst the general W-zone activity. We shall show later that removal of W-zone earthquakes from plots of the seismicity of this section of the Peruvian subduction zone brings into prominence a group of earthquakes occurring over a five year period in the hypocentral region of 741003 prior to the occurrence of that earthquake.

AFTERSHOCK CLUSTERS WITHIN THE AFTERSHOCK ZONES

Teleseismically recorded aftershocks for one month following each of the three major earthquakes under study (661017, 700531, and 741003) show a tendency to occur in clusters rather than to be distributed evenly throughout their aftershock zones. These clusters are outlined in figures 4, 5, and 6. The clustering of teleseismically-recorded aftershocks makes it difficult to assess from teleseismic data how much of a seismic gap has broken in a particular earthquake. For example, the cluster 66c, figure 4, consists of two rather small earthquakes. Should one suppose that the main rupture of 661017 extended from the main shock hypocenter continuously southeast to 66c, or do the earthquakes at 66c represent a distinct source region that was triggered by 661017? If the latter, does the 50 km segment of the interface between the mainshock and 66c still have the potential to rupture in a damaging earthquake?

The clustering of aftershocks is especially noticeable in the case of the shock of 700531 which was followed within 31 days by numerous moderate aftershocks. We have defined two clusters, 70a and 70b (fig. 5). The mainshock occurred in the cluster 70a. The first teleseismically recorded aftershock occurred at the south end of cluster 70b, so it is likely that the whole region of 70a-70b was involved in the mainshock rupture. Lomnitz (1970) pointed out that the 1970 earthquake was very likely a "multiple event" and that prominent later occurring events in the earthquake appeared to have focal

mechanisms different than that of the first event. Stauder (1975) and Isacks and Barazangi (1977) have called attention to the fact that many of the aftershock focal mechanisms involve thrust faulting in contrast to the normal faulting mechanisms of the 1970 main shock. Clusters of aftershocks following great earthquakes are often considered to occur at asperities on the main fault surface involved in the mainshock. However, the focal mechanisms determined by Stauder (1972) and Isacks and Barazangi (1977) for aftershocks of 700531 rule out the possibility that these shocks could represent breakage of previously unruptured patches of the fault plane involved in the main shock. It is interesting to speculate that the main earthquake of 700531 may have involved several spatially separated "events," corresponding to the regions of the aftershock activity, unconnected by continuous dislocation surface. These individual events may reflect discrete stress concentrations and might be analogous to the isolated individual rock bursts postulated to be components of larger gold mine tremors in South Africa (Spottiswoode and McGarr, 1966).

The shocks of 661017 (fig. 4) and 741003 (fig. 6) were both followed by aftershocks in the W-zone. In the case of the 1974 shock, there is a cluster of aftershock activity (74b), well-located by the temporary stations of Langer and Spence (1978), inland from and deeper than the principal aftershock region 74a. Langer and Spence (paper in preparation) have studied the geometry of the 1974 aftershock zone in some detail and conclude that the 74b aftershocks probably represent internal deformation of the Nazca plate. The occurrence in the W-zone of aftershocks of interface thrust earthquakes is the strongest evidence we now have of a linkage between the seismicity of the two zones.

NO SHORT-TERM POSITIVE CORRELATION BETWEEN ACTIVITY IN THREE PRINCIPAL SOURCE REGIONS

Figures 4, 5, and 6 show that in the month following each of the major shocks of 661017, 700531, and 741003, there were no well-located teleseismically-recorded shocks from the other two source regions. Apparently the stress changes associated with each of the main shocks was insufficient to immediately trigger an earthquake in the other two aftershock zones.

SEARCH FOR UNUSUAL PATTERNS OF SEISMICITY BEFORE THE EARTHQUAKES OF 17 OCTOBER 1966, 31 MAY 1970 AND 03 OCTOBER 1974

Kelleher and Savino (1975) report from study of a number of great earthquakes that "extensive portions of the interior of the rupture zones commonly remained aseismic until the time of the main shock" and that "seismicity that did occur was confined to the vicinity of the epicenters * * * of the impending main shocks and/or the edges of the rupture zones. There are some indications that the level of this prior activity, particularly in the vicinity of the epicenter, increased as the time of the main shock approached" (*ibid.*, p. 260).

We have constructed plots of seismicity for periods since 1964 before and between the earthquakes of 661017, 700531, and 741003, in order to test whether the features noted by Kelleher and Savino may also be found prior to the Peruvian earthquakes from hypocenters of moderate earthquakes for periods of time from approximately three years to approximately eleven years before the respective earthquakes.

We find no evidence among well-located epicenters (b-quality) for enhanced seismicity in the hypocentral region of 661017 prior to that earthquake, and the time period considered is too short to have permitted recognition of a seismic gap (figure 7). The pattern of seismicity prior to the earthquake of 700531 (figures 7 and 8) is consistent with the pattern identified by Kelleher and Savino. There is one well-located shock, 651106, close to devoid of well-located hypocenters. Unfortunately, there are other regions in the study area that look similar to the source region of 700531 (figs. 7 and 8), in that one could have identified a number of seismic gaps and have associated with them hypocenters that could be considered possible sites of future main shock hypocenters.

The strongest case for enhanced seismic activity prior to a destructive earthquake lies with a group of moderate shocks that preceded the earthquake of 741003. The suggestion of precursory seismic activity occurs near the source region 74a, principally along the landward edge of the future principal zone of rupture. In order to enhance this zone of foreshock activity we make use of the observation that shallow-focus seismicity within the Peruvian Benioff zone appears to occur in at least two separate source regions, and we have plotted in figure 10 only the earthquakes occurring oceanward of the boundary (figure 3) between the interface thrust fault zone and the W-zone. There are 8 earthquakes of b-quality or better within the interface thrust zone near the hypocenter of 741003. There is a tendency for a south to north progression of these hypocenters toward the hypocenters within this 650 km segment of the interface thrust zone are aftershocks of the interface thrust earthquake of 661017. The location of most of the prior seismicity on the landward side of the 1974 source region may indicate incipient failure at the lower edge of the seismically-active interface zone that was to fail in the main shock.

The seismicity of the interface thrust region prior to 1974 gave little indication of the dimensions of the source that was to rupture on 741003. the interface thrust region had been quiescent north of the source region 70a for nearly 100km to source region 66c (fig. 10) and to the southeast (beyond the range of figure 10) for 70 km, during the period 1964-1974.

DISCUSSION

The greatest hinderance to determining whether or not bursts of earthquake activity such as the group that preceded the 1974 shock may in fact serve as warning to a future destructive earthquake, is the scanty instrumental record against which we can compare activity such as that plotted in figure 10. for example, we have emphasized that the prior shocks from the source region of 1974 earthquake were unique in the region of our principal interest, since they were the only well-located shocks in the interface thrust zone that were not part of a main-shock-aftershock sequence. However, from 70 km to 130 km

southeast of our region of study, where the trend of Nazca Ridge intersects the continent (fig. 1), we have reliably relocated twelve earthquakes that would fall within our interface thrust zone if they were plotted on the same coordinate system as that used in figures 2 through 10. These earthquakes appear to be part of a larger zone, extending still farther to the southeast (Barazangi and Isacks, 1976, fig. 6). The existence of this source region does not force us to abandon hope that distinctive precursory activity may be identified prior to future interface thrust earthquakes in the 650-km long segment of our principal interest. It may be, for example, that the seismotectonics of the Peruvian subduction zone are locally affected by the subduction of the Nazca ridge. The point is that a reliable prediction method must have a long enough instrumental record to identify sections of subduction zones, such as that near the intersection of the Nazca Ridge with the South American continent, where moderate interface thrust events may occur rather commonly without being followed by a major shock. We are now in the process of relocating the larger instrumentally-recorded earthquakes that occurred in the coastal region of Peru prior to 1964.

The existence of two distinct zones of shallow-focus earthquakes, which we have labeled the interface thrust zone and the W-zone, may be an effect of the unusually low angle at which the Nazca plate dips beneath Peru. Isacks and Barazangi (1977) have attributed earthquakes of the W-zone to stresses generated by an "elastic 'unbending' after plastic deformation" (*ibid.*, p. 109), and suggest that similar types of earthquakes in more steeply dipping Benioff zones may occur at greater depths. Kanamori (1977) suggests that extensive internal deformation of subducting lithospheric plates might be characteristic of regions of strong interplate coupling, as would be expected in a region of a shallowly-dipping Benioff zone.

REFERENCES

- Abe, K. (1972). Mechanisms and tectonic implications of the 1966 and 1970 Peru earthquakes, *Phys. Earth, Planet. Interiors*, 5, 367-379.
- Barazangi, M. and Isacks, B.L. (1976). Spatial distribution of earthquakes and subduction of the Nazca plate beneath South America, *Geology*, 4, 686-692.
- Dewey, J.W. (1971). Seismicity studies with the method of joint hypocenter determination, Ph. D. Dissertation, Univ. Calif. Berkeley.
- Gutenberg, B. and Richter, C.F. (1939). New evidence for a change in physical conditions at depths near 100 kilometers, *Bull. Seism. Soc. Am.*, 29, 531-538.
- Isacks, B.L., and Barazangi, M. (1977). Geometry of Benioff Zones: Lateral segmentation and downward bending of the subduction lithosphere, *Maurice Ewing Series*, 1, 99-114.
- Kanamori, H. (1977). Seismic and aseismic slip along subduction zones and their tectonic implications, *Maurice Ewing Series*, 1, 163-174.
- Kelleher, J. (1972). Rupture zones of large South American earthquakes and some predictions, *J. Geophys. Res.*, 77, 2087-2103.
- Kelleher, J., Savino, J., Rowlett, H., and McCann, W. (1974). Why and where great thrust earthquakes occur along island areas, *J. Geophys. Res.*, 79, 4889-4899.
- Kelleher, J. and Savino, J. (1975). Distribution of seismicity before large strike slip and thrust-type earthquakes, *J. geophys. Res.*, 80, 260-271.
- Langer, C. (1977). The locations and characteristics of the aftershocks of the October 3, 1974, Peruvian earthquake, M.S. thesis, Virginia Poly. Inst., Blacksburg, VA.
- Langer, C.J. and Spence, W. (1978). A study of aftershocks of the October 3, 1974, Peru earthquake (abs.). *Earthquake Notes*, 49, 54.
- Lomnitz, C. (1971), The Peru earthquake of May 31, 1970: some preliminary seismological results, *Bull. Seism. Soc. Am.*, 61, 535-542.
- Mendiguren, J.A. (1971). Focal mechanism of a shock in the middle of the Nazca plate, *J. Geophys. Res.*, 76, 3861-3879.
- Minister, J.B., Jordan, T.H., Molnar, P., and Haines, E. (1974). Numerical modelling of instantaneous plate tectonics, *Geophys. J.R. Astr. Soc.*, 36, 541-576.

- Silgado, E. (1973). Historia de los seismos mas notables ocurridos en el Peru (1555-1970), Geofisica Panamerica, 2, 179-243.
- Spence, W., Giesecke, A., Langer, C. J., Ocola, L., Dillinger, W. H., and Jordan, J. N. (1975). The Lima, Peru 1974 earthquake series [abs.], Earthquake Notes, 46, 38.
- Spottiswoode, S. M., and McGarr, A. (1975). Source parameters of tremors in a deep-level gold mine, Bull. Seism. Soc. Am., 65, 93-112.
- Stauder, W. (1975). Subduction of the Nazca Plate under Peru as evidenced by focal mechanisms and by seismicity, J. Geophys. Res., 80, 1053-1064.

ACKNOWLEDGMENTS

The P-wave arrival time data that form the basis for this study were ISC data cleaned up and reformatted by A. C. Johnston of CIRES.

Figure Captions

Figure 1. Reference map of region under study. The rectangle encloses the area shown in figures 2 through 10. Hypocenters of earthquakes originally estimated to lie on the reference map were redetermined. Broad arrow shows direction of motion of Nazca plate relative to South American plate (Minster *et al.*, 1974). The 3000-fathom isobath of the Peru trench is plotted.

Figure 2. All hypocenters of focal depth less than 150 km determined by the International Seismological Centre (ISC) for the region of study (Figure 1). Our inferences are based principally on the 37% of all shocks whose epicenters, after we had relocated them, were estimated to be precise to within 20 km at a 90% level of confidence. 27% of the epicenters that we determined had 90%-confidence ellipses whose semi-axes exceeded 50 km in length and are not plotted in figures 3-10. Stars indicate hypocenters of the major shocks of 661017, 700531, and 741003. Dotted lines on the section normal to the trench indicate source regions defined by the b-quality relocated hypocenters of figure 3.

Figure 3. Hypocenters of b-quality (semi-axes of 90%-confidence ellipses on epicenter less than 20 km long) and d-quality (semi-axes of 90% confidence ellipses on epicenter between 20 km and 50 km long) occurring in the outlined region in figure 1. We define the dashed line (in the section normal to the trench) as the boundary between the interface thrust zone and the W-zone. Focal depths are likely to be least reliable near the trench.

Figure 4. Aftershocks occurring within one month of the earthquake of 661017. Map region as in figure 1 and symbols as in figure 3. Principal aftershock zones are outlined on the basis of b-quality hypocenters. Also plotted are the aftershock regions of the yet-to-occur earthquakes of 700531 (70a, 70b) and 741003 (74a, 74b). Main shock hypocenter indicated by star.

Figure 5. Aftershocks occurring within one month of the earthquake of 700531. Map region as in figure 1 and symbols as in figure 3. Principal aftershock zones are outlined on the basis of b-quality hypocenters. Also plotted are the aftershock zones of the previous shock of 661017 and the yet to occur shock of 741003. Main shock hypocenter indicated by star.

Figure 6. Aftershocks occurring within one month of the earthquake of 741003. Map region as in figure 1 and symbols as in figure 3. Principal aftershock zones are outlined on the basis of b-quality hypocenters. Also plotted are the aftershock zones of the previous shocks of 661017 and 700531. Main shock hypocenter indicated by star.

Figure 7. Seismic activity for 1964 through 661016. Map region as in figure 1 and symbols as in figure 3. Main shock hypocenters (stars) and aftershock zones of 661017, 700531, and 741003 are shown.

Figure 8. Seismic activity occurring from 661118 through 700530. Mainshock hypocenters and aftershock zones of 700531 and 741003, and aftershock zone of 661017 are shown. Map region as in figure 1 and symbols as in figure 3.

Figure 9. Seismic activity occurring from 700702 through 741002. Aftershock zones of 661017, 700531, and 741003 are shown as is the hypocenter of 741003 (star). Map region as in figure 1 and symbols as in figure 3.

Figure 10. Shocks occurring from 1964 through 741003, seaward from the postulated boundary between the interface thrust zone and the W-zone. Aftershock zones of 661011 and the 74a aftershock zone of 741003 are shown, as is the hypocenter of the mainshock of 741003 (star). Dates of shocks near the 74a focal region are also marked. Map region as in figure 1 and symbols as in figure 3.

151

Figure 1

Figure 2

Figure 3

661017 -- 661117

Figure 4

Figure 5

741003 - 741103

Figure 6

Figure 7

Figure 8

700702 - 741002

Figure 9

Figure 10

640101-741002

SEISMICITY PATTERNS PRECURSORY TO LARGER EARTHQUAKES IN THE ADAK REGION

E. R. Engdahl

Branch of Global Seismology
U. S. Geological Survey
Denver, Colorado 80225

The purpose of this study is to examine the space-time history of earthquakes occurring within the Adak region, with particular emphasis on precursors to the magnitude 6.7 Adak earthquake of November 4, 1977. We will discuss first the distribution of magnitude 3.0 and greater shallow earthquakes that have been located using data from the Adak microearthquake network since mid-1974. We will then examine the historical seismicity magnitude 4.5 and greater to search for similarities in the patterns of occurrence of these larger events. Finally, we will describe a prominent seismic gap for larger teleseismically detected earthquakes that is presently being monitored by the network.

In Figure 1 are plotted all magnitude 3.0 and greater events between 0 and 50 km in depth that have been located by the Adak network since mid-1974. Solid circles are magnitude 4.5 and above that have been detected teleseismically. Previous studies have shown that the most seismically energetic zone along the arc in this region occurs at the intersection of the Aleutian ridge and terrace as can be seen in this figure. Hence, an examination of the space-time history of the seismicity of this seismotectonic feature may well reveal clues about seismological precursors to major events in the region. The magnitude 6.7 earthquake of November 4, 1977, aftershocks, and possible premonitory seismic activity are visible as the large cluster of small and larger magnitude events between about 175.25° and 176.25° W longitude. It is useful to present these

data in the form of the space-time diagram shown in Figure 2. Distance here is arc length with respect to a center of curvature defined by the trench and volcanic arc, referenced at a radius approximately along the band of seismicity. The magnitude 6.7 event and aftershocks are clearly seen as a linear feature in the upper right hand side of this figure. Solid circles again indicate magnitude 4.5 and greater events. The main shock appears to have broken a segment of the Aleutian arc about 40 km in length. This arc segment had been a seismic gap for larger, teleseismically detected earthquakes from at most mid-1974, when the network began operating, until August 1976, when a group of larger earthquakes occurred within the gap over a six-month period. The first two earthquakes in this group were magnitude 4.6 and 4.8 events and were not accompanied by any other significant seismic activity. The third earthquake in the group, however, was accompanied by a dramatic surge in microearthquake activity, which included the ultimate source region of the magnitude 6.7 main shock, that included many magnitude 3.0 and greater events. This was the last major activity within the rupture zone of the main shock, with the exception of a less intense surge of activity characterized by the group of events occurring just two months before the main shock. At about the time of this second surge some larger earthquakes also occurred at both ends of the gap; the largest of these were an m_b 5.0 and 4.5 on the west end and an m_b 5.4 on the east end. Since the November earthquake is the largest event to have been "captured" by the network since its installation, it is appropriate to search the historical record for similar patterns of teleseismically detected earthquakes.

The occurrence of a number of teleseismically detected earthquakes, that have been located independently using only local network data, offers a

unique opportunity to resolve historical seismicity in greater spatial detail than previously obtainable. This premise is based on the assumption that residuals to stations at teleseismic distances from locally determined events describe path effects that normally bias hypocenters located with only teleseismic data. Hence, these residuals can be used as "station corrections" to relocate historical seismicity by the "master event" technique. To test the method a well-recorded earthquake centrally located within the network was used as a master to iteratively relocate magnitude 4.5 and greater historical seismicity using data from the ISC bulletins. The relocated events are shown in Figure 3. Solid circles indicate events located using only local network data. Readings from the Adak (ADK) station, which has operated since about 1965, were also used but without correction. For this test the results were not very satisfactory. Historical seismicity that is now relocated on the Aleutian terrace has apparently been biased about 20-25 km southward. We know this bias exists because the Adak residual is highly negative (the P wave arriving earlier than predicted) from these hypocenters, but is only 0.1 sec and less from hypocenters lying within the previously identified seismotectonic band of activity along the arc. This problem will be examined in considerable detail in future studies, since it appears that teleseismic station corrections can vary extremely rapidly across the shallow thrust zone, especially for North American stations. Nevertheless, these locations of larger earthquakes have been used to prepare a space-time diagram similar to Figure 2. In Figure 4 are plotted all relocated earthquakes magnitude 4.5 and greater that have occurred within the Adak region since 1964. The horizontal line marks the point at which the Adak network began operating, so that all the locations shown above that line have been locally determined. The cluster located in

the upper right-hand corner of this figure consists of the teleseismic events previously described in relation to the magnitude 6.7 earthquake of November 4, 1977. Solid circles indicate magnitude 5.0 and greater earthquakes. The historical record shows a very low level of seismicity over the 40 km arc segment broken by this earthquake since at least 1964. Other larger events appear to occur along the arc either in clusters over a period of several months or over a short time interval. The only other earthquake of a size comparable to the 6.7 is a 7.1 event that occurred in May 1971, and is located in about the center of the figure. Over a 2-3 month period before this event eight magnitude 5 or greater earthquakes occurred within about 30 km of the main shock location. Prior to that the arc segment had not been very seismically active. The patterns of earthquake occurrence shown here are very difficult to decipher and will probably take a careful case-by-case study to gain any measure of understanding. Additional relevant data will be obtained by examination of historical Adak station records and seismograms for evidence of microearthquake swarms or increased levels of local seismicity in the region.

Finally, we wish to address the prominent seismic gap for larger teleseismically detected earthquakes located in the top center of Figure 4. We have discussed this gap previously in a special study of the seismicity beneath Adak Canyon. Along about 60 km of this arc segment not a single teleseismically detected earthquake has occurred since the network was installed in mid-1974. From Figures 1 and 2 we can see that magnitude 3 to 4.5 earthquakes also appear to occur at a slightly lower level compared to the regions east and west of the gap. If the size of the gap is any measure of the potential rupture zone for a future earthquake, then on the basis of arc segments in this region

that have broken in the past (Figure 4), we can expect a future earthquake here to be magnitude 7 or larger.

REFERENCES

- Engdahl, E. R., Seismicity and plate subduction in the central Aleutians, in Island Arcs, Deep Sea Trenches, and Back-Arc Basins, ed. M. Talwani and W. C. Pitman III, Amer. Geophys. Union, Maurice Ewing Series I, 259-272, 1977.
- Engdahl, E. R., and C. Kisslinger, Seismological precursors to a magnitude 5 earthquake in the central Aleutian Islands, J. Phys. Earth, Proc. U.S.-Japan Seminar Earthquake Precursors, v. 25 supplement, 5243-5250.
- La Forge, R. C., Tectonic implications of seismicity in the Adak Canyon region, central Aleutians, M.S. thesis, University of Colorado, Boulder, CO, 1977.

Figure 1. Adak seismicity since July 1974, all events magnitude 3 and greater, less than 50 km deep. Solid circles are magnitude 4.5 and greater.

Figure 2. Space-time diagram of Adak seismicity since July 1974, same data and symbols as for Figure 1.

Figure 3. Relocated teleseismic events, magnitude 4.5 and greater, since 1964. Solid circles are events located using only local network data.

Figure 4. Space-time diagram of all events in Adak region since 1964, magnitude 4.5 and greater. Solid circles are magnitude 5 and greater.

THE OAXACA GAP: A CASE HISTORY

Tomas Garza and Cinna Lomnitz¹

An attempted use of seismic gap observations to predict a large earthquake in Oaxaca, Mexico is discussed. The observations were initially published in a scientific journal and were subsequently distorted by nonscientists, who predicted a major earthquake and tsunami to take place at Pinotepa Nacional, Oaxaca on 23 April 1978. Public reaction and property losses sustained by individuals and communities were comparable to those expected from an actual earthquake. No decrease in the number of felt events had been detected by the population of Pinotepa Nacional in recent years. A revision based on epicenter data from the NOAA data file showed that the probability that the number of earthquakes in two consecutive time intervals of a stationary Poisson process differs by an amount which would be reported as a "seismic gap" is of the order of 5% for Oaxaca. This means that "seismic gap" would be frequently observed even in relatively limited data runs. The possibility of detecting a true interval of abnormal quiescence in a random earthquake sequence appears to be fairly remote, unless other methods including a long-term seismicity analysis are adopted.

¹ Instituto de Investigaciones en Matematicas Aplicadas y en Sistemas, Universidad Nacional Autonoma de Mexico, Apartado 20-726, Mexico 20, D.F.

INTRODUCTION

The State of Oaxaca, Mexico, is among the most highly seismic regions in the world. Damaging large earthquakes, such as the shock which was destructive in the town of Pinotepa Nacional on 2 August 1968 ($M=7.5$), are not infrequent. The discovery by Ohtake et al. (1977) of a very clearly outlined seismic gap "starting in mid-1973, followed by complete quiescence" until the present time, was therefore believed highly significant. This was the first time that an announcement of an impending large earthquake was published in a scientific journal; however, this step seemed justified by the striking evidence presented by the authors (fig. 1). The "most probable characteristics" of the earthquake predicted by Ohtake et al. were as follows:

$$M = 7\frac{1}{2} \pm \frac{1}{4}$$

$$\phi = 16.5^\circ \pm 0.5^\circ\text{N}$$

$$\lambda = 96.0 \pm 0.5^\circ\text{W}$$

An occurrence time was not predicted, but it was stated that the current seismicity gap was already "substantially longer than the precursor times of the previous Oaxaca earthquake". In speaking of "the impending earthquake", the authors evidently believed the predicted shock to be imminent.

EFFECT OF PUBLICITY ON LOCAL POPULATION

On 7 February 1978 two residents of Las Vegas, Nevada sent a letter to the President of Mexico which contained the following prediction "based on demonstrated scientific facts":

"EARTHQUAKE IN THE STATE OF OAXACA IN THE TOWN OF PINOTEPA ON 23 APRIL 1978, AND LARGE QUANTITIES OF WATER CAUSING FLOODING".

A copy of this letter reached the office of the Mayor of Pinotepa a few days later. Some of the effects of the "prediction" were described in the local Pinotepa newspaper as follows: "After this announcement there has been a tremendous commotion on the Oaxaca Coast, to the point where many persons are fleeing their homes to emigrate to other towns in Mexico... The psychosis caused by the alarming news has induced them to sell their properties to the highest bidder thus destroying their homes... At first it was a speculative news item, but so much has been written about it that it has brought damage to all of Oaxaca as well as to the neighboring States of Guerrero, Michoacan, Puebla and others... Unfortunately there has been panic, particularly in Pinotepa and nearby coastal towns, and this is understandable because no one wishes to endanger their families; some local people have already sold their properties and people with money are buying land... One wonders: Who are these people picking up cheap real estate along the Oaxaca coast?" (Costa, Pinotepa, 21 April 1978.)

Meanwhile, a UPI press release from Austin, Texas, headlined in the Mexican dailies ("Texas U Predicts Big Mexico Quake", front page of The News, Mexico City, 10 April 1978), appeared to lend legitimacy to the "prediction" made for April 23. The press report did not contain a specific date, but it was studded with phrases such as "A massive earthquake will occur soon in the state of Oaxaca", "UT researchers expect the quake to be stronger than those that shook Managua and Guatemala", and so on. Verbatim quotes attributed to a reputable U.S. scientist appeared to confirm the urgency of the earthquake threat to Oaxaca.

The American press release caused speculation to flare up from Acapulco to Salina Cruz. One Acapulco local newspaper hatched a fantastic story, complete with "geological sections", claiming a foreign power had buried half a dozen nuclear charges in a fault located off the Oaxaca coast, to be detonated on April 23 by remote control from a plane flying at an altitude of 15,000 feet. Unfortunately, this story was widely circulated if not actually believed. A surprisingly large number of people thought that oil or uranium had been discovered on the Oaxaca coast and that cheap leases were being sought by foreign nationals. The questions asked by press reporters reflected similar beliefs.

THE ANTICLAMAX

Sunday, April 23 was a hot, sunny day in Pinotepa. The Governor of the State of Oaxaca, Eliseo Jimenez Ruiz, had announced that he would be in attendance to preside over special festivities organized "to reassure the people of the Oaxaca Coast that nothing was going to happen on that date, with reference to the said earthquake" (Costa, Pinotepa, 21 April 1978). Folk dancing groups, musicians and politicians had been brought in from the State Capital.

The Mayor of Pinotepa, Mr. Moises Cabrera Tellez, told us that there had been no letup in seismic activity in recent years. "We feel some 50 or 60 earthquakes every year," he said. "If there had been a lull, people would notice and talk about it. We just had another shock four days ago." He proceeded to tell us that the two strongest earthquakes announce the beginning and the end of the rainy season every year.

The Mayor was indignant about the prediction, which he claimed had caused more damage to Pinotepa than the 1968 earthquake. Though he used strong and

profane language, he also claimed that the reports of widespread panic were exaggerated. "Those who left were mainly out-of-towners", he stated. "Only about 15% of the citizens of Pinotepa are economically well off and can afford to leave." A stroll through the town revealed that perhaps 20% of the homes were shuttered, indicating that the residents were out of town.

The Govenor arrived at 5 p.m. and proceeded to the Town Hall, where a special exhibit (including a tent used for emergency housing) had been prepared by the Office of Urban Emergencies of the federal government. At 17:40:02 local time, while the Govenor was being shown around the exhibit, and earthquake of intensity III to IV shook Pinotepa and startled the crowd inside the Town Hall. The shock felt like a nearby local and was not immediately recognized as an earthquake, though it caused the metal doors to vibrate audibly. The Govenor and his party were unperturbed and later denied they had felt an earthquake. The shock was located by NEIS about 100 km inland ($m=4.2$); actually, it felt considerably closer to Pinotepa, perhaps 30 km away at most.

The festivities proceeded as planned. Around 10 p.m. a merry public dance began in the town square. Shortly after midnight, the Govenor looked at his watch and decided that it was time to return to Oaxaca City, as the prediction had lapsed. While some local residents still took the small shock in the afternoon as proof that the prediction had been partially successful, everyone was relieved that no disastrous earthquake and no tsunami had occurred.

AN INDEPENDENT CHECK

Pinotepa residents had been unanimous in denying that the frequency of local earthquakes had significantly decreased in recent years. This informa-

tion induced us to repeat the gap-detection procedure used by Ohtake et al. (1977), omitting the depth restriction ($h < 60$ km) which they had arbitrarily imposed on the data. Our source, the NOAA earthquake file, was presumably compatible with the PDE information utilized by the Texas group. The results are shown in fig. 2 and Table 1.

Because of the errors in epicentral location and particularly in focal depth determination, very large in this region, the fact that nine events appear within the alleged gap area though below the 60 km depth limit is not necessarily significant. Some of the newly found events are close to the gap boundary and could easily be excluded by means of a minimal shift of this boundary. On the other hand, the locations could easily be in error by 50 km or more. The focal depths could be in error by similar amounts, thus making the 60-km restriction meaningless. In order to clarify our argument we decided to apply some statistical tests to the following questions:

- (a) Did a seismic gap in fact occur during the period of June 1973 to date?
- (b) Was the occurrence of seismic gaps correlated with the occurrence of strong earthquakes in this region?

Test 1. Using the NEIS data file from January 1960 through December 1975, we find the mean number of events per two-year period in the alleged gap area:

$$\lambda = 21.4$$

Ohtake et al. implicitly defined a seismic gap as a time interval of abnormal seismic quiescence preceded by an equal interval of seismic activity. The discovery of 9 events in the "gap" interval (June 1973-May 1975) does not necessarily disprove the existence of a seismic gap according to this definition, since the seismic activity has undoubtedly been quite low. For example, the earthquake of 23 August 1965, which is claimed as an example of gap-induced

event in the same area, was also preceded by 9 shocks in the two-year period of 1963-1965. However, the definition used by Ohtake et al. is unorthodox because earlier authors have defined gaps in terms of the occurrence or non-occurrence of large earthquakes (potentially capable of rupturing the entire gap) and not of very small earthquakes.

Following the suggestion made by Ohtake et al., we assume random occurrence of small earthquakes in the gap area at all times except before the occurrence of a large earthquake. Ohtake et al. stated that "the probability that this gap could occur by chance in a given 2-year period is about 1 in 36,000". In other works, the probability that probability that such a period of decreased activity would occur by chance was claimed to be insignificant. In the following we show that it is very substantially higher than assumed by Ohtake et al.

Suppose the seismicity in the area obeys a Poisson process and that the expected number of earthquakes in a 2-year period is λ . Denote by N_1 , and N_2 the number of earthquakes in two consecutive non-overlapping 2-year intervals. Then

$$P(N_1=j) = P(N_2=j) = e^{-\lambda} \frac{\lambda^j}{j!}, j = 0, 1, 2, \dots \quad (1)$$

and it is well known that

$$P(N_1 - N_2 = j) = e^{-2\lambda} \sum_{k=0}^{\infty} \frac{\lambda^{j+2k}}{k!(j+k)!} = e^{-2\lambda} I_j(2\lambda) \quad (2)$$

where $I_j(x)$ is the modified Bessel function of order j .

We have computed the values of $P(N_1 - N_2 < j \mid N_1 > N_2)$ for different values of λ and j , in the range of experimental values found for the Oaxaca "gap". The results are shown in Table 2. For $N_1 = 25$ (1971-1973) and $N_2 = 9$ (1973-1975) the observed difference ($j > 15$) will occur with probability 0.020 when the mean is 21.4 events per 2-year period. On the other hand, if one accepted the supposition made by Ohtake et al. that the seismicity during the 1971-1973 period

was about average for the region ($\lambda = 25$), the probability of observing a difference of 16 events or more with the subsequent interval rises to 0.030; the difference with the previous case is not significant enough to allow a good degree of resolution. Finally, if the activity during the first interval equals the mean of the process ($N_1 = 21$) and the next interval contains 9 events or less the probability of such an occurrence is 0.08.

TABLE 2

Probabilities of occurrence of differences
 $N_1 - N_2 > j \mid N_1 > N_2$ for a Poisson Process of mean λ

$j =$	10	11	12	13	14	15
$\lambda =$						
14	0.0511	0.0324	0.0199	0.0119	0.0069	0.0039
15	0.0595	0.0387	0.0245	0.0151	0.0090	0.0053
16	0.0681	0.0453	0.0294	0.0186	0.0114	0.0069
17	0.0768	0.0521	0.0346	0.0224	0.0141	0.0087
18	0.0855	0.0591	0.0400	0.0264	0.0170	0.0107
19	0.0943	0.0663	0.0456	0.0307	0.0202	0.0130
20	0.1030	0.0735	0.0513	0.0351	0.0235	0.0154
21	0.1116	0.0807	0.0572	0.0398	0.0271	0.0181
22	0.1202	0.0880	0.0632	0.0446	0.0308	0.0209
23	0.1286	0.0953	0.0693	0.0495	0.0347	0.0239
24	0.1370	0.1025	0.0754	0.0545	0.0387	0.0270
25	0.1452	0.1097	0.0815	0.0593	0.0428	0.0302

In conclusion, the probability of random occurrence of a "gap" such as the one observed in Oaxaca is of order 1% to 8% for a Poisson sequence: these figures are not so small as to eliminate the possibility of attributing the "gap" to a random fluctuation. The likelihood of such a random fluctuation is certainly much higher than the figure of 1:36,000 quoted by Ohtake et al. The occurrence of periods of "high" seismicity followed by periods of "low" seismicity is not uncommon in a random process: so much so that the possibility of discriminating between a random quiescence and a quiescence due to non-random causes is uncertain. Additional statistical criteria would probably have to be used, as the likelihood of finding a seismic gap on the basis of low activity level alone is unpromising.

Test 2 Ohtake et al. suggested that the length of the interval of quiescence preceding a large earthquake should be correlated with the magnitude of the earthquake.

Thus, we compile the statistics of time intervals to the preceding event for all earthquakes in the alleged gap area during the period 1963-1975. The correlation with magnitude is shown in Table 3.

Table 3

Magnitude	<u>Mean interval to preceding quake</u>
Less than 5.0	36.5 days
5.0-5.9	14.5 days
More than 5.9	22.3 days
Mean interval between events (all magnitudes)	34.8 days

Observe that there is no indication of an absence of seismicity preceding the larger events. Quite the opposite, the interval preceding major earthquakes is shorter than the mean interval. This may be due to the occurrence of foreshocks, but it would also be expected if the large events are inserted at random in a Poisson sequence of smaller events.

Test 3 Finally, let us test the hypothesis of the "periods of quiescence". As implicitly defined by Ohtake et al., a "period of quiescence" is an interval of the order of two years containing few events (up to 9, in their article), and followed by a large earthquake. According to these authors, the 1965 and 1968 earthquakes were both preceded by such "periods of quiescence".

We select all events of magnitude 5 and above in the alleged gap area, for the period 1963 - 1975 in which USGS magnitudes are available. For each of these events we find the duration of the "period of quiescence", i.e. the time interval of occurrence for the preceding nine events. The results are displayed in Table 4 (obvious aftershocks have been excluded).

Table 4

<u>Magnitude m</u>	<u>mean "period" of quiescence"</u>
5.00 (three events)	19 months
5.20 (one event)	14 months
5.50 (three events)	13 months
5.60 (three events)	14 months
6.00 (one event)	15 months
6.30 (one event)	11 months
6.90 (one event)	6 months

The maximum "period of quiescence" which occurred inside the alleged gap was 32 months, between 24 January 1962 and 10 September 1964. The next large earthquake ($m=5.50$) occurred four months and 6 events later. A major earthquake ($M_S=6.25$, Pasadena) occurred on 3 April 1965, ten events after the end of the maximum "period of quiescence". In conclusion, no regularities are observed in respect to "periods of quiescence"; in particular, there appears to be no correlation between "periods of quiescence" and the occurrence of large earthquakes in the Oaxaca region. Strictly speaking, the most recent major earthquakes in this region occurred on 15 January 1931 ($M=7,9$) and 17 June 1928 ($M=8$, with tsunami).

CONCLUSIONS

The evidence for a true seismic gap having developed after 1973 in the area bounded by 95.5°W to 98°W along the coast of Oaxaca, Mexico is inconclusive. Periods of "seismic quiescence" are expected to occur in any random sequence, and have in fact occurred on the coast of Oaxaca. There appears to be no significant correlation between intervals of quiescence in the low-magnitude range and the occurrence of large earthquakes.

The suggestions advanced by Ohtake et al. (1977) that "such a prolonged period of quiescence is clearly unusual for this region", and "the relative seismic quiescence near Oaxaca may be signaling the eventual occurrence of a strong earthquake", are therefore unsupported by evidence. Random fluctuations in seismicity are normal in all seismic regions and should not be construed a priori as portents of seismic disaster. The data show that 1972 was a highly seismic year for the Oaxaca coast and that it has been followed by six years of very low seismicity. But the significance of such an isolated observation can only be perceived in the context of a detailed analysis of

the long-range background seismicity of the region. There are many examples of periods of high background seismic activity preceded by long periods of quiescence: the large-magnitude events do not necessarily appear at the end of the period of quiescence. A complete analysis based on all available data for the region would appear to be required.

In attempting a prediction based on the seismic gap hypothesis, care should be taken in letting the data define the gap and not vice versa. It seems arbitrary, for example, to cut off the data at a certain depth without justifying such a decision on the basis of an analysis of the depth distribution of earthquakes in the region and the errors involved. If the earthquakes occur near the center of the "gap" area (as they did in early 1978 off the Oaxaca coast) there is little argument but when they occur near the border of the gap it makes some difference whether the gap is defined on the basis of normal lines to the trench axis or simply of a pair of meridians. Thought should also be given to ensure completeness and accuracy of the data. Local station bulletins should be checked for consistency, as the focal depths computed from distant stations are known to be unreliable.

Unless seismic gaps can be defined by stringent statistical criteria their unqualified use as predictors of large seismic events seems premature. Predictions based on the sole evidence of seismic gaps may lead to serious disruption of normality in large areas, with associated property damage comparable to that caused by an actual earthquake.

Finally, the interpretation of forerunner phenomena requires some use of basic statistical concepts, including an analysis of errors in location

and focal depth. The statistical implications of the seismic gap hypothesis have been insufficiently recognized or explored. As seismologists we cannot doubt that a large earthquake will occur in Oaxaca sooner or later; but as statisticians we must seriously question the propriety of attaching a predictive significance to observations based on loosely defined concepts such as "gaps" or "quiescence", unless and until more specific supporting evidence becomes available. In particular, the relevance of the occurrence or non-occurrence of small earthquakes to the prognosis of major shocks has not been demonstrated, either in Oaxaca or in other regions. More extensive research is required before tentative forecasts based on this kind of observations can be discussed in the mass media.

ACKNOWLEDGEMENTS

We thank Arq. Mario Sosa, Director General of Urban Emergencies in the Department of Human Settlements and Public Works, for his valuable information and support. Access to important documents was kindly provided by Mr. Moises Cabrera Tellez, Mayor of Pinotepa Nacional.

REFERENCE

Ohtake, M., Matumoto, T., and Latham, G.V. (1977). Seismicity gap near Oaxaca, Southern Mexico as a probably precursor to a large earthquake, Pure and Appl. Geoph. 115, 375-385.

(a) Jun. 1971 - May 1973

(b) Jun. 1973 - May 1975

ALL DEPTHS

DEVIATORIC STRESSES AND EARTHQUAKE OCCURRENCE AT THE OUTER RISE

Thomas C. Hanks

U. S. Geological Survey
345 Middlefield Road
Menlo Park, California 94025

ABSTRACT

A straightforward, if approximate comparison between the strain energy (in bending) input rate to the continuously developing outer rise system and the elastic strain energy release rate occurring in the form of normal-faulting earthquakes in the lithosphere offshore of oceanic trenches requires that the extensional strains of $\sim 10^{-2}$ that exist in the upper half of the "elastic plate" indeed be elastic--and thus the existence of kilobars of deviatoric stress in the outer rise system. This result is conditioned to the assumption of a steady, flexure origin of the outer rise, as is often taken in modeling studies of bathymetry and gravity data. For the purposes to which it is put here, this assumption, at best, is likely to be valid only on a time scale corresponding to several cycles of strain accumulation and release associated with great underthrusting earthquakes onshore of the trench axis; the decade or so of reasonably well-determined and complete seismic observations that are presently available for the outer rise region represents only a small fraction of this time scale. Secondly, and in any case, "lithospheric normal-faulting earthquakes", of which the principal examples are the Sanriku (Mar. 3, 1933) and Java Trench (Aug. 19, 1977) earthquakes, cannot arise from tectonic stresses due to bending and thus certainly argue against the flexure origin of the outer rise locally and

perhaps globally as well. Whether or not these two earthquakes constitute compelling evidence against the general validity of the flexure origin of the outer rise is as yet moot, even if separate bathymetric, gravimetric, and seismologic evidence provide substantial support for it.

INTRODUCTION

The magnitude and orientation of deviatoric stresses in the Earth's crust and uppermost mantle is certainly one of the major outstanding problems in solid earth geophysics. The central issue of whether these deviatoric stresses, as averaged over the outer several tens of kilometers of the Earth, are a kilobar or perhaps somewhat greater or are 100 bars or perhaps somewhat less has been addressed, at least implicitly, in numerous and highly diverse studies, including investigations of regional gravity anomalies (Jeffreys, 1952; Banks *et al.*, 1977), earthquake stress drops (Chinnery, 1964; Hanks, 1977), the absence of a heat flow anomaly localized to the San Andreas fault (Brune *et al.*, 1969, Lachenbruch and Sass, 1973), deformation of the outer rise (Hanks, 1971; McKenzie, 1977), rock failure at mid-crustal conditions of pressure and temperature (Stesky and Brace, 1973; Byerlee, 1978), plate motion modeling (Forsyth and Uyeda, 1975; Davies, 1978), high pressure - high temperature flow laws (Carter, 1976; Kirby, 1977), petrofabric piezometry (Kohlstedt, 1978), and *in situ* stress measurements (McGarr, 1978); the issue has not, however, been decided. As a direct consequence of this uncertainty, so too are such fundamental issues as the driving mechanism of plate tectonics and the energetics of crustal faulting also unresolved.

Most of the evidence in support of the high-stress alternative is, in one sense or another, dependent on rheological assumptions. One may naturally and legitimately question whether inferences drawn from short-term laboratory studies of the fracture and flow of rocks constitute a correct description of the long-term material deformation occurring in the Earth's crust and uppermost mantle. More specifically, are inferences drawn from regional-scale gravity anomalies and deformation patterns at the outer rise correct in view

of the recent suggestion of Banks et al. (1977) that long-term elastic moduli may be substantially less than those measured seismically?

On the other hand, the principal line of evidence in support of the low-stress alternative, the absence of a heat flow anomaly localized to the San Andreas fault, is not dependent on rheological assumptions. So too, the constancy of earthquake stress drops is also independent of rheological assumptions, but while this important result is an oft used argument for the low-stress alternative, it is no less consistent with the high-stress hypothesis (Hanks, 1977).

This study develops a rheologically independent argument that elastic extensional strains of $\sim 10^{-2}$ exist in the upper 20 or so kilometers of the oceanic lithosphere offshore of oceanic trenches. The essence of the argument is its simplicity: the rate of elastic strain energy released by normal-faulting earthquakes within the oceanic lithosphere offshore of the trench axis must be less than the rate of strain energy invested in the continually developing outer rise.

THE OUTER RISE

The outer rise is a local bathymetric maximum (e.g. Chase et al., 1970), with a highly coincident gravimetric (e.g. Watts and Talwani, 1974) and structural (e.g. Talwani et al., 1978) expression, that occurs in the oceanic lithosphere adjacent to most oceanic trenches. There is general agreement that the outer rise results from deformation of the oceanic lithosphere as it is deflected from the Earth's surface and subducted at the trench. There is also general agreement that the strains associated with the outer rise deformation are large, of the order of 10^{-2} , if not somewhat larger in localized regions. Because of this, there has been a great deal of disagreement about whether the response of the oceanic lithosphere to

this deformation is entirely or even mostly elastic and by consequence about the loading conditions and material parameters that describe this deformation. In a substantial literature developed in the past decade, a range of disparate opinions on these matters has been voiced (Walcott, 1970; Hanks, 1971; Le Pichon *et al.*, 1973; Watts and Talwani, 1974; Sleep, 1975; Caldwell *et al.*, 1976; Parsons and Molnar, 1976; De Bremaecker, 1977; McKenzie, 1977; McNutt and Menard, 1978; Chen and Forsyth, 1978; Liu and Kosloff, 1978; McAdoo *et al.*, 1978; and Melosh, 1978).

The outer rise, presumably, is a steady feature of the region off-shore of oceanic trenches, but because of the mass transport associated with subduction of the oceanic lithosphere, it requires a constant input of strain energy to deform the initially undeformed material further off-shore. An estimate of this input rate of extensional strain energy is

$$\frac{1}{2} \sigma \epsilon L \dot{W}, \quad (1)$$

the result of collapsing the strain energy integral per unit time to the simple geometry of the problem: σ is an average extensional stress and ϵ an average extensional strain through a depth d , nominally one-half the effective elastic thickness in the usage of Hanks (1971); L is the length of all oceanic trench-outer rise systems and W is the width of the outer rise region normal to the trench axis. The $(\dot{})$ indicates a reciprocal time scale of millions of years for a particle to pass from the undeformed state several hundred kilometers offshore of the trench axis to subduction there, passing through ϵ in doing so; thus $\dot{W}$ is the convergence velocity.

Beyond the use of simple geometry and average (both spatially and temporally) values, (1) is general, at least insofar as no restriction on elastic strains has been made. On the short-term and in local regions,

however, (1) is unlikely to be a complete description of the strain energy existing in the outer rise system. In particular, the strain accumulation and release cycles associated with great underthrusting earthquakes will significantly affect the strain energy input rate (1) for the offshore region, since the rates of these processes are not at all small, even if the incremental strain energy associated with any one underthrusting earthquake is indeed small compared to the total strain energy existing in the outer rise system. Presumably, when averaged over many such underthrusting cycles, the net contribution of this process to the outer rise system strain energy is zero, if on the long-term the outer rise is indeed a steady feature. This assumption shall be taken here, but it is also appropriate to be cautious about the implications of the available seismic observations for a much shorter time period, which we compare with (1) below.

Extensional elastic strain energy is released in the form of normal-faulting earthquakes that occur in the deformed oceanic lithosphere offshore of the trench axis. Its magnitude is, per year,

$$\frac{1}{2\mu k}(\sigma_1 + \sigma_2) (\Sigma M_0/\text{year}), \quad (2)$$

the result of differencing the elastic strain energy integrals over the earthquake source volume V before (average stress σ_1) and after (average stress σ_2) the earthquake and using the relation

$$M_0 = k\Delta\sigma V \quad (3)$$

where $\Delta\sigma$ is the earthquake stress drop ($\sigma_1 - \sigma_2$) and k is a constant that depends on the geometry of faulting but is approximately 1. M_0 is the seismic moment of any earthquake, and the summation is taken for all normal-faulting earthquakes along the outer rise system of length L and width W .

At least on the long-term, (1) must be greater than (2). In (2), it has also been assumed that the quantity $(\sigma_1 + \sigma_2)$ is the same, or at least does not vary greatly, for all earthquakes considered. This has been done on the basis of the gross similarity of the inferred deformation for many outer rises (e.g. Watts and Talwani, 1974) and the constancy of earthquake stress drops (Aki, 1972; Thatcher and Hanks, 1973; Kanamori and Anderson, 1975; Hanks, 1977), despite the considerable scatter in the observations. Then the unknown stresses cancel, at least explicitly, from the problem. That is, for

$$k \frac{\sigma}{(\sigma_1 + \sigma_2)} \simeq 1 \quad (4)$$

the condition that (1) $\geq$ (2) requires that

$$\Sigma M_0/\text{year} \leq \mu \epsilon d \dot{W} \quad (5)$$

For $\mu = 5 \times 10^{11}$ dyne-cm, $\epsilon = 10^{-2}$, $L = 50,000$ km, $d = 20$ km, and $\dot{W} = 10$ cm/year

$$\Sigma M_0/\text{year} \lesssim 5 \times 10^{26} \text{ dyne-cm/year.} \quad (6)$$

The right-hand side of (6) corresponds to about one $M_s \simeq 7$ earthquake each year for the entire length of the outer rise system. Subject to several particulars concerning outer rise seismicity addressed in the next section, this is about the cumulative seismic moment per year that occurs. Because the elastic strain energy released by these earthquakes is, by definition, derived from elastic strain fields, this rate of elastic strain energy release requires that the strains of $\sim 10^{-2}$ used to obtain (6) be elastic--which therefore requires the existence of kilobars of extensional stress at the outer rise.

OUTER-RISE EARTHQUAKES

In the seventh decade of this century, normal-faulting mechanisms for $M_s \gtrsim 6$ earthquakes have been reported for the outer rise region offshore of most of the major oceanic trenches including the Chile (Stauder, 1973), Peru (Stauder, 1975), Aleutian (Stauder, 1968), Kuril (Stauder and Mualchin, 1976), Mariana (Katsumata and Sykes, 1969), Philippine (Fitch, 1970), Java (Fitch, 1970), New Hebrides (Johnson and Molnar, 1972), and Tonga (Johnson and Molnar, 1972) trenches. Depending on the judgment exercised in the counting of normal-faulting earthquakes that occur close to but onshore of the trench axis--but still reflect extensional elastic strain energy release in the subducting oceanic lithosphere--there have been approximately 30 $M_s \gtrsim 6$, normal-faulting earthquakes along the oceanic trench-outer rise system in the 1960's. This literature-search accounting certainly does not include all such earthquakes but presumably includes most if not all of the largest ones. At a level of uncertainty consistent with that existing in (6) (a factor of 2 or so) and in the conversion of M_s to M_0 (also a factor of 2 or so), these earthquakes certainly represent $\Sigma M_0/\text{year} \simeq 5 \times 10^{26}$ dyne-cm/year, if not somewhat more, over this decade.

There are three aspects of this set of earthquakes, however, that complicate this apparently straightforward if approximate comparison made just above. First, the normal-faulting earthquakes of the outer rise region are not distributed evenly through it, nor are they even uniformly distributed through the more strongly deformed region between the outer rise crest and the oceanic trench, typically a distance of approximately 120 km (Chase et al., 1970; Watts and Talwani, 1974) across which, given the bathymetry, strains of 10^{-2} should be the typical condition. Rather, these earthquakes almost always occur within several tens of kilometers of the trench axis.

There are two possible reasons for this. First, the ambient strains due to bending may simply be the largest closest to the trench axis (although this does not follow from all postulated models for outer rise deformation). Second, the effects of strain accumulation and release associated with the underthrusting cycle will be greatest closest to the trench axis, and, as we have noted previously, these effects may and occasionally do (see below) play an important role in spatially and temporally local regions of the outer rise system. Just as importantly, however, the almost-so-strongly deformed region between 50 and 150 km offshore of the trench axis, where strains of $\sim 10^{-2}$ almost certainly exist, is notably aseismic, at least at the $M \gtrsim 6$ level.

Second, a significant number of these earthquakes have occurred in close spatial and temporal relationship with great underthrusting earthquakes onshore of the trench axis. The effect is clearest in the case of the normal-faulting mechanisms reported for the area offshore of the Aleutian trench by Stauder (1968); eight of those twelve earthquakes occurred within $1\frac{1}{2}$ years of the Rat Island (Feb. 4, 1965; $M_0 = 1.2 \times 10^{29}$ dyne-cm) earthquake, although two of these occurred well east of the onshore aftershock zone. Table 1 indicates how closely the offshore seismicity is related in time to the onshore seismicity. The offshore seismicity decays even more rapidly than the onshore aftershocks, strongly suggesting a causal relationship to the Rat Island earthquake, a possibility noted previously by Sykes (1971) and Melosh (1976).

Evidently, the Rat Island earthquake, the result of underthrusting the oceanic lithosphere beneath the Aleutian Island arc, imparted a pulse of extensional stress to the oceanic lithosphere in the offshore region, resulting in a significant but transient release of extensional strain energy

in the outer rise region. As noted in the last section, this source of strain energy has not been accounted for in (1), which may be remedied by not counting those earthquakes which result from it in (2) or (6). Otherwise, one must conclude on the basis of (6) and for the much smaller length of oceanic trench-outer rise system involved in this burst of seismicity that perhaps a century or so of earthquakes at the rate of (6) occurred in a single 18-month period. Nevertheless, this latter conclusion is probably the correct one, because the extensional stress increment should be much smaller than the bending stresses in the oceanic lithosphere; that is, the offshore earthquakes were indeed triggered by the Rat Island earthquake but nevertheless drew their elastic strain energy release principally from the ambient bending stresses. But if this is indeed the case, there are obvious hazards in defining the left-hand-side of (5) on the basis of only a decade of reasonably well-determined seismic observations. To some extent, however, this uncertainty is probably offset by combining data for many oceanic trench-outer rise elements, each in different stages of the underthrusting cycle.

Similarly, in the Chile trench, five normal-faulting mechanisms between 1962 and 1968 have been reported (Stauder, 1973) for the area offshore of the great Chile (May 22, 1960; $M_0 = 2 \times 10^{30}$ dyne-cm) earthquake. Further north in the Chile trench (between 20° and 30° S), however, only one such event was reported by Stauder (1973) and this in the midst of a number of shallow events indicating horizontal compression both onshore and offshore of the trench axis. It is perhaps significant that much of this latter area, for which an abundance of evidence for ambient horizontal compression approximately normal to the trench axis both close to the plate boundary (Stauder, 1973; Hussong *et al.*, 1975; Prince and Schwelller, 1978) and far

away from it (Mendiguren, 1971; Mendiguren and Richter, 1978) is one of the world's principal seismic gaps (Kelleher, 1972). Other, isolated events in this class of earthquakes may also be associated with other great under-thrusting earthquakes, most notably the Fox Island (Mar. 9, 1957; $M_0 = 5.9 \times 10^{29}$ dyne-cm), Alaskan (Mar. 28, 1964; $M_0 = 8.2 \times 10^{29}$ dyne-cm) and Kuril Islands (Oct. 13, 1963; $M_0 = 6.7 \times 10^{28}$ dyne-cm) earthquakes.

A third important feature of these offshore normal-faulting earthquakes is their magnitudes are, with two important exceptions, less than approximately 7 to $7\frac{1}{2}$. One of these is the Sanriku (Mar. 2, 1933) earthquake in the Japanese trench; it has been assigned M_0 of 4.3×10^{28} dyne-cm by Kanamori (1971). The second earthquake occurred near the eastern end of the Java trench on Aug. 19, 1977; it has $M_0 = 1.2 \times 10^{29}$ dyne-cm according to Stewart (1978). Both of these earthquakes have been described as lithospheric normal-faulting earthquakes; that is, for both, an extensional fracture occurred over most if not all of the lithospheric thickness of approximately 70 km.

The significance of these two earthquakes lies in the facts that (1) normal-faulting mechanisms that fracture the lithosphere to this depth cannot have their origin in tectonic stresses that arise solely from bending and (2) these two earthquakes dominate, in terms of ΣM_0 , the combined contributions of all the offshore normal-faulting mechanisms for the past 50 years, however infrequent they may be. If, for example, the Java trench-size event has a recurrence time, again for the entire length of the outer rise system, significantly less than 240 years, it can only be concluded that the stresses due to bending are of not much significance, either in the deformation of the oceanic lithosphere offshore of oceanic trenches or in its subduction. Conversely, one may argue, on the basis of existing bathymetric, gravimetric,

and seismologic data that a flexure origin for the outer rise is generally applicable. If this is the case, these two earthquakes, as important as they are in terms of subduction in proximity to their locations, must nevertheless represent quite special conditions in these areas that are not generally applicable elsewhere.

In the case of the Sanriku earthquake, Kanamori (1971) estimated $M_0 = 4.3 \times 10^{28}$ dyne-cm on the basis of teleseismic surface waves and local seismograms in Japan and a fault length $l = 185$ km and fault width $w = 100$ km on the basis of the aftershock distribution (for a fault plane dip of 45° fracture to depth of 70 km is implied). From these estimates and the relation between M_0 and the static faulting parameters

$$M_0 = \mu \bar{u} l w, \quad (7)$$

an average faulting displacement $\bar{u} = 3.3$ meters is obtained. Considerably larger faulting displacements have been inferred from tsunami observations, however, 11-12 meters by Hatori (1966) and 6.6 meters by Aida (1977). Provided that Kanamori's (1971) estimates for M_0 and l are correct--they have been questioned by Ben-Menahem (1977)--, these estimates for $\bar{u}$ suggest w and depth of faulting 2 to 4 times smaller than Kanamori's (1971) estimates for these quantities. Inasmuch as tsunami generation is particularly sensitive to the surface and near-surface faulting displacements and to the extent these displacements are generally less than faulting displacements at depth (often by a considerable margin), an even smaller w is possible.

Moreover, the Sanriku earthquake was preceded by a major episode of underthrusting in the onshore region in 1896 to 1898. This earthquake began with a "tsunami earthquake" in 1896 for which M_0 of 6.8×10^{28} dyne-cm has been given by Kanamori (1972). It was followed by four large aftershocks

(all with $M \geq 8$; Richter, 1958, p. 560). It seems likely that ΣM_0 for this sequence is at least 1×10^{29} dyne-cm, if not somewhat larger. This sequence of interplate earthquakes (see also Shimizaki, 1978) must have had some effect on the offshore region, in a manner analogous to the Rat Island earthquake, although the delay of 35 to 37 years and the size of the Sanriku earthquake set this situation somewhat apart. Finally, the recurrence time of this earthquake is 86 years, using (6) as a measure of it, but then it has been 45 years since it occurred. Given the uncertainty in the numerical estimate of (6), that this earthquake is nominally a once every 86 year event is of not much consequence.

In all of this, it would seem that the Sanriku earthquake, while certainly an unusual earthquake, is not an impossible one to accommodate within the framework developed here. The Java Trench earthquake of 1977 is something else again, having M_0 almost 3 times larger than the Sanriku earthquake but having a fault length, inferred from aftershocks, of 160 km, approximately the same for the Sanriku earthquake. Perhaps the Java Trench earthquake is a very infrequent earthquake. Or perhaps its occurrence is specially related to its location near the eastern end of the Java Trench, beyond which a continental collision is occurring. Or perhaps it indicates that deformation of the outer rise due to bending is far less important than has previously been suspected.

SUMMARY AND CONCLUSIONS

The special circumstance of the outer rise is that it permits an estimate of the average, absolute extensional strain existing in the outer rise region, given a steady, flexure origin of the outer rise. A decade of seismic observations suggests, through a comparison of the elastic strain energy release rate associated with normal-faulting earthquakes at or offshore of oceanic

14

trenches with the strain energy input rate that must be continually invested in the outer rise system, that the strains of $\sim 10^{-2}$ that exist in the outer rise region be elastic. In turn, this suggests that kilobars of extensional stress are an average, ambient condition of the upper 15-25 km of the oceanic lithosphere in the outer rise region.

This result has important implications for the strength of the Earth's crust and uppermost mantle on time scales of millions of years and thus on the mechanism for driving plate motions across the Earth's surface (Hanks, 1977), but it is conditional on the assumption of a steady, flexure origin of the outer rise in two important ways. First, even if the steady, flexure model is correct on the long-term, strain accumulation and release associated with the underthrusting earthquake cycles suggest that several hundred years of seismic observations should be used to define numerically the left-hand-side of (5). Second, the interpretation of the Sanriku and Java Trench earthquakes as lithospheric normal-faulting earthquakes is fundamentally inconsistent with the flexure origin of the outer rise, steady or otherwise, at least in those localities affected by these earthquakes. Whether or not these two earthquakes are compelling evidence against the flexure origin of the outer rise is as yet moot, a situation that is not likely to change soon. Conversely, if the separate bathymetric, gravimetric, and seismologic arguments for the flexure origin of the outer rise are accepted, these two earthquakes, as important as they must be for the subduction process in proximity to their locations, must nevertheless represent quite special conditions in these areas that are not generally applicable elsewhere along the outer rise system.

ACKNOWLEDGMENTS

I have enjoyed useful conversations with H. Kanamori, P. Molnar, J.C.

Savage, and W. Thatcher in developing the ideas herein. Unless otherwise specified, seismic moments associated with specific earthquakes have been taken from Kanamori (1977).

REFERENCES

- Aida, I., Simulations of large tsunamis occurring in the past off the coast of the Sanriku district, Bull. Earthq. Res. Inst. U. Tokyo, 52, 71-101, 1977.
- Aki, K., Earthquake mechanism, Tectonophysics 13, 423-446, 1972.
- Banks, R.J., R.L. Parker, and S.P. Huestis, Isostatic compensation on a continental scale: local versus regional mechanisms, Geophys. J. R. astr. Soc. 51, 431-452, 1977.
- Ben-Menahem, A., Renormalization of the magnitude scale, Phys. Earth Planet. Interiors, 15, 315-340, 1977.
- Brune, J.N., T.L. Henyey, and R.F. Roy, Heat flow, stress, and rate of slip along the San Andreas fault, California, J. Geophys. Res. 74, 3821-3827, 1969.
- Byerlee, J.D., Friction of rocks, PAGEOPH, in press, 1978.
- Caldwell, J.G., W.F. Haxby, D.E. Karig, and D.L. Turcotte, On the applicability of a universal elastic trench profile, Earth Planet. Sci. Lett. 31, 239-246, 1976.
- Carter, N.L., Steady state flow of rocks, Rev. Geophys. Space Phys., 14, 301-310, 1976.
- Chase, T.E., H.W. Menard, and J. Mammerrickx, Bathymetry of the North Pacific, The Regents of the University of California, 1970.
- Chinnery, M.A., The strength of the Earth's crust under horizontal shear stress, J. Geophys. Res. 69, 2085-2089, 1964.
- Chen, T., and D.W. Forsyth, A detailed study of two earthquakes seaward of the Tonga trench: implications for mechanical behavior of the oceanic lithosphere, J. Geophys. Res., in press, 1978.
- Davies, G.F., The roles of boundary friction, basal shear stress and deep

- mantle convection in plate tectonics, Geophys. Res. Lett. 5, 161-164, 1978.
- De Bremaecker, J.-C., Is the oceanic lithosphere elastic or viscous?, J. Geophys. Res. 82, 2001-2004, 1977.
- Fitch, T.J., Earthquake mechanisms and island arc tectonics in the Indonesian-Philippine region, Bull. Seismol. Soc. Amer. 60, 565-591, 1970.
- Forsyth, D., and S. Uyeda, On the relative importance of the driving forces of plate motion, Geophys. J. Roy. astr. Soc. 43, 163-200, 1975.
- Hanks, T.C., The Kuril trench - Hokkaido rise system: large shallow earthquakes and simple models of deformation, Geophys. J. Roy. astr. Soc. 23, 173-189, 1971.
- Hanks, T.C., Earthquake stress drops, ambient tectonic stresses and stresses that drive plate motions, PAGEOPH 115, 441-458, 1977.
- Hatori, T., Vertical displacement in a tsunami source area and the topography of the sea bottom, Bull. Earthq. Res. Inst. U. Tokyo 44, 1449-1464, 1966.
- Hussong, D.M., M.E. Odegard, and L.K. Wipperman, Compressional faulting of the oceanic crust prior to subduction in the Peru-Chile trench, Geology 3, 601-604, 1975.
- Jeffreys, H., The Earth, pp. 392, Cambridge University Press, 1952.
- Johnson, T., and P. Molnar, Focal mechanisms and plate tectonics of the southwest Pacific, J. Geophys. Res. 77, 5000-5032, 1972.
- Kanamori, H., Seismological evidence for a lithospheric normal faulting - the Sanriku earthquake of 1933, Phys. Earth. Planet. Interiors 4, 289-300, 1971.
- Kanamori, H., Mechanism of tsunami earthquakes, Phys. Earth. Planet. Interiors 6, 346-359, 1972.
- Kanamori, H., The energy release in great earthquakes, J. Geophys. Res. 82,

2981-2987, 1977.

- Kanamori, H., and D.L. Anderson, Theoretical basis of some empirical relations in seismology, Bull. Seismol. Soc. Amer. 65, 1073-1096, 1975.
- Katsumata, M., and L.R. Sykes, Seismicity and tectonics of the western Pacific: Izu-Mariana-Caroline and Ryuku-Taiwan regions, J. Geophys. Res. 74, 5923-5948, 1969.
- Kelleher, J.A., Rupture zones of large South American earthquakes and some predictions, J. Geophys. Res. 77, 2087-2103, 1972.
- Kirby, S.H., State of stress in the lithosphere: inferences from the flow law of olivine, PAGEOPH 115, 245-258, 1977.
- Kohlstedt, D.L., Deformation-induced microstructures: a powerful new measure of geologic stress levels, EOS 59, 375, 1978 (abstract).
- Lachenbruch, A.H., and J.H. Sass, Thermomechanical aspects of the San Andreas fault system in Proceedings of the Conference on Tectonic Problems of the San Andreas fault system, R.L. Kovach and A. Nur, eds., Stanford University Publications, 206-213, 1973.
- Le Pichon, X., J. Francheteau, and J. Bonnin, Plate Tectonics, pp. 300, Elsevier, Amsterdam.
- Liu, H.-P., and D. Kosloff, Elastic-plastic bending of the lithosphere incorporating rock deformation data, with application to the structure of the Hawaiian archipelago, Tectonophysics, in press, 1978.
- McAdoo, D.C., J.G. Caldwell, and D.L. Turcotte, On the elastic - perfectly plastic bending of the lithosphere under generalized loading with application to the Kuril trench, Geophys. J. Roy. astr. Soc., in press, 1978.
- McGarr, A., and N.C. Gay, State of stress in the Earth's crust, Ann. Rev. Earth Planet. Sci., in press, 1978.

- McKenzie, D., Surface deformation, gravity anomalies and convection, Geophys. J. Roy. astr. Soc. 48, 211-238, 1977.
- McNutt, M., and H.W. Menard, Lithosphere flexure and uplifted atolls, J. Geophys. Res. 83, 1206-1212, 1978.
- Melosh, H.J., Nonlinear stress propagation in the Earth's upper mantle, J. Geophys. Res. 81, 5621-5632, 1976.
- Melosh, H.J., Dynamic support of the outer rise, Geophys. Res. Lett. 5, 321-324, 1978.
- Mendiguren, J.A., Focal mechanism of a shock in the middle of the Nazca plate, J. Geophys. Res. 76, 3861-3879, 1971.
- Mendiguren, J.A., and F.M. Richter, On the origin of compressional intra-plate stresses in South America, Phys. Earth Planet. Interiors 16, 318-326, 1978.
- Parsons, B., and P. Molnar, The origin of outer topographic rises associated with trenches, Geophys. J. Roy. astr. Soc. 45, 707-712, 1976.
- Prince, R.A., and W.J. Schweller, Dates, rates and angles of faulting in the Peru-Chile trench, Nature 271, 743-745, 1978.
- Shimizaki, K., Correlation between intraplate seismicity and interplate earthquakes in Tohoku, northeast Japan, Bull. Seismol. Soc. Amer. 68, 181-192, 1978.
- Sleep, N.H., Stress and flow beneath island arcs, Geophys. J. Roy. astr. Soc. 42, 827-857, 1975.
- Stauder, W., Tensional character of earthquake foci beneath the Aleutian trench with relation to sea-floor spreading, J. Geophys. Res. 73, 7693-7701, 1968.
- Stauder, W., Mechanism and spatial distribution of Chilean earthquakes with relation to subduction of the oceanic plate, J. Geophys. Res. 78, 5033-5061, 1973.
- Stauder, W., Subduction of the Nazca plate under Peru as evidenced by focal

- mechanisms and by seismicity, J. Geophys. Res. 80, 1053-1064, 1975.
- Stauder, W., and L. Mualchin, Fault motion in the larger earthquakes of the Kurile-Kamchatka arc and of the Kurile-Hokkaido corner, J. Geophys. Res. 81, 297-308, 1976.
- Stesky, R.M., and W.F. Brace, Estimation of frictional stress on the San Andreas fault from laboratory measurements, in Proceedings of the Conference on Tectonic Problems of the San Andreas fault system, R.L. Kovach and A. Nur, eds., Stanford University Publications, 206-213, 1973.
- Stewart, G.S., Implications for plate tectonics of the Aug. 19, 1977, Indonesian decoupling normal-fault earthquake, EOS, 59, 326, 1978 (abstract).
- Sykes, L.R., Aftershock zones of great earthquakes, seismicity gaps, and earthquake prediction for Alaska and the Aleutians, J. Geophys. Res. 76, 8021-8041, 1971.
- Talwani, M., P. Buhl, P.L. Stoffa, and T.K. Kan, The application of multi-channel seismics to the study of the outer gravity high near the Japan Trench, EOS, 59, 321, 1978.
- Thatcher, W., and T.C. Hanks, Source parameters of southern California earthquakes, J. Geophys. Res. 78, 8547-8576, 1973.
- Walcott, R.I., Flexural rigidity, thickness, and viscosity of the lithosphere, J. Geophys. Res. 75, 3941-3954, 1970.
- Watts, A.B., and M. Talwani, Gravity anomalies seaward of deep-sea trenches and their tectonic implications, Geophys. J. Roy. astr. Soc. 36, 57-90, 1974.

SEISMICITY OF THE ALEUTIAN TRENCH
 Feb. 4, 1965 - June 30, 1973; 160°W to 170°E
 $h \leq 60$ km, $m_b \geq 5.5$

	'65	'66	'67	'68	'69	'70	'71	'72	'73($\frac{1}{2}$)	2/65-12/66	1/67-6/73
All Earthquakes	56	9	9	4	13	8	14	7	6	65	61
Trench Axis/Offshore	10(5)	3	0	0(1)	3	2	0	1	1	13	7

2/65-12/66 - 20% of events are Trench/Axis/Offshore (another 8% close)

1/67- 6/73 - 11% of events are Trench/Axis/Offshore (another 2% close)

... .. 1954-1955
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

Some Characteristics of Foreshocks

by

Lucile M. Jones

Peter Molnar

Department of Earth and Planetary Sciences
Massachusetts Institute of Technology
Cambridge, MA 02139

This is a copy of a paper submitted to Journal of Geophysical Research, March 1978. For citation, please use the JGR.

ABSTRACT

Foreshocks occur before a large fraction of the world's major ($M \geq 7.0$) earthquakes. Teleseismically located events before major earthquakes from 1914 to 1973 were considered together to examine average foreshock patterns. There is a suggestion that about three months before the mainshocks the seismicity in the general vicinity of them (within 100 km) increases to about twice the previous level. About five days before the mainshocks and apparently nearer the epicenters of them the activity begins to increase culminating in a final rapid acceleration of activity in the last day. The acceleration continues up to the time of the mainshocks, except for a possible temporary decrease around six hours before them. The number of foreshocks per day is approximately proportional to the inverse of time before mainshock. This relationship is essentially unrelated to the magnitude of the mainshock. The magnitude of the largest foreshock is also unrelated to the magnitude of the mainshock. In addition, pairs of major events are common. Ten percent of the world's major events are preceded by other major events within 100 km and three months. For foreshocks within each of three sequences studied, the ratio of the amplitudes of the P and S waves were approximately the same, suggesting that the faulting mechanisms are the same for events in each sequence. By assuming an inhomogeneous fault plane on which asperities fail by static fatigue, we derived an equation for accelerating premonitory slip as a function of time, which agrees with the observed time dependence of foreshocks.

INTRODUCTION

When present, foreshocks are the most obvious premonitory phenomenon preceding major earthquakes. If they could be recognized as such before a main event, they would be a very useful tool in earthquake prediction because they accurately pinpoint the time and location of the forthcoming earthquake. This was demonstrated by the successful prediction of the 4 February 1975 Haicheng earthquake wherein foreshocks played a crucial role [Raleigh et al., 1977; Wu Kai-tong et al., 1976; Zhu Fung-ming, 1976]. But even beyond their use in earthquake prediction, foreshocks are the most convincing evidence that we have of deformation of the earth before major earthquakes. Therefore an improved understanding of foreshocks may provide important information about precursory deformation. Motivated both by the possibility of using foreshocks as a short-term predictor and by the need to better understand the mechanical behavior of the nearby rocks immediately preceding major earthquakes, we studied the characteristics of foreshocks that occur in the few hours, days, and weeks before large events.

In presenting the results, we first discuss the temporal and spatial distribution of seismicity prior to mainshocks to define short-term foreshock patterns and to determine how common foreshocks are. We also consider the possible effect of various characteristics of the mainshocks on these patterns. We then examine the relative magnitudes of foreshocks and mainshocks and the radiation patterns of three foreshock sequences for possibly recognizable characteristics. Finally we compare the data with a simple mathematical expression derived for accelerating premonitory fault slip.

DATA

We considered foreshocks before three groups of mainshocks. The first is all major ($M \geq 7.0$) shallow (depth ≤ 100 km) earthquakes in the world from 1914 to 1949 (511 events), listed by Duda [1965]. We searched for foreshocks in the Bulletins of International Seismological Summary (ISS). The second group is all major shallow earthquakes in the world from 1950 to 1964 [Duda, 1965] and from 1965 to 1973 (250 events) [U.S. Earthquakes, 1965 to 1973]. For this group, earthquakes were taken from the catalogue of world events prepared by the National Oceanic and Atmospheric Administration (NOAA) and bulletins of the International Seismological Center (ISC). When they were available, published reports discussing foreshocks before individual earthquakes were also used. Except for the foreshocks mentioned in the published reports, all of the locations had been computed from teleseismic data. Because of the location capability of world-wide networks, the magnitudes of essentially all of these foreshocks are greater than 4 and usually greater than $4 \frac{1}{2}$ [Evernden, 1970, 1971]. The third group of mainshocks includes earthquakes occurring near Greece, with $M \geq 5.6$ for events between 1911 and 1965 and $M \geq 5.1$ between 1966 and 1973, and with foreshock sequences listed by Papazachos et al. [1967; Papazachos, 1975a].

In some situations, two or more major earthquakes occurred very close to each other in time and space. In such cases, it did not seem reasonable to call these events separate mainshocks. When the events occurred within three months and 100 kilometers of each other (within the most extreme limit of foreshock activity by criteria considered

below) we considered the series of events to be only one sequence. If the first of the large events had the largest magnitude its origin time was naturally used as that of the mainshock. If the magnitude of the first event was no more than 0.2 units of magnitude less than the later events, the series was considered to be a double event, and again the origin time of the first event was used. Only if the second event had a magnitude more than 0.2 units greater than the magnitude of the first event was the first event considered to be a foreshock. In such a case, the smaller events occurring between the two large events were not considered to be foreshocks. These events were most likely aftershocks of the large foreshock and including them as foreshocks could distort the more general patterns of foreshock activity. In some situations, there were so many such aftershocks that inclusion of them would have masked the pattern in the other data.

These multiple major events occurred quite frequently. Of 939 major sequences recorded since 1897, 92 included more than one event with magnitude greater than 7.0 and occurring within 3 months of each other. Of these 92 sequences, 67 were not mainshock-aftershock sequences as the first event was either smaller than, or no more than .4 units of magnitude greater than a later event.

TIME AND SPACE DISTRIBUTION OF FORESHOCKS

In studying the temporal variation in foreshock occurrence, it is necessary to define foreshocks. To obtain a crude impression of the frequency of occurrence, we first considered any earthquake a foreshock if its computed location was within 100 km and it occurred within 40 days of

the main event [see Jones and Molnar, 1976].

This definition is only a starting point for the discussion and other definitions of foreshocks are discussed below. One hundred kilometers was chosen because it allows inclusion of essentially all events that occurred within the rupture area of mainshocks with $M \sim 7$, given the errors in the locations. All locations were taken from the sources cited above. Forty days was arbitrarily chosen because it seemed to correspond to the common usage of the term foreshock in the literature. By these criteria, the percentage of mainshocks preceded by foreshocks increases steadily until the present (Figure 1). This observation is almost surely a result of poorer recording capability earlier in the century.

To recognize foreshock patterns it is necessary first to establish levels of background seismicity. To do this we examined the seismicity in the year before each of the mainshocks with foreshocks from the second data set. One year is obviously insufficient to determine long term trends in seismicity, but our purpose here is to analyze only the activity in the few weeks before the mainshocks, not years. For this situation, a year of data is enough to define a background level and to allow recognition of short term changes in earthquake occurrence. The Greek earthquakes were not used because Papazachos [1975a] did not list locations for the foreshocks and the earlier major earthquakes were excluded because of poor locations. Although the average patterns might be more difficult to discern with a smaller group of events, to minimize further the effects of mislocation, we also examined

a subset of the second group, those earthquakes occurring from 1964 to 1973, which are more reliably located than those before 1964.

The events occurring during the year before the mainshocks were separated into three groups by the distance between their computed locations and those of the mainshocks - 30, 60, and 100 kilometers. Since no individual sequence had enough events recorded to define temporal and spatial patterns of occurrence, all of the sequences were combined into one sequence. This was done by putting all of the sequences onto the same time axis where the origin is the time of occurrence of each mainshock. A histogram was made by plotting the total number of foreshocks occurring in each 24-hour period before the mainshocks (Figure 2).

There is a suggestion of three changes in level of seismic activity preceding major earthquakes. The most notable is the very large number of events that occur in the day immediately preceding the mainshocks very near the mainshock epicenters (mostly within 30 km). This large number of events is the culmination of an increasing frequency of occurrence near to the mainshocks beginning a few days before them. There is also a suggestion of an increase in activity that starts about three months before the mainshocks. This increase is not as pronounced as the other increases and seems to occur at greater distances from the mainshocks. The rate of activity in the earlier part of the year could be the normal rate of activity. Alternatively it could be an abnormally low rate that is part of a longer term foreshock sequence so

that the apparent increase in activity three months before might be a return to the normal level. This variation in activity with time shown in Figure 2 might represent an increase in activity in the rupture areas of the mainshocks three months before them, and a subsequent migration towards the epicenters of the mainshocks. Thus, three possible definitions of foreshocks are suggested - events occurring within three months and sixty kilometers, five days and thirty kilometers, or one day and thirty kilometers of the mainshocks. All events fulfilling any of these criteria and their mainshocks are listed in Table I, available on microfiche.

A plot by hour of foreshock activity in the last few days shows that the frequency of occurrence continues to increase towards the time of the mainshocks (Figure 3). This average pattern continues except for a lull in activity five to eight hours before the mainshocks, or, alternatively, a peak nine to twelve hours before. The peak is defined by foreshocks before 18 events, not by just a few mainshocks. This peak followed by a lull is of interest because Chinese seismologists reported a drop in foreshock activity four to eight hours before the Haicheng earthquake [Wu Kai-tong et al., 1976].

There are many factors that could distort the temporal pattern of foreshock activity in figures 2 and 3.

Dominance by a few sequences. One possible factor is that a large number of foreshocks from a few mainshocks could be dominating the plot so that it is not representative of "average" foreshock sequences. To test for this possibility, histograms were made for mainshocks with only one, two and three, four and five, and more than five foreshocks (Figure 4).

The peak in foreshock activity on the last day exists for all cases and clearly does not depend upon how many foreshocks occurred before any event. Although the increase in activity approximately five days before the mainshocks is less well defined (perhaps due in part to the smaller data base involved) and is clear only for events with more than one foreshock, the pattern in Figure 2 is not dominated by a few sequences with many foreshocks.

Dependence on Magnitude. A dependence of foreshock activity on magnitude also might affect the reliability of the patterns in Figure 2. To examine this possibility, sequences were plotted separately for mainshocks with $M < 7.0$ (Greek events only), $7.0 \leq M < 7.8$ and $M \geq 7.8$ (Figure 5a-c). The time dependences of these three sequences are very similar to each other and also to that of the combined sequence (Figure 5d). Power law equations fitted to these data (Figure 5) indicate that the largest earthquakes ($M \geq 7.8$) might have slightly longer foreshock sequences but the difference is slight. All of the sequences show the same pattern of seismicity - an increase in activity beginning about five days before the main events culminating in a prominent maximum on the day before the mainshocks.

This pattern of foreshock activity appears in the temporal pattern of all the groups of mainshocks considered in figures 2 through 5. Although variation in the physical properties of materials from region to region is one of the reasons why earthquakes are not all alike, we think that the consistency of the patterns shows that they are typical of foreshocks or at least representative of an average pattern.

Because there are changes in the level of seismic activity at different times prior to the mainshocks, there are different possible definitions of the term foreshock. Thus to establish how common foreshocks are, more than one time period of foreshock activity should be considered. Table 2 lists the percentage of mainshocks which were preceded by earthquakes within the time limits of the above stages of foreshock activity. Table 2 includes mainshocks with foreshocks which were too small to be located with teleseismic data, but were mentioned in published reports. For instance, we included the 1966 Congo [Lahr and Pomeroy, 1970], the 1952 Kern County and the 1954 Dixie Valley - Fairview Peak earthquakes [Richter, 1958]. There probably are many other major events with foreshocks too small to be located. It is noteworthy that none of the more than 500 foreshocks of the 1975 Haicheng earthquake [Wu Kai-tong et al., 1976] were located teleseismically. To minimize the effect of poor location capability, Table 2 also considers a subset of the mainshocks, from 1950 to 1973, those with more than two aftershocks located teleseismically.

We were unable to recognize any dependence of the presence or absence of foreshock activity on tectonic setting [Jones and Molnar, 1976]. There is a suggestion, however, that foreshocks are more common in some regions than others (see Figure 2 of Jones and Molnar, [1976]). For instance, all of the earthquakes occurring in the section of Japan trench off of Hokkaido were preceded by foreshocks while none of the earthquakes in the neighboring part of the trench off of Honshu were. Thus knowing the previous history of foreshock activity on a given fault could enhance the use of foreshocks as a predictive tool for major events in such regions.

TIME FUNCTIONS

The data in Figure 5 were fitted to both exponential and power law functions by linear regression. Unlike the data of Pho et al. [1976], exponential functions fit poorly and are probably not a good representation of the physical processes involved. The data fit much better to power law equations. The combined data showed that foreshock activity is proportional to time raised to the minus one power which agrees with the findings of Kagan and Knopoff [1978] and Papazachos [1974a, 1974b, 1975b]. Although the calculated exponent is more negative for larger magnitude mainshocks, the difference is slight and probably negligible.

RELATIVE MAGNITUDES OF FORESHOCKS AND MAINSHOCKS

The time dependence of foreshock activity does not appear to be sensitive to the magnitude of the mainshocks (Figure 5). If foreshocks are to be used for earthquake prediction, some other characteristic must be used to estimate the magnitude of the mainshock. Papazachos [1974a, 1974b, 1975b] and Wu Kai-tong et al. [1976] suggested that, in spite of a great deal of scatter, there is a linear relationship between the magnitudes of mainshocks and those of their largest foreshocks. Figure 6 explores this possibility using all of the data available to us. Even allowing for an error of 0.5 units in magnitude, we can see no relationship between the magnitudes of mainshock and largest foreshock. The only limits on the distribution of the magnitudes of the largest foreshocks appear to be that a foreshock be smaller than its mainshock and large enough to be located. We suspect

that the crude relations in Papazachos' and Wu Kai-tong et al.'s data are the result only of the limited magnitude range of their data. We thus conclude that foreshocks cannot be used to predict reliably the magnitude of an impending earthquake. This lack of correlation can be interpreted to mean that neither the area that slips nor the amount of slip during foreshocks is a constant fraction of the fault area or displacements caused by the mainshock.

AMPLITUDE RATIOS OF P AND S WAVES FROM FORESHOCKS

Jin Yan et al. [1976] suggested that the fault plane solutions of foreshocks are similar and that this aspect would be diagnostic of them. Because most foreshocks are too small to determine a fault plane solution, they suggested that the similarity could be examined by comparing the amplitudes of P and S waves recorded at one station. The ratio of these amplitudes depends upon three factors - the propagation path, the fault plane solution and the position of the station with respect to the fault plane. If the amplitudes are measured at only one station and the foreshock hypocenters are very close to each other, the amplitude ratios will depend only on the fault plane solution. Jin Yan et al [1976] found that the foreshocks of the 1975 Haicheng earthquake did have very similar amplitude ratios, as did Lindh et al [1978] for two of the three foreshock sequences that they studied, and Bolt et al [1977] for the Briones Hills earthquake, but not Engdahl and Kisslinger [1977] for an Aleutian event.

We studied three mainshocks with many foreshocks, the only ones that occurred since the World Wide Standardized Seismograph Network (WWSSN) was installed and that occurred sufficiently

close to a WWSSN station to be well recorded. Records from the station at Baguio (BAG) were used for the August 1, 1968 earthquake in the Philippines [Su, 1970] and records from the station at Athens University (ATU) were used for the December 5, 1968 earthquake near Greece and the March 28, 1969 earthquake in western Turkey [Papazachos, 1975a]. In all cases, the events preceding the mainshocks are too small to be located reliably. They are thought to be foreshocks from their approximate locations determined from recordings from a single station. Their times of occurrence and P and S wave amplitudes are listed in Table III. The maximum amplitudes of the P and S waves, plotted versus one another for the three sequences in Figures 7, 8 and 9, define straight lines passing through the origin. In fact, not only are the amplitude ratios the same, but in general the shapes of the signals are remarkably similar. The only points that deviate from the lines in Figure 7, 8 and 9 are those for a few foreshocks occurring more than 3 weeks before the earthquake in Greece. All of the other foreshocks occur within four days of the mainshocks.

Although not proof, these data are consistent with identical fault plane solutions for the foreshocks occurring within about four days of their mainshocks. The different amplitude ratios of the earlier foreshocks of the event in Greece could be due either to differing fault plane solutions of different locations, as Engdahl and Kisslinger [1977] report for the Aleutian event.

DISCUSSION

One purpose of this study is to consider possible implications of foreshock activity for precursory deformation. We think the characteristics of foreshocks mentioned here are suggestive of accelerating premonitory fault slip. Foreshocks, being earthquakes, definitely represent slip somewhere; the spatial distribution and identical amplitude ratios (for the few events that we could study) suggest that in general the slip takes place on the same plane for each. We presume that this plane is the one that ruptures during the mainshock, although the data of Bolt et al. [1977] and Engdahl and Kisslinger [1977] suggests that this is not always the case. The temporal distribution implies that wherever the slip is, it is accelerating. The lack of correlation between the magnitudes of the mainshock and foreshocks suggests that the area of slip during the foreshocks is not a constant fraction of the rupture area of the mainshock. The foreshocks probably involve slip only in the vicinity of the mainshock hypocenter where the mainshocks are triggered. To illustrate how accelerating premonitory fault slip can explain these properties of foreshocks, we derive a simple expression for the development of an instability to compare with these data. Although the particular instability that we discuss was motivated by the observed characteristics of foreshocks, we do not claim that the data prove that it is an accurate description of what actually occurs.

Assume that the fault plane is held in place at a

number of asperities. This could also represent a plane of inhomogeneous stress where the points of highest stress are modeled by asperities. During the short time period involved (a few months or less) the tectonic stress acting on the whole fault plane can be assumed to be constant. If A_o is the rupture area, A_u is the portion of A_o that has not yet slipped, σ_o is the tectonic shear stress and σ is the average shear stress acting on the unslipped portion, these assumptions can be expressed as

$$A_o \sigma_o = A_u \sigma \quad (1)$$

The number of asperities, N , can be related to A_u by

$$A_u = N\bar{A} \quad (2)$$

where $\bar{A}$ is the average area of an asperity. As slip occurs on some segments of the fault (rupture of some asperities), the stress on the remaining unslipped segments will increase and the increased stress will cause further slip on other asperities. Eventually, the rate of failure of asperities increases rapidly causing an instability. With some additional, simple assumptions we can derive a differential equation relating the change in unslipped area to time.

Presumably, the lifetime of an asperity depends upon the stress acting on it. For instance, in the static fatigue of quartz in uniaxial compression Scholz [1972] showed that such a dependence could be expressed in terms of the mean time before failure

$$\langle t \rangle = T \exp(-k\sigma) \quad (3)$$

where σ is the differential stress and T and k are constants that can be determined experimentally [see also,

Scholz and Martin, 1971]. T is a time constant that depends upon the fluid pressure in the rock. Scholz [1972] did not evaluate T and it is difficult to calculate it for conditions in the earth. Scholz did evaluate k , which is a measure of the rock's strength. We assume that the lifetime of asperities on a fault obeys an equation like (3).

If static fatigue is a random process, then for a given stress we might expect that the number of asperities that break per unit time will be proportional to the number of unbroken asperities. Therefore,

$$\frac{dN}{dt} = - \frac{N}{t_0}$$

where t_0 is the average lifetime of an asperity. If $t_0 = \langle t \rangle$ defined in (3), then

$$\frac{dN}{dt} = - \frac{N}{T \exp(-k\sigma)}$$

Combining this equation with (2) to express this failure rate in terms of area, we have

$$\frac{dA_u}{dt} = - \frac{1}{T} A_u \exp(k\sigma) \quad (4)$$

Introducing a dimensionless variable, $x = \frac{A_0}{A_u} = \frac{\sigma}{\sigma_0}$,

$$(4) \text{ becomes } \frac{dx}{dt} = \frac{x}{T} \exp(k\sigma_0 x).$$

which can be integrated to give

$$\frac{t}{T} = \ln(k\sigma_0 x) + \sum_{n=1}^{\infty} \frac{(-k\sigma_0 x)^n}{n \cdot n!} - \frac{C}{T} \quad (5)$$

While the tectonic stress on the fault plane remains constant, as slip occurs on some asperities, the actual stress on the locked sections of the fault increases as shown in Figure 10.

This equation can also be expressed in terms of area.

In Figure 11, a nondimensional area variable, $(\frac{1}{k\sigma_0 x})$ is plotted as a function of a non-dimensional variable, time $\frac{t+C}{T}$. The former is related to the unslipped area, A_u , by

$$\frac{1}{k\sigma_0 x} = \frac{1}{k\sigma_0} \frac{A_u}{A_0} \quad \text{and to the area that has slipped, } A_s = A_0 - A_u, \text{ by}$$

$$\frac{1}{k\sigma_0 x} = 1 - \frac{A_s}{A_0}$$

The curve in Figure 11 is applicable only for a range of values of x : $1 \leq x < \infty$, corresponding to $A_0 \geq A_u \geq 0$. Notice that

$$\lim_{k\sigma_0 x \rightarrow \infty} \frac{\frac{\ln(k\sigma_0 x)}{\sum_{n=1}^{\infty} \frac{(-k\sigma_0 x)^n}{n \cdot n!}}}{\frac{\ln(k\sigma_0 x)}{\sum_{n=1}^{\infty} \frac{(-k\sigma_0 x)^n}{n \cdot n!}}} = -1 \quad (6)$$

Therefore as x approaches ∞ and A_u approaches zero, t approaches $-C$. The choice of C arbitrarily sets the origin or completion time of the process. The instability begins when $A_u = A_0$, or $x = 1$, for which $\frac{t}{T} = \ln(k\sigma_0) + \sum_{n=1}^{\infty} \frac{(-k\sigma_0)^n}{n \cdot n!} - \frac{C}{T}$ (7)

Below, we set $t = 0$ to correspond to the time of the mainshocks, so that the value of t in (7) gives the duration of the instability. Because of the limit in (6), the duration of the instability is less for larger values of σ_0 or k (higher tectonic stress or weaker asperities), than for smaller values of them.

If foreshocks are a manifestation of premonitory slip and the dependence on time of premonitory slip is approximately that of foreshock activity, then the time dependence of $\frac{A_s}{A_0}$ should match the cumulative plot of foreshock activity with time (Figure 11). These two curves are remarkably similar. Expressing the time scale in terms of days allows values of $T = 3000$ days and the constant of integration, $C = 1732$ days to be assigned. This value C corresponds with setting the time of the mainshock at $t = 0$.

Because of the dependence of T on fluid pressure, temperature and pressure, it is probably futile to attempt to compare this estimate of T with those calculable from laboratory data.

One feature of the fit of (5) to the data in Figure 11 is that the mainshock does not occur until almost 100% of the area has slipped. This need not mean that almost 100% of the total rupture area slips before the mainshock. It appears that in the few days before the mainshock, foreshocks occur very close to the epicenter of the mainshock (Figure 2) [eg. Wu Kai-tong et al., 1976]. Perhaps premonitory slip occurs only over a small fraction of the total area, which could be termed a triggering area. A rupture of the triggering area then propagates through to the rest of the fault plane [eg., Brune, 1978].

The duration of the instability is related to the tectonic stress and the strength of the asperities. Using the fit of (5) to the foreshock data (Figure 11) for $k\sigma_0 = 4$, the instability would begin to develop ten days before the mainshock but for $k\sigma_0 = 1$ or 0.2 , the instability would begin to develop 330 days or 2576 days (~ 7 years) before it. Scholz [1971] obtained a value of $k = 3.7 \text{ kbar}^{-1}$ for the a -axis of uniaxially loaded single quartz crystals. This value of k is for the case where σ is differential stress: if σ represents shear stress as in (5) then the value of k should be doubled. Using $k = 7.4 \text{ kbar}^{-1}$ in (5) we obtain estimates of the tectonic stress of 1081, 270 and 54 bars for $k\sigma_0 = 4, 2$ and 1 respectively. If the asperities were weaker than Scholz's single quartz crystals, such estimates of the tectonic stress would be lower.

Figure 11 and (5) describe a simplistic model strictly valid

only for a static situation. If this sort of instability takes place, the assumptions of static fatigue and setting $t_0 = \langle t \rangle$ in (3) are probably invalid as the ruptured area grows rapidly near the time of the mainshock. Nevertheless, the remarkably good fit encourages us to pursue it further.

CONCLUSION

Foreshocks occur before a large fraction of major earthquakes. There is a suggestion of a heightened level of activity in the general area of mainshocks about three months before them. Activity increases only slightly until about five days before the mainshock when it begins to increase rapidly, apparently near the mainshock epicenters, culminating with a peak in activity on the last day. On the last day the activity continues to increase until the time of the main event, except for a possible slight, and apparently temporary drop in activity a few hours before it. Neither the magnitudes of foreshocks nor the time dependence of foreshock activity seem to depend significantly on the magnitudes of the mainshocks. The magnitude of the largest foreshock can range from nearly the magnitude of the mainshock (twenty percent of the world's major earthquakes occur in series with at least one other major event) to being too small to locate. Judging by a few sequences studied, it seems likely that faulting mechanisms are usually the same for foreshocks within one sequence.

These findings suggest that foreshocks alone are insufficient for earthquake prediction. In many cases, they will not occur at all. When they do occur they might be recognized by their

temporal distribution or in some cases, by the similarity of their radiation patterns and therefore be of use in predicting the location and time of the impending mainshock. However, they may not provide information about the magnitude of the mainshock.

The frequency of foreshock occurrence shows that precursory deformation does occur before many major earthquakes. Many characteristics of foreshock activity could be explained if the deformation occurs by accelerating premonitory fault slip.

ACKNOWLEDGEMENTS

Some of the materials incorporated in this work were developed with support from the United States Geological Survey, Department of the Interior, Contract 14-08-0001-16758, and the Alfred P. Sloan Foundation Grant in Aid Research Fellowship, #7350. The views and conclusions contained in this paper are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of either the U.S. Government or the Alfred P. Sloan Foundation.

We would like to thank both Bernard Minster and Tom Heaton from the California Institute of Technology for valuable suggestions, and J. Ebel, T.C. Hanks, and A. Lindh for constructive criticisms of the manuscript.

REFERENCES

- Bolt, B.A., J. Stifler and R. Uhrhammer, The Briones Hills Earthquake Swarm of January 8, 1977 Contra Costa County, California, Bull. Seismol. Soc. America, 67, 1555-1564, 1977.
- Brune, J.N., Implications of earthquake triggering and rupture propagation for earthquake prediction based on premonitory phenomena, to be published in Proceedings of the Conference for "Fault Mechanics and its Relation to Earthquake Prediction," U.S. Geological Survey, 1978.
- Duda, S., Secular Seismic Release in the circum-Pacific belt, Tectonophysics, 2, 409-452, 1965.
- Engdahl and Kisslinger, Phenomena precursory to a magnitude 5 earthquake in the central Aleutian Islands, EOS Trans. Amer. Geophys. Un. 58, 433, 1977.
- Evernden, J.F., Study of regional seismicity and associated problems, Seismol. Soc. America Bull., 60, 393-446, 1970.
- Evernden, J.F., Location capability of various seismic networks, Seismol. Soc. America Bull., 61, 241-256, 1971.
- Jin Yan, Zhao Yi, Chen Yong, Yan Jia-quan, and Zhou Yu-ru, A characteristic feature of the dislocation model of foreshocks of the Haicheng earthquake, Lianoning Province, Acta Geophys. Sinica, 19, 156-164, 1976.
- Jones, L., and P. Molnar, Frequency of foreshocks, Nature, 262, 677-679, 1976.
- Kagan, Y. and L. Knopoff, Statistical study of the occurrence of shallow earthquakes, preprint, 1977.
- Lahr, J. and P.W. Pomeroy, The foreshock-aftershock sequence of the March 20, 1966 earthquake in the Republic of Congo, Seismol. Soc. America Bull., 60, 1245-1258, 1970.

- Lindh, A., G. Fuis and C. Mantis, Foreshock Amplitudes and fault plane changes: a new earthquake precursor?, submitted to Science, December 1977.
- Papazachos, B., On certain aftershock and foreshock parameters in the area of Greece, Annali di Geofis. 27, 497-515, 1974a.
- Papazachos, B., On the time distribution of aftershocks and foreshocks in the area of Greece, Pageoph. 112, 627-631, 1974b.
- Papazachos, B., Aftershock and foreshock sequences in the area of Greece during the period 1911-1973, Bull. Scientific Group for Space Res., 3, 1-44, 1975a.
- Papazachos, B., Foreshocks and earthquake prediction, Tectonophysics, 28, 213-226, 1975b.
- Papazachos, B., M. Delibasis, N. Liapis, G. Moumoulidis, and G. Purcaru, Aftershock sequences of some large earthquakes in the region Greece, Annali di Geofisica, 20, 1-93, 1967.
- Pho, H.-T., G. Wittinger and H. Haessler, Enregistrement et distribution temporelle des repliques, 80h apres le séisme destructeur du Frioul (Italie) du mai 1976, C.R. Acad. Sci. Paris 283 Sér. D., 611-614.
- Raleigh, C.B., et al., Prediction of the Haicheng Earthquake, EOS, Trans. Amer. Geophys. Un. 58, 236-272, 1977.
- Richter, C.F., Elementary Seismology, Freeman, New York, 1958.
- Scholz, C.H., Static Fatigue of Quartz, J. Geophys. Res., 77, 2104-2114, 1972.
- Scholz, C.H. and R.J. Martin, Crack growth and static fatigue in quartz, J. Amer. Ceram. Soc., 54, 474, 1971.
- Su, S.S., The Luzon Earthquake of 7 August 1968: A preliminary

report, Bull. Seismol. Soc. America, 59, 459-472, 1969.

United States Earthquakes, 1965 to 1973, (U.S. Government Printing Office, Washington D.C., 1967-1975).

Wu Kai-tong, Yue Ming-shang, Wu Huan-ying, Cao Xin-ling, Chen Hai-tong, Huang Wei-giong, Tian Kang-yuan, and Lu Shou-de, Certain characteristics of Haicheng earthquake ($M = 7.3$) sequence, Acta Geophys. Sinica, 19, 109-117, 1976.

Zhu Fung-ming, An outline of prediction and forecast of Haicheng earthquake of $M = 7.3$, Proceedings of the lectures of the seismological delegation of the People's Republic of China, (Jet Propulsion Laboratory, 1976).

TABLE I

Year		Date	G.M.T.	Location 1	Location 2	Mag
1914		none				
1915	m	Nov 01	07:23:43	38.0N 144.0E	38.0N 144.0E	7.8
	1	Oct 12	21:28:51		38.0N 145.0E	
1916	m	Mar 25	23:52:17	24.0N 12.4.0E	25.6N 126.0E	7.4
	1	Mar 01	18:01		25.3N 126.8E	
1917	m	Jul 29	14:32:15	41.0N 144.0E	41.0N 144.0E	7.0
	1	Jul 29	01:57:00		41.0N 144.0E	
1917	m	Aug 31	11:36:18	5.0N 75.0W	5.0N 75.0W	7.2
	1	Aug 30	13:24:10		5.0N 75.0W	
1918	m	Aug 15	12:18:12	5.5N 123.0E	5.4N 125.2E	8.3
	1	Aug 15	02:37:58		5.4N 125.2E	
	2	Aug 14	09:17:17		5.4N 125.2E	
	3	Aug 11	23:28:00		5.4N 125.2E	
1919	m	Dec 20	20:37:27	22.0N 122.0E	23.0N 121.7E	7.0
	1	Dec 20	19:34:00		23.0N 121.7E	
	2	Oct 15	15:45:40		23.3N 122.0E	
1920	m	Feb 02	11:22:18	4.0S 152.5E	7.0S 150.0E	7.7
	1	Feb 01	13:24:40		7.0S 150.0E	
	2	Jan 14	14:38:20		7.2S 150.0E	
1920	m	Mar 20	18:31:25	35.0S 110.0W	35.8S 109.4W	7.0
	1	Feb 08	05:24:30		35.0S 111.0E	
1922	m	Nov 11	04:32:36	28.5S 70.0W	29.0S 71.0W	8.4
	1	Nov 07	23:00:09	28.0S 72.0W	27.5S 72.8W	7.0
1923	m	Feb 03	16:01:41	54.0N 161.0E	54.0N 161.0E	8.4
	1	Feb 02	05:07:38	53.5N 162.0E	50.5N 164.0E	7.2
	2	Feb 02	01:06:15		50.5N 164.0E	
1923	m	Apr 19	03:09:08	2.5N 117.5E	2.5N 118.5E	7.0
	1	Apr 17	16:55:50		2.5N 118.5E	
1923	m	Nov 05	21:27:53	29.3N 130.0E	29.0N 130.0E	7.2
	1	Nov 03	16:19:12		29.0N 130.0E	
1924	m	Mar 04	10:07:42	9.8N 84.0W	9.5N 84.0W	7.0
	1	Mar 04	02:06:12		9.5N 84.0W	
1924	m	Aug 14	18:02:37	36.0N 142.0E	36.0N 142.0E	7.0
	1	Aug 14	17:53:36		36.0N 142.0E	
1925	m	Apr 16	19:52:35	22.0N 121.0E	22.0N 120.5E	7.1
	1	Mar 01	12:25:26		22.0N 120.5E	
1926	m	Sep 02	01:21:52	33.5S 59.0E	34.0S 57.0E	7.0
	1	Jul 22	22:54:10		34.0S 57.0E	
	2	Jul 08	07:16:10		34.0S 57.0E	
1926	m	Sep 10	10:34:29	9.0S 111.0E	9.0S 111.0E	7.0
	1	Jul 07	11:41:24		8.7S 111.0E	
	2	Jul 07	02:38:57		8.7S 111.0E	
	3	Jul 03	03:46:54		8.7S 111.0E	
1926	m	Oct 13	19:08:07	52.0N 176.0W	51.2N 176.0W	7.9
	1	Aug 09	03:39:22		52.0N 176.0W	
1926	m	Oct 26	03:44:41	3.5S 138.5E	2.7S 138.8E	7.9
	1	Oct 26	01:59:00		2.7S 138.8E	
1927	m	Jan 24	01:05:43	16.5S 167.5E	18.5S 168.5E	7.1
	1	Jan 05	13:24:00		18.5S 168.5E	
1927	m	Feb 16	01:35:20	47.0N 153.5E	46.0N 154.0E	7.0
	1	Nov 23	00:20:10		46.0N 154.0E	
1927	m	Mar 07	09:27:36	35.8N 134.8E	35.7N 134.8E	7.9
	1	Jan 15	14:21:16		36.2N 134.5E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1927	m	May 22	22:32:42	36.8N 102.0E	36.8N 102.8E	8.3
	1	May 22	21:43:00		36.8N 102.8E	
1927	m	Aug 05	21:12:55	37.5N 142.5E	38.5N 142.5E	7.1
	1	Jun 09	03:23:42		39.3N 142.4E	
1927	m	Aug 10	11:36:15	1.0S 131.0E	0.5S 130.0E	7.1
	1	Jun 11	02:49:25		1.0S 130.5E	
	2	Jun 11	02:32:06		1.0S 130.5E	
1927	m	Dec 28	18:20:23	55.0N 161.0E	54.0N 161.0E	7.3
	1	Dec 28	08:54:45		54.0N 161.0E	
1928	m	Mar 22	04:17:00	16.0N 96.0W	16.0N 96.0W	7.5
	1	Jan 25	21:06:40		16.0N 96.0W	
1928	m	May 27	09:50:26	40.0N 142.5E	40.0N 142.5E	7.0
	1	May 27	05:40:24		40.0N 142.5E	
1929	m	May 26	22:39:54	51.0N 131.0W	50.2N 130.7W	7.1
	1	Mar 01	08:52:36		50.2N 130.7W	
	2	Mar 01	07:31:00		50.2N 130.7W	
1929	m	Jul 07	21:23:12	52.0N 178.0W	50.5N 178.3W	7.3
	1	Jul 07	09:27:02		50.5N 178.3W	
	2	Jul 07	06:16:10		50.5N 178.3W	
	3	Jul 06	02:03:40		50.5N 178.3W	
	4	Jul 05	23:09:34		50.5N 178.3W	
	5	Jul 05	22:36:09		50.5N 178.3W	
	6	Jul 05	14:35:05		50.5N 178.3W	
	7	Jul 05	14:19:02	51.0N 178.0W	51.0N 179.5W	7.0
1930	m	May 06	22:34:23	38.0N 44.5E	37.0N 44.0E	7.2
	1	May 06	07:03:22		37.0N 44.0E	
	2	Apr 16	21:25:42		38.0N 43.8E	
1930	m	Nov 25	19:02:47	35.0N 139.0E	35.1N 139.0E	7.1
	1	Nov 21	12:55:50		35.1N 139.0E	
	2	Nov 21	12:17:30		35.1N 139.0E	
	3	Nov 21	10:16:56		35.1N 139.0E	
	4	Nov 21	09:49:58		35.1N 139.0E	
	5	Sep 28	09:51:51		34.5N 139.5E	
1930	m	Dec 03	18:51:44	18.0N 96.5E	18.2N 96.4E	7.3
	1	Dec 03	16:36:20		17.3N 96.5E	
1931	m	Feb 10	06:34:25	5.3S 102.5E	5.3S 102.5E	7.1
	1	Feb 10	05:58:42		5.3S 102.5E	
1931	m	Aug 27	15:27:17	29.8N 67.3E	29.8N 67.3E	7.4
	1	Aug 25	18:52:57		30.2N 67.7E	
	2	Aug 25	15:45:35		30.2N 67.7E	
	3	Aug 25	03:06:25		30.2N 67.7E	
	4	Aug 25	00:54:09		30.2N 67.7E	
	5	Aug 25	00:31:08		30.2N 67.7E	
	6	Aug 24	23:30:15		30.2N 67.7E	
	7	Aug 24	21:35:30		30.2N 67.7E	
1933	m	Jan 21	19:21:10	33.0S 57.5E	34.0S 57.5E	7.0
	1	Jan 17	18:47:45		34.0S 57.0E	
1934	m	Mar 05	11:46:15	40.5S 175.5E	40.4S 175.6E	7.5
1933	1	Dec 13	08:48:30		40.0S 175.5E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1935	m	Apr 20	22:01:54	24.3N 120.8E	24.0N 121.0E	7.1
	1	Apr 13	13:55:08		24.2N 121.8E	
	2	Apr 04	06:30:27		24.5N 121.5E	
		34 aftershocks of Feb 23 through Mar 03				
	3	Feb 23	20:52:29		24.0N 121.6E	
	4	Feb 22	09:05:08		24.2N 121.8E	
	5	Feb 22	08:55:28		24.2N 121.8E	
	6	Feb 17	16:13:30		24.0N 121.6E	
	7	Feb 09	19:19:42		24.2N 121.8E	
1935	m	Sep 04	01:37:41	22.3N 121.3E	22.3N 121.3E	7.2
	1	Aug 06	20:45:00		22.0N 122.0E	
	2	Jul 24	15:10:42		22.9N 121.5E	
	3	Jul 19	06:17:00		21.8N 121.5E	
	4	Jul 19	06:07:17		21.8N 121.5E	
1935	m	Sep 09	06:17:30	6.0N 141.0E	6.1N 140.7E	7.0
	1	Sep 09	06:11:47		6.1N 140.7E	
1935	m	Oct 18	00:11:56	40.5N 143.8E	40.2N 143.4E	7.2
	1	Oct 15	10:30:33		40.3N 144.0E	
	2	Oct 13	01:57:34		40.2N 144.0E	
	3	Oct 12	18:14:14		40.2N 144.0E	
	4	Oct 12	17:00:32		40.2N 144.0E	
	5	Oct 12	16:45:25	40.3N 143.3E	40.2N 144.0E	7.0
1936	m	Jul 05	18:55:13	6.3N 126.8E	5.9N 126.8E	7.3
	1	Apr 10	16:53:43		6.5N 126.0E	
1936	m	Aug 22	06:51:35	22.3N 120.8E	22.3N 121.3E	7.2
	1	May 28	12:28:25		22.3N 121.3E	
1937	m	Jan 23	10:55:51	4.5S 153.0E	5.1S 153.1E	7.0
1936	1	Dec 29	14:47:49		4.5S 153.9E	
	2	Dec 27	08:43:04		5.0S 153.0E	
1937	m	Feb 21	07:02:35	44.5N 149.5E	44.6N 149.4E	7.4
	1	Feb 17	09:15:24		44.6N 149.4E	
1938	m	May 30	14:29:50	20.5S 169.5E	19.9S 169.0E	7.0
	1	May 22	08:22:10		21.0S 169.5E	
	2	May 22	07:45:52		21.0S 169.5E	
1938	m	Sep 07	04:03:18	23.8N 121.5E	23.9N 121.7E	7.0
	1	Jun 16	22:50:30		24.6N 121.1E	
1938	m	Nov 05	08:43:21	36.8N 141.8E	37.1N 141.8E	7.7
	m	Nov 05	10:50:15	37.3N 141.8E	37.1N 141.8E	7.7
	m	Nov 06	08:53:53	37.3N 142.3E	37.1N 141.8E	7.6
	1	Sep 21	18:52:03		36.4N 141.1E	
	2	Sep 03	23:58:12		36.4N 141.1E	
1939	m	Jan 30	02:18:27	6.5S 155.5E	7.2S 155.3E	7.9
	1	Jan 22	18:40:48		6.2S 154.8E	
1939	m	Apr 05	16:42:40	19.5S 168.0E	19.9S 169.0E	7.1
	1	Apr 04	10:10:56		19.9S 169.0E	
	2	Apr 03	15:46:06		19.9S 169.0E	
1939	m	Apr 30	02:55:30	10.5S 158.5E	9.2S 159.5E	8.1
	1	Mar 07	17:16:22		9.2S 159.5E	
	2	Feb 04	06:07:26		9.2S 159.5E	
	3	Feb 04	05:19:24		9.2S 159.5E	
	4	Feb 03	05:26:20	10.5S 159.0E	9.2S 159.5E	7.1
1939	m	Jun 02	03:33:15	5.0N 127.0E	5.0N 126.8E	7.0
	1	Mar 05	02:13:45		5.0N 126.8E	
	2	Mar 04	20:03:58		5.5N 126.8E	
1940	m	Oct 04	07:54:42	22.0S 71.0W	20.5S 70.5W	7.3
	1	Oct 03	04:56:10		20.5S 70.5W	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1941	m	Jan 13	16:27:38	4.5S 152.5E	4.7S 152.7E	7.0
1940	1	Oct 21	20:15:22		5.1S 153.1E	
1941	m	Nov 08	23:37:22	0.5N 122.0E	0.1N 122.7E	7.3
	1	Sep 17	06:48:04		0.1N 122.7E	
	2	Aug 20	13:16:08		0.1N 122.7E	
1941	m	Nov 24	21:46:23	28.0S 177.5W	28.2S 177.7W	7.3
	1	Sep 16	21:39:03	28.8S 177.5W	29.1S 177.7W	7.0
1942	m	Aug 01	12:34:03	41.0S 175.8E	40.9S 175.8E	7.1
	1	Aug 01	04:47:48		40.9S 175.8E	
1942	m	Nov 15	17:12:00	37.0N 141.5E	36.4N 141.1E	7.0
	1	Nov 07	14:03:53		36.4N 140.6E	
	2	Nov 05	12:42:35		37.3N 141.3E	
	3	Sep 21	05:52:29		36.5N 141.6E	
	4	Sep 08	16:07:25		36.5N 141.6E	
	5	Sep 01	20:27:14		36.5N 141.6E	
1943	m	Mar 14	17:11:00	22.0S 169.5E	22.0S 170.3E	7.1
	1	Mar 11	09:34:05		21.5S 170.2E	7.3
1943	m	Mar 21	20:35:43	5.8S 152.3E	5.6S 150.5E	7.3
	1	Feb 16	16:50:53		5.6S 150.5E	
1943	m	Jun 09	03:06:22	1.0S 101.0E	1.2S 101.0E	7.6
	1	Jun 08	20:42:46	1.0S 101.0E	3.0S 101.0E	7.4
1943	m	Jun 13	05:11:49	42.8N 143.3E	40.9N 142.7E	7.4
	1	Apr 05	01:33:14		42.0N 142.8E	
1944	m	Mar 09	22:12:58	44.0N 84.0E	42.5N 82.5E	7.2
	1	Mar 09	22:03:43		42.5N 82.5E	
1944	m	Apr 27	14:38:09	0.5S 133.5E	1.1S 133.1E	7.4
	1	Apr 26	01:54:15	1.0S 134.0E	0.8S 133.5E	7.2
1944	m	May 25	12:58:05	2.5S 152.8E	2.5S 152.5E	7.5
	1	May 19	00:19:19	2.5S 152.8E	2.5S 152.3E	7.2
	2	May 18	04:43:17		2.0S 152.3E	
1944	m	Jun 28	07:58:54	15.0N 92.5W	13.8N 93.1W	7.0
	1	Jun 28	05:46:18		13.8N 93.1W	
	2	Jun 28	05:31:45		13.8N 93.1W	
	3	Jun 25	01:08:12		13.8N 93.1W	
1944	m	Sep 23	12:13:20	54.0N 160.0E	53.5N 160.5E	7.4
	1	Sep 19	13:06:00		53.5N 160.5E	
1945	m	Feb 01	12:13:40	22.0S 170.0E	21.4S 169.3E	7.25
	1	Feb 01	10:35:48	22.0S 170.0E	21.4S 169.3E	7.0
	2	Feb 01	10:24:44		21.4S 169.3E	
	3	Feb 01	10:19:53		21.4S 169.3E	
1945	m	Feb 10	04:57:56	41.3N 142.5E	41.1N 142.2E	7.3
	1	Feb 01	23:12:10		41.2N 142.5E	
1945	m	Mar 11	21:37:50	37.0N 142.0E	37.3N 142.0E	7.2
	1	Feb 04	15:46:25		37.3N 142.0E	
	2	Jan 06	22:01:38		36.7N 142.5E	
	3	Jan 01	08:50:56		37.8N 141.4E	
1945	m	Dec 28	17:48:45	6.0S 150.0E	6.1S 150.5E	7.8
	1	Dec 27	04:41:05	6.0S 151.0E	6.1S 150.5E	7.0
	2	Dec 08	01:04:02	6.5S 151.0E	6.1S 150.5E	7.1
	3	Nov 11	09:22:12		6.2S 151.5E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1946	m	Sep 12	15:20:20	23.5N 96.0E	23.9N 96.2E	7.75
	1	Sep 12	15:17:15	23.5N 96.0E	23.9N 96.2E	7.5
1946	m	Sep 29	03:01:55	4.5S 153.5E	5.1S 153.1E	7.75
	1	Jul 24	11:00:07		5.1S 153.1E	
	2	Jul 01	22:35:34		5.1S 153.1E	
1946	m	Dec 21	10:18:49	44.0N 149.0E	44.1N 148.2E	7.2
	1	Dec 21	03:39:15		43.5N 148.4E	
1947	m	Mar 02	19:09:26	5.0S 144.5E	5.0S 144.5E	7.0
1946	1	Dec 05	06:44:33		5.5S 144.5E	
1947	m	Mar 25	20:32:14	38.8S 178.5	38.4S 176.5E	7.0
	1	Feb 17	09:48:54		38.4S 176.5E	
1947	m	Oct 06	19:55:37	37.0N 22.0E	36.9N 22.0E	7.0
	1	Jul 21	09:36:21		36.7N 22.5E	
1947	m	Oct 16	02:09:47	64.5N 147.5W	64.2N 148.3W	7.0
	1	Oct 15	19:34:37		64.2N 148.3W	
	2	Oct 15	04:09:28		64.2N 148.3W	
	3	Oct 07	02:57:29		64.2N 148.3W	
	4	Oct 07	01:53:21		64.2N 148.3W	
1948	m	Jan 06	17:25:58	17.0N 98.0W	16.0N 98.4W	7.0
	m	Jan 06	17:23:26	17.0N 98.0W	16.0N 98.4W	7.0
1947	1	Dec 24	17:36:39		16.9N 98.7W	
	2	Dec 24	16:36:53		16.9N 98.7W	
1948	m	Jan 28	03:47:21	1.5N 126.5E	1.1N 126.4E	7.2
	m	Mar 13	20:02:35	1.5N 126.5E	1.5N 126.5E	7.1
1947	1	Nov 25	12:24:58		1.5N 126.0E	
1948	m	Aug 25	06:09:24	24.5S 65.0W	24.4S 64.6W	7.0
	1	Aug 16	10:45:11		24.4S 65.6W	
1948	m	Sep 02	23:34:50	10.0N 125.5E	10.3N 126.0E	7.0
	1	Jul 18	20:06:30		9.7N 125.7E	
1948	m	Sep 08	15:09:11	21.0S 174.0W	21.0S 174.0W	7.9
	1	Aug 28	12:24:26		20.0S 174.0W	
1949	m	Feb 23	16:08:08	41.0N 83.5E	42.2N 84.1E	7.3
	1	Feb 21	16:38:32		42.2N 84.5E	
1949	m	Jul 10	03:53:36	39.0N 70.5E	39.2N 70.7E	7.6
	1	Jul 08	08:02:15		39.2N 70.7E	
	2	Jul 08	07:50:37		39.2N 70.7E	
1949	m	Oct 19	21:00:19	5.5S 154.0E	6.5S 153.2E	7.25
	1	Sep 26	22:32:00		6.7S 153.0E	
1949	m	Dec 29	03:03:54	18.0N 121.0E	17.0N 121.5E	7.2
	1	Nov 22	06:26:59		17.0N 122.5E	
1950	m	Jan 30	00:56:32		53.5S 71.5W	7.0
	1	Dec 17	15:07:55		54.0S 71.0W	
	2	Dec 17	06:53:30		54.0S 71.0W	
1950	m	Oct 23	16:13:20		14.5N 91.5W	7.1
	1	Oct 23	15:03:40		14.5N 92.0W	
	2	Oct 23	14:55:36		14.5N 92.0W	
1950	m	Nov 08	02:18:12		10.0S 159.5E	7 1/4
	1	Oct 15	15:59:52		9.7S 159.6E	
1950	m	Dec 02	19:51:49		18.3S 167.5E	8.1
	m	Dec 02	19:55:27		18.3S 167.5E	7.25

Year		Date	G.M.T.	Location 1	Location 2	Mag
	1	Nov 08	01:59:06		18.0S 168.0E	
1951	m	Nov 06	16:40:05		47.8N 154.3E	7.2
	1	Nov 06	14:57:14		47.5N 153.9E	
1951	m	Nov 18	09:35:47		30.5N 91.0E	7.9
	1	Nov 18	09:26:32		30.5N 91.5E	
	2	Nov 17	04:46:00		31.0N 90.5E	
1952	m	Mar 04	01:22:43		42.5N 143.0E	8.6
	m	Mar 09	17:03:47		42.5N 143.0E	7.1
	1	Feb 14	19:20:36		42.3N 143.1E	
1952	m	Jul 21	11:52:14		35.0N 119.0W	7.7
	1	Jul 21	09:	Richter, p. 67		
1952	m	Aug 17	16:02:07		30.5N 91.5E	7.5
	1	Jun 02	10:33:34		30.5N 91.5E	
	2	Jun 02	10:08:23		30.5N 91.5E	
1952	m	Dec 24	18:39:38		5.5S 152.0E	7.0
	1	Dec 24	08:33:30		5.1S 151.3E	
1953	m	Jan 05	10:06:30		49.0N 156.0E	7.0
	1	Jan 04	06:43:15		48.0N 156.0E	
1952	2	Dec 31	21:43:49		49.5N 156.2E	
	3	Dec 24	15:49:30		49.5N 155.5E	
	4	Dec 16	06:16:00		48.6N 155.4E	
	5	Dec 11	13:11:47		48.6N 155.4E	
	6	Dec 11	08:58:18		48.6N 155.4E	
	7	Dec 04	10:49:32		49.3N 156.8E	
	8	Nov 21	03:19:21		49.5N 156.2E	
	9	Nov 12	00:40:56		49.5N 156.2E	
	10	Nov 09	05:32:13		49.5N 156.2E	
	11	Nov 09	00:22:19		49.5N 156.2E	
	12	Nov 08	19:33:21		49.2N 156.1E	
	13	Nov 07	15:40:42		49.5N 155.5E	
1953	m	Feb 26	11:42:28		11.0S 164.5E	7.0
	1	Jan 20	10:11:23		11.5S 165.0E	
1953	m	Aug 12	09:23:53		38.3N 20.8E	7.1
	1	Aug 12	06:08:01		38.3N 20.8E	
	2	Aug 11	13:11:04		38.3N 20.8E	
	3	Aug 11	12:43:22		38.3N 20.8E	
	4	Aug 11	03:32:22		38.3N 20.8E	
	5	Aug 09	07:41:07		38.3N 20.8E	
	6	Jun 21	08:11:26		37.5N 20.8E	
	7	Jun 01	20:17:33		37.5N 21.0E	
1953	m	Nov 17	13:29:53		13.8N 92.0W	7.1
	1	Sep 05	15:24:46		14.5N 92.3W	
	2	Aug 24	13:21:02		14.1N 91.4W	
1954	m	Jan 13	00:13:10		49.0S 165.8E	7.0
	1	Jan 12	22:26:30		49.5S 165.8E	
	2	Jan 12	14:20:26		48.5S 166.0E	
	3	Jan 12	14:16:22		49.0S 165.0E	
1954	m	Feb 19	19:07:48		30.0S 177.8W	7.0
	1	Jan 15	03:24:16		30.1S 177.8W	
1954	m	Mar 31	18:25:47		12.5N 58.0E	7.25
	1	Mar 16	14:24:16		11.5N 58.0E	
1954	m	Apr 29	11:34:31		29.2N 112.8W	7.0
	1	Apr 29	10:49:27		29.5N 112.5W	
1954	m	Apr 30	13:02:37		39.3N 22.2E	7.0
	1	Apr 25	20:03:46		39.5N 22.2E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1954	m	Jul 03	22:31:26		6.5S 105.3E	7.0
	1	Jun 18	17:54:41		6.3S 105.2E	
1954	m	Dec 16	11:07:10		39.3N 118.0W	7.1
	1	Dec 16	06:30		Richter, p. 73	
1955	m	Mar 18	00:06:43		54.0N 161.0E	7.4
	1	Feb 05	14:08:15		53.5N 161.5E	
	2	Jan 05	15:28:20		54.5N 161.0E	
1955	m	Mar 31	18:17:03		8.0N 124.0E	7.6
	1	Mar 11	23:32:46		8.0N 124.0E	
1955	m	Oct 10	08:57:47		5.0S 153.0E	7.3
	1	Oct 09	17:40:09		5.0S 153.0E	
	2	Aug 04	10:53:01		5.0S 152.5E	
1956	m	Jan 08	20:54:16		19.0S 70.0W	7.1
1955	1	Dec 12	02:32:58		19.5S 70.0W	
1956	m	Jun 09	23:13:52		35.0N 67.5E	7.6
	1	Jun 08	04:07:26		35.0N 67.5E	
1956	m	Jul 09	03:11:40		37.0N 26.0E	7.8
	1	Jul 08	13:05:22		36.9N 26.0E	
1957	m	Jan 02	03:48:44		53.0N 168.0W	7.0
	m	Mar 09	20:39:15		52.5N 169.5W	7.1
	m	Mar 11	09:58:42		53.0N 169.3W	7.0
	m	Mar 22	14:21:06		54.0N 166.0W	7.0
	1	Jan 02	03:41:08		52.5N 168.5W	
	2	Jan 02	03:30:34		52.5N 168.5W	
	3	Jan 02	03:12:52		53.0N 168.0W	
	4	Jan 02	02:17:15		52.5N 168.0W	
	5	Jan 02	00:39:22		53.0N 168.5W	
1956	6	Dec 09	05:19:06		53.0N 169.0W	
	7	Dec 03	07:20:06		52.6N 168.6W	
	8	Dec 03	07:12:44		53.0N 169.0W	
1957	m	Apr 14	19:18:01		15.5S 173.0W	7.5
	1	Apr 11	06:44:29		15.3S 172.3W	
	2	Apr 10	13:43:43		15.0S 173.0W	
	3	Jan 30	09:46:05		15.0S 173.0W	
	4	Jan 28	08:16:19		15.5S 173.0W	
1957	m	Apr 25	02:25:36		36.5N 28.5E	7.1
	1	Apr 24	19:10:05		36.0N 28.5E	
1957	m	Jun 22	23:50:23		1.5S 137.0E	7.3
	1	Apr 07	10:14:08		1.0S 137.5E	
1957	m	Jul 28	08:40:14		17.0N 99.0W	7.9
	1	Jun 09	01:16:14		16.6N 99.5W	
	2	May 30	17:24:12		16.4N 98.7W	
1957	m	Sep 24	08:21:05		5.5N 127.5E	7.6
	1	Jul 27	15:37:30		5.5N 127.5E	
	2	Jun 27	04:03:00		5.0N 127.5E	
1958	m	Apr 07	15:30:38		66.5N 157.0W	7.3
	1	Mar 31	05:50:	AGU Trans. v. 39, no. 5, p. 941		
1958	m	May 31	19:32:30		15.3S 168.5E	7.2
	1	Apr 26	09:25:54		15.0S 168.0E	
	2	Apr 14	19:21:54		14.5S 168.0E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
	3	Mar 03	04:06:16		14.5S 168.5E	
1958	m	Jul 10	06:15:51		58.6N 137.1W	7.9
	1	Jul 09	15:15		BSSA, v. 50, p. 217	
1958	m	Nov 06	22:58:06		44.5N 148.5E	8.7
	m	Nov 12	20:23:26		44.5N 148.5E	7.3
	1	Oct 27	18:16:53		44.5N 147.5E	
	2	Sep 05	10:59:06		45.0N 148.0E	
1959	m	May 04	07:15:42		52.5N 159.5E	8.25
	1	Mar 04	00:52:49		51.8N 159.3E	
	2	Feb 23	10:31:14		52.5N 159.0E	
1959	m	Aug 17	21:24:40		7.5S 156.0E	7.25
	1	Jun 29	07:16:07		7.3S 155.2E	
	2	Jun 05	05:58:40		7.0S 155.5E	
	3	May 30	14:18:30		7.0S 156.5E	
1959	m	Aug 18	06:37:15		44.8N 111.1W	7.1
	1	Aug 09			BSSA, v. 52, p. 156	
	2	Jun 14	11:23:00		45.0N 110.5W	
1959	m	Aug 24	21:30:46		10.5S 161.0E	7.0
	1	Aug 24	16:40:04		10.5S 161.5E	
	2	Aug 24	15:41:40		10.5S 161.3E	
	3	Aug 20	01:59:06		10.5S 161.0E	
	4	Aug 09	20:29:28		10.2S 161.0E	
1959	m	Sep 14	14:09:39		28.7S 177.7W	7.7
	1	Aug 14	03:47:30		29.0S 178.0W	
1959	m	Dec 14	23:21:56		59.8S 26.5W	7.0
	1	Oct 30	11:27:35		59.5S 27.0W	
1960	m	Mar 20	17:07:30		40.0N 143.5E	7.0
	1	Mar 20	13:44:25		39.8N 143.3E	
	2	Mar 20	13:36:49		39.8N 143.3E	
1960	m	May 22	19:11:17		39.5S 74.5W	8.3
	1	May 22	10:32:43		37.5S 73.0W	7.2
	2	May 21	10:02:50		37.5S 73.5W	7.25
1960	m	Jul 03	20:20:46		50.5N 177.0W	7.0
	1	Apr 18	11:15:54		51.0N 177.5W	
1960	m	Nov 01	08:45:59		38.5S 75.1W	7.2
	1	Oct 06	16:16:35		38.5S 74.9W	
	2	Sep 11	07:55:55		39.0S 74.8W	
	3	Aug 26	00:44:06		38.0S 74.3W	
	4	Aug 07	16:17:30		38.3S 74.5W	
1960	m	Nov 13	09:20:32		51.4N 168.8W	7.0
	1	Sep 22	22:47:01		51.7N 168.8W	
1960	m	Nov 24	06:52:41		24.4S 176.1W	7.0
	1	Nov 23	20:11:22		24.2S 176.2W	
	2	Nov 23	17:56:34		24.3S 176.3W	
	3	Nov 23	17:28:55		24.2S 176.4W	
1961	m	Jan 22	03:24:05		12.0S 166.2E	7.0
	1	Jan 18	09:65:44		12.3S 166.2E	
	2	Jan 02	10:11:58		12.5S 166.3E	
1960	3	Dec 19	22:46:24		12.2S 166.0E	
	4	Dec 19	22:13:59		12.7S 166.8E	
1961	m	Feb 26	18:10:49		31.6N 131.9E	7.2
	1	Jan 06	12:52:15		31.8N 131.6E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1961	m	Jul 23	21:51:08		18.5S 168.3E	7.2
	1	Jul 23	15:30:17		18.5S 168.3E	
	2	Jul 23	15:28:52		18.5S 168.2E	
	3	Jul 23	14:16:34		18.5S 168.1E	
	4	Jul 23	14:03:40		18.6S 168.2E	
	5	Jul 19	19:31:55		18.1S 167.7E	
	6	Jun 22	03:22:56		18.0S 168.9E	
	7	Jun 03	03:40:20		18.0S 167.9E	
	8	May 31	05:12:13		18.6S 168.7E	
	9	May 26	06:06:54		18.9S 168.9E	
	10	May 07	19:22:48		17.7S 168.0E	
1961	m	Aug 11	15:51:35		43.0N 145.0E	7.0
	1	Aug 10	12:15:46		42.9N 144.9E	
	2	Aug 10	12:05:24		42.6N 145.2E	
	3	Aug 04	07:17:47		42.5N 143.3E	
	4	Jun 12	01:37:10		42.7N 145.3E	
1962	m	Feb 14	06:36:05		37.8S 72.5W	7.25
1961	1	Dec 11	15:06:50		37.5S 72.8W	
1962	m	Apr 23	05:58:05		42.9N 143.4E	7.2
	1	Apr 01	05:01:56		41.9N 143.4E	
1962	m	May 06	19:00:14		60.2S 33.5W	7.0
	1	May 03	03:34:49		60.0S 32.9W	
	2	May 03	17:00:33		60.5S 33.2W	
1962	m	May 11	14:11:54		17.0N 99.6W	7.0
	m	May 19	14:58:15		17.2N 99.5W	7.2
	1	Mar 28	10:53:59		17.6N 99.0W	
	2	Mar 27	21:19:31		17.0N 99.8W	
1962	m	May 15	05:23:56		7.3S 128.3E	7.2
	1	Apr 23	14:51:26		6.9S 128.4E	
	2	Apr 17	15:15:10		7.0S 129.1E	
1962	m	Jul 30	17:16:44		3.4S 143.7E	7.0
	1	May 07	04:56:38		4.1S 143.7E	
1962	m	Aug 03	08:56:12		23.2S 67.5W	7.1
	1	Jul 23	07:19:35		22.9S 67.8W	
1962	m	Sep 18	00:29:05		7.5N 82.3W	7.0
	1	Jul 26	08:14:45		7.4N 82.7W	
	2	Jun 21	05:13:17		8.1N 82.8W	
1963	m	Feb 13	08:50:02		24.5N 121.8E	7.25
	1	Jan 15	13:08:53		25.0N 122.0E	
1963	m	Mar 16	08:44:48		46.5N 154.7E	7.0
	1	Jan 18	12:33:51		46.5N 154.3E	
	2	Jan 04	08:47:26		46.5N 153.9E	
1963	m	Mar 26	09:48:20		29.7S 177.8W	7.0
	1	Feb 18	14:51:40		30.4S 177.2W	
	2	Jan 11	17:05:42		29.5S 178.6W	
	3	Jan 09	02:02:39		29.0S 177.3W	
1963	m	Apr 16	01:29:19		0.8S 128.0E	7.0
	1	Apr 15	20:19:30		0.8S 128.0E	
1963	m	Sep 15	00:46:54		10.3S 165.6E	7.25
	m	Sep 17	19:20:08		10.1S 165.3E	7.25
	1	Sep 06	05:29:54		10.9S 164.8E	
1963	m	Sep 24	16:30:16		10.6S 78.0W	7.0
	1	Sep 20	04:00:36		11.3S 77.6W	
	2	Sep 17	05:54:34		10.6S 78.2W	
	3	Aug 15	14:36:01		10.0S 78.6W	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1963	m	Oct 13	05:17:58		44.8N 149.5E	8.25
	m	Oct 20	00:53:07		44.7N 150.7E	7.0
	1	Oct 12	11:26:58		44.8N 149.0E	7.0
	2	Oct 05	23:50:46		45.3N 150.0E	
	3	Sep 27	18:04:38		45.6N 149.5E	
1964	m	Mar 28	03:36:13		61.1N 147.6W	8.5
	1	Mar 22	06:22:15		61.3N 147.8W	
1964	m	Jul 24	08:12:40		47.2N 153.8E	7.0
	1	Jul 24	07:13:01		46.9N 153.8E	
	2	Jul 24	06:50:52		46.9N 153.9E	
	3	Jun 13	20:42:18		46.4N 153.8E	
	4	Jun 13	08:28:39		46.4N 153.8E	
1964	m	Nov 17	08:15:39		5.7S 150.7E	7.6
	1	Oct 14	12:06:38		5.7S 150.5E	
	2	Sep 28	20:06:46		5.2S 150.5E	
	3	Sep 27	22:01:40		5.5S 151.6E	
	4	Aug 16	11:08:00		5.9S 151.4E	
1965	m	Jan 24	00:11:12		2.4S 126.0E	7.6
USCGS preceded by week of activity						
1965	m	Feb 04	05:01:22		51.3N 178.6E	7 3/4
	m	Mar 30	02:27:07		50.6N 177.9E	7.3
	1	Feb 04	04:53:53		51.1N 178.4E	
	2	Feb 01	06:49:01		51.4N 178.6E	
	3	Feb 01	02:20:52		51.3N 178.5E	
	4	Jan 31	23:50:28		51.1N 178.2E	
	5	Jan 31	23:36:13		51.2N 178.6E	
	6	Nov 25	07:35:41		51.4N 178.8E	
	7	Nov 16	20:35:06		51.6N 178.5E	
	8	Nov 07	11:20:46		50.7N 179.1E	
1965	m	Mar 28	16:33:16		32.4S 71.3W	
	1	Mar 23	04:05:44		32.6S 72.1W	
	2	Mar 22	22:56:26		31.9S 71.5W	
1965	m	May 20	00:40:11		14.7S 167.4E	7.0
	m	Aug 11	03:40:55		15.5S 166.9E	7
	m	Aug 11	22:31:46		15.8S 167.2E	7.3
	m	Aug 13	12:40:37		15.9S 167.6E	7
	1	May 06	09:52:23		15.5S 167.6E	
	2	Apr 22	14:52:05		14.6S 167.3E	
	3	Apr 22	01:05:50		14.3S 167.3E	
	4	Apr 20	02:03:33		14.6S 166.9E	
	5	Apr 17	09:46:42		14.6S 167.5E	
	6	Apr 14	22:53:00		15.4S 167.5E	
	7	Apr 02	16:43:06		14.8S 167.3E	
	8	Mar 25	07:17:21		14.3S 167.4E	
1965	m	Aug 23	19:46:02		16.3N 95.8W	7.25
	1	Jul 08	07:20:41		15.9N 96.2W	
	2	Jun 02	12:30:44		16.0N 96.7W	
1966	m	Mar 12	16:31:22		24.1N 122.6E	7 1/2 - 7 3/4
	1	Mar 02	14:09:02		24.5N 122.6E	
	2	Feb 03	17:57:39		24.2N 121.9E	
	3	Feb 03	17:21:08		24.2N 121.8E	
	4	Feb 03	17:11:19		24.2N 122.1E	
1966	m	Mar 20	01:42:50		0.6N 30.2E	7

35 f.s. 1.8 days before BSSA v. 60, p. 1245

Year		Date	G.M.T.	Location 1	Location 2	Mag
1966	m	Mar 22	08:19:34		37.6N 115.2E	6 3/4-
	1	Mar 22	08:11:35		37.6N 115.2E	
	2	Mar 22	05:57:46		37.6N 115.2E	
	3	Mar 19	16:59:42		37.4N 114.9E	
	4	Mar 11	06:22:44		37.7N 114.8E	
	5	Mar 11	06:20:49		37.1N 115.1E	
	6	Mar 07	21:29:17		37.3N 114.9E	
with 6 aftershocks to 03/07 on 03/07 and 03/08						
1966	m	Jun 15	00:59:46		10.4S 160.8E	7 1/2
	m	Jun 15	01:32:55		10.2S 161.1E	7 1/4
	1	May 05	05:42:49		10.1S 161.1E	
	2	May 03	01:15:58		10.5S 161.6E	
	3	May 01	01:09:57		10.2S 161.2E	
	4	Apr 18	09:08:13		10.6S 161.6E	
	5	Apr 08	06:23:39		10.2S 161.3E	
	6	Mar 30	01:26:36		10.3S 161.6E	
1966	m	Jul 04	18:33:36		51.7N 179.8E	7
	1	Jul 04	01:36:29		51.7N 179.9E	
	2	Jul 03	18:55:39		51.9N 179.3W	
	3	Jul 03	17:45:35		52.0N 179.7W	
	4	Jul 03	15:21:31		51.7N 179.4E	
	5	Jul 03	01:33:01		51.6N 179.9E	
	6	Jul 02	07:04:37		51.6N 179.9E	
	7	Jul 01	16:38:19		51.8N 179.8E	
	8	Jul 01	04:53:26		51.5N 179.1E	
	9	Jun 30	17:07:10		51.9N 179.7W	
	10	Jun 27	04:11:03		52.1N 179.6W	
	11	May 07	09:04:55		51.1N 179.8E	
	12	Apr 20	07:57:10		50.9N 179.5E	
	13	Apr 18	15:07:34		51.2N 179.7W	
	14	Apr 11	23:45:16		51.1N 179.9E	
1966	m	Aug 07	02:13:05		50.6N 171.3W	7.0
	1	May 25	16:58:56		51.4N 170.9W	
1966	m	Aug 19	12:22:10		39.2N 41.6E	7
	1	Jul 12	00:04:09		39.2N 41.3E	
1966	m	Sep 08	21:15:53		2.4N 128.4E	7.0
	1	Aug 28	19:00:55		2.5N 128.0E	
	2	Aug 28	18:47:53		2.5N 128.1E	
	3	Aug 28	02:11:21		2.5N 129.0E	
1966	m	Oct 17	21:41:56		10.7S 78.7W	7.5
	1	Aug 23	01:37:15		10.8S 78.1W	
1966	m	Dec 28	08:18:07		25.5S 70.7W	7 3/4
	1	Nov 25	02:18:75		25.6S 70.7W	
1966	m	Dec 31	18:23:04		11.8S 166.5E	7.5
	m	Dec 31	22:15:14		11.3S 164.8E	7.3
	1	Dec 26	17:16:36		11.3S 164.3E	
	2	Dec 15	13:42:06		11.2S 165.7E	
	3	Dec 14	23:21:08		11.2S 166.6E	
	4	Dec 11	17:34:57		11.8S 166.4E	
	5	Nov 29	03:17:22		11.5S 166.3E	
	6	Nov 13	03:03:33		11.1S 166.3E	
	7	Nov 05	16:03:49		11.0S 166.9E	
	8	Oct 19	11:22:15		12.5S 166.9E	
	9	Oct 18	20:49:52		11.1S 166.7E	

Year	Date	G.M.T.	Location 1	Location 2	Mag
	10	Oct 18	20:06:51	11.0S 167.0E	
	11	Oct 18	02:37:47	11.1S 166.7E	
	12	Oct 17	13:48:54	11.1S 166.8E	
	13	Oct 17	12:09:23	11.1S 167.1E	
	14	Oct 17	10:15:39	11.0S 166.8E	
	15	Oct 17	09:13:25	11.2S 167.0E	
	16	Oct 17	08:51:45	11.2S 167.1E	
	17	Oct 17	03:57:48	11.1S 166.9E	
	m	Mar 04	17:58:06	39.2N 24.6E	6 3/4-7
	1	Mar 04	17:55:11	39.1N 24.6E	
	2	Mar 04	17:46:05	39.1N 24.6E	
1967	m	Mar 13	16:06:54	40.1S 74.5W	7.3
	1	Feb 10	03:17:43	39.2S 74.6W	
1967	m	Jul 22	16:56:53	40.7N 30.8E	7.25
	1	May 25	03:35:29	40.3N 30.9E	
1967	m	Sep 03	21:07:31	10.6S 79.8W	6 3/4-7
	1	Jul 24	04:45:23	11.3S 79.5W	
	2	Jun 09	05:44:37	11.3S 79.4W	
1967	m	Dec 21	02:25:22	21.8S 70.0W	7.0
	1	Oct 18	06:38:55	22.0S 69.2W	
1967	m	Dec 24	20:03:11	17.4N 61.1W	6 3/4-7
	1	Oct 28	11:28:34	17.9N 61.2W	
	2	Oct 26	13:44:45	17.6N 61.0W	
	3	Oct 26	12:21:33	17.7N 60.9W	
	4	Oct 17	06:05:37	17.0N 61.0W	
	5	Oct 17	01:51:27	16.9N 61.0W	
	6	Oct 14	03:31:05	17.3N 60.8W	
1967	m	Dec 25	01:23:24	5.3S 153.7E	6 3/4-7
1968	m	Feb 12	05:44:48	5.5S 153.2E	7
1967	1	Dec 19	02:58:27	6.0S 153.5E	
	2	Dec 18	12:27:38	6.1S 153.5E	
	3	Dec 12	04:24:42	5.2S 152.8E	
	4	Dec 11	16:18:20	5.2S 152.9E	
	5	Dec 09	17:00:24	4.4S 153.6E	
	6	Nov 29	04:07:37	4.9S 153.2E	
	7	Oct 31	13:36:03	6.0S 154.2E	
	8	Oct 31	13:04:01	6.0S 154.0E	
	9	Oct 24	09:50:15	5.8S 153.0E	
	10	Oct 16	13:29:16	5.1S 154.0E	
	11	Oct 14	01:15:22	5.9S 153.5E	
	12	Oct 12	01:22:12	5.6S 153.3E	
	13	Oct 11	13:14:21	5.6S 153.8E	
	14	Oct 10	05:15:13	5.6S 153.9E	
	15	Oct 9	13:27:57	5.7S 154.0E	
	16	Oct 8	18:08:18	5.6S 154.0E	
	17	Oct 8	13:13:46	5.8S 153.9E	
	18	Oct 7	20:46:46	5.0S 153.9E	
	19	Oct 7	19:52:07	5.6S 153.7E	
	20	Oct 7	19:11:02	6.0S 154.1E	
	21	Oct 5	12:35:39	5.2S 154.1E	
	22	Oct 5	04:07:42	5.8S 154.0E	
	23	Oct 4	23:14:33	5.8S 153.8E	
	24	Oct 4	17:21:21	5.7S 153.9E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1968	m	Feb 19	22:45:41		39.4N 25.0E	7 1/4 - 7 1/2
	1	Feb 19	22:45:03		39.3N 25.0E	
	2	Feb 06	21:32:04		39.7N 25.2E	
1968	m	May 16	00:48:55		40.8N 143.2E	7.9
	m	May 16	10:39:02		41.5N 142.7E	7.0
	1	May 01	19:16:39		40.9N 142.4E	
	2	May 01	19:12:53		40.8N 142.5E	
	3	Apr 26	00:29:53		40.8N 143.3E	
	4	Feb 20	17:30:35		41.2N 142.6E	
1968	m	May 20	20:05:49		30.7S 178.4W	7.0
	m	Jul 25	07:23:08		30.8S 178.4W	7 1/4
	1	May 20	07:13:03		30.9S 178.3W	
	2	Mar 26	21:24:59		30.3S 178.0W	
	3	Feb 21	19:27:30		30.2S 179.0W	
1968	m	Aug 01	20:19:22		16.5N 122.2E	7.3
	10	in 24 hr before BSSA, v. 59, p. 459				
1968	m	Aug 02	14:06:44		16.6N 97.7W	7.1
	1	Jul 21	08:59:16		16.5N 97.1W	
1969	m	Jan 06	15:39:01		10.5S 164.5E	7.1
1968	1	Dec 14	00:48:44		10.9S 164.7E	
1969	m	Jan 30	10:29:40		4.8N 127.4E	7.2
1968	1	Nov 25	18:36:53		5.0N 126.9E	
1969	m	Feb 28	02:40:32		36.0N 10.6W	8.0
	1	Feb 23	15:26:47		36.0N 10.5W	
1969	m	Mar 27	12:41:36		4.8N 127.5E	7.0
	1	Mar 01	07:52:17		3.9N 127.8E	
1969	m	Mar 31	07:15:54		27.7N 34.0E	7-7 1/4
	1	Mar 27	06:15:30		27.5N 33.9E	
	2	Mar 24	12:50:50		27.6N 33.8E	
	3	Mar 24	11:54:15		27.5N 33.8E	
1969	m	May 14	19:32:54		51.3N 179.9W	7.0
	1	Apr 08	02:44:01		51.3N 179.3W	
1969	m	Aug 05	02:13:10		1.3N 126.2E	7.2
	m	Aug 11	23:52:56		1.7N 126.5E	7.0
	1	Jul 24	05:03:27		1.6N 126.5E	
	2	Jul 08	18:43:57		1.4N 126.2E	
	3	Jul 07	03:37:54		1.8N 126.4E	
	4	Jul 05	00:37:33		1.2N 126.1E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1969	5	May 13	06:00:07		1.5N 126.3E	
	m	Aug 11	21:27:39		43.5N 147.4E	7.8
	1	Aug 11	21:27:36		43.5N 147.8E	
	2	Aug 11	21:27:26		43.6N 147.7E	
	3	Aug 11	21:26:38		43.4N 147.8E	
	4	Aug 11	21:21:47		43.7N 147.6E	
	5	Aug 11	21:07:58		44.0N 147.8E	
	6	Aug 11	21:02:15		44.0N 147.8E	
	7	Aug 11	20:58:53		44.0N 147.7E	
1969	8	Aug 10	21:09:26		43.6N 147.6E	
	9	Jul 13	05:41:25		43.3N 147.4E	
	m	Dec 25	21:32:27		15.8N 59.7W	7.2
1970	1	Dec 19	08:02:46		16.2N 60.4W	
	m	Jan 10	12:07:09		6.8N 126.7E	7.3
1970	1	Nov 07	09:13:12		6.6N 125.8E	
	m	Apr 29	14:01:33		14.5N 92.6W	7.3
	1	Apr 29	14:01:19		14.5N 92.6W	
	2	Apr 29	12:52:53		14.5N 92.8W	
	3	Apr 29	12:29:19		14.5N 92.7W	
	4	Apr 29	12:07:31		14.2N 92.8W	
	5	Apr 29	11:46:34		14.7N 92.6W	
	6	Apr 29	11:42:38		14.4N 92.6W	
	7	Apr 29	11:22:36		14.6N 92.7W	
	8	Apr 17	16:20:06		14.5N 93.1W	
	m	May 31	20:23:27		9.2S 78.8W	7.8
	1	May 31	20:23:07		BSSA, v. 61, p. 535	
	m	Jun 24	13:09:08		51.8N 131.0W	7.0
	1	Jun 24	07:30:31		51.8N 130.8W	
	m	Aug 11	10:22:20		14.1S 166.6E	7.0
1970	1	Aug 11	10:09:28		14.1S 166.6E	
	2	Aug 10	15:15:20		13.9S 166.8E	
	3	Aug 01	10:36:05		14.8S 166.7E	
	4	Jun 13	21:25:28		13.4S 166.8E	
	5	Jun 09	16:15:23		14.0S 166.9E	
	6	Jun 07	04:20:08		13.8S 167.2E	
	7	May 24	02:18:58		14.3S 167.1E	
	m	Oct 31	17:53:09		4.9S 145.5E	7.0
	1	Oct 31	06:19:41		4.9S 145.5E	
	2	Aug 24	10:07:47		5.5S 145.7E	
	3	Aug 22	12:58:21		5.4S 145.7E	
	m	Dec 02	15:54:20		11.0S 163.3E	7.0
	1	Dec 02	15:25:47		10.9S 163.5E	
	2	Dec 02	15:17:22		11.0S 163.6E	
	3	Dec 01	18:51:19		11.0S 163.4E	
1970	4	Dec 01	18:14:39		11.0S 163.4E	
	5	Dec 01	21:35:34		11.1S 163.3E	
	m	Jan 10	07:17:04		3.1S 139.7E	7.0
	1	Dec 31	02:30:24		2.8S 139.4E	
	m	Feb 04	15:33:29		0.7N 98.8E	7.1
	1	Dec 30	08:11:10		1.4N 99.1E	
	m	Feb 08	21:04:22		63.5S 61.2W	7.0
	1	Feb 08	20:37		63.3S 61.6W	
	m	Jan 10	07:17:04		3.1S 139.7E	8.1
	1	Dec 31	02:30:24		2.8S 139.4E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1971	m	May 02	06:08:27		51.4N 177.2W	7.1
	1	Apr 12	12:10:35		51.3N 177.1W	
	2	Mar 30	11:30:39		51.2N 177.5W	
	3	Mar 02	09:42:13		51.8N 176.8W	
	4	Feb 28	07:21:38		51.3N 177.0W	
	5	Feb 24	23:22:16		50.9N 177.4W	
	6	Feb 16	09:37:06		51.3N 177.6W	
	7	Feb 10	10:07:05		51.3N 177.0W	
	8	Feb 08	19:21:12		51.2N 176.8W	
	9	Feb 08	06:39:18		51.2N 176.5W	
	10	Feb 08	04:43:01		51.3N 177.0W	
	11	Feb 08	04:00:05		51.4N 176.7W	
	12	Feb 08	01:03:14		51.5N 176.7W	
	13	Feb 08	00:04:31		51.2N 176.7W	
	14	Feb 07	23:45:26		51.5N 176.8W	
	15	Feb 07	23:34:13		51.2N 176.7W	
	16	Feb 07	08:19:20		51.5N 176.2W	
	17	Feb 07	05:08:45		51.3N 176.9W	
	18	Feb 07	04:01:59		51.2N 177.0W	
	19	Feb 07	03:39:07		51.1N 177.1W	
	20	Feb 07	03:21:00		51.1N 177.0W	
	21	Feb 07	03:19:12		51.2N 177.0W	
	22	Feb 07	03:18:57		51.1N 177.0W	
	23	Feb 07	03:12:06		51.2N 176.9W	
	24	Feb 07	03:03:18		51.7N 177.3W	
	25	Feb 07	03:00:39		51.4N 177.1W	
	26	Feb 07	02:53:09		51.2N 176.9W	
	27	Feb 07	02:50:49		51.1N 177.0W	
	28	Feb 07	02:42:05		51.2N 177.1W	
	29	Feb 07	02:29:28		51.4N 176.7W	
1971	m	Jul 14	06:11:29		5.5S 153.9E	7.9
	m	Jul 19	00:14:45		5.7S 153.8E	7.1
	m	Jul 26	01:23:21		4.9S 153.2E	7.9
	1	Jul 10	05:23:52		6.0S 154.6E	
	2	Jul 09	11:50:34		6.0S 154.4E	
	3	Jul 09	09:49:43		5.9S 154.4E	
	4	May 22	05:01:18		5.3S 154.1E	
	5	May 08	08:30:06		5.5S 153.3E	
1971	m	Aug 02	07:24:57		41.4N 143.5E	7.0
	1	Jul 15	10:57:02		41.6N 142.6E	
1971	m	Oct 27	17:58:37		15.5S 167.2E	7.1
	1	Oct 10	21:24:59		15.5S 167.7E	
	2	Oct 05	16:35:22		14.6S 167.4E	
	3	Sep 29	05:34:47		15.3S 167.8E	
	4	Aug 30	21:12:52		14.8S 167.3E	
	5	Aug 28	09:45:03		14.7S 167.6E	
	6	Aug 14	03:36:07		14.8S 167.3E	
	7	Aug 14	00:15:18		14.9S 167.2E	
1972	m	Jan 23	21:17:52		13.2S 166.4E	7.1
	1	Jan 23	18:04:00		13.2S 166.4E	
	2	Jan 23	17:18:39		13.1S 166.3E	
	3	Jan 13	08:10:04		12.9S 166.7E	
1971	4	Dec 17	04:57:50		13.9S 166.2E	
	5	Nov 30	15:07:42		13.0S 167.2E	
	6	Nov 26	14:40:24		12.8S 166.4E	
	7	Nov 23	08:48:34		13.4S 166.6E	
	8	Nov 20	22:59:17		13.2S 166.5E	

Year	Date	G.M.T.	Location 1	Location 2	Mag
	9	Oct 30	22:35:18	13.5S 166.3E	
	10	Oct 29	23:09:32	13.6S 166.4E	
	11	Oct 29	20:07:13	13.5S 166.3E	
	12	Oct 29	20:04:50	13.6S 166.4E	
	13	Oct 28	18:00:01	13.4S 166.4E	
	14	Oct 28	14:20:30	13.3S 166.5E	
	15	Oct 26	22:32:22	13.0S 166.5E	
	16	Oct 26	22:23:51	12.8S 167.2E	
	17	Oct 26	20:39:48	13.0S 166.5E	
	18	Oct 25	03:46:36	13.1S 166.5E	
1972	m	Jan 25	02:06:23	22.5N 122.3E	7.5
	m	Jan 25	03:41:24	23.0N 122.2E	7.0
	1	Jan 04	03:16:54	22.6N 122.1E	
1972	m	Feb 29	09:23:00	33.3N 140.8E	7.2
	1	Jan 04	00:45:45	33.9N 141.3E	
1972	m	Apr 25	19:30:09	13.4N 120.3E	7.2
	1	Mar 28	08:29:21	13.5N 120.8E	
1972	m	Aug 17	23:44:06	6.0S 152.9E	7.1
	1	Aug 03	11:18:40	5.8S 152.0E	
	2	Aug 02	06:06:00	6.0S 152.7E	
	3	Jul 22	19:47:34	5.8S 153.0E	
	4	Jun 28	08:36:02	5.8S 153.3E	
	5	Jun 27	18:56:53	5.1S 152.7E	
1972	m	Sep 04	18:11:12	11.7S 166.2E	7.0
	1	Aug 12	13:42:32	10.9S 165.9E	
	2	Jul 07	05:07:30	11.1S 165.5E	
	3	Jun 17	12:44:27	11.0S 166.2E	
1972	m	Nov 02	19:55:22	20.0S 168.8E	7.0
	1	Oct 27	09:23:08	20.0S 169.0E	
	2	Sep 24	14:23:21	20.7S 168.4E	
	3	Sep 23	02:59:33	20.9S 168.5E	
	4	Sep 15	17:46:51	20.4S 168.7E	
	5	Sep 15	13:78:45	20.1S 168.8E	
	6	Sep 14	09:24:12	20.5S 168.9E	
	7	Sep 13	14:55:46	20.4S 168.8E	
	8	Sep 13	14:33:38	20.4S 168.8E	
	9	Sep 13	13:51:59	20.2S 168.8E	
	10	Sep 13	13:32:34	20.5S 168.8E	
	11	Sep 13	13:06:56	20.3S 168.8E	
	12	Sep 13	12:58:56	20.5S 168.8E	
	13	Sep 13	12:52:18	20.4S 168.7E	
	14	Sep 13	06:22:23	20.3S 168.7E	
	15	Sep 12	02:17:22	20.5S 168.9E	
	16	Sep 11	11:50:26	20.5S 168.8E	
	17	Sep 10	21:48:47	20.5S 168.8E	
	18	Sep 10	11:07:31	20.1S 168.8E	
	19	Sep 10	10:48:28	20.3S 168.8E	
	20	Sep 10	10:07:13	20.4S 169.0E	
	21	Sep 10	09:16:36	20.3S 168.6E	
	22	Sep 10	06:03:26	20.2S 168.8E	
	23	Sep 05	18:03:35	19.8S 169.0E	
	24	Sep 05	17:30:36	19.9S 169.0E	
	25	Aug 30	16:24:47	19.5S 169.0E	

Year		Date	G.M.T.	Location 1	Location 2	Mag
1972	m	Dec 02	00:19:47		6.5N 126.0E	7.4
	1	Oct 28	11:57:53		5.9N 126.6E	
1972	m	Dec 04	10:16:12		33.3N 140.7E	7.4
	1	Dec 04	05:19:32		33.2N 140.9E	
	2	Dec 02	20:32:59		33.2N 141.0E	
	3	Dec 02	03:57:54		33.2N 141.0E	
	4	Nov 12	06:00:45		33.4N 140.8E	
	5	Oct 28	17:54:16		33.3N 140.1E	
	6	Oct 23	21:25:17		33.2N 141.0E	
	7	Sep 30	10:41:13		33.1N 140.6E	
1973	m	Feb 28	06:37:49		50.5N 156.6E	7.2
	1	Feb 08	02:35:09		50.5N 156.4E	
	2	Feb 04	01:16:08		50.9N 157.2E	

Major ($M > 7.0$) shallow ($d < 100$ km) earthquakes preceded by foreshocks 1914 to 1973. The mainshocks (m) are listed in chronological order with the foreshocks for each (1, 2, etc.) listed after it. For 1914 to 1949, Location 1 refers to the location given by Duda (1965) and Location 2 refers to the location given by the International Seismological Summary.

TABLE II

Percentage of Mainshocks having foreshocks within

	90 days	40 days	5 days	1 day
1914-1949 (511 events)	14.7	11.3	6.6	3.9
1950-1973 (250 events)	49.8	42.2	20.0	14.8
1950-1973 (162 events) with 2 aftershocks	66.0	55.6	28.5	22.8

Table III

Maximum amplitudes of events preceding major earthquakes. Amplitudes recorded on short period instruments are given in millimeters (accuracy $\pm 10\%$). Parentheses around a number indicate that the wave trace was too faint to allow accurate readings.

OCCURENCE time	VERTICAL		NORTH		EAST	
	P	S	P	S	P	S
Earthquakes preceding the 01 August 1968 event in the Philippines						
01 Aug 20:13	18	(70)	15	(70)	-	-
20:08	3	18	3	15	5	17
20:00	8	38	6	33	9	-
18:58	5	30	4.5	28	6	-
18:23	(25)	(70)	24	-	-	-
17:16	2	13	2	10	3	(15)
16:38	3	18	3	15	4	35
11:17	5	30	10	55	-	-
Earthquakes preceding the 05 December 1968 event in Greece						
04 Dec 20:48	2.5	6	2	7	2.5	6
20:30	6	24	6	22	7.5	18
20:11	5	16	5	20	6	15
19:37	40	(75)	(24)	(100)	(40)	(86)
18:52	20	(48)	(14)	(60)	20	50
18:42	7	22	6	25	10	22
26 Nov 04:30	15	(120)	13	-	21	53
13 Nov 17:48	12	40	10	38	20	70
12 Nov 23:09	(2)	15	2	10	5	10
06:52	(3)	30	(4)	36	28	52
06:09	(52)	(100)	(30)	(84)	(30)	(76)
04:01	8	50	8	(30)	10	38
03:42	15	42	9	(40)	19	(60)
01:20	2	11	2	8	4	9
01:17	2	20	2	16	3	18
03 Nov 14:08	4	20	2	10	3	10
Earthquakes preceding the 28 March 1969 event in Turkey						
27 Mar 18:07	4	6			3	6
26 Mar 09:00	2	3			2	4
03:31	7	12			7	14
25 Mar 16:13	10	25			15	31
14:40	4	10			5	14
14:19	16	30			16	37
24 Mar 12:13	6	13			7	16
11:35	6	24			6	14
08:13	9	21			8	20
02:59	3	6			3	7
23 Mar 03:51	5	8			4	9
00:16	5	9			4	9

FIGURE CAPTIONS

Figure 1. Percentage of major earthquakes ($M \geq 7$) in each year from 1914 to 1973 that were preceded by recorded foreshocks. For 1950 to 1973, the hatched area is the percentage of earthquakes that had at least two aftershocks recorded by NOAA that were preceded by foreshocks.

Figure 2. Foreshock activity as a function of time before mainshocks. a - earthquakes between 1950 and 1973 and b - earthquakes between 1964 and 1973. The number of foreshocks are given separately for three different ranges of distance between the computed foreshock and mainshock epicenters. Inserts show the number of foreshocks averaged over ten day periods for a one-year period preceding the mainshocks.

Figure 3. Foreshock activity as a function of time before mainshocks for the three days before the mainshocks.

Figure 4. Number of foreshocks per day as a function of number of the foreshocks recorded for different earthquakes. Separate plots are shown for foreshock sequences consisting of only 1 event, 2 or 3, 3 or 4 or greater than 5 events. Note that the increase about five days before the mainshocks and the peak on the last day do not depend upon how many foreshocks preceded each event.

Figure 5. Foreshock activity as a function of time before mainshocks for mainshocks of different magnitudes (a, b, c) and for all mainshocks from all three data sets (d). Also shown are the exponential (solid line)

and power law (dotted line) equations that best fit these data. Note that the temporal variation does not depend upon the magnitude of the mainshock.

Figure 6. The magnitudes of the largest foreshocks as a function of the magnitude of their mainshocks. No pattern is obviously discernable.

Figure 7. The maximum amplitude of the P-wave vs. the maximum amplitude of the S-wave of both the vertical and north-south records of the 01 August 1968 earthquake in the Philippines recorded at Baguio (BAG)

Figure 8. The maximum amplitude of the P-wave vs. the maximum amplitude of the S-wave of the 05 Dec 1968 earthquake near Greece recorded at Athens University (ATU).

The closed circles are foreshocks that occurred within 4 days of the mainshock while the open circles are foreshocks occurring more than 20 days before it.

Figure 9. The maximum amplitude of the P-wave vs. the maximum amplitude of the S-wave of the 28 Mar 1969 earthquake in western Turkey recorded at Athens University (ATU).

Figure 10. A plot of equation (5) in terms of the non-dimensional variables $k\sigma (=k\sigma_0 x)$ and $(t+C)/T$. This shows the increase of stress on the increasing unbroken asperities with increasing time to the left.

Figure 11. A plot of equation (5) in terms of the non-dimensional variables $1/k\sigma_0 x$ and $(t+C)/T$.

Smooth curve shows the decrease of unslipped area with time.

The fraction of the area that has slipped is shown

at the right for three values of $k\sigma_0$. Dots connected by line segments show the total, integrated number of foreshocks that have occurred up to the given time (in days) before the mainshock.

Figure 1

Figure 2 a

Figure 2 b

NUMBER OF FORESHOCKS PER HOUR

60
50
40
30
20
10

6 12 18 24 30 36 42 48 54 60 66 72

TIME BEFORE MAINSHOCK IN HOURS

Figure 3

NUMBER OF EARTHQUAKES PRECEDING MAINSHOCKS

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

$\frac{A_s}{A_0}$	1.00	1.00
— 0.60	0.90	0.98
— 0.20	0.80	0.96
— 0.0	0.75	0.95
if $k\sigma_0 = 4.0$	1.0	0.2

STATISTICAL STUDY OF THE OCCURRENCE OF SHALLOW EARTHQUAKES^{*}

Yan Kagan and Leon Knopoff

Institute of Geophysics and Planetary Physics

University of California, Los Angeles, California 90024 (U.S.A.)

Abstract

The time-space-magnitude interaction of shallow earthquakes has been investigated for three catalogs: worldwide ($M \geq 7.0$), Southern and Northern California ($M \geq 4.0$) and Central California ($M \geq 1.5$). The earthquake sequences are considered as a multi-dimensional stochastic point process; the estimates of the parameters for a branching model of the seismic process are obtained by a maximum-likelihood procedure. After applying magnitude-time and magnitude-distance scaling, the pattern of relationship among earthquakes of different magnitude ranges is almost identical. The number of foreshocks diminishes as the magnitude difference between the main shock and the foreshocks increases, while the magnitude distribution of aftershocks has the opposite property. The strongest aftershocks are likely to occur at the beginning of the sequence; later they migrate away with velocities of the order of km/day. The sequences which are composed of smaller aftershocks last longer and there are indications that they remain essentially in the

^{*}Publication No. 1694 of the Institute of Geophysics and Planetary Physics, University of California, Los Angeles, California 90024, U.S.A.

focal region. Foreshocks also appear to migrate, but in this case, toward the main shock. The rate of occurrence of dependent shocks increases as t^{-1} as the origin time of the main shock is approached, effectively making every earthquake a multi-shock event. This interaction of earthquakes was modelled by a Monte-Carlo simulation technique. The statistical inversion of simulated catalogs was undertaken to derive the information we would be able to retrieve from actual data, as well as possible errors of estimates. The possibility of using these results as a tool for seismic risk prediction is discussed and evaluated.

1. Introduction

This paper is a continuation of our efforts to investigate the interaction among earthquakes using statistical methods (Kagan and Knopoff, 1976; Kagan and Knopoff, 1977; hereafter referred to as I and II respectively). In I a stochastic model of time-space-magnitude interrelationships among earthquakes was proposed and some properties of the seismicity of strong earthquakes were studied. In II we discussed the possibility of making statistical predictions of earthquakes, and evaluated the predictability of the seismic process of strong earthquakes. In I and II we used a corrected catalog of large earthquakes ($M \geq 7.0$) (Duda, 1965) extended to date, as raw data. This catalog contains less than 1000 shallow shocks; the narrow magnitude span and the small size of this catalog has prevented us from investigating some important problems of earthquake interaction, the answers to which are necessary for more accurate seismic risk calculations, as well as for understanding some aspects of the earthquake focal process. Some of these questions are

- 1) Is seismicity self-similar for collections of earthquakes having remotely different magnitudes?
- 2) Are there significant differences in the time-distance-magnitude distributions for foreshocks and aftershocks? How is the major shock anticipated? What is the detailed history of seismicity after the earthquake?
- 3) How much information can be extracted from both local and global earthquake catalogs to permit the prediction of future

seismic risk? To answer these questions we need much more complete and diverse data than those used in I and II.

In this paper we extend, with some modifications, the investigations made with the Duda catalog to three available catalogs of California earthquakes: Southern California (Allen et al., 1965; Hileman, et al., 1973), Northern California (Bolt and Miller, 1971, 1975) and Central California (Lee et al., 1971; Lester et al., 1976 which also has references to earlier reports). These catalogs contain many thousands of earthquakes and are probably the best documented, most complete, and most accurate catalogs now in existence. Because of modifications of the procedure to be described below, we have also repeated the calculations for the Duda catalog using the same model on all the catalogs.

2. The catalogs.

In order to apply our statistical model (I) to an earthquake catalog, certain conditions must be imposed, the most important of which is completeness of the catalog. By this we mean that all or almost all the earthquakes, within specified time-space-magnitude limits should be known. It is especially important that regular gaps of activity not be present in the catalogs such as those, for instance, which might occur by omitting smaller shocks immediately after strong earthquakes; such gaps could easily introduce significant bias

in the estimates of parameters and distortion of the calculated patterns of earthquake occurrence. A second condition is that the accuracy of hypocenter or epicenter determination be uniform for the entire catalog. Because the distances between epicenters are used in the distance correlations (see I), the relative error of epicenter location is of concern to us; the distance errors should be essentially constant over space and time, i.e., for different parts of the area under investigation, as well as over the time span of the catalog.

These requirements restrict the amount of earthquake data which is useful. Thus, in a given catalog, many shocks are outside the 'uniform' region of space and time. We can use these latter earthquakes to try to reduce edge effects associated with neglect of the contribution of events outside the catalog. We perform the suppression of edge effects by considering earthquakes outside the catalog interval as main events which influence earthquakes inside the catalog interval; on our model this would compensate for the lack of contribution from interactions between main shocks within the catalog and dependent earthquakes outside it, since weaker events outside the catalog are, in the main, absent. Of course, only those earthquakes outside the space and time region of the catalog which interact with shocks inside, according to the model, will be taken into consideration.

What are the spans of homogeneity of space, time and magnitude of the catalogs? The catalog which regularly reports the

lowest magnitude shocks is the Central California catalog prepared by the National Center for Earthquake Research (NCER), U.S. Geological Survey. The NCER network consists of more than one hundred stations in an area shown as region 3 in fig. 1. The network began operation in 1969 and had roughly its present-day configuration after 1970; thus the time limits of our reduced catalog have been taken as 1971-1975. With regard to space limits, we have used the boundaries given by Lee et. al. (1972b, fig. 1). These 1971 boundaries lie within enlarged perimeters given for later years (Lester et. al., 1976, and earlier reports), but to ensure uniformity, we use the smaller region. The earthquakes within these boundaries are uniformly well located, i.e., spatial variations of standard errors for hypocenter locations are small in this area. As to the magnitude cutoff of the catalog, although the lower limit of magnitudes may be as low as zero, coverage at this level is irregular; a more appropriate cutoff magnitude threshold is probably around $M=1.0$. Magnitude-frequency plots of shocks inside the network area for each year in the interval 1969-1975 are approximately parallel to each other down to magnitude 1.5; below this value, they are no longer linear and differ substantially from each other. Indeed, the plots for 1970 and 1971 show a decrease in the number of shocks below $M=1.2$ or 1.3. As a further test, we compared the frequency distribution of two sets of earthquakes: 1) shocks which have epicentral distances less than 20 km from at least 4 stations, and 2) all

the other shocks within the boundaries of the region. These distributions also appear to differ for $M \leq 1.5$. Of course, these differences may be caused by the irregular location of the stations in the network, or for geophysical reasons; as an example of the latter, variations in magnitude distributions before 1972 might have been associated with the Bear Valley earthquake of 24 February, 1972 (Wesson and Ellsworth, 1973). For reasons of caution, we use earthquakes with $M \geq 1.5$ in our catalog. In this case the catalog contains 5689 'inside' earthquakes for 1971-1975 and located within the boundaries of the region, and about 130 'outside' shocks, namely those from 1969-1970 plus those located outside the boundaries of the region up to 1975. This catalog is called USGS in this paper.

The second data set combines the Southern and Northern California catalogs. The catalogs available to us covered the periods from January 1, 1932 to December 31, 1975 for Southern California and from January 1, 1910 to June 30, 1974 for Northern California. During these time intervals the number of seismographic stations, recording equipment, methods of data processing and data presentation changed significantly. Thus the completeness and the uniformity of the catalogs is expected to be highly irregular. Both catalogs were combined to increase the total amount of data. The lower time limit for this combined catalog was chosen to be January 1, 1944; before this date magnitudes were rounded off to the nearest half-unit, while in our compu-

tational model a round-off of 0.1 of a magnitude unit has been used (I). The upper time limit was taken to be June 30, 1974 from the mutual time span of both subsets. The spatial boundaries of the combined regions, thus defining the "inside" region (fig. 1), were selected to ensure roughly uniform coverage of the area by the seismic networks after 1944 (see above mentioned references and Gutenberg, 1955). The magnitude limits are variable over the time interval between 1944 and 1974. Until the time of the Kern County Earthquake of July 21, 1952, the reliability threshold for the Southern California network was around $M=4.0$ (Knopoff and Gardner, 1969). During the Kern County aftershock sequence the cutoff magnitude remained about 4.0 for more than 100 days (Richter, 1955). Subsequent to this earthquake an improvement in the instrumental density of the network lowered the threshold to about $M=3.0$ (Knopoff and Gardner, 1969). A significant increase in the density of stations in the Northern California network plus the start of a transition to computer techniques for locating epicenters began around 1961 (Bolt and Miller, 1975). We have taken $M_L=4.0$ as a magnitude threshold for the first part of the combined catalog; from January 1, 1962 to the end of the catalog we have used a magnitude threshold of $M_L=3.5$. The combined California Catalog (CACA) thus defined contains 997 events inside the interval and 246 events outside it.

With regard to the worldwide catalog of large shallow earthquakes the Duda list (1965) forms the bulk of our catalog; the lower magnitude limit is $M_S=7.0$. We have extended this catalog

to April, 1975 using the PDE data file (Meyers and von Hake, 1976). As a lower time limit we have used January 1, 1905; before this date, listings of weaker events ($7.0 < M < 8.0$) are incomplete. December 31, 1972 was taken as the upper time limit. This catalog (called DUDA below) contains 914 'inside' events and 81 'outside' events.

In the following discussion, a single magnitude scale is used which incorporates, without change of scale, both the M_S of DUDA and the M_L of CACA and USGS.

3. Moments

The second-order moments of a seismic process represent a convenient exploratory tool for studying interactions among earthquake events (I). In I we studied the conditional intensity function or conditional moment $m(\underline{x}/\underline{x}_1)$ of a seismic process, i.e. the number of earthquakes in a certain space-time interval $\underline{x}_1$ given the occurrence of an earthquake at a point $\underline{x}$. As raw data the DUDA catalog was used. When we tried to apply a similar technique to the Southern California catalog, we found that the seismic history of Southern California of the past 30 years is dominated by two large earthquakes: Kern County (1952) and San Fernando (1971). Thus a large number of the shocks listed in the catalog are stochastically dependent on these earthquakes, and hence many of the events are statistically connected, although not necessarily directly.

The conditional time-distance moment will consist of contributions from both direct interactions and secondary interrelationships; the latter are represented by convolutions in the time-space domain of the former. The most interesting features of earthquake interactions are likely to be smoothed out by strong secondary interference effects in such convolutions. For now, we indicate that the second-order moments from the Southern California catalog give approximately the same information as obtained from DUDA (I); below we will return to this problem.

Before proceeding to the analysis of these catalogs using a stochastic model, we discuss the conditional spatial moment of the earthquake process. In I we found that the quantity $N(dr)/r$ is uniformly distributed over the distance range 200 - 2000 km for large worldwide earthquakes, where $N(dr)$ is the number of shocks in the distance interval between r and $r+dr$ from a given epicenter. We have checked whether the property of uniformity of this function applies to an earthquake process on a much smaller scale through the study of the earthquake catalogs for California. Catalog USGS provides us with a unique opportunity to study the three-dimensional, or hypocentral, spatial moments, since it has estimates of focal depths which are probably as reliable as can be obtained routinely today. When we try to study the spatial distribution of USGS hypocenters we find that we must take into account edge effects due to boundaries of the region (see fig. 1) and the non-uniform distribution of seismicity with depth: about one-half

of all USGS earthquakes are concentrated in a depth interval 4 to 7 km. To take these effects into account we have first calculated the (conditional) number of USGS hypocenters $N(V_r)$, which are situated at distances less than r from a given earthquake. These computations were then repeated with a spatially randomized catalog, where the same total number of shocks was distributed uniformly over the horizontal plane, but the depth distribution was taken to be the same as the USGS catalog. The number of shocks in the simulated, i.e. Poissonian catalog, for small distances $r < 3.2$ km., was too small for reliable comparison, but we can calculate these quantities, using the formulas

$$N_p(V_r) = \frac{4\pi}{S} \int_0^r R dR \int_0^R K(h) dh$$

where depth covariance function $K(h)$ is defined as:

$$K(h) = \frac{1}{(\Delta z)^2} \int N(z, z+\Delta z) \cdot N(z+h, z+h+\Delta z) dz$$

Here $N_p(V_r)$ is the number of pairs of hypocenters in a sphere of radius r centered on one shock focus; S is the surface area spanned by the USGS catalog which is approximately, 19200 km^2 , $N(z, z+\Delta z)$ is the number of shocks in the depth interval $(z, z + \Delta z)$.

The ratio $N(V_r)/N_p(V_r)$ gives the spatial moment behavior as if the shocks were distributed uniformly in all three directions. For display purposes this ratio is multiplied by r (fig. 2). This product is essentially uniform over distances from 1 km to more than 20 km. For distances less than 1 km the hypocenter distribution is dominated by location errors; independent estimates of these errors are approximately 1 km, (Lee et. al.,

1971, Lester et. al., 1976).

The increase in this curve (see fig. 2) for $r > 20$ km. is most probably connected with bias in choosing the regional boundaries of USGS with respect to the seismicity pattern. If, for example, we were to rotate the region of USGS so that its longer side is perpendicular to the San Andreas fault, the number of long distance shocks would decrease significantly; a random orientation of the polygon would probably yield a uniform distribution of $r \cdot N(V_r) / N_p(V_r)$ over the entire long-distance range.

Taking these considerations into account, we shall assume that the curve of spatial moment is proportional to r^{-1} over the entire range of distances. Since this result arose in I as well, we propose the conjecture that this is a general feature of the spatial distribution of earthquakes, pointing to some property of the geometry of faulting. For example, our conjecture rules out a spatially isotropic uniform distribution of earthquake faults. For the purposes of comparison of results later on, we have used only two-dimensional spatial moments in our calculations of the likelihood function for USGS, shown also in the lower curve of fig. 2; in this figure (S_r) signifies a circle of radius r .

4. Likelihood function calculations

To analyze catalogs such as the two from California, we need to apply a stochastic model of the earthquake process which takes all possible interactions into account. In this paper we

have applied essentially the same branching model of earthquake occurrence used earlier (I, II); this is a restricted version of the model of complete interaction. In particular, the function describing the interaction between earthquakes is taken to be uniform within each of several small time-distance intervals surrounding the main event. This type of function has greater versatility in comparison with algebraic functions (I, p. 308). The time-distance plot around each earthquake was divided into 9 time and 4 distance intervals, i.e., 36 space-time intervals both before and after the main event (fig. 3). The equally spaced parallel diagonal lines in the left-hand diagram of this figure show the boundaries of the time intervals as a function of magnitude of main event. Each of these intervals is subdivided into four distance intervals as shown on the right side of the figure. We number the intervals of time sequentially from the shortest to the longest time interval, and number the intervals of distance similarly from the shortest to the longest range. We assume the magnitude distribution of dependent events varies as $c^{\Delta M}$, where c is an adjustable parameter in the fit and ΔM is the difference of magnitude between the main event and a dependent shock. Here, as in previous investigations, it is more convenient to identify each 0.1 magnitude interval by an integer k

$$k_i = 10(M_i - M_{\text{cutoff}}) + 1 \quad (1)$$

The logarithmic likelihood function for the branching model may be calculated by the expression (formula 6 in I, formula 3

in II).

$$\begin{aligned}
 \ell = & -A \cdot \frac{\gamma^{k_{\max}+1} - 1}{\gamma - 1} - \sum_{i=1}^n \sum_{s=1}^4 \left(\sum_{t=1}^9 \mu_{ts}^- \cdot \frac{c_i^- - c_-}{c_- - 1} + \sum_{t=1}^9 \mu_{ts}^+ \cdot \frac{c_i^+ - c_+}{c_+ - 1} \right) \\
 & + \sum_{i=1}^n \log \frac{v(h_i) d\tau \cdot dh}{\Pi(H) \cdot T} + \sum_{i=1}^n \log \left[A \cdot \gamma^{k_i-1} + \right. \\
 & \left. + \sum_{k_i < k_j} \mu_{ts} \cdot \frac{\Pi(H) \cdot T}{\Pi_j(\Delta H_s) \cdot \Delta T_t} \right] + \text{const } (d\tau, dh) \quad (2)
 \end{aligned}$$

In this formula the subscripts $\pm$ refer to aftershocks and foreshocks respectively, t and s are the indices of the time and distance intervals, T is total time span of the catalog, ΔT_t is the time-span of the t^{th} time interval, $v(h)$ is the level of seismicity of independent shocks, and $\Pi(\Delta H_s)$ is the average number of independent shocks in the s^{th} ring around the main shock, $\Pi(A) = \int_A v(h) dh$ and n is the total number of earthquakes in the 'inside' catalog. A and γ correspond roughly to the parameters "a" and "b" in the magnitude frequency relations; and μ the branching coefficient defining the number of dependent shocks per 0.1 magnitude interval.

The logarithm of the likelihood function for a Poissonian model of the process ℓ_0 may be calculated by setting the value of the branching coefficients μ in formula 2 to be zero. Actually we are interested in the difference $(\ell - \ell_0)$ and for this quantity, the third and fifth terms in formula (2) are absent.

In order to intercompare the three catalogs some scaling must be applied. The differences in the principal magnitudes of shocks in each of the three catalogs means that the duration of after-shock sequences and the distances over which aftershocks might be expected to occur should be an increasing function of magnitude. Such a model was proposed by Knopoff and Gardner (1972) and Gardner and Knopoff (1974).

The determination of the appropriate distance-magnitude scaling law presents a difficult problem. For the smallest earthquakes in a catalog, the size of the focal zone, or the zone where the shock influences the stress pattern, is much less than the error in locating the earthquakes. A similar problem arises when we use the epicentral, rather than the hypocentral, distance between the shocks, since depth estimates are much less accurate than those for epicentral coordinates. Thus the details of the interaction pattern of the smallest earthquakes in a given catalog are smoothed due to location errors. As noted above the relative errors are of importance because we use the distances between events to compute the likelihood function; as a consequence, the average values of the standard errors in epicenter location should roughly define the distance interval scale for small shocks. For USGS these errors are 0.5-1.0 km; for CACA the errors are 1.5-4.0 km.

With regard to the upper magnitude limit of a catalog, empirical magnitude-distance relations have been well-studied (King and

Knopoff, 1968; Kanamori and Anderson, 1975). A combination of the magnitude-fault length relation, together with the increasing importance of uncertainties in epicenter location at small magnitudes, has been considered by Knopoff and Gardner (1972) for the NOAA catalog 1964-71 where it is assumed that the uncertainties in location overtake the distance-magnitude formula around $M=5$. The Knopoff-Gardner (1972) relation, which is intended as an upper bound to isolate aftershock sequences, is shown in fig. 3 as line 5. In this paper we use a similar relation with logarithmic subdivisions of the distance intervals as shown in fig. 3. Several calculations have been made with different sets of parameters to define the intervals, and those shown in fig. 3 have been found appropriate for this calculation by being close to the maximum of the likelihood function.

To define the magnitude-time scaling parameters (I, pp. 302-303), we have calculated the value of $\ell-\ell_0$ for different values of the scaling factor θ , which is defined so that if the magnitude of the main shock increases by one unit, the time interval is changed by a factor of θ , i.e. ΔT_t in formula 2 will be changed to $\Delta T_t \cdot \theta^{\Delta M}$ (see fig. 3). For values of θ between 2.0 and 4.0 there is very little change in the likelihood function (I, p. 303); very likely this means that, for our model (equation 2), θ is closely correlated with some of the other parameters and hence is ill-defined.

There are two alternative ways to compute the factor θ . In the first alternative we adjust the parameter θ to match the relationship between the duration of the earthquake signal and the magnitude of the shock. During a large earthquake it is difficult to determine arrival times of close aftershocks of the event; hence a catalog is incomplete.

The data specifying the magnitude-signal duration dependence for small shallow earthquakes (Lee et. al., 1972a, Real and Teng, 1973), plotted in fig. 3 as lines 1 and 2 respectively suggest that $\theta=3.0$ would be appropriate for our purposes (see also Herrmann, 1975). The long-period (M_S) data (Evernden, 1976, p. 550) also support this value, though not so definitively. The durations in line 3 were calculated using an amplitude level corresponding to $M_S=5.5$. This cutoff amplitude limit seems to be appropriate for the largest part of DUDA.

The second alternative uses a postulated relation between an upper bound for the duration of aftershock sequences and the magnitude of the main shock given by Gardner and Knopoff (1974) for earthquakes in Southern California; this curve is plotted in fig. 3 as line 4 and also gives a value of $\theta=3.0$. The corresponding curve from the NOAA list of worldwide earthquakes also gives the same value (Knopoff and Gardner, 1972). The value $\theta=3.0$ was used in our computations.

To avoid the inhomogeneities introduced into a catalog at short times after a large event, we have deleted all earthquakes

from the catalogs which lie to the left of line A in fig. 3, which is our estimate of the duration of the earthquake signal. The number of earthquakes removed for the 'inside' USGS catalog is 103 or about 2%; for the CACA catalog it is 160 or about 16%, and for the DUDA catalog it is 20, also about 2%. In section 8 we discuss possible bias that might be introduced by such removal of close, dependent shocks.

5. Results

The values of the parameters obtained by maximizing the likelihood function of formula 2 and some characteristics of the catalogs are shown in table 1. The estimates of errors were obtained by inversion of the matrix of second derivatives $\partial^2 \ell / \partial P_i \partial P_j$, where P_i is the i^{th} parameter.

We have the relation $b = 10 \log_{10} \gamma$ for independent earthquakes, and for the Poissonian model $b_0 = 10 \log_{10} \gamma_0$. The results show a slight increase of b_0 with M_{max} ; this may reflect a weak non-linearity of the magnitude-frequency curve plot over the whole range of magnitudes or it might be due to our use of the different magnitude scales M_S and M_L (see end of section two). With regard to the parameter a in the magnitude-frequency law, we write this law in the form

$$\text{Log}_{10} \dot{N} = a - bM_r - b (M - M_r)$$

The usual value of the activity coefficient a corresponds to the reference magnitude $M_r = 0$. However, $M_r = 0$ is usually a

magnitude outside the span encompassed by most catalogs. Hence the value of $\underline{a}$ is statistically correlated with values of $\underline{b}$. A statistically independent estimate of $(a - bM_r)$ would be to take M_r near the "center of gravity" of the magnitude-frequency data. We have, for convenience used a value close to this, namely,

$$\alpha = a - b (M_{\text{cutoff}} + 0.5)$$

In other words, α is the ordinate at $M_r = M_{\text{cutoff}} + 0.5$. The parameter $\underline{A}$ in equation (2) is the ordinate at the upper magnitude limit of the catalog.

The parameters (γ, α) describe the distribution of independent shocks for our branching model. The number of these shocks is significantly smaller in comparison with the Poissonian model. The parameters $\Sigma\mu^-$ and $\Sigma\mu^+$ describe dependent shocks: the former gives the total number of foreshocks and the latter the number of aftershocks in the magnitude interval 0.1 unit as the difference of magnitude between these shocks and the main event approaches zero. In the calculations we have not considered the interaction of earthquakes with equal integer magnitudes k but,

if this interaction were taken into account, it would only increase the value of the likelihood function by about 3%. The values of $\Sigma\mu^-$ and $\Sigma\mu^+$ are approximately equal for all three catalogs; it should be noted that we have used 8 time intervals for DUDA instead of 9 so we expect lower values for this catalog. The quantities c_- and c_+ give the magnitude

distributions of the foreshocks and aftershocks respectively (Eq. 2). For example, the value of $c_-^{10}=0.51$ means that for each change of one unit of magnitude, the number of foreshocks changes 0.51 - fold. Once again there is great similarity among the c_+ , c_- values for all three catalogs. The similarity of the c_+ and the μ_+ values may be a fortuitous consequence of close correlation between estimates of these parameters; the correlation coefficient is of the order 0.1 - 0.4. Hence we have calculated the number of dependent shocks v for a magnitude difference of one unit:

$$v = \frac{c^{11} - c}{c - 1} \cdot \Sigma \mu \quad (3)$$

and once again we have good agreement among all three catalogs. The N-values (Table 1) are the total number (and the percentage) of dependent shocks throughout the catalog.

The c-values are significantly different with regard to the magnitude distribution of the fore- and aftershocks: there are many aftershocks which are much weaker than the main shock, while the number of foreshocks diminishes as the magnitude difference between the main shock and the foreshocks increases. It is interesting to note that $c_- \approx 1/c_+$ and $\Sigma \mu^- \approx \Sigma \mu^+$ for each catalog. This means that in reality there are no strict differences between fore- and aftershocks; they are simply descriptions of the same mutual interaction of earthquakes. We discuss this notion further in the next section.

The last row of Table 1 gives the information rate per earth-

quake, i.e. how much information we can obtain about an earthquake, on the average, from a given catalog, for this particular model of the earthquake process (this calculation uses base 2 logarithms); see II for more detail. This rate depends strongly on the magnitude span of the catalog; another influence, as will be seen below, is the presence of a very strong earthquake in the catalog. With the data presently at our disposal, we are able to obtain information about future earthquakes in the amount of 0.5 to 1.0 bits per shock, i.e., to reduce the uncertainty of the predicted event by a factor of $2^{0.5}$ to $2^{1.0}$ (II).

The similarity of earthquake precesses in the different magnitude ranges is exemplified further by Table 2, which shows the time-distance interaction of earthquakes. Values of μ such as 10 ± 16 should not be taken to mean that μ can take on a negative value: by definition μ is non-negative. As noted in I, these $\pm$ values give only a crude estimate of uncertainty. Values of μ that are most reliable are those for times closest to the origin time of the main earthquake, i.e. corresponding to the smaller sequence numbers of time intervals. These estimates of μ are also less dependent on the specific assumptions regarding the model and these estimates are virtually uncorrelated statistically with the estimates of the other parameters than $\underline{c}$, the values of these correlation coefficients are generally less than 0.1. The values of μ for longer time intervals, i.e. larger sequence numbers show much larger scatter. Taking these

observations into account, we assert that the values of μ are similar for all three catalogs. With increasing distance, the larger values of μ are found among the greater time intervals. This means that aftershocks are migrating with velocities of from 2 to 10 km/day at the beginning of the sequence and from 0.1 to 0.5 km/day for remote aftershocks. Foreshocks seem to move toward the main shock.

The time-distance interaction of earthquakes in different magnitude ranges can be investigated by constructing four-dimensional distributions, with the magnitude of the main event, the magnitude of the dependent event, the distance and the time as the variables. Indeed, by putting the values of the parameters we have already derived into the computational scheme, it is possible to calculate the number of dependent shocks in any given 4-dimensional interval; this has all the advantages of the moment approach (see section 3) and differs only in that all the major interactions will have been taken into account automatically. From this calculation we find for example that the strongest earthquakes in the California catalogs (especially in CACA) were not preceded by significant numbers of foreshocks; the non-zero values of μ corresponding to foreshocks, are connected mostly with earthquakes of small and medium size. This does not invalidate the concept of foreshocks as a common and frequent occurrence (I; Jones and Molnar, 1976). In CACA there are only two major earthquakes, so the statistics are poor; DUDA

gives many examples of strong earthquakes preceded by foreshocks. However, the annoying possibility exists that California may be anomalous in regard to the presence of foreshocks; we simply do not have sufficient data.

Another feature we have investigated is the behavior of different sizes of aftershocks. For a specified time interval, we compared the numbers of aftershocks in the region that is most distant from the focal zone, distance intervals 3 and 4, with the total number of aftershocks; the latter number is the sum of the entries in the four distance intervals. The time intervals we have chosen are grouped by threes to increase the population in the entries. These ratios are shown in Table 3 for different magnitude ranges of main and dependent earthquakes. The values of $\Delta M = M_1 - M_2$ between the centers of each magnitude band are also listed; entries for the same values of ΔM with different values of M_1 are connected by braces.

Four significant observations can be made with regard to Table 3. These are 1) the increase in the percentages from left to right across the top row, which can be interpreted to mean that strong aftershocks, with $\Delta M = 0$, are concentrated close to the epicenter in the early part of the time sequence and later large aftershocks begin to migrate into the regions more remote from the epicenter; 2) the decrease in the percentages from left to right across the bottom row, which can be interpreted as implying that weaker aftershocks do not appear in large numbers

at long range and at short times after the parent event and indeed this number becomes even smaller at long range in later times; 3) the decrease in values from top to bottom in the last column implies that we find a greater percentage of the total large aftershock population than the small aftershock population at large distances and large times from a parent earthquake; 4) the rather close agreement between the two pairs of rows connected by braces which testifies to the stability of the analysis. Unfortunately the number of aftershocks with $M_2 > 3.5$ is too small to draw reliable conclusions regarding the extension of this pattern. Partial confirmation may also be seen in Table 2, where the μ -values for DUDA soon achieve a zero value at the epicenter (distance intervals 1 and 2), while the μ -values for USGS, with a much greater magnitude span, remain non-zero for all time.

Similar results are obtained when we replace the two values of $\underline{c}$ in formula 2 by twelve values, each $\underline{c}$ -parameter being assigned to a region consisting of 3×2 time-distance intervals (fig. 1). The gain in the likelihood function for the three catalogs was 12.8, 11.2 and 2.2 respectively; for USGS and CACA the results are significant at a level greater than 98%; the results for DUDA of course are not statistically significant.

From the analysis of the USGS data (Table 4), we see that the values of $\underline{c}$ for foreshocks are in general lower near the epicenter than farther away; for aftershocks the pattern is changed in the opposite

sense. As indicated above, low values of $\underline{c}$ show the prevalence of strong dependent shocks and large values of $\underline{c}$, their absence. Similar observations can be made for CACA aftershocks, but the pattern is not confirmed for CACA foreshocks. In any case the effect is small and its statistical reliability has not been fully tested. With these restrictions, we see that strong earthquakes are taking place at the site of the future main shock; after the main earthquake they migrate to the outer fringes of the focal area. In the focal area, weak shocks are suppressed before the main shock, but after the main shock the number of small shocks increases significantly. The changes of our parameter $\underline{c}$ correspond to changes of the $\underline{b}$ -value of the magnitude-frequency law before and after a strong earthquake (Wyss and Lee, 1973). A low value of $\underline{c}$ corresponds to a decrease of the $\underline{b}$ -value, but this decrease depends also on the ratio of dependent to independent shocks in the sample. Near the time of the main shock when the rate of dependent shocks is high, the changes in $\underline{b}$ are great, etc.

6. Close interaction of shocks and pattern of earthquake occurrence

In the last row of Table 2 we give the sum over all distance intervals of the average values of the ratio μ . These values are roughly uniform over time. Since the time intervals we have used (fig. 3) increase logarithmically, the rate of occurrence of dependent shocks increases as t^{-1} as we approach the time

of the main shock. This is in fact the Omori law of aftershock occurrence (Richter, 1958, Utsu, 1969). A similar picture was obtained for foreshocks by Jones and Molnar (1976).

If we try to extrapolate these values of μ toward $t \rightarrow 0$ we will get an infinite number of foreshocks and aftershocks. Of course this is meaningless, since our notion of earthquakes as a point process, i.e., that earthquakes occur instantly at one point of the space, ceases to be valid as we approach the origin time of the earthquake. The duration and the dimensions of an earthquake are not infinitesimal. We can avoid this contradiction by assuming, for example, that the earthquake does not generate dependent shocks during the process of rupture. For reasonable estimates of the focal dimensions and velocity of rupture we get a rough estimate of the rupture duration shown as line B in fig. 3. Those dependent events which take place in the time interval between lines A and B (fig. 3) usually are not distinguishable (see also discussion in section 4), but we may assume that their rate of occurrence increases continually with decreasing time interval to the origin time of the main shock. Recently there has been some discussion concerning these short range multiple shocks (Trifunac, 1972; Fukao and Furumoto, 1975; Burdick and Mellman, 1976).

Assuming the value of μ is constant in the interval between lines A and B, we can try to estimate the total number of aftershocks v generated by the main shock and its aftershocks (Kagan, 1973),

$$v = c\mu \frac{[c(1+\mu)]^k - 1}{c(1+\mu) - 1} \quad (4)$$

Taking $c=1.195$, and $\mu=0.014$ we get for $\Delta M=1.0$ the value $v=0.46$; for $\Delta M=2.0$ we get $v=3.60$. From the relation $E \sim 10^{1.5M}$ we find that close aftershocks on the average have 10% of the energy of the main shock, if the extrapolation of our model toward $t \rightarrow 0$ is valid. These estimates are approximate because we have not taken into account foreshocks and the possibility that a second-order aftershock, i.e. an aftershock of an aftershock, might fall outside the time interval. More exact calculations will be described in the next section.

The following picture of earthquake occurrence can be constructed. A forthcoming strong main shock suppresses the creation of small earthquakes in the future focal area; thus we have a decrease of seismic activity, or a seismic gap, before this earthquake. As the time of the main shock approaches, foreshocks migrate toward it and their rate of occurrence increases. The foreshocks are strong relative to the main shock. Finally one of them triggers an event which ruptures the entire focal region, thus becoming the main shock. Near the origin time there is actually a merger of foreshocks, main event and close aftershocks. During this time the c-value becomes larger than 1.0, reflecting the creation of many smaller earthquakes. Strong aftershocks soon migrate beyond the focal area, leaving behind mostly weaker earthquakes.

Unfortunately, the available data do not permit the evaluation of the pattern of earthquake occurrence in the time intervals far from strong earthquakes.

7. Monte Carlo simulation of an Earthquake 'family'

As indicated above, we may use Monte Carlo techniques to estimate more precisely the number of close dependent events, which may occur during the main phase of an earthquake. For these calculations we have applied a continuum model of the interaction function between earthquakes, which is consistent with our results. The magnitude distribution of dependent shocks was taken to be $\exp(c \Delta M)$ and the rate of occurrence to be μt^{-1} . We take $c_- = -1.55$, $\mu^- = 0.0315$, $c_+ = 1.78$, $\mu^+ = 0.0205$; (these values are approximately equal to those obtained in the previous section). The total number of dependent shocks in the time interval between lines A and B (fig. 3) was calculated both before and after $t=0$; that is, both close foreshocks and aftershocks were taken into account. For $\Delta M=1.0$ this number is equal to 0.59, and increases 7.5-fold for each increase of ΔM by one unit, in agreement with our previous calculation.

We can check to see if our conclusions that a shallow earthquake can be considered to be a multiple event (Miyamura, 1965) are compatible with the available data, i.e., is the extrapolation of our model toward $t=0$ warranted? The most complete data for the time interval immediately after an earthquake are obtained from the accelerometer record of the

1971 San Fernando earthquake, during the first six minutes of the earthquake. Fig. 4a displays the time sequence for all shocks with $M \geq 4.5$ (Trifunac, 1972) during this period. In Figs. 4b and 4c we show two examples of realizations of the aftershock sequence obtained by a Monte Carlo procedure using the values of parameters in the preceding paragraph. In addition, similar simulations may be made as a first step of generating synthetic seismograms. By averaging 2000 such realizations, we calculate that in only about 4% of the cases, are 7 or more aftershocks expected in this time interval; the average number of aftershocks is about 2.5. So we face the dilemma that either the 1971 earthquake was a relatively rare event, having 7 aftershocks in 6 minutes, or the values of the parameters should be greater. Similar results can be obtained for the other major California earthquake, the 1940 Imperial Valley event (Trifunac and Brune, 1970).

8. Monte Carlo simulations of catalogs

A similar simulation technique can be used to generate artificial catalogs of earthquake data (cf. Knopoff and Kagan, 1977). Several problems are solved by creating the catalogs by Monte Carlo methods and then inverting them by the maximum likelihood method. Among these 1) we can test the general validity of our concepts and the computational scheme, 2) we can estimate biases intro-

duced by removing shocks at short time intervals or to simulate the absence of them in the catalog and 3) we can evaluate the statistical errors associated with our estimates of the parameters.

To reduce computing time we restrict this calculation to one in a time-magnitude state space, i.e., we ignore the spatial distribution of the earthquakes. In its other details, the procedure was similar to that shown in fig. 4, or to that described in II. To create a catalog similar to CACA, earthquakes with magnitude 4.0 and greater were generated over a time span of 60 years; the central 40 years form the 'inside' catalog. The interaction function was taken to be uniform over all 15 time intervals: nine of them are shown in fig. 3 and the remaining six are located between lines A and B in fig. 3; we take $\mu = 0.00260$, $c_+ = 1.17$, $c_- = 1/c_+$, $\gamma = 1.236$, $\alpha = 0.55$ earthquake/year. These values are roughly equivalent to those obtained for CACA: α is the number of earthquakes with magnitude $4.5 \pm .05$. The principal results for five realizations of the 'inside' catalog are given in table 5. The number of independent shocks varies slightly, but the number of dependent shocks is strongly connected to the largest magnitude of the earthquake in the catalog.

We have inverted the catalog data to derive the values of parameters. A first round of calculation was made with a catalog in its original form (column 'a' in table 6) and a second round with the close dependent shocks removed (column 'b'). The first three catalogs in table 5 were processed; the results

are presented in table 6 in a format compatible with that of table 1. The parameters α and γ have a great deal of scatter, especially in case b, where the removal of close dependent shocks seems to cause an increase of c and to a lesser degree μ , so that the estimate of the number of independent shocks N_{ind} is drastically reduced. We propose that the close strong shocks which are removed are 'parents' of some of the remaining shocks, so these missing parents should be replaced by a greater "productivity", as reflected in the values of the parameters μ and $\underline{c}$ of the remaining ⁱⁿ strong shocks. A closer investigation shows that the remote time intervals have the greatest change of the values of the parameters, up to 50%; for closer intervals the bias is about 20%. This may explain the slight increase of the average μ -values for larger time intervals (see rows $\Sigma\mu$ of Table 2). The estimates of parameters for 15 time intervals (case 'a') show a rather good agreement with the original values: $\bar{c}_+ = 1.173 \pm 0.00048$. The values of N_{ind} display the largest differences; these values are all less than the expected value of 332.5. This bias is probably due to our use of outside shocks (see section 2); if we do not use outside shocks it will result in a negative bias for the estimates of the parameters $\underline{c}$ and μ . It is possible, of course, to make more exact calculations, as we did in II, taking into account all the time and distance intervals between the shocks and the edges of a catalog, but this will not be done here.

The values of the likelihood function and the information rate show significant dependence on the maximum magnitude of the earthquake in the catalog (section 5). These values are in general lower than similar estimates for real catalogs (table 1), reflecting the absence of spatial information in the simulated catalogs. The large difference between the values for cases 'a' and 'b' show once again (II) that most of the information in the catalogs is contained in interactions of close events.

9. Discussion and Summary

A coherent quantitative pattern of earthquake occurrence has been derived from the statistical analysis of three catalogs of shallow earthquakes. This picture synthesizes most of the known results of the time-space-magnitude interaction of earthquakes: the seismic gaps before earthquakes, the migration of epicenters, the change of b-values before and after a strong earthquake, the increase in the rate of foreshock occurrence before strong shocks, the complexity of the earthquake source function, etc. It has also been shown that the earthquake process is uniform over a broad range of magnitudes, extending from $M=1.5$ to the largest earthquakes.

We can use our results in two ways: the first is an extrapolation of the catalog in time, i.e., seismic risk prediction as in (II). As we have shown the catalogs of earthquakes now available supply us with about 0.5-1.0 bits of information per earthquake. It is possible that different models of earthquake occurrence may raise this value somewhat, but it is doubtful

that substantial improvement is possible. To show what improvement we need, consider the following example. From the values of parameters listed in Table 1, we calculate, using Poissonian model, that the average return period of an earthquake with magnitude $8.0 > M \geq 7.5$ is about 100 years, in the part of California spanned by CACA. If we had 10 bits of information per earthquake, we could reduce the uncertainty by 2^{10} times, resulting in an uncertainty in the conditional return period of a little more than one month. If an information rate of 20 bits were available, the earthquake could be pinpointed with an accuracy of about one hour.

In order to make this improvement possible, the information content of the catalog must be enlarged significantly. Since we cannot increase the duration span of catalogs, an alternative would be to expand the description of an earthquake source, which is given at present in terms of only one parameter, the magnitude, to a more complicated, multi-dimensional description which may involve retrieving the complex earthquake source-time function from seismographic records (Gilbert and Dziewonski, 1975). Here our results relating to the interaction of close earthquakes might be of help by providing an estimate of the a priori distribution of dependent shocks. Hence, a modified stochastic theory of earthquake occurrence can in principle contribute to the solution of one of the major inverse problems of seismology, namely the retrieval of the complex

earthquake moment tensor function from seismograms.

A second application is the extrapolation of our results toward the origin time of the main shock, by the simulation of the complex source model of an earthquake. The forward problem involves the construction of synthetic seismograms for strong earthquakes, where the stochastic interaction of shocks is taken into account. The implications for engineering seismology are obvious.

Acknowledgements

This research was supported by Grant ENV76-01706 of the RANN program of the National Science Foundation. The authors wish to thank W. M. K. Lee, J. Pfluke, M. Friedmann, and B. A. Bolt for providing us with relevant catalogs. The authors appreciate useful comments received from D. Vere-Jones and K. Aki.

References

- Allen, C. R., P. St. Amand, C. F. Richter, and J. M. Nordquist, 1965. Relationship between seismicity and geologic structure in the Southern California region. Bull. Seism. Soc. Am., 55, 753-797.
- Bolt, B. A., and R. D. Miller, 1971. Seismicity of Northern and Central California, 1965-1969. Bull. Seism. Soc. Am., 61, 1831-1847.
- Bolt, B. A., and R. D. Miller, 1975. Catalogue of Earthquakes in Northern California and adjoining areas, 1 January 1910-31, December 1972. Seismographic stations, University of California, Berkeley, pp. 567.
- Burdick, L. J., and G. R. Mellman, 1976. Inversion of the body waves from the Borrego mountain earthquake to the source mechanism. Bull. Seism. Soc. Am., 66, 1485-1499.
- Duda, S. J., 1965. Secular seismic release in the circum-Pacific belt. Tectonophysics, 2, 409-452.
- Evernden, J. F., 1976. Study of seismic evasion, III. Bull. Seism. Soc. Am., 66, 549-592.
- Fukao, Y., and M. Furumoto, 1975. Foreshocks and multiple shocks of large earthquakes. Physics of Earth and Planetary Interiors, 10, 355-368.

- Gardner, J. K., and L. Knopoff, 1974. Is the sequence of earthquakes in Southern California with aftershocks removed, Poissonian? *Bull. Seism. Soc. Am.*, 64, 1363-1367.
- Gilbert, F., and A. M. Dziewonski, 1975. An application of normal mode theory to the retrieval of structural parameters and source mechanisms from seismic spectra. *Phil. Trans. Roy. Soc. London, Ser. A.*, 278, 187-269.
- Gutenberg, B., 1955. Seismograph stations in California in: Earthquakes in Kern county, California during 1952, California division of mines, Bull. 171, pp. 153-156.
- Herrmann, R. B., 1975. The use of duration as a measure of seismic moment and magnitude. *Bull. Seism. Soc. Am.*, 65, 899-914.
- Hileman, J. A., C. R. Allen, and J. M. Nordquist, 1973. Seismicity of the Southern California region, 1 January 1932 to 31 December 1972. Seismological Laboratory, California Institute of Technology, Pasadena, 490 pp.
- Jones, L. and P. Molnar, 1976. Frequency of foreshocks. *Nature*, 262, 677-679.
- Kagan, Y. Y., 1973. A probabilistic description of the seismic regime. *Izv. Acad. Sci. U.S.S.R., Phys. Solid Earth*, pp. 213-219 (English translation).
- Kagan, Y. and L. Knopoff, 1976. Statistical search for non-random features of the seismicity of strong earthquakes. *Physics of Earth and Planetary Interiors*, 12, 291-318.

- Kagan, Y. and L. Knopoff, 1977. Earthquake risk prediction as a stochastic process. *Physics of Earth and Planetary Interiors*, 14, 97-108.
- Kanamori, H., and D. L. Anderson, 1975. Theoretical basis of some empirical relations in seismology. *Bull. Seism. Soc. Am.*, 65, 1073-1095.
- King, C. -Y., and L. Knopoff, 1968. Stress drop in earthquakes. *Bull. Seis. Soc. Am.*, 58, 249-257.
- Knopoff, L., and J. K. Gardner, 1969. Homogeneous catalogs of earthquakes. *Proc. Nat. Acad. Sciences*, 63, 1051-1054.
- Knopoff, L., and J. K. Gardner, 1972. Higher seismic activity during local night on the raw-world-wide earthquake catalog. *Geophys. J. Roy. Astron. Soc.*, 28, 311-313.
- Knopoff, L., and Y. Kagan, 1977. Analysis of the theory of extremes as applied to earthquake problems. Accepted by *J. Geophys. Res.*, 82, 5647-5657.
- Lee, W. H. K., M. S. Eaton, and E. E. Brabb, 1971. The earthquake sequence near Danville, California, 1970. *Bull. Seism. Soc. Am.*, 61, 1771-1794.
- Lee, W. H. K., R. E. Bennett, and K. L. Meagher, 1972a. A method of estimating magnitude of local earthquakes from signal duration. U.S. Geological Survey, Open File Report, 28 p.

- Lee, W. H. K., K. L. Meagher, R. E. Bennett, and E. E. Matamoros, 1972b. Catalog of earthquakes along the San Andreas fault system in Central California for the year 1971. U.S. Geological Survey, Open File Report, 67 p.
- Lester, F. W., S. L. Kirkman, and K. L. Meagher, 1976. Catalog of earthquakes along San Andreas fault system in Central California, October-December, 1973. U.S. Geological Survey, Open File Report, 37 p.
- Meyers, H. and C. A. von Hake, 1976. Earthquake data file summary. National Geophysical and Solar-Terrestrial Data Center, Boulder, Colorado, 56 pp.
- Miyamura, S., S. Omote, R. Teisseyre, and E. Vesanen, 1965. Multiple shocks and earthquake series pattern. Bull. Int. Inst. Seism. Earthq. Eng., 2, 71-92.
- Real, C. R., and T. L. Teng, 1973. Local Richter magnitude and total signal duration in Southern California. Bull. Seism. Soc. Am., 63, 1809-1827.
- Richter, C. F., 1955. Foreshocks and aftershocks, in: Earthquakes in Kern county, California during 1952, California division of Mines, Bull., 171, San Francisco, pp. 177-198.
- Richter, C. F., 1958. Elementary Seismology. W. H. Freeman & Co., San Francisco, 768 pp.

- Trifunac, M. D., 1972. Stress estimates for the San Fernando, California earthquake of February 9, 1971: Main event and thirteen aftershocks. *Bull. Seism. Soc. Am.*, 62, 721-750.
- Trifunac, M. D., and J. N. Brune, 1970. Complexity of energy release during the Imperial Valley, California, earthquake of 1940. *Bull. Seism. Soc. Am.*, 60, 137-160.
- Utsu, T., 1969. Aftershocks and earthquake statistics (I). *J. Fac. Sci., Hokkaido Univ., Japan, Ser. VII*, 3, 129-195.
- Wesson, R. L., and W. L. Ellsworth, 1973. Seismicity preceding moderate earthquake in California. *J. Geophys. Res.*, 78, 8527-8546.
- Wyss, M., and W. H. K. Lee, 1973. Time variation of the average earthquake magnitude in Central California, in: Proceedings of the conference on tectonic problems of the San Andreas fault system, Geological Sciences, XIII, School of Earth Sciences, Stanford Univ., 24-42.

Table 1

Values of the parameters in the
stochastic model

Parameters	CATALOG		
	USGS	CACA	DUDA
γ (b_0)	1.220 + 0.003 (0.86)	1.225 + 0.008 (0.88)	1.270 + 0.011 (1.04)
α_0	375.0 + 1.0	89.5 + 2.5	57.0 + 2.0
γ^0	1.224 + 0.004	1.235 + 0.012	1.263 + 0.012
α	251.6 + 4.6	46.0 + 2.6	46.7 + 2.5
$\Sigma\mu_+$	0.02025	0.02592	0.01119
$\Sigma\mu_-$	0.01834	0.01979	0.02365
c_- (c^{10})	0.936 + 0.023 (0.51)	0.856 + 0.078 (0.21)	1.003 + 0.077 (1.03)
c_+ (c^{T0})	1.169 + 0.006 (4.75)	1.188 + 0.007 (5.66)	1.116 + 0.045 (2.99)
v_-	0.143	0.122	0.114
v_+	0.477	0.575	0.453
N_- (%)	379.5 (6.8%)	44.5 (5.3%)	37.1 (4.2%)
N_+ (%)	1432.1 (25.6%)	363.8 (43.5%)	137.8 (15.4%)
$M_{\max}$	5.2	7.7	8.9
M_{cutoff}	1.5	4.0 (3.5)	7.0
n	5586	837	894
$\ell_{\max} - \ell_0$	4670.8	1665.3	145.4
I/n (bits)	1.21	2.87	0.23

Table 2

Values of Branching Ratio μ

Distance Intervals	Catalog	Foreshock time intervals								
		9	8	7	6	5	4	3	2	1
1	USGS	134+58	67+38	141+40	153+39	154+37	203+41	133+32	78+23	29+14
	CACA	0	10+112	146+136	12+113	179+128	167+125	217+135	0	117+91
	DUDA	—	0	0	0	0	88+65	48+47	47+44	120+74
	average	67	26	96	55	111	153	133	42	89
2	USGS	0	115+51	44+34	150+41	82+32	71+25	51+21	40+16	76+22
	CACA	0	40+155	112+111	38+97	256+167	4+141	64+82	145+112	174+115
	DUDA	—	0	0	0	0	75+61	56+47	82+59	0
	average	0	52	52	63	113	50	57	89	83
3	USGS	14+61	0	44+33	54+28	26+19	35+17	46+20	16+10	0
	CACA	0	0	0	110+116	113+145	103+109	82+116	58+62	0
	DUDA	—	110+312	0	0	0	7+32	20+32	27+43	0
	average	7	37	15	55	46	48	49	34	0
4	USGS	0	26+53	1+31	12+24	0	0	11+10	5+7	14+10
	CACA	0	94+145	4+104	45+150	153+151	0	85+89	64+67	0
	DUDA	—	0	360+337	0	0	0	29+46	24+33	26+32
	average	0	40	122	19	51	0	42	31	13
$\Sigma \mu$	average	74	155	285	192	321	251	281	196	185

All values of μ should be multiplied by 10^{-5}

Table 2 continued
Values of Branching Ratio μ

Distance Intervals	Catalog	Aftershock time intervals								
		9	8	7	6	5	4	3	2	1
1	USGS	62+10	65+10	72+12	85+13	78+13	127+17	103+16	86+18	102+22
	CACA	78+23	111+29	178+39	91+28	154+39	90+29	58+27	57+27	9+29
	DUDA	86+44	63+38	141+61	93+49	122+63	0	0	0	—
	average	75	80	130	90	118	72	54	48	56
2	USGS	32+7	54+10	48+10	81+13	121+17	88+15	108+18	81+21	58+27
	CACA	56+20	101+29	163+39	148+40	77+33	77+31	127+39	34+30	10+40
	DUDA	0	34+32	30+31	76+46	19+48	140+78	0	51+156	—
	average	29	63	80	102	72	102	78	55	34
3	USGS	11+4	28+7	8+4	22+7	25+8	27+10	28+14	52+20	51+30
	CACA	23+13	0	0	15+13	45+21	12+17	58+28	0	0
	DUDA	0	10+17	11+26	12+34	63+58	55+82	2+132	56+233	—
	average	11	13	6	16	44	31	29	36	26
4	USGS	4+2	6+3	6+4	13+5	8+6	9+8	4+13	44+23	37+32
	CACA	0	8+9	26+23	0	19+16	31+24	23+33	0	100+63
	DUDA	9+17	10+18	13+25	14+39	68+73	119+127	158+197	910+379	—
	average	4	8	15	9	32	53	62	318	69
$\Sigma \mu^+$	average	119	164	231	217	266	258	223	457	185

All values of μ should be multiplied by 10^{-5}

Table 3

Percentages of aftershocks in distance
intervals 3 and 4 in USGS

Magnitude interval of main shock, M_1	Magnitude interval of aftershocks, M_2	$\Delta M = M_1 - M_2$	Time Intervals		
			1-3	4-6	7-9
2.5 to 3.4	2.5 to 3.4	0	2.7	19.2	45.9
	1.5 to 2.4	1	19.2	16.3	29.0
3.5 to 4.4	2.5 to 3.4	1	23.9	16.8	35.0
	1.5 to 2.4	2	8.9	14.7	11.0
4.5 to 5.4	2.5 to 3.4	2	18.5	15.7	9.7
	1.5 to 2.4	3	12.0	6.6	2.4

Table 4
 Values of the Coefficient c
 for Catalog USGS

Distance Intervals	<u>Foreshock time intervals</u>		
	<u>9-7</u>	<u>6-4</u>	<u>3-1</u>
1-2	0.722	0.893	0.988
3-4	0.880	0.915	1.092
Distance Intervals	<u>Aftershock time intervals</u>		
	<u>1-3</u>	<u>4-6</u>	<u>7-9</u>
1-2	1.181	1.171	1.188
3-4	1.173	1.141	1.042

Table 5
Simulated California Catalogs

Number of Realization	Number of events				Maximum magni- tude
	Independent	Dependent		Total	
		all	close		
1	318	217	89	535	6.7
2	336	889	382	1225	7.6
3	302	287	123	589	7.1
4	339	221	94	560	6.6
5	341	518	191	859	7.4

Table 6

The parameter values of simulated catalog

Parameters	CATALOG NUMBER					
	1		2		3	
	a	b	a	b	a	b
γ_0	1.285	1.290	1.240	1.235	1.265	1.270
α	33.5	28.0	80.8	55.9	38.0	29.9
γ^0	1.285	1.285	1.213	1.165	1.252	1.251
α	20.0	19.8	19.7	8.84	15.1	14.8
$\Sigma\mu^-$	0.03332	0.02154	0.03699	0.01990	0.04723	0.02974
$\Sigma\mu^+$	0.03856	0.02828	0.04139	0.02356	0.04249	0.03022
c_-	0.852	0.856	0.851	0.855	0.851	0.855
c_+	1.164	1.171	1.174	1.216	1.181	1.208
v_-	0.153	0.101	0.169	0.093	0.216	0.139
v_+	0.976	0.745	1.110	0.805	1.186	0.986
N_-	35.0	18.9	98.8	37.6	56.7	28.1
N_+	183.9	115.8	831.9	671.5	297.9	208.8
N_{ind}	316.1	311.3	294.3	133.9	234.4	229.1
M_{max}	6.7	6.7	7.6	7.6	7.1	7.1
n_{max}	535	446	1225	843	589	466
$\ell_{max} - \ell_0$	828.0	173.8	3568.0	756.3	1163.2	195.7
$I/n(\text{bits})$	2.23	0.56	4.20	1.29	2.85	0.61

a-15 time intervals, b-9 time intervals

List of Captions

- Fig. 1. Map of California, showing the regions spanned by the catalogs: 1. - Northern California.
2. - Southern California.
3. - Central California (USGS).
- Fig. 2. Spatial moment of the earthquake process.
- Fig. 3. Time and distance intervals. The numbers are explained in the text. A is the estimate of signal duration of an earthquake, B is the estimate of rupture duration of an earthquake.
- Fig. 4. Magnitude-time plots of 1971 San Fernando earthquake sequences: a. Actual sequence (Trifunac, 1972). b, c. Monte Carlo simulations.

Nature of Seismic Gaps and Foreshocks

Hiroo Kanamori

Seismological Laboratory

California Institute of Technology

Pasadena, California 91125

In this report we will discuss several problems on seismic gaps from the point of view of identifying ready-to-break gaps.

1. Seismic gaps at subduction zones

Seismic gaps are usually identified as a spatial gap of activity along subduction zones. Whether the identified gap is seismic or aseismic (i.e. permanent gap) has an important bearing on the problem of capturing a gap earthquake in the near future. For example, most gaps to the north of 40°N in the western Pacific region seem to be seismic, but the nature of the gaps to the south is unclear (Figure 1). For the region between 36°N and 37°N , Abe (1977) concluded that there was no major earthquake in this region at least for the past 800 years. The most recent activity in this region is represented by the 1938 Shioya-Oki earthquake sequence. Although the magnitudes of the earthquakes in this sequence are close to 8, they are significantly smaller than those to the north of 40°N .

Further to the south along the Izu-Bonin-Mariana arc, there is no record of large shallow earthquakes. Although earthquake data go back only several decades, historical tsunami records strongly indicate a lack of great shallow earthquakes in this region at least for the past several hundred years.

Several possibilities have been suggested for the lack of great earthquakes: decoupling between the oceanic and continental lithospheres (Kanamori, 1971, 1977), hang-up of subduction caused by buoyant oceanic lithospheres (Vogt, 1973; Kelleher and McCann, 1976) and receding upper-plates (Uyeda and Kanamori, 1978). In order to distinguish seismic gaps from aseismic gaps, it is important to make detailed studies of sea-bottom topography of the subducting

Figure 1a. Seismic fault zones in the northwestern Pacific. The upper figure shows a sequence from 1763 to 1918. Note the missing event in E. The lower figure shows the last sequence (after Fukao and Furumoto, 1978).

Figure 1b. Epicenters of large earthquakes in and around northern Japan and Izu-Bonin trench to the south of 38°N (after Abe, 1977).

lithosphere and the mechanisms of earthquakes in the adjacent regions. The former would provide information on the buoyancy of the oceanic lithosphere, and the latter, the nature of the lithospheric coupling.

It is also important to note that a recent study of Fukao and Furumoto (1977) suggests that the 1958 Kurile islands earthquake (Figure 1) occurred in a region which remained as a gap during the previous cycle of activity. This result indicates that the fracture strength along a subduction zone varies substantially and the repeat time of great earthquakes may vary greatly from place to place even along the same arc.

Another interesting feature related to the spatial transition from seismic to aseismic gap in the north-western Pacific is the foreshock activity. The 1963 and the 1969 Kurile islands earthquakes had pronounced foreshock activity, and the 1973 Nemuro-Oki earthquake had two teleseismically located foreshocks (Figure 2). However, the 1952 and the 1968 Tokachi-Oki earthquakes which are located between the seismic and aseismic arcs did not have pronounced foreshock activity (see Figure 1). This difference probably reflects the difference in the mechanical property of the subduction zone. Since foreshock activity is one of the most definitive precursors, this difference is important for designing an experiment for capturing a large gap earthquake.

Many great earthquakes at subduction zones were preceded by pronounced foreshock activities (e.g. 1963 Kurile islands earthquake, 1965, Rat Island earthquake, 1960 Chilean earthquake) (Mogi, 1969). These foreshock activities are relatively short-term (days and weeks) (see Figure 2b) but it is not clear whether they represent a distinct clustering of events within the individual gap during the inter-seismic period or not. The existing earthquake

Figure 2a. Foreshocks of the 1963 Kurile Islands, the 1969 Kurile Islands, the 1973 Nemuro-Oki earthquake, and the 1965 Rat Islands earthquake.

Figure 2b. Time sequence of the foreshocks of the 1963 Kurile Islands earthquake (a) and the 1965 Rat Islands earthquake (b).

catalogs are too incomplete and inaccurate to resolve this problem.

2. Other seismic gaps

Wesson and Ellsworth (1973) and Kelleher and Savino (1975) discussed seismicity prior to several subduction zone, as well as non-subduction zone earthquakes. Here we will focus on the 1971 San Fernando earthquake and the 1952 Kern County earthquake because these earthquakes have fairly complete earthquake data during the pre-seismic period.

Figures 3 and 4 show the seismicity pattern prior to the 1971 San Fernando earthquake. It appears that a quiet period from 1965 to 1968 (gap) was followed by foreshock activity from 1969 to 1971. This pattern is similar to that found for several other earthquakes. However, the space-time plot (Figure 3) shows that a similar "gap-foreshock" pattern is seen at least three times prior to 1965 (i.e., 1940 to 1949, 1950 to 1956, and 1957 to 1965), and, therefore, the pattern seen immediately before the San Fernando earthquake may not be a unique one. It is unclear whether the gap during the period from 1965 to 1969 has any significance as precursor. However, the foreshocks that occurred during the period from 1969 to 1970 have distinct wave forms (Ishida and Kanamori, 1978a). This feature, together with the seismicity pattern may be a more useful diagnostic pattern of a ready-to-break gap.

Figure 5 shows the seismicity prior to the Kern County earthquake of 1952 (Ishida and Kanamori, 1978b). Figure 5a, which includes a substantial part of the data represents the pattern already suggested by Wesson and Ellsworth (1973) and Kelleher and Savino (1975). A large part of the fault plane seems to have been seismically quiet for nearly 15 years before the Kern County earthquake. During this period the activity near the epicentral area was very high. As shown by Figures 5b and 5c, this activity became very low during the two-year period from

Figure 3. Space-time plot of the seismicity before the 1971 San Fernando earthquake (Ishida and Kanamori, 1977).

Spatial distribution of relocated epicenters

Figure 4. Distribution of epicenters before the 1971 San Fernando earthquake for four periods (Ishida and Kanamori, 1978a)

Figure 5. Distribution of epicenters before the 1952 Kern County earthquake for three periods (Ishida and Kanamori, 1978b).

January 1949 to December 1950. This low activity was followed by a clustering of events near the main shock epicenter which may be considered to be foreshock activity of the Kern County earthquake. This pattern is very similar to that found for the 1971 San Fernando earthquake. However, a space-time plot (Figure 6) indicates that this pattern of "gap -foreshocks" appeared several times before 1950. Thus, it appears that this pattern alone may be of little use as a definitive seismic precursor, unless the events just before the main shock can be distinguished on the basis of another criterion.

One possibility is that this gap corresponds to that seen prior to large subduction zone events (Kelleher and Savino, 1975) and the activity near the epicentral area corresponds to the foreshock activity immediately before large subduction zone events. If this is the case, the foreshock activity before the Kern County earthquake is considered to have lasted for at least 20 years. In this case, the usefulness of foreshock activity for practical earthquake prediction is questionable, unless another criterion is used to distinguish immediate foreshocks from the rest.

3. Discussion

For the purpose of capturing a large earthquake, it is essential to identify a gap which is most likely to break in ten years or so. When a gap in seismic activity is found on the basis of geological data and historical earthquake catalog (e.g., Suruga Bay gap, Oaxaca gap (Ohtake et al., 1977)), it is important to examine whether it is a seismic gap or a permanent gap. Even if the gap is identified as a seismic gap, geological and historical data alone cannot provide information regarding when it is going to break. For

Figure 6. Seismicity in the epicentral area of the 1952 Kern County earthquake as a function of time. Each figure represents seismicity for a consecutive two-year period, except the last figure that represents the period from January 1951 to July 1952 (Ishida and Kanamori, 1978b). Aftershock area (after H. Benioff *et al.*, 1953). White Wolf fault (after B. Gutenberg, 1955).

very large ($M_s \geq 8$) earthquakes, the repeat time is usually longer than 100 years, with very large uncertainty, so that a simple statistical estimate is too uncertain to be useful for identifying a ready-to-break gap. For smaller events ($M_s \sim 7$), the repeat time may be short enough to make statistical approach useful (e.g. the New Hebrides). However, the source size of $M_s \sim 7$ event is about 50 km so that it is more difficult to capture them than $M_s \geq 8$ events whose dimension is 200 km or larger.

Short-term precursors such as anomalous crustal deformation and foreshock activity appear most useful, if they occur at all, for identifying a ready-to-break gap. Although geodetic data are very sparse, foreshock activity has been observed for many gap events. Since the time of occurrence of an earthquake is ultimately controlled by distribution of stress on the fault plane, it is essential to know how spatio-temporal distribution of stress on the fault plane changes. In view of the lack of a simple method for direct determination of stresses at depths, it appears most promising to use spatio-temporal variation of seismicity (including spectra, wave form, mechanism etc.) in seismic gaps and in the surrounding areas for determining whether the gap is soon-to-break or not.

Contribution No. 3131, Division of Geological and Planetary Sciences,
California Institute of Technology, Pasadena, California 91125.

REFERENCES

- Abe, K., Tectonic implications of the large Shioya-Oki earthquakes of 1938, Tectonophysics, 41, 269-289, 1977.
- Fukao, Y., and M. Furumoto, Anomalous stress drop of the great Kurile earthquake of Nov. 6, 1958, EOS, 58, 446, 1977.
- Ishida, M., and H. Kanamori, The spatio-temporal variation of seismicity before the 1971 San Fernando earthquake, California, Geophys. Res. Lett., 4, 345-346, 1977.
- Ishida, M., and H. Kanamori, The foreshock activity of the 1971 San Fernando earthquake, California, Bull. Seismol. Soc. Am., (in press), 1978a.
- Ishida, M., and H. Kanamori, Seismicity before the 1952 Kern County earthquake, California (in preparation), 1978b.
- Kanamori, H., Great earthquakes at island arcs and the lithosphere, Tectonophysics, 12, 187-198, 1971.
- Kanamori, H., Seismic and aseismic slip along subduction zones and their tectonic implications, in Island Arcs, Deep Sea Trenches and Back-Arc Basins, Maurice Ewing Series 1, 163-174, American Geophysical Union, Washington D.C., 1977.
- Kelleher, J., and W. McCann, Buoyant zones, great earthquakes, and unstable boundaries of subduction, J. Geophys. Res., 81, 4885-4896, 1976.
- Kelleher, J., and J. Savino, Distribution of seismicity before large strike slip and thrust-type earthquakes, J. Geophys. Res., 80, 260-271, 1975.
- Mogi, K., Source locations of elastic shocks in the fracturing process in rocks (1), Bull. Earthquake Res. Inst., Tokyo Univ., 46, 1103-1125, 1968b.
- Ohtake, M., Matumoto, T., and G. V. Latham, Seismicity gap near Oaxaca, Southern Mexico as a probable precursor to a large earthquake, Pageoph, 115, 375-385, 1977.

Uyeda, S., and H. Kanamori, Back-arc opening and the mode of subduction,
submitted to J. Geophys. Res., 1978.

Vogt, P.R., Subduction and aseismic ridges, Nature, 241, 189-191, 1973.

Wesson, R., and W. Ellsworth, Seismicity preceding moderate earthquakes in
California, J. Geophys. Res., 78, 8527-8546, 1973.

Keichi KASAHARA,

Earthquake Research Institute, University of Tokyo.

ABSTRACT

This technique assumes lateral migration of a narrow linear zone of seismic activity at a uniform velocity across a seismic area, and identifies groups of earthquakes which satisfy the kinematic criterion of uni-directional migration. If a group of earthquakes occur in the above-stated manner, their distribution in the time - space domain defines a plane surface in the (x, y, t) coordinates. Let Δt denote the time interval of a pair of events, and R and θ , the distance and azimuth from one epicenter to another, respectively; then we have a geometrical relation:

$$\Delta t / R = A_0 \cos (\theta - \theta_0),$$

where θ_0 is the direction to the steepest slope of the plane. Consequently, the $\Delta t / R$ versus θ plot of the seismic data will reveal a cosine trend if a considerable number of events follow the above criterion. For easier recognition of the trend, we may introduce a second diagram, which compares theoretical curves of various sets of A_0 and θ_0 with respect to their score, i.e. the probability that a curve will involve the observational points in its vicinity. A spot (or spots) of high score will be observed in the diagram when migration exists, from which we estimate possible direction and slowness of seismicity migration.

In practice, a program reads a set of seismicity data and displays sets of $\Delta t / R - \theta$ and score diagrams successively, by scanning suspicious ranges of epoch, area, and magnitude, following a preset schedule. Major earthquakes in the Anatolia fault, Turkey and Tohoku District, Honshu, Japan, have been studied in order to explain the application of the present technique to migration studies.

INTRODUCTION

Earthquake foci in a seismic zone sometimes appear to migrate systematically in one direction. The most convincing evidence for this is perhaps the last sequence in the Anatolia fault zone, Turkey, which occurred during the two or three decades following 1939. Richter (1958), and later, Mogi (1968), recognized a clear trend of migration from the eastern to the western extremity of the fault, from which an apparent velocity of about 80 km/yr is estimated. More examples of a similar kind have been discovered in various areas, sometimes on the basis of seismic sequences, but also on the basis of land movements which tend to migrate slowly at velocities 10 - 100 km/yr (see a review paper by Kasahara, 1978). The good agreement between these two migration

velocities may not be accidental. Thus several theoretical models have been proposed for the purpose of explaining the physical mechanisms of excitation and slow migration of such crustal activities (earthquake occurrence and/or anomalous land movements) as mentioned above. The Bott - Dean model (1973) introduces, for example, an elastic plate underlain by a viscous layer, and is applicable to the lithosphere - asthenosphere system responding to periodic stress change at a trench axis. Anderson (1975) has employed a similar layered model of a plate system beyond the trench in order to interpret lateral migration of earthquakes along the trench axis.

Another version of migration studies has been emphasized by successful prediction of the Haicheng earthquake, NE China (1975, $M = 7.3$). As Scholz (1977) has shown in his review, seismic activity and anomalous land movements there tended to migrate northeastward across the Pohai Bay, so that the Chinese seismologists could promptly intensify their prediction work in the area of the coming earthquake. This example suggests that migration of crustal activity may be a useful clue to long-term prediction of earthquakes, on the basis of a proper evaluation of a soon-to-break gap.

Until now, recognition of migration has been done manually, by hypothesizing a possible direction of migration to examine the time - space relation of events to be studied. It is a simple technique, but the initial choice of the migration direction is not easy if the migration is contaminated by random events. The purpose of the present work is to search for a more efficient and objective technique of discrimination of seismicity migration which might lie in a given data file.

TECHNIQUE

Let us suppose, in the x, y - space, a linear front (or stripe) of high seismicity, which migrates with an azimuth θ_0 at a uniform velocity v_0 . Figure 1 compares the seismicity maps at three successive moments, and illustrates that the migrating front defines a plane surface in the x, y, t - coordinate system. This suggests that seismic events responding to migration satisfy the mutual time - space relation

$$\Delta t / R = A_0 \cos (\theta - \theta_0), \quad (1)$$

where Δt denotes the time interval of a pair of events, and R and θ , the distance and azimuth from one epicenter to another, respectively. θ_0 , the direction of migration, represents the direction of steepest slope of the plane, and A_0 , the steepest slope, may be understood as the slowness of migration ($1 / v_0$). Consequently, if we plot the $\Delta t / R$ versus θ relation of all the possible pairs of events which share a common migration plane, then we shall observe a cosine trend in the diagram. This is very similar, in principle, to the technique of Miyabe (1931), which derives the trend of land tilting from leveling data. Figure 2 (top) illustrates the cosine curve which will be revealed in this manner.

The least - squares method may be used conveniently for this purpose, if we know, a priori, that the given file of seismic data bears a single mode of migration. If this is unknown, as is mostly the case,

we must use an alternative method to deal with plural modes of migration in an area. Figure 2 (bottom) illustrates the diagram for this purpose, which represents a score of the respective set of the parameters (A_0, θ_0) as an array. Suppose, in the first diagram, that a theoretical curve $A_0 \cos(\theta - \theta_0)$ involves $n(A_0, \theta_0)$ observational points in its vicinity, i.e. in a zone bounded by two parallel curves, $A_0 \cos(\theta - \theta_0) + \Delta A$ and $A_0 \cos(\theta - \theta_0) - \Delta A$. Then, the points (A_0, θ_0) in the second diagram may be assigned the score $n(A_0, \theta_0) / n_0$, where n_0 denotes the total number of points plotted in the first diagram. Practically, we take discrete values of A_0 and θ_0 and assign all the grid points (A_0, θ_0) in the second diagram with their respective score. A spot (or spots) of high score will be observed where migration exists, from which we may estimate the possible direction and slowness of the events under consideration.

EXAMPLES

Let us first take the seismic sequence in the Anatolia fault zone to test the present technique. Various authors have studied this region from the seismological and tectonic viewpoints (e.g. Richter, 1958; Mogi, 1968; Allen, 1969). Recently, Toksöz, Shakal and Michael (1978) have examined the existing earthquake catalogs of this region and have studied its seismic history, migration and seismic gap problems. The following test will refer to their catalog of earthquakes (see Table 1 of their paper).

A computer program reads the earthquake catalog, including the time, location and magnitude of earthquakes in the list, and calculates Δt , R and θ for all possible pairs of events (those pairs with $\Delta t > 10$ yr are excluded). The output is an array of frequency distribution of pairs, with respect to their slowness $(\Delta t / R)$ and azimuth θ , as illustrated in Figure 3 (bottom). The top part of the figure displays the slowness - azimuth relation for magnitude $M \geq 6.5$. Scarcity of data for azimuths excluding W and E is attributed to linear direction of epicenters along the fault zone, yet we observe, in the figure, small numbers of clusters of the plotted points, with various slowness about the azimuths W and E. Broken-line cosine curves, which are adopted in reference to the next figure, may interpret the present data, although we do not have good control of azimuth due to the above-stated condition.

The top and bottom parts of Figure 4 correspond respectively to the previous pictures, and display the score arrays as explained in the previous section. The zone width $\pm \Delta A$ is taken as ± 0.00222 yr/km, i.e. $1/30$ of the full scale of the ordinate. Several areas of high score are recognizable; but they are characterized by concave patterns in the diagram. This is again attributed to the epicenter distribution being restricted to the fault zone, and suggests that the present data may specify only an apparent velocity along the extension of the seismic zone. The apices of the concave patterns represent this velocity in the respective migration group. They are,

	direction	velocity (km/yr)
trend(predominant)	southwest	25
(faint)	west	80 25
	east	70 50 20

Toksöz et al. (1978) have recognized two modes of migration along the Anatolia fault zone, indicating average velocities of about 50 km/yr to the west and 10 km/yr to the east, for a 38 year period since 1939. The above-stated results seem to correspond to these velocities, respectively.

Figures 5 and 6 illustrate another example of a test for seismic sequence in the Tohoku district, Honshu, Japan. A catalog of disastrous earthquakes in the Japan area, 1872 - 1977 (Rika-Nenpyo, 1978) is used in order to examine migrational features in the successive periods since 1872. The previous program has been improved for multiple run processing. That is to say, it first reads a batch of seismicity data and repeat the process successively, by scanning suspicious ranges of epoch, area, and magnitude, following a preset schedule. In the present case, for example, tests are worked out for two magnitude ranges ($M \geq 5.5$ and ≥ 6.5), and for four epochs (1880 - 1920, 1900 - 1940, 1920 - 1960, 1940 - 1980), regarding the area, 37 - 42° N, 138 - 143° E, as shown in Figure 6. Let us study the score diagrams in Figure 5 (the slowness - azimuth diagrams are not shown). We may recognize, in the top figure (a), predominance of a northeastward migration for the period 1880 - 1920, but this mode seems to be replaced by a westward one in the later periods (figures (b) - (d)).

The southward migration of large offshore earthquakes in the period 1931 - 1938 has been discovered by Mogi (1969), and more recently, Ishii (1976) and Tanaka (1978) have respectively recognized the northwestward (or westward) migration of crustal deformation and seismic activity in the same district. Therefore, some of the present findings may not be new, in themselves. More notable, however, is the fact that the present diagram enables us to observe a comprehensive picture of migration in the area, from which we learn that various modes of migration have appeared and disappeared in a transient manner. For example, the northward (or northwestward) mode is recognized in (a) and (b), but not in the later two pictures. The southward mode, which is invisible in (a), appears slightly in (b) and more obviously in (c), then disappears in (d). On the other hand, the northwestward one becomes predominant after period (c). The westward mode seems most recent, appearing only in the last period. Figure 6 summarizes transient features of these migrational modes, together with their minimum possible velocities as estimated from the apex points of the respective concave patterns.

CONCLUDING REMARKS

The basic function of this technique is to identify a group (or groups) of earthquakes which answer to the kinematic criterion as previously proposed. In other words, it is concerned with the necessary conditions for migration, but may not necessarily satisfy the sufficient conditions. Perhaps its findings are only preliminary and must undergo further statistical and seismological examination for more definite recognition of migration. In spite of these characteristics, however, this technique will be a useful tool, especially for quick and sensitive discrimination of migration modes which might exist in a large amount of seismicity data. The present paper has dealt with horizontal migration of shallow earthquakes, migration of deep earthquakes (if it exists) may be studied equally well,

when a tilted seismic plane is recognizable.

As the case of the 1975 Haicheng earthquake has proved, migration of crustal activity, like local seismicity and crustal deformation, may sometimes be a promising working hypothesis for earthquake prediction. If our future routine system could identify migrational components in an area and monitor their behavior continuously, then it would certainly be advantageous for a better understanding of the seismic triggering mechanism and for proper evaluation of soon-to-break gaps in the area of interest. The present method might play a role in this kind of data processing system.

I acknowledge with thanks the U. S. Geological Survey and Dr. Jack F. Evernden, Convenor of the Conference, for their kind invitation to the meeting. Dr. Paul Somerville reviewed the manuscript, to whom I express my thanks, too.

REFERENCES

- Allen, C. R., Active faulting in northern Turkey, Contr. No. 1577, Div. Geol. Sci., Calif. Inst. Tech., 32, 1969.
- Anderson, D. L., Accelerated plate tectonics, Science, 187, 1077 - 1079, 1975.
- Bott, M. H. P. and Deans, D. S., Stress diffusion from plate boundaries, Nature, 243, 339 - 341, 1973.
- Ishii, H., Characteristics of crustal movement observed at wide area, Symposium on Earthquake Prediction, Proceedings, 116 - 126, 1976 (in Japanese with English abstract).
- Kasahara, K., Migration of crustal deformation, 1977 RCM Symposium, Tectonophysics, 1978 (in press).
- Miyabe, N., Blocks in the Earth's crust and their movements, Part I, Bull. Earthq. Res. Inst., 9, 256 - 270, 1931.
- Mogi, K., Migration of seismic activity, Bull. Earthq. Res. Inst., Tokyo Univ., 46, 53 - 74, 1968.
- Mogi, K., Some features of recent seismic activity in and near Japan, (2) Activity before and after great earthquakes, Bull. Earthq. Res. Inst., 47, 395 - 417, 1969.
- Richter, C. F., Elementary Seismology, Freeman and Cooper, San Francisco, pp. 611 - 616, 1958.
- Rika-Nenpyo (Annual Table of Scientific Constants*) — 1978 —, Maruzen, Tokyo, 1977 (in Japanese).
- Scholz, C. H., A physical interpretation of the Haicheng earthquake prediction, Nature, 267, 121 - 124, 1977.
- Tanaka, K., Migration of seismic and volcanic activity*, Abstracts, Ann. Meeting, Seismol. Soc. Japan, No. 1, 1978 (in Japanese).
- Toksöz, M. N., Shakal, A. F. and Michael, A. J., Space-time migration of earthquakes along the North Anatolian fault zone and seismic gaps, Methodology for Defining Seismic Gaps and Soon-To-Break Gaps, Open-File Report, USGS, No. , 1978 (in press).

*) provisional translation by the present author.

FIGURE CAPTIONS

- Figure 1 A plane surface in the time - space domain, as defined by uni-directional migration of seismicity.
- Figure 2 (top): a slowness-azimuth diagram of seismicity data, in which a cosine trend represents a mode of migration, and (bottom): a score diagram, which displays, as an array, scores of the fit of the respective sets of parameters (A_0 , θ_0) to the observational data.
- Figure 3 Migration of seismicity in the Anatolia fault zone, Turkey (1930 - 1970), as seen in slowness - azimuth diagrams. (top): $M \geq 6.5$, (bottom): $M \geq 5.5$. Thick and thin broken lines represent the predominant and faint trends of migration as inferred from the next figure.
SYMBOL	.	:	;	/	x	+	=	⊕	%
FREQUENCY (N)	1	2	3	4	5	6	7	8	9 OR MORE
- Figure 4 Migration of seismicity in the Anatolia fault zone, Turkey (1930 - 1970), as seen in score diagrams. (top): $M \geq 6.5$, (bottom): $M \geq 5.5$.
- Figure 5 Migration of seismicity in the Tohoku district, Japan, as seen in score diagrams ($M \geq 6.5$). (a) 1880 -1920, (b) 1900 - 1940, (c) 1920 -1960, (d) 1940 -1980.
- Figure 6 Possible modes of seismicity migration in the Tohoku district, Japan, in the successive four periods ($M \geq 6.5$). Velocities represent the lowest possible values in the respective mode.

Figure 2

Figure 1

Figure 3

$M \geq 6.5$

3 SLOWNESS (YR / KM)

Figure 3 (continued)

ANATOLIA, TURKEY

(1930 - 1970)

Figure 4

Figure 5 (a)

Figure 5 (b)

TOHOKU DISTRICT, JAPAN

(1920 - 1960)

C

Figure 5 (c)

Figure 6

BURSTS OF SEISMICITY AS LONG-TERM PRECURSORS
OF STRONG EARTHQUAKES

V. I. Keilis-Borok,^{*} L. Knopoff,^{**} I. M. Rotvain,^{*}
and T. M. Sidorenko^{*}

^{*}Institute of the Physics of the Earth, Academy of Sciences, Moscow, USSR.

^{**}Institute of Geophysics and Planetary Physics, University of California at
Los Angeles, California.

An edited version of this paper is being submitted to the Journal of Geophysical Research. Please make reference to JGR, submitted.

ABSTRACT

Three seismicity patterns premonitory to strong earthquakes are described and analyzed. These are: (1) occurrence of an earthquake of medium magnitude with an abnormally large number of aftershocks at the beginning of the aftershock sequence (pattern "B," or "bursts of aftershocks"); (2) a cluster of earthquakes at a time when seismicity is not below average (pattern "S," or "swarms"); (3) clusters of earthquakes in the time-energy domain (pattern Σ , or "sigma," represented by the sum of energies to the power $2/3$).

These patterns, after formalization, are successfully tested by retrospective prediction of the strongest earthquakes in Southern California and several other regions. The first pattern is introduced in this paper; the two others were formalized by Caputo et al., 1978; Keilis-Borok and Rotvain, 1978; and Keilis-Borok and Malinovskaya, 1964.

The three patterns seem to be different projections of a general pattern of bursts of seismicity, i.e., abnormal clustering of earthquakes in time, space, and energy. These patterns can be explained by, and inferred from, the model of earthquake sequences described by Knopoff et al., 1969, 1972, 1976, and 1978.

Prologue

The first part of this paper, which is concerned with the construction of a model, came after much of the data processing that we describe in the second part of this paper had been performed. Many of the physical relationships that are to be described had been indicated by the earthquake catalogs first. The introduction of a possible model to explain the observations has been inserted at the beginning of this paper as a matter of convenience and certain amount of a posteriori reasoning was used.

Introduction

The logic of a search for premonitory seismicity patterns to identify precursors of large earthquakes is based on certain simple assumptions which take the form of platitudes.

- 1) The presence of causally connected earthquakes in a catalog implies that the statistics of the earthquakes in the catalog do not exhibit a Poissonian distribution.
- 2) The study of very large earthquakes in a certain region leads us to a listing of such a small number of large events that statistical methods probably do not apply.

The existence of precursors to earthquakes implies that triggering relationships may exist. If conditions exist leading to an eventual earthquake, then we may suppose these processes or conditions may be accelerated by the proper agent. We search to identify such patterns in earthquakes sequence, which represent triggering events of other, larger, earthquakes. With regard to the first of the assumptions above, we note the additional platitude that triggering relationships among earthquakes are all too abundant: aftershocks, in the conventional sense, are most obviously events that are triggered by other earthquakes. We have not yet defined an aftershock; any rigorous attempt to do so (i.e. without arbitrary assumptions involving parameterization) must ultimately force us to describe the large events being triggered as being themselves aftershocks of other events; in short, the definition cannot accomodate differences between aftershocks and foreshocks. Hence we may appeal to conventionality in our definition; put in still other terms, we indicate that we will search for triggering relationships in certain convenient intervals of time. This is a form of parameterization.

Let us suppose an aftershock can be defined by parameterization, even if the parameterization is arbitrary. If these aftershocks are removed from the catalog, the residual catalog is found to be Poissonian in character (Gardner and Knopoff, 1974). What this means is that any identifiable triggering of large, relatively infrequent earthquakes, if it is present at all, is a small effect and is likely to be lost by embedding in a larger catalog from which aftershocks have been removed; we call such a catalog, a residual catalog.

Kagan and Knopoff (1976, 1977, 1978) have devised one method of increasing the sensitivity of the search within a mostly Poissonian residual catalog to try to identify causally connected events preceding large earthquakes. Indeed, the method can be applied to complete catalogs and need not be applied to residual catalogs only. Kagan and Knopoff have computed the second moment of the occurrence of earthquakes in a catalog in which the number of earthquakes occurring within a given magnitude interval, time interval and distance interval, of any event in the full catalog is tabulated. In their study, as could have been anticipated, the most obvious feature of the second moment of the catalog is the aftershocks which are clustered at short distance intervals and short time intervals. But there are several other features in their work that show considerable promise for our task. These features, albeit much weaker than that of aftershocks, are still statistically significant. (This latter remark implies that there is an abundant enough catalog to test the statistical significance). Among these features are

- 1) The identification of aftershock "migration." It is found that, if a large event occurred, a later weaker event, can occur in a relatively well-defined more-or-less linear time-distance band.
- 2) Foreshocks in the range of $7.5 < M \leq 8.0$ are strongly and abundantly suppressed on a time scale of about 8 years before a large earthquake of magnitude $M > 8.0$ and within a distance of about 2,000 km from the epicenter of the subsequent large event. Such a rarefaction is called a "seismic gap."
- 3) Weaker foreshocks also exhibit a "migration" toward the epicenter of a subsequent larger event. Once again, the word migration is used to define a pair relationship in which the earlier, weaker event occurs in a relatively well-defined, more-or-less linear time-distance band.

It is the latter feature that interests us in this paper since this is evidence for a weak, but statistically significant triggering of large earthquakes by other slightly smaller earthquakes. In this paper we present the results of an exploration of a hypothesis that such triggering effects are non-randomly distributed over the entire globe. If our hypothesis is correct, then in certain areas of the world, the effect of foreshock migration should be enhanced, and in other regions, the effect should be suppressed, perhaps even into the level of insignificance. If the accentuated effect can be identified and is sufficiently strong, then this may become a useful precursor for the forecasting of strong earthquakes. Here we face the ominous barrier of the second assumption. If we reduce the scope of our investigations to a smaller geographic scale, we lose the advantages of testing our conjectures by statistical methods, since the number of strong earthquakes occurring in the time span of modern, well-documented catalogs is generally small for most regions of the world on the distance scale of a few hundreds of kilometers. Because of this, we can only construct our model as conjecture and display the results. But we cannot substantiate these results, if any, on statistical grounds; a similar situation has arisen in the case of prediction of locations of epicenters of large earthquakes (Gelfand, et al, 1976).

I. Physical Influences on Statistical Models

To improve our chances of success, we have added an abstraction of a physical model to our processing procedures. Let us suppose that the triggered event, i.e., the subsequent stronger event to be predicted, is about to take place whether triggered or not, in a locality that is "prepared." That is to say, the event is likely to occur in a time, let us say, of the order of a few years in any case. Then, if we invoke a stress redistribution model for a distant earthquake, which is our triggering event, the stress level at the site of the forthcoming triggered event may be raised by an amount which is sufficient to cause the latter event to occur somewhat prematurely.

The increase in stress at distance due to a triggering event is likely to be weak. The stress redistribution following an earthquake is of the order of $\sigma(L/r)^2$ at large distances where σ is the average stress drop on the ruptured segment, $2L$ is a characteristic fault dimension such as the fault length, and r is the distance from the nearest edge of the crack (Starr, 1928; Knopoff, 1958;

Keilis-Borok, 1959). Dynamical faulting models yield the same result (Burridge and Halliday, 1971). If we take the triggering event to have magnitude 6.0 for example, then L is about 10 km (King and Knopoff, 1968); let us take σ to be about 100 bars. Then at a distance of 100 km, the increase in stress is only of the order of 1 bar due to the remote earthquake. If the rheological properties of the rocks are those of a material which supports pre-fracture creep (Griggs, 1940), an earthquake, which would normally occur after a significant delay due to creep processes, would now be likely to occur after a somewhat shorter interval. The effect is small, but may be significant; in any case we focus our attention on events of intermediate size as possible triggering events of larger earthquakes.

In the procedure we describe below, we have chosen to study as precursors, only those intermediate magnitude events that have an abundance of aftershocks occurring within a very short time of the main event, and to reject, for purposes of analysis, those events that are not followed by many aftershocks. By a certain amount of a posteriori reasoning, we are led to the following steps toward this additional feature of our model.

We assume that both of the possible models for dynamic fracture, that are described in the literature, can occur in nature. Burridge and Halliday (1971) have described a model of rupture and extension without any influence of cohesive forces on the dynamical history of extension and healing of the crack. In their model, the fault reaches its maximum length due to extension into a region where the dynamical stress drop, which drives the fault extension in its earlier stages, becomes negative and hence causes slowing of the rate of extension. Ultimately it causes the crack to stop tearing. If, for the purposes of discussion, the dynamical frictional shear stress were zero, this would mean that the crack had extended into a region where the pre-stress has reversed sign. One of the consequences of this model is that there is no major stress concentration near the edges of the fault. In this region, stress varies $\sigma(r/L)^{1/2}$ just beyond the edges of the crack. The far-field stresses are $\sigma(L/r)^2$ as described above.

The alternative model is that in which the fault tears under the influence of cohesive forces at the tips of the crack (Knopoff, et al., 1973). In this case, the crack extends at a velocity which is moderated by the cohesive forces; in general, the stronger the cohesion, the slower

the rate of extension. Indeed, if the cohesion is too great, the crack stops. After suturing of the moving crack stress concentrations are found at the edges of the crack which vary as $\sigma(r/L)^{1/2}$, just as in the static solutions. The far-field is as in the other case.

There are two related features in which the two models differ. First, there is the presence of an extremely large stress concentration at the edges of the torn segment of the crack in the case of tearing in the presence of cohesive forces; this stress concentration is absent in the cohesion-free model. It is very likely that relaxation of these large stresses is associated with the origin of aftershocks and post-shock creep processes. Second, the model of extension under the influence of cohesion, stop tearing because it encounters an "unbreakable" obstacle, i.e., one with an excessive cohesion; in this case the dynamical stress drop near the edge of the crack need not be negative. In the cohesion-free case, the crack stops because it has entered a region of negative stress drop. It is not possible for earthquakes to originate in regions where negative stress drops are present. Thus, in the immediate neighborhood of earthquakes which cause aftershocks, the stress drop favors the occurrence of other earthquakes; if aftershocks are absent, the possibility of having a second large earthquake in the neighborhood of the first is small. Thus, regions that are "prepared" over a broad area, at least favor the possibility of triggering of large events by other earthquakes associated with aftershocks and scaled at moderate distances.

II. The Burst Hypothesis (B)

Earthquakes with abnormally large number of aftershocks, concentrated at the beginning of aftershocks sequence, is the pattern premonitory to strong earthquake. It will be referred to as pattern B.

We now proceed to the analysis of catalogs under the qualitative constraints of the above model. Let us suppose that we wish to study seismicity patterns premonitory to earthquakes with magnitudes $M \geq M_0$. Such earthquakes will be called strong. In the first example we have used below, namely that of earthquakes in Southern California we have set $M_0 = 6.5$. There are five independent events with $M \leq 6.5$ in the Southern California catalog 1932-1978. As indicated, we study the earthquakes with magnitude less than M_0 as possible triggering events; these are presumed to be relatively strong shocks in the next lower band of magnitudes. It is our assumption that these can be

identified in some way as precursors of strong earthquakes. Earthquakes with $M_1 \leq M < M_0$ are called main shocks. In the Southern California example we restrict main shocks to be in the band $5.4 \leq M < 6.5$. Shocks in this magnitude band that are aftershocks of strong earthquakes are those specifically excluded from the class of main shocks.

We now proceed to the definition of aftershocks, which is necessary for two reasons. First we wish to delete some shocks from the class of main shocks, even though they are in the correct magnitude range, due to the circumstances above. Second, we shall count aftershocks in order to define those main events which have large numbers of aftershocks associated with them.

To avoid bias in the counting process due to inhomogeneity in the catalog, we truncate the catalog at some lower magnitude threshold m , rejecting all shocks with magnitudes less than m . This is done for several reasons including 1) a possible bias against reporting small and moderate sized earthquakes immediately after a large event, 2) the inclusion of earthquake sequences in a catalog which have been specially studied by the laboratory compiling the catalog, including microearthquake studies, and especially 3) a bias against inadequate coverage of small earthquakes due especially to the discrete character of any seismic network. Knopoff and Gardner (1969) have given an empirical procedure for determining the lower-magnitude threshold of homogeneity of the catalog. For Southern California it has been found that 1934 to 1952 to, $m \approx 3.8$; from 1952-1967, $m \approx 2.8$. For the Southern California catalog 1932-1978 that we have processed, we take $m=3.8$.

An aftershock is defined as an event occurring within a time interval $\Delta T(M)$ after a main shock and within the spatial distance $\Delta \phi(M)$, $\Delta \lambda(M)$ of the epicenter (ϕ_i, λ_i) of the i th main shock. For Southern California earthquakes we need not assume that a depth window $\Delta h(m)$ need be applied. Examples of catalog-filtering windows of this type have been given by Knopoff and Gardner (1972) and Gardner and Knopoff (1974).

As our measure of the propensity of an earthquake to the cause of aftershocks, we count the number of aftershocks occurring within a time e following a main shock. In the example of Southern California, we have chosen $e=2$ days. We let b_i be the number of aftershocks in the window of duration e days following the i th main shock.

We now define a threshold $\bar{b}(t)$. If $b_i > \bar{b}$, then we declare that a main shock has occurred associated with a burst of aftershocks. An alarm is declared in that case. The alarm is cancelled after the first strong earthquake in the region or after τ years, whichever comes first. In the Southern California case we take $\tau=2.5$ years.

Several definitions of the threshold $\bar{b}$ are possible. For example let us define $b(t)$ to be the maximum value of b_i in a sliding time window of width s , i.e., $b(t)$ is the largest value of b_i in the interval from $t-s$ to t . Then we may take $\bar{b}(t)$ to be the average value of $b(t)$ over the interval from the start of the catalog to time t ; or we may take $\bar{b}(t)$ to be the average value of $b(t)$ in the interval from $t-D$ to t . Or we may take $\bar{b}$ to be a constant proportional to the long term average of $b(t)$.

Another measure of the strength of a burst of aftershocks could be the parameter B_1 in the Omori Law of aftershock occurrence

$$\dot{n}(t) = B_1/t$$

where $\dot{n}(t)$ is the number of rate of occurrence of aftershocks at a time t after the main shock. According to our hypothesis of bursts of aftershocks, if a value of B_1 is found which falls above a certain threshold, we assume that such an event is a precursor of a strong earthquake. If the number of aftershocks after a main shock is large enough, this measure may be preferable to the preceding one. We have not used this measure of the presence of aftershock bursts in present analysis.

III. Analysis of the Southern California Catalog

To test the hypothesis that bursts of aftershocks of main events are precursors of strong shocks, we first make some arbitrary decisions regarding the values of the thresholds and windows M_0 , m_1 , m , e , ΔT , $\Delta\lambda$, $\Delta\phi$, s , etc. There are so many adjustable parameters, that we run the risk of self-deception by having a large number of degrees of freedom, and it is difficult to perform the job of data fitting honestly. Furthermore, since most of the catalogs are numerically dominated by the smallest events, it is clear that only the events with magnitudes closest to m , i.e., to the effective lower detection magnitude sensitivity of the network, will contribute significantly to the tabulation. Below we will provide a modification of our

hypothesis which gives a stronger weighting factor to larger aftershocks of main events.

We have processed our updating of the Southern California catalog 1932-1978 (Hileman, et al., 1973; Friedman et al., 1976) using parameters M_0 , M_1 , m as above. We used the filter-window algorithm of Gardner and Knopoff (1974) to define aftershocks (Table I). We used also other definition of aftershocks, to show that our results are not strongly sensitive to the choice of windowing algorithm over a rather wide range of variation. We chose $\theta=2$ days. The window τ was chosen to be 2.5 years, a posteriori. We have set the threshold for identification of strong main shocks with bursts of aftershocks as $M_1=5.4$, $\bar{b}=13$ aftershocks.

The results we have obtained are shown in Fig. 1. We have plotted the numbers of aftershocks within 2 days following a precursory main event with $5.4 \leq M \leq 6.4$ as black dots in the figure. We have not plotted the numbers of aftershocks of strong events, $M \geq 6.5$. The times of the strong events are shown with vertical strokes. The windows associated with the precursors are shown as horizontal bars. Of the five strong earthquakes ($M \geq 6.5$) in the Southern California geographic area, whose times are indicated in the figure, four were "predicted" by pattern B. We failed to "predict" one earthquake with $M=6.7$ occurring on 19 May, 1940; that is to say this earthquake was not preceded by any event within 2.5 years before that had more than 13 aftershocks within two days and had a magnitude within the range of M_1 . One "false alarm" is associated with the main events of 14 February 1956 ($M=6.3$) and 15 February 1956 ($M=6.4$).

Another false alarm would appear to be associated with the event 20 March 1969 ($M=5.7$). This event forecasts a strong earthquake within 2.5 years. In point of fact, we may identify the San Fernando earthquake ($M=6.4$) as one which occurred within the specified time window. However, the magnitude of this event is 0.1 too small to fit the threshold of M_0 . If we were prepared to lower the threshold of M_0 to 6.3, we would have been able to "predict" the San Fernando event, but in this case we must add the Borrego Mountain earthquake (9 April 1968, $M=6.4$) to the list of strong earthquakes to be predicted. The Borrego Mountain event was not preceded by pattern B. Furthermore, if M_0 is now lowered, then the San Fernando earthquake generated a false alarm, since it was not followed in the requisite 2.5 years by a strong earthquake. The Borrego Mountain earthquake is also an acceptable precursor under

hypothesis B but the window for this event is kept open by the succession of the two later events, each with magnitude acceptable in the range of M_1 . Thus our score of success for the sequence of three earthquakes spanning the Borrego Mountain and San Fernando events, is one failure to predict, if $M_0=6.5$. If we lower the threshold for M_0 to 6.4, then our success score for this triplet of events is one failure to predict and one successful prediction. If, however, the magnitude of the San Fernando earthquake were to have been underestimated by 0.1 magnitude or more, then our success score becomes one successful prediction. This discussion is given here to show how dependent our assessment is on a digitization process for magnitudes, which of itself must have large errors associated with it. Vanity causes us to count the San Fernando event as a successful prediction under the assumption that the magnitude is greater than 6.4; in this case, it was predicted by the Borrego Mountain and the other precursory event. This discussion illustrates the weak point of our algorithm, which is excessive discretization.

If we were to lower the threshold of $\bar{b}$ from 13 to 10, then two additional events become precursors. (If we lower the threshold still more, we get a number of additional precursory main shocks which are imbedded as dots in the shaded region of the bottom of Figure 1). In the case $\bar{b}=10$, we find that neither of the two new main events predicts a strong succeeding earthquake within its allocated time window. However, the situation is not quite as bleak as this implies. The precursor of 1 July 1941 ($M=5.9$) actually was followed, within schedule, by an event with $M=6.5$. (21 October 1942). The aftershock window of Appendix I has identified the latter event as a late aftershock of the strong event with $M=6.7$ (19 May 1940), itself not predicted. If we were to redesign the later part of the aftershock window algorithm of Appendix I, we will then have predicted the event with $M=6.5$, but we should continue to keep in mind that we fail to predict if $\bar{b}=13$.

The second main event that is introduced if we lower the threshold for $\bar{b}$ (1 December 1958, $M=5.8$) does no damage to our score since it occurred within our alarm period and it serves merely to extend the alarm period. Once again, our vanity suggests that we should choose $\bar{b}=13$ and identify the $M=6.5$ event of 1942 as an aftershock. Once again, we are the victims of a scheme which depends on the use of digitized magnitudes, measured with imprecision, rather than some more precise method of measures such as moment.

Alarms were sounded for 13.7 years or 34% of the 40

year period of the catalog. Let us assume the alarm, ended by the San Fernando earthquake was not a false alarm. Since the algorithm predicted 6 events for which the windows were potentially open for 15 years, the probability of capturing 5 events and failing to capture a sixth, or capturing all six events, by a random binomial process would be

$$6 \cdot \left(\frac{15}{40}\right)^5 \left(\frac{25}{40}\right) + \left(\frac{25}{40}\right)^6 = 0.03$$

Therefore, this method has only a 3% chance of being a random predictor. We must be careful here since we do not know how many degrees of freedom have been lost due to the introduction of a posteriori constraints. Only the test of pattern B with the same values of parameters on other regions, and/or future can overcome this difficulty. Furthermore, we have not evaluated the effects of keeping windows open beyond the 2.5 year period due to a succession of precursors. Evaluations for other numerical cations of numbers of successes, failures and false alarms are calculable similarly.

We have made an alternative assessment of the same catalog using different threshold criteria. In the second exercise, we have reduced m to zero, i.e., we have used all earthquakes as listed, without regard to threshold magnitude homogeneity of the catalog. The results are as shown in Fig. 2. We have used different definition of $\bar{b}(t)$, the new function is indicated by the solid curve. In this case, we have calculated $\bar{b}$ by listing the number of aftershocks occurring in two days that occur in a two day interval following all main shocks $5.4 \leq M \leq 6.4$ and have selected the largest number from this list in any given year. The list of annual maximum numbers of aftershocks was then averaged over time by a running average from the start of the catalog. The Long Beach earthquake of 1933 ($M=6.3$) had 102 aftershocks in a two day interval and this leads to the large spike in $\bar{b}(t)$ at the beginning of the time interval. The curve of $\bar{b}(t)$ stabilizes after a long time near a value of about 8.

In this case aftershocks were defined by a magnitude-time-distance window which is listed in Table II. By the use of this window and the definition of $\bar{b}(t)$, 12 events in the magnitude range $5.4 \leq M \leq 6.4$ had aftershocks that were above threshold $\bar{b}(t)$. The event of 21 October 1942 ($M=6.5$), identified as a late large aftershock in our first study, now emerges as an independent strong event. Neither this event nor that of its parent in the first study (19 May 1940, $M=6.7$) are now "predicted" by this algorithm, in which a

15

window is opened for 2.5 years following the precursor. The remaining four strong earthquakes are predicted, as before. There is one false alarm as before.

With regard to the Borrego Mountain-San Fernando pair, there are four precursors, including these two events themselves. None of these "precursors" was followed by a shock with $M \geq M_0 = 6.5$. Since these events occur in sufficiently close succession to keep the alarm window open, this sequence must be counted as generating one false alarm. If however we lower the threshold to $M_0 = 6.4$, both Borrego Mountain and San Fernando events are successfully predicted. Our score, for this sequence is therefore

one false alarm if $M_0 = 6.5$,
two successful predictions if $M_0 = 6.4$,
one successful prediction if $M_0 = 6.5$ and the magnitude of the San Fernando event is 6.5 or greater.

We have already discussed above the perils of discrete digitization of both thresholds and magnitudes.

We have noted that the Long Beach earthquake of 1933 keeps the threshold $\bar{b}$ rather higher than its asymptote for about 20 years due to the method of providing a running average from the start of the catalog. This is an initial transient; later strong events such as 1969 and 1971 did not produce such a dramatic effect. If we lower $\bar{b}$ to a value close to its asymptotic value of 8, then one additional main shock (1941) emerges above threshold and becomes a predictor for the strong earthquake of 21 October, 1942 ($M = 6.5$).

A curiosity is the main event with 8 aftershocks which occurred on 18 May 1940. This number of aftershocks is at the asymptotic threshold. This event preceded the strong earthquake with $M = 7.1$ by only 23 hours. Due to the distance between the shocks of about 160 km the weaker event cannot be considered to be a local foreshock in the conventional sense.

The threshold number of aftershocks was not attained within 23 hours. This earthquake is therefore not a precursor to the $M = 7.1$ event. Whether this is to be counted as a precursor or not depends on a) whether we use a literal interpretation of the threshold and b) whether the model accommodates the maintenance of a regional alarm after a strong earthquake has occurred. By our definition the main event is not a precursor to the $M = 7.1$ event.

With a uniform threshold of $\bar{b}=8$, with $M_0=6.4$ our score in this variation is 7 successes, 1 failure to predict and 1 false alarm.

IV. The Swarm Hypothesis (S)

A second premonitory pattern, S, is a variation of the first. (Caputo et al. 1978; Keilis-Borok and Rotvain, 1978; Evison, N.F. 1976). A swarm is defined as a group of earthquakes concentrated in time and in space occurring at a time when the seismicity of the region is not below average. In this case we relax the restriction of activity within two days and instead analyze seismic activity over a longer time span. In the example we shall use a year long span.

We make the following definitions:

- 1) Let $N(t)$ be the total number of earthquakes with $M \geq m$ occurring within the time interval from $t-s$ to t and within an entire region, such as Southern California. The count includes aftershocks of strong earthquakes.
- 2) Let $\bar{N}(t)$ be the running average of $N(t)$ using a time window which either starts at the beginning of the catalog or extends from $t-D$ to t .
- 3) Let $n(t)$ be the number of earthquakes with $M \geq m$ occurring within the time interval extending from $t-s$ to t and within the same entire region, but in this time excluding aftershocks of strong earthquakes.
- 4) $r(t)$ which is the maximal number of earthquakes with $M \geq m$ in the rectangle within the sliding window extending from $t-s$ to t rectangle slides over the map of epicenters.

The magnitude m is some lower cutoff magnitude which we may take as the threshold of homogeneity

Under the swarm hypothesis, an alarm is declared if

$$n(t) > C_1 \bar{N}(t)$$

and

$$r(t) > C_2 n(t) \text{ for any rectangle}$$

where

C_1 and C_2 are some constants.

The results for Southern California are shown in Figure 3. The cutoff magnitude has been chosen to be $M=3.8$ as before. The curve of $n(t)$ is computed using a one-year window;

adjacent points are overlapping evaluations taken 6 months apart. The sizes of the rectangles may be taken formally as fixed, e.g. 40 x 40 km; in practice we find we may use the window algorithm of Tables I or II with equal ease. We have used the window algorithm of Table I. The dashed curve is that of $\bar{N}(t)$ computed as a running average from the start of the catalog. As before, there is a huge spike due to the effect of the Long Beach earthquake, which occurs shortly after the start of the catalog. The much more irregular curve is the graph of $n(t)$ and shows periods of elevated seismicity in Southern California, not associated with large earthquakes. Some of the elevated seismicity is of a broad regional character and some of it is local. We have identified the locally intense regions from the calculation of $r(t)$. We choose $C=\frac{1}{2}$ as before. The times for which $2r(t)$ exceeds $n(t)$ are shown by the short vertical strokes indicating the start of an alarm. The value of $2r$ is shown next to the short vertical stroke. The score for our analysis is the same as before. All earthquakes predicted before are predicted once again. The Borrego Mountain sequence acts as a precursor for the San Fernando sequence. We continue to have one false alarm starting in 1959. We fail to predict the 1940 ($M=6.7$) earthquake. The magnitude 6.5 aftershock of this event is "predicted" by a precursor which is near the threshold. We note that this earthquake was not predicted in one of the earlier attempts and was predicted by the other. Finally, the same events that serve as precursors in the hypothesis S also serve as precursors under hypothesis B. This means that short term aftershock activity, i.e. within 2 days of a main event, necessary for success of pattern B, also generates enough annual activity to provide pattern S.

V. The Sigma Hypothesis (Σ)

Some unease is certain to occur regarding the counting of numbers of earthquakes in the preceding exercises. Models of cessation of rupture in the dynamical earthquake process depend on inhomogeneities in the rheological environment (Knopoff, 1972). A given budget of stress-energy can be released by a wide variety of individual frequency-magnitude sequences, each presumably reflecting the local variations in strength of rocks. The question arises whether we should give more attention to a sequence of small earthquakes which presumably mirrors the presence of a series of "unbreakable objects" in the fault region with a relatively short spacing, in comparison with a single large earthquake. For example should we count two associated events of magnitude 5.0 as two in a counting scheme and hence give them twice as much weight as a single event of magnitude 6.0, even though the

latter releases more energy than the two smaller events combined? One solution to this problem is to sum the energies instead of the numbers of earthquakes. But, as is well known, this sum is completely dominated by the largest earthquakes, and we may as well neglect the smaller events. Benioff proposed a compromise between these two extremes by summing the square roots of the energies, which he called "strain."

The summation formulas are essentially of the form:

$$\Sigma(t) = \sum_i C 10^{dM_i}$$

If we choose $d = 1.5$, we have the sum of energies according to Gutenberg's magnitude-energy relation. The case of $d = 0.75$ sums the Benioff strains. The case $d = 0$ is that we have already considered, of counting shocks without regard to magnitudes. Keilis-Borok and Malinovskaya (1964) proposed the value equivalent to summation of energy in the power of about $2/3$.

We may use the above formula in the manner already described. We construct a running sum of $\Sigma(t)$, taking all events recurring within $t-s < t_i < t$ for some window interval s ; we also restrict M_i to some band of magnitudes below M_0 and eliminate from summation all aftershocks of events with $M_0 \geq 0$ even if their magnitudes are within this band. An alarm is sounded if $\Sigma(t)$ lies above some threshold; the threshold

$$.5 C 10^{dM_0}$$

was assumed as before.

VI. Two Other Regions

Premonitory patterns, described above, need and apparently deserve to be tested in other seismic regions of the world. At the time this paper was to be submitted we have some results for New Zealand, Italy, Japan, and Himalaya-Tibet.

The results for New Zealand are shown on the Figures 6-8 and in the Table 4. Earthquakes within the region, indicated on fig. 6, and focal depth ≤ 100 km were considered. The revised catalog of New Zealand earthquakes by W. Smith from 1940 to 1977 was used. The choice of the region and depth was made a priori, on the basis of possibilities, offered by the catalog, and some tectonic considerations. Most of the arbitrary parameters were chosen the same as for Southern California. However, we have to raise the threshold M_0 up to 6.7, and to increase e up to 7 days. The threshold for b_i is 12. The result for pattern B seems good, and the result for pattern Σ satisfactory, if we allow for the fact, that both patterns were formulated a priori and confirmed elsewhere. Two earthquakes with $M > 6.7$ just outside the chosen region and the only earthquake with b_i just below the threshold (78/8/28) are added to illustrate the stability of results. The main difference between the results in the last column is due to m .

The results for three regions of Italy are shown in Figure 9. These regions were chosen a priori in the paper (Caputo et al, in press). The threshold M_0 was lowered to 6.0, accordingly to the seismicity of Italy. The results may be considered as a confirmation of existence of premonitory Pattern B. But its actual use for prediction with the catalog in our disposition would be premature. For the southern region of Italy - Sicilia and Southern Appenines, the test of pattern B gave negative results. It may be associated with much larger focal depth of many strong earthquakes in this region. Pattern S for Italy was described in above mentioned paper. Pattern Σ so far was not tested.

VII. Conclusions

Bursts of seismicity, i.e. anomalous clustering of earthquakes in time and in space, may be regarded as patterns of seismicity that are precursory to strong earthquakes. These bursts may be diagnosed by constructing certain projections of the catalogs, three of which have been considered in this paper. In all likelihood, these projections are strongly overlapping and contain little independent information. These projections are:

Pattern B-bursts of aftershocks. We search for earthquakes of medium magnitude and having large numbers of aftershocks within a short time interval.

Pattern S-swarms. We identify high levels of concentration of earthquake activity in a relatively small part of time and space, on the background of sufficiently large general activity.

Pattern Σ -Exponential energy summation. We identify a sharp increase in the sum $\sum_i E_i^n$ where E_i is the energy of an earthquake occurring within a limited magnitude range and within a moving time-window. In practice, we have chosen the exponent n to be about $2/3$. We have provided parameterizations of these definitions; these parameterizations are most certainly non-unique, and are probably not optimal as well.

An application of these patterns to the catalog of earthquakes of Southern California for the years 1932-1978 gave 7 periods of alarm with a total duration of about $13\frac{1}{2}$ years. At least 5 out of 7 strong earthquakes of this region occurred during these alarm periods. The study of catalogs of Italy and New Zealand showed that pattern B also preceded the strongest earthquakes of these regions.

These premonitory patterns take place in almost all cases at considerable distance from the subsequent strong earthquakes. This leads us to form a conjecture that small, but significant, stress redistribution due to medium sized earthquakes at moderate distances can act as triggering agents; to account for the delays between the triggering and the triggered events, we must invoke a creep rheology for the rocks in the vicinity of the triggered event.

Acknowledgements

This research was performed as part of Project III of the U.S.-U.S.S.R. cooperative program in Earthquake Prediction. Support for this work, and the coverage of computing and other expenses, was provided by the Office of Earthquake Studies, United States Geological Survey. Consultation and assistance of Dr. J. K. Gardner, Dr. Y. Y. Kagan, and D. Eckelmann is gratefully acknowledged.

20

References

- R. Burridge and G.S. Halliday, Dynamic shear cracks with friction as models for shallow focus earthquakes, *Geophys.J.Roy.Astron.Soc.*, 35(1971) 261-283.
- M. Caputo, P. Gasperini, V. Keilis-Brook, L. Marcelli, E. Ranzman and I. Rotwain, Earth swarms as precursors of strong earthquakes in Italy, In press, *Annali di Geofisica*.
- F. Evison, The precursory earthquake swarms, *PEPI*, 15(1977) 19-23.
- M.E. Friedman, J.H. Whitcomb, C.R. Allen and J.A. Hileman, Seismicity of the Southern California Region, 1 January 1972 to 31 December 1974, Seismological Laboratory, California Institute of Technology, 1976.
- J.K. Gardner and L. Knopoff, Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, *Bull.Seismol.Soc.Amer.*, 64 (1974) 1363-1367.
- D.T. Griggs, Experimental flow of rocks under conditions favoring recrystallization, *Bull.Geol.Soc.Amer.*, 51 (1940) 1001-1022.
- J.A. Hileman, C.R. Allen and J.M. Nordquist, Seismicity of the Southern California Region, 1 January 1932 to 31 December, 1972, Seismological Laboratory, California Institute of Technology, 1973.
- Y. Kagan and L. Knopoff, Statistical search for non-random features of the seismicity of strong earthquakes, *Phys.Earth and Planet.Interiors*, 12 (1976) 291-318.
- Y. Kagan and L. Knopoff, Statistical study of the occurrence of shallow earthquakes, *Geophys.J.Roy. Astron.Soc.*, In press, 1978.
- V. Keilis-Borok, On estimation of the displacement in an earthquake source and of source dimensions, *Annali di Geofisica*, 12(1959) 205-214.
- V. Keilis-Borok and L.N. Malinovskaya, One regularity in the occurrence of strong earthquakes, *J.Geophys.Res.*, 69(1964) 3019-3024.

- V. Keilis-Borok and I. Rotvain, Two long-term precursors of strong earthquakes, In press, Computational Seismology, Vol. 12, 1978.
- V. Y. King and L. Knopoff, Stress drop in earthquakes, Bull.Seismol.Soc.Amer., 58(1968) 249-257.
- L. Knopoff, Energy release in earthquakes, Geophys.J. Roy.Astron.Soc., 1(1958) 44-52.
- L. Knopoff, Model for aftershock occurrence in Flow and Fracture of Rocks, Amer.Geophys.Union Monograph, 16, H.C. Heard, Ed. (1972) 259-263.
- L. Knopoff and J.K. Gardner, Homogeneous catalogs of earthquakes, Proc.Nat.Acad.Sciences, 63(1969) 1051-1054.
- L. Knopoff and J.K. Gardner, Higher seismic activity during local night on the Worldwide Earthquake Catalog, Geophys.J.Roy.Astron.Soc., 28(1972) 311-313.
- L. Knopoff, J.O. Mouton and R. Burridge, The dynamics of a one-dimensional fault in the presence of friction, Geophys.J.Roy.Astron.Soc., 35(1973) 169-184.
- A.T. Starr, Slip in a crystal and rupture in a solid due to shear, Proc.Camb.Phil.Soc., 24(1928) 489-500.

Table I

Windows for identification of aftershocks

M	R (km)	T (days)
2.5	19.5	6.
3.0	22.5	11.5
3.5	26.	22.
4.0	30.	42.
4.5	35.	83.
5.0	40.	155.
5.5	47.	290.
6.0	54.	510.
6.5	61.	790.
7.0	70.	915.
7.5	81.	960.
8.0	94.	985.

Table II

Window for identification of aftershocks,
another version

	R (km)	T (days)
2.4 $\geq$ M $\geq$ 2.0	20	1.43
6.4 $\geq$ M $\geq$ 5.5	40	182.5
5.4 $\geq$ M $\geq$ 5.0	40	91.25
4.9 $\geq$ M $\geq$ 4.5	40	45.63
4.4 $\geq$ M $\geq$ 4.0	40	22.81
3.9 $\geq$ M $\geq$ 3.5	40	11.41
3.4 $\geq$ M $\geq$ 3.0	20	5.70
2.9 $\geq$ M $\geq$ 2.5	20	2.85
M $\geq$ 6.5	80	365.3

Table 3

STRONG EARTHQUAKES AND PATTERN B
SOUTHERN CALIFORNIA

N	Date	Epicenter		M	b_1 (2 days)
		λ	ϕ		
1*	3 11 33	118.0	33.6	6.3	103
	12 31 34	114.8	32.0	7.1	
2	5 19 40	115.5	32.7	6.7	
3 ⁺	7 1 41	119.6	34.4	5.9	11
	10 21 42	116.0	33.0	6.5	
4	3 15 46	118.1	35.7	6.3	14
	4 10 47	116.6	35.0	6.2	14
	7 24 47	116.5	34.0	5.5	15
	12 4 48	116.4	33.9	6.5	
5	7 28 50	115.6	33.1	5.4	18
	7 21 52	119.0	35.0	7.7	
6	3 19 54	116.2	33.3	6.2	19
	11 12 54	116.0	31.5	6.3	19
	2 9 56	115.9	31.8	6.8	
7	12 1 58	115.8	32.3	5.8	11
	4 9 68	116.1	33.2	6.4	13
8	3 21 69	114.2	31.2	5.8	42
	2 9 71	118.4	34.4	6.4	69

* Preceded by strong foreshock: 12/30/34; 115.5, 32.3; 6.5.

⁺ Aftershock of 2.

Table 4. Strong earthquakes and pattern B in
New Zealand

Year	Mo/Day	ρ (S)	λ (E)	h	M	B(7 days)**	B(7 days)*	No. on Fig. 6
1942	6/24	40.9	175.9	12.0	7.0	3	25	
1942	8/1	41.0	175.8	55.0	7.1	0	5	
1943	2/17	45.2	167.9	95.0	6.9	1	3	
1943	8/2				6.7		0	
1943	9/22	36.0	177.0	33.0	6.8	0	0	
1946	6/26	43.2	171.5	30.0	6.2	4	33	1
1947	8/27	39.7	179.2	33.0	6.7	4	6	I
1951	1/10	42.8	173.2	12.0	5.6	5	35	2
1951	4/23	37.8	178.2	12.0	6.7	0	0	II
1952	8/28	39.9	176.9	12.0	5.8	1	11	
1960	5/24	44.2	167.7	12.0	7.0	7	64	III
1962	5/10	41.7	171.3	12.0	5.9	0	40	
1966	3/4	38.8	178.2	33.0	6.2	0	14	
1968	5/23	41.8	172.0	12.0	7.1	9	102	IV
1974	2/28	36.6	177.0	12.0	5.6	6	68	5
1974	9/20	44.4	168.0	12.0	5.9	1	34	5
1974	11/5	39.5	173.5	12.0	6.1	2	26	5
1976	3/20	39.3	177.5	12.0	5.6	3	75	5
1976	5/4	44.7	167.5	12.0	7.0	1	62	$\bar{V}$

Note: Data on the strong earthquakes are underlined. Two extreme versions of aftershocks identifications are compared in last columns:

* windows from the table 1, $m=5.5$

** windows from the table 2, $m=2$

In both versions lower limit for the magnitudes of the main shocks is $M_1=5$

Earthquake of 8/28/52 has B below the threshold.

Earthquake at 5/10/62 gives false alarm.

FIGURE CAPTIONS

- Fig. 1. Pattern B and strong earthquakes in Southern California. Aftershocks identified by the windows in Table 1. Black dot-value of $b_i(2)$ - number of aftershocks in the first two days. Most of main shocks have $b_i \leq 10$ and are not included. Vertical stroke - moment of strong earthquake with magnitude indicated at the top. Horizontal bar - alarm. Horizontal line - a threshold for alarm.
- Fig. 2. Pattern B and strong earthquakes in Southern California. Aftershocks are identified by the windows in Table 1. Notations the same as on Fig. 1. Threshold $\bar{b}(t)$ is shown by continuous curve.
- Fig. 3. Pattern S and strong earthquakes of Southern California. Vertical stroke-moment of strong earthquake with magnitude indicated at the top. Horizontal bars - alarms, their start at the moment, when pattern S ($n > \bar{N}$, $2r \geq n$) is diagnosed. Number at the beginning of alarm - double value of r .

Fig. 4. Pattern Σ and strong earthquakes of Southern California. Vertical stroke-moment of strong earthquake with magnitude indicated at the top. Horizontal bar - alarm. It starts when pattern Σ is diagnosed (Σ above threshold, indicated by dotted line).

Fig. 5. Strong earthquakes and pattern B.

Fig. 6. Strong normal earthquakes of New Zealand, and the earthquakes, which generated pattern B.

1. epicenter of strong earthquake. Roman number the same as in Table 4
2. epicenter of the earthquake, which generated pattern B (abnormal number of aftershocks in the first 7 days). Number the same as in Table 4.
3. false alarm
4. boundaries of the region

Fig. 7. Pattern B and strong earthquakes of New Zealand.

Circles- b_i for the windows in Table 2 and $m=2$

Triangles - b_i for the windows in Table 1 and $m = 1$.

Open signs - $M \leq 6.5$. Solid signs - $M \geq 6.7$.

Dashed column - the end of the catalog.

Vertical stroke-moment of strong earthquake with magnitude indicated at the top.

Fig. 8. Function Σ and strong normal earthquakes of New Zealand. Dashed column - the end of the catalog.

Fig. 9. Pattern B and strong earthquakes in three regions of Italy. Open circle-values of b_i . VIII - Maximal intensity (magnitude unknown). Vertical bars - moments of the strong ($M \geq 6$) earthquakes. Circle-value of b_i (x- for strong earthquake)

Figure 1

Figure 2

Figure 3

Figure 4

- $M \geq 7.0$
- $6.9 \geq M \geq 6.5$
- PATTERN B

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

On Soon-to-Break Seismic Gaps

by

V. Keilis-Borok and J. Rotwain

ABSTRACT

The premonitory seismicity pattern AQ (activation-quiescence) is formally defined as a combination of activation and quiescence on opposite sides of earthquake-prone areas. In other words, soon-to-break seismic gap is a gap, near which activation began recently. Retrospective application of this pattern to Anatolia and adjacent regions for 1900-1976 gives the average (per earthquake-prone area) duration of alarm of 13 years. Seven out of nine earthquakes, with $M \geq 7.2$ (not used in data-fitting) happened during alarm time. Joint application of this pattern with pattern "swarm" reduces average alarm time to six years without the loss of earthquakes during alarms. Pattern AQ should be tested on more regions, before its practical value could be estimated.

1. Hypothesis

Scheme of lineaments and their earthquake-prone segments are assumed to be determined. An example of such determination can be found in [3]. Let us consider the areas of the following types [Fig. 1]: an earthquake-prone area, where epicenters of the earthquakes with $M \geq M_0$ may be situated (1 on Fig. 1); lineaments between this area and the next such area inclusive (2,3,4 on Fig. 1); lineaments between this area and the end of the region (5 on Fig. 1). Hypothesis: earthquake with $M \geq M_0$ is being expected in area (1), if three following conditions are satisfied: this area is in the state of quiescence ("Q") or was in this state $\leq f$ years ago, at

least one of adjacent areas (2,3,4 or 5) is in the state of quiescence, and the opposite area is in the state of activation ("A"). In other words, earthquake is expected at the end of seismic gap, if the adjacent part of seismoactive lineament is activated. We data-fitted the following definitions of the states Q and A: area is in the state A (activation), if the last earthquake with $M \geq M_0$ was in this area $\leq a(M)$ years ago; area is in the state Q (quiescence), if the last earthquake with $M \geq M_q$ was in this area $\geq q(M)$ years ago.

The assumed functions $a(M)$, $q(M)$ are shown on Fig. 2. These definitions and even more the thresholds on Fig. 2 are by no means the only possible ones.

2. Pattern AQ, before earthquakes of $M > 7.2$ in Anatolia and adjacent regions

The region considered is shown on Fig. 3. 23 areas, where epicenters of earthquakes with $M \geq 7.0$ may be situated, were recognized for this region in [8]. We assumed that $M \geq 7.2$ are also possible in this region, since this magnitude threshold was established a priori, in a study of migration of epicenters. Functions $a(M)$, $b(M)$ were data-fitted, using 5 earthquakes on Large Anatolian fault (Table 1). Other strong earthquakes will fall into periods of alarm, if our hypothesis is correct.

The results for some earthquake-prone areas are shown on Fig.4. They seem satisfactory: 14 earthquakes with $M \geq 14$ since 1900 are known in this region [Table 1]. Eleven of them happened during alarms. Data on 5 of these 11 were used in fitting the parameters of $a(M)$, $q(M)$ (Fig.1): one of these 5 falls outside the alarm.

So, we got into .17% alarm period 7 events out of 9 independent ones. The probability of getting such results by random process with binomial distribution is negligible. However, actual confidence level is not so great, since pattern AQ was first found in the same region.

We determined also premonitory pattern S ("swarms") [1,2]. Its formulation was specified in the study for Italy [1] so that data-fitting was avoided.

If we declare an alarm, when patterns "AQ" and "S" are observed simultaneously, the average duration of alarms in one earthquake-prone area is reduced to 6 years, without the loss of earthquakes within the periods of alarm. With random binomial distribution we should increase the alarm time up to 40%, in order to have 1% probability of such result (7 out of 9 earthquakes during alarms).

Practical applications of such results are strongly reduced by the fact that two out of three alarms are false. However, they are not annulled by this fact, since they can be used for intensification of observations in the area of alarm, for weighting other precursors and for some practical low key safety measures [10].

A larger set of precursors seems necessary to get better results from the data base so far available [7,11]. Present study is described in more details in [6].

REFERENCES

1. Caputo, M. et al., Earthquake swarms, as forerunners of strong earthquakes in Italy, *Annali di Geofisica*, in print.
2. Evison, R., The precursory earthquake swarm, *PEPI*, 15, (1977).
3. Geltand, I. et al., Pattern recognition, applied to epicenters of strong earthquakes in California, *PEPI*, 13, (1976).
4. Karnik, V., *Seismology of the European Area*, Part 1, Praha, 1968.
5. Kazim, E. et al., A catalog of earthquakes for Turkey and surrounding area.
6. Keilis-Borok, V. and Rotvain, J., Two long-term precursors of strong earthquakes, *Vichislitel'naya seismologiya*, (Computational seismology), 12, (in Russian), in print.
7. Keilis-Borok, V., L. Knopoff, I. Rotvain and T. Sidorenko, Burst of aftershocks, as premonitory seismicity pattern, in print.
8. Kosobockov, V. and I. Rotvain, Pattern recognition of earthquake prone areas, VI. Magnitude $M \geq 7$, *Vichislitel'naya seismologiya*, (Computational seismology), 10, Moskva, Nauka, 1977 (in Russian).
9. Preliminary determination of epicenters, NOAA-USGS.
10. UNESCO interdisciplinary working group on seismic risk, *Resolutions*, Paris, 1977.
11. Wyss, U., Personal communication.

FIGURE 1 Division of lineaments.

1 - earthquake-prone area

2-4 - segments of adjacent lineaments up to the next earthquake-prone area inclusive

5 - adjacent lineament up to the end of the seismic region

FIGURE 2 Data-fitting of the functions $a(M)$, $b(M)$.

t - the time between the last earthquake of magnitude M and the subsequent strong earthquake (with magnitude 7.2 or more)

Points indicate the actual data for 5 earthquake-prone areas on Figure 3.

FIGURE 3 Lineaments of the first three ranks in Anatolia and adjacent regions, after [8].

- 1, 2 & 3 - lineaments of the first, second and third ranks
- 4 - transverse lineaments
dashed lines indicate hypothetical lineaments
- 5 - epicenters of the earthquakes with $M \geq 7.2$
- 6-8 - earthquake-prone areas determined by pattern recognition [8]
- 7 - Pattern S
- 8 - Patterns S and AQ

FIGURE 4 Identification of Pattern AQ for some earthquake-prone areas (number indicates the earthquake-prone area on Figure 3).

- 1-3 - states Q (solid line) or A (dashed line)
- 1 - earthquake-prone area (type 1 on Figure 1)
- 2,3 - adjacent lineaments (types 2-5 on Figure 1): 2 - to the East and West; 3 - to the North and South
- 4 - swarms
- 5 - moment and magnitude of strong earthquake
- 6 - Pattern S
- 7 - Patterns AQ and S

A POINT PROCESS ANALYSIS OF CHINESE EARTHQUAKE HISTORY

W. H. K. Lee

Office of Earthquake Studies

U.S. Geological Survey

Menlo Park, CA

and

D. R. Brillinger

Department of Statistics

University of California

Berkeley, CA

ABSTRACT

Since the 1950's, the Academia Sinica in Peking has carried out extensive research on the Chinese earthquake history. With a recorded history dating back some 3,000 years, there is a wealth of information on earthquakes in China. Despite the monumental tasks by the Academia Sinica, much work is still necessary to correct the existing data for historical changes in population, customs, communication, and dynasties. In this paper we will report on the status of our investigation on the Chinese earthquake history and present some preliminary results. By applying point process analysis in our study, we found suggestions that higher seismic activity occurred in "Central China" at an interval of about 300 years.

INTRODUCTION

Although seismicity is one of the primary data sources in making long-term earthquake prediction and hazard evaluations, knowledge of seismicity is rather limited. Modern seismographs began operation at the end of the 19th century, and adequate instrumental records for locating earthquakes on a worldwide basis date back only to 1904 for events of about magnitude $6\frac{1}{2}$ and greater. Indeed, accurate location of earthquakes (magnitude equal to or greater than 5) around the world became possible only after the establishment of the World-Wide Standardized Seismograph Network in 1963. Even in a well-studied area such as California, regional seismic networks were established in the 1930's and local networks in the 1960's. Consequently, knowledge of instrumentally determined seismicity covers only a short period of several tens of years.

In the framework of plate tectonics, tens or hundreds of years may be required to accumulate sufficient strains to generate large earthquakes. To identify seismic gaps and to determine how often a large earthquake occurs, accurate seismicity data covering periods of hundreds or thousands of years are required. Except for extremely active seismic regions, the instrumentally determined seismicity covers too short a period, and we must therefore rely upon historical records to estimate long-term seismicity.

With the exception of the Middle East, only China has a continuously recorded history extending back some 3,000 years. In this paper we report on the status of our investigation on Chinese earthquakes, and

present some preliminary results. We will also discuss a variety of difficulties which we encountered in the course of our study.

SOURCE MATERIALS ON CHINESE EARTHQUAKES

The first reliable description of a Chinese earthquake (in the 12th century B.C.) appeared in Lushi Chunqiu (Annals of Mr. Lu) compiled in the third century B.C. Several earthquakes were mentioned in Shiji (Records of the Grand Historian) written by Si-ma Qian in the first century B.C. The first Chinese earthquake catalog appeared as a subsection of natural omens in Hanshu (Standard History of the Former Han Dynasty) by Ban Gu in the first century A.D. A list of 13 earthquakes, including damage and casualties, were summarized. In all of the subsequent 24 standard histories or dynasty records, earthquakes were reported.

In addition to normal reportings of earthquakes in the dynasty records, catalogs of Chinese earthquakes in various details also appear in numerous encyclopaedias and collected works. For example, earthquakes were summarized in a section of Tongzhi (Comprehensive Encyclopaedia of Institutions) by Zheng Qiao in 1149 A.D., and of Wenxian Tongkao (Critical Examination of Documents and Studies) by Ma Duan-lin in 1224 A.D.

With the advent of printing in the 10th century A.D., it became popular in China to compile local gazetteers describing details of local history and geography, especially in the Ming and Ching dynasties (14th-20th centuries). About 7,500 local gazetteers, with a total of 110,000 chapters, are now in existence. Detailed descriptions of earthquakes were often found in these voluminous documents.

At the turn of the last century, many western seismologists were fascinated by the rich historical accounts of Chinese earthquakes. Several catalogs of Chinese earthquakes were published in the west, and these activities were reviewed by Drake (1912). All of these catalogs suffer two main defects in that references to the original sources are not specific, and that the epicenter and size of earthquakes are not adequately quantified.

In the 1950's, the Seismological Committee of the Academia Sinica initiated a systematic study of Chinese historical earthquakes in order to provide data on earthquake hazards in support of industrialization planning in China. The first task the Committee did was to compile, as completely as possible, all descriptions of earthquakes in Chinese literature and documents. In 1956, two large volumes totaling 1653 pages were published under the title of "Chronological Tables of Earthquake Data of China" (Academia Sinica, 1956), and covered the period from 1189 B.C. to 1955 A.D. Descriptions of earthquakes were extracted from over 8,000 documents (many of them multi-volume publications) and were arranged by provinces. Within each province, entries were arranged in chronological order. Source references were cited, and notes were added to clarify ancient geographic names and errors in reporting. A summary of earthquake activity for each province was given, and an index by county and city of felt earthquakes was also included.

Attempts to quantify the historical descriptions of earthquakes were also carried out by the Academia Sinica. The results were published under the title "Catalog of Chinese Earthquakes" (Academia Sinica, 1970a), and

covered the period from 1177 B.C. to 1949 A.D. In this volume, the date, epicenter, magnitude, and epicentral intensity were provided for selected earthquakes noted in the 1956 compilation. Selections were based on whether or not sufficient information existed to deduce an earthquake location and magnitude. A brief summary of large earthquakes in China with magnitudes greater than or equal to 6 was also published (Academia Sinica, 1970b) and revised later (Academic Sinica, 1974, 1976). It covered the period from 780 B.C. to the present.

Lee, Wu, and Jacobsen (1976) presented a catalog of historical earthquakes in China from 1177 B.C. to 1899 A.D. in a form suitable for computers. Their source materials are publications in Chinese of the Institute of Geophysics, Academia Sinica as described in the previous paragraph. A similar catalog for instrumentally determined earthquakes in China after 1900 A.D. (magnitude ≥ 6) was compiled by Lee, Wu, and Wang (1978) from an extensive search of literature in Chinese, Japanese, and western languages.

LIMITATIONS OF CHINESE EARTHQUAKE HISTORY

The accuracy and completeness of the Chinese historical earthquake data are difficult to evaluate. The questions are: How accurately and completely were the data recorded, passed down through history, and interpreted in the publications by Academia Sinica? Although fairly complete dynasty records began at about 200 B.C., printing did not begin in China until about 1000 A.D. so that very few original documents

(written before 1000 A.D.) exist and many ancient records are lost.

However, the dynasty records and some ancient literature that survive are fairly well preserved through hand copying. Recent editions of dynasty records and ancient literature indicate only minor discrepancies among different earlier and independent versions, and small parts may be missing but could be filled in from other sources that quoted the missing text.

The accuracy of Chinese dynasty records may be subject to debate.

Earthquakes, like other disasters or unusual phenomena such as eclipses, were often regarded as punishments to the emperor and/or the people for their sins. As a result, the record of their occurrence could have been misrepresented by court historians.

Another consideration is that ancient China does not cover as much territory as modern China. Earthquakes would be recorded mostly within the territory that was under the control of the dynasty, and could have been noticed only if there was a sufficient number of people living near the epicentral region. Consequently, one would expect fewer and fewer earthquake reports as one goes back in time. In addition, many historical earthquakes (especially ancient ones) could not be interpreted and quantified because their descriptions are vague. There is also a tendency for the magnitudes of the older earthquakes to be underestimated because they are based on estimates of epicentral intensity and extent of damaged and felt areas. Because older documents tend to get lost and ancient population size was smaller than current, one would underestimate these parameters. The intensity scale was not described in Academia Sinica (1970a, 1970b, 1974, 1976), but we suspect that the intensity scale used

is that of Hsieh (1957) which consists of 12 degrees similar to the modified Mercalli scale. The magnitude (M) was determined by the formula

$$M = 0.58 I_0 + 1.5$$

where I_0 is the epicentral intensity, and adjusted according to the extent of damaged and felt areas (Lee, 1958; Academia Sinica, 1970a). However, the method of adjustment is not described in the above-mentioned sources.

HISTORICAL AND INSTRUMENTALLY DETERMINED SEISMICITY

The historical earthquakes (before 1900 A.D.) compiled from recent Chinese publications by Lee, Wu, and Jacobsen (1976) were plotted on a fault-map base as shown in Figure 1. In Figure 2, we show the instrumentally determined earthquakes (after 1900 A.D.) compiled from various publications by Lee, Wu, and Wang (1978). Comparison of these two figures indicates that there is pronounced difference in the distribution of earthquakes--the apparent seismic quiescence of western China in the pre-instrumental period. This must be due virtually to the lack of historical data there because western China has a very low population and was not under the firm control of China until very recently.

To put the historical seismicity in proper perspective, one must consider the population distribution in time and space. Population distribution is in turn influenced by topography of the land. Figure 3

shows the general topography of China indicating that most of western China has an elevation in excess of 1,000 meters. The modern population distribution is shown in Figure 4. It is clear that most of the Chinese population is concentrated in land below 1,000 meters. The boundaries of the past Chinese dynasties vary greatly as shown in Figure 5 for a few examples. With these factors in mind, we may select a region of China that appears approximately homogeneous spatially in earthquake reportings. This area is taken to be from 30° N to 42° N, and from 100° E to 125° E as shown in Figure 6. We call this study area "Central China" (recognizing that it is not geographically the central region).

As discussed in the previous section, the magnitudes of historical Chinese earthquakes are probably underestimated. Although the historical catalog includes earthquakes of magnitude less than 6, we think that most of the events are probably of magnitude 6 or greater. Therefore we consider these historical earthquakes together with the instrumentally determined ones of magnitude 6 or greater.

In Table 1, we list the number of Chinese earthquakes by time and region. The reason to choose the "Central China" region becomes obvious. From 1177 B.C. to 0 B.C., there were 12 events in the Chinese catalogs of which 11 occurred in "Central China". For the next thousand years, "Central China" accounted for 90% of all Chinese earthquakes. From 1001 A.D. to 1900 A.D., only 55% of all Chinese earthquakes occurred in "Central China". During the instrumental period from 1901 to 1976, only 11% of all Chinese earthquakes occurred in "Central China".

If we consider that the instrumentally determined earthquakes are complete and that the recent seismicity is representative of the ancient one, then we must conclude that about 2/3 of all earthquakes in "Central China" which would have occurred did not appear in the catalog for the period from 1001 A.D. to 1900 A.D. As noted in the previous section, printing and local gazetteers began in the 10th century but did not become widespread until the 15th century. If we take this into account, we have 231 earthquakes in "Central China" from 1501 A.D. to 1900 A.D., yielding a rate of 58 earthquakes/century (versus the modern rate of about 86 earthquakes/century as shown in Table 3 later).

The above analysis clearly suggests that the present earthquake catalog is not complete even in "Central China". Despite a history of some 3,000 years, more and more earthquakes are absent from our present catalog as we go back in time.

CORRECTIONS TO THE EARTHQUAKE CATALOG IN "CENTRAL CHINA"

A list of earthquakes in "Central China" was extracted from the earthquake catalogs given by Lee, Wu, and Jacobsen (1976) and Lee, Wu, and Wang (1978), and is given in Table 2. A breakdown of the number of earthquakes by century is presented in Table 3, and it is evident that there are considerable variations. As discussed in the previous sections, there are many factors leading to fewer and fewer earthquakes in an earthquake catalog as one goes back in time. We shall now make an effort to correct for these factors.

One major factor is the "stability" of the dynasties. If there were wars and famines, the chance for an earthquake to be recorded and the information to be preserved would decrease. A measure of the dynasty stability is the census data which were collected for taxation. The taxation census is probably not a true measure of the actual population, but rather reflects how well the dynasty is in control of the country. Detailed taxation census appeared in dynasty records, and a summary may be found in Chang (1959). A plot of the census data is given in Figure 7. It is rather surprising that the taxation census show an apparent population of about 30 million from 0 to 1700 A.D. with large fluctuations. The apparent population increased more than 10-fold in the 18th century for reasons that scholars are still debating.

There are four major "troughs" in Figure 7 which are related to wars and major dynasty changes. The first decrease is related to the change from the Former Han dynasty to the Later Han dynasty. The second decrease is related to the breaking up of China into several warring factions at the end of the Later Han dynasty. The third decrease occurred during the middle of the Tang dynasty when there were extensive influxes of western tribes into "Central China". The fourth decrease is due to the change from the Ming dynasty to the Ching dynasty when the Manchu tribe from northeastern China crossed the Great Wall and took over "Central China".

It is not clear to us how to best utilize the taxation census to correct for earthquakes that were not recorded. We use a simple model here for our initial investigation. The probability that an earthquake will be recorded at time t is modelled by:

$$\Pi_1(t) = 1 - C \left[(S - P(t))/S \right]; \quad \Pi_1(t) \leq 1$$

where $P(t)$ is the apparent population according to taxation census, S is a saturation population, and C is a constant. Both S and C are model constants, and we choose $S = 50$ million and $C = 1/3$. Obviously $\Pi_1(t)$ cannot be greater than 1 and we impose this condition. The resulting probability as a function of time is given in Figure 8.

Another major factor is how well the historical earthquake records are preserved in the present literature. It is obvious that printing plays an important part in preserving historical records, and the advent of printing occurred in the 10th century and became widespread by the 15th century. Therefore, we may construct a simple model for the probability of an earthquake record to be preserved by specifying:

$$\begin{aligned} \Pi_2(t) &= C_1 & \text{at} & \quad t = 0 \text{ A.D.} \\ &= C_2 & \text{at} & \quad t = 1000 \text{ A.D.} \\ &= 1 & \text{at} & \quad t = 1500 \text{ A.D.} \end{aligned}$$

where C_1 and C_2 are constants, and Π_2 linearly extrapolated between values at given times. The resulting probability is given by Figure 9 for $C_1 = 0.05$ and $C_2 = 0.1$. (These constants were chosen arbitrarily.)

We may combine the above two probabilities into a probability that an earthquake will appear in present literature. Specifically

$$\Pi(t) = \Pi_1(t) \Pi_2(t) .$$

The resulting probability function is shown in Figure 10. By dividing the observed number of earthquakes by this probability we may estimate the number of earthquakes in each century and the results are shown in Table 3 and Figure 11. Because there are very few earthquakes in our catalog before 1000 A.D., corrections by the probability function tend to introduce large fluctuations and are unreliable. Corrections to data after 1000 A.D. seem to bring out a relatively high seismicity in the 11th and 14th centuries more clearly. This simple exercise supports the well-known argument that there is a temporal pattern of seismicity in "Central China". Earthquake activity perhaps increases at an interval of about 300 years as evidenced by the relatively large numbers of earthquakes in the 11th, 14th, 17th, and 20th centuries in comparison to their adjacent centuries.

Another way to count the number of earthquakes is through a moving time window. We shall restrict ourselves to the data from 1000 A.D. to the present. If we count the number of earthquakes in a time window of 50 years at increments of 20 years, we have the curve as shown in Figure 12. If we use the probability function (Figure 10) to correct the earthquake data as described above, we have the curve as shown in Figure 13. The results are suggestive that seismicity peaked at the 10th, 14th, 17th and perhaps the 20th century.

POINT PROCESS ANALYSIS

In recent years, point process analysis has been introduced into statistics to analyze a sequence of times of events in manners similar to those of ordinary time series analysis. General introductions to point process analysis may be found in Cox and Lewis (1966), Daley and Vere-Jones (1972) and Brillinger (1976). Applications of the theory to seismology may be found in Vere-Jones (1970), Lomnitz (1974), Udias and Rice (1975), Kagan and Knopoff (1976, 1978).

By a point process is meant a sequence of times, say $\tau_1, \tau_2, \tau_3, \dots$, at which certain events of interest occur. By a stochastic point process is meant a point process in which the times at which the events occur is random. A stochastic point process may be characterized by giving the probability distribution of the number of events in particular time intervals or collections of intervals. It may alternately be characterized by giving the probability distribution of the times between successive events. In the seismological case the events are earthquakes and the times are those at which certain earthquakes of interest occurred.

A point process is often described by a counting function $N(t)$, with $N(t)$ giving the number of events in the time interval $(0, t]$. This is a nondecreasing step function that increases by 1 every time a new event occurs. Using the Dirac delta function one may give a symbolic representation of the derivative of the point process as

$$\frac{dN(t)}{dt} = \sum_j \delta(t - \tau_j)$$

where the τ_j are the times of the events. In a variety of circumstances this derivative may be viewed as an ordinary time series and suggestive expressions set down. This series will be 0 for almost all t , but infinite at times at which events occur.

A key descriptor of a stochastic point process is provided by its rate, or intensity, function. This is defined as

$$h_N(t) = \lim_{h \rightarrow 0} \text{Prob} \left\{ \text{event in the time interval } (t, t+h) \right\} / h$$

and is analogous to the mean function of an ordinary (stochastic) time series. An important second-order parameter is provided by the autointensity function

$$h_{NN}(t_1, t_2) = \lim_{h \rightarrow 0} \text{Prob} \left\{ \text{event in } (t_1, t_1 + h) \mid \text{event at } t_2 \right\} / h$$

$$t_1 \neq t_2,$$

measuring the degree of serial dependence of the process in a certain sense.

A stochastic point process is called stationary if its probabilistic properties are unaffected by shift of the time origin. In this situation the rate function is constant and the (above) autointensity function depends on $t_1 - t_2$ alone. In our application of

point process analysis to the Chinese earthquake data, a basic assumption is that the earthquake-generating process is not changing in time, and that the earthquakes generated correspond to a realization of a stationary stochastic point process. This seems reasonable in view of the theory of plate tectonics suggesting that the plate motions which generate earthquakes do not change in a time span of a few thousand years.

Let $\tau_1, \tau_2, \tau_3, \dots$ denote the times of the earthquakes occurring in our earthquake catalog and let $N(t)$ denote the corresponding counting function. We further assume that the probability of an earthquake occurring at time t being listed in our catalog is $\Pi(t)$ and that the events being listed are statistically independent for each event. (We recognize that this is not totally realistic as whole stretches of events may have been lost or unrecorded together, but it seems a reasonable first approximation.) For simplicity in this initial study we shall take the function of Figure 10 for $\Pi(t)$. The derivative of the process $N(t)$ now has the representations

$$\frac{dN(t)}{dt} = \sum_j \delta(t - \tau_j) = \sum_k I_k(t - \sigma_k)$$

with the I_k independent 0-1 variates having

$$\text{Prob} \{I_k = 1\} = \Pi(\sigma_k)$$

$$\text{Prob} \{I_k = 0\} = 1 - \Pi(\sigma_k)$$

where the σ_k are the times at which earthquakes actually occurred. Let the point process $M(t)$, of actual earthquake times, have rate h_M and autointensity function $h_{MM}(t)$, then the rate of the process $N(t)$ is

$$h_N(t) = \Pi(t) h_M$$

and its autointensity is

$$h_{NN}(t_1, t_2) = \Pi(t) h_{MM}(t_1 - t_2) .$$

Estimates of the rate and autointensity function of a stationary point process are discussed in Cox and Lewis (1966) and Brillinger (1976). (Examples are given in Udias and Rice [1975].)

Supposing the catalog covers the time interval $(0, T]$ with $N(T)$ events occurring in total at times $\tau_1, \tau_2, \dots, \tau_{N(T)}$, the rate of the basic process M may be estimated by

$$\hat{h}_M = T^{-1} \sum_{j=1}^{N(T)} 1 / \Pi(\tau_j)$$

Further, the autointensity of the M process may be estimated by

$$\hat{h}_{MM}(t) = \frac{1}{\beta T \hat{h}_M} \sum_{i \neq j} \left| \tau_i - \tau_j - t \right| < \beta/2 \Big / [\Pi(\tau_i) \Pi(\tau_j)]$$

with the symbol $\{E\}$ defined to be 1 if the event E is true and to be 0 if E is false and with β , a cell width. (In fact in our calculations the above estimate $\hat{h}_{MM}(t)$ is weighted together with $\hat{h}_M$ in order that the estimate have the proper behavior for large t .)

We have estimated the autointensity function for the earthquake sequence in our catalog from 1000 A.D. to 1976 A.D. The result is shown in Figure 14 where the first peak of the curve suggests a periodicity of about 300 years in the occurrence of earthquakes in "central China". The result is in agreement with the eariler findings described in the previous section. The central peak suggests the clustering together of the events partly due to aftershocks. For our present data ($T = 977$ years, $N(t) = 349$), we obtain the rate of earthquake occurrence, $\hat{h}_M$, equal to 0.449 events/year from the formula given. This corresponds to about 45 events per century.

In the case that all events are observed (i.e., $\Pi(t) = 1$), the standrad error of the estimate of the autointensity function may be estimated by

$$\epsilon = \sqrt{\frac{\hat{h}_{MM}(t)}{\beta T \hat{h}_M}}$$

(see Brillinger, 1976).

For a point process with missing events, a formula for the standard error of the estimate of the autointensity function has been developed. It is rather complex and simplifications are now being sought.

DISCUSSION

It is obvious that the results from analyses of historical earthquakes depend critically on the quality and completeness of the earthquake catalog. In "Central China", the present catalog is inadequate before 1500 A.D. and we are now in the process of improving the existing catalog and developing methods of dealing with its incompleteness. We also plan to develop an extensive set of statistical tools to analyze the data, and would welcome suggestions and comments.

REFERENCES

- Academia Sinica, Seismological Committee (1956). Chronological Tables of Earthquake Data of China, Science Press, Peking, 2 vol., 1653 pp. (in Chinese).
- Academia Sinica, Institute of Geophysics (1970a). Catalog of Chinese Earthquakes, Academia Sinica, Peking, 361 pp. (in Chinese).
- Academia Sinica, Institute of Geophysics (1970b). Summary of Large Earthquakes in China (from 780 B.C. to February 1970, with magnitude ≥ 6), Academia Sinica, Peking, 29 pp. (in Chinese).
- Academia Sinica, Institute of Geophysics (1974). Summary of Large Earthquakes in China (from 780 B.C. to 1973, with magnitude ≥ 6). Academia Sinica, Peking, 31 pp. (in Chinese).
- Academia Sinica, Institute of Geophysics (1976). Summary of Large Earthquakes in China (from 780 B.C. to 1976, with magnitude ≥ 6), Academia Sinica, Peking, 29 pp. (in Chinese).
- Brillinger, D. R. (1976). Measuring the association of point processes: a case history, Am. Math. Monthly, 83, 16-22.
- Chang, C. Y. (1959). China's Population Problem, Chinese Population Society, Taipei, 444 pp. (in Chinese).
- Cox, D. R., and P. A. W. Lewis (1966). The Statistical Analysis of Series of Events, Barnes & Noble, New York.
- Daley, D. J., and D. Vere-Jones (1972). A summary of the theory of point processes, in "Stochastic Point Processes", edited by P. A. W. Lewis, Wiley-Interscience, New York, p. 299-383.

- Drake, N. F. (1912). Destructive earthquakes in China, Bull. Seism. Soc. Am., 2, 45-91.
- Fessler, L. (1963). China, Time Inc., New York, 176 pp.
- Hsieh, C. M. (1973). Atlas of China, McGraw-Hill, New York, 282 pp.
- Hsieh, Y. S. (1957). A new scale of seismic intensity adapted to the conditions in Chinese territories, Acta Geophys. Sinica, 6, 35-47 (in Chinese with English abstract).
- Kagan, Y. and Knopoff, L. (1976). Statistical search for non-random features of the seismicity of strong earthquakes, Phys. Earth Planetary Interiors, 12, 291-318.
- Kagin, Y. and Knopoff, L. (1978). Statistical study of the occurrence of shallow earthquakes, Publ. 1694, Inst. Geophys. Planetary Physic., UCLA.
- Lee, S. P. (1958). A practical magnitude scale, Acta Geophys. Sinica, 7, 98-102 (in Chinese with English abstract).
- Lee, W. H. K., F. T. Wu, and C. Jacobsen (1976). A catalog of historical earthquakes in China compiled from recent Chinese publications, Bull. Seism. Soc. Am., 66, 2003-2016.
- Lee, W. H. K, F. T. Wu, and S. C. Wang (1978). A catalog of instrumentally determined earthquakes in China (Magnitude ≥ 6) Compiled from various sources, Bull. Seism. Soc. Am., 68, 383-398.
- Lomnitz, C. (1974). Global Tectonics and Earthquake Risk, Elseiver, Amsterdam.

- Udias, A. and J. Rice, (1975). Statistical analysis of microearthquake activity near San Andreas Geophysical Observatory, Hollister, California, Bull. Seis. Soc. Am., 65, 809-827.
- Vere-Jones, D. (1970). Stochastic models for earthquake occurrence, J. R. Stat. Soc. B., 32, 1-62.

Table 1. Number of Chinese earthquakes by time and region

<u>Time</u>	<u>Time Period (years)</u>	<u>Number of Earthquakes in</u>	
		<u>China</u>	<u>"Central" China</u>
1177 B.C. to 0 B.C.	1177	12	11
1 A.D. to 1000 A.D.	1000	42	38
1001 A.D. to 1900 A.D.	900	512	284
1901 A.D. to 1976 A.D.	76	579	65

TABLE 2. List of earthquakes in "Central China".

YEAR	DATE	LAT.	LONG.	MAG.	INT.	PROVINCE
-1177		34.5 N	107.8 E	4.5		SHENSI
-780		34.5 N	107.8 E	6.5		SHENSI
-231		36.5 N	111.5 E	6.5	8.	SHANSI
-193	2	35.4 N	103.9 E	6.5		KANSU
-186	2 22	33.4 N	104.8 E	6.5		KANSU
-159	6	32.2 N	110.4 E	5.	6.	HUPEH
-154		32.2 N	110.4 E	5.	6.	HUPEH
-142?		32.2 N	110.4 E	5.	6.	HUPEH
-70	6 1	36.3 N	119.0 E	7.0	9.	SHANTUNG
-47	4 17	35.1 N	104.6 E	6.75	9.	KANSU
-35	7	34.4 N	109.0 E	5.	6.5	SHENSI
46	10 23	33.0 N	112.5 E	6.5	8.	HONAN
128	2 23	34.7 N	105.4 E	6.5	8.	KANSU
138	3 1	35.5 N	104.0 E	6.75	9.	KANSU
143	10	34.7 N	105.3 E	7.0	9.	KANSU
294	7	32.6 N	116.8 E	5.5	7.	ANHWEI
294	9	40.3 N	116.0 E	5.5	7.	HOPEH
319	6 17	34.0 N	105.2 E	4.5		KANSU
344		36.3 N	114.5 E	5.5		HOPEH
373	8	36.6 N	101.8 E	4.75	6.	TSINGHAI
406	6	36.3 N	104.5 E	5.5		KANSU
408		36.8 N	118.3 E	5.	6.5	SHANTUNG
408		39.0 N	100.5 E	4.75	6.	KANSU
416		34.3 N	105.5 E	5.	6.	KANSU
421		41.6 N	121.4 E	5.	6.	LIAONING
462	8 16	35.6 N	116.8 E	5.5		SHANTUNG
495	3 31	37.5 N	121.2 E	5.5	7.	SHANTUNG
506	8 30	37.9 N	102.6 E	4.75	6.	KANSU
512	5 21	39.0 N	113.0 E	7.5	10.	SHANSI
575	1 14	37.9 N	102.6 E	5.5	7.	KANSU
600	12 13	34.3 N	108.9 E	5.5	7.	SHENSI
638	2 11	32.6 N	103.6 E	5.	6.	SZECHWAN
649	9 12	36.1 N	111.5 E	5.5	7.	SHANSI
692	3?	37.5 N	117.5 E	5.	6.	SHANTUNG
734	3 19	34.7 N	106.3 E	7.0	9.	KANSU
756	11 27	39.0 N	100.5 E	6.0	7.	KANSU
777		37.8 N	115.2 E	6.0	8.	500 HOPEH
788	3 8	32.5 N	109.2 E	6.5	8.	SHENSI
793	5 27	34.5 N	109.7 E	6.0	7.5	SHENSI
835	4 11	34.3 N	108.9 E	4.75	6.	SHENSI
836	2 25	34.3 N	108.9 E	4.75	6.	SHENSI
839		34.4 N	104.0 E	6.5		KANSU
865	12	35.9 N	111.4 E	5.5	7.	SHANSI
867	2 14	35.9 N	111.4 E	5.5	7.	SHANSI
876	7 14	37.8 N	105.9 E	6.5	8.	NINGSIA
879	3	34.2 N	109.3 E	4.75	6.	SHENSI
880	2	34.5 N	107.8 E	4.75	6.	SHENSI
953	11	36.3 N	115.1 E	4.75	6.	HOPEH
999	10	31.8 N	119.9 E	5.5	7.	KIANGSU
1010	11	38.1 N	106.4 E	5.5	7.	NINGSIA
1011	8	38.2 N	114.6 E	4.75	6.	HOPEH
1022	4	39.7 N	113.1 E	6.5	8.	SHANSI
1038	1 9	38.4 N	112.9 E	7.25	10.	23000 SHANSI
1046	4 18	37.8 N	120.7 E	5.	6.	SHANTUNG
1057		39.5 N	116.3 E	6.75	9.	50000 HOPEH
1068	8 14	38.5 N	116.1 E	6.0	8.	HOPEH
1069	1 18	38.3 N	116.2 E	4.75	6.	HOPEH
1092	12?	37.9 N	102.6 E	4.75	6.	KANSU
1125	8 30	36.0 N	103.9 E	7.0	9.	KANSU
1143	4	38.5 N	106.3 E	6.5	8.	NINGSIA
1169	1 24	31.9 N	104.4 E	4.75	6.	SZECHWAN
1209	12 04	36.0 N	111.8 E	6.5	8.5	SHANSI
1219	5 21	36.0 N	106.2 E	6.5	8.5	NINGSIA
1290	9 27	41.5 N	119.3 E	6.75	9.	LIAONING *
1291	8 25	36.1 N	111.5 E	6.5	8.	SHANSI
1303	9 17	36.3 N	111.7 E	8.0	11.	SHANSI
1304	2	36.1 N	111.5 E	5.5	7.	SHANSI
1304	9 3	37.5 N	112.6 E	4.75	6.	SHANSI
1305	5 3	39.8 N	113.1 E	6.5	8.5	SHANSI
1306	9 12	35.9 N	106.1 E	6.5	8.5	NINGSIA
1314	10 5	36.5 N	113.8 E	6.0	8.	340 HOPEH
1316		36.4 N	111.1 E	5.5	7.	SHANSI
1322		40.6 N	115.0 E	4.5		HOPEH
1336	3?	30.1 N	115.9 E	4.7	6.	HUPEH
1337	9 8	40.4 N	115.7 E	6.5	8.	HOPEH
1338	8 2	40.4 N	115.2 E	5.	6.	HOPEH
1342	5 5	37.9 N	112.6 E	5.5	7.	SHANSI
1346	4	37.1 N	118.0 E	4.75	6.	SHANTUNG
1351	5 14	37.3 N	113.0 E	5.5	7.	SHANSI
1352	4 18	35.6 N	105.3 E	7.0	9.	KANSU
1368	7 8	37.6 N	112.5 E	6.0	7.	SHANSI
1372	8 16	32.0 N	118.8 E	4.75	6.	KIANGSU
1378	4 30	38.5 N	106.3 E	5.75	7.	NINGSIA
1399	4 29	32.0 N	118.8 E	4.75	6.	KIANGSU
1407	11	31.2 N	112.6 E	5.5	7.	HUPEH
1425	3 7	31.7 N	116.5 E	5.75	7.	ANHWEI
1433		30.5 N	115.2 E	4.75	6.	HUPEH
1440	10 26	36.2 N	103.4 E	6.25	8.	KANSU
1448	9 30	38.3 N	109.7 E	5.	6.5	SHENSI
1467	6 9	39.6 N	112.3 E	5.5	7.	SHANSI
1469	11 4	31.2 N	112.6 E	5.	6.5	HUPEH
1474	12 11	38.0 N	106.3 E	5.5	7.	NINGSIA
1477	3 19	35.2 N	104.2 E	5.	6.	KANSU
1477	5 13	38.5 N	106.3 E	6.5	8.	NINGSIA
1484	1 29	40.4 N	116.1 E	6.75	8.5	HOPEH
1485	1 17	34.8 N	110.4 E	5.	6.5	SHANSI
1485	5 27	40.2 N	118.0 E	5.	6.	HOPEH
1487	8 10	34.3 N	109.1 E	6.25	8.	SHENSI
1488	9 15	31.7 N	103.9 E	5.5	6.5	SZECHWAN
1495	4 10	37.6 N	105.6 E	6.25	8.	NINGSIA
1497	2 17	36.3 N	112.9 E	4.75	6.	SHANSI
1511	1 19	34.8 N	110.1 E	7.	9.	SHENSI
1502	1 17	34.8 N	110.1 E	5.	6.	SHENSI
1502	10 17	35.7 N	115.3 E	6.5	8.	HONAN *
1505	7 10	37.8 N	105.9 E	5.5	7.	NINGSIA
1505	10 16	35.3 N	110.8 E	5.	6.	SHANSI
1506	3 19	35.3 N	110.1 E	5.5	7.	SHENSI
1506	8 28	36.3 N	120.7 E	4.75	6.	SHANTUNG
1522	1 28	34.2 N	114.1 E	5.75	7.	HONAN
1523	8 14	30.0 N	122.1 E	5.5	7.	CHEKIANG
1524	2 4	33.8 N	113.9 E	5.75	7.	HONAN
1527		39.8 N	118.1 E	5.5	7.	HOPEH
1528	6?	37.9 N	114.7 E	5.	6.	HOPEH
1532	11 6	39.9 N	116.9 E	5.5	7.	HOPEH
1536	10 22	39.8 N	116.8 E	6.	7.5	HOPEH
1537	5 13	33.6 N	117.6 E	5.5	7.	ANHWEI
1538		38.0 N	115.6 E	5.	6.	HOPEH

1542	11	19	39.0	N	111.0	E	5.	6.	SHANSI
1542	11	19	34.7	N	104.9	E	5.	6.	KANSU
1543	9	13	37.8	N	120.7	E	6.	8.	SHANTUNG
1549	6		37.0	N	111.9	E	5.	6.	SHANSI
1556	1	23	34.5	N	109.7	E	8.	11.	SHENSI
1556	12	3	38.8	N	101.1	E	5.	6.	KANSU
1558	6		37.5	N	112.2	E	5.	6.	SHANSI
1558	11	21	34.5	N	109.7	E	5.5	7.	SHENSI
1561	2	21	38.8	N	101.1	E	5.	6.	KANSU
1561	7	25	37.4	N	106.0	E	7.25	9.5	NINGHSIA
1562			38.5	N	106.3	E	5.	6.	NINGHSIA
1562	6		39.7	N	118.7	E	5.	6.	HOPEH
1567			39.7	N	119.2	E	4.75	6.	HOPEH
1568	1		34.2	N	109.3	E	5.	6.	SHENSI
1568	4	1	38.5	N	106.3	E	5.75	7.	NINGHSIA
1568	4	2	34.4	N	109.2	E	5.5	7.	SHENSI
1568	4	12	33.1	N	107.0	E	5.	6.	SHENSI
1568	4	25	39.0	N	119.0	E	6.		HOPEH
1568	5	15	34.4	N	109.0	E	6.75	9.	SHENSI
1569			32.7	N	109.0	E	5.	6.	SHENSI
1569			34.6	N	110.3	E	5.	6.	SHENSI
1573	1	10	34.4	N	104.0	E	6.75	8.5	KANSU
1578	7	17	40.4	N	115.7	E	5.	6.	HOPEH
1580	9	5	39.5	N	112.3	E	5.5	7.	SHANSI
1581	5	18	39.8	N	114.5	E	6.	7.5	HOPEH
1581	7		33.0	N	104.6	E	5.5	7.	KANSU
1582	3		40.1	N	113.2	E	5.	6.	SHANSI
1583	5	18	39.7	N	113.8	E	5.5	7.	SHANSI
1584	3	6	30.8	N	115.7	E	5.5	7.	HUPEH
1584	4		38.4	N	111.9	E	4.75	6.	SHANSI
1585	3	6	31.2	N	117.7	E	6.	7.	ANHWEI
1586	5	26	39.9	N	116.3	E	5.	6.	HOPEH
1587	4	10	35.3	N	113.5	E	6.	7.	HONAN
1587	10	4	35.2	N	110.8	E	5.	6.	SHANSI
1588	8		38.4	N	112.2	E	5.	6.	SHANSI
1590	7		36.5	N	102.7	E	5.	6.	TSINGHAI
1590	7	7	35.4	N	103.5	E	5.5	7.	KANSU
1591			36.6	N	110.2	E	5.	6.	SHENSI
1591	11	21	38.8	N	101.1	E	5.	6.	KANSU
1597	2	14	31.9	N	104.3	E	5.	6.	SZECHWAN
1599			35.6	N	109.2	E	5.	6.	SHENSI
1603	5	30	31.2	N	112.6	E	5.	6.	HUPEH
1604	10	25	34.2	N	105.0	E	6.	7.5	KANSU
1608	9	23	37.5	N	105.7	E	5.5	7.	NINGHSIA
1610	3	9	32.5	N	104.5	E	5.5	7.	SZECHWAN
1614	10	23	37.2	N	112.5	E	6.	7.5	SHANSI
1615	3	1	32.0	N	120.9	E	5.	6.	KIANGSU
1615	7	20	38.8	N	106.3	E	5.5	7.	NINGHSIA
1616	2	10	37.8	N	105.9	E	5.75	7.	NINGHSIA
1616	10	10	40.9	N	116.0	E	5.	6.	HOPEH
1618	5	20	37.0	N	111.9	E	6.5	8.	SHANSI
1618	11	16	39.8	N	114.5	E	6.	7.5	HOPEH
1620	10	19	37.1	N	117.5	E	4.75	6.	SHANTUNG
1621	3		39.4	N	116.8	E	5.5	7.	

1623	12	26	36.0	N	115.1	E	4.75	6.75	HONAN
1624	2	10	32.5	N	119.5	E	6.5	7.0	KIANGSU
1624	3?		38.4	N	112.8	E	5.0	6.0	SHANSI
1624	4	17	39.7	N	118.7	E	6.25	8.0	HOPEH
1624	7	4	35.4	N	105.9	E	6.0	7.5	KANSU
1624	7	19	38.8	N	115.5	E	5.5	7.0	HOPEH
1624	9	1	31.1	N	121.4	E	5.5	7.0	KIANGSU
1624	9		33.2	N	107.5	E	5.5	7.0	SHENSI
1625	4		38.3	N	116.8	E	5.0	6.0	HOPEH
1626	6	28	39.4	N	114.2	E	7.0	9.0	SHANSI
1627	2	16	37.5	N	105.5	E	6.0	8.0	NINGZIA
1627	6?		37.6	N	113.7	E	5.5	7.0	SHANSI
1628	12?		33.0	N	104.6	E	5.5	7.0	KANSU
1629	3		36.0	N	103.9	E	5.0	6.0	KANSU
1630			30.7	N	113.5	E	5.0	6.0	HUPEH
1630	1	16	32.6	N	104.1	E	6.25	8.0	SZECWAN
1630	10	14	30.4	N	113.5	E	5.0	6.0	HUPEH
1631			33.7	N	106.2	E	5.0	6.0	KANSU
1631	7	21	35.3	N	104.3	E	5.0	6.0	KANSU
1631	7		35.5	N	107.8	E	5.0	6.0	KANSU
1633			32.4	N	109.7	E	5.0	6.0	HUPEH
1633			37.3	N	111.8	E	5.0	6.0	SHANSI
1634	1		34.0	N	105.2	E	6.0	7.5	KANSU
1634	3		30.7	N	115.1	E	5.5	6.5	HUPEH
1634	12		35.1	N	107.7	E	5.5	7.0	KANSU
1635	10	26	33.2	N	107.5	E	5.5	7.0	SHENSI
1636			33.1	N	107.0	E	5.5	7.0	SHENSI
1636			37.0	N	108.9	E	4.75	6.0	SHENSI
1638	1		36.6	N	105.7	E	5.5	7.0	NINGZIA
1640	4		34.7	N	112.5	E	5.0	6.0	HONAN
1640	9		30.4	N	114.9	E	5.0	6.0	HUPEH
1641	6	21	34.3	N	105.5	E	4.75	6.0	KANSU
1642			33.0	N	118.4	E	5.0	6.0	KIANGSU
1642	6	30	34.9	N	111.1	E	6.0	8.0	SHANSI
1652	2	10	31.4	N	115.3	E	5.0	6.0	ANHWEI
1652	3	23	31.5	N	116.5	E	6.0	7.5	ANHWEI
1652	8		33.4	N	104.6	E	5.5	7.0	KANSU
1653			33.1	N	107.0	E	5.0	6.0	SHENSI
1654	7	21	34.3	N	105.5	E	7.5	10.0	KANSU
1654	9	15	36.1	N	115.6	E	5.5	7.0	SHANTUNG
1657	4	21	31.5	N	103.7	E	6.0	8.0	SZECWAN
1658	2	3	39.4	N	115.7	E	6.0	7.5	HOPEH
1662			33.4	N	120.1	E	4.75	6.0	KIANGSU
1662	10	11	33.2	N	114.8	E	5.5	7.0	HONAN
1664			38.7	N	112.7	E	5.5	7.0	SHANSI
1664	4	1	39.9	N	116.7	E	4.75	6.0	HOPEH
1665			37.9	N	102.6	E	5.0	6.0	KANSU
1665	4	16	39.9	N	116.7	E	6.5	8.0	HOPEH
1668	7	25	35.3	N	118.6	E	8.5	12.0	SHANTUNG
1671	9		35.3	N	118.0	E	5.0	6.0	SHANTUNG
1673	3	29	31.8	N	117.3	E	5.0	6.0	ANHWEI
1673	10	18	40.5	N	114.1	E	5.5	7.0	SHANSI
1673	10	18	39.0	N	111.0	E	5.5	6.5	SHANSI
1675			34.1	N	114.8	E	5.5	7.0	HONAN
1675	6		34.8	N	111.1	E	4.75	6.0	SHANSI
1675	6		35.6	N	115.9	E	5.0	6.0	SHANTUNG
1677	9		33.4	N	104.6	E	5.5	7.0	KANSU
1678	6?		40.7	N	115.3	E	5.0	6.0	HOPEH
1679	9	2	40.0	N	117.0	E	8.0	11.0	HOPEH
1679	10		37.5	N	112.5	E	5.5	7.0	SHANSI

1683	11	22	38.7	N	112.7	E	7.	9.
1686			37.1	N	106.4	E	5.	6.
1695	5	18	36.0	N	111.5	E	8.	10.
1698			41.5	N	121.2	E	5.	6.
1704	9	18	38.0	N	116.5	E	5.5	7.
1704	9	28	34.9	N	106.8	E	6.	7.5
1708	10	26	36.7	N	114.7	E	5.5	7.
1709	10	14	37.4	N	105.3	E	7.5	9.5
1713	9	4	32.0	N	103.7	E	6.5	8.
1718	6	19	35.0	N	105.2	E	7.5	10.
1720	7	12	40.4	N	115.5	E	6.75	9.
1724			40.4	N	115.2	E	5.	6.
1725	7		30.0	N	102.0	E	5.5	7.
1730	9	30	40.0	N	116.2	E	6.5	8.
1730	12?		36.9	N	117.9	E	5.	6.5
1731	11		31.4	N	121.0	E	5.	6.
1734	3		30.3	N	103.5	E	5.	6.
1737	9	30	35.3	N	113.8	E	5.5	6.
1738	5	19	33.2	N	104.2	E	5.5	7.
1739	1	3	38.9	N	106.5	E	8.	10.
1739	2	13	38.5	N	106.3	E	5.5	7.
1742			32.0	N	110.8	E	5.	6.
1743	6	29	30.7	N	110.4	E	5.5	7.
1746	7	29	40.2	N	116.2	E	5.	6.
1748	2	23	31.3	N	103.4	E	5.5	7.
1748	8	29	30.5	N	101.5	E	5.5	7.
1748	11	21	36.4	N	106.1	E	5.5	7.
1754	5		37.7	N	112.5	E	5.	6.
1765	3	15	41.8	N	123.4	E	5.5	7.
1765	5	1	35.3	N	103.9	E	5.25	7.
1765	9	2	34.8	N	105.0	E	6.5	8.5
1772	3	1	38.3	N	114.4	E	5.	6.
1785	12?		30.5	N	101.5	E	5.	6.
1787	12	13	31.0	N	103.7	E	4.75	6.
1789	11	7	34.6	N	110.3	E	5.	6.
1791	2	11	38.0	N	115.5	E	5.	6.
1792	9	7	30.5	N	101.5	E	6.	8.
1792	11	30	30.5	N	101.5	E	5.5	7.
1793	5	15	30.5	N	101.5	E	6.	7.5
1795	8	5	39.7	N	118.7	E	5.25	6.5
1796	3		36.0	N	119.4	E	5.	6.
1805	6	25	37.1	N	114.5	E	5.	6.
1805	8	5	39.7	N	119.2	E	5.5	7.
1811	9	27	31.7	N	100.3	E	6.	7.5
1812	4	2	34.6	N	110.6	E	5.	6.
1813			36.0	N	111.4	E	5.25	6.5
1814	2	4	35.8	N	114.4	E	5.25	7.
1815	8	6	39.0	N	117.5	E	5.	6.
1815	10	23	34.8	N	111.2	E	6.75	9.
1819	2	24	36.3	N	102.3	E	5.75	7.
1820	8	3	34.1	N	113.9	E	6.	8.
1820	10		34.8	N	111.2	E	5.	6.
1822	6	18	33.0	N	104.6	E	5.5	7.
1823	8		32.5	N	107.9	E	5.	6.
1827	3	23	34.9	N	111.1	E	5.25	6.5
1829	4		37.5	N	111.2	E	5.25	6.5
1829	6	18	34.6	N	110.6	E	5.	6.
1829	11	18	33.2	N	117.9	E	5.5	7.
1829	11	19	36.6	N	118.5	E	6.	8.
1830	6	12	36.2	N	114.2	E	7.5	10.

SHANSI
NINGSIA
SHANSI
LIAONING
HOPEH
SHENSI
HOPEH
NINGSIA
SZECHWAN
KANSU
HOPEH
HOPEH
SZECHWAN
HOPEH
SHANTUNG
KIANGSU
SZECHWAN
HONAN
SZECHWAN
NINGSIA
NINGSIA
HUPEH
ANHWEI
HOPEH
SZECHWAN
SZECHWAN
NINGSIA
SHANSI
LIAONING
KANSU
KANSU
HOPEH
SZECHWAN
SZECHWAN
SHENSI
HOPEH
SZECHWAN
SZECHWAN
SZECHWAN
HOPEH
SHANTUNG
HOPEH
HOPEH
SZECHWAN
HONAN
SHANSI
HONAN
HOPEH
SHANSI
TSINGHAI
HONAN
HONAN
KANSU
SHENSI
SHANSI
SHANSI
HONAN
ANHWEI
SHANTUNG
HOPEH

1000

1000

6000

1831	9	28	35.9	N	117.8	E	5.25	6.
1831			32.8	N	116.9	E	5.	6.
1835	6	6	36.3	N	116.4	E	5.	6.
1837	9	15?	34.6	N	103.7	E	6.	8.
1846			31.6	N	106.0	E	5.5	7.
1847	3		34.8	N	111.8	E	5.	6.
1850			34.7	N	104.9	E	5.	6.
1852	5	26	37.5	N	105.2	E	6.	8.
1855	2	27	30.1	N	120.0	E	5.	6.
1855	12	11	39.1	N	121.7	E	5.5	7.
1856	4	10	39.1	N	121.7	E	5.25	6.5
1859	9	19	40.7	N	122.2	E	5.	6.
1861	7	19	39.1	N	121.7	E	6.	8.
1862	12	23	35.5	N	111.5	E	5.5	7.
1867	9		30.5	N	120.8	E	5.	6.
1868	10	30	32.4	N	117.8	E	5.5	6.5
1879	5	12	31.8	N	104.8	E	5.	6.
1879	7	1	33.2	N	104.7	E	7.5	10.
1880	9	30	39.7	N	119.7	E	5.	6.
1881	7	20	33.6	N	104.6	E	6.5	8.
1882	12	2	38.1	N	115.5	E	6.	8.
1885	1	14	34.5	N	105.7	E	6.	7.5
1885	2	21	40.7	N	122.2	E	5.	6.
1887	7		37.0	N	103.8	E	5.	6.
1888	6	13	38.5	N	119.0	E	7.5	
1888	11	2	37.1	N	104.2	E	6.25	8.
1889	9		38.1	N	106.3	E	5.	6.5
1889	10		36.3	N	115.1	E	5.	6.
1890			36.9	N	112.9	E	5.5	7.
1890	2	17	36.6	N	101.8	E	5.	6.
1891	4	17	37.0	N	111.9	E	5.75	7.
1893	2	23	38.3	N	116.8	E	5.	6.
1893	6	1	36.6	N	101.8	E	5.5	7.
1893	8	29	30.5	N	101.5	E	6.	8.
1898	9	22	39.1	N	113.0	E	5.5	7.
1904	8	30	31.2	N	100.9	E	6.	8.
AAA 1910	1	8	35.0	N	122.0	E	6.75	
GGG 1917	1	24	31.3	N	116.3	E	6.25	8.
IAA 1919	5	29	31.5	N	100.5	E	6.25	
IAA 1919	8	25	32.0	N	100.0	E	6.25	
AAG 1920	12	16	36.5	N	105.7	E	8.5	12.
AAA 1920	12	25	35.6	N	106.3	E	7.	
AAA 1920	12	28	35.5	N	105.5	E	6.25	
IIL 1921	1	6	38.0	N	107.0	E	6.	
IIA 1921	1	7	38.0	N	107.0	E	6.	
IIA 1921	1	7	35.8	N	106.2	E	6.5	
AAA 1921	4	12	31.3	N	100.8	E	7.3	10.
GAG 1923	3	24	33.5	N	121.0	E	6.5	
GGG 1927	2	3	37.6	N	102.6	E	25	8.0
AAG 1927	5	22	37.7	N	102.2	E	6.	11.
AAA 1927	5	23	37.6	N	102.2	E	6.	
AAG 1928	3	7	40.7	N	111.3	E	6.	8.
AAA 1929	1	17	30.1	N	101.8	E	6.	8.
IAA 1932	3	6	31.4	N	115.0	E	6.	8.
AAG 1932	4	6	36.1	N	121.6	E	6.25	
AAG 1932	8	22	32.0	N	103.7	E	7.4	10.
AAG 1933	8	25	41.1	N	108.3	E	6.25	8.
AAG 1934	1	20	33.3	N	101.1	E	6.	
AAG 1935	7	26	35.4	N	103.4	E	6.75	
AAG 1936	2	7	34.2	N	105.7	E	6.	8.
AAG 1936	8	1						

SHANTUNG
ANHWEI
SHANTUNG
KANSU
SZECHWAN
HONAN
KANSU
NINGSIA
CHEKIANG
LIAONING
LIAONING
LIAONING
LIAONING
SHANSI
CHEKIANG
ANHWEI
SZECHWAN
KANSU
HOPEH
KANSU
HOPEH
KANSU
LIAONING
KANSU
SHANTUNG
KANSU
NINGSIA
HOPEH
SHANSI
TSINGHAI
SHANSI
HOPEH
TSINGHAI
SZECHWAN
SHANSI
SZECHWAN
HUANGHAI*
ANHWEI
SZECHWAN
SZECHWAN
NINGSIA
NINGSIA
KANSU
NINGSIA
NINGSIA
NINGSIA
SZECHWAN
HUANGHAI*
KANSU
KANSU
KANSU
INNER M.
SZECHWAN
HUPEH
HUANGHAI*
SZECHWAN
INNER M.
TSINGHAI
KANSU
KANSU

AC
AC G U
AC I
AC I
AC I
ACDGIU
AC I
AC I
C I
AC I
AC I
ACDGIU
AC GIU
ACDGIU
AC I
AC GIU
AC I
AC I
AC GIU
ACDGIU
AC GIU
AC GIU
AC GIU
AC GIU
AC GIU

*

*

*

*

*

AAG	1937	7	31	20	35	43.	35.2	N	115.3	E	6.9	9.	AC	GIU	SHANTUNG	
AAG	1937	9	1	10	41	03.	35.3	N	115.4	E	6.75	8.	AC	GIU	SHANTUNG	
AAA	1938	3	14	05	14	21.	32.3	N	103.6	E	6.		AC	IU	SZECHWAN	
AAG	1941	6	11	23	13	31.	30.1	N	102.5	E	6.		AC	GIU	SZECHWAN	
AAA	1941	10	8	15	24	20.	32.1	N	103.3	E	6.	8.	AC	I	SZECHWAN	
AAG	1944	12	19	14	09	04.	39.7	N	124.3	E	6.75		AC	GIU	LIAONING*	
AAG	1945	9	23	15	34	23.	39.7	N	118.7	E	6.25	8.	AC	GIU	HOPEH	
AAA	1948	5	23	09	13	18.	37.7	N	121.9	E	6.		AC	I	SHANTUNG*	
IIA	1949	6	15	09	42	42.	33.3	N	100.0	E	6.		AC	I	TSINGHAI	
AAA	1952	10	31	23	51	40.	33.3	N	101.0	E	6.		A	IU	TSINGHAI	
AAD	1954	2	11	30	30	15.	39.0	N	101.3	E	7.25	10.	A	D IU	KANSU	
AAA	1954	2	11	04	53	46.	39.0	N	101.5	E	6.		A	IU	KANSU	
AAA	1954	7	31	01	00	00.	38.8	N	104.2	E	7.		A	IU	KANSU	
AAA	1955	4	14	01	29	02.	30.0	N	101.8	E	7.5	9.	A	D IU	SZECHWAN	
III	1955	10	1	06	29	54.	30.0	N	101.4	E	6.		A	IU	SZECHWAN	
AAA	1958	2	7	23	23	36.	31.5	N	104.0	E	6.2		A	IU	SZECHWAN	
AAA	1960	11	9	10	43	42.	32.78N		103.67E		6.75	9.	A	IU	SZECHWAN	
AAA	1966	3	7	21	29	14.	37.35N		114.92E	10	6.8	9.5	A	IU	HOPEH	
AAA	1966	3	22	08	11	36.	37.50N		115.08E		6.7		A	IU	HOPEH	
AAA	1966	3	22	08	19	46.	37.53N		115.05E		7.2	10.	A	IU	HOPEH	
AAA	1966	3	26	15	19	04.	37.60N		115.27E	15	6.2	7.5	A	IU	HOPEH	
AAA	1966	3	29	06	11	59.	37.47N		114.88E	25	6.		A	IU	HOPEH	
AAA	1967	3	27	08	58	20.	38.5	N	116.5	E	30	6.3	7.	A	IU	HOPEH
AAA	1967	8	30	04	22	09.	31.62N		103.33E	8	6.8	9.	A	IU	SZECHWAN	
AAA	1967	8	30	11	08	51.	31.70N		100.33E		6.		A	IU	SZECHWAN	
AAA	1969	7	18	05	24	49.	38.2	N	119.4	E	35	7.4		A	IU	PO HAI *
AAA	1970	2	24	02	07	34.4	30.65N		103.28E	15	6.2	7.5	A	IU	SZECHWAN	
AAA	1973	2	6	10	37	08.3	31.3	N	100.5	E	17	7.9	10.	A	IU	SZECHWAN
AAA	1973	2	7	16	06	27.0	31.5	N	100.3	E	8	6.0		A	IU	SZECHWAN
AAA	1973	8	11	07	15	34.6	32.88N		104.00E	8	6.5		A	IU	SZECHWAN	
YYR	1975	2	4	11	36	05.0	40.65N		122.80E	12	7.3			U	LIAONING	
UBB	1976	4	5	16	54	40.1	40.2	N	112.2	E	6.2			U	INNER M.	
UBB	1976	7	27	19	42	54.6	39.4	N	118.1	E	7.8			U	HOPEH	
UBB	1976	7	27	23	17	31.4	39.2	N	117.9	E	6.2			U	HOPEH	
UBB	1976	7	28	10	45	35.2	39.7	N	118.7	E	7.0			U	HOPEH	
UBR	1976	8	16	14	06	45.9	32.9	N	104.1	E	7.1			U	SZECHWAN	
UBB	1976	8	21	21	49	54.2	32.7	N	104.2	E	6.6			U	SZECHWAN	
UBP	1976	8	23	03	30	07.6	32.5	N	104.1	E	6.6			U	SZECHWAN	
UBL	1976	9	22	20	07	03.2	40.1	N	106.4	E	6.			U	INNER M.	
URL	1976	11	15	13	53	00.6	39.4	N	117.9	E	6.5			U	HOPEH	

Table 3. Number of earthquakes in "Central China" with corrections

<u>Century</u>	<u>Number of Earthquakes</u>	<u>Probability</u>	<u>Corrected Number of Earthquakes</u>
1	1	0.05	21
2	3	0.06	54
3	2	0.05	41
4	3	0.05	56
5	7	0.06	118
6	3	0.07	44
7	4	0.08	50
8	5	0.08	64
9	8	0.07	111
10	2	0.08	26
11	10	0.18	56
12	3	0.37	8
13	4	0.55	7
14	19	0.73	26
15	17	0.91	19
16	57	1.00	57
17	83	0.87	100
18	37	0.96	39
19	54	1.00	54
20	(86)*	1.00	(86)

* There are actually 65 earthquakes from 1900-1976, so we extrapolate it to 100 years.

FIGURE 1. Map showing epicenter distribution of historical earthquakes in China (1177 B.C. to 1899 A.D.). Figure taken from Lee, Wu & Jacobsen (1976, Fig. 3).

FIGURE 2. Map showing epicenter distribution of instrumentally determined earthquakes in China (1901 A.D. to 1976 A.D.). Figure taken from Lee, Wu & Wang (1978, Fig.2).

FIGURE 3. Generalized topography of China.

FIGURE 4. Population density map of modern China.

FIGURE 5. Approximate boundaries of Chou, Han, Tang, and Ching dynasties. Modified from Fessler (1963).

FIGURE 6. Provinces of China. The boxed area is "Central China".

FIGURE 7. Apparent population in China according to taxation census.

FIGURE 8. Probability function according to apparent population.

FIGURE 9. Probability function of document survival.

FIGURE 10. Probability function that an earthquake occurring in time t will appear in our catalog.

FIGURE 11. Number and corrected number of earthquakes per century in "Central China".

FIGURE 12. Number of earthquakes in a moving window of 50 years.

FIGURE 13. Corrected number of earthquakes in a moving window of 50 years.

FIGURE 14. Autointensity function for the earthquake sequence in "Central China" from 1000 A.D. to 1976 A.D.

SEISMIC GAPS AND PLATE TECTONICS: SEISMIC POTENTIAL
FOR MAJOR PLATE BOUNDARIES

By William R. McCann^{2,3}), Stuart P. Nishenko^{2,3}),
Lynn R. Sykes^{2,3}) and Janet Krause^{2,3})

Abstract

The theory of plate tectonics provides a basic framework for evaluating the potential for future great earthquakes to occur along major plate boundaries. Along the transform and convergent plate boundaries considered in this paper, the majority of seismic slip occurs during large earthquakes, i.e., those of magnitude 7 or greater. In general, rupture zones, as delineated by aftershocks, tend to abut rather than overlap, and large events occur in regions with histories of both long and short-term seismic quiescence. These techniques are used in this paper to delineate major seismic gaps.

In detail, however, the distribution of large shallow earthquakes along convergent plate margins is not always consistent with a simple model derived from plate tectonics. Certain plate boundaries, for example, appear in the long term to be nearly aseismic with respect to large earthquakes. The identification of specific tectonic regimes, as defined by dip of the

² Lamont-Doherty Geological Observatory of Columbia University, Palisades, New York 10964

³ Also, Department of Geological Sciences, Columbia University

inclined seismic zone, the presence or absence of aseismic ridges and seamounts on the downgoing lithospheric plate, the age contrast between the overthrust and underthrust plates, and the presence or absence of back-arc spreading, have led to a refinement in the application of plate tectonic theory to the evaluation of seismic potential.

The term seismic gap is taken to refer to any region along an active plate boundary that has not experienced a large thrust or strike-slip earthquake for more than 30 years. A region of high seismic potential is a seismic gap that, for historic or tectonic reasons, is considered likely to produce a large shock during the next few decades. The seismic gap technique provides estimates of the location, size of future events and origin time to within a few tens of years at best.

The accompanying map summarizes six categories of seismic potential for major plate boundaries in and around the margins of the Pacific Ocean and the Caribbean, South Sandwich and Sunda (Indonesia) regions for the next few decades. These categories range from what we consider high to low potential for being the site of large earthquakes during that period of time. Categories 1, 2 and 6 define a time-dependent potential based on the amount of time elapsed since the last large earthquake. The remaining categories, 3, 4, and 5, are used for areas that have ambiguous histories for large earthquakes; their seismic potential is inferred from various tectonic criteria. These six categories are meant to be interpreted as forecasts of the location and size of future large shocks and should not be considered to be predictions in which a precise estimate of the time of occurrence is specified.

Several of the segments of major plate boundaries that are assigned the highest potential, i.e., category 1, are located along continental margins,

adjacent to centers of population. Several of them are several hundred kilometers long. High priority should be given to instrumenting and studying several of these major seismic gaps since many are now poorly instrumented. The categories of potential assigned here provide a rationale for assigning priorities for instrumentation, for future studies aimed at predicting large earthquakes and for making estimates of tsunami potential.

1. Introduction

This paper attempts to review and synthesize information about the potential of certain parts of plate boundaries to rupture in future large earthquakes. Over the last ten years it has become clear that some segments of major plate boundaries that have not been the sites of large earthquakes for tens to hundreds of years, i.e., have been seismic gaps for large shocks, are more likely to be sites of future large shocks than segments that have experienced major rupture during, say, the last 30 years. FEDOTOV (1965), MOGI (1968), TOBIN and SYKES (1968), SYKES (1971), KELLEHER (1972), KELLEHER et al. (1973) and others delineate major seismic gaps for several of the major plate boundaries of the circum-Pacific and Caribbean regions. Several of these gaps have since been "filled in" by rupture during large earthquakes.

The seismic gap technique provides estimates of the location and maximum possible size of shocks, but it does not provide an estimate to better than tens of years of the time of occurrence of future large shocks. In this paper the word "forecast" is used for estimates of location and size (magnitude or seismic moment) of large shocks by the method of seismic gaps. The term "prediction" is reserved for estimates that involve a more precise calculation of the time of occurrence and probability as well as the size and location.

Although the concept of seismic gaps was applied by FEDOTOV (1965) to the active Kuril-Kamchatka seismic zone prior to the advent of plate tectonics, the plate tectonic model provides a foundation for understanding the build up and release of strain energy in large shocks and for understanding why the gap hypothesis works as well as it appears to. Most of the world's earthquakes occur along rather well defined belts that coincide with plate boundaries. Relatively little deformation occurs in most of the interiors of plates.

Most of the world's great (magnitude, $M \geq 7 \frac{3}{4}$) earthquakes (category a of GUTENBERG and RICHTER, 1954) occur at shallow depths (depths less than about 40 km) along plate boundaries of the thrust (convergent) type where subduction of one lithospheric plate is presently occurring. Most of the earth's other great shocks occur along long transform faults where relative plate motion involves strike-slip faulting. Along these two types of seismic zones, the slip during earthquakes reflects the relative motion of two interacting lithospheric plates. Active ridge crests where sea-floor spreading is occurring, however, do not appear to be sites of shocks larger than about magnitude 7. Along ridge crests, short transform faults, and along those parts of transform faults near spreading centers the lithosphere is young, hot, weak, and thin, and the area capable of releasing seismic energy is apparently insufficient for earthquakes larger than about magnitude 7.

This study is restricted to large shallow shocks along simple plate boundaries of the thrust or transform type. It attempts to include many (but not all) of the zones of this type in and around the margins of the Pacific Ocean as well as the major plate boundaries of the Caribbean, South Sandwich, and Sunda (Indonesian) regions. The zone of complex, multibranched deformation extending from the Mediterranean across the Middle East, Central Asia, and China is not covered here.

In this paper the term "large" is used in a general sense for shocks of magnitude 7 or greater. The term "major" is employed for $7 \leq M \leq 7 \frac{3}{4}$ (class b of GUTENBERG and RICHTER, 1954). Great shocks are those of $M \geq 7 \frac{3}{4}$.

Most of the cumulative seismic energy release, cumulative seismic moment or cumulative seismic slip along major plate boundaries occurs in large or great earthquakes (BRUNE, 1968). Shocks of magnitude smaller than 7 may be regarded as noise or as accessory to the major plate movement, which occurs

mainly in conjunction with large earthquakes or which may occur in part by a seismic deformation. The strain energy that is released in large shallow shocks is believed to be built up slowly along simple plate boundaries for tens to hundreds of years. This strain comes from the movement of plates, which varies from about 2 to 12 cm/year for the major plate boundaries discussed here. Friction along plate boundaries prevents many major seismic zones from moving continuously on a scale shorter than tens to hundreds of years. Once stresses build up to a critical level, the plate interface moves suddenly about 1 to 20 m during the rupture associated with a large earthquake.

Plate tectonic theory indicates that plate boundaries are continuous and that they do not end suddenly unless the motion is transformed into some other type of tectonic feature such as a spreading ridge, subduction zone or transform fault. Hence, parts of a plate boundary that have not experienced large earthquakes for tens to hundreds of years are likely to be either: 1) sites of future large shocks, or 2) regions where plate movement is accommodated either aseismically or by the occurrence of only small to moderate-size shocks. In the long term, which is here taken to be thousands of years, plate movement is thought to be fairly even along the entire length of a simple plate boundary. For more complex, multibranched zones of deformation as in Central Asia, western China or Central Alaska, strain build up may occur less regularly and in a more complex manner which is less suitable for forecasts based on the concepts of seismic gaps.

While the concept of a seismic gap probably is applicable to shocks smaller than magnitude 7, we do not do so in this paper. The resolution in defining rupture zones from either the distribution of teleseismically-located aftershocks, felt reports or tsunami data is no better than a few tens of kilometers. Hence, the size of the smallest rupture zones that can be defined with confi-

dence from data of those types is equivalent to an event of magnitude near 7. In areas where surface breakage is common or a local network has existed for a long time (as along parts of the San Jacinto fault zone in southern California), seismic gaps can be defined for moderate-size earthquakes.

In subduction zones great earthquakes and at least some major shocks appear to rupture that portion of the interface between the two plates from the surface to a depth of about 40 km DAVIES and HOUSE (1979). In great events like the 1964 Alaskan and 1960 Chilean earthquakes, aftershocks extend from the surface at a point near the trench to a depth of about 40 km beneath the arc. Below this depth fault plane solutions of the thrusting type are less common; many of the solutions exhibit either down-dip tension or compression parallel to the downgoing lithospheric slab. They are inferred to represent stresses in the interior of the downgoing plate (ISACKS and MOLNAR, 1971). This transition from intraplate to interplate motion may not be sharp and the thickness of the lithosphere varies among the methods used to define it. One common assumption that is often made in analyses of seismic gaps is that large shocks rupture the entire plate boundary in depth. There are, nonetheless, a few examples where the deeper part of the plate boundary ruptures during a shock (usually a major and not a great event) and the shallow portion ruptures in a second later event. Rupture in great shocks is commonly initiated at a depth of about 40 km and usually near one of the ends of a seismic gap. Rupture then propagates both up dip and along the strike (KELLEHER et al., 1973) of the interface.

As pointed out later in the text, the forecasts made here are for a specific set of assumptions, types of plate boundaries and range of magnitudes. The paper does not attempt to forecast events smaller than magnitude 7, large shocks of intermediate depth, earthquakes involving normal faulting in the

bottom or along the outer (seaward) walls of deep-sea trenches, events adjacent to a nearby plate boundary as in Central Alaska, shocks along a multiple plate boundary (for example, along the Hayward fault in the San Francisco Bay area adjacent to the San Andreas fault), or seismic events along the set of grabens that are often found along the chain of volcanoes near subduction zones. Large shocks are less common in those areas than they are along the main plate boundaries. Nevertheless, earthquakes of those types, including ones of only moderate size can be destructive when they occur near centers of population.

If an area is identified as a seismic gap in this paper, this means that the region in question has not been ruptured by a large shock in the last 30 years or more. This identification should not be interpreted to mean, however, that the site necessarily has a high potential for being the location of a large shock during the next few decades. Other additional information, such as the seismic history or tectonic regime, are needed before the area can be designated as one of high seismic potential. In this paper major plate boundaries are classified into six categories of seismic potential. Areas along plate boundaries of lowest assigned potential for the next few tens of years (category 6) are those that ruptured in large shocks during the last 30 years. Of course, after many decades regions now assigned to category 6 will have a higher potential. Gaps assigned the highest potential (category 1) are those in which a great event is known to have occurred in the area but not within the last 100 years. Gaps that were the sites of large shocks more than 30 but less than 100 years ago (category 2) also appear to have a high potential for the occurrence of future large shocks. Many of the latter gaps, however, probably will not be sites of large shocks for the next few tens of years since a long time is needed to build up high tectonic stresses. In some places where the repeat time appears to be about 40 years as in parts of

Middle America, however, large shocks may occur in areas designated as category 2 within the next few tens of years. Hence, for categories 1, 2 and 6 the potential for large shocks within the next few decades is fairly clear.

The potential for large shocks is not as clear for the regions designated in categories 3, 4 and 5. These areas have either an ambiguous history of great earthquakes or no record of large shocks. For categories 3, 4 and 5 we resort to assessing the potential by comparing them with other regions that appear to have a similar tectonic setting and, we presume, a similar seismic potential. It should be remembered, however, that designations in these three categories are more hypothetical and that much work remains to be done in checking the validity of the bases for those hypotheses. It does appear, nonetheless, that certain parts of plate boundaries are not typified by the occurrence of great shocks and that they probably will remain areas of low or negligible potential for thousands of years.

In the future it may be possible to parameterize the seismic potential according to more physically understandable quantities such as the long-term rate of plate motion, repeat time of large shocks or the configuration of the interface between the two interacting plates. At the present time the uncertainty in these quantities, especially in repeat time, is large enough that we have simply used cutoffs of 30 and 100 years in assigning regions to specific categories.

We find a relatively small number of regions that are assigned the highest seismic potential, i.e., category 1. In many of those areas, however, local seismic networks and strong-motion instruments are either non-existent or nearly so. Many of these gaps are located largely in submarine areas adjacent to land masses. These gaps clearly deserve high priority for instrumentation and intensive study both on land and at sea.

2. Map of Seismic Potential for Major Plate Boundaries

(a) Six Categories of Potential

Figure 1 displays our conclusions about the relative seismic potential for the major plate boundaries of the Pacific, Caribbean, Indonesian and South Sandwich areas. Before discussing these categories, it is important to remember that the forecasts are made subject to several assumptions and limitations which are as follows:

1. only shallow earthquakes of magnitude 7 or greater are considered or forecast;
2. some future events of magnitude near 7 may fail to be forecast because the error in mapping aftershock zones and rupture zones is comparable to the size of the rupture zone of an event near magnitude 7;
3. earthquakes of intermediate and deep focus are not considered;
4. only simple plate boundaries of the thrust (convergent) or transform type are considered;
5. events near a major plate boundary as in central Alaska or along the grabens and volcanic zone of Central America are excluded;
6. shocks involving normal faulting of the down-bent lithosphere along the inner walls of trenches or in the bottoms of deep-sea trenches are excluded;
7. zones of multibranch deformation (i.e., 2 or more major subparallel faults) are not considered;

Figure 1

Seismic potential for events with $M \geq 7$ for the next few decades along certain major plate boundaries. The six categories presented are based on historic and tectonic criteria. Shaded areas are those portions of plate boundaries about which we have the most data, historic and/or tectonic, and hence the most confidence in our evaluation. Dark areas (category 1) have not ruptured in a great earthquake in over 100 years, and are considered likely candidates for major or great shocks within the next decade or few decades. Similarly, lightly shaded areas (category 6) have ruptured in a great earthquake within the past 30 years, and are considered to have the lowest seismic potential for the present time. Some regions that are not shaded (categories 3, 5) are areas where more information is needed in order to more accurately assess their seismic potential. Areas that are cross-hatched (category 4) are characterized by plate motion subparallel to the arc, and appear to be tectonically similar. The seismic potentials presented on this map are meant as general forecasts, not specific predictions of the time of occurrence.

8. large shocks are assumed to rupture the plate boundary from about 0 to 40 km in depth;
9. a second large event cannot occur along the same plate interface for many decades after a large earthquake until the stress is slowly built up again by plate movements;
10. the rupture zone is accurately reflected by the extent of the aftershock zone, by the extent of the area of intense shaking or damage, or by the area of tsunami generation. One case is known (NANKAIDO, JAPAN 1946) in which the rupture zone is larger than the area inferred from aftershocks;
11. the forecasts are valid only for the next few tens of years;
12. the designation of high seismic potential is not a prediction in the sense that a precise origin time of a future shock is estimated. Each segment of these plate boundaries is assigned to one of six categories.

These categories are designed to reflect our assessment of: (1) the relative seismic potential of the region; (2) the completeness of its seismic history; (3) our understanding of the tectonic regime. The areas thought to be of the highest potential are assigned to category 1; those of successively lower potential are assigned higher numbers up to six. The categories of seismic potential are as follows:

1. The region (portion of a plate boundary) has experienced at least one large shock in the historic past with the most recent event occurring prior to 1879, i.e., more than 100 years ago. This category represents the highest seismic potential.
2. The region has experienced at least one large shock in the past with the most recent event occurring between 1879 and 1949, i.e., more than 30 years ago, but less than 100 years ago.

3. The region has an incomplete history of large earthquakes. No historic event is clearly documented as having ruptured the plate boundary. There is no evidence, however that would indicate that the region may not be the site of a future large earthquake. A comparison of the tectonic framework with that of other areas known to be sites of historic large shocks may also suggest that the region is capable of being the site of a future large shock.
4. Motion between the plates is parallel or nearly parallel to the local strike of the subduction zone (trench). This category applies to the Puerto Rico-Virgin Island region, the Commander Islands in the westernmost Aleutians, and the Andaman-Nicobar region in the Indian Ocean. All appear to have a similar tectonic setting. A resolution of the question of seismic potential for one area may be useful in assessing the potential for the other two.
5. The region does not have a history of great earthquakes. Several tectonic hypotheses, which are proposed by various investigators, suggest that these regions will not be the sites of great shocks in the future.
6. The region has been ruptured by a large earthquake during the last 30 years (since January 1, 1949). This category is considered to represent the lowest seismic potential for the next few decades.

Regions assigned to category 1 or 6 are from our most clearly defined data set. In each case the regions have clearly ruptured during a large quake. Those in category 1 have been accumulating strain energy for at least 100 years. They are believed to be highly stressed and should be considered likely sites for great earthquakes in the next few to few tens of years.

We have used 100 years as an arbitrary cutoff for category 1. Repeat times of large shocks, even in areas with a long history, vary enough that they do not appear to be useful for forecasts of time of occurrence to better than a factor of about 1.5 to 2. The repeat time might be expected to be a function of the dip of the interface between plates, the age of the interacting lithosphere and the rate of relative plate motion. As yet these factors have not been taken into consideration in a quantitative sense. Most of the plate boundaries considered here have relative movements between 5 and 12 cm/year. Repeat times might be expected to be longer for boundaries with slow relative motion, such as the Caribbean plate where the relative motion is about 2 cm/year. The motion of the Philippine Sea plate relative to surrounding plates is also poorly known. Nevertheless, since the repeat time is poorly determined for most areas, some regions in category 1 may still not be sites of large shocks for many years to come.

Category 4 denotes regions that have relative motion between plates that is parallel or nearly parallel to the axis of the trench. Only three areas are placed in this category. None of these regions (Puerto Rico-Virgin Islands, Commander Islands, Andaman-Nicobar Islands) has experienced a great shallow earthquake during this century that has ruptured an extensive portion of the plate boundary. A major or great earthquake may have occurred in the Commander Islands during the 19th century. The 400 year history of Puerto Rico, however, yields no record of a great event even though shocks of about magnitude 7 have occurred several times. The seismic potential for great earthquakes in these regions should be considered poorly constrained; major events are more likely. The physical mode of plate motion is also not well understood for these three areas. Focal mechanism solutions indicate thrust faulting along nearly horizontal planes with the slip vector nearly parallel

to the trench (ISACKS, OLIVER and SYKES, 1968; MOLNAR and SYKES, 1969; CORMIER, 1975).

Category 2 contains regions that experienced a large shock more than 30 but less than 100 years ago. Some of these regions may be the sites of large shocks in the next few years to tens of years as their recurrence times may be as short as 30 to 50 years (i.e., Central America, Solomon Islands).

Category 3 represents those regions for which the historic record appears to be incomplete. There is no clear and unambiguous indication of the past occurrence of large shocks. The record of large earthquakes in these regions is such that historic reports are absent, too short in length, or yet to be researched. They may have a seismic potential either in categories 1, 2 or 5 when more analysis is performed.

Category 5 also includes regions that have not been the site of a great earthquake in recorded history. In each case there are reasons for inferring that the area may never be the site of a great shock. These inferences stem from either the presence of a tectonic regime along the plate boundary that is thought to be the site of only occasional large shocks or the region is thought to be experiencing some modification of the subduction process such that large shallow shocks are rare or absent. A reduction in the size and number of large shocks is commonly observed in many regions where aseismic ridges and other bathymetric highs on the seafloor encounter subduction zones (KELLEHER and McCANN, 1976). Category 5 includes subduction zones where the density contrast between the two interacting plates is thought to be unusually large, and the plates may be moving aseismically (McCANN and NISHENKO, 1978; FRANKEL and McCANN, 1979). These regions correspond, in general, with the "Mariana type" of subduction of UYEDA and KANAMORI (1978). The density contrast hypothesis, the decoupling idea of KANAMORI (1971, 1977a), and the inference

that the presence or absence of back-arc spreading plays an important role in the state of stress and mode of occurrence of large earthquakes need further investigation. Therefore, assignment of regions to category 5 should be considered temporary pending more detailed research on these hypotheses and on their possible role in governing the occurrence of large earthquakes. In several areas, like the Marianas, each of these hypotheses predicts that few if any great shocks are to be expected in the future.

(b) Other Phenomena Shown on Map of Seismic Potential

Tsunami. In Figure 1 wavy lines are used to denote a possible tsunami risk in those regions in category 1 that have experienced an historic, destructive tsunami. This is not to indicate that these areas will have a destructive tsunami associated with the next great shock but only that a potential exists. Conversely, those regions on convergent plate boundaries with no tsunami potential indicated, nevertheless, may produce tsunamis during future large earthquakes.

Focal Mechanism. Typical focal mechanisms are presented (lower hemisphere projection, where shaded areas denote compressions) for shallow earthquakes along some of the major plate boundaries in Figure 1. These solutions indicate the probable sense of motion during the next large earthquake on a particular plate boundary.

Volcanoes. Triangles are shown in Figure 1 where active volcanoes occur in conjunction with the subduction process. These volcanoes may show fluctuations in activity before the next nearby great earthquake as discussed

by NAKAMURA (1975), CARR (1977), and KIMURA and McCANN (1979).

Successful Forecasts. Stars are placed in Figure 1 on those segments of plate boundaries that experienced large shocks after the region was cited in the literature as a major seismic gap. Some of the successful forecasts are indicated in Table 1.

TABLE 1

FORECAST		EARTHQUAKE		
Investigator	Year Published	Location	Year	Magnitude
Fedotov	1965	near Hokkaido, Japan (Tokachi-oki)	1968	8.2 M_w
Fedotov	1965	near southern Kuril Islands	1969	8.2 M_w
Fedotov	1965	near central Kamchatka	1971	7.8 M_s
Fedotov Mogi	1965 1968	near Hokkaido, Japan (Nemuro-oki)	1973	7.7 M_s
Sykes	1971	near Sitka southeastern Alaska	1972	7.6 M_s
Kelleher	1972	near Lima, Peru	1974	8.1 M_w
Kelleher & Others	1973	near Colima, Mexico	1973	7.5 M_s

3. Historic Review and Basic Concepts

(a) Initial Research on Seismic Gaps

The first work on seismic gaps appeared in 1965 by S. A. FEDOTOV (Fig. 2). His study covered a portion of the seismically active margin of the northwest Pacific extending from central Japan to the Kamchatka Peninsula near the Commander Islands. His examination of the occurrence of great ($M_s \geq 7 \frac{3}{4}$), shallow earthquakes led him to conclude that great earthquakes tend to occur in regions that were not the sites of great earthquakes for at least several decades. He proposed that the rupture zones of large shocks were delineated by aftershock activity and observed that the aftershock zones of nearby large earthquakes tend to abut without overlapping. FEDOTOV (1965) also specifically mentioned several areas that are probable locations for future large earthquakes. Since 1965 several of the seismic gaps he delineated near the southern Kuril Islands and central Kamchatka were "filled" by the occurrence of recent large earthquakes (see Table 1).

Fig. 3 (from MOGI, 1968 and SYKES, 1971) shows an example of how aftershock zones near Japan tend to abut rather than overlap. Double circles are locations of main shocks; single circles represent aftershocks. The 1968 and 1952 zones abut, as do the 1968 and 1897 zones. The 1968 and 1952 zones were seismic gaps since the turn of the century. The 1968 zone was specified as a gap by FEDOTOV (1965). The 1933 Sanriku event was not associated with underthrusting beneath the Japanese arc but involved normal faulting in the deepest part of the Japan trench (KANAMORI, 1971).

Figure 2

Focal regions of great ($M \geq 7 \frac{3}{4}$) earthquakes in the Kuril-Kamchatka region from 1904 to 1963 and the probable areas of future earthquakes of that strength (from FEDOTOV, 1965).

- 1 - instrumental epicenters of great earthquakes
- 2 - borders of the focal regions of great earthquakes
- 3 - uncertain part of the boundary or possible variations in the outline of the focal region
- 4 - possible focal region
- 5 - proposed (probable) focal regions of the strongest earthquakes of the past century; aftershocks of earthquakes
- 6 - May 1, 1915
- 7 - January 30, 1917
- 8 - September 7, 1918
- 9 - February 3, 1923
- 10 - strong foreshocks of the 1923 and 1963 earthquakes
- 11 - outline of the region of tsunami origin
- 12 - the most probable locations of future earthquakes with $M \geq 7 \frac{3}{4}$
- 13 - less likely locations of future earthquakes with $M \geq 7 \frac{3}{4}$
- 14 - distance along the Kuril-Kamchatka trench
- 15 - axes of the trench
- 16 - axes of the Kuril-Kamchatka volcanic belt

Figure 3

Large earthquakes near northern Japan (after MOGI, 1968 and SYKES, 1971). Double circles are epicenters of main shocks, single circles are aftershocks which define the rupture area (solid line). The dashed line marks the zone affected by a series of shocks occurring between 1897-1901. No reliable instrumental data are available for these events. Note how the rupture areas of the 1968 and 1952 zones abut as do the 1968 zone and that inferred for the 1897-1901 events. The 1933 Sanriku event is not related to the underthrusting of seafloor beneath the Japanese arc.

ALLEN et al. (1965), TOBIN and SYKES (1968) and MOGI (1968, 1969) also discuss ideas similar to those proposed by FEDOTOV. In these papers, however, smaller magnitude events are considered. They find that small-scale earthquake activity is generally very low for several years, perhaps decades, before a large event occurs in a seismic gap. Thus, seismic gaps often appear to be regions of reduced activity for moderate and small shocks as well as regions of relative quiescence for large shocks. MOGI (1969) notes that activity often increases in a ring-shaped region immediately surrounding a quiescent zone, that coincides with the rupture zone of its coming large shock. Together these form a "donut" pattern of small and moderate earthquakes. The level of seismic activity before great earthquakes is examined in detail by KELLEHER and SAVINO (1975); their conclusions generally confirm the observations of low activity in areas that later rupture in large earthquakes. They do find, however, forerunning activity near the epicenter of the coming shock and in the region surrounding its rupture zone as found by MOGI (1969).

KELLEHER (1970) examined the spatio-temporal relationship of large earthquakes in the Alaska-Aleutian area, among others. Several areas exhibited migrations of activity along their seismic zones. In some cases the main shock of a large earthquake occurred at one end of the aftershock zone and fault rupture progressed unilaterally along the seismic zone. In many cases the main shock occurred near the aftershock zone of a former large event and rupture propagated away from that aftershock area.

SYKES (1971) examined the seismic history of the Alaska-Aleutian seismic zone using plate tectonic theory. Aftershocks of large events were relocated by computer to increase the location accuracy. He also finds that aftershock

zones of large earthquakes do not overlap but tend to abut and that large events tend to occur in regions that were quiescent for several decades prior to the main shock. In these nearly quiescent zones strain energy, which appears to result from the large-scale motion of rigid lithospheric plates, has been accumulating for tens to hundreds of years prior to the occurrence of a large shock. If the plates are to maintain their rigidity on a time scale of thousands of years, large events must occur in these gaps and release the strain energy that is stored.

(b) Criteria for Determining Seismic Gaps

Expanding on the work of SYKES (1971) and KELLEHER (1972) for individual subduction zones, KELLEHER et al. (1973) analyzed seismic zones along several regions of plate convergence (subduction) and transform faulting. Their study included the northern, eastern and western margins of the Pacific and most of the border of the Caribbean plate. It was directed solely at large, shallow, interplate events, i.e., shocks in which rupture involves the relative motion of two lithospheric plates.

The initial criteria used by KELLEHER et al. (1973) to determine a seismic gap were:

1. "The segment is part of a major seismic belt characterized predominantly by strike-slip or thrust faulting."
2. "The segment has not ruptured for at least 30 years."

These criteria appear to be very simple, but experience since 1973 shows that they are very effective tools for determining possible sites of future large events. Several segments of the plate boundaries studied by KELLEHER

et al. (1973), however, have no historic record of ever being affected by a great thrust earthquake. Hence, either the historic quiescence is representative of a long-term stationary process or it is a temporary phenomenon in which the repeat times of large shocks exceed the length of the historic record. It is necessary, then, to distinguish among different types of seismic gaps; supplementary criteria must be developed to indicate whether a given gap either has a record of previous large shocks or some other evidence indicates that large events could be expected to occur there.

These supplementary criteria include:

1. an historic record of one or more large earthquakes for the region in question;
2. the recurrence interval for large events in a given segment is nearly equal to the time interval since the last large event and hence a large event could be expected;
3. the site appears to be the next event in a series of earthquakes in a regular space-time progression of large events along a plate boundary; and
4. the region is tectonically similar to other areas in which large shocks either have or have not occurred in the historic record.

As will be discussed later, supplementary criterion 1 is not trivial, as several regions not meeting this criterion appear to be similar tectonically. Some of these regions probably will be the source regions of future great earthquakes, while others may be permanently aseismic for large shocks. Some possible tectonic criteria are developed later in an attempt to distinguish these two types of gaps.

(c) Sizes of Future Shocks and the Geometry of the Subduction Zone

KELLEHER et al. (1974) find that for many of the subduction zones of the Pacific the source dimensions of the large shallow earthquakes have a characteristic maximum length. This characteristic maximum appears to be strongly influenced by the geometry of the interface zone between the two interacting plates. It appears to be particularly influenced by the down-dip width of the interface between the overthrust and underthrust slabs of lithosphere. KELLEHER et al. (1974) find that variations in the inferred width, w , of the interface, as defined by the zone of shallow (< 70 km) hypocenters, correlate with source size (magnitude and dimensions) of many large shallow earthquakes of this century. Regions with wide interfaces ($w \geq 100$ km) are the source areas for great thrust earthquakes in which the maximum lengths of ruptures along a subduction zone may exceed 400 km. Along transform fault zones this interface is nearly vertical and interplate seismic slip extends to a depth of about 15 km. Along subduction zones the interface has a shallow dip and appears to extend to a depth of about 40 km (rather than 70 km as assumed by KELLEHER et al., 1974). This depth is ascertained from the depths of aftershocks of great earthquakes and the transition from interplate thrust faulting found at shallower depths to deformation within the downgoing plate at greater depths (DAVIES and HOUSE, 1979).

Fig. 4a presents a diagrammatic model of a subduction zone. Oceanic lithosphere (lower plate) approaches the trench from the upper left-hand corner of the diagram. Upon entering the trench region the oceanic plate

Figure 4a

Typical rupture propagation during a large thrust earthquake along a convergent plate boundary. Hypocenter of the main shock (largest hatched star) is landward and usually deeper than most aftershocks (smaller hatched stars). Rupture propagates upward and outward (solid arrows) along the plate interface (thick black line). For some great earthquakes the rupture propagates hundreds of kilometers laterally along the plate interface (dashed line). Epicentral locations of aftershocks (solid ellipses) define the surface projection of the rupture zone (shaded area).

is deflected or bent downward beneath the overriding plate. Normal faults that develop on the oceanic plate result from this deformation. The zone of contact (i.e., the interface) between the overthrust and underthrust plates accumulates energy in the form of elastic strain during the long interval between great earthquakes. When the shear stress across the boundary exceeds the static friction on the fault, rupture occurs. Rupture initiates at the hypocenter of the main shock (largest star pattern), its epicenter (largest ellipse) is the position of the hypocenter projected onto the surface of the overthrust plate. Motion initiated at the hypocenter of the main shock then generally spreads both updip and along the strike of the zone of contact between the two plates. Aftershocks (smaller star patterns and ellipses) occur along the rupture zone during the first few weeks or months after the main shock. As these aftershocks represent readjustments occurring on the ruptured fault surface, the spatial extent of the aftershock activity can be used to infer the configuration of the rupture zone of the main shock (shaded area).

Fig. 4b, from KELLEHER et al. (1974), presents a schematic diagram of the interface between the overthrust and underthrust lithosphere. It illustrates the relationships among the width of interface (w), the width of the zone of shallow earthquakes and the angle of dip (θ) of the shallow portion of the underthrust lithosphere. It can be seen that a wide interface can result from either a low angle of dip or an increased thickness of the overthrust plate of lithosphere.

Differences in the width of the interface are inferred using the epicenters of shallow earthquakes. The width of the zone of shallow earth-

Figure 4b

Interface area between overthrust and underthrust slabs of lithosphere at a subduction zone (from KELLEHER et al., 1974). This is a side view showing relationships among the width of interface, w , the zone of shallow earthquakes and the angle of dip (θ).

quakes is taken to be the distance between the trench axis and the contour defined by hypocenters 70 km deep. The 70 km contour is determined using the locations of hypocenters of both shallow (< 70 km) and intermediate (> 70 km) depth. It is employed here as it was by KELLEHER et al. (1974) since calculated depths are subject to greater errors for foci near 40 km. The upper limit of the zone of plate contact occurs either near the axis of the trench or along the inner wall of the trench. In detail, regional variations in the development of the arc, such as the presence of an accretionary sedimentary wedge, may influence the dimensions of rupture zones. In general, however, the aftershock zones of great earthquakes extend to near the trench axis. This information is an important aid in understanding the generation of large events at thrust boundaries, for if the width of interface is reasonably well known, then estimates of the largest possible shock in any particular gap can be made.

(d) Permanent Gaps

Returning now to supplementary criterion 1, KELLEHER and McCANN (1976, 1977) find that the distribution of large shallow earthquakes along subduction boundaries does not always agree with the distribution pattern that might be predicted from a simple model derived from plate tectonics. For extensive sections of some island arcs large shocks have occurred infrequently or not at all during recorded history. Most of these zones of long-term quiescence (category 5) are nearly coterminous with segments of plate boundaries where groups of seamounts, aseismic ridges or other topographic

features on the underthrust lithospheric slab appear to be interacting with and modifying the subduction process. Besides a decrease in the level of large shocks, other manifestations of a modified subduction process include gaps and offsets in the line of active volcanoes, wide diffuse zones of deformation or abrupt jumps in the plate margin, infrequent shallow earthquakes with thrust-type mechanisms and the diminution or disappearance of hypocenters at intermediate depths. In contrast, nearly all segments of subduction zones landward of smooth, low-lying ocean floor have a history of at least one large shock.

Differences in the seismic and tectonic regimes among various island arcs suggest that the lack or near lack of large shocks in some regions probably results from specific differences in the tectonic framework and not from the brevity of the seismic record. KELLEHER and McCANN (1976) hypothesize that these variations are related to the density contrast between the overthrust and underthrust lithospheric plates. Major topographic rises and seamount provinces, for example, represent regions with lithospheric densities intermediate between those of typical oceanic and continental lithosphere. Buoyant material on the downgoing plate may lead to marked changes (and in the most extreme case possibly even to a cessation) in the subduction process. Variations in geological and geophysical parameters suggest that this buoyant interaction is also capable of producing modifications in the subduction process at deeper levels.

These complexities in the subduction process, which appear to affect the occurrence of large shocks, make it necessary to use the terms gap and seismic potential with care. We use the term seismic gap to refer to any

region along an active plate boundary that has not experienced a large thrust or strike-slip earthquake for at least a minimum period of time, which here is taken to be 30 years. In most regions 30 years is the minimum time between the recurrence of large events; in many active zones the repeat time is 100 to 300 years. The term seismic gap does not imply the potential for a large event to occur in that region. A region of seismic potential, however, is here taken to be a seismic gap that, for tectonic or historic reasons, is thought to be capable of producing large shocks in the future.

(e) Aseismic Slip

The quiescence observed for great earthquakes and the near quiescence for moderate-size shocks is observed along several arc segments that are not presently subducting bathymetric features. Each of these regions subducts relatively old (> 60 m.y.) seafloor beneath a young island arc reared by an active marginal basin. These regions have been assigned to category 5. A large density contrast could be inferred from the observed differences in the ages of the abutting lithospheric plates. This large density contrast may be responsible for a reduction of the stresses necessary for the generation of large earthquakes (McCANN and NISHENKO, 1978; FRANKEL and McCANN, 1979). That is, large density contrasts between the overthrust and underthrust lithospheric plates may result in poor coupling or low normal stresses along the interface, or zone of contact between the plates. Hence, motion between the plates may occur with little or no stick-slip motion, which would result in a reduced number of large shocks. Thus, the full range of density contrasts from large to moderate and small may have a strong influ-

ence on the frictional coupling along the plate interface and on the forces responsible for the generation of large earthquakes.

UYEDA and KANAMORI (1978) propose that the presence or absence of an active marginal basin on the overthrust plate may be responsible for the observed variation in the number of large shocks occurring along various convergent plate boundaries. They delineate two modes of subduction using criteria based on the spreading activity in the region behind the arc. They associate Chilean-type subduction with the occurrence of great, shallow earthquakes, the absence of spreading centers behind the arcs, and little or no aseismic slip on the thrust boundary. Mariana-type subduction is characterized by a lack of great earthquakes along the thrust boundary at shallow depths, and active back-arc spreading, with a substantial portion of the relative motion between the plates possibly occurring aseismically. UYEDA and KANAMORI (1978) propose two possible mechanisms for the presence of marginal basins and the absence of great earthquakes on the thrust boundary.

1. the contact between the converging plates changes, through an evolutionary process, from tight coupling to decoupling;
2. if the subducted slab is anchored at its bottom to the mantle, a retreating overthrust plate would create marginal basins near the trench; an actively converging overthrust plate would not lead to the development of marginal basins.

At present it is not possible to choose among these models of aseismic and seismic behavior. Aseismic slip appears to play a larger role in plate motion than was generally assumed a few years ago. Hence, these models

deserve more attention. The occurrence or non-occurrence of large earthquakes along convergent plate boundaries may be strongly influenced not only by the density contrast between the plates but also by the absolute plate motions and by the mode of interplate coupling. Until we understand why large shocks do not occur along some parts of plate boundaries, it is not possible to accurately forecast whether some areas that have been aseismic for the last 100 to 300 years will remain so in the future.

(f) Basins on the Upper Slopes of Trenches, Deep-Sea Terraces, and Large Earthquakes

Variations in the structures on the inner wall of trenches appear to reflect changes in both the lengths of rupture zones and in the source areas of tsunamis that are associated with large shallow earthquakes (NISHENKO and McCANN, 1979). Crustal deformation that occurs during large earthquakes and related tsunamic activity play important roles in the development and maintenance of topographic features on the inner walls of deep-sea trenches. A regional comparison of the average length of upper slope basins and terraces with the maximum length of earthquake rupture zones shows that longer basins and terraces characteristically occur in regions with larger rupture zones. Examples in Japan, Alaska and the Aleutians clearly show how these topographic features reflect the size and spatial distribution of seismic-tsunamic source areas. In many cases, this relationship may be explained by the coseismic reactivation of structural units on the inner wall of the trench. The two principal areas of

coseismic reactivation are the trench slope break and the frontal arc region. Thus, upper slope basins, deep-sea terraces and other topographic features may serve as indicators of the tectonic regime and seismic-tsunami risk along convergent plate margins. Also, gross features of the seismic regime (i.e., sizes of rupture zones) for at least the last tens of thousands of years appear to be recorded on the inner wall of deep-sea trenches. This gives us a valuable tool for further studying the seismic interactions between plates.

Variations of the dimensions of rupture zones also appear to be influenced by the block-like behavior of the overthrust plate. In many instances, the dimensions of upper slope basins and terraces are nearly equivalent to those of the crustal blocks. Regions with basin or terrace lengths greater than 100 km have histories of large ruptures involving adjacent segments of the arc (i.e., rupture lengths of several hundred kilometers). Regions with basin or terrace dimensions less than 100 km tend to rupture independently of adjacent segments (NISHENKO and McCANN, 1977).

(g) Temporal-Spatial Relationships Between Volcanism and Large Thrust Earthquakes

NAKAMURA (1975) studies the short-term relationship between volcanic eruptions and the occurrence of nearby large earthquakes. He finds that volcanic activity may increase during the years immediately surrounding the time of large shocks. The study of individual volcanoes, however, has not yet been shown to be a useful tool for forecasting the location of large

shocks. CARR (1977) examines regional volcanic activity before several great earthquakes. He observes a marked reduction in volcanic activity about ten years before a main shock. KIMURA and McCANN (1979) relate long-term volcanic activity and its temporal-spatial relationship to great earthquakes of the thrust type. Volcanoes standing landward of converging plates show increased eruptive activity up to 30 years before several main shocks. The volcanic activity, which is often characterized by lava flows or effusion of great amounts of volcanic material, either ceases or markedly decreases after the occurrence of large shocks. In some cases volcanic activity decreases before the earthquake occurs. After the occurrence of the earthquake, volcanic activity in neighboring seismic gaps tends to increase. This relationship between volcanic activity and the occurrence of large shocks is most clearly seen along the arcs of the northern Pacific (e.g., Aleutians, Kamchatka, Kuriles) and suggests that the timing of regional volcanic activity is at least in part controlled by regional tectonic movements associated with the generation of large earthquakes of the thrust type. Hence, changes in volcanic activity should be carefully monitored and studied along some of the plate boundaries we identify as having the greatest seismic potential for the occurrence of future large shocks.

4. Discussion of Seismic Regions

(a) Alaska-Aleutians

The Alaska-Aleutian seismic belt forms part of the northern and eastern boundary of the Pacific plate. Right-lateral strike-slip motion occurs along the Fairweather and Queen Charlotte transform faults as the Pacific plate moves northward relative to the North American Plate (TOBIN and SYKES, 1968). In southern Alaska, the Pacific seafloor underthrusts the Alaska peninsula and Aleutian Islands, but the direction of convergence becomes more oblique to the arc and trench in the central and western Aleutians. In the Commander Islands, at the far western end of the Aleutian chain, the motion between the Pacific and North American plates is nearly parallel to the axis of the trench (CORMIER, 1975).

Figure 5 shows the rupture areas of great earthquakes along the Alaska-Aleutian seismic zone since about 1920. The seismic history of this region is only complete to about the turn of the century. Nevertheless, the instrumental record indicates that the release of seismic energy in the region is dominated by infrequent, great earthquakes that rupture extensive (> 500 km) portions of the plate boundary (KELLEHER, 1970; SYKES, 1971). Three great earthquakes (1957, 1964 and 1965) have recently ruptured most of the zone of underthrusting along the arc. One seismic gap lies at the western end of the Aleutian arc.

No large earthquakes have been recorded on the thrust boundary during this century. A large earthquake did occur in this region in 1849 (MEDVEDEV, 1968) and was accompanied by a locally destructive seawave. An independent source (ELLA, 1890) indicates that this event probably occurred on the south-

Figure 5

Recent large earthquakes of the Alaska-Aleutian seismic zone. Note that the rupture zones (hatched regions) tend to extend for several hundred kilometers. Year and magnitude of the events are shown. Arrows give direction of motion of the Pacific plate relative to North America (MINSTER et al., 1974). Three large gaps are stippled regions. Lines in Gulf of Alaska are 2000 fathom contours.

44

ern side of the island chain as a seawave was observed in the Samoan and New Hebrides Islands in the southwest Pacific about ten hours after the main shock. A second line of evidence also points to a source to the south of the islands. Events occurring on the northern flank of the island arc typically have strike-slip mechanisms with little or no vertical motion (CORMIER, 1975). It is, therefore, unlikely that a large event on the northern flank of the island chain would produce a large seawave capable of being observed in the southwest Pacific. Although we cannot be sure that this event was not one involving normal faulting in the trench, the sketchy historical record does indicate that this region might be capable of storing enough strain energy to generate large destructive earthquakes with recurrence times of 130 years or more. This region is assigned to category 4. This category has been assigned to three regions that have slip vectors nearly parallel to the trench and that show evidence of a downgoing seismic zone. None of these regions can be definitely associated with a great earthquake that ruptured an extensive portion of the plate boundary. At the present time we do not know if these regions which have a similar tectonic style, have the potential for generating major or great shocks. Hence, all three areas are placed in a single category, category 4.

The second gap lies between the 1957 and 1964 rupture zones. Important activity near this gap includes a great earthquake in 1938 (M_w 8.2). The western limit of the aftershock zone of the 1938 event is uncertain, as only a few poorly located events lie to the west of the main shock. The existing aftershock data suggest that only about half of the plate boundary between the 1964 and 1957 shocks ruptured in 1938. Hence, that region appears to be a major gap. This area has been noted by KELLEHER (1970) as the possible location for the next large event in a space-time progression of large events

moving north and then west along the Alaskan coast (1949, 1958, 1964).

Historically only one large shallow event is reported for this region. It occurred in 1788 and was associated with a locally destructive tsunami on Unga Island in the Shumagin Islands and on Sanak Island about 200 km to the southwest (COX and PARARAS-CARYANNIS, 1976). The intensities reported for this event, however, only reached Modified Mercalli intensity V (MEYERS et al., 1976). The intensity and tsunami damage associated with this event are similar to the macroseismic effects of the 1946 earthquake. Although the 1946 event was only moderate in size ($M_s = 7.4$), it was associated with an extremely destructive seawave (EPPLEY, 1965). More thorough documentation of the 1788 event is necessary. Until those data are collected, however, the region between the 1957 and 1938 rupture zones should be considered one of high risk and possibly the site of the next large earthquake in a space-time progression.

Recent work in this region has focused on two earthquakes that occurred in 1974. ARCHAMBEAU (1978) calculated a stress drop of several hundred bars for one of these events. Both earthquakes (m_b 5 1/2 - 6) triggered a strong-motion instrument in the Shumagin Islands. Detailed analysis of the first motions of P-waves and the strong-motion records indicate that the earthquakes were of the thrust type and each event is associated with a stress drop of about 500 bars (HOUSE and BOATWRIGHT, 1979). These high values of stress drop may be indicative of a broad region that is under considerable tectonic stress. ARCHAMBEAU (1978) also reports high stresses for other nearby events in this area. Because the seismic history is unclear, this region has been assigned a potential of category 3.

A third gap lies between the rupture zones of the 1964 and 1958 earthquakes in southern Alaska. This region is near the site of several large

events that occurred around the turn of the century (RICHTER, 1958). TARR and MARTIN (1912) found that uplift in those shocks was restricted to Yakutat Bay. THATCHER and PLAFKER (1977) conclude that the region affected by those shocks was limited in extent, with motion possibly occurring on a small portion of an east-trending thrust fault and a small segment of the Fairweather fault. As no tsunami was recorded outside the immediate area of the events, it is unlikely that they ruptured a great distance along the coast of Alaska. The region affected by these events lies in or near the area ruptured by the 1958 event. The remaining gap, therefore, lies in a region of transition from transform to thrusting motion and has no clear history of being associated with a large earthquake. We have no evidence, however, that would indicate that this region could not be the site of a future large earthquake. Therefore, it is assigned to category 3.

There are other faults near the Alaska-Aleutian plate boundary, such as those of central Alaska and the northwestern extension of the Fairweather fault. These faults have been the sites of large earthquakes in the past and can be expected to be the sites of large shocks in the future. Because these regions do not lie on clearly defined plate boundaries, however, the categories used here may not apply and we do not consider their seismic potential in this paper. Also, to the south of the 1949 Queen Charlotte event the maximum size of instrumentally recorded earthquakes is considerably smaller than that to the north. KELLEHER and SAVINO (1975) relate this change in rupture size to the thinner lithosphere that lies to the south, adjacent to the spreading centers that lie off the Washington-Oregon coast. The region to the north of these spreading centers of thin lithosphere has been the source of large earthquakes (KELLEHER and SAVINO, 1975; GUTENBERG and RICHTER, 1954), however, because of the locally thin lithosphere most of the events in

this region will fall below the threshold of $M_s \geq 7.0$. Thus, we will not consider the area in detail.

(b) California

The San Andreas fault system extends along much of the coast of California (Figure 6). Right-lateral strike-slip motion has been well documented along much of the fault system. Great earthquakes with extensive rupture zones have occurred along two sections of the fault. The San Francisco earthquake of 1906 broke the northern segment of the fault from near Shelter Cove to San Juan Bautista (REID, 1910). The Fort Tejon earthquake in 1857 ruptured from north of Cholame Valley to near San Bernardino in the south (ALLEN, 1968). The fault segment that lies between these rupture zones does not appear to have the potential for being the site of future large shocks as the creep rate along the fault in much of that segment is nearly equal to the rate of plate movement (SAVAGE and BURFORD, 1973).

Historic records for California are not very complete for more than about 100 years. There is no clear evidence of a great earthquake prior to the 1857 event. ALLEN (1968) indicates that events in 1838 near San Francisco are not associated with ground breakage as extensive as that of the 1906 event. JAHNS (1977) recently completed a study of Holocene displacement along the southern portion of the San Andreas fault. Average recurrence times from the two sites studied are rather different: 225 years for one and 160 or 323 years for the other, depending on the interpretation of the smaller offsets at that site. There is much scatter in the observed return intervals themselves, 57 to 255 years. Because the reported recurrence times scatter a great deal, we do not believe that they can be used at

Figure 6

Location and rupture zones for the great earthquakes of 1906 and 1857 that broke the San Andreas fault. Fault segment between these rupture zones is creeping and thus may not be the site of future great earthquakes.

present to provide much control on the times of future large earthquakes. That portion of southern California has not experienced a great earthquake during the last 100 years. Thus, it has been assigned to category 1.

The area of the 1906 rupture, which appears to have extended to the Mendocino fracture zone, is assigned to category 2. The Gorda ridge and the Mendocino fracture zone represent the continuation of the Pacific-North American plate boundary to the northwest of the San Andreas fault system. They have experienced large shocks in the past (GUTENBERG and RICHTER, 1954). Although none of these shocks are as large as the great earthquakes to the south, future major shocks along the eastern parts of these features could be very damaging in northern California. Those features are also assigned to category 2.

The region of southern California to the south of the zone that ruptured in 1857 is characterized by a series of faults trending parallel or sub-parallel to the trend of the San Andreas fault. Some of these fault segments have experienced large shocks, although most sizeable events in the region are in the $6 \leq M \leq 7$ range (THATCHER et al., 1975). The largest size of events may be related to the distance from spreading centers in the Imperial Valley and the northern Gulf of California. This area is not discussed further in this paper.

(c) Middle America

The Middle America seismic zone extends along western Mexico and Central America between 82.5° and 105°W , defining the northeastern boundary of the Cocos plate. Focal mechanism solutions for shallow earthquakes along this zone indicate thrusting of the Cocos plate beneath the Caribbean and North

American plates (MOLNAR and SYKES, 1969; DEAN and DRAKE, 1978) with about a N30°E direction of convergence (CARR, 1974).

Both the seismic gaps and rupture zones of earthquakes of this century are relatively small (100-200 km) compared to the dimensions of rupture zones along the Alaska-Aleutian or Kurile-Kamchatka seismic zones. Northwest of 95°W, the Middle America trench is subducting somewhat younger seafloor than the segment to the southeast. Consequently, the inferred dip of the seismic zone northwest of 95°W is generally shallower than in the southeast (KELLEHER and McCANN, 1976; DEAN and DRAKE, 1978). The Tehuantepec ridge divides these two regions of seafloor and intersects the trench near 95°W. Rupture zones of large earthquakes occurring along the segment of the arc that is subducting young seafloor tend to be somewhat larger than those occurring southeast of 95°W. CARR and STOIBER (1977) suggest that variations in the geometry of the inclined seismic zone influence the spatial location of large earthquakes and other subduction-related phenomena. Along the inner wall of the trench northwest of 95°W, the lengths of fore-arc basins correlate well with the rupture lengths of the largest known shocks. Several basins are bounded by submarine canyons, which may be fault controlled (FISHER, 1961). The alignment of these canyons with other transverse breaks mapped by CARR et al. (1974) further supports Fisher's interpretation.

The Middle America seismic zone has seven major areas that have not experienced shallow earthquakes larger than magnitude 7 for 45 years or more; these seismic gaps are presented in Figure 7. The northwestern end of the Middle America seismic zone last ruptured during 1932-1934. The short recurrence times for this arc, 40-50 years (KELLEHER et al., 1973), indicate that some of these segments have a high potential for being the sites of future large earthquakes. The northwestern end of the arc appears to have

Figure 7

Rupture zones (ellipses) and epicenters (triangles and circles) of large shallow earthquakes (after KELLEHER et al., 1973) and bathymetry (CHASE et al., 1970) along the Middle America arc. Note that six gaps which have earthquake histories have not ruptured for 40 years or more. In contrast, the gap near the intersection of the Tehuantepec ridge has no known history of large shocks. Contours are in fathoms.

that potential and is placed in category 2.

A major gap is located where the Orozco fracture zone and other associated features intersect the trench. These features may be interfering with the normal subduction process; hence, this region may not have rupture zones or recurrence intervals similar to the rest of the arc. A series of major earthquakes occurred in this gap in 1907-1909, 1911 and 1943; however, there are insufficient data to accurately determine the size and precise location of the rupture zones. This area is assigned to category 2. Note that the 1973 Colima earthquake ($M = 7.5$) apparently ruptured the same area as the 1941 event and skipped the adjacent long-term gap to the southeast (REYES *et al.*, 1978).

A third seismic gap, which has recently attracted much attention, is located near the coast of Oaxaca, Mexico. It is assigned to category 2. The last large earthquake in this region occurred in 1928 ($M = 7.9$). A decrease in local seismic activity in mid-1973 is noted by OHTAKE *et al.* (1977). They think the decrease is similar to patterns of seismic quiescence preceding previous large earthquakes. Observations of variations in seismic activity are crucial in efforts to predict earthquakes; however, as GARZA and LOMNITZ (1978) note, these variations need to be evaluated with respect to the long-term seismic history. That is, do changes of this type occur without being followed by large shocks or are they only associated with the occurrence of large earthquakes?

A major long-term seismic gap is located near 95°W , where the Tehuantepec ridge intersects the Middle America trench. This segment has no clear association with large historic earthquakes (KELLEHER *et al.*, 1973). Hence, we conclude that it has a relatively low potential for being the site of a future large earthquake and assign it to category 5.

Three seismic gaps are located southeast of 95°W . The large gap near the coast of El Salvador experienced large events in 1921 ($M = 7.3$) and 1926 ($M = 7.1$). It is not clear that these shocks ruptured the entire gap, however, their magnitudes would suggest that they did not. Prior to those dates, the historic record indicates that the region offshore from El Salvador may have ruptured in two space-time progressions in 1847-1850 and 1898-1902 (CARR and STOIBER, 1977). This gap has been seismically quiescent for shallow events of $M \geq 4$ since at least 1950. Like the Oaxaca gap, this region deserves more attention as a potential site for a future large earthquake. This gap has the potential (category 2) for being the site of one to a few large shocks or a great earthquake of magnitude exceeding 8. A smaller gap (category 2) is also located between the 1950 and 1956 rupture zones in southern Nicaragua and Costa Rica.

The southeastern end of the Middle America trench, near the west coast of Costa Rica, is located in a rather complicated tectonic regime. It is in this region that the Cocos ridge intersects the trench. The collision of this topographic feature with the subduction zone undoubtedly complicates and thus reduces the reliability of any analysis of this area using the techniques of seismic gaps. Large events occurring in 1934 ($M = 7.7$) and 1941 ($M = 7.5$) may have only partially ruptured the plate boundary. Prior to that, the last great earthquake occurred in 1851 (CARR and STOIBER, 1977). Thus, this area has the potential for a future great earthquake and is assigned to category 2.

Several moderate-size earthquakes that have occurred along the grabens and volcanic zone of Central America during the past 400 years have been quite damaging to nearby centers of population. These shocks, however, are not located along the main plate boundary and hence are not included in the forecasts made here. It can be expected that similar shocks will pose a hazard in

the future to cities like Guatemala City and Managua.

(d) South America

The South American seismic zone extends along western South America from 5°N to 46°S , forming the eastern boundary of the Nazca plate. Typical focal mechanism solutions for shallow earthquakes are of the thrust type with the direction of relative plate motion about $\text{N}80^{\circ}\text{E}$ (STAUDER, 1973). The maximum dimensions of rupture zones of large South American earthquakes vary considerably along the strike of the active seismic zone. This large variation correlates strongly with the width of the zone of contact between the overthrust and underthrust plates (KELLEHER *et al.*, 1974), which may be influenced by the segmentation of the underthrust plate (CARR *et al.*, 1974; BARAZANGI and ISACKS, 1976). Variations of the morphology of the inner wall of the trench (KULM *et al.*, 1977) also appear to correlate with the segmentation of the seismic zone as well.

Five regions have been designated as seismic gaps. Many other areas have not ruptured in the last 30 years. Historic records, however, indicate that recurrence intervals along the South American seismic zone tend to be larger than about 60 years (KELLEHER, 1972). The northernmost gap, in northern Ecuador - southern Colombia (Figure 8), includes the northern portion of the rupture area of the 1906 event; the southern portion of this zone ruptured during events in 1942 ($M = 8.3$) and 1958 ($M = 7.8$). The northern portion of the zone is assigned to category 2. If the recurrence interval is as short in the northern part as it is in the southern region (40 to 50 years), then this region is a prime location for a future large earthquake.

An extensive gap exists from 0° to 9°S . Historic records give no indi-

Figure 8

Bathymetry (CHASE et al., 1970), rupture zones of large earthquakes (cross hatching) and active volcanoes (solid triangles) along western South America (from KELLEHER and McCANN, 1976). Note the absence of great shocks near the intersection of the Carnegie ridge complex with Peru-Ecuador. Discontinuities in the distribution of active volcanoes reflect the intersection of ridges with the continent and discontinuities in the already-subducted Nazca plate. Contours are in fathoms.

cation of a great earthquake occurring in this region during the last 400 years (KELLEHER, 1972); hence, this gap is assigned to category 5. Unless the recurrence times for great events in this region are unusually long (more than 400 years), one would not expect any change from the past pattern of relative seismic quiescence for large shocks. This zone corresponds to that portion of the South American trench that is consuming the Carnegie ridge, a large aseismic feature that is thought to have been formed at the Galapagos hot spot. The interaction of both the Carnegie and Nazca ridges with the subduction zone may explain the extremely shallow dip of the seismic zone from 0° to 15°S , as well as the absence of appreciable Neogene volcanism and historic great earthquakes. Major structural changes along the Andes are noted where these ridges intersect the coast (GANSSEER, 1973). The Amotape zone, which forms the boundary between the northwest-trending central Andes and the northeast-trending northern Andes, occurs near the intersection of the Carnegie ridge with the coast. In addition, the Amotape zone may continue eastward into the Amazon depression, which has been suggested to follow a reactivated lineament in the shield (DE LOCZY, 1968).

The region from 9° to 15°S has a history of great earthquakes; however, the extent of rupture is usually limited to 150 km or less (KELLEHER and McCANN, 1976). A previously noted gap near 12°S was the site of a large event in 1974 ($M = 8.1$). The aftershocks of that event nearly fill the seismic gap. The southern boundary of this region, near 15°S , occurs near a major tear in the descending Nazca plate (BARAZANGI and ISACKS, 1976), which may be associated with the collision of the Nazca ridge with the trench. On land, the coastal cordillera of the central Andes and Neogene volcanics terminate abruptly near the intersection of the Nazca ridge and the coast (GANSSEER, 1973).

Another extensive seismic gap is located along the coast of southern Peru

and northern Chile. This region, along the big bend in the South American coastline, has a history of highly destructive earthquakes and tsunamis. This region is delimited by a high concentration of Neogene volcanism and a relatively steeply dipping seismic zone. Along the northern portion of this gap, great earthquakes in 1604 and 1868 completely destroyed Arica and other coastal towns and generated damaging Pacific-wide tsunamis. The only great event known in the southern portion of this gap occurred in 1877 and was also accompanied by a destructive tsunami (LOMNITZ, 1970). The earthquake history for the northern region suggests a recurrence time for great tsunamic earthquakes of about 260 years. Hence, this area may continue to accumulate tectonic strain for many years before a great shock occurs. Nevertheless, the town of Arica has been destroyed five times in the historic record (LOMNITZ, 1970), indicating the destructive potential of large earthquakes in this region as well. This region is assigned to category 1 based on the long time interval since the last great events in 1868 and 1877.

Variations in population density suggest a rather incomplete historic record along segments of western South America. The regular occurrence of earthquakes near populated areas is suspect; however, they do provide an estimate of recurrence times. Earthquakes occurring in regions between large cities are not well documented and may provide a misleading interpretation of the seismic history, i.e., the inferred repeat times are larger than the actual recurrence intervals.

Historic earthquake activity near Copiapo, Chile (25°S) suggests a rather short recurrence interval of about 40 years. Since the last large earthquake occurred in 1918 ($M = 7.7$), the area is assigned to category 2. The region between Copiapo and Valparaíso (Transverse Valley region) is located along a segment of the arc that is characterized by a shallow dipping seismic zone

and by a lack of Neogene volcanism. Both the 1922 ($M = 8.4$) and the 1943 ($M = 8.3$) events indicate that the region has a seismic potential; however, the apparent lack of historic earthquake activity necessitates placing this region in category 3.

The region near Valparaíso, Chile is marked by the intersection of the Juan Fernandez ridge with the trench, an increase in the dip of the Chilean seismic (Benioff) zone and the abrupt appearance of Neogene volcanism to the south. A moderate-size gap exists south of Valparaíso and is placed in category 2. A recurrence time of approximately 85 years is suggested from the historic data. During this century the region first ruptured in 1906 ($M = 8.2$), and the northern segment ruptured again in 1971 ($M = 7.9$). As shown in Figure 8, large events tend to migrate south along this section of the seismic zone. If this pattern continues, then the southern portion of the 1906 zone should be the site of the next large earthquake in central Chile.

South of the triple junction between the Peru-Chile trench and the Chile rise at 46°S the oceanic portion of the Antarctic plate is being subducted beneath the South American plate at a rate of about 2 cm/year (FORSYTH, 1975). Magnetic anomalies in the southeast Pacific indicate that segments of the Chile rise collided with the southern portion of the South American continent 26 m.y. ago (HERRON and TUCHOLKE, 1976). Since that time, there has been a great diminution of volcanic activity on the continent and a cessation of folding in sedimentary basins (WINSLOW, 1978). The seismicity of the region between 46° and 52°S is very low, with only a few events reported in the magnitude 4.5 to 6 range; hence, this region is assigned to category 5.

Two convergent left-lateral shear zones transect the southernmost South American continental margin near 53°S (Fig. 9). The main branch of the more easterly-trending shear zone includes the Magellan fault of KATZ (1964) which

Figure 9

Seismic activity for southern South America since 1948. Modified after WINSLOW (1978). Events are from GUTENBERG and RICHTER (1954), ISS (1950), and FORSYTH (1975). Shaded area is inferred rupture zone of 1949-50 events, based on historical data at both extremities and by field mapping in the region in between. Aftershocks, as recorded by local sources, define the rupture zone. PA = Punta Arenas, Chile; U = Ushuaia, Argentina; TF = Tierra del Fuego; SM = Strait of Magellan.

can be followed east into the North Scotia ridge. Both the Shackleton fracture zone and the North Scotia ridge correspond to the western and northern transform boundaries of the Scotia plate (FORSYTH, 1975).

Since the settlement of this region has been very recent, the earliest record of earthquake activity is the Magellan Strait earthquake of 1879 ($M = 7-8?$) (LOMNITZ, 1970). Two large earthquakes occurred within a 10 hour period in 1949 ($M = 7.5$, GUTENBERG and RICHTER, 1954). Field mapping of ground displacements associated with the 1949 events indicates a rupture length of about 300 km (WINSLOW, 1976, 1979). Damage to structures occurred in Punta Arenas, Chile as well as north of Ushuaia, Argentina, which is 300 km to the southeast (INTERNATIONAL SEISMOLOGICAL SUMMARY, 1949, 1950). The inferred rupture zone is shown in Figure 9. This region has the potential for large earthquakes and is assigned to category 6, having ruptured 30 years ago or less.

(e) Caribbean

Large segments of the seismic zone surrounding the Caribbean plate have not ruptured in the last 100 years (Figure 10). The rates of motion between the Caribbean and the North and South American plates is estimated to be approximately 2 cm/year (JORDAN, 1975); therefore recurrence intervals of at least 100 years, and probably more typically hundreds of years, may be more commonplace for this region. Since the rate of plate motion is relatively low, it is very difficult to estimate recurrence times for the Caribbean. In many areas only one large shock is reported during the last 200 years. In other areas no shocks are known historically. The quiescence observed at many sites along the plate boundary, therefore, may not be related to an incomplete

Figure 10

Estimated rupture zones of Caribbean shocks since 1800 (after KELLEHER et al., 1973, with additions). Three great shocks of the 18th century are also included. Estimated rupture zones for the earthquakes of 1943, 1946, 1948, 1953, 1969 and 1974 are based on aftershock data. Other estimates are based on isoseismal areas where intensity IX or greater was experienced. Rupture zones of events during the 19th century along the Lesser Antilles were probably seaward of the areas indicated in the figure. Arrow shows the relative motion of the Caribbean plate with respect to the North and South American plates. Dashed line is Puerto Rico trench; dotted line is the Anegada trough.

historical record but rather to the long intervals between large shocks.

The northern Lesser Antilles are, perhaps, the only classical subduction zone in the region in that they contain historically active volcanoes and subduction is occurring nearly perpendicular to the arc, i.e., in a WSW direction. This arc segment ruptured in great shocks in 1690 and 1843; the zones affected by each event appear to be very similar (KELLEHER et al., 1973). There is, then, a crude indication of a repeat time of about 150 years for this segment of the arc. It has been assigned to category 1.

To the south of this region the earthquake history does not indicate that extensive great earthquakes have recurred along any portion of the arc. In fact, it appears that some segments of this arc have not ruptured in historic times. A large sedimentary wedge, a part of the delta of the Orinoco River, lies seaward of the central and southern Lesser Antilles. Large amounts of sediment, may interfere with the subduction process (JACOB et al., 1977). Therefore, the relatively quiescent past for the central and southern Lesser Antilles may continue for some time in the future.

The coast of Venezuela has felt the effects of several major and great earthquakes, the most extensive event being the great earthquake of 1812 (KELLEHER et al., 1973). Its destruction was spread inland along the Bocono fault zone for several hundred kilometers. Undoubtedly much strain has accumulated along this fault zone and other areas that have experienced large earthquakes, but the complicated tectonics and relatively low rate of plate motion in this region make it difficult to further identify regions of high seismic potential, with the possible exception of a zone between Caracas and Cumana (64° - 67° W, 10° N, Figure 10). That zone may not have ruptured during the 1900 event, which RICHTER (1958) estimated to have a magnitude of 8.4.

The Greater Antilles are also a region of complex tectonics. Fault plane solutions near these islands indicate plate motion trending WSW. This motion occurs on fault planes that dip gently to the south beneath the inner wall of the Puerto Rico trench (MOLNAR and SYKES, 1969). The motion in this region is therefore characterized by oblique thrusting. Great earthquakes have not occurred north of much of Puerto Rico and the Virgin Islands for at least the last 400 years (KELLEHER *et al.*, 1973). Large destructive shocks of magnitude near 7 1/4, such as that in 1844, apparently originated to the north of Puerto Rico, but great events from this region are unknown. The presence of recently-deformed sediments in the accretionary wedge north of these islands indicates that the possibility of future great earthquakes cannot be ruled out. A study of the relationship between the observed seismicity and these deformed sediments and associated sediment ponds (fore-arc basins?) is presently underway (McCANN, MURPHY and FRANKEL, 1979). This region of oblique slip is similar to the Commander Islands in the western Aleutian Islands. Although the westernmost Aleutians have also been a seismic gap for this century, they may have been the site of a large interplate event in 1849 (MEDVEDEV, 1968). If so, by analogy the Puerto Rico-Virgin Islands region may be one of considerable tsunamic-seismic risk. These regions and the Andaman arc, which appear to have a similar tectonic setting, are assigned to an individual category, 4.

The Anegada passage, an extensional trough in the Virgin Islands was the source of a large tsunamic earthquake in 1867 (REID and TABER, 1920). The felt area would suggest a magnitude of about 7 3/4 to 8. Although the tectonic history and present framework of the passage are yet to be unraveled, the trough has the potential for being a source of destructive earthquakes. A local network has recorded several microearthquakes along the north wall of the passage indicating that it is presently an active tectonic feature

(MURPHY and McCANN, 1979).

Large earthquakes have occurred along the western end of the Puerto Rico trench in 1911 ($M = 7.0$), 1917 ($M = 7.0$), 1918 ($M = 7.5$), 1943 ($M = 7 \frac{3}{4}$), 1946 ($M = 8.1$) and 1948 ($M = 7.3$) (GUTENBERG and RICHTER, 1954). These events were centered in either the Mona passage off the northwest coast of Puerto Rico or along the northeast coast of Hispaniola, where subduction may be occurring more nearly perpendicular to the plate boundary than it is farther to the east or west.

The history of large shocks in Hispaniola stands in marked contrast to the quiescence of much of the Puerto Rico-Virgin Islands region. Great earthquakes have occurred on both the north and south coasts of the island (KELLEHER *et al.*, 1973). Slip between the plates in this region may be accommodated by two or more fault zones striking easterly. The north and south coasts of the central and western portions of the island have not experienced large shocks for several scores of years and in some places for over two centuries. Therefore, any of these regions may have stored sufficient energy to generate a large shock.

Jamaica and southeastern Cuba have been the sites of several major earthquakes since the 17th century (TABER, 1920, 1922). Destructive shocks occurred in Jamaica in 1692 and 1907 with another strong shock in 1812. Seawaves from each of these events indicate sources to the north of the island (TABER, 1920). It is interesting to note that these shocks that are very destructive on Jamaica are not regarded as strong shocks on the southeastern coast of Cuba less than 200 km away. In fact, there are no reports of damage on Cuba from the "great" Jamaican earthquake of 1692 (TABER, 1922), nor is the "great" Cuban earthquake of 1678 reported on Jamaica. As each of these areas has experienced at least locally destructive shocks, they have the potential to

Figure 11

Relocated epicenters (circles) of shallow earthquakes (depths ≤ 70 km) for the western boundary of the Caribbean plate from 1954 to 1962 (after MOLNAR and SYKES, 1969, with additions). Larger symbols are more precise locations. Note the general lack of activity south of Cuba and thence west to Guatemala. Solid circles are sites of major earthquakes of this century. Solid arcs show regions affected by large earthquakes or tsunamis. Solid line shows inferred position of boundary of Caribbean plate.

be the future sites of major earthquakes. Preliminary analysis of the historic record indicates major Cuban shocks occur about every 100 years. The last occurred in 1852. Southeastern Cuba is assigned to category 1. Major shocks have occurred near Jamaica in 1692, 1812 and 1907 with the 1812 event being of lesser intensity. The events of 1692 and 1907 are associated with widespread destruction, and are probably great ($M > 7.75$) in size. The event in 1812 appears not to have been that large. The area near Jamaica is assigned to category 2.

Very little information exists about large shocks to the west of Jamaica. The area is largely submarine and the islands are sparsely populated; therefore, only instrumental locations are available. The area of contact along this transform boundary is probably of sufficient strength to be the source of future large shocks involving strike-slip motion. The thin, weak crust near the Cayman spreading center near 82°W , however, will probably not be the site of large shocks in the future.

(f) South Sandwich Arc

In the central and southern parts of the South Sandwich arc, the subduction of young, oceanic lithosphere generated at an adjacent spreading center (Fig. 12) results in a sizeable region that lacks known shallow earthquakes of magnitude 6 or larger (Figs. 13 and 14; FRANKEL and McCANN, 1979). Subduction of young buoyant lithosphere is also accompanied by uplift of the platform on which the southern islands are located and by shoaling of the trench. Bathymetric features on this incoming plate may further modify the subduction process in the far southern portion of the arc, as it is a region of intraplate seismicity. In the northern part of the arc where older (70 m.y.)

Figure 12

Generalized configuration of plate boundaries in the South Sandwich region (from FRANKEL and McCANN, 1979). Single lines represent transform boundaries, double lines are spreading centers, serrated border is subduction zone (teeth are on the overthrust plate), and dashed lines indicate plate boundaries of undetermined character. SSFZ is the South Sandwich fracture zone. NSR and SSR are segments of the North Scotia ridge and the South Scotia ridge, respectively. The arrows indicate the sense of motion along these features. The vertical lines with the numbers below them represent the oldest magnetic anomalies identified behind the South Sandwich arc and their anomaly numbers.

Figure 13

Large ($M \geq 7$) earthquakes in the South Sandwich region, 1910–August 1977 (from FRANKEL and McCANN, 1979). Error bars indicate the 95% confidence limit on relocated epicenters. Bathymetry is in fathoms (1 fathom $\sim$ 2 meters), shaded areas being deeper than 4000 fathoms.

Figure 14

Moderate-sized earthquakes of the South Sandwich region, 1923-1976 (from FRANKEL and McCANN, 1979).

lithosphere is being subducted, more events of moderate size ($6 \leq M \leq 7$) occur in the region of underthrusting along the inner wall of the South Sandwich trench. Conspicuously absent, however, are large or great events involving thrust faulting. Perhaps the density contrast between the young overthrust plate (age approximately 7 m.y.) and the old underthrust plate is so large that only weak coupling exists between the plates. This may result in motion that is mostly aseismic. Another possibility is that back-arc spreading, which is occurring at a fast rate behind the arc, may lead to decoupling or low stress along the entire plate interface at the subduction zone.

(g/i) Western Pacific

There is a remarkable variations in the size and number of large thrust earthquakes that occur along the western margin of the Pacific plate (Fig. 15). Much of this variation can be explained by changes in the geometry and geophysical parameters that determine the tectonic regime of a particular segment of an arc; these include:

- 1) changes in the interface or zone of contact between the overthrust and underthrust plates;
- 2) modification of the subduction process by the interaction of bathymetric features on the incoming seafloor with the overthrust plate [this condition is believed to lower the density contrast between the converging plates and hence to inhibit the subduction process (KELLEHER and McCANN, 1976)];
- 3) large density contrasts between the overthrust and underthrust plates which may lead to poor seismic coupling and a reduction in the ratio of seismic to aseismic slip (KANAMORI, 1977a; McCANN and NISHENKO, 1978).

Figure 15

A comparison of bathymetry (CHASE et al., 1970), rupture zones of large earthquakes (cross hatching) and active volcanoes (solid triangles) along the western Pacific (from KELLEHER and McCANN, 1976). Note the smooth ocean floor, large rupture zones and nearly continuous chain of volcanoes toward the north (Kamchatka-Kuril-Japan). Contrast these features with the irregular bathymetry, absence of great earthquakes and gaps and irregularities in the line of active volcanoes along the Bonin and Marianas arcs.

(g) Izu-Bonin-Mariana Arcs

The Izu-Bonin-Mariana arcs form the eastern edge of the Philippine Sea plate. In those areas the subducted Pacific plate appears to be reflected in a downgoing seismic zone which extends to depths of almost 700 km (KATSUMATA and SYKES, 1969). Neither GUTENBERG and RICHTER (1954) nor ROTHE (1969) report any shallow events of magnitude greater than 7.3 along these arcs between latitudes 10° and 35° N. The remarkable absence of great earthquakes and the small number of reported large shocks along this plate boundary may be attributed to one of the following factors:

- 1) There is little or no relative motion between the Philippine Sea plate and the Pacific plate.
- 2) Motion along the entire plate boundary occurs aseismically as a result of a decoupling of the plates (KANAMORI, 1977a). This could result from a large density contrast between the overthrust and underthrust plates (McCANN and NISHENKO, 1978; FRANKEL and McCANN, 1979) or from a weakening of the zone of interplate motion by frictional (or other) heating (KANAMORI, 1971).
- 3) Segments of the Pacific seafloor are too light to be subducted beneath the overthrust plate. Deformation of the plates then occurs; intraplate activity may increase, but large thrust events become rare as subduction is interrupted.
- 4) The recurrence time is extremely long (i.e., larger than the historic record which is complete only for about the last 75 years for most of the region).

At present the first hypothesis cannot be ruled out. The paucity of shallow activity and the lack of focal mechanism solutions of the thrust type can be

used to argue for this possibility as well as for the second and third hypotheses. Until more information is gathered on this region we cannot resolve this matter. The deformation of the Mariana arc where the Marcus-Necker and Carolina ridges intersect the trench may indicate that the third possibility may apply there. KELLEHER and McCANN (1976) and VOGT et al. (1976) favor that interpretation. KANAMORI (1977a) proposes that the absence of great earthquakes along this margin may be the result of poor coupling between the plates. He shows that a significant portion of the motion between the Pacific and Eurasian plates along that segment of the plate boundary off Honshu and the southern Kuril Islands may occur aseismically. He proposes that the decoupling is greater farther to the south, which results in nearly total aseismic motion on the zone of shallow thrusting in the Marianas, Bonin and Izu arcs.

Large shocks do occur occasionally along this boundary. The magnitude of one event in 1902, which is listed by RICHTER (1954) as a great shock ($M = 8.1$) is probably over estimated (KANAMORI and ABE, 1978). The macro-seismic effect of that shock did not include any indication of a seawave, although two islands 250 km apart are claimed to have exhibited effects similar to intensity IX on the Modified Mercalli scale (ALGUÉ, 1909). Guam has been the site of historic destructive earthquakes (ALGUÉ, 1909) but is has not experienced destructive seawaves. It is possible that these events are intermediate in depth and are therefore not directly related to the under-thrusting process. The instrumental record and the short historic record of the Marianas give no indication of the occurrence of large events of the thrusting type that are typical of so many other subduction zones.

It is possible that recurrence times in this region are longer than the historic record. If this is so, then this region is very unusual.

Convergent zones with relatively narrow interfaces, i.e., thin zones of plate contact, are found in the Central America and along the Solomon arc, where short recurrence times (30-50 years) are typical. The Mariana arc also has a narrow interface. Given the present instrumental record, recurrence times would have to be at least 75 years. While this could be the case, it is difficult to explain why the whole of the Izu-Bonin-Mariana arc system from 10° to 35°N is presently in a quiescent stage. This type of synchronicity is unparalleled in any other arc with the possible exception of Java, the Lesser Sunda Islands (which will be discussed later), and the South Sandwich arc.

Certain tectonic similarities between these arcs and others with no history of great shocks suggest to us that the 75 years of instrumental data are typical of the longer pattern of seismic energy release. Hence, we conclude that these arcs have a low potential (category 5) for being the sites of large or great shallow earthquakes of the thrusting type.

(h) Northern Honshu and Hokkaido

Fig. 16a, b, c from KELLEHER and McCANN (1977) shows bathymetry, recent tsunamis and moderate and large ($M \geq 6.5$) earthquakes near the east coast of Honshu, Japan from 1926 to 1973. Smaller tsunamis and earthquakes have occurred in the region south of 38°N near the section of trench that is interacting with a group of seamounts on the Pacific plate (KELLEHER and McCANN, 1977). To the north, smooth seafloor is being subducted into a region that has been the site of several great earthquakes and destructive tsunamis. The observed spatial pattern of occurrence of large shocks is consistent with the historic record. Hence, we expect this pattern to continue in the future.

Figure 16

Seismicity and tectonics near the Japan trench (from KELLEHER and McCANN, 1977). a) Epicenters and focal regions of shallow earthquakes of magnitude 6.5 and larger from 1926 to 1973 after UTSU (1974). Note the characteristically smaller focal regions near the southern half of the Japan trench. b) Tsunami sources near Japan for 76 years (1893-1968) after HATORI (1969). Distribution during past centuries was similar (HATORI, 1975). Note large tsunami sources near northern half of Japan trench. c) Bathymetry near Japan trench after CHASE et al. (1970). Note zone of seamounts near southern half of trench and smooth ocean floor seaward of northern half of trench where great earthquakes and tsunamis have originated.

Fig. 17 summarizes the known seismic gaps near the southern and eastern coasts of Japan. These gaps are nearly identical to those identified by MOGI (1968) and ANDO (1975).

(i) Kuril-Kamchatka

Much of this portion of the Pacific plate boundary has ruptured during the last 80 years. Fig. 18 from KIMURA and McCANN (1979) shows the rupture zones of great earthquakes and active volcanoes of this region. Two prominent seismic gaps are also indicated. The northernmost gap last ruptured during a great earthquake in 1923 (FEDOTOV, 1965). It has been assigned to category 2. The southern gap, near the central Kuril Islands, may have ruptured, in part, during an event in 1915. Since very little instrumental information is available for that event, the extent of rupture is uncertain. Earthquake catalogs (MEDVEDEV, 1968; KONDORSKYA and SHEBALIN, 1977) indicate that this region was less active for major shocks than the adjoining regions as far back as the early 1800's. More analysis of the geophysical data for this region is necessary to properly assess its potential as a source of future shocks. It is tentatively assigned to category 3.

Fig. 19, also from KIMURA and McCANN (1979), displays the history of eruptions for various volcanoes along the Kuril-Kamchatka arc. Heavy bars indicate the locations in time and space of all great earthquakes since 1950. For each volcano, bars are shown indicating the time of a major (full bar) or moderate (one-half bar) eruption. Volcanic activity adjacent to the rupture zones of the 1971 and 1952 earthquakes was clearly higher for several decades before each of those events compared to the years after those events. Therefore, observations of volcanic activity may provide a valuable tool for the

Figure 17

Seismic gaps and recent large earthquakes near the eastern and southern coasts of Japan. Hatched regions are rupture areas determined from after-shocks. Year and magnitude of the events are indicated next to the affected region.

Figure 18

Great earthquakes along the Kuril-Kamchatka arc (from KIMURA and McCANN, 1979). Solid lines represent rupture areas of recent great events. Dashed lines represent rupture zones inferred from felt reports. Dots and X's represent poor and good epicentral locations, respectively, for main shocks. Dates and magnitudes of main shocks are indicated. Numbers next to volcanoes (solid triangles) are for reference to Fig. 19. Arrows denote directions of plate motion taken from MINSTER et al. (1974).

Figure 19

Space-time plot of the seismic and volcanic history of the Kuril-Kamchatka region (from KIMURA and McCANN, 1979). Thick, long bars represent time and location of great earthquakes indicated on the previous figure. Small and full bars represent the time of small or moderate (m) to great (g) eruptions. Lava flows are indicated by wavy arrows.

KURIL

the long-term forecasting of future large earthquakes.

(j) Western Margin of the Philippine Sea

The western margin of the Philippine Sea plate can be divided into five zones, based on the tectonic character: the Nankai trough, the Ryukyu arc system, Taiwan, the Luzon Strait between Taiwan and the northern Philippines, and the Philippines. The Nankai trough is a young trench that has a historic record of great earthquakes, with extensive rupture zones and tsunamis, dating back to A.D. 684 (ANDO, 1975). The Ryukyu arc is a well-developed trench, with a well-defined dipping seismic zone. Taiwan, the Philippines, and the region between are complex island arc systems, which have undergone (and may be presently undergoing) polarity reversals. Parts of the latter experience frequent larger shocks, but lack well-developed subduction zones.

ANDO (1975) examined historic great earthquakes in the Nankai trough which meets the Ryukyu arc at the Palau-Kyushu ridge (Fig. 20). He divided the region into four blocks, west to east, A through D. He finds that great earthquakes tend to occur in cycles, which migrate from west to east, with large shocks occurring at a given place every 100 to 200 years. The latest cycle seems to have begun with the 1944-46 Tonankai and Nankaido earthquakes, which ruptured sections A through C. The cycle has not yet reached section D, which last ruptured in the Ansei I earthquake of 1854 (ANDO, 1975). Hence, a great earthquake may be expected east of the Tokai district. That region has been assigned to category 1, i.e., highest seismic potential. The great historic earthquakes of the Nankai trough studied by Ando all have rupture zones that extend for several hundreds of kilometers, and terminate at the intersection of the Palau-Kyushu ridge with the Nankai trough and northern end of the Ryukyu arc (KELLEHER and McCANN, 1976).

Figure 20

Bathymetry and active tectonics along the Ryukyu arc and Southwest Japan (after KELLEHER and McCANN, 1976). Seismic record near Southwest Japan is based on ANDO (1975). Triangles represent epicenters located by GUTENBERG and RICHTER (1954). Depth contours are in fathoms (1 fathom = 1.83 m). Solid triangles are Quaternary volcanoes.

The Ryukyu arc is typical of a subducting margin: earthquakes are concentrated in a well-developed, planar zone that dips 35° to 45° to the northwest, and reaches depths of 250 to 300 km (KATSUMATA and SYKES, 1969); ROWLETT and KELLEHER, 1976). Behind the frontal arc and volcanic chain, the Okinawa trough represents a slowly-extending marginal basin (KARIG, 1973).

Historical data indicate that the southwestern portion of the Ryukyu arc has had a greater occurrence of large earthquakes than other portions of the arc. ROWLETT and KELLEHER (1976) note at least one severely destructive shock in 1771, which was accompanied by a destructive tsunami (Fig. 20). The northeastern portion, from Okinawa to Kyushu (Fig. 20), by contrast, has experienced only infrequent large shocks, with no reports of destructive tsunamis (ROWLETT and KELLEHER, 1976). Seaward of the southwestern portion of the arc, the bathymetry is relatively flat, whereas seaward of the portion from Okinawa to Kyushu a series of aseismic ridges intersect the trench and may be interfering with the subduction process (KELLEHER and McCANN, 1976). KATSUMATA and SYKES (1969) cite a region from 24° to 25° N, 126° to 128° E as being nearly aseismic for small to moderate size earthquakes of shallow depth. In this region the Miyako Depression (Fig. 21) intersects the Ryukyu trench. To the northeast and southwest of this "gap" KATSUMATA and SYKES (1969) and ROWLETT and KELLEHER (1976) noticed a clustering of activity since 1961. Similar patterns of gaps in short-term seismicity, with clustering of events at the edges, were observed by MOGI (1968) and KELLEHER and SAVINO (1975) along subduction zones prior to the occurrence of large earthquakes. Such areas may be likely to experience large shocks in the future. The northeast portion of the Ryukyu arc, which has no historic record of great earthquakes with extensive rupture zones or destructive tsunamis, is assigned to category 5. Because prominent bathymetric features seaward of the trench may be hindering the subduction process in this area, we do not expect great earthquakes for this region. The southeast portion, which may be exhibiting

Figure 21

Epicenters of large shallow earthquakes and aftershock zones for the Luzon Strait-Taiwan-Ryukyu Islands region since 1918 (after ROWLETT and KELLEHER, 1976). Relocated epicenters (solid circles), epicenters located by GUTENBERG and RICHTER (1954) (solid triangles), aftershock data determined from relocated data (hachured regions), and Longitudinal fault of Taiwan (L.F., solid line) are shown. Note the intense activity for large earthquakes near Taiwan as compared to that near either the Ryukyu Islands or the Luzon Strait. Much of this activity is located on the "oceanic" side of the Longitudinal fault.

activity observed along other subduction zones prior to large earthquakes, and has not had a great shock since 1771, has been assigned to category 1, the highest seismic potential.

During this century Taiwan (Fig. 21) has experienced frequent large ($7 \leq M \leq 7.5$) shocks, with ruptures extending tens (as opposed to hundreds) of kilometers (ROWLETT and KELLEHER, 1976). Much of this activity tends to cluster along the Longitudinal Valley fault zone near eastern Taiwan. This feature probably represents the collisional boundary between the Asian continental plate and the northern part of the Luzon arc system (KARIG, 1973; ROWLETT and KELLEHER, 1976).

East of the Longitudinal Valley, Miocene to Pliocene melange and andesitic volcanic terrane comprise the Coast Range (KARIG, 1973). Karig notes the continuation of "arc structural units" from the Batanes-Babuyan Islands to the south within the Coast Range, as well as a continuation of the east-dipping seismic zone of shallow (mostly less than 100 km) foci (KATSUMATA and SYKES, 1969) as evidence that the eastern section of Taiwan is part of the Luzon arc system. Taiwan is thought to have collided with the west-facing Luzon arc in the Late Pliocene (KARIG, 1973), and may be undergoing a reversal in polarity at the present time. Left-lateral offset along the Longitudinal Valley and along faults to the west are related by KARIG (1973) to intraplate shortening, which results from continued collision.

The present seismic regime of Taiwan is characterized by: 1) large, frequent shocks with small rupture zones; 2) focal mechanisms with scattered orientation; and 3) a poorly-defined seismic zone with an easterly-to-vertical dip beneath the east coast of Taiwan and maximum depths less than 200 km, and a tendency for shocks to be associated with the Longitudinal Valley fault (KATSUMATA and SYKES, 1969; ROWLETT and KELLEHER, 1976). The rather widespread distribution of seismic activity in Taiwan and as far south as the Philippines may be related to internal plate deformation associated with the collision of the Luzon arc system with

continental material and to a possible reversal of polarity of the arc (KARIG, 1973; ROWLETT and KELLEHER, 1976). Taiwan is assigned to categories 1 and 2. Even though it has experienced large events less than 100 years ago, the repeat time of large shocks appears to be less than 100 years. Large shocks, however, have been a relatively rare phenomenon on Taiwan since about 1957. That relative quiescence for large shocks appears to be a temporary phenomenon.

The Luzon Strait (Figs. 21, 22), between the northern Philippines and Taiwan, consists of a series of ridges and troughs that are broken into en echelon segments, which suggest that left-lateral shear is operating in this region (KARIG, 1973). KARIG (1973) interprets these features as a continuation of the west-facing Luzon arc system, and traces them into eastern Taiwan. No large earthquakes are known to have occurred in this region during this century, but moderate ($M = 6-6.9$) shallow events are observed across a 200 km-wide zone between Luzon and Taiwan (ROWLETT and KELLEHER, 1976). West of Luzon, the Manila trench (Fig. 22) is continuous as a depression to 20.5°N , where it merges with the series of north-northeast trending ridges and troughs of the Luzon Strait. KARIG (1973) and ROWLETT and KELLEHER (1976) suggest that plate motion in the region between southern Taiwan and the northern Philippines is accommodated by regional release of tectonic strain and that sharp plate boundaries have not yet developed. If this is the case, the absence of large earthquakes may be a permanent feature of this region (ROWLETT and KELLEHER, 1976). Hence, it has been assigned to category 5.

The Philippines are characterized by frequent large, shallow earthquakes that occur throughout the islands. The active features with which the seismicity is associated are the Manila trench, Philippine fault system and the Philippine trench (Fig. 22).

The present tectonic pattern of northwestern Luzon is that of a west-facing island arc system (KARIG, 1973). The Manila trench is associated with a poorly developed seismic zone that dips east. The deepest foci lie between 150 and 200 km.

Figure 22

Large earthquakes near the Philippine Islands (after ROWLETT and KELLEHER, 1976). Relocated epicenters (solid circles), epicenters located by GUTENBERG (1956) or GUTENBERG and RICHTER (1954) (solid triangles), aftershock zones determined from relocated data (hachured regions), estimated rupture zones of several earthquakes prior to 1900 (dotted regions), and trace of the Philippine fault (solid and dotted lines) as mapped by ALLEN (1962) are shown. Dashed boundaries and queries indicate uncertain boundaries of aftershock zones or less reliable epicentral locations. Data prior to 1918 are not complete. Note that several large earthquakes have occurred during the past 105 years near the trace of the Philippine fault.

Seamounts and other bathymetric features on the underthrusting plate to the west of Luzon may be modifying the subduction process along the northern portion of the Manila trench (ROWLETT and KELLEHER, 1976). KARIG (1973) and ROWLETT and KELLEHER (1976) suggest that subduction may be in the process of terminating along the Manila trench. Focal mechanism solutions of shallow-focus events along the east coast of northern Luzon, which do not appear to be associated with the Philippine fault system, indicate westerly-directed underthrusting. These mechanisms are cited by KARIG (1973) as evidence that subduction may be developing in northeastern Luzon.

The Philippine fault system extends from Mindanao to central Luzon, where it branches into several subparallel faults which extend into northwestern Luzon. FITCH (1972) proposes that oblique convergence of the Philippine and Asian plates may result in thrust motion at the Philippine trench and left-lateral and strike-slip motion along the Philippine fault. Epicenters of large earthquakes for this century appear to lie close to the fault trace, and sections of the fault 75 km or more in length (Fig. 22) which have not ruptured for several decades appear to have the potential for being sites of future large shocks (ROWLETT and KELLEHER, 1976).

The Philippine trench has typical features of an active subduction zone from about 3°N to 12.5°N (ROWLETT and KELLEHER, 1976). Near 15°N it swings west as a shallower trough, which is almost sediment free, is characterized by shallow seismicity, and is presumed to be a zone of underthrusting and left-lateral shear (KARIG, 1973). Large earthquakes have occurred along most of the trench during the past century. South of Mindanao, the seismic zone is characterized by fault plane solutions involving thrust faulting (ROWLETT and KELLEHER, 1976). KARIG (1973) suggests that subduction may be beginning along this seismically active section. Scattered activity throughout the lesser Philippine Islands may indicate the boundaries of a small plate in the western Philippines (KATSUMATA and SYKES, 1969) or internal deformation (FITCH, 1972).

Most of the Philippines are assigned to category 2; large, frequent shocks are

characteristic. Many large earthquakes have occurred there within the past 30 to 100 years.

(k) Sunda Arc

The seismotectonic nature of the Sunda arc changes throughout its length as the direction of relative plate motion changes from normal near Java to oblique near the Andaman Islands. East of the Sunda Strait, in Java and the Lesser Sunda Islands, there is a lack of historic great earthquakes and associated tsunamis, whereas to the west, in Sumatra, the historic record indicates great earthquakes and extensive damage by tsunamis. From the historic record there appears to be a higher potential for great earthquakes on and near Sumatra than near Java and the islands to the east (Fig. 23). At present the tectonic reasons for this are poorly understood and more work needs to be done in this area.

VISSER (1922), reporting on historic Sunda earthquakes from the eighteenth, nineteenth and early twentieth centuries states, "Most earthquakes felt on the Isle of Java are weak". KELLEHR and McCANN (1976) note a prominent gap in activity for large shocks during the twentieth century between 105° and 122°E . This lack of earthquakes may be related to the interaction of the Christmas Islands ridge, a prominent bathymetric feature, with the subduction zone off central Java and to the interference of the Australian continent with the subduction process near the Lesser Sunda Islands. CARDWELL and ISACKS (1978) note a reduced number of shallow earthquakes as well as a paucity of focal mechanism solutions involving underthrusting for the eastern Sunda arc near Timor. They relate these factors to subduction of the leading edge of the Australian continental shelf. On the basis of 41 new and old focal mechanism solutions for the eastern Sunda arc, CARDWELL and ISACKS (1978) propose a model of a slab that is

Figure 23

Large earthquakes ($M \geq 7.5$), of shallow focus, of the twentieth century along the Sunda arc (from KELLEHER and McCANN, 1976). Note the gap in activity for large shocks from 106° to 122°E . This region has no history of large destructive earthquakes. The region offshore from Sumatra, however, has been the site of great tsunamic earthquakes of extensive length and should be considered as a potential site for the occurrence of future large and great events.

contorted and bent at depth as a result of curvature of the eastern end of the arc. North-northwesterly convergence would, according to their model, translate the bend of the slab at the eastern end from the surface down to depths of 600 km. The north-northwesterly direction of convergence between the Australian plate and the Sunda arc inferred by Cardwell and Isacks contradicts the northeasterly direction computed by LePICHON (1968), MORGAN (1972), and MINSTER et al. (1974). Their poles of rotation were based on magnetic anomalies and hot spot traced (MINSTER et al., 1974), with no instantaneous slip vectors derived from focal mechanism solutions. There is a dearth of shallow underthrusting mechanisms for the entire Sunda arc; only four have been reported, all of which lie near Sumatra (FITCH, 1972; CARDWELL and ISACKS, 1978). CARDWELL and ISACKS (1978) cite seven volcanoes at the eastern end of the arc near the Weber basin as evidence of north-northwesterly convergence. They propose that Southeast Asia is not part of the Eurasian plate, but that there is a separate Southeast Asian plate with a southeasterly direction of motion relative to the Australian plate. Their model is consistent with the direction of oblique convergence to the north-northwest between the Australian plate and Sunda arc near the Andaman Islands. Other evidence that may support the model of Cardwell and Isacks is the observed dextral slip on the Semangko fault in Sumatra and south-southeasterly extension in the Andaman basin (RODOLFO, 1969).

The tectonics of the eastern Sunda arc are complex and still poorly understood. The lack of great thrust earthquakes at shallow depths may be related to the contortions of the dipping slab as it bends around the eastern Banda Sea and to the interference of the Australian continental shelf, Christmas Island ridge, and other bathymetric features with the subduction process. Although more work is needed to understand the tectonic regime, there is

clearly a lack of historic great thrust earthquakes and associated tsunamis for the eastern Sunda arc. Hence, on the map of seismic potential this region is shown as category 5. Under the present tectonic regime the lack of great earthquakes recorded for the past 200 years may be expected to prevail during the foreseeable future. Nevertheless, the reasons we give for believing this region differs tectonically from subduction zones with a record of great shocks should be regarded as a hypothesis and not proven fact. The basins north of Java and the Lesser Sunda Islands do not appear to be characterized by high heat flow and recent extension as they do behind other arcs such as the Marianas and South Sandwich, which also do not have a history of great shocks. If current spreading along marginal basins is the important factor in governing the occurrence of great shocks, then the eastern end of the Sunda arc may, in fact, have the potential for great shocks to occur.

At the Sunda Strait the arc undergoes a sharp bend from an easterly to northwesterly strike, and the direction of relative plate motion becomes increasingly oblique. The maximum depth of earthquakes changes abruptly from 600 km east of the Sunda Strait to about 200 km on the west side. FITCH (1972) proposes that the Sunda Strait may be undergoing extension on the seaward (convex) side and compression on the landward side, which may be related to recent shallow activity in the area. The Semangko fault of Sumatra shows evidence of historic and recent right-lateral movement (RODOLFO, 1969), while the Lambang fault east of the strait, in Java, which strikes parallel to the arc, may indicate left-lateral offset (KATILI, 1970).

FITCH (1972) proposes a model for island arcs with oblique convergence in which there is decoupling along a pre-existing zone of weakness. According to his model, strike-slip movement occurs along a fault between the trench and volcanic front, while underthrusting takes place at the trench. In the

western Sunda arc underthrusting is occurring along the inner wall of the Java trench, while horizontal shear is simultaneously occurring along the Semangko fault.

Great shallow events that are thought to involve thrust faulting are recorded historically, as well as large inland shocks associated with the Semangko fault. Rupture lengths for great historic earthquakes for the nineteenth and twentieth centuries, as estimated by reports of tsunami destruction by VISSER (1922) tend to be several hundred kilometers long (Fig. 24). Also shown in Fig. 24 are locations of events with reported intensities of VIII or IX from SOEDATI (1962) for the period 1900 to 1960, which appear to be associated with the Semangko fault, as well as events of magnitude $M \geq 7.5$ that occurred during the twentieth century. [Intensities reported by Soedati are based on the Rossi-Forel scale; equivalent Modified Mercalli intensities for R.-F. VIII to IX are VII to IX (RICHTER, 1958)]. It is evident that Sumatra, unlike Java and the eastern Sunda Islands, has had a history of large to great events, both shallow thrusting (associated with the trench), and strike-slip (associated with the Semangko fault).

The extent of the rupture zone of the 1861 event, as inferred from reports of tsunami inundation by VISSER (1922) may be as great as 800 km (Fig. 24). If this estimate is credible, then that event is the most recent great event that can be said to have ruptured a major portion of the plate boundary and relieved significant tectonic strain. Hence, the western portion of the Sunda arc off Sumatra has not experienced a great event for over 100 years, and is assigned a seismic potential of category 1.

The portions of the Semangko fault with historically recorded activity all appear to have ruptured between 30 and 100 years ago, and hence are assigned to category 2. [The oldest reported large event along the fault

Figure 24

Large historic earthquakes in the Sunda region. Dashed regions are areas of reported damage by tsunamis associated with historic great earthquakes (VISSER, 1922). The Semangko fault in Sumatra is shown as a sinuous line. Heavily lined portions along the Semangko fault are areas reported by VISSER (1922) to have ruptured in the years shown. Squares are reported intensities VIII and IX (Rossi-Forel) by SOEDATI (1962). Solid circles are relocated epicenters originally reported by GUTENBERG and RICHTER (1954); open circles are Gutenberg and Richter epicenters which have not been relocated. Diamond with dot in insert is the only historically active volcano in the Andaman-Nicobar Islands region, reported by RODOLFO (1969). Thin arrow is direction of relative motion between Australian and Eurasian plates of MORGAN (1972); thin dashed arrow is that of LePICHON (1968). Heavily blocked arrow is direction of motion of hypothesized Southeast Asian plate with respect to Australian plate (CARDWELL and ISACKS, 1978).

occurred in 1892 (VISSER, 1922), which resulted in 2 meters of offset (RODOLFO, 1969).]

The 1941 earthquake in the Andaman Islands, although given a magnitude of 8.7 by RICHTER (1958), was probably not of comparable size as is discussed later. Its rupture zone does not appear to be several hundred kilometers long.

The portion of the Sunda arc along the Andaman and Nicobar Islands is characterized by oblique convergence. There are no great earthquakes or extensive tsunamis reported historically. The earthquake of June 26, 1941, which occurred west of the Andaman Islands, is believed to have been the largest event of this century recorded along the western Sunda arc; however, it is probably not a great event. SINDVAHL et al. (1978), using RICHTER'S (1958) magnitude of 8.7, infer a rupture zone of 800 km, but these estimates of magnitude and rupture length appear to be far too high.

GELLER and KANAMORI (1977) re-examined the methods by which RICHTER (1958) and GUTENBERG and RICHTER (1954) computed magnitudes. An appendix to the paper of Geller and Kanamori contains corrected values of M_s . The 1941 event has a corrected M_s of 7.7. KANAMORI (1977b) calculated revised magnitudes, M_w , based on seismic moment for great earthquakes with extensive rupture zones. His new magnitude scale, M_w , extends the M_s scale, which saturates at magnitudes near 8.0, so as to accurately measure the moment and energy release of large shocks. Events with rupture lengths of about 800 km have M_w values near 9.0 on his scale. The M_s and M_w scales coincide up to approximately magnitude 8.0. Thus, an event such as the 1941 shock of M_s near 7.7, is unlikely to have a rupture zone of hundreds of kilometers, as its magnitude would have been re-calculated, by Kanamori's method, near 9.0. Also, there are no reports of tsunamis associated with the 1941 event, and few reported aftershocks. The lower re-calculated magnitude, general lack of aftershock activity and macro-

seismic effects for the 1941 event lead us to believe that SINDVAHL et al. (1978) over estimated the extent of the rupture of this event. Hence, there may not appear to be great earthquakes associated with the Sunda arc in the Andaman-Nicobar region.

The northwestern part of the Sunda arc near the Andaman and Nicobar Islands is assigned to category 4 because plate convergence occurs subparallel to the subduction zone in that region. A dipping seismic zone extends to depths of 100 to 200 km beneath the Andaman-Nicobar ridge, but the dominant sense of motion as derived from the few available focal mechanism solutions for this region is dextral strike-slip, with nearly vertical nodal planes striking northwest (FITCH, 1972). RODOLFO (1969) notes only one historically active volcano in the area near the island of Narcondam (Fig. 24). He cites evidence that extension has been occurring in the Andaman basin since the Late Miocene. High heat flow measured in the Nicobar rift, subparallel north-northeast trending magnetic anomalies in the Andaman basin, and relatively thin (1.5 km) sediment cover in the southern Andaman basin are presented by Rodolfo as evidence of extensional tectonics, trending NW. FITCH (1972) compared the Andaman basin to the model proposed by SCHOLZ et al. (1971) for the western United States. As compressive stress was relieved off the coast of California in the late Tertiary, and the dominant plate motion changed to strike-slip along the San Andreas fault zone, extension was triggered in the Basin and Range province. FITCH (1972) suggests that the submarine continuation of the Semangko fault might be comparable to the San Andreas. The evidence for southeasterly extension in the Andaman basin cited by RODOLFO (1969) appears to support CARDWELL and ISACK'S (1978) northwesterly motion of the Australian plate with respect to the Sunda arc and the existence of separate Eurasian and Southeast Asian plates.

The Andaman portion of the Sunda arc appears to be tectonically similar to the westernmost Aleutians, where plate convergence is subparallel to the arc and high heat flow is present behind the arc (COOPER et al., 1977).

(1) Southwest Pacific

During this century large shallow earthquakes have not been evenly distributed along the subduction zones of the southwest Pacific. Figs. 25, 26 and 27 are from a recently-completed study of large earthquakes occurring along the shallow thrust boundaries of the southwest Pacific arcs (McCANN and KELLEHER, 1979). The details of the tectonics in this complicated region will not be presented here. These figures show either the clustering or the absence of known large events along various convergent boundaries. Since these variations appear to be spatially related to the regional tectonic setting, they may represent long-term features of the seismicity.

A triple junction between the South Bismark, Solomon Sea and Pacific plates is located near 152°E, 6°S (JOHNSON and MOLNAR, 1972) (Fig. 25). The intense clustering of large events in this region appears to result from the rapid subduction of the Solomon Sea plate near this triple junction. This intense activity, therefore, may continue indefinitely as it appears to stem from a tectonic regime characterized by high stress. The large gap in the central portion of the Solomon arc (near 157°E) occupies that segment of the arc that intersects the Woodlark rise, a series of spreading centers that forms the boundary between the Solomon Sea and Australian plates. Lithosphere in that area appears to be too weak to store the energy sufficient to generate large earthquakes. The region between 159° and 161°E recently ruptured during a series of large events. The remaining segment of the Solomon arc from 161°

Figure 25

Large ($M \geq 7.0$) earthquakes of the New Guinea-Solomon area for the twentieth century (from McCANN and KELLEHER, 1979). The large gap in activity near 157°E coincides with the segment of the arc subducting the Woodlark rise, a spreading center. This gap may therefore persist indefinitely. Contours are in fathoms. Solid circles are locations taken from GUTENBERG and RICHTER (1954). Open circles are relocated epicenters; hatched regions are rupture zones; triangles are events with uncertain depth determinations. Small circle pattern in the region of the trench delineates water depths greater than 3000 fathoms.

Figure 26

Large ($M \geq 7.0$) earthquakes of the twentieth century and their associated aftershock zones for the New Hebrides arc (after McCANN and KELLEHER, 1979). Note the lack of recent large shocks to the south of 20°S . Heavily hatched regions is that portion of the New Hebrides trench with water depths greater than 300 fathoms.

Figure 27

Large ($M \geq 7.0$) earthquakes of the twentieth century along the Tonga and Kermadec arcs (from McCANN and KELLEHER, 1979). Note the sparse activity for large earthquakes in the Tonga arc region and the smaller number of large shocks from 32° to 38°S and from 23° to 28°S . These gaps appear to be of tectonic origin and may persist indefinitely. Symbols same as in Figure 26.

to 163°E, which has not ruptured in over 30 years, is probably the most likely site for future large shocks along this arc.

The rupture lengths of large shallow earthquakes are only moderately long (maximum dimensions less than 150 km) along all of the subduction zones of this region. These limited dimensions of rupture zones may result from the narrow width of interface, the zone of contact between the underthrust and overthrust plates, along the convergent boundaries. There is an observable variation in the maximum dimension of rupture zones from north to south along the New Hebrides arc. Rupture zones with large maximum dimensions occur in the northern portion of the arc. Heat flow values in this region are lower on both the incoming seafloor and on the overthrust plate compared to values observed to the south. The very southern portion of the arc has a poorly-developed island arc platform and may only be Quaternary in age (KARIG and MAMMERICKX, 1972). In this region rupture zones are smallest when compared with the rest of the arc. The change in the rupture dimensions may result from a narrower interface between the younger plates of the southern portion of the arc.

The distribution of large shocks along the Tonga and Kermadec arcs is very uneven. In the Kermadec arc, which is about 1000 km long, large shocks of this century are clustered along a 300 km segment. In the Tonga arc earthquakes are distributed more evenly but are less frequent than in the Kermadec arc. The last large shock in the central portion of the Tonga arc occurred in 1949. This uneven distribution may not be random as the gaps in activity lie landward of regions subducting relatively shallow seafloor. As discussed earlier, the subduction of bathymetric highs may inhibit the subduction process; regions interacting with bathymetric highs typically lack large shocks, and in some cases gaps appear in the volcanic chain. Note the

good correlation among the presence of volcanoes, large shocks and low lying seafloor ($> 3,000$ fathoms) for the Tonga and Kermadec arcs.

5. Discussion

(a) Successes and Problems

Seismic gap analysis has led to the successful forecasting of the sites of a number of large earthquakes (Table 1). In each case the area was discussed in the literature prior to occurrence of the shock. No great earthquakes have occurred in at least the last 5 years on segments of plate boundaries that were thought to be of low seismic potential. The general success of this method can probably be attributed to the regular manner in which large shocks occur along the shallow portions of simple plate boundaries.

Two unusual events, however, occurred in the Kuril Islands in 1975 and 1976. Their rupture zones are located near a segment of the arc that ruptured during a somewhat deeper-than-normal (about 50 km) event in 1958. Although each event was of magnitude $M = 7.0$, they were associated with a large number of aftershocks that were spread over about a 400 km segment of the plate boundary. Events of this size cannot be forecast for this region because most of the seismic energy release occurs during great earthquakes ($M \geq 7 \frac{3}{4}$). Thus, despite the large aftershock areas, these events probably did not contribute significantly to the cumulative seismic motion between the plates.

KANAMORI (1977a) examined seismograms of the 1975 event and found an excess of long-period energy (20 sec waves) compared to that for shorter periods. Such an excess of long-period energy appears to be characteristic of a long rise time for motion at a given point on the rupture zone, which appears to be associated with larger than normal tsunamis. It may reflect a reduced coupling between the oceanic and continental lithospheric plates in that area (KANAMORI, 1977a). Kanamori also reports that a significant amount of plate motion in the southern Kuril Islands occurs aseismically.

Historic records of the southern Kuril Islands indicate that a portion of the plate boundary may not have ruptured during the sequence of great earthquakes in the 19th century. The one segment that may not have ruptured was the site of the large shock in 1958. This event is anomalous since the hypocenter of the main shock and the depths of aftershock activity (40-60 km) were usually deep. Slippage during the 1958 event was greater than that during each of the other events of the 20th century that ruptured segments affected in the 19th century (KANAMORI, 1978). The excess slippage experienced during that event, therefore, may have released strain energy stored during the 18th and 19th centuries but not released when the region was skipped during the great earthquakes of the 19th century. This region may then be capable of accumulating varying amounts of strain energy. This may reduce the reliability of the seismic gap techniques used here. In that area very old oceanic lithosphere is being subducted. That may account for the greater depth at which earthquakes are reflecting relative motion between plates. Hence, in that region the assumption that great shocks break the entire depth range of the plate interface may not always hold.

Problems are presented in applying gap theory to regions in which the relative plate motion occurs along a series of faults rather than a single boundary. Strain accumulation on these faults may not be a steady state process, and motion along these faults may be episodic, jumping from one fault to another in time. Although primary faults can be assigned to various categories of seismic potential, their splay faults cannot be accurately classified, and the presence of splay faults reduces our confidence in forecasts for these regions. Thus, some of the regions assigned to categories 2 or 6 may have a potential for large shocks to occur along splay faults that are subparallel to the main plate boundary.

(b) Future Research Using Existing Data

Much work still needs to be done to improve the historic record of many regions along plate boundaries. Those areas shown in category 3 in Fig. 1 could probably benefit most from a more complete historical record, as they could be placed into a more clearly defined category of seismic potential. Much of the historic record needs to be re-examined in the light of our present understanding of plate tectonics, and the hypotheses of aseismic plate movement and modified subduction. We also need a clearer understanding of the parameters involved in the generation of tsunamis. In some cases the damage from a tsunami is more extensive than that from the earthquake itself. If tsunamic earthquakes (KANAMORI, 1977a) could be regionally and tectonically categorized, tsunami risk could be more definitiely forecast. An examination of historic records appears to be one of the most effective methods for achieving a better understanding of tsunamis.

The larger and generally more complete historical record of earthquakes and tsunamis in Japan yield important information about the recurrence times of large shocks for particular segments of the plate boundary near Japan. Some information about repeat times of historic events or those deduced from geologic movements along particular faults during the past few thousand years is, of course, very much needed for most other plate boundaries. More precise repeat times would be preferable, in determining the seismic potential of a region, to the use of an arbitrary cutoff of 100 years. At present there is much scatter in many of the reported recurrence times even for regions with a long historical record. We must gain confidence in the completeness of the historical records; only then can we be certain that the apparent scatter in observed repeat times is not caused by the missing of events.

Variations in intraplate activity near active seismic zones need to be studied more thoroughly. CARR (1977) examined large shocks in Central America, differentiating between inter- and intraplate earthquakes. However, no conclusions were drawn from the variation in activity. The occurrence of intraplate events in either the frontal arc or volcanic arc regions may reflect the state of stress on the thrust zone. A recognizable spatial pattern may be observed before large thrust events as in Japan (SHIMAZAKI, 1976, 1978).

Since the advent of the World-Wide Standardized Seismograph Network in 1962, a large body of epicentral data has been collected for small and moderate size shocks. The spatial-temporal variations in this data set should be examined for each region of high seismic potential. Marked quiescence for small and moderate size shocks has also preceded many large shocks, examination of this data set may well reveal a similar behavior for some of the gaps in Fig. 1. A search of this and other types of premonitory phenomena should be made, especially for the regions assigned to category 1. One set of data that is easily researched is the level of volcanic activity for regions near seismic gaps. Variations in eruptive activity and in the level of lava lakes (NAKAMURA, 1975; KIMURA, 1976; CARR, 1977; KIMURA and McCANN, 1979) were observed before some great and major earthquakes. More research is needed to properly document the strength of known volcanic eruptions so as to reduce the large amount of what appears to be noise in the historic record of volcanic activity.

(c) Collection of New Data in Seismic Gaps

A concentrated effort is recommended to collect a variety of possible premonitory data for regions assigned to category 1 in Fig. 1. This would undoubtedly increase our chances of "catching" pre-seismic changes associated with large earthquakes. Detailed studies may also help us to identify those regions

100 years ago, the area is assigned to category 1. During this century large ($M \geq 7$) shallow shocks along the coast of Sumatra clustered into two groups separated by some 500 km. The northern half of the zone of quiescence that lies between these clusters was the site of moderate (i.e., $6 < M < 6.9$) events until 1955. A single event recently occurred in this area in 1970. The southern portion of the quiescent region has not had any shallow events of magnitude 6 or above since at least the 1930's and possibly much longer. Small earthquakes (i.e., $M < 6$) recorded since 1962 have distributed themselves rather evenly along the shallow portion of the seismic zone.

A zone of quiescence bounded by a region of high seismic activity has been observed prior to large thrust and strike-slip type earthquakes. The observations of the earthquake activity along the Sumatran coast are similar in several respects to those of previous researchers but there are some differences.

The zone of quiescence for moderate and large shocks has only persisted since 1955; for larger shocks the quiescence extends back to the beginning of instrumental coverage. Shallow shocks of the last 15 years appear to cover the region rather evenly despite the clustering of the larger events. Thus, more work is needed to fully understand the significance of the zone of quiescence for large shocks between 0° and 5°S , near the coast of Sumatra, as the spatial pattern of occurrence of large and moderate earthquakes is similar to those observed prior to great earthquakes.

Northern Lesser Antilles. Great earthquakes ruptured an extensive portion of the northern Lesser Antillean arc in 1690 and 1843. In the absence of other data it is not unreasonable to take the time interval between these events as an indication of a recurrence time of about 150 years. In 1969 and 1974 large intraplate events occurred on the edges of the rupture area of the 1843 event. An increase of shallow intraplate activity within Japan was observed prior to

in category 1 that are highly stressed. In-situ stress measurements using the hydrofracturing technique and determinations of stress drops for moderate size shocks could provide data pertinent to understanding the state of stress in seismic gaps or of possible stress concentrations near the edges of seismic gaps. HOUSE and BOATWRIGHT (1979) infer stress drops of about 500 bars for two moderate size shocks in the seismic gap near the Shumagin Islands and the Alaska Peninsula. Their study brings up the exciting possibility of ascertaining the seismic potential for large shocks from the state of stress in major seismic gaps.

Many of the regions assigned to the highest categories of risk in Fig. 1 are situated largely in submarine areas along active continental margins. There is a great need to develop the capability of measuring crustal movements (both co-seismic and premonitory) and strong ground motion as well as to record small shocks on the ocean floor in some of the critical seismic gaps of Fig. 1.

Many seismic gaps, including those in category 1, are only sparsely instrumented. Several strong-motion instruments could be deployed and level lines constructed in many of the areas designated as category 1 with a relatively small expense.

(d) Regions of Particular Interest

Several of the regions assigned high seismic potentials in Fig. 1 (categories 1 to 3) are of particular interest since some observations indicate that these regions may be exhibiting some types of long-term premonitory behavior which may be indicative of future large shocks.

Sumatra. The last great earthquake along the Sumatran coast occurred in 1861. It was associated with a seawave that destroyed coastal villages for several hundred kilometers. Since the last great earthquake occurred more than

several large earthquakes along the nearby plate boundary (SHIMAZAKI, 1976, 1978). These events in Japan may be indicative of higher stress within the overriding plate prior to the occurrence of large interplate shocks. The intraplate shocks in the Lesser Antilles may represent a similar phenomenon.

Grande Soufrière on Guadeloupe in the northern Lesser Antilles has been active during the past few years. CARR (1977) suggests that this increase in volcanic activity may be indicative of a forthcoming great earthquake. That volcano erupted a few years before the last great shock in 1843.

Shumagin Islands, Alaska. This region has an unclear seismic history but may be the site of a future large shock. As discussed previously this region experienced events with high stress drops as inferred from the ratio M_s/M_b and from information derived from local strong-motion records. This area was cited by KELLEHER (1970) as the next in a space-time progression of events that has moved north along the southeast coast of Alaska and then west along its southern coast. The volcano Pavlof (near 162°W, on the Alaskan Peninsula), which is adjacent to the gap, has shown increased eruptive activity in the last few years. Thus, despite the incomplete seismic history of this region there are several observations that could be interpreted as indicators of a high level of tectonic stress.

6. Summary

Plate tectonics has given us a fuller understanding of the mechanism by which strain energy is stored on plate boundaries and then suddenly released as large earthquakes. This understanding has led to successful forecasts of the sites of several large shocks along convergent and strike-slip boundaries. In this paper simple, major plate boundaries of the Pacific, Caribbean, Sunda and South Sandwich regions are classified into six categories of seismic potential for large earthquakes for the next few decades. These categories reflect our knowledge and assessment of the historic record for large earthquakes, our understanding and speculations about tectonic regimes and the length of time since the last large earthquake. Several hypotheses are used to infer the seismic potential for regions that have an ambiguous history of great shocks. Since categories 3 and 5 are based on hypotheses and not on proven fact, the assignment of regions to those categories should be regarded as tentative. The degree of seismic potential is more certain for areas that have experienced large earthquakes, i.e., for categories 1, 2 and 6. Several of the areas assigned the highest risk are poorly instrumented and obviously deserve high priority for intense study.

Acknowledgements

We would like to thank K. Jacob and T. Johnson for reviewing the manuscript. The first two authors wish to thank John Kelleher for his inspiration and support during their apprentice years with him. We also wish to thank Margaret Winslow for furnishing unpublished information about southern South America. Linda Murphy typed the manuscript and Kazuko Nagao drafted several of the figures. This research was supported by the Division of Earth Sciences, National Science Foundation, NSF Grant EAR 75-03640; U.S. Department of the Interior Geological Survey Number USGS-14-08-0001-16748; and the Energy Research and Development Authority Contract Number EY-76-S-02-3134B.

- ALGUE, REV. JOSE (1909), Mirador Observatory Baguio, Benguet: A new meteorological-geodynamic station of the weather burea, Manila, Philippines, Bureau of Printing, 26-27.
- ALLEN, C. R. (1962), Circum-Pacific faulting in the Philippines-Taiwan region, J. Geophys. Res. 67, 4795-4812.
- ALLEN, C. R. (1968), The tectonic environments of seismically active and inactive areas along the San Andreas fault system, Proceeding of a conference "Geological Problems of the San Andreas Fault", Stanford Univ. Publs. Geol. Sci. 11, 70-83.
- ALLEN, C. R., ST. ARMAND, P., RICHTER, C. F. and NORDQUIST, J. M. (1965), Relationships between seismicity and geologic structure in the southern California region, Bull. Seism. Soc. Amer. 55, 753-797.
- ANDO, M. (1975), Source mechanisms and tectonic significance of historical earthquakes along the Nankai Trough, Japan, Tectonophys. 27, 119-140.
- ARCHAMBEAU, C. (1978), Estimation of non-hydrostatic stress in the earth by seismic methods: Lithospheric stress levels along Pacific and Nazca plate subduction zones, this volume.
- BARAZANGI, M. and ISACKS, B. L. (1976), Spatial distribution of earthquakes and subduction of the Nazca plate beneath South America, Geology 4, 686-692.
- BRUNE, J. (1968), Seismic moment, seismicity and rate of slip along major fault zones, J. Geophys. Res. 73, 777-784.
- CARDWELL, R. K. and ISACKS, B. L. (1978), Geometry of the subducted lithosphere beneath the Banda Sea in eastern Indonesia from seismicity and fault plane solutions, J. Geophys. Res. 83, 2825-2838.
- CARR, M. J. (1974), Tectonics of the Pacific margin of northern Central America, Ph.D. dissertation, Dartmouth College, Hanover, N.H. 159 p.
- CARR, M. J. (1977), Volcanic activity and great earthquakes at convergent plate margins, Science 197, 655-657.

- CARR, M. J., STOIBER, R. E. and DRAKE, C. L. (1974), The segmented nature of some continental margins, in C. A. Burk and C. L. Drake, eds., The Geology of Continental Margins, Springer-Verlag, 105-114.
- CARR, M. J. and STOIBER, R. E. (1977), Geologic setting of some destructive earthquakes in Central America, Geol. Soc. Amer. Bull. 88, 151-156.
- CHASE, C., MENARD, H. and MAMMERICKX, J. (1970), Bathymetry of the North Pacific, Charts 1-6, Scripps Inst. of Oceanogr. Inst. of Mar. Resour., LaJolla, Calif.
- COOPER, A K., MARLOW, M. S. and SCHOLL, D. W. (1977), The Bering Sea - A multifarious marginal basin, Maurice Ewing Series No. 1, Amer. Geophys. Union, Washington, D. C., 437-450.
- CORMIER, V. (1975), Tectonics near the junction of the Aleutian and Kuril-Kamchatka arcs and a mechanism for middle Tertiary magmatism in the Kamchatka Basin, Geol. Soc. Amer. Bull. 86, 443-453.
- COX, D. and PARARAS-CARYANNIS, G. (1976), Catalog of Tsunamis in Alaska, U.S. Dept. of Commerce, NOAA/EDS, Asheville, N.C., 78p.
- DAVIES, J. N. and HOUSE, L. (1979), Aleutian subduction zone seismicity, volcano-trench separation, and their relation to great thrust-type earthquakes, in preparation.
- DEAN, B. W. and DRAKE, C. L. (1978), Focal mechanism solutions and tectonics of the Middle America arc, J. Geol. 86, 111-128.
- DE LOCZY, L. (1968), Geotectonic evolution of the Amazon, Parnaiba and Parna basins, An. Acad. brasil. Cieñc 40, 231-249.
- ELLA, Rev. SAMUEL (1890), Some physical phenomena of the South Pacific Islands, in Report of the Second Meeting of the Australian Association for the Advancement of Science, Melbourne, Australia, 559-572.
- EPPLEY, R. A. (1965), Earthquake History of the United States Part I, U. S. Dept. of Commerce, Env. Sci. Serv. Admin. Coast and Geod. Surv., 120 p.

- ALGUE, REV. JOSE (1909), Mirador Observatory Baguio, Benguet: A new meteorological-geodynamic station of the weather burea, Manila, Philippines, Bureau of Printing, 26-27.
- ALLEN, C. R. (1962), Circum-Pacific faulting in the Philippines-Taiwan region, J. Geophys. Res. 67, 4795-4812.
- ALLEN, C. R. (1968), The tectonic environments of seismically active and inactive areas along the San Andreas fault system, Proceeding of a conference "Geological Problems of the San Andreas Fault", Stanford Univ. Publs. Geol. Sci. 11, 70-83.
- ALLEN, C. R., ST. ARMAND, P., RICHTER, C. F. and NORDQUIST, J. M. (1965), Relationships between seismicity and geologic structure in the southern California region, Bull. Seism. Soc. Amer. 55, 753-797.
- ANDO, M. (1975), Source mechanisms and tectonic significance of historical earthquakes along the Nankai Trough, Japan, Tectonophys. 27, 119-140.
- ARCHAMBEAU, C. (1978), Estimation of non-hydrostatic stress in the earth by seismic methods: Lithospheric stress levels along Pacific and Nazca plate subduction zones, this volume.
- BARAZANGI, M. and ISACKS, B. L. (1976), Spatial distribution of earthquakes and subduction of the Nazca plate beneath South America, Geology 4, 686-692.
- BRUNE, J. (1968), Seismic moment, seismicity and rate of slip along major fault zones, J. Geophys. Res. 73, 777-784.
- CARDWELL, R. K. and ISACKS, B. L. (1978), Geometry of the subducted lithosphere beneath the Banda Sea in eastern Indonesia from seismicity and fault plane solutions, J. Geophys. Res. 83, 2825-2838.
- CARR, M. J. (1974), Tectonics of the Pacific margin of northern Central America, Ph.D. dissertation, Dartmouth College, Hanover, N.H. 159 p.
- CARR, M. J. (1977), Volcanic activity and great earthquakes at convergent plate margins, Science 197, 655-657.

- CARR, M. J., STOIBER, R. E. and DRAKE, C. L. (1974), The segmented nature of some continental margins, in C. A. Burk and C. L. Drake, eds., The Geology of Continental Margins, Springer-Verlag, 105-114.
- CARR, M. J. and STOIBER, R. E. (1977), Geologic setting of some destructive earthquakes in Central America, Geol. Soc. Amer. Bull. 88, 151-156.
- CHASE, C., MENARD, H. and MAMMERICKX, J. (1970), Bathymetry of the North Pacific, Charts 1-6, Scripps Inst. of Oceanogr. Inst. of Mar. Resour., LaJolla, Calif.
- COOPER, A K., MARLOW, M. S. and SCHOLL, D. W. (1977), The Bering Sea - A multifarious marginal basin, Maurice Ewing Series No. 1, Amer. Geophys. Union, Washington, D. C., 437-450.
- CORMIER, V. (1975), Tectonics near the junction of the Aleutian and Kuril-Kamchatka arcs and a mechanism for middle Tertiary magmatism in the Kamchatka Basin, Geol. Soc. Amer. Bull. 86, 443-453.
- COX, D. and PARARAS-CARYANNIS, G. (1976), Catalog of Tsunamis in Alaska, U.S. Dept. of Commerce, NOAA/EDS, Asheville, N.C., 78p.
- DAVIES, J. N. and HOUSE, L. (1979), Aleutian subduction zone seismicity, volcano-trench separation, and their relation to great thrust-type earthquakes, in preparation.
- DEAN, B. W. and DRAKE, C. L. (1978), Focal mechanism solutions and tectonics of the Middle America arc, J. Geol. 86, 111-128.
- DE LOCZY, L. (1968), Geotectonic evolution of the Amazon, Parnaiba and Parna basins, An. Acad. brasil. Cieñc 40, 231-249.
- ELLA, Rev. SAMUEL (1890), Some physical phenomena of the South Pacific Islands, in Report of the Second Meeting of the Australian Association for the Advancement of Science, Melbourne, Australia, 559-572.
- EPPLEY, R. A. (1965), Earthquake History of the United States Part I, U. S. Dept. of Commerce, Env. Sci. Serv. Admin. Coast and Geod. Surv., 120 p.

- FEDOTOV, S. A. (1965), Regularities of the distribution of strong earthquakes in Kamchatka, the Kuril Islands, and northeast Japan, Trudy Inst. Fiz Zemli., Acad. Nauk. SSSR 36, 66-93.
- FISHER, R. L. (1961), Middle America trench: Topography and structure, Geol. Soc. Amer. Bull. 72, 703-720.
- FITCH, T. J. (1972), Plate convergence, transcurrent faults, and internal deformation adjacent to Southeast Asia and the Western Pacific, J. Geophys. Res. 77, 4432-4460.
- FITCH, T. J. and MOLNAR, P. (1970), Focal mechanisms along inclined earthquake zones in the Indonesian-Philippine region, J. Geophys. Res. 75, 1431-1444.
- FORSYTH, D. W. (1975), Fault-plane solutions and tectonics of the South Atlantic and Scotia Sea, J. Geophys. Res. 80, 1429-1443.
- FRANKEL, A. and McCANN, W. (1978), Seismicity of the South Sandwich arc: Consequences of a ridge-trench collision, J. Geophys. Res. (in press).
- GANSSEER, A. (1973), Facts and theories on the Andes, J. Geol. Soc. London 129, 93-131.
- GARZA, T. and LOMNITZ, C. (1978), The Oaxaca gap: A case history, in this volume.
- GELLER, R. and KANAMORI, H. (1972), Magnitudes of great shallow earthquakes from 1904 to 1952, Bull. Seism. Soc. Amer. 67, 587-598.
- GUTENBERG, B. and RICHTER, C. F. (1954), Seismicity of the Earth (2nd edit. Princeton Univ. Press, New Jersey 1954), pp. 161-234.
- HATORI, T. (1969), Dimensions and geographical distribution of tsunami sources near Japan, Bull. Earthq. Res. Inst., Univ. Tokyo 47, 185-214.

- HATORI, T. (1975), Sources of tsunamis generated off Boso Peninsula, Bull. Earthq. Res. Inst., Univ. Tokyo 50, 83-91.
- HERRON, E. M. and TUCHOLKE, B. E. (1976), Sea-floor magnetic patterns and basement structure in the southeastern Pacific in Initial Reports of the Deep Sea Drilling Project 35 (U. S. Govt. Printing Office, Washington, D.C. 1976), pp. 263-278.
- HOUSE, L. and BOATWRIGHT, J. (1979), Investigation of two high-stress drop earthquakes in the Shumagin seismic gap, Alaska (in preparation).
- International Seismic Summary (Kew Observatory, Richmond, Surrey, England 1949, 1950).
- ISACKS, B., OLIVER, J. and SYKES, L. (1968), Seismology and the new global tectonics, J. Geophys. Res. 73, 5855-5899.
- JACOB, K., NAKAMURA, K. and DAVIES, J. N. (1977), Trench-volcano gap along the Alaska-Aleutian arc: Facts and speculations on the role of terrigenous sediments for subduction in Island Arcs, Deep Sea Trenches and Back-Arc Basins (Talwani, M. and Pitman III, W. C., eds., Maurice Ewing Series I, Amer. Geophys. Union), pp. 243-258.
- JAHNS, R. (1977), Study of Holocene displacement history along the south-central reach of the San Andreas fault, USGS Final Report #14-08-0001-15225, 1-219.
- JOHNSON, T. and MOLNAR, P. (1972), Focal mechanisms and plate tectonics of the southwest Pacific, J. Geophys. Res. 77, 5000-5032.
- JORDAN, T. (1975), The present-day motions of the Caribbean plate, J. Geophys. Res. 80, 4433-4439.
- KANAMORI, H. (1971), Great earthquakes at island arcs and the lithosphere, Tectonophysics 12, 187-198.
- KANAMORI, H. (1977a), Seismic and aseismic slip along subduction zones and their tectonic implications in Island Arcs, Deep Sea Trenches and Back-Arc Basins (Talwani, M. and Pitman III, W. C., eds., Maurice Ewing Series I, Amer. Geophys. Union), pp. 273-284.

- FEDOTOV, S. A. (1965), Regularities of the distribution of strong earthquakes in Kamchatka, the Kuril Islands, and northeast Japan, Trudy Inst. Fiz Zemli., Acad. Nauk. SSSR 36, 66-93.
- FISHER, R. L. (1961), Middle America trench: Topography and structure, Geol. Soc. Amer. Bull. 72, 703-720.
- FITCH, T. J. (1972), Plate convergence, transcurrent faults, and internal deformation adjacent to Southeast Asia and the Western Pacific, J. Geophys. Res. 77, 4432-4460.
- FITCH, T. J. and MOLNAR, P. (1970), Focal mechanisms along inclined earthquake zones in the Indonesian-Philippine region, J. Geophys. Res. 75, 1431-1444.
- FORSYTH, D. W. (1975), Fault-plane solutions and tectonics of the South Atlantic and Scotia Sea, J. Geophys. Res. 80, 1429-1443.
- FRANKEL, A. and McCANN, W. (1978), Seismicity of the South Sandwich arc: Consequences of a ridge-trench collision, J. Geophys. Res. (in press).
- GANSSEER, A. (1973), Facts and theories on the Andes, J. Geol. Soc. London 129, 93-131.
- GARZA, T. and LOMNITZ, C. (1978), The Oaxaca gap: A case history, in this volume.
- GELLER, R. and KANAMORI, H. (1972), Magnitudes of great shallow earthquakes from 1904 to 1952, Bull. Seism. Soc. Amer. 67, 587-598.
- GUTENBERG, B. and RICHTER, C. F. (1954), Seismicity of the Earth (2nd edit. Princeton Univ. Press, New Jersey 1954), pp. 161-234.
- HATORI, T. (1969), Dimensions and geographical distribution of tsunami sources near Japan, Bull. Earthq. Res. Inst., Univ. Tokyo 47, 185-214.

- HATORI, T. (1975), Sources of tsunamis generated off Boso Peninsula, Bull. Earthq. Res. Inst., Univ. Tokyo 50, 83-91.
- HERRON, E. M. and TUCHOLKE, B. E. (1976), Sea-floor magnetic patterns and basement structure in the southeastern Pacific in Initial Reports of the Deep Sea Drilling Project 35 (U. S. Govt. Printing Office, Washington, D.C. 1976), pp. 263-278.
- HOUSE, L. and BOATWRIGHT, J. (1979), Investigation of two high-stress drop earthquakes in the Shumagin seismic gap, Alaska (in preparation).
- International Seismic Summary (Kew Observatory, Richmond, Surrey, England 1949, 1950).
- ISACKS, B., OLIVER, J. and SYKES, L. (1968), Seismology and the new global tectonics, J. Geophys. Res. 73, 5855-5899.
- JACOB, K., NAKAMURA, K. and DAVIES, J. N. (1977), Trench-volcano gap along the Alaska-Aleutian arc: Facts and speculations on the role of terrigenous sediments for subduction in Island Arcs, Deep Sea Trenches and Back-Arc Basins (Talwani, M. and Pitman III, W. C., eds., Maurice Ewing Series I, Amer. Geophys. Union), pp. 243-258.
- JAHNS, R. (1977), Study of Holocene displacement history along the south-central reach of the San Andreas fault, USGS Final Report #14-08-0001-15225, 1-219.
- JOHNSON, T. and MOLNAR, P. (1972), Focal mechanisms and plate tectonics of the southwest Pacific, J. Geophys. Res. 77, 5000-5032.
- JORDAN, T. (1975), The present-day motions of the Caribbean plate, J. Geophys. Res. 80, 4433-4439.
- KANAMORI, H. (1971), Great earthquakes at island arcs and the lithosphere, Tectonophysics 12, 187-198.
- KANAMORI, H. (1977a), Seismic and aseismic slip along subduction zones and their tectonic implications in Islands Arcs, Deep Sea Trenches and Back-Arc Basins (Talwani, M. and Pitman III, W. C., eds., Maurice Ewing Series I, Amer. Geophys. Union), pp. 273-284.

- KANAMORI, H. (1977b), The energy release in great earthquakes, J. Geophys. Res., 82, 2981-2987.
- KANAMORI, H. (1978), Nature of seismic gaps and foreshocks, in this volume.
- KANAMORI, H. and ABE, K. (1978), Re-evaluation of the turn-of-the-century seismicity peak using an undamped Milne seismograph, EOS, Spring AGU Abstracts, April, p. 327.
- KARIG, D.E., (1973), Plate convergence between the Philippines and the Ryukyu Islands, Mar. Geol., 14, 153-168.
- KARIG, D.E. and MAMMERICKX, J. (1972), Tectonic framework of the New Hebrides Island arc, Mar. Geol., 12, 187-205.
- KATILI, J. A. (1970), Large transcurrent faults in Southeast Asia with special reference to Indonesia, Geologische Rundschau, 59, 581-600.
- KATSUMATA, M. and SYKES, L.R. (1969), Seismicity and tectonics of the western Pacific: Izu-Mariana-Caroline and Ryukyu-Taiwan regions, J. Geophys. Res., 74, 5923-5948.
- KATZ, H.R. (1964), Some new concepts on geosynclinal development and mountain building of the southern end of South America, Proc. 22nd Internat. Geol. Cong., New Dehli, 4, 241-255.
- KELLEHER, J.A. (1970), Space-time seismicity of the Alaska-Aleutian seismic zone, J. Geophys. Res., 75, 5745-5756.
- KELLEHER, J.A. (1972), Rupture zones of large South American earthquakes and some predictions, J. Geophys. Res., 77, 2087-2103.
- KELLEHER, J.A., SYKES, L.R. and OLIVER, J. (1973), Possible criteria for predicting earthquake locations and their applications to major plate boundaries of the Pacific and Caribbean, J. Geophys. Res., 78, 2547-2585.

- KELLEHER, J.A., SAVINO, J., ROWLETT, H. and McCANN, W. (1974), Why and where great thrust earthquakes occur along island arcs, J. Geophys. Res., 79 4889-4899.
- KELLEHER, J.A. and SAVINO, J. (1975), Distribution of seismicity before large strike-slip and thrust-type earthquakes, J. Geophys. Res., 80, 260-271.
- KELLEHER, J.A. and McCANN, W. (1976), Buoyant zones, great earthquakes and unstable boundaries of subduction, J. Geophys. Res., 81, 4885-4908.
- KELLEHER, J.A. and McCANN, W. (1977), Bathymetric highs and the development of convergent plate boundaries, in (ed. M. Talwani and W.C. Pitman III) Maurice Ewing Series I: Island Arcs, Deep Sea Trenches and Back Arc Basins, AGU, 115-122.
- KIMURA, M. (1976), Major magnetic activity as a key to predicting large earthquakes along the Sagami trough, Japan, Nature, 260, 131-133.
- KIMURA, M. and McCANN, W. (1979), Relation between great earthquakes and eruptive activity in the circum-Pacific area, in preparation.
- KONDORSKYA, N.V. and SHEBALIN, N.V. (eds.) (1977), New Catalog of Strong Earthquakes in the Territory of the USSR, Nauka, Moscow, USSR, p. 374-433.
- KULM, L.D., SCHWELLER, W.J. and MASIAS, A. (1977), A preliminary analysis of the subduction process along the Andean continental margin, 6° to 45°S, in (ed. M. Talwani and W.C. Pitman III) Maurice Ewing Series I: Island Arcs, Deep Sea Trenches and Back Arc Basins, AGU, 285-302.
- LePICHON, X. (1968), Sea floor spreading and continental drift, J. Geophys. Res., 73, 3661-3697.
- LOMNITZ, C. (1970), Major earthquakes and tsunamis in Chile during the period 1535 to 1955, Geol. Rundschau, 59, 938-960.
- McCANN, W. and NISHENKO, S. (1978), Large earthquakes and the tectonics of the western Pacific - past, present and future, Internat. Geodynamics Conf. 'Western Pacific and Magma Genesis', Tokyo, 100-101.

- McCANN, W. and KELLEHER, J. (1979), Moderate and large earthquakes and their relationship to the tectonics of the southwest Pacific (in preparation).
- McCANN, W., MURPHY, A. and FRANKEL, A. (1979), Small earthquakes and the tectonics of the Virgin Islands region (in preparation).
- MEDVEDEV, S. V. (ed.) (1968), Seismic Zoning of the USSR (Izdatel'stvo Nauka, Moscow 1968), pp. 460-467 (Trans. by Teteruk-Schneider, R. in Hardin, R. (ed.), Israel Progr. Sci. Trans., Jerusalem 1976).
- MEYERS, H., BROZEE, R., COFFMAN, J. and LESSIG, S. (1976), An analysis of earthquake intensities and recurrence rates in and near Alaska, NOAA Tech. Rept. EDS-NGSDC-3, 79 p.
- MINSTER, J. B., JORDAN, T. H., MOLNAR, P. and HAINES, E. (1974), Numerical modelling of instantaneous plate tectonics, Geophys. J. Roy. astronom. Soc. 36, 541-576.
- MOGI, K. (1968), Some features of recent seismic activity in and near Japan (1), Bull. Earthq. Res. Inst., Univ. Tokyo 46, 1225-1236.
- MOGI, K. (1969), Some features of recent seismic activity in and near Japan (2), Bull. Earthq. Res. Inst., Univ. Tokyo 47, 395-417.
- MOLNAR, P. and SYKES, L. R. (1969), Tectonics of the Caribbean and Middle America regions from focal mechanisms and seismicity, Geol. Soc. Amer. Bull. 80, 1639-1684.
- MORGAN, W. J. (1972), Plate motion and deep mantle convection in Studies in Earth and Space Science, GSA Memoir 132, Shagarn, R. (ed.).
- MURPHY, A. and McCANN, W. (1979), Preliminary results from a new seismic network in the northeast Caribbean (in preparation).
- NAKAMURA, K. (1975), Volcano structure and possible mechanical correlation between volcanic eruptions and earthquakes, Bull. Volcanol. Soc. Japan, Ser. 2, 20, 229-240.

- NISHENKO, S. and McCANN, W. (1977), Deep sea terraces: Indicators of great thrust earthquake history (abstract), EOS, Trans. Amer. Geophys. Union 58, 437.
- NISHENKO, S. and McCANN, W. (1978), Large thrust earthquakes and tsunamis: Implications for the development of fore-arc basins, J. Geophys. Res. (in press).
- OHTAKE, M., MATUMOTO, T. and LATHAM, G. V. (1977), Seismicity gap near Oaxaca, southern Mexico as a probable precursor to a large earthquake, Pure appl. Geophys 115, 375-385.
- REID, H. (1910), The California earthquake of April 18, 1906 in The Mechanics of the Earthquake, Report of the State Earthquake Investigation Comm. 2, pp. 16-19, Carnegie Inst. Washington, D.C.
- REID, H. and TABER, S. (1920), The Virgin Islands earthquakes of 1867-1868, Bull. Seism. Soc. Amer. 10, 9-30.
- REYES, A., BRUNE, J. N. and LOMNITZ, C. (1978), Source mechanism and after-shock study of the Colima, Mexico earthquake of January 10, 1973, in this volume.
- RICHTER, C. F. (1958), Elementary Seismology (W. H. Freeman, San Francisco), 768 p.
- RODOLFO, K. S. (1969), Bathymetry and marine geology of the Andaman Basin and tectonic implications for Southeast Asia, Geol. Soc. Amer. Bull. 80, 1203-1230.
- ROTHER, J. (1969), The Seismicity of the Earth 1953-1965 (Unesco, Paris), p. 336.
- ROWLETT, H. and KELLEHER, J. (1976), Evolving seismic and tectonic patterns along the western margin of the Philippine Sea plate, J. Geophys. Res. 81, 3518-3524.
- SAVAGE, J. C. and BURFORD, R. O. (1973), Geodetic determination of relative plate motions in central California, J. Geophys. Res. 78, 832-845.

- SCHOLZ, C. H., BARAZANGI, M. and SBAR, M.L. (1971), Late Cenozoic evolution of the Great Basin western United States as an ensialic inter-arc basin, Geol. Soc. Amer. Bull. 82, 2979-2990.
- SHIMAZAKI, K. (1978), Correlation between intraplate seismicity and interplate earthquakes in Tohoku, northeast Japan, Bull. Seism. Soc. Amer. 68, 181-192.
- SHIMAZAKI, K. (1976), Intra-plate seismicity gap along the Median Tectonic Line and oblique plate convergence in southwest Japan, Tectonophys. 31, 139-156.
- SINDVAHL, H., KHATTTRI, K. N., RAI, K. and GAUR, V. K. (1978), Neo-tectonics and space-time seismicity of the Andaman-Nicobar region, Bull. Seism. Soc. Amer. 68, 399-409.
- SOEDATI, K. (1963), Seismic zones in Indonesia, Geophys. Notes No. 2, Dept. Perhubungan Udara-Duawatan Meteorologi dan Geofisik, Duakarta, Indonesia, 10 p.
- STAUDER, W. (1973), Mechanism and spatial distribution of Chilean earthquakes with relation to subduction of the oceanic plate, J. Geophys. Res. 78, 5033-5061.
- SYKES, L. R. (1971), Aftershock zones of great earthquakes, seismicity gaps and earthquake prediction for Alaska and the Aleutians, J. Geophys. Res. 76, 8021-8041.
- TABER, S. (1920), Jamaica earthquakes and the Bartlett Trough, Bull. Seism. Soc. Amer. 10, 55-89.
- TABER, S. (1922), The Great Fault Troughs of the Antilles, J. Geol. 30, 89-114.
- TARR, R. and MARTIN, L. (1912), The earthquakes at Yakutat Bay, Alaska in September, 1899, U.S. Geol. Surv. Prof. Pap. 69, 1-131.
- THATCHER, W., HILEMAN, J. and HANKS, T. (1975), Seismic slip distribution along the San Jacinto fault zone, Southern California, and its implications, Bull. Geol. Soc. Amer. 86, 1140-1146.
- THATCHER, W. and G. PLAFKER (1977), The Yakutat Bay, Alaska, earthquakes: seismograms and crustal deformation, Geol. Soc. Amer. Abs. w. Prog. 9, p515.

- TOBIN, D. G. and SYKES, L. R. (1968), Seismicity and tectonics of the north-east Pacific Ocean, J. Geophys. Res. 73, 3821-3845.
- UTSU, T. (1974), Space-time pattern of large earthquakes occurring off the Pacific coast of the Japanese Islands, J. Phys. Earth 22, 325-342, 1974.
- UYEDA, S. and KANAMORI, H. (1978), Back-arc opening and the mode of subduction, this volume.
- VISSER, S. W. (1922), Inland and submarine epicenters of Sumatra and Java earthquakes, Koninklijk Magnetisch en Meteorologisch Observatorium te Batavia, Verhandelingen, 9, 14 p.
- VOGT, P., LOWRIE, A., BRACE, D. and HEY, R. (1976), Subduction of aseismic oceanic ridges: Effects on shape, seismicity, and other characteristics of consuming plate boundaries, Geol. Soc. Amer. Spec. Pap. 172, 59 p.
- WINSLOW, M. A. (1976), Active transcurrent shear zones in Southern Chile as landward expressions of transform plate boundaries, Geol. Soc. Amer. Abs. w. Prog. 1976 Ann. Meeting, Denver, Col. p 1173.
- WINSLOW, M. A. (1978), The structural evolution of the Magallanes Basin and neotectonics in the southernmost Andes, SCAR - Third Symp. on Antarctic Geol. and Geophys., Madison, Wisc. held Aug. 1977, p 113.
- WINSLOW, M. A. (1979), Late Cenozoic tectonic history of Tienadel Fuego, Chile, in preparation.

Patterns of Seismicity Preceding Earthquakes in Central America, Mexico, and California

by

Masakazu Ohtake⁽¹⁾

Tosimatu Matumoto⁽²⁾

Gary V. Latham⁽²⁾

INTRODUCTION

Patterns of seismic activity preceding the 15 largest earthquakes that have occurred since 1964 in Mexico, Central America, and California have been examined. The results, summarized in this paper, have been presented in two published papers (Ohtake *et al.*, 1977a and 1977b) and one oral presentation (Ohtake *et al.*, 1977c). We emphasize the California results in the body of this paper, describing the previously published material in summary form only.

The patterns found in the cases studied, include features recognized by many previous investigators. In general, quiescence developed in the region of the eventual shock, followed by resumption of activity leading up to the main shock. We refer to the quiescent period as the α stage, and to the interval of renewed activity as the β stage. The diameter of the region of anomalous seismicity ranged from 200 km to 300 km for the cases studied. The onset of quiescence (α stage) preceded the main shock by intervals ranging between 3 months and 5 years. Renewal of activity

University of Texas, Marine Science Institute, Contribution No. XXX

(1) National Research Center for Disaster Prevention, Sakura-mura, Niihari-gun, Ibaraki Pref, Japan

(2) University of Texas, Marine Science Institute, Geophysics Lab, Galveston, Texas 77550, U.S.A.

was detected in every case but one. The onset of β -stage activity preceded the main shock by intervals ranging between 8 days and 2 years. The case in which no β -stage activity was detected, was a magnitude 6.5 earthquake that occurred in northern Costa Rica in 1973.

There appears to be some correlation between the total interval of anomalous seismicity (α plus β) and the magnitude of the main shock; whereas little or no correlation is found between the dimensions of the zone of anomalous seismicity and the magnitude of the main shock. Our results suggest that resumption of activity in a previously quiescent zone may be one of the least ambiguous signals that failure is about to occur. The patterns of seismic activity will undoubtedly vary for differing tectonic regimes. However, if adequate consistency in the pattern for a given region can be established, the following surveillance strategy can be considered:

1. Detection of the onset of unusual quiescence by stations of a permanent regional network.
2. Deployment of a temporary network of sensors (seismometers, tiltmeters, etc.) within the zone of anomalous activity.
3. Intensification of monitoring efforts when significant resumption of activity occurs (or when other precursory phenomena, yet to be determined, are detected).

A region in southern Mexico that may be in the α stage of activity has been identified. An opportunity to test this strategy is offered

through this finding. A crucial point, of course, is whether large earthquakes do occur without precursory earthquakes. Kelleher and Savino (1975), for example, reported that prior seismic activity was not detected for the magnitude 7.6 (Ms) Sitka earthquake of 1972; although, they did find precursory seismic activity in other cases. Aftershocks with magnitudes as small as 2.5 were detected by the local seismic station following the Sitka earthquake. This, presumably would be the lower limit on detectable precursory activity. Experiments in rock mechanics reported by Mogi (1963) suggests that variations in foreshock activity might be a reasonable expectation. His results showed that foreshocks are much more numerous, and of greater intensity for heterogeneous media than for homogeneous media. This result suggests that the degree of structural irregularity in the affected zone may be the principal influence on precursory seismic activity. However, since there is no such thing as a homogeneous medium in nature, we must hold open the possibility that foreshocks do occur in all cases, but have been too small to be detected in some instances.

DATA FOR CALIFORNIA EARTHQUAKES

The locations of the four California earthquakes studied are shown in Figure 1. Pertinent parameters for these earthquakes are listed in Table 1. Several methods for displaying the patterns of seismic activity in the region of each epicenter have been tried. For the purposes of this discussion, we choose to show the space-time plots shown in Figures 2 through 5. Data for these plots were obtained from the

magnetic tape file of earthquake information issued by the Environmental Data Service of NOAA for the period 1900-1974. In each case, the epicentral distance from the main shock, up to a maximum range of 300 km, (ordinate) is plotted against the origin time (abscissa) of each event. The plots terminate at the time of occurrence of the main shock. Tentative identifications of the onsets of α and β stages of activity are indicated by arrows. The horizontal dashed line in each figure indicates the radius of the zone of anomalous seismicity. The corresponding gap dimensions and time intervals for α - and β -stages of activity are listed in Table 1. Two possible interpretations are shown for the San Fernando earthquake. Some aftershocks of the San Fernando earthquake, located about 70 km from the Oxnard (Pt. Mugu) epicenter, are included in the Oxnard plot.

THE OAXACA CASE

During the course of our study of the fluctuations in seismic activity in Mexico, we discovered an area of significant seismic quiescence in the state of Oaxaca, southern Mexico. The quiescence began in mid-1973 throughout a zone approximately 200 km long. Shallow earthquakes ($h < 60$ km) originating in this region during the 3-1/2 period preceding the onset of quiescence is compared with the 4 year period following that time in Figure 6. If we take the gap to be a zone 200 km wide, striking N22°E, i.e., perpendicular to the trend of the seismic belt, and centered within the present gap, the statistics of earthquakes originating within the zone are as given in Table 2. Complete quiescence

developed in the upper 60 km in this zone in mid-1973 and persists to the time of this writing: a total of five years. Activity also diminished at depths greater than 60 km, but the change is less pronounced.

Large earthquakes (magnitudes 7 to 7.5) occurred in the anomalous zone in 1965 and 1968. Based upon the focal mechanisms and aftershock zones of these earthquakes, we derived an estimate for the location and probable magnitude of a future earthquake. As yet, we have no reliable basis for estimating the time of occurrence. However, we note that the two preceding large earthquakes were separated by only three years, and that the time interval between onset of quiescence and main shock for earthquakes of Central America in the magnitude range from 7 to 7.5 (4 cases) was about 3 years. Quiescence in the Oaxaca zone persists to the present; a total of nearly 5 years.

Based upon the seismicity of Southern Mexico during the period 1963-1973, the mean return time for earthquakes of magnitude (M_s) 7.5, and larger, for a region of the size of the present gap, is 18 years (see Appendix II). Lomnitz (1974) lists 15 damaging earthquakes as having occurred in the Oaxaca region since 1800: an average rate of about one moderate to large earthquake every 12 years. Ten years have passed since the damaging earthquake of 1968.

We earlier estimated (Appendix I) that the probability that this gap could occur by chance in a 2-year period (the age of the gap at the time the estimate was made) was about 1 in 36,000. Considering the increase in period of quiescence to 5 years, the assertion of nonrandomness is further strengthened. We suggest that this zone represents an

excellent opportunity to apply the monitoring strategy outlined in the Introduction.

DISCUSSION

It is obvious that a large degree of subjectivity is involved in the assignment of parameters α and β to the data sets shown in figures 2-5, and in the cases previously examined (Ohtake *et al.*, 1977b). A strong case simply cannot be claimed from the limited data available. The obvious remedy for this problem is the installation of improved seismic networks to monitor active zones, coupled with improved processing methods. Despite these reservations, we believe that the results are suggestive enough that increased attention to the details of fluctuations in seismic activity is warranted.

Returning to the earthquake histories considered here, two additional points are worthy of note: (1) the epicenters of the main shocks are usually located near the center of the zone of anomalous seismicity; and (2) the aftershock zones do not fill the anomalous zone. We have not recognized any pattern to the spatial distribution of β -stage events, such as clustering toward the center, or around the edges of the anomalous zone. However, in view of the uncertainties in epicenter locations, and the relatively small numbers of events available for analysis in most cases, the existence of such patterns cannot be ruled out.

Assuming that our estimates of α and β intervals are valid, we next seek possible correlations between these parameters and main-shock magnitudes. The total interval of anomalous seismicity (α plus β) vs magnitude is plotted in figure 7. Also shown is an empirical curve

of precursory times for various phenomena developed by Rikitake (1976), and additional data points compiled by Ohtake (1977). The Rikitake curve is quite close to that earlier given by Scholz et al. (1973). While the precursory times for the four California shocks fall reasonably close to the empirical curve, the Central American and Mexican cases fall well below the curve; particularly, at higher magnitudes. We offer no explanation for this difference at present, except to note that a difference is not an unreasonable expectation in view of the quite different tectonic regimes involved.

A plot of β -stage intervals vs. magnitude is shown in figure 8. In this case, the data points for the California earthquakes cluster with the Central American and Mexican earthquake data. The trend that is poorly defined in this plot for earthquakes of intermediate magnitude, falls apart completely for large magnitude earthquakes. However, in view of the large scatter throughout the data set, we attach no significance to this feature at present. We expect that the distribution of seismic stations relative to a particular region should have a strong influence on the detection of renewed activity; hence, on the β value. For the Mexican and Central American cases, locations are dependent mainly on the data from North American stations. This limits the minimum magnitude for locateable events to about 4.0.

The linear dimensions of the regions of anomalous seismic activity for the California cases are plotted vs. magnitude in figure 9. An empirical relation between magnitude and the linear dimensions of regions in which velocity changes were recognized (Anderson and Whitcomb,

1975), and additional points compiled by Ohtake (1977) are also shown. Two additional relationships of interest in this connection are shown for comparison: (1) the relationship given by Utsu (1961) between magnitude and length of aftershock zones, and (2) the relationship given by Dambarra (1966) between earthquake magnitude and the diameter of the zone of crustal deformation detected by geodetic measurements related to the earthquake. These two curves fall well below the Anderson and Whitcomb relationship and our data points. The dimensions of the zones of anomalous seismic activity for the Mexican and Central American cases all fall between 200 and 300 km. These comparisons suggest that the zone of anomalous seismicity is considerably larger than that in which other precursory phenomena are detected.

FIGURE CAPTIONS

- Figure 1. Epicentral locations of the four earthquakes considered in this paper.
- Figure 2. Space-time diagram of earthquakes preceding the Parkfield earthquake of 1966. The epicentral distance for each earthquake (ordinate) is the distance between the main shock and the given earthquake. Arrows mark possible onsets of α and β stages of activity. The terminus of the plot corresponds to the time of the main shock. A horizontal dashed line marks the radius of the zone of anomalous seismic activity.
- Figure 3. Space-time diagram of earthquakes preceding the Borrego Mountain earthquake of 1968. Symbols are described in the caption for figure 2.
- Figure 4. Space-time diagram for earthquakes preceding the San Fernando earthquake of 1971. The symbols are as defined in the caption for figure 2. Two interpretations are suggested: one in which quiescence begins in mid-1967, with renewal of activity early in 1969, and one in which quiescence begins in mid-1969 and renewal of activity begins in mid-1970.
- Figure 5. Space-time diagram for earthquakes preceding the Oxnard earthquake of 1973. Symbols are as defined in the caption for figure 2. The plot covers the period from 1971 to the origin time of the main shock.
- Figure 6. Updated plot of the Oaxaca gap. Earthquakes located throughout the region for the period 1970 through mid-1973 are shown in the upper panel. Earthquake epicenters located in this region

between mid-1973 and Day 211 1977 are shown in the lower panel.

Figure 7. Time intervals between the onset of quiescence and the main shock (α plus β) for cases presented in this paper. Also shown is an empirical curve for various precursory phenomena developed by Rikitake (1976). Full circles correspond to the California earthquakes studied; OX = Oxnard, PF = Parkfield, BM = Borrego Mountain, SF = San Fernando. Open circles are data points from previous studies. Data points for the Mexican and Central American shocks are marked by X's. Subscripts indicate the case number given in Ohtake et al. (1977b). Sources of data are indicated as follows: B = Borovik et al. (1971), E = Engdahl and Kisslinger (1977), K = Kelleher and Savino (1975), I = Ishida and Kanamori (1977), M = Mogi (1969), O_1 = Ohtake (1970), O_2 = Ohtake (1976), S = Su-Shaosein (1976).

Figure 8. Interval between the renewal of activity vs. magnitude for the cases presented in this paper. Symbols are as described in the caption for figure 7.

Figure 9. Linear dimensions of the zones of anomalous seismic activity for the four California earthquakes examined in this paper vs. magnitude of the main shock. Empirical relationships between: (A) magnitude and the linear dimension of zones in which P-wave velocity changes were detected prior to an earthquake (Anderson and Whitcomb, 1975); (U) magnitude vs. length of aftershock zone (UTSU, 1961); and (D) magnitude vs. diameter

of regions of crustal deformation detected by geodetic measurements prior to earthquakes (Dambara, 1966). Barred circles are for earthquakes associated with subduction. Open circles correspond to intraplate earthquakes.

Table 1. Pertinent Data for Four California Earthquakes Studied

(M_S = surface wave magnitude, L = length of quiescent zone)

Location	Date	M_S	L, km	Anomalous Period, days		
				α -stage	β -stage	Total
Parkfield	June 28, 1966	5.6	290	274	8	282
Borrego Mt.	April 9, 1968	6.4	280	357	395	752
San Fernando	Feb. 9, 1971	6.5	240	596 (386)	712 (211)	1308 (597)
Oxnard	Feb. 21, 1973	5.2	300	72	25	97

Table 2. Pre- and Post- Gap Earthquake Activity of the Oaxaca Gap

<u>Time Interval</u>	<u>Number of Earthquakes</u>		<u>Total</u>
	<u>Located in the Gap</u>		
	<u>h<60 km</u>	<u>h>60 km</u>	
Mid 1963 - Mid 1973	74	34	108
Mid 1973 - Mid 1977	0	6	6

1. h = focal depth
2. Data derived from the magnetic tape file of the National Earthquake Information Service, NOAA, Boulder, Colorado.

Appendix I

Since the seismic belt is narrow compared to its length, we consider the epicenters as a distribution of points along a line parallel to the trend of the belt. Thus, the distribution of points is defined by a single variable. The question we ask is the following: if we select a sample at random (e.g., distribution of epicenters in a 2-year period), and identify a segment of minimum activity (zero in the present case) of length equal to a given fraction of the total length, what is the probability that this could occur in a random process?

In the present case, the length of the excluded zone is about 1/6 of the total (Figure 2b). Thus, we divide the zone into 6 segments of equal length. We could have chosen any of 6 two-year intervals from the 12-year period investigated. Approximately 500 earthquakes of magnitude 4 and greater, with focal depths less than 60 km, occurred during this period throughout the zone. The probability of a gap of the observed proportion, or larger, occurring as a fluctuation in a random process is given by the binomial function

$$\left(1 - \frac{1}{6 \times 6}\right)^{500} \times (6 \times 6) = 2.74 \times 10^{-5}$$

or 1 chance in 36,385. This calculation ignores uncertainties in epicentral locations or clustering effects (foreshock and aftershock sequences). If we assume that as many as 20% of the located events are part of foreshock and aftershock sequences, the probability is reduced to 1 chance in 2,175.

APPENDIX II

The frequency-magnitude relationship for a belt 720 in length, including the Oaxaca gap, determined for the 10-year interval 1963-1973 is:

$$\text{Log } N = -1.00 m + 6.72 \quad (1)$$

where N is the number of earthquakes of body wave magnitude m and larger that occurred in the zone. The relationship between body wave magnitude, m , and surface wave magnitude, M , determined empirically from the NEIS data file for the same earthquakes used in obtaining equation (1) is:

$$m = 2.74 + 0.480M \quad (2)$$

From (2), $M = 7.5$ corresponds to $m = 6.4$. Inserting this value into (1), yields $N = 2$ earthquakes of $M = 7.5$ every ten years in a belt 720 in length. Thus, the mean return time for earthquake of magnitude $M = 7.5$ for a zone 200 km in length is

$$\frac{2}{10} \times \frac{720}{200} = 18 \text{ years.}$$

REFERENCES

- Anderson, D. and Whitcomb, J. (1975), Time-dependent seismology, J. Geophys. Res. 80, 1497-1503.
- Borovik, N., Misharina, L. and Treskov, A. (1971), On the possibility of strong earthquakes in Pribavakalia in the future, Izv. Acad. Sci. USSR, Phys. Solid Earth Engl. Transl. 13-16.
- Dambara, T. (1966), Vertical movements of the earth's crust in relation to the Matsushiro earthquakes, J. Geod. Soc. Japan, 12, 18-45 (in Japanese).
- Engdahl, E. and Kisslinger, C. (1977), Phenomena precursory to a magnitude 5 earthquake in the Central Aleutian Islands, E.O.S., 58, 433 (abstract).
- Ishida, M. and Kanamori, H. (1977), The spatio-temporal variation of seismicity before the 1971 San Fernando earthquake, California, Geophys. Res. Letters, 4, 345-346.
- Kelleher, J. and Savino, J. (1975), Distribution of seismicity before large strike slip and thrust-type earthquakes, J. Geophys. Res., 80, 260-271.
- Lomnitz, C. (1974), Global Tectonics and Earthquake Risk, Developments in Geotectonics 5, Elsevier Scientific Publishing Co., Amsterdam-London-New York, 192-193.
- Mogi, K. (1963), Some discussions on aftershocks, foreshocks and earthquake swarms --- the fracture of a semi-infinite body caused by an inner stress origin and its relation to the earthquake phenomena, Bull. Earthq. Res. Inst., Univ. Tokyo, 41, 615-658.

- Mogi, K. (1969), Some features of recent seismic activity in and near Japan (2), Activity before and after great earthquakes, Bull. Earthq. Res. Inst., Univ. Tokyo. 47, 395-417.
- Ohtake, M. (1970), Micro-structure of the seismic sequence related to a moderate earthquake, Bull. Earthq. Res. Inst., Univ. of Tokyo, 48, 1053-1067.
- Ohtake, M. (1976), Search for precursors of the 1974 Izu-Hanto-oki earthquake, Japan, Pure Appl. Geophys., 114 (in press).
- Ohtake, M., Matumoto, T., and Latham, G.V. (1977a), Seismicity gap near Oaxaca, Southern Mexico as a probably precursor to a large earthquake, Pure Appl. Geophys., 115, 375-385.
- Ohtake, M., Matumoto, T. and Latham, G. (1977b), Temporal changes in seismicity preceding some shallow earthquakes in Mexico and Central America, Bull. Int. Inst. of Seism. and Earthquake Eng., 15, 105-123.
- Ohtake, M., Matumoto, T., and Latham, G. (1977c), Seismicity gaps preceding some shallow earthquakes in California, Abstracts of Seism. Soc. Japan, Fall Meeting, 1977.
- Rikitake, T. (1976), Earthquake Prediction, Elsevier Scientific Publishing Co., Amsterdam-London-New York, p. 357.
- Scholz, C., Sykes, L. and Aggarwal, Y. (1973), Earthquake prediction: a physical basis, Science, 181, 803-910.
- Su-Shaosein (1976), Characteristics of seismic activity related to the Haicheng earthquake, Proc. Lectures by the Delegation of the People's Republic of China, Bull. Seism. Soc. Japan, 27-41 (in Japanese).
- Utsu, T. (1961), A statistical study on the occurrence of aftershocks, Geophys. Mag. 30, 521-605.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

73181-77211

Figure 6

Figure 7
608

Duration of β -stage, days

Figure 8
609

Figure 9

THE MAKRAN COAST OF IRAN,
A POSSIBLE SEISMIC GAP

W. D. Page, G. Anttonen, and W. U. Savage
Woodward-Clyde Consultants, #3 Embarcadero Center,
San Francisco, California 94111

ABSTRACT

The available record of historical seismicity in the Makran region is limited and indicates scattered seismic activity east of the Iran-Pakistan border. If only the Iranian portion of the Makran coast is considered, the seismic hazard may be incorrectly assessed as low. An analysis of the seismic hazard that includes consideration of regional tectonics in combination with the historical record of seismicity and geologic evidence of Quaternary faulting, suggests that the region may be a seismic gap.

The Makran region can be divided tectonically into two separate zones: the west Makran coastal zone from east of Bandar Abbas to Jask and the Makran Ranges and Coast from Jask, Iran to Karachi, Pakistan. The west Makran zone is characterized by high angle east dipping reverse faults of the Zendan fault system. Numerous of these faults displace Quaternary alluvial fan and stream deposits. Earthquakes in the northern part of the west Makran coastal zone and an evaluation of lengths of these faults suggest that the maximum expected earthquake is 7.5 and such an earthquake could occur as far south as Jask.

The main Makran coast appears to lie shoreward of an active subduction zone. The 27 November 1945 earthquake (M_s 8.3) epicenter, originally located offshore, has been relocated to Pasni, Pakistan, which indicates that this large earthquake is related to the near-coast subduction zone. Geologic field investigations of the effects of the 1945 earthquake documented 1 to 3 meters of uplift of the coast at Ormara, Pakistan. The coast at Ormara has a number of raised marine terraces and beaches, and similar raised terraces are found on the headlands along the entire Makran coast. At Konarak, Iran, for example, at least nine terraces were observed. The two lowest terraces at Konarak occur as small remnants, and radiocarbon dates indicate they are Holocene in age.

These raised marine terraces and other geologic evidence for the presence of a subduction zone indicate that the emergence of the Makran coast is probably associated with major earthquakes and that major earthquakes can be expected in the future along the entire coastline from Karachi to Jask.

The Makran region illustrates the importance of assessing the tectonic framework and regional geology and of reevaluating the locations of historical earthquakes in the region as part of an accurate assessment of seismic hazards.

INTRODUCTION

Modernization and industrialization of Iran is proceeding at a rapid pace and extending into regions of the country that have formerly been inhospitable desert with only few scattered villages. The southern coast of Iran, south and east of Bandar Abbas to the Pakistan border, is one of these regions. As plans for development are made and various structures and facilities built, evaluation of seismic risk for major installations becomes desirable. If only the historic and instrumental seismic data is used for this evaluation, the tendency would be to dismiss the risk because the instrumental seismic record in this region shows only a few small earthquakes, mostly located near Bandar Abbas and along the Pakistan Border.

If the geologic record is examined and related to the probable plate tectonic model for the region, it is apparent that the area is potentially seismic and a location where large earthquakes could occur in the future. This is a seismic gap as used in this paper; that is, an area with little seismicity but where other evidence suggests it should be seismic. The area in question along the coast can be divided geologically and tectonically into two parts, the west Makran coast along the Strait of Hormuz from 50 km east of Bandar Abbas to Jask and the main Makran coast along the Gulf of Oman and the Indian Ocean from Jask, Iran, to Karachi, Pakistan.

This paper discusses the seismic record in the area, presents a tectonic model and examines the geology and Quaternary faulting along the west Makran coast. A comparison is also made between geologic similarities of the Pakistan part of the Makran coast, where the 1945 Pasni, Pakistan, earthquake (M_s 8.3) occurred, with the Iranian portion of the Makran coast.

The field work was done as part of a siting study by Woodward-Clyde Consultants for the Atomic Energy Organization of Iran. Six weeks from May to July 1975 were spent in the field. Air photo analysis and aerial reconnaissance of the Iranian coastline in a fixed-wing aircraft preceded ground studies at Kuhestak, Tujak, Jask, and Konarak. A special investigation was made of the reported uplift associated with the 1945 earthquake at Ormara and Pasni, Pakistan. The authors wish to thank the other geologists who worked on parts of the study for their assistance and critical discussion of the problem, particularly Lloyd Cluff, Don Tocher, Jack Alt, Shakir Zuberi, Tom Freeman, Mike Perkins, and Paul Guptil.

SEISMICITY OF THE MAKRAN REGION

The seismicity of the Makran region (Fig. 1) shows an area of scattered, infrequent earthquakes in the period of record, May 1926 through April, 1977 (The NOAA Catalog). By contrast, in the Zagros Mountains to the northwest, moderate to large earthquakes are common, particularly along the eastern Zagros

front or fold belt. To the east of the Makran area moderate earthquakes are scattered along the Owens fracture zone and inland along or near the Ornach Nal and Chaman fault belts.

The Iranian segment of the Makran coast and Makran ranges has had only one third the number of earthquakes that the Pakistani segment has had. These earthquakes are concentrated in two places: east of Bandar Abbas and near the Pakistan border. East of Bandar Abbas and north of Minab three earthquakes with magnitudes greater than 5 were recorded. Near the Pakistan border and 200 km north of the coast, two earthquakes of magnitude 6.5 were recorded. East of Char Bahar, two earthquakes of magnitudes 5 to 5.5 have occurred and were the largest events along the Iranian segment of the Makran coast. The largest two earthquakes in the region: the November 27, 1945 earthquake (M_s 8.3) and the August 5, 1947 earthquake (M_s 7.3) occurred near Pasni, Pakistan. The small number of earthquakes and their wide variation in reported focal depths (near zero to more than 100 km) does not support any clear correlation with geologic structures in the Makran region. The correlation of instrumentally located earthquakes with specific geologic structures such as anticlines and faults requires location accuracies not obtainable with present limited regional seismographic installations in southern Iran. Nowroozi (1972) does not provide any focal mechanisms for events within the Makran region; but two events somewhat to the north appear to be produced by a stress regime similar to that in the Zagros region, that is, predominant northeast-southwest compression. Jacob and Quittmeyer (1978) show that the depths of the earthquakes increase in land from the coast to a depth of 80 km.

REGIONAL STRUCTURES

Southern coastal Iran can be divided into three structurally distinct regions: The Zagros Mountains, which lie between the Persian Gulf and the desert plains; the west Makran coastal zone which lies along the coast of the Strait of Hormuz from east of Bandar Abbas to Jask; and the Makran ranges which lie between the Gulf of Oman and the Jaz Murian Depression.

The Zagros Mountains trend northwest-southeast and consist of broad anticlines that are locally pierced by salt diapirs. The valleys between the anticlines are narrow synclines and thrust faults. The amount of thrusting increases to the northeast. The structures progress from broad gentle anticlines along the Persian Gulf to a discontinuous series of low-angle thrusts and overturned folds to the more continuous, main Zagros thrust along the mountain front. Rock types in the Zagros include marl, conglomerate, turbidite, and radiolarite-ophiolite, which have been interpreted as a former subduction zone (Dewey and Bird, 1971).

The main Zagros thrust bifurcates at its southeastern end and appears to continue southward and southeastward into the

Makran ranges as two major fault systems, the Zendan fault system along the west Makran coast, and the Jaz Murian system along the southern margin of the Jaz Murian Depression (Fig. 2). The Makran ranges have two distinct structural trends. The west Makran coastal zone is a narrow strip approximately 30 km wide of north-south folds and faults lying between the Strait of Hormuz and the north-south trending Zendan fault system. The main Makran range east of the Zendan fault system has east-west trending faults and folds extending from the Jaz Murian Depression to the Gulf of Oman.

The west Makran coastal zone generally consists of north-south trending folds of Miocene marine deposits of the Mishan and Agha Jari Formations (IOOC, 1969): siltstone, mudstone, and sandstone with local lenses of conglomerates. These are overlain by Pliocene (?) conglomerate.

In contrast to the broad anticlines and narrow synclines typical of Zagros-type folding, folding along the west Makran coast is characterized by broad, dish-shaped synclines separated by narrow, sharply defined anticlines with steeply dipping flanks. The anticlinal folds are from 20 to 70 km in length. High-angle, east-dipping reverse faults associated with overturned beds are common on the western anticlinal limbs. These structures appear to reflect westward-directed tangential compression and major overthrusting toward the west (or underthrusting toward the east) along the north-south trending Zendan fault system.

The Zendan fault system extends from the termination of the main Zagros thrust on the north to the coastal mountains west of Jask, a distance of over 250 km (Fig. 2). It is characterized by east-dipping faults produced by tangential compression and overthrusting toward the west. The fault system is relatively linear from its intersection with the main Zagros thrust to north of Senderk, approximately 50 km southeast of Minab. This segment of the fault system consists of only two or three main traces striking approximately N20°W. Fault traces in the southern part of this system are less linear, which could reflect lower-angle thrusting than along the northern segment. Branch faults east of the main thrust zone appear to change in strike and parallel the east-west structural trends of the main segment of the Makran ranges.

The older rocks in west Makran region are the most strongly deformed, indicating that deformation has been continuous since Pliocene or early Quaternary. Older Quaternary alluvium consisting of stratified sand and lesser conglomerate was deposited across the folded and eroded Tertiary strata with a marked angular unconformity. Renewed deformation in later Quaternary time later activated these structures, faulting and deforming the older Quaternary alluvium.

Folds in the Makran ranges east of the Zendan fault system generally parallel the southeasterly and east-west trends of the Makran ranges. Overturning toward the south associated with north-dipping thrust faults is common along the southern flanks of these structures. Ahmed (1969) describes similar structural relationships in the eastern Makran ranges of Pakistan, and Klitgord (1976) and Farhoudi and Karig (1977) discuss the geometry and evolution of these folds.

Thrust faults south of the Jaz Murian Depression appear to exhibit the same sense of displacement as the Zagros thrust, overthrusting toward the south and southwest.

Faults at the western end of the Jaz Murian Depression and along its southern border exhibit evidence of Quaternary activity. Linear scarps displace alluvial fans along both the eastern and western base of the mountain block at the western end of the Jaz Murian Depression (Fig. 2). The adjacent mountain block to the west also has a fresh scarp along the base of its eastern flank. These north-south-trending faults change in strike to the southeast and join the fault system south of Jaz Murian. The most easterly fault, which was examined for a distance of 150 km, exhibits signs of recent activity along this entire length, even to the southeast, where it distinctly joins the southeast-trending fault system south of the Jaz Murian Depression.

QUATERNARY FAULTING IN THE WEST MAKRAK AREA

Aerial and ground reconnaissance during this study verified late Quaternary activity of the Zendan fault system. Fault scarps in late Quaternary alluvial fans along the Senderk River, approximately 45 km southeast of Minab, range from 3 to 15 m in height and indicate uplift of the inland block to the east (see Fig. 4). A large fault scarp approximately 30 m high in alluvium 20 km east of Minab on the eastern side of the fault system also exhibits uplift to the east (Fig. 5). These displacements are consistent with the postulated sense of overthrusting toward the west. None of the scarps displace the youngest alluvium, and all of them exhibit signs of erosion such as gullying, rounded crests, slopes less than the angle of repose for unconsolidated alluvium, and the development of a patina on surface cobbles. The displaced alluvium appears to be Quaternary in age, and since the scarps are younger than the alluvium, they are also Quaternary in age. Vita-Finzi and Ghorashi (1978) describe 3.5 m of Holocene faulting (7000-1250 yr BP) 40 km north of Minab. This faulting is also compressional in nature, but with the southwestern block overthrust to the northeast.

Evidence of Quaternary fault activity along the system is less pronounced south of Senderk, but alluvial fans crossing faults in this zone, which would provide geologically youthful marker units to demonstrate offset, are few. The presence of west-facing, easily eroded bedrock scarps suggests late Quaternary

faulting involving uplift and overthrusting from the east, as do various other youthful geomorphic features along the southern segment of the main thrust.

Several smaller faults, found west of the main Zendan fault zone in an area southeast of Kuhestak were studied in detail and showed displaced Quaternary alluvium. This area is in the coastal plain and low inland hills. Though relief is only a few hundred meters, the topography is rugged where it is underlain by the tightly folded Miocene strata of the Mishan and Agha Jari Formations (IOOC, 1969).

The major faults in the Kuhestak area are east-dipping reverse faults associated with anticlinal folds. These faults are found in the steeply overturned, western limbs of anticlines. They include the 8-km-long Zeyarat fault, located 8 km southeast of Kuhestak; the 20-km-long Labani fault, located 10 km northeast of Kuhestak; a short west-dipping bedding fault, Fault 5, west of the Zeyarat fault; and a 10 km long fault, which offsets the axis of an anticline about 1 km right lateral, located 16 km east of Kuhestak. The Zendan fault system lies 22 km to the east.

Several of these faults have displaced the older Quaternary alluvium as well as all the bedrock units. The Zeyarat fault, which is a reverse fault, displaces older alluvial sand and gravel (Fig. 6). Fault 5 also displaces older Quaternary alluvium. Labani fault displaces older alluvial sands (Fig. 6).

UPLIFT OF THE MAKHRAN COAST

Pleistocene beaches and marine terraces along the Makhran coast from Jask to Karachi, which indicate a tectonically emerging coast (Snead, 1969, 1970; Little, 1972; and Vita-Finzi, 1975), have been related to earthquakes (Page and others, 1978).

The coast is marked by a series of prominent headlands separated by low areas. The headlands, which are generally flat topped or gently tilted and locally stepped with terraces, are preserved from rapid erosion by cemented, resistant conglomerates or coquina cap rocks 0.5-5 m thick. These landforms are erosional remnants protected by the resistant cap of Pleistocene terrace deposits. Where these deposits or other resistant rocks are not present, the siltstone and claystone bedrock is rapidly weathered and eroded to low lying pediments. Some terraces are 1 to 2 km wide and 10 to 15 km long. Wave-cut cliffs, notches, and related geomorphic features are found locally on the shore side of the wave-cut platform. The wave-cut terraces are locally tilted, and small faults with a few meters displacement cut some terraces. Landward from and between the headlands are accretionary beaches, lagoonal deposits, and tombolos up to 10 m in elevation.

November 27, 1945, Earthquake, M_s 8.3

The November 27, 1945 earthquake near Pasni, Pakistan is a key to the tectonics of the Makran coast and is discussed in some detail as an example of the type of earthquake possible in the seismically quiet Iranian sector of the Makran coast.

Location - Because the published locations of this earthquake were as much as 200 km apart, the earthquake was relocated using all available stations at epicentral distances less than 100° (but excluding International Seismological Summary published readings with residuals larger than 15 seconds). The solution using the teleseismic location method (Bolt, 1960) and Jeffreys-Bullen travel time curves did not appear to be biased by the large number of stations in one quadrant. The magnitude was redetermined from 20-second surface waves recorded by the Wood-Anderson seismographs at Mt. Hamilton, California. The focal depth as determined from pP readings was not clearly defined, and individual estimates varied over more than 100 km. However, the earthquake appeared to be of shallow focus. There were too few first-motion data available to constrain the focal mechanism.

<u>Epicenter</u>	<u>Origin Time</u>	<u>Depth</u>	<u>Magnitude</u>	<u>No. of Stations</u>
25.18°N 63.47°E	2156:50.5 GCT	Restricted to Normal	8.3	70

Intensity - Intensity of the shock was greatest in a 100 km-long stretch of the Pakistani coast may have reached XI on the modified Mercalli scale at Pasni and Ormara. Pasni is built on beach sands overlapping the Parkini Mudstone Formation. Ormara is on sand tombolo about 500 m wide. Grass mat hut suffered little or no damage from the shaking, but many one-story buildings, constructed of unreinforced masonry (sun-dried brick) and dressed stone and mud mortar, collapsed or were seriously damaged during the earthquake. Most deaths, about 50 in Pasni and 16 in Ormara, were caused by collapse of structures.

There was substantial ground failure at Pansia and Ormara. Ground cracks that issued water were numerous at both places. About 5 km to the north of Ormara, ground cracks were 60 or more centimeters wide. Displacement of about 1.6 m occurred along a long east-west crack, north side up. The earthquake also caused many landslides and rock falls along the steep bluffs south of Ormara (Fig. 7). In Pasni, the shoreward part of town was part of a submarine slide which submerged a zone along the shore 95 to 125 m wide.

The intensity at the east end of the Makran coast had decreased to no more than IV or V, and to the west at Jiwani the intensity had decreased to no more than IV.

Three tsunamis followed this earthquake, 5 to 15 minutes apart and 1 1/2 to 2 hours after the earthquake, around 5:00 a.m. The third wave was the highest and swept over the one-story (4-5 m high) structures at Pasni and Ormara.

At the time of the 1945 earthquake, four mud volcanoes rose near shore 8 to 30 m above water, 4 to 7 fathoms deep; one was in Quadar West Bay, two in Ormara West Bay, and one about 3 km southeast of the mouth of the Hingol River (Sondhi, 1947).

Tectonic uplift - No tectonic change in elevation accompanying the earthquake was documented at Pasni, but approximately two meters occurred at Ormara (Page and others, 1978). The Ormara area is characterized by a steep, rocky headland that is connected to the mainland by a low tombolo. The headland is underlain by siltstones, mudstones, sandstones, and conglomerates of the Ormara Formation (Plio-Pleistocene age) and capped by coquina, limestones, and sandstones of the Jiwani Formation (Pleistocene age) (Photographic Survey Corporation, 1958).

Pre-1945 beaches are now 1 to 3 m above highest tide and the tombolo broadened several hundred meters. A large sand flat, which was exposed to the east of the village of Ormara, is now inundated only at high tide, and the 1945 beach and beach berm has been buried by advancing sand dunes (Fig. 7). The formation of new sand spits and bars, particularly on the more sheltered east side of the tombolo, is still continuing in response to uplift associated with the 1945 event. Local fishermen report that the west side of the headland was impassable to foot traffic even at the lowest tides before the earthquake; afterwards, one could walk this part of the headland even at the highest tides. The tidal range for spring tides is 2.3-m and the maximum range is 3.7 m between the highest high and the lowest low (National Ocean Survey, 1973), indicating that the amount of uplift was greater than 3.7 m. This part of the headland at present shows an elevated wave-cut platform 2 to 3 m above the present one and the 1945 beach berm about 2 m above the present beach berm (Fig. 7).

The Ormara headland and vicinity had also been uplifted prior to the 1945 earthquake. Marine terraces are reported on the seaward side of the headland (Photographic Survey Corporation, 1958). A raised alluvial fan was found on the northeast side of the headland that has boulders with slightly weathered oysters in growth position on the fan's surface. These shells are 11 m above present low tide level and are greater than 37,000 radiocarbon years old. On the central part of the tombolo at 8 m elevation near the village of Ormara, sand beds containing unweathered marine shells (not in growth position, however) are exposed at 2 m depth in hand-dug wells. These shells are $2,710 \pm 135$ radiocarbon years old.

Raised Beaches in Iran

Konarak - The highest, most extensive sequence of raised beaches in Iran is present at Konarak (Page and others, 1978), and is summarized here. At least nine terraces are preserved on the south side of Konarak headland which is connected to the mainland by a tombolo, a situation similar to that at Ormara. Shallow water, marine deposits of late Quaternary age form terraces and overlie siltstone bedrock of Tertiary age with an angular unconformity. Some of the terraces are continuous for several kilometers while others are preserved as only scattered remnants. The terraces are marked by wave-cut platforms, wave-cut cliffs and boulder breccias, and swash troughs. With increasing elevation, the terraces also show an increase in dissection and weathering, and shells in the deposits show increasing recrystallization.

The lowest two terraces occur as small remnants along the southeast side of the headland (Fig. 8). The third and most extensive terrace covers most of the southeast part of the headland. It is separated from the fourth terrace by a scarp 10 m high that runs east-west parallel to the headland. The morphology of the scarp, the presence of swash troughs at regular intervals eroded into the rock in front of the scarp, and boulder breccia found at the base of the scarp indicate that it is a raised beach cliff. The deposits of the third terrace consist of about 4 m of limey sands and silts with numerous shells. Uranium series dates on shells range from 138,000 to 156,000 years old, and indicate that this terrace is Sangamon in age (Page and others, 1978). The fourth through ninth terraces are older but have not been dated.

Jask - A single terrace is located on the south side of the village of Jask and is 5 to 6 m above present sea level. The terrace deposits are 2 to 3 m thick and consist of horizontally bedded fossiliferous Quaternary sandstone and siltstone which overlie gray siltstone and sandstone Tertiary bedrock with an angular unconformity. Uranium Series dates on shells from this deposit range from 114,000 to 140,000 and suggest that the Jask terrace is Sangamon in age (Page and others, 1978).

LONG TERM AVERAGE RATES OF UPLIFT

Calculating the amounts of uplift and rates at a particular place require estimates of the relative ("eustatic") position of sea level during the late Quaternary at that place. Calculations of Sangamon high sea level stand (approximately 125,000 years ago) at plus six meters (Broecker and Van Donk, 1970) used in this study are based on the late Pleistocene sea levels recorded from Barbados (Stenen and others, 1973) and New Guinea (Bloom and others, 1974). Clark's (1977) curve of the Holocene rise of sea level along the Makran coast is used for Holocene sea level estimates (see Page and others, 1978; for a more complete discussion).

At Konarak the best Holocene data, shells in situ from the raised lagoonal deposits, give a Holocene rate of uplift the same as the post-Sangamon rate suggesting that the actual tectonic rate is 0.01-0.02 cm/yr, although rates range as high as 0.1 cm/yr. At Ormara, the Holocene rate is even higher: 0.18 cm/yr after uplift associated with the 1945 earthquake or 0.11 cm/yr, if calculated before the earthquake. There is no post-Sangamon rate for comparison, but the headland and associated raised terraces are higher than at Konarak (Photographic Survey Corp., 1958). This could indicate a faster longterm average rate of tectonic uplift for the Ormara area. At Jask the post-Sangamon uplift rate is zero.

The long term average uplift rate appears to increase from Jask on the west to Ormara on the east. This reflects different amounts of uplift at different places on the coast, a fact supported by the tilting and faulting and discontinuous remnants of the pre-Holocene terraces seen all along the coast. In general the terraces increase in number and elevation from one at Jask, to nine at Konarak up to 100 m in elevation and an unknown number at Ormara up to 500 m in elevation.

PLATE TECTONIC MODEL

Various models have been advanced to explain all or part of the tectonic framework of southern Iran (see Stocklin, 1968; Wells, 1969; Takin, 1972; Nowroozi, 1971 and 1972; Dewey et al, 1973; Stonely, 1974; Stocklin, 1974; Haynes and McQuillan, 1974; Jacob and Quittmeyer, 1978). The plate tectonic model (Fig. 2) used in this report is from Anttonen (manuscript, 1978). This model is drawn from Stonely (1974) with modifications based on data collected during this study and the work of White and Klitgord (1976) and Farhoudi and Karig (1977). In the proposed model the Afro-Arabian plate is moving northeastward with respect to the Eurasian plate and underthrusting beneath it. The Zagros thrust zone connects with the north-south-trending Zendan fault system which in turn connects with a subduction zone off the south Makran coast.

In the Zagros area the collision of the two plates has proceeded to where most of the oceanic lithosphere separating the two continental plates has been consumed, resulting in destruction of the old trench, collision of the continental segments, and underthrusting of the Afro-Arabian plate. A thick wedge of epicontinental sediments has been squeezed between the two continents as they approached one another, which has produced extensive shearing (probably involving basement rocks) in the Zagros thrust zone. Farther southwest, more moderate compression has produced the Zagros folded belt.

The northern part of the Zendan fault system apparently connects to the east end of the Zagros thrust and the western end of the subduction zone off the south Makran coast.

Stonely (1974) characterizes the Zendan fault system as a right-lateral strike-slip fault. Field reconnaissance during this study, however, revealed no positive evidence of right slip offset along this structure. Surface features associated with the fault system indicate overthrusting from the east, suggesting a more easterly direction of movement for the Arabian plate than that postulated by Stonely. The Zendan fault system could have a strong component of right-slip offset obscured by thrust displacement. White and Klitgord (1976) also suggest oblique subduction on the basis of increase in amplitude of frontal folds of the Makran ranges from west to east and the en echelon pattern of the folds.

McKenzie and others (1970) suggest a pole of rotation off the western coast of Greece to account for the opening of the Red Sea graben and northeastward movement of the Saudi Arabian Peninsula. A pole of rotation with this location should produce northeast underthrusting along the Zagros trend, oblique underthrusting with a strong component of right-lateral movement along the west Makran coast, and more moderately oblique subduction along the south Makran coast.

In this model, rapid subsidence of the Oman Peninsula would reflect depression of the northern margin of the Afro-Arabian plate as it approaches the Eurasian plate.

The postulated active subduction zone extends from the Zendan fault system to the Owens fracture zone and the Chaman fault. Evidence for this subduction zone consists of 1) active andesitic volcanoes north of the Jaz Murian Depression and 350 km north of the trench in the Gulf of Oman (Jacob and Quittmeyer, 1978); 2) an accretionary sequence of turbidite deposits constituting the Makran ranges which are being squeezed between subducting oceanic lithosphere on the south and the Eurasian continental lithosphere on the north (Farhoudi and Karig, 1977); 3) seismic events that increase in depth inland from the coast (Jacob and Quittmeyer, 1978); and 4) raised beaches between Jask, Iran, and Karachi, Pakistan (see Falcon, 1947; Little, 1972; Vita-Finzi, 1975; and Page and others, 1978), which indicate episodic uplift of the continental margin along the southern Makran coast (see Fig. 3). The active fault system trending along the southern edge of the Jaz Murian Depression is not a plate boundary in the proposed model, but is related to the subduction zone (Anttonen; manuscript, 1978).

POSSIBLE SEISMIC GAPS

In contrast to the low seismicity known from the historic and instrumental record, the geologic evidence and the tectonic setting show that the coastal area of Iran from 50 km east of Bandar Abbas to the Pakistan border is tectonically active. The geologic record can be used to estimate the size of potential earthquakes.

West Makran Coast

Numerous faults, including faults in the Zendan fault system, have displaced Quaternary alluvial units in the west Makran region. These Quaternary units have not been dated [other than locally by reconnaissance work of Vita-Finzi (1975, 1977)] nor has the faulting been studied in enough detail to assess the degree of activity. However, lengths of the faults studied for this report can be used to estimate maximum earthquake magnitudes. Using the curves of Patwardhan and others (1975) the Zendan fault system (200 + km long) would have a maximum earthquake magnitude of 7.5, the Labani fault (20 km long) a maximum magnitude of 6.5, and the Zeyarat fault (8 km long) a maximum magnitude of 6.0. Two earthquakes, magnitudes 5.3 and 5.6, and one with magnitude 6.25 (Fig. 1) have occurred in the northern area of the Zendan fault; these tend to confirm the estimates of magnitude based on fault length.

Makran Coast

If the 1945 Pasni earthquake is typical and assuming no aseismic uplift in this area, then uplift along the Makran coast can occur as discrete increments along different parts of the coast at different times. The lower two terraces at Konarak may have formed this way. Using the measured uplift rate at Ormara of 0.1-0.2 cm/yr and an uplift of 2 m for the Pasni-Ormara earthquake, then the recurrence of such an earthquake along this particular part of the coast is 1000-2000 years. At Konarak, where the uplift rate is slower (0.02-0.1 cm/yr), the recurrence of a Pasni-Ormara type earthquake would be 2,000-10,000 years. Assuming the Pasni earthquake reflects a strain release along the coast from Pasni to east of Ormara, a distance of 125 km, and measuring the length of the coastal subduction zone of 850 km, the recurrence for a similar Magnitude 8+ earthquake somewhere along the Makran coast is approximately 125-250 years. Because the terraces increase toward the east, the east end of the coast appears to more likely to have larger magnitude earthquakes than the west end.

REFERENCES CITED

- Ahmed, S. S., 1969, Tertiary geology of part of South Makran, Baluchistan, West Pakistan: American Association of Petroleum Geologists Bulletin, v. 53, no. 7, p. 1480-1449.
- Bloom, A. L., Broecker, W. S., Chappell, J. M. A., Matthews, R. K., and Mesolella, K. J., 1974, Quaternary sea level fluctuations on a tectonic coast--new $^{230}\text{Th}/^{234}\text{U}$ dates from the Huon Peninsula, New Guinea: Quaternary Research, v. 4, p. 185-205.
- Bolt, B. A., 1960, The revision of earthquake epicenter, focal depths and origin times using a high speed computer: Geophysical Journal of the Royal Astronomical Society, v. 3, p. 433-440.
- Broecker, W. S., and van Donk, J., 1970, Insolation changes, ice volumes and the 0^{18} record in deep-sea cores: Review of Geophysics and Space Physics, v. 8, p. 169-198.
- Clark, J. A., 1977, Global sea level changes since the last glacial maximum and sea level constraints on the ice sheet disintegration history: University of Colorado, Ph.D. thesis, 150 p.
- Dewey, J. F., and Bird, J. M., 1971, Origin and emplacement of the ophiolite suite--Appalachian ophiolites in Newfoundland: Journal of Geophysical Research, v. 76, no. 14, p. 3179-3206.
- Dewey, J. F., Pitman, W. C., III, Ryan, W. B. F., and Bonnin, J., 1973, Plate tectonics and the evolution of the Alpine system: Geological Society of America Bulletin, v. 84, no. 10, p. 3137-3180.
- Falcon, N. L., 1947, Raised beaches and terraces of the Iranian Makran Coast: Geog. Journal, v. 109-110, p. 149-151.
- Farhoudi, G., and Karig, D. E., 1977, The Makran of Iran and Pakistan as an active arc system: Geology, v. 5, p. 664-668.
- Haynes, S. J., and McQuillan, H., 1974, Evolution of the Zagros suture zone, southern Iran: Geological Society of America Bulletin, v. 85, p. 739-744.
- Iranian Oil Operating Companies (IOOC), 1969, Geologic map of southwest Iran: Geological and Exploration Division, Tehran, scale 1:1,000,000.

- Jacob, K. H., and Quittmeyer, R. C., 1978, The Makran region of Pakistan and Iran--trench-arc system with active plate subduction: manuscript in press.
- Little, R. D., 1972, Terraces of the Makran coast of Iran and parts of West Pakistan: University of Southern California, M.A. thesis, 151 p.
- McKenzie, D. P., Davies, D., and Molnar, P., 1970, The plate tectonics of the Red Sea and East Africa: *Nature*, v. 226, p. 243-248.
- National Ocean Survey, 1973, Tide tables, high and low water predictions, 1974--central and western Pacific Ocean and Indian Ocean: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Washington, D.C., 344 p.
- Nowroozi, A. A., 1971, Seismo-tectonics of the Persian plateau--eastern Turkey, Caucasus, and Hindu-Kush regions: *Seismological Society of America Bulletin*, v. 61, p. 317-341.
- _____, 1972, Focal mechanism of earthquakes in Persia, Turkey, West Pakistan, and Afghanistan and plate tectonics of the Middle East: *Seismological Society of America Bulletin*, v. 62, p. 823-850.
- Page, W. D., Alt, J. D., Cluff, L. S., and Plafker, G., 1978 (in press), Evidence for the recurrence of large magnitude earthquakes along the Makran coast of Iran and Pakistan: *Tectonophysics*.
- Patwardhan, A. S., Tocher, D., and Savage, E. D., 1975, Relationships between earthquake magnitude and length of rupture surface based on aftershock zones (abs): *Geological Society of America Abstracts with Programs*, v. 7, no. 3, p. 419.
- Photographic Survey Corporation, 1958, Reconnaissance geology of part of West Pakistan--a Colombo plan cooperative project: Photographic Survey Corporation, Toronto, Canada, Geological Maps 3 to 5.
- Shearman, D. J., 1977, The geological evolution of southern Iran--the report of the Iranian Makran Expedition: *Geographical Journal*, v. 142, p. 393-410.
- Snead, R. E., 1969, Physical geography reconnaissance--West Pakistan coastal zone: University of New Mexico Publications in Geography I, University of New Mexico, Albuquerque, 55 p.

- Snead, R. E., 1970, Physical geography of the Makran coastal plain of Iran: report to Office of Naval Research, Contract N00014 66 C D104, Task order NR388 082, University of New Mexico, Albuquerque, 715 p.
- Sondhi, U. P., 1947, The Makran earthquake, 28th November 1945--the birth of new islands: Geological Survey, Indian Minerals, v. 1, no. 3.
- Steinen, R. P., Harrison, R. S., and Matthews, R. K., 1973, Eustatic low stand of sea level between 125,000 and 105,000 B.P.--evidence from the subsurface of Barbados, West Indies: Geological Society of America Bulletin, v. 84, p. 63-70.
- Stocklin, J., 1968, Structural history and tectonics of Iran--a review: American Association of Petroleum Geologists Bulletin, v. 52, no. 7, p. 1229-1258.
- _____, 1974, Possible ancient continental margins in Iran, in Burk, C. A., and Drake, C. L. (eds.), The geology of continental margins: Springer-Verlag, New York, p. 873-887.
- Stonely, R., 1974, Evolution of the continental margins bounding a former southern Tehys, in Burk, C. A., and Drake, C. L. (eds.), The geology of continental margins: Springer-Verlag, New York, p. 889-903.
- Takin, M., 1972, Iranian geology and continental drift in the Middle East: Nature, v. 235, p. 147-150.
- Vita-Finzi, C., 1975, Quaternary deposits in the Iranian Makran: Geographical Journal, v. 141, p. 415-420.
- Vita-Finzi, C., and Ghorashi, M., 1978, A recent faulting episode in the Iranian Makran: Tectonophysics, v. 44, p. T21-T25.
- Wells, A. J., 1969, The crush zone of the Iranian Zagros mountains and its implications: Geol. Magazine, v. 106, no. 5, p. 385-514.
- White, R. S., and Klitgord, K., 1976, Sediment deformation and plate tectonics in the Gulf of Oman: Earth and Planetary Science Letters, v. 32, p. 199-209.

REGIONAL GEOLOGIC STRUCTURES

Figure 2

Figure 3
 SCHEMATIC CROSS-SECTION
 OF THE SOUTH MAKHRAN COAST

Modified after Farhoudi and Karig, 1977

A. **FAULT SCARP** — Oblique aerial view of west-facing fault scarp in Quaternary alluvium 20 kilometers east of Minab. Indicated displacement is up (U) to the east. Photo taken looking east.

B. **GROUND VIEW** — Looking south at scarp shown in (A) above. Arrow points to face of scarp, which is approximately 25 meters high.

Figure 4

QUATERNARY FAULT EAST OF MINAB

A. FAULT SCARP — In Quaternary alluvium 5 kilometers north-east of Senderk. Oblique aerial view looking southwest. Displacement is up (U) to east.

B. GROUND VIEW — Of fault scarp shown in (A). Note the rounded crest and gully erosion on this west-facing scarp. Height is approximately 10 meters. Viewed looking east.

A. LABANI FAULT — Miocene sediments (Tm) thrust over older alluvium (Qoa) along north-trending Labani fault (arrows). View is to east.

B. ZEYARAT FAULT — Faulted older alluvial sands and gravel (Qoa) at north end of the Zeyarat fault. The fault (dotted line) parallels bedding of steeply overturned Miocene sediments (Tm) seen east (right) of fault. Miocene sediments west (left) of this fault are upright and gently dipping.

Figure 6

QUATERNARY FAULTS NEAR KUHSTAK

- A. The 1945 beach (center and left) on the east side of the tombolo now overlain by sand dunes; flat to right is exposed except during extreme high tides. Photo looking north.

- B. The 1945 wave-cut platform (arrow) covered by landslide debris caused by the 1945 earthquake at the west end of the Omara headland. The present wave-cut platform, which is still being cut, is approximately 2m below the 1945 platform. Photo looking south.

Figure 7

UPLIFT FEATURES AT ORMARA,
ASSOCIATED WITH THE 1945 EARTHQUAKE

A. Raised wave-cut platforms and cliffs for Terraces 1, 2, and 3 (arrows), southeast part of Konarak headland. Photo looking east.

B. Raised wave-cut terraces and cliffs (paired numbered arrows) for Terraces 1, 2, and 3 on southeast end of Konarak headland. Terraces 1 and 2 are Holocene and Terrace 3 is Sangamon in age. Photo looking northwest.

Figure 8
TERRACES, KONARAK HEADLAND

Figure 1. A map of the study area showing the location of the study sites. The map includes the names of the study sites and the names of the study sites.

Figure 2. A map of the study area showing the location of the study sites. The map includes the names of the study sites and the names of the study sites.

A SEMI-MARKOV MODEL FOR CHARACTERIZING
RECURRENCE OF GREAT EARTHQUAKES

By

Ashok S. Patwardhan, Ram B. Kulkarni, and Don Tocher
Woodward-Clyde Consultants, San Francisco, California

Paper Presented at the
EHRP Conference on Methodology for Defining
Seismic Gaps and Soon-to-Break Gaps

Sponsored by

U. S. Geological Survey
Office of Earthquake Studies
345 Middlefield Road
Menlo Park, California 94025

At

Massachusetts Institute of Technology
May 25, 26, and 27, 1978

A SEMI-MARKOV MODEL FOR CHARACTERIZING RECURRENCE OF GREAT EARTHQUAKES

By

Ashok S. Patwardhan, Ram B. Kulkarni, and Don Tocher*

1. INTRODUCTION

The objective of this paper is to describe an analytical model for characterizing the recurrence of great earthquakes (defined as earthquakes of magnitude $M \geq 7.8$) consistent with the general physical processes contributing to their occurrence. Recurrence characterization includes estimates of size and holding times between successive great earthquakes at a given location. Because uncertainties exist in the various factors contributing to the underlying physical processes, the characterization is probabilistic in nature.

During a great earthquake, major tectonic deformation occurs and a substantial amount of seismic energy is released over an extensive area which can cause injury and widespread economic loss. Therefore, the occurrence of great earthquakes is of interest both from a scientific point of view and for the assessment and mitigation of earthquake damage hazards. Non-occurrence of great earthquakes (often referred to as "seismic gap") is also of interest because it represents a non-uniform spatial and temporal distribution of earthquakes and a temporally varying seismic risk function.

Available historical seismicity data suggest that great earthquakes exhibit patterns of non-randomness in location, size, and time of occurrence. From a physical standpoint, the occurrence of great earthquakes can be represented by a continuous, gradual process of strain accumulation interrupted intermittently by episodes of sudden release. Several factors are believed to influence the size of great earthquakes in a given area; for example, accumulated strain, shearing resistance, slip rates, tectonic stress, and displacement over the interface area. An illustration of this process is

* Woodward-Clyde Consultants, 3 Embarcadero Center, Suite 700,
San Francisco, California 94111

provided by Figures 1(a) and 2, which show the extent of rupture surfaces due to great earthquakes in the southern Alaskan-Aleutian area and temporal distribution of earthquakes ($M \geq 5$) along three cross sections, respectively. The clustering in time and variation in location of great earthquakes is evident. The lack of great earthquakes in cross section A-A (Gulf of Alaska region) for approximately 78 years has classified it as a seismic gap (Sykes, 1971). Similar evaluations have been made in regard to other areas (e.g., Kelleher et al, 1974; Mogi, 1968).

Several statistical models have been proposed to represent the process of earthquake occurrence. The most common model is the Poisson model, which assumes spatial and temporal independence of all earthquakes including great earthquakes; i.e., the occurrence of one earthquake does not affect the likelihood of a similar earthquake at the same location in the next unit of time. Other models such as those proposed by Esteva (1976) and Shlien and Toksoz (1970) consider the clustering of earthquakes in time. A few other probabilistic models have been used to represent earthquake sequences as strain energy release mechanisms. Hagiwara (1975) has proposed a Markov model to describe an earthquake mechanism simulated by a belt-conveyor model. A Weibull distribution is assumed by Rikitake (1975) for the ultimate strain of earth's crust to estimate the probabilities of earthquake occurrences. Earthquake magnitudes, however, are not represented in this model. Knopoff and Kagan (1977) have used a stochastic branching process that considers a stationary rate of occurrence of main shocks and a distribution function for the space-time location of foreshocks and aftershocks. These models are useful in the broad context of predicting earthquake sequences over large tectonic regions.

However, these models are not adequate to characterize the location-specific occurrences of great earthquakes. While a Poisson process does provide estimates of the probability of occurrence of great earthquakes of any size or the formation of a seismic gap which may be characteristic of a whole region, the estimates are independent of the size and time elapsed since the last great earthquake, invariant in time, and insensitive to location. The physical model outlined above would suggest, on the contrary, a dependence on at least two initial conditions--the size of and the time elapsed since the

last great earthquake. Since both of these conditions will vary from location to location, the probability of occurrence of a great earthquake or continuation of a seismic gap can be expected to vary from location to location even within the same seismic region.

A need exists, therefore, for establishing an analytical model that is more consistent with the underlying physical processes and that can characterize the recurrence of great earthquakes on a more location-specific basis.

2. FORM OF THE SELECTED MODEL

In this paper, we have utilized a semi-Markov process which can model the spatial and temporal dependencies of great, main-sequence earthquakes. A semi-Markovian representation of earthquake sequences is consistent with the above generalized understanding of earthquake generation consisting of gradual, uniform accumulation and periodic release of significant amounts of strain energy in the earth's crust. Since the buildup of strain energy sufficient to generate another great earthquake would take some time, the occurrence of a great earthquake at the same location is less likely within short periods of time following an earthquake of similar size than within an area which has not experienced a similar earthquake for a long time. As the time elapsed without the occurrence of another great earthquake increases, so does the probability of its occurrence. It is reasonable to assume that both the size and waiting time to the next earthquake is influenced by the amount of strain energy released in the previous earthquake (related to the magnitude of that earthquake) and the length of time over which strain has been accumulating. For instance, assuming a constant strain rate, the strain buildup required to generate a magnitude 8.6 earthquake is likely to take longer than the strain buildup to generate a magnitude 7.8 earthquake. These considerations are well modelled by a semi-Markovian representation of earthquake sequences.

A semi-Markov process has the basic Markovian property of one-step memory (i.e., the probability that the next earthquake is of a given magnitude depends on the magnitude of the previous earthquake). However, an additional feature of a semi-Markov process is that it provides for the distribution of a holding time between successive earthquakes, which depends on the magnitudes

of the previous and the next earthquake. Consideration of the holding time in effect provides a multi-step memory for the semi-Markov process.

The following sections describe the development and application of the semi-Markov model.

3. DEVELOPMENT OF THE MODEL

The theoretical development of a semi-Markov process is discussed in the literature (e.g., Howard, 1971). Summarized below are the principal theoretical considerations for modelling the process of great earthquake occurrences. The primary objective is to establish a procedure to evaluate the likelihoods of different numbers of various magnitude earthquakes during a specific period of interest.

The model is described by two parameters: state, i , and holding time, τ . A state is defined by the magnitude of a great earthquake. The continuous magnitude scale can be divided into appropriate intervals to specify discrete states of the system. Figure 3 is a schematic representation of the semi-Markov process. It shows the present conditions at a given location given by the magnitude of the last great earthquake, M_0 , and the time elapsed since its occurrence, t_0 . In the next unit of time, the system may either experience no great earthquake or make a transition to any of the other discrete states, M_1 , M_2 , or M_3 . The representation of earthquake occurrences by a semi-Markov process implies that the likelihood of the next great earthquake being of a particular magnitude (i.e., transition to state j) depends on the magnitude of the previous great earthquake (present state i). The holding time, τ , represents the time period for which the system holds in a given state i . As discussed by Howard (1971), the successive state occupancies will be governed by the transition probabilities of a Markov process, but the stay in any state (holding time) will be described by an integer-valued random variable that depends on the state presently occupied and on the state to which the next transition will be made.

The Formal Model

Let p_{ij} be the probability that a semi-Markov process which entered state i on its last transition will enter state j on its next transition. The transition

probabilities must satisfy the following properties:

$$p_{ij} \geq 0 \quad i = 1, 2, \dots, N; j = 1, 2, \dots, N \quad (1)$$

$$\text{and} \quad \sum_{j=1}^N p_{ij} = 1 \quad i = 1, 2, \dots, N \quad (2)$$

where N is the total number of states in the system.

Whenever the process enters a state i , the likelihood that it will go to state j at some future time is determined by the transition probability p_{ij} . However, after j has been selected, but before making the transition from state i to state j , the process "holds" for a time τ_{ij} in state i . The holding times τ_{ij} are positive, integer-valued random variables each governed by a probability mass function $h_{ij}(m)$ called the holding time mass function for a transition from state i to state j . Thus,

$$\begin{aligned} P(\tau_{ij} = m) &= h_{ij}(m) & m &= 1, 2, 3, \dots \\ & & i &= 1, 2, \dots, N \\ & & j &= 1, 2, \dots, N \end{aligned} \quad (3)$$

We assume that a system entering a state i at time 0 will not make another transition at time 0; i.e.,

$$h_{ij}(0) = 0 \quad (4)$$

After holding in state i for τ_{ij} , the process makes the transition to state j and then immediately selects a new destination state k using the transition probabilities $p_{j1}, p_{j2}, \dots, p_{jN}$. It next chooses a holding time τ_{jk} in state j according to the probability function $h_{jk}(m)$ and makes its transition at time τ_{jk} after entering state j . The process continues developing its trajectory in this way indefinitely. A possible trajectory of such a process is shown in Figure 3.

The time a semi-Markov process spends in state i given that it enters i at time 0 without knowing the destination state is called the waiting time τ_i in state i . Let $W_i(\cdot)$ be the probability mass function of τ_i ; i.e.,

$$W_1(m) = P(\tau_1 = m) = \sum_{j=1}^N P_{ij}(m) \quad (5)$$

The cumulative distribution function (CDF) and complementary CDF of τ_{ij} are denoted by $^<h_{ij}(\cdot)$ and $^>h_{ij}(\cdot)$, respectively. The same functions of τ_i are denoted by $^<W_i(\cdot)$ and $^>W_i(\cdot)$, respectively.

If the system has already spent some time (say, t_0) in a particular state (say, i), its first transition out of state i will not be governed by the functions p_{ij} and $h_{ij}(\cdot)$. These functions apply for a system which has just entered state i ; they do not apply for a system which is observed in state i at present time. Let the transition and holding time probability functions for the first transition out of state i given that it has stayed in state i for the previous t_0 time periods be p_{ij}^1 and $h_{ij}^1(\cdot)$, respectively. By using Bayes' theorem, we find the following expression for p_{ij}^1 :

$$\begin{aligned} p_{ij}^1 &= P(\text{transition to } j|i, t_0) \\ &= \frac{P(\text{holding time at least } t_0|i, j) P(\text{transition to } j|i)}{\sum_j P(\text{holding time at least } t_0|i, j) P(\text{transition to } j|i)} \\ &= \frac{^>h_{ij}(t_0) p_{ij}}{\sum_j ^>h_{ij}(t_0) p_{ij}} \end{aligned} \quad (6)$$

An expression for $h_{ij}^1(\cdot)$ can be derived as follows:

$$\begin{aligned} h_{ij}^1(m) &= P(\text{holding time in } i \text{ is } m | \text{holding time in } i \\ &\quad \text{is at least } t_0 \text{ and the next transition is to } j) \\ &= \frac{P(\tau_{ij} = m + t_0)}{P(\tau_{ij} > t_0)} = \frac{h_{ij}(m + t_0)}{^>h_{ij}(t_0)} \end{aligned} \quad (7)$$

A superscript 1 is used to distinguish the probability functions for the first transition from those for the subsequent transitions; e.g., $^<h_{ij}^1(\cdot)$ is the CDF of holding time for the first transition.

When t_0 is equal to zero (i.e., the system has just entered state i), p_{ij}^1 and $h_{ij}^1(\cdot)$ become p_{ij} and $h_{ij}(\cdot)$, respectively, as they should.

The transition probabilities, the holding time probability distributions, and the initial conditions (i.e., the most recent state of the system and time elapsed since then) are the basic parameters of a semi-Markov process. These parameters can be assessed from historical seismicity and geological data as well as experience and judgments of experts. The assessment procedure is illustrated in the next section.

The next step in the development of a semi-Markov process of earthquake occurrences is to evaluate the probability distribution of the number of times the process enters different states (different magnitude earthquakes) in a specified time period.

Let $\omega_i(K_1, K_2, \dots, K_N | n)$ be the probability that a system which has just entered state i will make K_1 transitions to state 1, K_2 transitions to state 2, $\dots$, K_N transitions to state N in n time periods. We term this the joint probability distribution of state occupancies. For a system which has been in state i for t_0 time periods, the joint probability distribution of state occupancies is denoted by $\omega_i^1(K_1, K_2, \dots, K_N | n)$. Expressions for $\omega_i(\cdot)$ and $\omega_i^1(\cdot)$ are obtained below.

Joint Probability Distribution of State Occupancies

To evaluate $\omega_i(\cdot)$ and $\omega_i^1(\cdot)$, we note that the two sequences of events which lead to K_1 transitions to state 1, K_2 transitions to state 2, $\dots$, and K_N transitions to state N through time n are:

1. the first transition out of state i to a state j at time m and then K_1 transitions to state 1, $\dots$, $K_j - 1$ transitions to state j , $\dots$, and K_N transitions to state N in the remaining time $n - m$, and
2. when $K_1 = K_2 = \dots = K_N = 0$, the first transition out of state i after time n .

By summing the probabilities of these sequences of events over appropriate destination states j and time periods m , we obtain:

$$\begin{aligned} \omega_i(K_1, K_2, \dots, K_N | n) \\ = \sum_{j=1}^N \sum_{m=1}^n p_{ij} h_{ij}(m) \omega_j(K_1, K_2, \dots, K_{j-1}, \dots, K_N | n-m) \\ + \delta(K_1, K_2, \dots, K_N) >_{W_i}(n) \end{aligned} \quad (8)$$

$$\begin{aligned} {}^1_i(K_1, K_2, \dots, K_N | n) \\ = \sum_{j=1}^N \sum_{m=1}^n p_{ij}^1(m) h_{ij}^1(m) \omega_j(K_1, K_2, \dots, K_{j-1}, \dots, K_N | n-m) \\ + (K_1, K_2, \dots, K_N) >_{W_i^1}(n) \end{aligned} \quad (9)$$

$$\begin{aligned} i &= 1, 2, \dots, N \\ j &= 1, 2, \dots, N \\ K_1 &= K_2 = \dots = K_N = 0, 1, 2, \dots, n \\ n &= 0, 1, 2, \dots \end{aligned}$$

where: $\delta(K_1, K_2, \dots, K_N) = 1$ if $K_1 = K_2 = \dots = K_N = 0$
 $= 0$ otherwise.

Equations 8 and 9 provide convenient recursive relationships for computing joint probabilities of state occupancies.

4. APPLICATION OF THE MODEL

The model described above can be applied to any area subject to great earthquakes. In this paper, the model was applied to selected areas in the circum-Pacific belt (see Figure 4) in which the primary process of occurrence of great earthquakes is one of subduction and which have a fairly complete record of great earthquakes dating back approximately 80 years (in some areas such as Japan, it is longer). The areas are extensive in length, within which differences exist in the characteristics of relative plate motions--rates and direction, size of interface areas, stress conditions, size of rupture surfaces in individual events, etc.--that probably influence the transition

probabilities and holding times at different locations. To account for these differences, the areas were subdivided into a number of subareas designated as A through G in Figure 4. Each subarea was next subdivided into several smaller zones in which a sequence of historical great earthquakes could be identified.

Great earthquakes are generally associated with rupture surfaces extending over areas in the order of one thousand to tens of thousands of square kilometers (Tocher, 1958; Benioff, 1951; Mogi, 1968; Kelleher, 1974; Patwardhan et al, 1975). The size (length) of these rupture surfaces depends on earthquake magnitude and, for a given magnitude, show a scatter. It has been observed that in many cases the rupture surfaces associated with historical great earthquakes in the same general area show little overlap, and the rupture surfaces of subsequent great earthquakes tend to fill the "gaps" or available spaces between previous great earthquakes. In some other cases, the boundaries of the rupture surfaces coincide with transverse geologic structures. These observations provide a reasonable basis for delineation of zones or spaces for the occurrence of the next set of great earthquakes within a given area. It is not necessary to assume that these spaces are "permanent" and earthquakes of only up to the historical maximum size could recur in the future. It is possible that the rupture surface associated with the next great earthquake will include the rupture surfaces of more than one previous great earthquake. The probability of such an occurrence is not excluded in the model.

The rupture surfaces of the largest magnitude among the recent great earthquakes were utilized to delineate each of the areas shown in Figure 4 into zones for which calculation of transition probabilities and holding times is desired for a limited period of interest (5 to 80 years). The available history of rupture in each zone is reviewed and used as part of the input data.

The delineation of zones was based on the locations and magnitudes of earthquakes given by USGS (1977). Where available, rupture lengths associated with individual great earthquakes were identified based on aftershock zones. For example, in the Alaska-Aleutian area, five zones were identified as shown

in Figure 1(a). Only independent great earthquakes were included, and clearly identifiable aftershocks were excluded. Where the rupture surface dimensions could not be determined, average dimensions based on magnitude-length relationships based on aftershock zones (Patwardhan et al, 1975) were utilized to estimate the size of a zone. Figure 1(b) shows an example of the procedure used for establishing zone boundaries and holding times.

The data from the USGS Hypocenter Data File gives locations and magnitudes of earthquakes. The magnitudes are given in different scales such as Richter magnitudes, surface wave magnitude, and body wave magnitude. It is sometimes difficult to assess the correctness of the magnitude and location of an individual earthquake. Both parameters can influence the delineation of zones and calculation of holding times. Also, in case of some of the older great earthquakes, the threshold of detection of small magnitudes may not have been low enough to permit a delineation of the aftershock zones. Also, some great earthquakes do not appear to have had identifiable aftershock zones. In such cases, it is difficult to establish the extent of the rupture surface associated with that earthquake.

Since the primary objective of this paper is to demonstrate the feasibility of applying the semi-Markov model to the recurrence of great earthquakes and not to establish specific parameters for a given area, possible inaccuracies in the delineation of zones and calculation of holding times are not significant. When recurrence parameters are to be established for a given area, appropriate data should be evaluated carefully before the model is applied. The evaluation should be supplemented by a parametric analysis to assess the significance of uncertainties in the data.

Considerable scatter exists in the estimated lengths of aftershock zones of earthquakes of a given magnitude (by a factor of three to seven; e.g., Patwardhan et al, 1975; Kelleher and McCann, 1976). These differences may stem from a variety of reasons including the differences in the source characteristics of earthquakes. Some authors also suggest that these differences are regional in nature (e.g., Kelleher and McCann, 1976) and an upper limit may exist for each area. Such differences influence the delineation of boundaries of zones and the assessment of holding times. The significance of

the variable magnitude-rupture length relationships to the holding times and results of the semi-Markov model are discussed in greater detail in Section 7.

The procedure of calculating probabilities of different magnitude earthquakes in a given zone within a specific period of interest Y using a semi-Markov process consists of the following steps:

1. Define states and unit time for the semi-Markov process.
2. Define the initial seismicity condition of the zone in terms of the magnitude of the last great earthquake (greater than 7.8) in the zone and the time elapsed since then.
3. Assess the model parameters consisting of the transition probabilities p_{ij} and probability distribution of holding times $h_{ij}(\cdot)$ on the basis of available historical seismicity and geologic data and subjective assessments.
4. Use equations 8 and 9 recursively to calculate the probabilities of K earthquakes ($K = 0, 1, 2, \dots$) of different magnitudes ($M \geq 7.8$) during the time period Y .

The following paragraphs describe the procedure in each of the above steps.

Step 1 - Defining States and Unit Time

The following discrete states were defined for the system of occurrence of great earthquakes ($M \geq 7.8$):

<u>State Index</u>	<u>Magnitude</u>
1	8 ± 0.2
2	8.4 ± 0.2
3	8.75 ± 0.15

A unit time for a semi-Markov process should be small enough so that the probability of two or more transitions (great earthquakes) is negligible and

large enough so that only a limited number of transitions need to be studied during the selected period of interest. On the basis of these considerations, a unit time of five years was selected. Historical seismicity data suggest that this unit time is reasonable although shorter unit times have been observed in some cases (e.g., in zone 1 of Figure 1).

Step 2 - Defining Initial Conditions of the System

Two initial conditions need to be defined for each zone: the size of the last great earthquake (M_0) and the time elapsed since its occurrence (t_0). Both conditions can be established relatively easily for zones which have had at least one great earthquake in historical times. In such cases, known great earthquakes are arranged in a chronological sequence, and the last earthquake in the sequence is identified as M_0 (e.g., $M_0 = 8.3$ for zone 1 and 8.2 for zone 4 in Table 1). Similarly, the time elapsed since the last great earthquake (t_0) is estimated from the sequence (e.g., for zone 1 in Figure 1, $t_0 = 78$ years, and for zone 4, $t_0 = 21$ years). A compilation of magnitudes of the last great earthquakes (M_0) and holding times (t_0) for the Alaska-Aleutian area (derived from Table 1) is shown in Tables 3 and 2, respectively. Similar values of M_0 and t_0 for all areas (excluding Area B of Figure 4) are shown in Tables 4 and 5, respectively. Area B exhibits preponderant recurrence of a specific state ($M = 8 \pm 0.2$). Therefore, sample data for M_0 and t_0 for Area B were tabulated separately in Tables 6 and 7. If a zone has not had any great earthquake during historical times, it is not included in the present analysis. Zones that have had only one great earthquake in historical times are given in the last column of Tables 2 and 4 and, in general, include some of the longest seismic gaps.

Table 1. GREAT EARTHQUAKES IN THE ALASKA-ALEUTIAN AREA (1899-Date)

Zone 1		Zone 2		Zone 3		Zone 4		Zone 5	
Date	M	Date	M	Date	M	Date	M	Date	M
9-4-1899	8.3	3-28-1964	8.5	11-10-1938	8.7	1-1-1902	7.8	9-2-1907	7.8
9-10-1899	8.6					2-5-1905	7.9	2-4-1965	7.9
10-9-1900	8.3					8-17-1906	8.3		
						3-7-1929	8.6		
						3-9-1957	8.2		

Table 2. ALASKA-ALEUTIAN AREA, SUMMARY OF
HOLDING TIMES BETWEEN GREAT EARTHQUAKES

<u>Magnitude of Last Earth- quake (M_i)</u>	<u>Magnitude of Following Earthquakes (M_j)</u>	<u>Holding Time (Years)</u>	<u>Number of Years as Gap Following M_i</u>
8 ± 0.2	8 ± 0.2	3, 59	
	8.4 ± 0.2	1	21, 13
	8.75 ± 0.15	--	
8.4 ± 0.2	8 ± 0.2	28	
	8.4 ± 0.2	1, 1, 23	78, 13
	8.75 ± 0.15	--	
8.75 ± 0.15	8 ± 0.2	--	
	8.4 ± 0.2	--	
	8.75 ± 0.15	--	40

Table 3. ALASKA-ALEUTIAN AREA, SUMMARY OF
TRANSITIONS OF MAGNITUDES BETWEEN GREAT EARTHQUAKES

<u>Magnitude of Last Earthquake</u>	<u>Magnitude of Following Earthquake</u>
8 ± 0.2	8.3, 7.9, 8.3, 7.9
8.4 ± 0.2	8.6, 8.3, 8.6, 8.2
8.75 ± 0.15	--

Step 3 - Assessment of Model Parameters

Since the upper range of holding times between great earthquakes may be up to a couple of hundred years, the historical seismicity data alone of about 80 years are not sufficient to provide reliable estimates of the parameters of a semi-Markov process; namely, transition and holding time probability distributions. Therefore, a Bayesian procedure which utilizes both historical seismicity data as well as subjective inputs based on judgment in a formal statistical format was used. A schematic representation of the Bayesian procedure is shown in Figure 5. The main steps involved are:

- calculation of sample likelihood for historical seismicity data,
- assessment of prior distribution based on subjective inputs, and
- calculation of posterior distribution by combining information from both sources.

(a) Calculation of Sample Likelihood

The sample likelihood is obtained from the historical seismicity data assuming an appropriate probability distribution. For the present model, two types of variables are involved: (1) the holding time, τ_{ij} , between earthquakes of magnitudes corresponding to states i and j ; and (2) the magnitude of the earthquake, M_i , following the earthquake of a magnitude corresponding to state i . Since three discrete states of earthquake magnitudes have been defined, a total of nine τ_{ij} variables and three M_i variables are required in the analysis. A lognormal probability distribution is assumed for τ_{ij} and $(M_i - 7.8)$. (Note that 7.8 is the smallest earthquake magnitude considered in the model.) The lognormal distribution has a range of 0 to ∞ and is skewed to the right. Both these properties are reasonable for the variables under consideration.

If a random variable X is lognormally distributed, it follows that $Y = \ln X$ is normally distributed (Benjamin and Cornell, 1970). The sample likelihood for data points drawn from a normal process can be specified in terms of the following parameters:

$$\text{sample mean:} \quad \bar{Y} = \frac{1}{n} \sum Y_i \quad (10)$$

$$\text{sample variance:} \quad S^2 = \frac{1}{n-1} \sum (Y_i - \bar{Y})^2 \quad (11)$$

$$\text{sample size:} \quad n = \text{number of data points} \quad (12)$$

$$\text{degrees of freedom:} \quad \nu = n - 1 \quad (13)$$

The data on τ_{ij} and M_i were obtained from earthquake sequences tabulated from NOAA catalogs. To illustrate the procedure, consider the Alaska-Aleutian region. Based on the rupture lengths of the largest events, this area was divided into five zones (Figure 1). In each of the zones, sequences of great earthquakes were listed with respect to their magnitude and year of occurrence as shown in Table 1.

From these earthquake sequences, the holding times and transition earthquake magnitudes were estimated as summarized in Tables 2 and 3, respectively.

A similar procedure was used for the remaining areas shown in Figure 4. The holding times and transition magnitudes obtained from analysis of the great earthquakes in these areas are summarized as sample data in Tables 4 and 5, respectively. All holding times were calculated to present (1978).

(b) Assessment of Prior Distribution

For combining the data with the subjective inputs in an analytically convenient manner, a conjugate prior distribution is often assumed (Raiffa and Schlaifer, 1960). If the data are assumed normally distributed, the appropriate conjugate prior distribution for the model parameters is the student-t. This distribution can be specified in terms of four parameters: $\bar{Y}'$, S'^2 , n' , and ν' (the prime indicates a prior parameter). In order to calculate the prior parameters, the probability distribution function of the corresponding variable (holding time, τ_{ij} , or earthquake magnitude, M_i) was assessed using the fractile method discussed by Raiffa (1968).

The subjective assessments used as prior fractiles were developed based on the trends indicated by the general physical model of earthquake generation discussed in Section 1. For all initial states, the likelihood of the following state being $M_j = 8 \pm 0.2$ was assumed to be higher than the other two states, $M_j = 8.4 \pm 0.2$ or 8.75 ± 0.15 . The transition probabilities for different states decrease with increasing magnitude. The probability of transition to state 1 ($M = 8 \pm 0.2$) is relatively insensitive to the initial state, but the distribution of transition probabilities is considered to be somewhat sensitive to the initial state (see Table 5). The holding times were considered to increase with increasing magnitude but were also considered to be relatively less sensitive to the initial state (see Table 4). The distributions of holding times for all areas and Area B were assumed to be the same in all cases (see Table 6). However, in the case of Area B, the transition probabilities were considered to be high for state 1 and significantly smaller for other states (see Table 7).

The prior parameters are related to the moments of the prior distribution as follows:

$$\nu' = \frac{6 \times (\text{variance})}{\text{fourth moment} - 3 (\text{variance})} + 4 \quad (14)$$

$$\bar{Y}' = \bar{Y} \quad (15)$$

$$S'^2 = \frac{\text{variance } (v'^2 - 4)}{(v'^2 + 1) v'} \quad (16)$$

$$n' = v' + 1 \quad (17)$$

(c) Calculation of Posterior Distribution

If the observations on a variable Y are drawn from a normal distribution and a conjugate prior distribution is assessed for the model parameters, it follows that the parameters of the posterior distribution can be obtained from the following equations (Raiffa and Schlaifer, 1960):

$$n'' = n + n' \quad (18)$$

$$v'' = v + v' + 1 \quad (19)$$

$$\bar{Y}'' = (n' \bar{Y}' + n \bar{Y}) / (n + n') \quad (20)$$

$$S''^2 = [(v' S'^2 + n' \bar{Y}'^2) + (v S^2 + n \bar{Y}^2) - n'' \bar{Y}''^2] / v'' \quad (21)$$

Fractile values for posterior distributions of the holding times and state transitions are given in Tables 4 and 5, respectively. Tables 6 and 7 give the prior and posterior fractile values for holding times and transitions for Area B, respectively. Figure 6 depicts an example of the prior and posterior distributions for holding times in transition from state 1 ($M = 8 \pm 0.2$) to state 2 ($M = 8.4 \pm 0.2$) and state 3 ($M = 8.75 \pm 0.15$), respectively. Figure 7 shows an example of the prior and posterior distributions for transition from state 1 to any of the three states. For purposes of analysis, the continuous probability functions on τ_{ij} and M_i were discretized to obtain the model parameters.

5. CALCULATION OF RESULTS

The primary result obtained from the model is the set of probabilities of occurrences of different magnitude earthquakes in a given zone during a specified period of interest. Equations 8 and 9 are used recursively to

calculate the probability of K_1 earthquakes of magnitude $8(\pm 0.2)$, K_2 earthquakes of magnitude $8.4(\pm 0.2)$, and K_3 earthquakes of magnitude $8.75(\pm 0.15)$. The probabilities are location and time specific; i.e., they are dependent on the initial conditions of the zone and are applicable for the duration of real time. For example, if a period of interest of 40 years is specified, the probabilities apply to the next 40 years, rather than to any 40 years.

Table 8 shows an example of the results for a zone with initial conditions of $M_0 = 8$ and $t_0 = 5$ years. The probabilities of occurrences of different combinations of earthquake magnitudes during the next 40 years are shown. For example, the probability that no great earthquake ($M_s \geq 7.8$) will occur in the next 40 years is 0.19, while the probability of occurrence of one $8.4(\pm 0.2)$ magnitude earthquake is 0.056. The holding time and transition probabilities used in obtaining these results are based on the assumption of a constant magnitude-rupture length relationship for all the areas included in the analysis (data shown in Tables 4 through 7).

The basic output of the semi-Markov model (i.e., the probabilities such as those shown in Table 8) provide one of the inputs necessary for seismic risk calculation at a given location.

6. UTILIZATION OF RESULTS

The set of probabilities of different magnitudes of great earthquakes within a given time period yielded by this model can be readily applied for a number of purposes. The model is particularly helpful in characterization of a $N(M)$ relationship, in the magnitude range where the Poisson model has difficulty due to lack of sufficient data; i.e., near the tail of a distribution. It can also be utilized to assess the probability of lack of earthquakes; i.e., the probability of continuation of a "seismic gap". Both determinations can be location-specific and on a "real-time" basis. Discussed below is the methodology for evaluation of "seismic gaps" and characterization of tails of magnitude distributions.

Characterization of Seismic Gaps

There is no well-accepted definition of a "seismic gap". The term "gap" has

been applied both in a spatial and/or temporal sense. When used only in a spatial sense, it refers to a zone in which no great earthquake has occurred for sometime since the last great earthquake (usually for more than 20 to 30 years), while the adjacent zones have experienced great earthquakes within the same time period. When used in a spatial-temporal sense, it refers to the non-occurrence of a great earthquake in a given zone or part of it for a period of time since the last great earthquake. In this paper, the term is used in the latter sense (i.e., spatial-temporal sense). The minimum time period for the identification of a gap is taken as one unit time; i.e., 5 years. The model can be utilized to estimate the probabilities that there will be no earthquake within a period of interest (say, 40 years) for different initial conditions. Figure 8(a) shows an example in which the relationship between the probability of continuation of a gap for the next 40 years is plotted against the number of years for which there have been no earthquakes following earthquakes of $M = 8 \pm 0.2$ and $M = 8.75 \pm 0.15$. Figure 8(b) shows similar relationships for Area B. The relationships in Figure 8(a) indicate that the probability that a gap will be formed following a magnitude 8 ± 0.2 earthquake is approximately 20 percent in the first 5 years. If no great earthquake occurs for 20 years, the probability of continuation of the gap for another 40 years decreases to approximately 10 percent; this result differs from a Poisson model, which indicates a constant probability of formation or continuation of a gap. In a different seismic environment such as Area B, the corresponding probabilities for $M = 8 \pm 0.2$ for the continuation of a gap for periods of 5 and 20 years are 10 and 4 percent, respectively; i.e., considerably smaller than the probabilities for the continuation of gaps in other areas. In other words, gaps in Area B have a greater likelihood to be filled than the gaps in other areas. The relationships in Figure 8(a) also indicate that the probability of continuation of a gap after the lapse of any given time period increases with the magnitude of the last earthquake; this result is to be anticipated. The trends for higher magnitudes show some interesting differences especially when the elapsed time is greater than approximately 50 years. As seen from the posterior probability distribution for $\tau_{ij} > 40$ years, the probability of having a magnitude 8 ± 0.2 or 8 ± 0.4 earthquakes is very small, and the probability of continuation of a gap is primarily the probability of having no earthquakes of magnitude 8.75 ± 0.15 . This result is discussed further in the next paragraph.

The probabilities of continuation of a gap for different initial conditions (M_0 , t_0) and different time periods of interest (t) are shown in Tables 9 and 10 for all areas except Area B and in Table 11 for Area B. Table 9 gives values of probabilities ($M_0 = 8 \pm 0.2$) estimated from the model and corresponding relative frequency values estimated by directly using the sample data and data on the events with no earthquakes for a number of years following M_1 shown in Table 4. As seen from the table, the values calculated from the model show very good agreement with the data. For example, to estimate the probabilities from the data in Table 4 for $M_0 = 8$, $t_0 = 5$, and $t = 5$, the sample data and the data on number of years for different events with no earthquake following $M_1 = 8 \pm 0.2$ were examined to count the number of events with at least one earthquake in 5 years (5 events) and the number of events with no earthquakes between 5 and 10 years (22 events), which yield a relative frequency of 22/27 or 0.81 for the continuation of a gap. This value compares favorably with the probability of 0.74 estimated from the model. Similar estimates for other initial conditions show good agreement with the model except where the data are too scanty (e.g., $M_0 = 8 \pm 0.2$, $t_0 = 30$ years, $t = 20$ years).

Table 10 shows a similar comparison for $M_0 = 8.4 \pm 0.2$. As in the previous case, the agreement between estimates of probability of continuation of gaps from the data and the probabilities estimated from the model is reasonable. This agreement between the probabilities of formation of gaps also suggests that the prior distributions based on subjective assessment are reasonable. The comparisons discussed above are also shown in Figure 9.

Table 11 shows similar comparisons for Area B. Good agreement is observed between the relative frequencies estimated from the data and the probabilities calculated from the model.

Characterization of Tails of Earthquake Magnitude Distribution

A Poisson model of earthquake occurrences based on a typical $N(M)$ curve (i.e., a and b values) generally does not give reasonable estimates of probabilities of great earthquakes (e.g., $M_s \geq 7.8$). We believe that the semi-Markov model developed in this study provides a better characterization of the tails of the probability distribution of earthquake magnitudes since the model takes into

account the interaction between the length of holding time and the magnitude of the next great earthquake, and the influence of recent energy releases on the magnitude and time of the next great earthquake.

Figure 10 shows the probability of at least one earthquake of magnitude M or greater in the next 40 years for different initial conditions. For comparison, the probabilities calculated from a Poisson model using an average number of 2.44 earthquakes ≥ 7.8 in 40 years have a b -value of 1.0. These values approximately represent the level of seismic activity for great earthquakes in zone 4 of the Alaska-Aleutian area. It is seen that the results of a Poisson model are close to those of the semi-Markov model with initial conditions $M_0 = 8$ and $t_0 = 5$ years; i.e., the case in which a magnitude 8 ± 0.2 earthquake occurred 5 years ago. For areas where a great earthquake has not occurred for a relatively long period of time (e.g., $M_0 = 8$, $t_0 = 80$ years), the probability of occurrence of another great earthquake in, say, 40 years is significantly higher than that estimated by a Poisson model.

To examine further the differences in the results obtained by using a Poisson and a semi-Markov model, the probability distribution of number of earthquakes greater than or equal to $M = 7.8$ is plotted for both models in Figure 11. For the Poisson model, the historical seismicity data for the Alaska-Aleutian area is used with a b -value of 1.0. The data show five great earthquakes ($M \geq 7.8$) in 82 years in this area, which gives an average number of great earthquakes per year of $5/82$.

The Poisson model seems to give higher probabilities for the occurrences of multiple earthquakes ($n \geq 3$) in 40 years than the semi-Markov model for the initial conditions of $M_0 = 8$ and $t_0 = 5$ years. For the latter model, the probabilities of occurrences of a given number of earthquakes are influenced by the initial conditions (recent releases of strain energy); these probabilities provide inputs to a seismic risk model that are more consistent with the postulated earthquake mechanism of strain accumulation and release. For example, the results would indicate that to achieve the same level of risk, a facility in a zone with a seismic "gap" of, say, 50 years after the occurrence of a magnitude 8 earthquake should be designed for a higher seismic

loading than a facility in a zone where a magnitude 8 earthquake occurred, say, 5 years ago; this inference appears to be reasonable.

7. PARAMETRIC ANALYSES

To assess the effect of variation in input parameters on the results provided by the model, two analyses were made. In one, the holding times in the prior distributions were increased by factors of 1.5 and 6. The probabilities of continuation of a seismic gap for the next 40 years for both cases of increased holding time for $M_0 = 8 \pm 0.2$ and 8.75 ± 0.15 and different time periods (t_0) after the previous great earthquakes is shown in Figure 12. When the holding times are increased by a factor 1.5, the probability of continuation of a seismic gap following a magnitude $M_0 = 8 \pm 0.2$ earthquake increases by a factor of 2 to 3; while for an increase by a factor 6, the corresponding probability increases by a factor 10. In the case of higher magnitudes ($M_0 = 8.75 \pm 0.15$), a different trend is observed. If the holding time is increased by a factor 1.5, the probability of continuation of a gap for 40 years increases by a factor 7. If the holding time is increased by a factor 6, the probability of a gap increases by a factor 14. This is so because, as the average holding times increase, the probability of having earthquakes of magnitudes 8 ± 0.2 or greater are not insignificant even after 40 years and the probability of continuation of a gap is not small. The historical seismicity data indicate a similar trend.

The second parametric analysis of the zones were defined based on variable rupture lengths. Considerable uncertainty exists in the rupture lengths appropriate for different areas. Kelleher and McCann (1976) give estimates of maximum rupture lengths but do not suggest magnitude-length relationships for different areas. Assuming that these lengths are associated with the largest magnitude from historical data in a given area, the rupture lengths of lesser magnitudes were selected by judgment. Thus, zones in the Alaska-Aleutian area are based on a rupture length of $800 \pm$ km for $M_0 = 8 \pm 0.2$, while those in the Honshu area are based on rupture lengths of 150 to 200 km. Table 12 shows the sample data, prior and posterior estimates of holding times, and lengths of gaps in the various areas. Table 13 shows the sample data and prior and posterior estimates of transition states. A comparison of Table 12 with Table 4 and Table 13 with Table 5 is instructive. Use of variable rupture lengths

decreases the size of sample data, increases the number of gaps, and suggests generally longer holding times. The probability of continuation of seismic gaps for various initial conditions are expected to be higher, as the case in Figure 13 illustrates.

The parametric analyses provide a useful insight into the effect of rupture sizes and holding times on the formation and continuation of gaps. Thus, in areas where the rupture lengths are higher or holding times are longer, the probability of formation and continuation of a gap is higher. The results illustrated in Figures 8 and 13 provide an approximate quantitative assessment of the degree of variation. These results can be applied for differentiating between the characteristics of gaps in different areas; e.g., between Alaska-Aleutians area and Japan or between Central America and New Guinea.

8. SUMMARY AND CONCLUSIONS

A semi-Markov model is developed to estimate the likelihoods of occurrences of great earthquakes (magnitude ≥ 7.8) at a given location during a specified period of interest. The model takes into account the influence of the length of time over which strain energy is accumulating since the most recent great earthquake in a zone on the magnitude and time of the next great earthquake in the zone.

The basic parameters of the model are (1) probability distribution of holding times between earthquakes of magnitudes M_1 and M_j , (2) transition probabilities (i.e., the probabilities that the next earthquake will be of specified magnitudes following an earthquake of magnitude M_1), and (3) initial seismicity conditions of a zone (i.e., magnitude M_0 of the most recent great earthquake and time t_0 since that earthquake). These parameters were obtained by combining historical seismicity data and expert judgments through the use of a Bayesian procedure. This procedure provided better reliability in the estimation of the parameters than using only the limited historical seismicity data.

The values of probabilities of different magnitudes and holding times is influenced in part by the accuracy and completeness of the historical seismicity record with respect to location and magnitude. Careful

reevaluation of the data should be made before applying the model to a specific area.

The model is based on a qualitative assessment of strain accumulation and intermittent release. The possibility of making quantitative assessments in terms of seismic moments should be explored.

The application of the model was discussed for high seismicity areas in the circum-Pacific belt in which the primary process of earthquake generation is that of subduction. The basic output of the model is the set of probabilities of occurrence of a different number of various magnitude earthquakes (≥ 7.8) in a given zone during selected periods of interest. This output can be used for a variety of purposes in seismicity evaluation problems:

(1) Characterization of a seismic gap. The likelihood that a given zone will form a seismic gap (i.e., will not experience a great earthquake) during a specified period of time can be determined. Furthermore, for a series of adjacent zones, one can estimate the most likely zone which will be "filled" by a great earthquake. This will be useful in identifying the likely spatial pattern for the occurrences of great earthquakes.

(2) Definition of real time inputs to the seismic risk model.

(3) Characterization of tails of earthquake magnitude distributions.

The semi-Markov model provides several advantages over other models in that it is location specific, can take into account initial conditions, and is flexible enough so that its parameters can be adjusted to represent any regimen of great earthquake occurrences such as predominance of certain magnitudes and variations in holding times. The probabilities of earthquake occurrence (and gaps) estimated from the model show reasonably good agreement with the values obtained from available data.

9. ACKNOWLEDGEMENTS

This work is part of an ongoing study supported by the Professional Development Program of Woodward-Clyde Consultants. The authors gratefully

acknowledge this support from WCC, especially Dr. I. M. Idriss, Mr. Douglas C. Moorhouse, and Dr. Keshavan Nair. Dr. Chhote Saraf assisted in the computer analyses, and Mr. Robert L. Nowack assisted in the compilation of earthquake data. Dr. William U. Savage reviewed the manuscript and made several useful suggestions. The help by these colleagues is acknowledged.

REFERENCES

- Benioff, H., 1951, Global strain accumulation and release as revealed by great earthquakes: Geological Society of America Bulletin, v. 62, p. 331-338.
- Benjamin, J., and Cornell, C., 1970, Probability, statistics, and decision for civil engineers: McGraw-Hill, New York.
- de Finetti, R., 1964, Foresight: Its logical laws, its subjective sources (Henry E. Kyburg, Jr., trans.), in Studies in Subjective Probability, Henry E. Kyburg, Jr., and Howard E. Smokler (eds.): John Wiley & Sons, New York.
- Esteva, L., 1976, in Seismic Risk and Engineering Decisions; Developments in Geotechnical Engineering 15: Elsevier Scientific Publishing Company, New York.
- Hagiwara, Y., 1975, A stochastic model of earthquake occurrence and the accompanying horizontal land deformations: Tectonophysics, v. 26, no. 1/2, March, p. 91-101.
- Howard, R. A., 1971, Dynamic probabilistic systems, Vol. II: John Wiley & Sons, Inc.
- Kelleher, J., and McCann, W., 1976, Buoyant zones, great earthquakes, and unstable boundaries of subduction: Journal of Geophysical Research, v. 81, no. 26, p. 4885-4896.
- Kelleher, J., Savino, J., Rowlett, H., and McCann, W., 1974, Why and where great thrust earthquakes occur along island arcs: Journal of Geophysical Research, v. 79, no. 32, p. 4889-4899.
- Knopoff, L., and Kagan, Y., 1977, Analysis of the theory of extremes as applied to earthquake problems: Journal of Geophysical Research, v. 82, no. 36, December.
- Mogi, K., 1968, Some features of recent seismic activity in and near Japan: Bulletin of the Earthquake Research Institute, v. 46, p. 1225-1236.
- Nair, K., and Cluff, L. S., 1977, An approach to establishing design surface displacements for active faults: Sixth World Conference on Earthquake Engineering, New Delhi, India.
- Patwardhan, A. S., Tocher, D., and Savage, E. D., 1975, Relationship between earthquake magnitude and length of rupture surface based on aftershock zones: Geological Society of America, Abstracts with Programs, v. 7, no. 3, February.
- Raiffa, H., 1968, Decision analysis: Addison-Wesley, Reading, Massachusetts.

- Raiffa, H., and Schlaifer, R., 1960, Applied statistical decision theory: Harvard University.
- Rikitake, T., 1975, Statistics of ultimate strain of the earth's crust and probability of earthquake occurrence: *Tectonophysics*, v. 26, no. 1/2, March, p. 1-21.
- Savage, L. J., 1954, The foundations of statistics: John Wiley & Sons, New York.
- Schlaifer, R. O., 1969, Analysis of decisions under uncertainty: McGraw-Hill, New York.
- Selvidge, J., 1972, Assigning probabilities to rare events: Ph.D. Dissertation, Graduate School of Business Admin., Harvard University.
- Shlien, S., and Toksoz, M., 1970, A clustering model for earthquake occurrences: *Seismological Society of America Bulletin*, v. 60, no. 6, p. 1765-1787.
- Stael von Holstein, C.-A.S., 1970, Measurement of subjective probability: *Acta Psychologica*, v. 34, p. 146-159.
- Sykes, L. R., 1971, Aftershock zones of great earthquakes, seismicity gaps, and earthquake prediction for Alaska and the Aleutians: *Journal of Geophysical Research*, v. 76, no. 32, p. 8021-8041.
- Tocher, D., 1958, Earthquake energy and ground breakage: *Seismological Society of America Bulletin*, v. 48, p. 147-153.
- U. S. Geological Survey, 1977, National Oceanic and Atmospheric Administration Hypocenter Data File.
- Winkler, R. L., 1967, The assessment of prior distributions in Bayesian analysis: *Journal of the American Statistical Association*, v. 62, p. 776-800.
- Winkler, R. L., and Murphy, A. H., 1968, Evaluation of subjective precipitation probability forecasts: *Proceedings of the First Conference on Statistical Meteorology*, Boston, Massachusetts.

Table 4. SUMMARY OF HOLDING TIMES USED IN ANALYSIS

Initial State M_i	Following State M_j	Sample Data (Yrs)	Prior Fractiles (Yrs)				Posterior Fractiles (Yrs)				Number of Years for Different Events as Gap Following M_i
			0.25	0.50	0.75	1.0	0.25	0.50	0.75	1.0	
8 ± 0.2	8 ± 0.2	1,2,3,5,17,17,22,26,27, 32,41,1,2,2,3,5,12,14	12	20	30	66	4	10.3	26.6	66	3,2,77,35,21,46, 21,13,10,7,76, 74,20,29
	8.4 ± 0.2	1,1,6,19,27,3,4,6	15	25	34	70	3.9	11.1	31.8	70	
	8.75 ± 0.2	15,5,28	20	35	50	105	13.6	23.6	40.9	105	
8.4 ± 0.2	8 ± 0.2	2,8,13,29,32,4,7,11,11,22	12	20	30	66	8.3	13.9	23.3	66	32,58,30,46,74,36, 78,14,34,26,34,18
	8.4 ± 0.2	1,4,27,4,6,11,29,43	20	30	41	80	4.9	13.5	37.4	80	
	8.75 ± 0.15	1,27,2,10,15	34	50	67	145	3.2	16.6	85.3	145	
8.75 ± 0.15	8 ± 0.2	42,16	18	30	43	85	17.4	25.6	37.5	85	40,20,36
	8.4 ± 0.2	1,2,28,11,16,36	28	40	54	98	4.3	16.5	62.4	98	
	8.75 ± 0.15	19,25	82	100	134	200	39	65.2	109	200	

Table 5. SUMMARY OF TRANSITION STATES USED IN ANALYSIS

Initial State <u>M_i</u>	Sample Data (Mag.)	Prior Fractiles (Mag.)				Posterior Fractiles (Mag.)			
		<u>0.25</u>	<u>0.50</u>	<u>0.75</u>	<u>1.0</u>	<u>0.25</u>	<u>0.50</u>	<u>0.75</u>	<u>1.0</u>
8 ± 0.2	8.7, 7.8, 8, 7.9, 7.9, 8.4, 7.8, 8.3, 7.9, 7.9, 7.9, 7.9, 7.9, 8.3, 8.3, 7.9, 8.3, 7.9, 8.1, 8.3, 8.9, 8.1, 8.4, 7.8, 8.9, 7.8, 8.4, 7.9, 7.8	7.9	8	8.4	8.8	7.8	8	8.5	8.8
8.4 ± 0.2	7.9, 8.3, 8.3, 8.3, 7.8, 7.9, 8.1, 8.6, 8.6, 7.9, 8.9, 8.9, 7.9, 8.3, 7.8, 8.4, 8, 7.8, 8.6, 7.9, 8.3, 8.3, 8.3	7.9	8	8.4	8.8	7.9	8	8.6	8.8
8.75 ± 0.15	8.4, 7.8, 8.3, 8.2, 8.3, 8.7, 8.6, 8.9, 8.3, 8.4	7.9	8	8.4	8.8	7.9	8.1	8.7	8.8

Table 6. SUMMARY OF HOLDING TIMES FOR AREA B

Initial State M_1	Following State M_j	Sample Data (Yrs)	Prior Fractiles (Yrs)				Posterior Fractiles (Yrs)				Number of Years for Different Events as Gap Following M_1
			0.25	0.50	0.75	1.0	0.25	0.50	0.75	1.0	
8 ± 0.2	8 ± 0.2	34,7,40,3,52,6,26,8,16, 10,9,36	12	20	30	66	9.8	16.1	26.4	66	44,28,7,7,39,64,7
	8.4 ± 0.2	15,25,34,70	15	25	34	70	14.3	20.9	30.6	70	
	8.75 ± 0.2	0	20	35	50	105	18.7	28.9	44.6	105	
8.4 ± 0.2	8 ± 0.2	19	12	20	30	66	11.5	17.6	27.1	66	58,72
	8.4 ± 0.2	0	20	30	41	80	22.4	27	32.5	80	
	8.75 ± 0.15	0	34	50	67	145	38.2	45.8	54.7	145	
8.75 ± 0.15	8 ± 0.2	0	18	30	43	85	17.2	25.5	37.8	85	40
	8.4 ± 0.2	0	28	40	54	98	31.7	36.8	42.8	98	
	8.75 ± 0.15	0	82	100	134	200	94.8	101.1	107.7	200	

Table 7. SUMMARY OF TRANSITION STATES FOR AREA B

Initial State <u>M₁</u>	Sample Data (Mag.)	Prior Fractiles (Mag.)				Posterior Fractiles (Mag.)			
		<u>0.25</u>	<u>0.50</u>	<u>0.75</u>	<u>1.0</u>	<u>0.25</u>	<u>0.50</u>	<u>0.75</u>	<u>1.0</u>
8 ± 0.2	8.1,7.9,8.1,8.1,7.9,8.1, 7.8,7.9,8.1,7.9,7.9,8.1	7.9	8	8.1	8.4	7.9	8	8.1	8.4
8.4 ± 0.2	8.3	7.9	8	8.1	8.4	8	8	8.2	8.4
8.75 ± 0.15	7.8	7.9	8	8.1	8.4	7.9	8	8.1	8.4

Table 8. EARTHQUAKE PROBABILITIES FOR THE NEXT 40 YEARS
CALCULATED FROM THE MODEL ($M_0 = 8 \pm 0.2$; $t_0 = 5$ Years)

<u>Probability</u>	<u>Number of Great Earthquakes of Each Magnitude</u>		
	<u>8 ± 0.2</u>	<u>8.4 ± 0.2</u>	<u>8.75 ± 0.15</u>
0.1933	0	0	0
0.1691	1	0	0
0.9840×10^{-1}	2	0	0
0.9539×10^{-1}	0	0	1
0.5852×10^{-1}	1	1	0
0.5627×10^{-1}	0	1	0
0.5593×10^{-1}	1	0	1
0.4684×10^{-1}	3	0	0
0.3966×10^{-1}	2	1	0
0.2371×10^{-1}	2	0	1
0.1981×10^{-1}	3	1	0
0.1849×10^{-1}	4	0	0
0.1770×10^{-1}	0	1	1
0.1724×10^{-1}	1	1	1
0.9683×10^{-2}	1	2	0
0.8271×10^{-2}	2	1	1
0.7256×10^{-2}	3	0	1
0.7229×10^{-2}	2	2	0
0.6807×10^{-2}	0	2	0
0.5505×10^{-2}	0	0	2
0.2764×10^{-2}	1	2	1
0.2490×10^{-2}	1	0	2
0.2451×10^{-2}	0	2	1
0.1828×10^{-2}	0	1	2
0.1081×10^{-2}	1	1	2
0.1067×10^{-2}	1	3	0
0.7314×10^{-3}	2	0	2
0.6698×10^{-3}	0	3	0
0.3025×10^{-3}	0	2	2
0.2575×10^{-3}	0	3	1
0.1035×10^{-3}	0	0	3
0.5808×10^{-4}	0	1	3

Table 9. COMPARISON OF RELATIVE FREQUENCY OF GAP FROM HISTORICAL SEISMICITY DATA WITH PROBABILITIES CALCULATED FROM MODEL ($M_0 = 8 \pm 0.2$)

<u>M_0</u>	<u>t_0, years</u>	<u>Period of Interest, t years</u>	<u>Number of Events with at Least One Earthquake in Time t</u>	<u>Number of Events with No Earthquake in Time t</u>	<u>Relative Frequency of Gap from Data</u>	<u>Probability of Gap from Model</u>
8 ± 0.2	5	5	5	22	0.81	0.74
8 ± 0.2	5	20	12	12	0.50	0.42
8 ± 0.2	30	20	2	3	0.60	0.42
8 ± 0.2	5	40	18	4	0.18	0.19

Table 10. COMPARISON OF RELATIVE FREQUENCY OF GAP FROM HISTORICAL SEISMICITY DATA WITH PROBABILITIES CALCULATED FROM MODEL ($M_0 = 8.4 \pm 0.2$)

<u>M_0</u>	<u>t_0, years</u>	<u>Period of Interest, t years</u>	<u>Number of Events with at Least One Earthquake in Time t</u>	<u>Number of Events with No Earthquake in Time t</u>	<u>Relative Frequency of Gap from Data</u>	<u>Probability of Gap from Model</u>
8.4 ± 0.2	5	5	4	24	0.86	0.77
8.4 ± 0.2	5	20	10	16	0.62	0.40
8.4 ± 0.2	30	20	2	3	0.60	0.58
8.4 ± 0.2	5	40	16	4	0.20	0.24
8.4 ± 0.2	30	40	2	2	0.50	0.31
8.4 ± 0.2	30	5	1	6	0.86	0.88

Table 11. AREA B, COMPARISON OF RELATIVE FREQUENCY OF GAP FROM HISTORICAL SEISMICITY DATA WITH PROBABILITIES CALCULATED FROM MODEL ($M_0 = 8 \pm 0.2$)

M_0	t_0 , years	Period of Interest, t years	Number of Events with at Least One Earthquake in Time t	Number of Events with No Earthquake in Time t	Relative Frequency of Gap from Data	Probability of Gap from Model
8 ± 0.2	5	5	5	10	0.67	0.78
8 ± 0.2	5	20	6	9	0.60	0.29
8 ± 0.2	30	20	3	2	0.40	0.34
8 ± 0.2	5	40	10	2	0.16	0.11

Table 12. SUMMARY OF HOLDING TIMES USED IN PARAMETRIC ANALYSIS

Initial State M_i	Following State M_j	Sample Data (Yrs)	Prior Fractiles (Yrs)				Posterior Fractiles (Yrs)				Number of Years for Different Events as Gap Following M_i
			0.25	0.50	0.75	1.0	0.25	0.50	0.75	1.0	
8 ± 0.2	8 ± 0.2	3,2,46,17,41,2,3,5,60	12	20	30	66	5.4	13.3	32.5	66	19,20,29,34,13,21, 2,30,3,63,15,78, 10,37,78,74,47,20, 46,77,74,51,67,35, 74,25,20
	8.4 ± 0.2	27,17,41,6,6,2,4,2	15	25	34	70	6.2	13.6	29.8	70	
	8.75 ± 0.2	57,26,5	20	35	50	105	14.7	26.2	46.8	105	
8.4 ± 0.2	8 ± 0.2	45,42,29,11,4,32,13, 25,2,55,24,26,28	12	20	30	66	10.7	18.3	31.3	66	55,26,38,36,56,56, 35,28,13,77,26,55, 34,74,76,36,58,32, 30,60,73,46,38,36
	8.4 ± 0.2	22,11,4,2,2	20	30	41	80	4.1	11.9	34.3	80	
	8.75 ± 0.15	29	34	50	67	145	34.6	42.4	52	145	
8.75 ± 0.15	8 ± 0.2	58	18	30	43	85	17.7	27.3	42.1	85	40,20,45
	8.4 ± 0.2	36	28	40	54	98	31.7	36.7	42.4	98	
	8.75 ± 0.15	0	82	100	134	200	94.8	101.1	107.7	200	

Note: Zones were defined by using variable magnitude-rupture length relationships (all areas except Area B; see Figure 4).

Table 13. SUMMARY OF TRANSITION STATES USED IN PARAMETRIC ANALYSIS

Initial State <u>M₁</u>	Sample Data (Mag.)	Prior Fractiles (Mag.)				Posterior Fractiles (Mag.)			
		<u>0.25</u>	<u>0.50</u>	<u>0.75</u>	<u>1.0</u>	<u>0.25</u>	<u>0.50</u>	<u>0.75</u>	<u>1.0</u>
8 ± 0.2	8.4, 8, 8.7, 8.8, 7.9, 7.9, 7.9, 8.3, 8.4, 7.8, 8.3, 8.8, 8.1, 8.4, 8.4, 8.4, 7.8, 8.3, 7.9, 8.3, 7.9	7.9	8	8.4	8.8	7.9	8.1	8.5	8.8
8.4 ± 0.2	7.9, 7.8, 8.1, 7.9, 7.9, 8.1, 7.8, 7.9, 7.8, 8.4, 8, 7.9, 8.3, 8.3, 8.3, 8.6, 8.3, 8.6, 8.2	7.9	8	8.4	8.8	7.9	8	8.4	8.8
8.75 ± 0.15	7.8, 8.3	7.9	8	8.4	8.8	7.8	8	8.6	8.8

Note: Zones were defined by using variable magnitude-rupture length relationships (all areas except Area B; see Figure 4).

(Based on Sykes, 1971)

(a) ALASKA ALEUTIAN AREA, RUPTURE SURFACES OF RECENT GREAT EARTHQUAKES

(b) DELINEATION OF ZONES, AREA A

Fig. 1 EXAMPLE OF ZONE IDENTIFICATION

Fig. 2 ALASKA ALEUTIAN AREA, CROSS-SECTIONS SHOWING TEMPORAL DISTRIBUTION OF EARTHQUAKES GREATER THAN $M = 5$

Fig. 3 SCHEMATIC REPRESENTATION OF THE TRAJECTORY OF A SEMI-MARKOV PROCESS

EXPLANATION

(A) Tonga, Kermadec	(D) Ryukyu, Philippines	(G) Alaska—Aleutians
(B) New Hebrides—New Guinea	(E) South America	
(C) Kurile, Kamchatka, NE Honshu, Marianas	(F) Central America	

Fig. 4 AREAS OF CIRCUM PACIFIC BELT INCLUDED IN THE STUDY

Fig. 5 SCHEMATIC REPRESENTATION OF BAYESIAN PROCEDURE

Fig. 6 EXAMPLE OF PRIOR AND POSTERIOR PROBABILITY DISTRIBUTIONS FOR HOLDING TIMES USED IN ANALYSIS

Fig. 7 EXAMPLE OF PRIOR AND POSTERIOR PROBABILITY DISTRIBUTIONS FOR TRANSITIONS USED IN ANALYSIS

Fig. 8 PROBABILITY OF CONTINUATION OF A SEISMIC GAP FOR DIFFERENT INITIAL CONDITIONS

Fig. 9 COMPARISON OF PROBABILITY OF CONTINUATION OF GAPS COMPLIED FROM DATA WITH THE PROBABILITIES COMPUTED FROM THE MODEL

M_0 = magnitude of last great earthquake
 t_0 = waiting time since last great earthquake

Fig. 10 COMPARISON OF MAGNITUDE DISTRIBUTION RELATIONSHIPS DERIVED FROM THE MODEL FOR DIFFERENT INITIAL CONDITIONS WITH RELATIONSHIPS BASED ON THE POISSON MODEL

Fig. 11 COMPARISON OF NUMBER OF EARTHQUAKES ≥ 7.8 , (N) IN 40 YEARS USING THE SEMI-MARKOV MODEL WITH THE POISSON MODEL (ZONE 4 ALASKA-ALEUTIAN AREA)

Fig. 12 PROBABILITY OF CONTINUATION OF A SEISMIC GAP FOR DIFFERENT INITIAL CONDITIONS (HOLDING TIME INCREASED APPROXIMATELY BY A FACTOR 1.5 AND 6)

Fig. 13 PROBABILITY OF CONTINUATION OF A SEISMIC GAP FOR DIFFERENT INITIAL CONDITIONS USING VARIABLE MAGNITUDE-RUPTURE LENGTH RELATIONSHIPS

M_0 = magnitude of last great earthquake
 t_0 = waiting time since last great earthquake

Fig. 14 INFLUENCE OF INITIAL CONDITIONS ON SELECTED DESIGN EARTHQUAKES

Fig. 14. EFFECT OF M. & S. ON THE PRESSURE ON M. & S. (lb./sq. in.)

UPLIFT HISTORY AND EARTHQUAKE RECURRENCE AS DEDUCED FROM MARINE
TERRACES ON MIDDLETON ISLAND, ALASKA

George Plafker

U.S. Geological Survey

Menlo Park, CA 94025

Meyer Rubin

U.S. Geological Survey

Reston, VA 22092

Prepared for Research Conference on Seismic Gaps
and Earthquake Recurrence, May 25-27, 1978

Abstract

Middleton Island, near the margin of the continental shelf in the northern Gulf of Alaska, has emerged from the sea during six major episodes of coseismic uplift from oldest to youngest of about 7 m, 8 m, 6 m, 9 m, 7.5 m, and 3.5 m which are recorded by marine terraces. All but the youngest uplift have been dated by radiocarbon methods at roughly 4,300, 3,800, 3,100, 2,390, and 1,350 years before present, respectively, and the most recent uplift occurred during the March 27, 1964 Alaska earthquake^{1/}. Recurrence time for these movements is on the order of 500 to 1,350 years. Average uplift rate is approximately 1 cm/yr since the island first emerged from the sea 4,300 years ago and there appears to be an abrupt decrease in the rate of uplift to 0.6 cm/yr in the interval preceding uplift of the 1964 terrace. Uplift of the terraces is believed to result from local imbrication along thrust faults within the upper plate during major earthquakes related to slip on the Aleutian megathrust. Available data from the Middleton Island terraces and 1964 coseismic displacements suggest that much of the interseismic strain accumulated in the eastern part of the 1964 Alaska earthquake focal region has yet to be released. Together with historic seismicity, the data suggest that the accumulated strain is most likely to be released during one or more future earthquakes along the 100- to 200-km-long segment of plate boundary extending eastward from the vicinity of Middleton Island to Icy Bay.

^{1/} All dates used in this paper are in radiocarbon years. Application of tree ring corrections will increase the ages of the four older terraces to about 4,900, 4,300, 3,400, and 2,500 years and result in slightly reduced average uplift rates and a longer uplift recurrence intervals. As these differences are within the limits of error of the data and do not affect the conclusions of this paper no attempt has been made to correct the radiocarbon dates.

INTRODUCTION

Middleton Island is located near the edge of the continental shelf in the northern Gulf of Alaska. Structurally, it is situated near the leading edge of the North American plate where it is being relatively underthrust by the Pacific plate along the Aleutian Trench. The island, which trends northeast-southwest, is about 8 km long, 1.6 km wide, and has a maximum elevation of 47.6 m (fig. 1).

During the March 27, 1964 Alaska earthquake the island was abruptly uplifted several meters with resultant exposure of a marine terrace. This terrace, together with five higher and older radiometrically dated marine terraces on the island provide an exceptionally clear record of intermittent earthquake-related tectonic movements during the past four millenia. It is the purpose of this paper to briefly outline data relevant to the nature and age of these terraces and to speculate on the tectonic history recorded by them.

Previous work

The marine terraces at Middleton Island were first described, on the basis of a brief air reconnaissance, by Capps (1933) who correctly inferred that the terraces recorded pulsating uplifts relative to sea level. In 1949, Don J. Miller visited the island for about 1 week during which time he carried out a remarkably thorough examination of both the bedrock geology and marine terraces (Miller, 1953). Despite the fact that he had no adequate topographic maps and radiocarbon dating of the terrace materials was not yet feasible, Miller succeeded in delineating the pre-1964 terraces and he speculated that they were probably of latest Pleistocene or Holocene age and tectonic in origin.

Present investigation

Our study of the Middleton Island terrace sequence was carried out by one or both of the authors during brief visits to the island in 1963, 1965, 1967, and 1970. The objectives were to date the terraces with radiocarbon methods, obtain precise data on the terrace elevations with particular emphasis on the heights of beach angles, and determine how much information on the terrace history could be deduced from the unconsolidated deposits on them. As part of this study a special base map at scale 1:10,000 with 5-ft contour intervals was prepared by photogrammetric methods by the Topographic Division of the U.S. Geological Survey from high-quality airphotos taken by the U.S. Coast and Geodetic Survey after the 1964 earthquake. Additional control on the elevation and configuration of the terraces is provided by three lines of levels surveyed across the island in 1963 by the senior author.

DESCRIPTION OF THE TERRACES

The most striking feature of Middleton Island is the steplike character of its surface which consists of a series of six terraces with average beach angle elevations above MLLW of about 6.4, 14, 23, 29, 37 and 44 m (figs. 1-3). Beach angle elevations for terraces I-V are estimated to be accurate within $1\frac{1}{2}$ m and the terrace heights are thus indicated as 3-m-wide bands on figure 4. This uncertainty is due to a combination of the possibility of slight tilting of the island which gives different beach angle elevations for the higher terraces across the island, mantling of the beach angles by talus, slope wash, and peat which conceal it in most places, and initial variations in the elevation of the beach angles. Steep sea cliffs, cut into moderately well indurated

early Pleistocene marine strata that dip northwest as much as 30° , are developed around much of the island as a result of rapid erosion by the prevailing southeasterly and southwesterly storm waves. The northwestern end of the island is largely constructional, being made up of sand and gravel derived by longshore drift from erosion of outcrop on the southern part of the island. In this area the lower terraces slope off gradually to merge with the present beach and an area of irregular dunes.

Terrace uplift history

Long-continued late Quaternary emergence and tilting of Middleton Island is indicated by the exposure on the island of a beveled early Pleistocene marine section roughly 1,000 m thick that dips $22-30^{\circ}$ northwest (Plafker and others, 1975; Miller, 1953). In addition, marine seismic reflection data on the adjacent shelf indicate that even the youngest Quaternary strata are deformed in a complex of structural highs and lows (Plafker and others, 1975). The marine terraces of Middleton Island, as interpreted on figures 2 and 3, record pulsating emergence of the Middleton platform in seven stages. Each terrace consists of a wave-cut bedrock surface overlain by stratified deposits of gravel, sand, and mud formed in the associated beach, lagoon, and shallow marine environments, and of peat formed in a freshwater or subaerial environment after emergence. Following the convention established by Miller (1953), the stages of emergence and the resulting terraces and associated features are designated by the Roman numerals I-VI with I representing the highest terrace and the earliest recorded stage of uplift, and VI representing the terrace formed in 1964. The following description of the sequence of terrace development is modified after Miller (1953). The major differences are in the

elevations. Miller's were from relatively inaccurate altimeter readings whereas ours are based on the large-scale topographic map, level lines, and altimeter readings. In addition, we recognize a narrow stage V terrace on the southeast coast of the island that was not delineated by the previous reconnaissance work.

Preservation of so many strikingly developed terraces in the small area of Middleton Island is the result of a fortuitous balance between uplift and rate of marine erosion. The marginal slope of steeper gradient below a depth of 27 m around the present Middleton platform may represent part of the surface of the ancestral island or submarine bank initially exposed to marine planation. Projection of this slope to the present level of the stage I terrace indicates that the island or submarine bank exposed to marine planation at the outset of stage I may have been as much as 4.8 km wide and 16 km long. Other stages of uplift may have intervened between any two successive stages represented by the terraces now preserved on Middleton Island, but, if so, they are not recorded. Evidence to indicate emergence of the Middleton platform above the present stand also is lacking.

At the outset of stage I, partly indurated bedrock forming the Middleton platform was placed within the limit of wave action, either by (1) differential upwarping accompanied by tilting of the bedrock; (2) regional uplift of the sea bottom on which the platform already stood as a topographic high; or (3) eustatic lowering of sea level. The platform did not necessarily emerge as an island at this stage but, instead, may have formed a submarine bank. The marine terrace formed during stage I is preserved on the highest part of Middleton Island. The average altitude

of the terrace surface ranges from about 44 m near the middle to about 40 m at either end. In the southern, broader part of the terrace (fig. 3) the wave-cut bedrock surface lies for the most part at an altitude of 40-41 m but rises abruptly to altitudes of up to 48 m in many conical to ellipsoidal mounds. These mounds, which are grouped in several rows that parallel the strike of the bedrock, represent erosional remnants formed along layers of more resistant bedrock. The unconsolidated deposits resting on the bedrock surface in the southern part of the stage I terrace range in thickness from about 30 cm (atop some of the mounds) to about 3 m. Auger holes, pits, and natural sections in the present sea cliff at the margin of the southern part of the terrace show as much as 1-1.3 m of interbedded sand and mud, locally gravelly, overlain by 0.3-2 m of peat. At many places the peat rests on the surface of the bedrock.

After stage I the ancestral Middleton Island, including the preserved part of the stage I terrace and an adjacent, probably much larger, area, emerged in pulses to the present stand. The remaining parts of the terraces and adjacent wave-cut cliffs of stages II, III, and IV form roughly concentric bands around the south and east margins of the stage I terrace. The stage V terrace, in contrast, is preserved only in two narrow bands on the northwest and northeast sides of Middleton Island. The wave-cut cliffs formed during stages V and VI cut all the higher terraces.

The beach angle of the stage II terrace (toe of the adjacent wave-cut cliff) is at an average altitude of 37 m, indicating vertical movement of the land with respect to sea level of at least 7 m between stages I and II. The inner margins of the lower uplifted terraces are at altitudes of about 29 m, 23 m, and 14 m, indicating additional uplift of at least 8 m between

stages II and III, 6 m between stages III and IV, 9 m between stages IV and V, and 7.6 m between stages V and VI. The most recent pulse of emergence relative to sea level at Middleton Island occurred at the time of the great earthquake of March 27, 1964. During this seismic event the island was tectonically uplifted 3.4 m (Plafker and Rubin, 1967). As a result, a terrace surface was permanently exposed above high tide level around the perimeter of the island. This terrace, labeled VI on figures 2 and 3, varies in width from about 20 m to 150 m and is the narrowest of the six terraces on the island. Below this terrace the littoral zone between the low tide and high tide shorelines (0 and 10-ft contours, fig. 1) is a broad platform that has an average width of about 200 m and a maximum width of 700 m near the southwest and northeast ends of the island.

Terrace deposits

In the northern, narrower part of the stage I terrace the altitude of the bedrock surface decreases northward, but the general level of the terrace surface is maintained by a corresponding increase in the thickness of the unconsolidated deposits. This part of the terrace is, therefore, a constructional feature.

The unconsolidated deposits of the stage II and stage III terraces, as exposed in natural sections and in a few pits and auger holes, are similar to the previously described deposits of the stage I terrace.

Near the north end of the stage IV terrace the bedrock surface falls from an altitude of 12 m to sea level within about 100 m. Here, as at the north end of the stage I terrace, the general level of the terrace is maintained for some distance north of the point where the bedrock surface

begins to fall off, by a corresponding increase in thickness of the unconsolidated deposits.

The hooked projection at the north end of the stage III terrace, which is entirely composed of gravel in a ridge that rises to 6 m above the level of the stage IV terrace, probably formed as a spit that merges at its south end with the stage III beach deposits.

The stage IV terrace, lying along the southeast side of Middleton Island, is mantled by thick deposits of predominantly sand and gravel with ridges of stratified sand and gravel that rise as much as 2 m above the general level of the terrace. These ridges are believed to have been a series of either spits or offshore bars behind which finer grained sediment was trapped.

The stage V terrace at the northwestern end of the island and along the northeastern side of the island is mantled with a sequence of lagoonal deposits that accumulated behind offshore gravel bars. The lagoonal deposits are especially well exposed along the northeastern shore of the island where they are incised by the stage VI sea cliff. A section of these deposits, which are up to 4 m thick, is shown on figure 4.

Around most of the shore of Middleton Island the wave-cut bedrock surface being formed at the present stage either is exposed or is covered by not more than a few tens of centimeters of unconsolidated deposits that are constantly being reworked and redistributed by wave and current action. These deposits consist of sand and pebble gravel that form the narrow beach and well-rounded cobbles and boulders that form spits or cobble-boulder pavements or are scattered at random on the bedrock surface. The bedrock surface probably slopes off at the north end of the island in the

vicinity of the sand-gravel spit, for it is not exposed there even at the lowest tide stage. Notations on U.S. Coast and Geodetic Survey hydrographic chart 5422 show predominantly rock bottom, with scattered patches of sand and gravel, out to and at some places beyond, the 27 m depth contour around the island.

RADIOCARBON DATA

The ages of the pre-1964 terraces have been determined by radiocarbon dating of 24 samples of fossil driftwood in the deposits on the terraces or peat from within or on the terrace deposits. Sample localities are shown on figure 3 and the analytical data are given in table 1. All ages are given in radiocarbon years.

Although Middleton Island is almost treeless, driftwood is abundant on the present beaches and has been found on all terraces except for stage II. Radiocarbon dates on driftwood are believed to provide the most reliable times for terrace emergence where they are incorporated into beach deposits or lag gravels immediately above the bedrock surface.

Peat deposits develop readily in the cool, moist climate that prevails at Middleton Island. The peat, however, may begin to accumulate at any time after terrace emergence so that the basal peat can be considered to provide only a minimum age for the surface on which it occurs. Furthermore, because the basal peat is always subject to contamination with roots from above, the radiocarbon ages may be younger than the initial time of peat formation.

Figure 4 shows the ages of radiometrically dated wood and peat samples plotted against the average shore angle elevation of the five older terraces. Also shown is the height of the 1964 terrace and a eustatic sea

level curve after Coleman and Smith (1964) for the past 5,000 years.

The data suggest the following ages for the older terraces:

I - $4,300 \pm 250$ b.p., II - approximately 3,800 b.p., III - $3,100 \pm 250$ b.p., IV - $2,390 \pm 200$ b.p., and V - $1,350 \pm 200$. The lowest terrace, VI, formed in 1964 and the lower part of this surface is approximately at the mean higher high tide level (10-ft contour on figs. 1 and 3).

Terrace I is well dated by two wood samples (3, 5) and peat from the base of a section 238 cm thick (4). Samples 1 and 2 from this same surface yield ages that are as much as 1,000 years younger than terrace emergence.

The age of terrace II is poorly constrained as no driftwood was found in the deposits on its surface. Sample 7 provides a minimum age of 3,590 years and the second peat sample (6) formed long after emergence of the terrace. We have inferred a linear uplift rate on the basis of the better data obtained from the adjacent terraces (fig. 4) which yields an extrapolated age of about 3,800 years for this surface.

The age of terrace III is based on the two oldest peat samples (12 and 13). A driftwood sample (14) from this terrace is anomalous in that it is roughly 1,000 years older than the deposits in which it occurs. Because the age of the sample falls in the age range of dated driftwood from the stage I terrace, we believe that it was eroded out of the highest terrace and subsequently incorporated into the stage III beach deposits. This is analogous to the present process whereby abundant driftwood from the older terraces--especially stages I and V--has been deposited on the modern beaches.

The age of the stage IV terrace is closely controlled by two dates on driftwood (15, 16) that are essentially the same within the limits of the dating method.

The stage V terrace is very well dated by three wood samples (21, 23, 24) and one peat sample (22) that yield reasonably concordant ages. This terrace is unique in that it has a sequence approximately 4 m thick of unconsolidated deposits containing abundant driftwood and some peat on it along the northeast side of the island. These deposits are believed to represent beach and lagoonal deposits that were deposited after emergence of the terrace approximately 1,350 years ago as indicated by the columnar section on figure 4. Sample 24, in lag gravel on the terrace surface, dates initial uplift as do samples of comparable age from this terrace on the opposite side of the island (22, 23, 24). Subsequently, a lagoon developed that was gradually filled in with sediment; the age of some of these deposits is given by samples 18 (660 ± 250 b.p.) and 17 (460 ± 250 b.p.).

COSEISMIC MOVEMENTS AND POSSIBLE INTERSEISMIC MOVEMENTS

The Middleton Island terraces were cut during long periods of stability or slight submergence relative to sea level and were elevated during brief intervals of coseismic uplift. Two curves depicting the most probable uplift histories are depicted on figure 5. For the pre-1964 stages, it is not known whether the indicated uplifts represent single events or more than one event that are relatively closely spaced in time. Because sea level was at or near its present stand during the period in which the terraces were formed (fig. 4), eustatic sea level changes did not play a significant part in their formation. Minimum coseismic uplifts

are 7 m for stage I, 8 m for stage II, 6 m for stage III, 9 m for stage IV, 7.5 m for stage V, and 3.5 m for stage VI.

The solid curve on figure 5 assumes no interseismic recovery, and the dashed line shows a more probable situation in which there is a significant, but unknown, component of interseismic submergence. Interseismic submergence has been observed over an extensive region of Prince William Sound, the mainland coast, and at Kayak and Wingham Islands where as much as 4.7 m submergence occurred in the interval between uplift of the stage V and VI terraces at Middleton Island (Plafker and Rubin, 1967; Plafker, 1969). It is uncertain whether the submergence extended as far out on the continental shelf as Middleton Island although submergence may be indicated by two lines of evidence. One is that the pre-earthquake high sea levels reached to the base of the prominent sea cliff that encircles much of the island, which could be due to either a very long period of relative stability and erosion or to some submergence since the stage V uplift of the island. The second is that unconsolidated deposits, such as those on the stage V terrace (fig. 4), were deposited in a lagoon on the lower part of the terrace long after emergence. It is not known whether the observed relationships are indicative of interseismic submergence or normal marine processes that resulted in accumulation of these lagoonal deposits. However, the time lag between coseismic uplift and accumulation of the lagoonal deposits, the high sedimentation rate of 2 m/200 years indicated by the ages and stratigraphic positions of samples 17 and 18, and the evidence for widespread coastal submergence elsewhere in the region are all suggestive of interseismic submergence. From figure 5, it should be apparent that if interseismic submergence did occur, the true coseismic uplift for the pre-1964 terraces is equal to the indicated uplift plus the amount of interseismic subsidence.

RECURRENCE INTERVALS AND UPLIFT RATES

The 1964 coseismic uplift, together with available radiometrically dated samples from the five higher terraces indicate recurrence intervals on the order of 500 years between stages I and II, 700 years between II and III, and III and IV, 1,040 years between IV and V, and 1,350 years between V and VI (fig. 5). As noted earlier, it should be emphasized that the indicated coseismic uplift could represent either a single earthquake or more than one earthquake that are relatively closely spaced in time.

The average uplift rate for the Middleton Island terrace sequence, which is slightly more than 1 cm/yr (1.05 cm/yr), is among the highest documented rates of tectonic uplift in the world. In detail, however, the rate of uplift appears to show significant and systematic changes with time as indicated by the light solid line on figure 5. Within the limitations of the data, the uplift rates averaged about 1.25 cm/yr for the 1,200 years between stages I to III, 0.87 cm/yr for 1,650 years between stages III to V, and 0.59 cm/yr for 1,350 years between the stage V and VI uplifts.

FUTURE EARTHQUAKES

The likelihood of another great earthquake and related uplift at Middleton Island cannot be predicted from the available data. It is clear that the 1964 uplift of 3.5 m, which followed the longest interseismic interval recorded in the marine terraces at Middleton Island, is significantly less than the steps between the older terraces which range from 6 to 9 m and average 7.5 m (fig. 5). As a consequence, the resulting 1964 terrace is less than half the width of the older ones (fig. 2).

If the uplift rate at Middleton Island was constant over the last few

thousand years, it would be possible to predict that more than half of the accumulated strain is not represented by the 3.5 m of uplift in 1964. This could be taken to imply that the region may be overdue for another tectonic earthquake involving additional uplift of the island of about 3.5 m (Plafker, 1972). These data, together with the historic seismic record (Sykes, 1971) have been used to identify the segment of the Gulf of Alaska continental margin extending east of Middleton Island as a region where one or more major earthquakes are highly probable (Plafker and others, 1975). The causative plate boundary fault for the postulated event could overlap the Middleton Island area and result in enough uplift to bring it up to the apparent long-term trend.

On the other hand, one could argue that the rate of uplift, as shown on figure 4, has been decreasing progressively over the last 4,000 years and that the 3.5 m of uplift in 1964 has relieved most of the accumulated strain. If so, the uplift rate must have abruptly diminished by about 0.25 cm/yr during the last 1,350 years. Although such a change is possible, it does not appear to be likely in view of the long history of active uplift, tilting, and truncation that is recorded in the bedrock and marine terrace sequence at Middleton Island.

Another alternative possibility is that part of the accumulated strain not released during the 1964 earthquake will be released by postseismic creep deformation during a time interval that will be short relative to the earthquake recurrence time. The combined coseismic and postseismic uplift could be comparable to the amount of uplift that has affected the older terraces. Northwestward postseismic tilt of Middleton Island from 1966 to 1975 suggests that at least local adjustments to the coseismic deformation have apparently persisted for a period

of 10 years (Prescott and Lisowski, 1977). The measured northwestward tilt, which is in the same sense as the long term tilt recorded in the dipping Pleistocene strata that underlie the island, has been tentatively interpreted by Prescott and Lisowski (1977) as indicative of either 2 1/2 m/yr slip on the flat-dipping Aleutian megathrust or 1 m/yr slip on the more steeply dipping thrust fault that is inferred to bound the Middleton Shelf on the southeast (fig. 8C). No data on post-1964 elevation changes relative to sea level are available for Middleton Island to corroborate the observed tilting. However, the closest permanent tide gage, located at Cordova 100 km north of Middleton Island on the mainland, has shown a 6 percent decrease in the 2 m coseismic uplift during the decade following the earthquake (table 2). Unfortunately, no other tide gage data are available within the area that was uplifted during the 1964 earthquake. However, observations by one of us (George Plafker) along the mainland coast since the earthquake indicate that there have been no measurable postseismic vertical movements on the scale of those required by Prescott and Lisowski (1977) to account for the tilting at Middleton Island.

An independent check on the recurrence intervals indicated by the terrace data is from the measured horizontal strains associated with the 1964 earthquake and the relative plate motions in the region as deduced from oceanic paleomagnetic data. Assuming that convergence in the eastern Aleutian Trench is uniform at a rate of 5.5 cm/yr (Minster and others, 1974) and that the surface component of horizontal displacement measured by triangulation surveys is at least 20 m (Plafker, 1969), the strain released could have accumulated in only 360 years if the strain were

purely elastic. Thus, even if we assume significant amounts of aseismic creep and permanent deformation, it is clear that the 1964 event could not have released all the horizontal strain accumulated during the period of 1,350 years preceding the earthquake. Gradual postseismic horizontal extension in the area of the triangulation survey should presumably result in elastic thinning and further subsidence relative to sea level. In fact, however, all four permanent tide gages in the region that subsided have shown the opposite effect during the decade after the earthquake, with recoveries on the order of 12 to 34 percent of the amount of earthquake-related subsidence (table 2).

In summary, the accumulated Middleton Island terrace data suggest recurrence intervals of 500 to 1,350 years for large arc-related events of the 1964 type. The data from terrace uplift steps and rates at Middleton Island, together with the results of triangulation resurveys in the earthquake-affected region suggest that at least half of the strain accumulated during the 1,350 years that preceded the 1964 earthquake has yet to be released, assuming no significant aseismic prequake creep. The accumulated strain could be released either by aseismic creep or in one or more large earthquakes over a time interval that is short, relative to the interval between successive terrace uplifts. Because the tide gage data indicate recovery rather than continued gradual strain release in the decade following the 1964 earthquake, it appears more likely that any residual accumulated strain will be released during future earthquakes along the Pacific-North American plate boundary.

A SEISMIC GAP IN THE NORTHERN GULF OF ALASKA?

The general tectonic setting of Middleton Island is on the upper plate above a low-angle thrust fault along which the Pacific plate is relatively underthrusting the North American plate in a northwesterly direction (fig. 7). As a consequence of this relative motion, nearly orthogonal convergence is presently occurring along the eastern part of the Aleutian megathrust and Pamplona fault zone, whereas dextral slip characterizes the Fairweather and Queen Charlotte transform faults. The structurally complex area between the transform and underthrust zones in the vicinity of Malaspina Glacier is the focal region of the series of great earthquakes and tectonic displacements that occurred in the vicinity of Yakutat Bay in 1899 (Tarr and Martin, 1912; Thatcher and Plafker, 1977). Marine geophysical data indicate that an oblique underthrust fault system, the Transition fault, lies along the base of the continental slope between the eastern end of the Aleutian Trench and the northern end of the Queen Charlotte fault. This is a zone along which late Cenozoic deformation has occurred locally and which may still be weakly active seismically. It can be considered as a subhorizontal zone of oblique subduction between the Pacific plate and North American continental crust. The present shift of the plate boundary to the Fairweather-Pamplona system is interpreted as an evolutionary process in which a complicated oblique arc juncture is being converted into a much simpler right-angle juncture (Plafker and others, 1978).

According to the present interpretation, there should be no major seismic gap in the area east of Icy Bay because the transform segment of the plate boundary has been largely relieved of accumulated elastic strain

energy by the 1958 Lituya Bay earthquake on the Fairweather fault (Sykes, 1971) and by the 1972 Sitka earthquakes on its presumed offshore extension to the southeast (Page, 1974). Furthermore, the 1899 Yakutat Bay earthquakes relieved strain in the complex zone near Malaspina Glacier and possibly on the northern part of the Fairweather fault (Thatcher and Plafker, 1977). Thus, the major remaining seismic gap along the Gulf of Alaska margin appears to be the segment, about 100-200 km long, between the offshore Pamplona zone and the eastern part of the 1964 earthquake focal region to the west (fig. 7).

MECHANISM OF TERRACE UPLIFTS

A major unresolved question is how are terraces uplifted in steps during major tectonic earthquakes? Clearly, the worldwide historical record shows that this phenomenon is primarily related to thrust-type earthquakes in arc environments. In the Gulf of Alaska, geochronologic studies of terrace sequences on the mainland west of Icy Bay and near Lituya Bay (Hudson and others, 1976) together with the Middleton Island data indicate that local diastrophic structural growth, rather than regional tectonic deformation, is probably the dominant mechanism of terrace formation. This conclusion is based on the differences in tectonic setting between the three areas of terraces, the general lack of correlation of terrace ages and heights along the Gulf of Alaska margin, and the occurrence of all three terrace sequences on the flanks of growing anticlinal structures (Hudson and others, 1976).

Figure 8 illustrates three general cases of coseismic deformation across the area affected by tectonic deformation during the 1964 earthquake and the resultant inferred or observed vertical surface displacements.

The section line is oriented in a northwest-southeast direction through Middleton Island (line A-A', fig. 6).

In figure 8A deformation within the upper plate is considered to be entirely elastic and the coseismic slip is along the megathrust and equal in amount to the preseismic horizontal strain. The resultant coseismic surface displacements would be primarily subsidence due to elastic extension and attenuation of the upper plate with a small component of uplift at the leading edge of the upper plate due to updip thrusting. Such movement clearly could not result in uplift of terraces such as those at Middleton Island.

In figure 8B part of the stored elastic strain is dissipated by imbrication on the thrust faults that broke the surface at Montague Island (Plafker, 1967) and the remainder results in warping of the leading edge of the upper plate due to a "hang-up" on the megathrust. In this model the upper plate is tectonically shortened by approximately the amount of the preseismic horizontal strain. The 1964 coseismic vertical surface displacements are compatible with this model, but the high rate of uplift and tilting locally in the Middleton Island area are not satisfactorily explained by regional warping.

The preferred model, shown in figure 8C, attributes the observed uplift to the Montague Island faults and two or more submarine thrust faults within the upper plate. According to this model most, if not all, of the preseismic horizontal strain is taken up by imbricate thrust faulting within the upper plate. Such a model can adequately account for the observed coseismic uplift as well as the long-term deformation at Middleton Island. It is also compatible with marine geophysical data

indicating that the continental margin in the region of near-orthogonal convergence is characterized by a series of fault-bounded asymmetrical anticlinal folds and broad intervening synclines that involve deformation of late Cenozoic strata (Bruns and Plafker, 1975; Plafker and others, 1975).

In conclusion, the combined data on structure, terraces, and 1964 coseismic displacements suggest that terrace formation is related to growth of local anticlinal structures within the upper plate and that the positions of these structures change with time. A consequence of this conclusion is that terraces are not likely to be correlatable from place to place in this type of tectonic environment and it is unwarranted to assume constant uplift (or subsidence) rates over long periods of time.

References cited

- Bruns, T. R., and Plafker, George, 1975, Preliminary structural map of part of the offshore Gulf of Alaska Tertiary Province: U.S. Geol. Survey Open-File Map 75-508.
- Capps, S. R., 1933, An air reconnaissance of Middleton Island, Alaska: Jour. Geology, v. 41, p. 728-736.
- Coleman, J. M., and Smith, W. G., 1964, Late Recent rise of sea level: Geol. Soc. America Bull., v. 75, no. 9, p. 833-840.
- Hudson, Travis, Plafker, George, and Rubin, Meyer, 1976, Uplift rates of marine terrace sequences in the Gulf of Alaska: U.S. Geol. Survey Circ. 722, p. 11-13.
- Miller, D. J., 1953, Late Cenozoic marine glacial sediments and marine terraces of Middleton Island, Alaska: Jour. Geology, v. 61, no. 1, p. 17-40.
- Minster, J. B., Jordan, T. H., Molnar, P., and Haines, E., 1974, Numerical modelling of instantaneous plate tectonics: Royal Astron. Soc. Geophys. Jour., v. 36, p. 541-576.
- Page, R. A., 1974, Evaluation of seismicity and shaking at offshore sites: Offshore Technology Conf., 7th, Houston, Texas Proc., v. 3, p. 179-190.
- Plafker, George, 1967, Surface faults on Montague Island associated with the 1964 Alaska earthquake: U.S. Geol. Survey Prof. Paper 543-G, p. G1-G42.
- _____, 1969, Tectonics of the March 27, 1964 Alaska earthquake: U.S. Geol. Survey Prof. Paper 543-I, 74 p.
- _____, 1972, Alaskan earthquake of 1964 and Chilean earthquake of 1960: Implications for arc tectonics: Jour. Geophys. Research, v. 77, p. 901-925.

- Plafker, George, Bruns, T. R., and Page, R. A., 1975, Interim report on the petroleum resource potential and geologic hazards in the Outer Continental Shelf of the Gulf of Alaska Tertiary Province: U.S. Geol Survey Open-File Report 75-592, 74 p.
- Plafker, George, Hudson, Travis, Rubin, Meyer, and Bruns, Terry, 1978, Late Quaternary offsets along the Fairweather fault and crustal plate interactions in southern Alaska: Canadian Jour. Earth Sci. [in press].
- Plafker, George, and Rubin, Meyer, 1967, Vertical tectonic displacements in south-central Alaska during and prior to the great 1964 earthquake: Jour. Geosciences (Osaka City Univ.), v. 10, art. 1-7, p. 1-14.
- Prescott, W. H., and Lisowski, M., 1977, Deformation at Middleton Island, Alaska, during the decade after the Alaska earthquake of 1964: Seismol. Soc. America Bull., v. 67, no. 3, p. 579-586.
- Sykes, L. R., 1971, Aftershock zones of great earthquakes, seismicity gaps, and earthquake prediction for Alaska and the Aleutians: Jour. Geophys. Research, v. 75, p. 8021-8041.
- Tarr, R. S., and Martin, Lawrence, 1912, The earthquakes at Yakutat Bay, Alaska in September, 1899: U.S. Geol. Survey Prof. Paper 69, 135 p.
- Thatcher, Wayne, and Plafker, George, 1977, The Yakutat Bay, Alaska earthquakes; seismograms and crustal deformation: Geol. Soc. America Abs. with Programs, v. 9, no. 4, p. 515.

FIGURE CAPTIONS

Figure 1. Location and topography of Middleton Island.

2. Profiles across Middleton Island showing marine terraces (Roman numerals I-VI). Arrows indicate the beach angle location for the 1964 terrace; datum is mean lower low water.
3. Map of marine terraces on Middleton Island, and locations of radiometrically dated samples listed in table 1.
4. Diagram of terrace height versus age. Locations of numbered samples are shown on figure 3 and date information is presented in table 1. The inset at right is a stratigraphic column for the sequence of unconsolidated deposits on the stage V terrace. Eustatic sea level curve is after Coleman and Smith (1964).
5. Generalized diagram showing average terrace height and minimum tectonic uplift per event(s), versus terrace age and approximate recurrence interval. The solid curve shows an inferred uplift sequence assuming no interseismic vertical movement; the dashed line assumes some interseismic subsidence. The light solid line indicates the average uplift rate between terraces.
6. Map showing the distribution of tectonic uplift and subsidence in south-central Alaska related to the March 27, 1964 Alaska earthquake (Plafker, 1969). Isobases are in meters.
7. Map showing locations of earthquake epicenters and late Cenozoic faults in the northern Gulf of Alaska region. Large arrow indicates the inferred movement direction of the Pacific crustal plate relative to North America. From Plafker and others, 1975.

Figure 8. Diagrammatic vertical displacement profiles (above) and structure sections through the crust and upper mantle (below) across the northern part of the region affected by the 1964 earthquake (line A-A', fig. 6). Figure A assumes all slip occurs along the underthrust plate boundary (Aleutian megathrust) at time of earthquake and the profile above shows the predicted vertical displacements at the surface which would be associated with such movement. Figure B shows the vertical displacements resulting from a combination of imbricate faulting within the upper plate and warping of the upper plate due to a "hang-up" on the megathrust. Figure C shows the step-like displacement pattern that would result from multiple imbrication on thrust faults within the upper plate.

Table 1.--Radiocarbon-dated samples from marine terraces on Middleton Island, Alaska

No. (fig.)	Lab. Sample No.	Terrace	Age	Collector	Year	Material dated	Comments
1	W-2190	I	3,340 $\pm$ 250	Plafker	1967	Peat	
2	W-2058	I	3,760	Plafker	1967	Peat	Same locality as sample no. 5
3	W-2185	I	4,100 $\pm$ 250	Plafker	1967	Wood	
4	W-2191	I	4,300 $\pm$ 250	Plafker	1967	Peat	
5	W-1405	I	4,470 $\pm$ 250	Plafker	1963	Wood	Driftwood from 1.3-ft bed of sand. Overlain by 2.5 ft clay and sand and 4 ft peat. Underlain by 2.1 ft beach gravel over bedrock surface
6	W-1728	II	940 $\pm$ 350	Plafker	1965	Peat	Peat from discontinuous patches in lag gravel 6 in. thick on bedrock surface. Overlain by 1-1/2 ft gravelly sand and 2 ft sandy peat
7	W-2062	II	3,590	Plafker	1967	Peat	From base of 5-ft-thick peat section
8	W-1732	III	550 $\pm$ 200	Plafker	1965	Peat	Peat from discontinuous patches in lag gravel 5 in. thick on bedrock surface. Overlain by 6 in. pebbly mud and 6 in. peat
9	W-1205	III	600 $\pm$ 250	Wilamovsky	1956	Peat	
10	W-2187	III	2,370 $\pm$ 250	Plafker	1967	Peat	
11	W-2060	III	2,480	Plafker	1967	Peat	From base of 3.5-ft-thick peat section overlying bedrock surface
12	W-2546	III	3,020 $\pm$ 250	Plafker	1970	Peat	
13	W-2548	III	3,260 $\pm$ 250	Plafker	1970	Peat	
14	W-1797	III	4,185 $\pm$ 250	Plafker	1965	Wood	Probably washed out of stage I terrace and redeposited. Driftwood from peaty and pebbly sand 20 in. thick on bedrock surface. Overlain by 20 in. peaty sand and muskeg
15	W-2059	IV	2,220	Plafker	1967	Wood	
16	W-1404	IV	2,390 $\pm$ 200	Plafker	1963	Wood	Driftwood at base of 4.3-ft-thick peat section overlying 1.5 ft of beach gravels on bedrock surface
17	W-1261	V	460 $\pm$ 250	Karlstrom	1962	Wood	Driftwood in lagoonal deposits
18	W-1259	V	660 $\pm$ 250	Karlstrom	1962	Wood	Driftwood in lagoonal deposits
19	W-1202	V	700 $\pm$ 250	Wilamovsky	1956	Peat	Peat in lagoonal deposits
20	W-2057	V	1,020	Plafker	1967	Wood	Driftwood from lagoonal deposits
21	W-1796	V	1,150 $\pm$ 500	Plafker	1965	Wood	Driftwood from top of 1 ft peaty lag gravel on bedrock surface. Overlain by 2 ft sandy peat
22	W-2064	V	1,250	Plafker	1967	Peat	
23	W-2063	V	1,380	Plafker	1967	Wood	
24	W-1724	V	1,350 $\pm$ 200	Plafker	1965	Wood	Driftwood from beach gravels 3.6 ft on bedrock surface. Overlain by 1-2 ft sandy peat and colluvium

Table 2.--Postseismic elevation changes at tide gages in the region affected by vertical tectonic movements during the March 27, 1964 Alaska earthquake. Locations of gages shown on figure 6.

Tide gage	1964 coseismic displacement (cm)	Postseismic Displacement (cm)	Postseismic Recovery (percent)	Years of record
Cordova	+189	-12	6	1964-74
Kodiak	-171	+58	34	do.
Seldovia	-119	+14	12	do.
Anchorage	- 79	+14	18	do.
Seward	-110	+15	14	do.

Figure 1.

Figure 2.

Figure 4.

Figure 5.

Figure 7.

Figure 8.

ANALYSIS OF SEISMIC MIGRATION

Albert T. Smith
Jacques Delsemme

Earth Sciences Board
University of California, Santa Cruz
Santa Cruz, Ca. 95064

ABSTRACT

Earthquake migration along linear seismic belts is investigated by analyzing space-time diagrams using spectral analysis. In order to sample the earthquakes in the space-time domain, they must first be convolved with a $(\sin x \sin t)/xt$ surface to obtain an unbiased and alias free two-dimensional Fourier spectrum. Further enhancements are provided by selectively stacking patterns (a pattern is defined as the distribution of earthquakes in space and time before a particular earthquake), thereby reinforcing the similarities within the various patterns. With these techniques, it is possible to quantitatively estimate the migration rates (from their spatial frequencies) and recurrence intervals (from their temporal frequencies) of large earthquakes in South America.

Preliminary examination of the spectra for South America indicates that a low frequency peak occurs at approximately 2500 km and 27 years for earthquakes with magnitudes greater than 7.7. The results suggest a migration rate of approximately 95 km/yr from south to north and a recurrence interval of 27 years.

INTRODUCTION

Seismologists have often searched for periodicities in the occurrence of earthquakes. These attempts have usually resulted in disappointment since the main characteristic of seismicity seems to be randomness. In most cases, part of the non-random component can be explained by aftershocks (Shlien and Toksacz, 1970). This is true when examining small regions (Knopoff, 1964), as well as the earth as a whole (Lomnitz, 1974). However, if the location of earthquakes is considered in addition to the time of occurrence, a different view emerges.

Several researchers have observed that large earthquakes migrate along seismic belts as time progresses (Mogi, 1968a,b; Kelleher and others, 1970, 1972, 1973, 1974, 1975, 1976). Mogi (1968a) observed migration for the Sanriku seismic activity between 1930 and 1935. In Anatolia, earthquakes migrate from east to west (Mogi, 1968a; Kelleher, 1970). In the Aleutians, two sequences are noted by Kelleher (1970). Earthquakes above magnitude 7.7 migrate from east to west at about 100 km/yr. In central California, earthquakes greater than magnitude 5.0 can be plotted on five linear trends with migration rates of 3.3 km/yr, where the average interval between these lines is about 7 years (Wood and Allen, 1973). This corresponds to the period observed for the cyclical variations of dilatational strain extending from San Francisco to San Bernardino (Greensfelder, 1972, 1973). The apparent correlation between dilatational strain and migrating trends in California would favor a deep-seated, migrating deformation front as an important factor in controlling the occurrence of earthquakes.

A similar mechanism has been suggested by Scholz (1977) for the 1975 Haichang earthquake. This would be the last major shock in a series of earthquakes triggered by a deformation front that moved through NE China at a velocity of 110 km/yr.

In these cases the description of migration has been

qualitative; however, a few investigators have attempted to quantify their results. Kagan and Knopoff (1976, 1977) computed the second order moment function of a catalog of large earthquakes for the years 1897 to 1972. For a point process, such as individual earthquakes, the first order moment is defined as the average number of seismic events in a given time and distance interval. The second order moment is "the number of earthquakes in different magnitude ranges that occur in a certain time-distance interval from [a given] shock" (Kagan and Knopoff, 1976). The second order moment corresponds to the correlation function in a continuous random process.

When Kagan and Knopoff considered all earthquakes (thereby not distinguishing between various regions), they observed several features for an earthquake of magnitude 8 or greater. Aftershocks are grouped around the main shock extending from 1 to 2 years and 250 to 1000 kilometers. Following the aftershocks, there is a decreasing number of earthquakes for about 5 years before the activity returns to its previous level. A seismic gap occurs about 8 years and 2000 kilometers prior to the main earthquake. The relationship between large earthquakes at long time-distance intervals is less clear, but may indicate a migration velocity of 250 to 500 km/yr.

Yang (1976) proposes a different method than Kagan and Knopoff to objectively distinguish between trends of earthquakes. In a two-dimensional plot of earthquakes where one axis is time and the other depth, he optimizes the clustering of events into a series of bands by rotating the bands and computing the angle that gives the highest contrast between bands. This angle defines a migration rate. The technique depends at least in part on the width of the band chosen to determine the contrast. It also assumes that earthquake migrations are linear. Instead of analyzing how the depths of earthquakes change with time, it is also possible to apply this method to the migration of earthquakes along seismic belts.

We attempt to quantify the relationship between earthquakes

in both time and space by directly computing the two-dimensional (space-time) fourier spectrum. Here we define space as the distance along seismic belts. In contrast to Yang's technique, spectral analysis directly gives the periodicities in a given space-time pattern; the only limitation is that the point process be stationary (Bartlett, 1964). For earthquakes it is likely that stationarity is satisfied since the driving force of plates is probably constant over short geological time periods, although it may deviate in recent history (Kagan and Knopoff, 1976; Kanamori, 1977). In addition, techniques from exploration seismology are adapted to better extract the non-random component contained within the space-time distribution of earthquakes.

METHOD OF ANALYSIS

Earthquakes tend to occur as belts along the surface of the earth. Usually these belts are roughly linear; consequently, a two-dimensional graph (space vs. time) lends itself to plotting the occurrence of an earthquake, both in time and space (Mogi, 1968a). To define the space axis on such a space-time graph, a line defined by segments of great circles is drawn through the belt of seismicity. If an earthquake is within a specified distance from the line, the earthquake is projected perpendicularly to the nearest great circle segment. Consequently, one coordinate, its distance along the belt, specifies the location of the earthquake. The other axis of the graph is the time axis.

Qualitative analysis of these space-time diagrams has been the traditional approach to determine seismic migration (e.g. Mogi, 1968a; Kelleher, 1970); however, the space-time distribution of earthquakes contains a strong random component making it difficult to recognize unbiased trends. In order to distinguish these trends with less ambiguity, a technique known as stacking has been adapted from exploration seismology.

In exploration seismology, seismic records of waves passing through the same region are summed or stacked to produce a single

seismic trace. This trace has a higher signal-to-noise ratio since the desired reflections are reinforced proportional to N , the number of traces, while the noise, being random, is added as the square root of N (Telford et al., 1976). Here instead of stacking seismic traces, we stack analogous space-time patterns.

In this study a pattern is defined as the distribution of earthquakes in space and time before a particular earthquake. Figure 1 is an example of patterns in a space-time graph. Common features in these patterns become more prominent when stacked, while random features are not reinforced. The number of stacked patterns within a diagram is the fold of the stack. Pattern size varies from earthquake to earthquake. The spatial axis of a pattern depends on the geometry of its seismic belt: Earthquakes toward the left end of the belt have patterns extending mostly toward the right and vice-versa. The pattern's time axis is limited by the list of earthquakes from which the data is taken. The pattern of the oldest earthquake in the list has a duration of zero; the pattern of the most recent earthquake in the list has a duration corresponding to the total time interval covered by the earthquake list. The remaining patterns will fall between the two extremes. Thus for more recent earthquakes, the observable patterns are longer. If we examine patterns that are 20 years long, only earthquakes occurring 20 years after the beginning of the list will have complete patterns. This limits which earthquakes may be stacked, since only those earthquakes with complete patterns may be stacked in order to avoid a bias in favor of earthquakes occurring just prior to the stacked earthquake.

The epicenter rather than the rupture zone represents the space-time location of each earthquake. The rupture zone would indeed be required if the epicenter represented only a place of structural weakness originating in heterogeneities along the fault; however, the epicenter is usually not randomly located in the rupture zone but occurs near the preceding earthquake (Kelleher, 1970). Since the epicenter is influenced by the strain field, we assume a point process as the initial location rather than a line

source.

Spectral analysis characterizes any phenomena in terms of periodicities. If the data is assumed to be periodic, the spectrum is a discrete distribution containing a finite number of frequencies. If the data is not periodic, the spectrum is continuous, each computed frequency actually representing a band of frequencies (Rayner, 1971). The latter case is more common, and earthquakes certainly fall into this category. If a spectrum has a fairly uniform distribution, it suggests a random process; however, significant deviations from a uniform distribution indicates that other processes are at work which may be distinguishable.

Earthquakes, however, present special problems. The space-time plane cannot be treated as a continuous function because it is a point process, each point defining the time and epicenter of an earthquake. Each event is a spike with essentially instantaneous rise time. A spike contains a uniform frequency distribution; however, a bandlimited signal is needed to avoid aliasing.

To avoid these difficulties, past researchers computed the spectrum of a related variable, rather than the spectrum of the point process itself. In this manner the process could be described as continuous and bandlimited. For example, the process could be considered a series of interspike intervals, a cumulative counting process, or an instantaneous frequency function. Unfortunately, each alternative has drawbacks severe enough to limit its usefulness.

These problems were elegantly solved by French and Holden (1971). Their method involves several steps which are to be simultaneously executed. Although they describe the solution for a one-dimensional case, their method can be extended to more than one dimension. In two-dimensions, the steps are:

(1) Truncate the data to a finite record. (2) Convolve each earthquake or spike with amplitude A with a function that is an ideal low-pass filter (Brigham, 1974):

$$A^2 \cdot \frac{\sin(\pi t / dt)}{\pi t} \cdot \frac{\sin(\pi x / dx)}{\pi x}$$

where t denotes time and x represents space with respective Nyquist frequencies of $(1 / 2 \, dt)$ and $(1 / 2 \, dx)$. (3) Truncate the data again with the same window used in step (1). (4) Sample the data with sampling rates of $(1 / dt)$, $(1 / dx)$. (5) Remove the mean to prevent the appearance of a DC spike in the final spectrum. In addition to these steps, the space-time diagrams are extended in all directions with zeroes in order to increase the resolution of the final spectra. This is done prior to step (2) to avoid truncation of the ideal low-pass filter and, thereby, minimize bias. A 25% split cosine bell taper is applied at the edges of the sampling grid, a region extending 64 by 64 points. The sampling intervals are approximately every 2.4 years in time and every 212 km in space. The two-dimensional Fast Fourier Transform is computed, and the sum of the square of its coefficients forms the raw power spectrum or the spectrum of the variance. The spectra are blanked near the origin, since these values represent periods longer than the length of the data. To better visualize the true spectrum, the raw spectrum is smoothed using a nine-term running average. This narrows the confidence bands around each value but decreases the resolution.

APPLICATIONS TO SEISMIC MIGRATION

Our concern is the recurrence interval and migration of major earthquakes greater than magnitude 7.7; thus, aftershocks are eliminated from the analysis (Kelleher, 1970). Duda's (1965) earthquake catalogue provides the source of data for earthquakes occurring from 1897 to 1964 with additional extensions to the end of 1974 (U.S. Earthquakes, 1965 to 1974). Because the Richter magnitude is no longer representative of the energy released (Kanamori, 1977, 1978), all earthquakes are considered to be equal.

South America

The spatial axis extends from the triple junction between the Cocos, Nazca, and South America plates along the western coast of South America to the triple junction between the Nazca, Antarctica, and South America plates. There are 30 earthquakes with magnitude above 7.7 included in the analysis. Figure 1a shows the space-time graph and Figure 1c shows the space-time stack.

The space-time diagram shown in Figure 1a suggests migration trends from south to north; however, the pattern is not obvious. A rapid trend could also be picked starting in the north near about 1900 and migrating south. If the space-time stack in Figure 1c is now examined, common features prior to each earthquake are reinforced in the pattern. Strong south to north migrations are now observed as a series of parallel bands. These bands suggest a recurrence interval of approximately 27 years, while migrating approximately 2500 km to the north. No clustering which is common to all the earthquakes is observed, suggesting that earthquakes do not occur at very regular intervals, even along the migrating trend.

The two-dimensional power spectrum reinforces this interpretation in both the stacked and unstacked space-time plots. The spectrum of the space-time plot (Fig. 2a) contains a single low frequency peak at 27 years and 2550 kilometers, implying a south to north migration. Although the raw spectrum is noisy, the same features can be seen when it is smoothed (Fig. 2b). They appear still more clearly in the raw (Fig. 3a) and smoothed (Fig. 3b) stacked versions. The values given above correspond to the intercepts of a linear band of earthquakes in the space-time stack (Fig. 1c). The intercept in time represents the recurrence interval (about 27 years) and the intercept in space represents the migration interval (about 2500 kilometers).

If the 7 deep earthquakes (with depths greater than 70 km) are removed from the analysis, the features observed within the spectra change very little except for a decrease in the signal-to-

noise ratio. This decrease could result from fewer earthquakes in the analysis, or because deep earthquakes actually play a role in the observed migration. Although deep earthquakes are not directly related to the interaction between the Nazca and South America plates, they most certainly contribute to the strain release at the surface. Such a rupture at depth could therefore be part of a sequence of earthquakes and reinforce the observed migration pattern.

The direction of propagation from south to north is opposite that observed by Kelleher (1972) for earthquakes south of 32°S . Since his space axis encompasses only a small (about one fifth) portion of the one used in our study, it is difficult to conclude whether there is a basic conflict between them.

MODELS

To assist in the interpretation of the space-time distribution of earthquakes and its associated spectra, simple artificially generated patterns are created. Earthquake epicenters are placed on a grid-like pattern consisting of parallelograms with a base of 2500 km, a height of 9 years, and offset of 830 km. The pattern then repeats itself every 27 years and every 2500 km. These figures correspond to those determined for South America. It is possible to predict where the spectral peaks should appear, even without actually computing the two-dimensional spectrum (see Papoulis, 1965, 1968). Like the epicenters, the spectral peaks fall on a parallelogram grid: The peaks closest to the origin indicate the fundamental periods, while the others correspond to their higher harmonics.

The epicenters are then randomized about their initial positions. This is accomplished by assuming that the "earthquake" deviate from their initial positions and form a Gaussian population whose standard deviations can be specified in both time and space. The deviations form an ellipse whose orientation can be specified.

First let us consider deviations with the axes parallel to

the time and space axes. A large temporal standard deviation effectively destroys the higher temporal harmonics in the frequency domain. The larger the deviation, the lower the frequency cut-off point. The same principle applies for the spatial frequencies. When both are used together, only the peaks closest to the origin remain, but no satisfactory model was produced with deviations parallel to the space-time axes.

By inclining the standard deviation ellipse with respect to the time and space axes, it becomes possible to orient it parallel to the observed migration trend. Only then is this simple model able to duplicate several features found among South American earthquakes. Figures 4 to 6 show a representative model. The standard deviations are 2.0 years and 225 km, rotated counterclockwise to match the observed migration trend. The rotation is analogous to factor analysis where the migration trend represents one independent direction or factor. The space-time plot of the model is shown in Figure 4a, and its space-time stack is shown in Figure 4b. Note that the latter has similar features as the space-time stack for the South American earthquakes shown in Figure 1c. The raw and smoothed spectra of the model are shown in Figure 5 for the space-time plot and in Figure 6 for the space-time stack. When compared to the real spectra of South America, Figures 2 and 3, the gross features are similar: the peak occurring at 27 years and 2500 km is prominent while other peaks tend to be suppressed.

CONCLUSIONS

The sampling procedure permits the analysis of any discontinuous point process, such as earthquakes, with standard techniques usually restricted to continuous process. We have presented here one such analysis: the two-dimensional spectral analysis of the space-time distribution of earthquakes in South America. It gives new insights into the basic process underlying the occurrence of large earthquakes. The simple model explains several

features exhibited by the distribution of earthquakes in South America. First, migration occurs at a rate of approximately 95 km/yr from south to north for earthquakes with Richter magnitude greater than 7.7. The recurrence interval for major earthquakes has a dominate period of 27 years using our limited historical record. The artificially generated model suggests, however, that the deviations in the migration rate may be large for individual earthquakes. A more complete catalogue of earthquakes may alter the conclusions of this preliminary analysis, but the spectral analysis performed here represents a step towards a quantitative analysis of seismic migration.

ACKNOWLEDGEMENTS

This research was supported by the Earthquakes Hazards Reduction Program of the United States Geological Survey, Grant No. 14-08-0001-G-399.

- M. S. Bartlett, The spectral analysis of two-dimensional point processes, *Biometrika* 51 (1964), 299-311.
- E. O. Brigham, *The Fast Fourier Transform* (Prentice-Hall, Englewood Cliffs 1974).
- S. J. Duda, Secular seismic energy release in the circum-Pacific belt, *Tectonophysics*, 2 (1965), 409-452.
- A. S. French and A. V. Holden, Alias-free sampling of Neuronal spike trains, *Kybernetik*, 8, 5 (1971), 165-171.
- R. W. Greensfelder, Crustal movement investigations in California: their history, data, and significance, *Calif. Div. Mines Geol., Special Publ. 37* (1972).
- R. W. Greensfelder and J. H. Bennett, Characteristics of strain variation along the San Andreas fault from geodimeter measurements, *Stanford Univ. Publ., Geol. Sci., XIII* (1973), 54-63.
- Y. Kagan and L. Knopoff, Statistical search for non-random features of the seismicity of strong earthquakes, *Phys. Earth Planet. Int.*, 12 (1976), 291-318.
- Y. Kagan and L. Knopoff, Earthquake risk prediction as a stochastic process, *Phys. Earth Planet. Int.*, 14 (1977), 97-108.
- H. Kanamori, The energy release in great earthquakes, *J. Geophys. Res.*, 82, 20 (1977), 2981-2988.
- H. Kanamori, Quantification of earthquakes, *Nature*, 271 (1978), 411-414.
- J. A. Kelleher, Space-time seismicity of the Alaska-Aleutian seismic zone, *J. Geophys. Res.*, 75, 29 (1970), 5745-5756.
- J. A. Kelleher, Rupture zones of large South American earthquakes and some predictions, *J. Geophys. Res.*, 77 (1972), 2087-2103.
- J. Kelleher, L. Sykes, and J. Oliver, Possible criteria for predicting earthquake locations and their application to major plate boundaries of the Pacific and the Caribbean, *J. Geophys. Res.*, 78, 14 (1973), 2547-2585.

- J. Kelleher, J. Savino, H. Rowlett, and W. McCann, Why and where great earthquakes occur along island arcs, *J. Geophys. Res.*, 79, 32 (1974), 4889-4899.
- J. Kelleher and J. Savino, Distribution of seismicity before large strike slip and thrust-type earthquakes, *J. Geophys. Res.*, 80, 2 (1975), 260-271.
- J. Kelleher and W. McCann, Buoyant zones, great earthquakes, and unstable boundaries of subduction, *J. Geophys. Res.*, 81, 26 (1976), 4885-4896.
- L. Knopoff, The statistics of earthquakes in Southern California, *Bull. Seismol. Soc. Am.*, 54 (1964), 1871-1873.
- C. Lomnitz, Global tectonics and earthquake risk (Elsevier Scientific Publishing Co., 1974).
- K. Mogi, Migration of seismic activity, *Bull. Earthquake Res. Inst.*, 46 (1968a), 53-74.
- K. Mogi, Sequential occurrences of recent great earthquakes, *J. Phys. Earth*, 16, 1 (1968b), 30-36.
- A. Papoulis, Probability, random variables and stochastic processes (McGraw-Hill Book Co., New York 1965).
- A. Papoulis, Systems and transforms with applications in optics (McGraw-Hill Book Co., New York 1968).
- J. N. Rayner, An introduction to spectral analysis (Pion Limited, London 1971).
- C. H. Scholz, A physical interpretation of the Haicheng earthquake prediction, *Nature* 267 (1977), 121-124.
- S. Schlien and M. N. Toksoz, A clustering model for earthquake occurrences, *Bull. Seism. Soc. Am.*, 60, 6 (1970), 1765-1787.
- W. M. Telford, L. P. Geldart, R. E. Sheriff, and D. A. Keys, Applied geophysics (Cambridge University Press, Cambridge 1976).
- U. S. Earthquakes, Principal earthquakes of the world, U. S. Government Printing Office, Washington, D.C. 20402 (1965 to 1974).

M. D. Wood and S. S. Allen, Recurrence of seismic migrations along the central California segment of the San Andreas fault system, *Nature*, 244 (1973), 213-215.

M. Yang, The effect of viscoelastic medium and earthquakes on plate motion, *Personal Communication* (1976).

TABLE I

Coordinates of great circle segments defining the spatial axes. North and east are positive, south and west are negative. Values are expressed in degrees.

REGION	LATITUDE	LONGITUDE
South America	6.0	-75.0
	-2.0	-79.0
	-10.0	-78.0
	-18.0	-69.0
	-24.0	-69.0
	-45.0	-76.0
	-49.0	-73.0

FIGURE CAPTIONS

Figure 1a: Space-time plot of earthquakes with magnitudes greater than 7.7 for western South America. The space axis goes from north to south; the origin is located at 6°N , -75°W .

Triangles are shallow earthquakes (less than 70 km), crosses are intermediate earthquakes (from 70 to 300 km), squares are deep earthquakes (deeper than 300 km).

Figure 1b: Space-time plot of earthquakes with magnitude above 7.7 for western South America as in Figure 1a but with stacked coordinate frame for each earthquake. The smaller rectangular boxes represent patterns 20 years long and 830 km wide. Only patterns complete in time are shown. The origin of each box is located at an earthquake; the relative positions of preceding earthquakes are plotted in the new coordinate frame. Later, the relative coordinates or boxes are stacked: the relative positions of the earthquakes are superimposed on one another. The actual stacking involves larger boxes (see text).

Figure 1c: Space-time stack of earthquakes with magnitudes above 7.7 for western South America (fold = 15). The space axis goes from north to south. The coordinates are relative to the position of the stacked earthquake.

Figure 2a: Raw power spectrum of space-time graph for South America in Figure 1a.

For all power spectra, values around the origin have been blanked since the periods they represent are longer than the length of the data itself. This occurs because the space and time axes were extended with zeroes to increase the resolution in the two-dimensional spectra.

Figure 2b: Smoothed power spectrum of space-time graph for South America in Figures 1a and 2a. The smoothing was done with a 9-term running average.

Figure 3a: Raw power spectrum of stacked space-time graph for South America in Figure 1c.

Figure 4a: Space-time plot of model. The parameters are: parallelograms with a base of 2500 km, a height of 9 years, and an offset of 830 km; the temporal and spatial standard deviations are 2 years and 225 km respectively, rotated counter-clockwise by $\arctan(1670 \text{ km}/9 \text{ years})$. Compare with Figure 1a, the space-time plot for South America.

Figure 4b: Space-time stack (fold = 13) of artificial model. The stack has a similar appearance to Figure 1c, the space-time stack of South America.

Figure 5a: Raw power spectrum of the space-time plot of the artificial model in Figure 4a. A dominate peak occurs at periods of approximately 27 years and 2500 km (approximately 0.04 yr^{-1} and $0.4 \times 10^{-3} \text{ km}^{-1}$), a similar location to the dominate peak in the power spectrum for South America in Figure 2a.

Figure 5b: Smoothed power spectrum of the space-time plot for the artificial model. Regularity generated by higher harmonics of the parallelogram structure is more evident in the artificial model than South America in Figure 2b.

Figure 6a: Raw power spectrum of space-time stack of model. The results are similar to Figure 3a, the corresponding diagram for South America.

Figure 6b: Smoothed power spectrum of space-time stack of artificial model. Again the structure is very similar to Figure 3b for South America.

Figure 1a

Figure 1b

SLAVE EMIT

SPACE (x1000 km)

-40
-3.33

0

-40
+3.33

Figure 2a

Figure 3a

Figure 5a

Figure 5b

Figure 6a

Figure 6b

A Model for the Earthquake Cycle in Underthrust Zones

Wayne Thatcher*
U.S. Geological Survey
Menlo Park, California 94025

John B. Rundle
Sandia Laboratories
Albuquerque, New Mexico 87115

Short Title: Crustal Deformation in Underthrust Zones

Submitted to Journal of Geophysical Research August 1978

*Address from September 1978 to March 1979:
Earthquake Research Institute
Tokyo University
Bunkyo-ku
Tokyo, Japan (113)

ABSTRACT

The time-dependent surface deformation due to thrust faulting in an elastic plate overlying a viscoelastic half-space is examined and compared with geodetic measurements of earthquake-related crustal movements from Japan. The fault is two-dimensional and the half-space rheology is that of a Maxwell solid, whose instantaneous response to applied loads is purely elastic, but which subsequently flows to relieve imposed shear stresses. A characteristic feature of shallow angle underthrusting shown by these computations is that buried slip produces predominantly land uplift, while asthenospheric relaxation due to surface faulting results principally in surface downwarping. These distinctive patterns, also reflected in the measured deformation, are particularly useful in formulating a viable model for the entire earthquake cycle. For the model constructed in this way, coseismic faulting in the upper part of the plate transfers a load to the lower lithosphere and asthenosphere, inducing postseismic slip downdip of the seismic rupture and relaxation in the asthenosphere, these transients eventually merging into the steady buried slip and smoother asthenospheric flow that characterizes the interseismic phase of the deformation cycle. Aseismic slip and asthenospheric relaxation cause the load supported by the plate-bounding fault to be gradually transferred back to the shallow locked segment of the fault, effecting the strain build-up for a subsequent earthquake, and the cycle is repeated. Energy for the process is ultimately supplied by the convergence of the plates, manifested in this kinematic model by the steady buried slip that persists throughout the cycle.

Characteristic features of the observed deformation in southwest Japan, site of the 1946 Nankaido earthquake, and in the South Kanto district, where the great 1923 earthquake occurred, are matched by this model using conventional values of lithospheric thickness (60 km) and asthenospheric viscosity (10^{21} poise, or 10^{20} N s/m²), although the model is not strongly tied to these precise values. Postseismic movements are adequately explained by episodic slip that occurs below the coseismic rupture, is 10 to 30% of the seismic slip, and acts in the same sense. Superposed on these movements are the lesser effects of asthenospheric relaxation, which in the South Kanto district contribute significantly to the postseismic vertical level changes because post-1923 buried slip is comparatively small and involves largely strike-slip movements. Interseismic subsidence in both regions is explained well by asthenospheric relaxation and less significant deformation effects due to steady aseismic slip on the lower part of the plate boundary.

1. INTRODUCTION

The plate tectonic view that the earth's surface is comprised of a mosaic of strong elastic plates that overlie a relatively weaker viscous asthenosphere provides a conceptual framework that successfully accounts for a varied and diverse assemblage of geologic and geophysical observations. Several important implications of this model for the crustal deformation occurring at plate boundaries are now being recognized, and a number of recent studies have suggested that the geodetically-measured deformation observed in

active seismic zones may be due to time-dependent flow in the asthenosphere induced by the sudden stress drop resulting from major earthquakes in the overlying lithosphere (Nur and Mavko, 1974; Smith, 1974; Rundle and Jackson, 1977; Spence and Turcotte, 1978; Savage and Prescott, 1978b). On the other hand, many features of the deformation can be adequately explained with purely elastic models in which both the time-dependent postseismic movements and the relatively steady interseismic secular deformation are attributed to episodic or steady aseismic slip on the downward extension of the fault plane that ruptured coseismically (Savage and Burford, 1970, 1973; Shimazaki, 1974; Thatcher, 1975; Kasahara, 1975; Brown et al., 1977; Scholz and Kato, 1978). These results suggest that both asthenospheric relaxation and buried aseismic slip are plausible mechanisms for plate boundary deformation. Recent work (e.g. Rundle and Jackson, 1977; Spence and Turcotte, 1978; Savage and Prescott, 1978b) demonstrates that for a strike-slip transform fault in the lithosphere, the effects of buried slip and asthenospheric relaxation are very similar, satisfy available data about equally well, and may be observationally indistinguishable. However, as we shall show here, the situation differs in the case of dip-slip faulting.

In accounting for the geodetically-observed surface deformation, the importance of relaxation processes in the asthenosphere relative to the movement of dislocations in the lithosphere depends significantly on assumed values for lithospheric thickness and asthenospheric viscosity. First of all, the surface deformation due to asthenospheric relaxation depends on both the downdip width of the coseismic fault plane and the effective elastic thickness of the lithosphere. Unfortunately, although the width and maximum depth of

rupture are reasonably constrained, there is no general agreement on the value of lithospheric thickness appropriate for modeling post-slip asthenospheric relaxation. Geodetic observations constrain coseismic rupture depths and indicate they are significantly greater in underthrust zones than at strike-slip plate boundaries: ~ 10 km depth on the San Andreas Fault in California (Kasahara, 1957; Chinnery, 1961; Thatcher, 1975) and 20–50 km deep at circum-Pacific subduction boundaries (Hastie and Savage, 1969; Plafker and Savage, 1970; Ando, 1971, 1975). Although the lithospheric thicknesses inferred from seismic velocity measurements in these regions generally range from 70 to 120 km, studies of lithospheric flexure (Hanks, 1971; Watts and Talwani, 1974; McNutt and Menard, 1978) suggest elastic thicknesses may be only 15–40 km. For the timescales of earthquake-related aseismic deformation, which vary from a few years to a few hundred years, neither estimate may be appropriate: the seismic estimates apply to wave periods of ~ 1 –100 seconds, and at these periods the asthenosphere is nearly perfectly elastic; on the other hand, the flexure calculations are relevant to loading intervals of 10^5 years or longer, and for these models the viscous effects of the asthenosphere are assumed to be negligible and the lithosphere is taken to be a free plate.

As several studies have emphasized (Nur and Mavko, 1974; Smith, 1974; Rundle and Jackson, 1977), in order to explain the relatively short timescale of the immediate postseismic effects of major crustal earthquakes with asthenospheric relaxation requires mantle viscosities that are rather low, more than an order of magnitude less than those estimated by other methods, even in tectonically active regions. For example, from the isostatic rebound

of Lake Bonneville in the Basin and Range Province of the western United States, Cathles (1975) estimates a lower bound of 5×10^{20} poise (5×10^{19} N s/m²) for the uppermost mantle, a viscosity that in turn is about a factor of ten less than viscosities obtained from postglacial uplift of stable shield areas. Postseismic relaxation times reported following several major Japanese earthquakes are 5 years or less (Fitch and Scholz, 1971; Scholz and Kato, 1978), and to explain such short timescales Nur and Mavko (1974) and Smith (1974) have suggested that viscosities of $\sim 5 \times 10^{19}$ poise (5×10^{18} N s/m²) or less are needed. Nonetheless, these viscosities are not grossly unreasonable and certainly are not unassailable grounds for rejecting the asthenospheric relaxation mechanism out-of-hand.

The work reported here focuses on the surface deformation effects of thrust faulting in an elastic plate overlying a viscoelastic (Maxwell) half-space. Put simply, the results obtained here show that for surface rupture of a shallowly inclined thrust fault, the predominant effect of asthenospheric relaxation is to produce downwarping at the earth's surface, while for buried aseismic slip, the principal surface deformation is crustal uplift. Several features of the measured deformation show correspondingly distinctive patterns, and characteristic features of the co-, post-, and inter-seismic deformation observed in underthrust zones often allow these three movement phases to be clearly distinguished. We refer to these phases collectively as the earthquake cycle, and make the case for the viability of our model by comparing the predicted effects of asthenospheric relaxation and fault slip with geodetic measurements from Japan.

This paper is organized into nine sections. Following the introduction, section 2 considers the time-dependent surface deformation for a faulting model in which the effects of a viscoelastic asthenosphere are included. In section 3 these results are used to construct a model for the complete deformation cycle, and the relation of this model to the broader scale processes of plate tectonics is discussed. Sections 4 through 7 justify the model by comparing its predictions with extensive observations from central and southwest Japan. Section 8 discusses and compares the viscoelastic modeling approach with a simpler formulation that considers only the purely elastic effects of faulting in a uniform half-space. Finally, section 9 is a concluding summary and discussion. We suggest that the more casual readers of this paper skip the rather lengthy discussions of sections 4 to 7 and study only the figures and figure captions related to this portion of the paper (Figures 4-14), which can be followed largely independent of the text.

2. MODELS WITH ASTHENOSPHERIC RELAXATION

The instantaneous elastic effects and time-dependent viscoelastic deformation due to surface faulting and to buried fault slip are compared in Figure 1. The results shown here use the formulation developed by Rundle (1978) for three-dimensional dislocations in layered elastic and viscoelastic media. Simplifications that result for two-dimensional faults and extension of Rundle's method to ramp time functions for fault slip are given in Appendices A and B. In each case the fault is infinitely long and dips at an angle of 30° , and the slip occurs abruptly as a step function in time at $t=0$.

Vertical displacement at the free surface is normalized by the reverse slip on the fault, and the distance perpendicular to the fault is in multiples of the lithospheric thickness. The assumed earth model consists of an elastic plate of thickness H that overlies a viscoelastic half-space with the properties of a Maxwell solid. With this constitutive law for the half-space, its immediate response to applied stresses is purely elastic, while its long term deformation is that of a Newtonian viscous fluid; a one-dimensional analogue to this material behaviour is an elastic spring and a viscous dashpot connected in series (for further details see Fung, 1965, pp. 22-25). The half-space has viscosity η , and its shear modulus μ is taken equal to that of the elastic plate. The viscoelastic post-slip response plotted in Figure 1 is for $t=2\tau$, where τ is the characteristic relaxation time of the Maxwell half-space ($\tau=2\eta/\mu$); elastic uplift ($t=0$ response, solid lines) is not included in the dashed curves for $t=2\tau$. Since the viscoelastic substrate has immediate elastic properties that are identical to those of the overlying plate, the $t=0$ deformation should be the same as that for faulting in an elastic half-space; that this in fact is the case is one check on the correctness of our calculations.

Figure 1a shows the coseismic and subsequent viscoelastic deformation for a fault that ruptures the free surface and extends through three-quarters of the lithosphere ($d/H=0.75$). The most important feature illustrated by the viscoelastic deformation profile (dashed curve) is that the effect of the asthenosphere is to produce post-slip downwarping over a broad region, even in portions of the profile that subsided coseismically (minor amounts of uplift occur on the hanging wall (right hand) block of the fault beyond the segment

of the profile plotted in Figure 1a). Physically, the reason for this subsidence is clear: shallow dip underthrusting downwarps the lithosphere-asthenosphere boundary asymmetrically from large negative values of y/H to $y/H \approx 1.5$, and displaced it upwards by lesser amounts for larger positive y/H ; then, relaxation of these coseismically-imposed stresses in the asthenosphere progressively increases the downwarping of the boundary for $y/H \gtrsim 1.5$ and enhances its upward displacement beyond this point, with surface displacements reflecting this pattern of asthenospheric flow. For a vertical dip-slip fault the post-slip vertical displacement is perfectly anti-symmetric about $y=0$; successively shallower dips then concentrate the deformation on the hanging wall block of the fault, accentuating downwarping near the fault and spreading out the compensating uplift over a wider region further away from the surface trace.

Figure 1b shows the elastic ($t=0$) and viscoelastic ($t=2\tau$) vertical displacement due to buried slip on the downdip extension of the fault plane assumed in Figure 1a. This slippage then ruptures the bottom one-quarter of the lithospheric plate. A comparison of the dashed curve in Figure 1a with both profiles in Figure 1b demonstrates what is perhaps the most significant result of our modeling experiments, that both the immediate elastic effects and the gradual viscoelastic deformations resulting from buried slip differ in a first-order way from the viscoelastic response to surface faulting: the predominant effect of buried slip is to produce uplift of the free surface above the dislocation and some depression beyond it, while the asthenospheric relaxation due to surface faulting results in almost universal downwarping throughout the region of the coseismic deformation. A possibly significant

exception to this general rule is for the case of surface faulting that ruptures the entire lithosphere, and we shall discuss this situation later.

Now, however, we return to the surface faulting model of Figure 1a and examine the spatial pattern and time dependence of the viscoelastic surface deformation in more detail. Figure 2a is a plot of the cumulative post-slip viscoelastic displacements for successive times from 0.1τ to 2.0τ . The figure shows that although the cumulative displacements increase monotonically with time, the general shape of the profile remains unchanged, i.e. there is little evidence for any lateral migration of the deformation with time away from the region of sudden fault slip at $t=0$. As Savage and Prescott (1978a) have pointed out, the absence of such a strain migration effect is due to the instantaneous elasticity of the Maxwell half-space: stresses imposed on the asthenosphere coseismically relax viscoelastically after $t=0$, whereas with a purely viscous rheology only the plate responds elastically, and the subsequent adjustment of the asthenosphere produces the wave-like migration effects discussed by Elsasser (1969), Bott and Dean (1973), Anderson (1975) and Melosh (1976). Since, for example, the earth's mantle efficiently transmits transverse seismic waves, we agree with Savage and Prescott (1978a) that for modeling earthquake-induced effects in the asthenosphere a Maxwell rheology is more nearly correct than is a purely viscous constitutive law.

The time-dependence of vertical movements at selected surface locations is shown in Figure 2b. The most striking feature of these curves is that in the region of large post-slip deformation (roughly from $y/H = -0.5 \rightarrow +2.0$) movements persist for many asthenospheric relaxation times and do not, for example, simply decay exponentially with characteristic time τ . It can be

seen from curves plotted in Spence and Turcotte (1978) that the same sort of long-lived deformation occurs for the corresponding strike-slip problem, verifying that this behavior is a general property of faulting models that include an elastic plate coupled to a viscoelastic substrate. In retrospect at least, this feature of the deformation is physically plausible: although the tendency of the asthenosphere is to damp out impulsively applied loads exponentially, elastic restoring forces in the plate will resist its flexure due to tractions applied at the lithosphere-asthenospheric boundary, retarding the flow of the asthenosphere and smoothing out the time-dependence of the movements. Note also from Figure 2b that the surface displacements show a time dependence that varies with position, so that if each curve were fit empirically to an exponential decay, inferred decay times would not only fail to reflect the true relaxation time of the asthenosphere, but would also vary significantly with position.

Figure 3 is similar to Figure 2 but shows the space and time dependence of the viscoelastic deformation that results from rupture of the entire lithosphere. The plate-wide thrust fault dips at 30° , and hence this model is simply a superposition of the models of Figures 1a and 1b. Comparison of Figure 3 with Figures 1a and 2 reveals some significant differences, the most striking of which is that for the plate-wide rupture there is substantial post-slip uplift centered near $y/H=2$ (Figure 3a), with this uplift continuing to grow steadily for long times after $t=0$ ($y/H = 2.0$ curve, Figure 3b). In addition, the post-slip deformation extends somewhat farther into the plate, and the time-dependence (Figure 3b) is more diverse and shows greater spatial variation than does the model in which only part of the plate is faulted coseismically.

The results shown in Figure 1a, both for the coseismic and post-slip deformation, differ substantially from those obtained previously by Nur and Mavko (1974). The model parameters used in Figure 1a are identical to those chosen by them with the exception of the assumed asthenospheric rheology, which we have taken to be the constitutive law for a Maxwell solid but which Nur and Mavko assume to be a standard linear solid. However, this single difference cannot explain the discrepancies, since computations we have carried out for a standard linear solid asthenosphere are similar to the results shown in Figure 1a but yet remain quite different from Nur and Mavko's results. Two lines of evidence support the correctness of our calculations. First, our coseismic uplift profile agrees, as it should, with that computed for faulting in an elastic half-space, while Nur and Mavko's does not (refer to their Figures 1b and 2a for the thrust faulting case). Furthermore, the elastic solution used by Nur and Mavko as the starting point in their analysis is an approximate one, while our results, applying Rundle's (1978) technique, begin with the exact elastic solution. We suspect that this latter difference accounts for the discrepancies between our results and theirs. Our exact formulation and the agreement it produces with the expected $t=0$ elastic half-space response gives us confidence that our results are indeed correct and that we have not merely generated a different but nonetheless still incorrect solution to the problem. Further confirmation of the correctness of our results comes by comparing our work with Smith's (1974), obtained using a finite element approach, and with a recent independent calculation made by G. M. Mavko (personal communication, 1978); although model parameters used by these investigators differ slightly from our own, their results agree with the

basic features of the asthenospheric relaxation due to reverse dip-slip faulting that we show here and differ significantly from the model calculations originally presented by Nur and Mavko.

3. THE EARTHQUAKE DEFORMATION CYCLE

The results of the previous section suggest a simple model for the seismic cycle at convergent plate boundaries, and the principal features of this model can be understood by again referring to Figure 1. Consider first the coseismic strain release, occurring as sudden slip on the shallower portions of the plate boundary megathrust and producing the vertical displacements at the earth's surface given by the solid curve in Figure 1a. This shallow stress release then transfers a shear load to the deeper portions of the plate and the asthenosphere, and the asthenosphere will flow to relax these stresses while episodic aseismic slip occurs downdip of the coseismic fault plane.

These movements at depth initiate the postseismic phase of the deformation, relieving strains imposed by the earthquake but at the same time transferring part of the shear load supported by the plate boundary fault back to the shallow portions of the lithosphere, beginning the strain accumulation process anew. During the postseismic interval, the surface deformation profile will be some combination of asthenospheric relaxation due to surface faulting (dashed curve, Figure 1a) and the elastic and viscoelastic responses to buried slip (solid and dashed curves, Figure 1b), the exact proportions of each depending on the d/H ratio, the amount of coseismic slip, the asthenospheric relaxation time, and the amount and extent of episodic postearthquake slip.

Once the more rapid effects of asthenospheric adjustment and episodic slip have died away, the postseismic movements should merge into the pattern of interseismic secular deformation, with surface straining occurring at approximately uniform rates that reflect both the nearly constant rate of asthenospheric relaxation and the more or less steady deep aseismic slip that is driven by the convergence of the plates. Once again, the surface displacements during this phase of the movement cycle will be a superposition of the dashed profile of Figure 1a and the two curves plotted in Figure 1b. Both processes increase shear stresses in the shallow lithosphere near the plate boundary, and once the static strength of the brittle coseismic fault segment is exceeded, a great earthquake occurs and the cycle begins once more.

In this model of the seismic cycle, the strain energy for the earthquake occurring at the lithospheric boundary is supplied by the motion of the plates, and their rate of convergence also largely determines the recurrence interval between major shocks. The plate movement is reflected in the steady aseismic slip that we postulate to occur beneath the shallow locked portion of the underthrust zone, and the rate of slip here is determined by the rate of relative plate motion. This steady buried slip continually loads the shallow locked seismic zone, and in this way the plate motion rate is related to the rate of strain accumulation and hence the recurrence interval. In addition, postearthquake flow in the asthenosphere also contributes to the reloading, transferring stresses that were imposed coseismically back into the overlying lithosphere at a rate that depends on the asthenospheric viscosity.

Viewed on a plate tectonic scale, the plate boundary deformation cycle we have formulated here is a very local process that can be understood

kinematically without actually knowing the forces that drive the plates and ultimately supply the energy to maintain the cycle. Whatever these forces may be, in our model their net effect on plate boundary deformation is due solely to the steady aseismic buried slip that persists throughout the earthquake cycle, concentrating strains within a few lithospheric thicknesses of the plate boundary (viz. Figure 1b). Similarly, the effects of asthenospheric relaxation are concentrated near the underthrust zone, and because of the finite viscosity of the asthenosphere, on the timescale of earthquake recurrence the plate boundary is isolated from its interior. Thus, the processes occurring at the plate boundary may be understood as an essentially local interaction between seismic and aseismic slip on the megathrust and the viscoelastic reaction of the asthenosphere to these loads.

In comparing the modeling results of Figures 1, 2, and 3 with observed deformation in underthrust zones, two details not included in our calculations deserve brief mention. First of all, our results consider the asthenospheric relaxation effects of only a single earthquake rather than the superposition of a periodic or irregular sequence of events. Thus our "one-shot" model will always tend to somewhat underestimate the interseismic subsidence due to asthenospheric relaxation, and the error incurred will depend on the ratio of recurrence interval, T , to asthenospheric relaxation time, τ , the approximation being best when T/τ and t/τ are small, and worst when these two ratios are large. In fitting data, the net effect of this approximation will be to slightly underestimate the true asthenospheric viscosity, although for the modeling done here the error is not large.

A second effect of the interseismic deformation that we have not modeled exactly is the response of the asthenosphere to steady rather than impulsive buried slip (recall that the results in Figure 1b are for slippage occurring as a step function in time). For a completely relaxed asthenosphere, an upper bound to this effect is given by steady slip in a free plate, and this in turn can be modeled to sufficient accuracy for our purposes by a buried two-dimensional edge dislocation in an elastic half-space. One such half-space calculation for buried slip on a 30° inclined fault is given by the dashed INTER curve in Figure 15. A comparison with Figure 1b shows that effects near the fault are quite similar in the two cases, both curves showing uplift predominating on the hanging wall block of the fault. However, further away the differences are significant, and the half-space model has large block offsets (uniform uplift and downwarping far from the fault) that are absent from both the solid and the dashed curves in Figure 1b. We expect the block offsets to be present in our model as well when the effects of steady buried slip that extends to the base of the plate are computed precisely. However, as we shall argue later (section 8), in the absence of stresses to maintain them, these offsets will be largely eliminated by the effects of isostatic compensation, which sufficiently far from the fault will become more important than the viscous stresses due to asthenospheric relaxation. Nearer to the fault, relaxation effects dominate, and for the interseismic data modeled in this paper, all of it within ~ 150 km from the trench, the differences between the uplift effects shown in Figure 1b and Figure 15 (INTER) are not great and do not affect our principal conclusions. Since we believe that the block offsets are not realistic in any case, we retain models like that shown in

Figure 1b to approximate the steady buried slip that contributes to the interseismic deformation. We are currently working on more refined models that include periodically recurring earthquakes, steady buried slip, and buoyancy effects.

4. OBSERVED DEFORMATION IN SOUTHWEST JAPAN

Active underthrusting of the Philippine Sea plate beneath southwestern Japan occurs along the Nankai Trough (Figure 4), and Seno (1978) has estimated the plate convergence rate here to be about 40 mm/yr, with the orientation of the slip vector indicating that the relative motion occurring across this part of the plate boundary is predominantly reverse dip-slip with a smaller component of right-lateral strike-slip movement. Subduction here is comparatively recent, and the absence of earthquakes deeper than about 100 km indicates that the underthrust plate has not yet penetrated below this depth (Shiono, 1974).

Recurrent destructive earthquakes are a characteristic feature of the Nankai Trough in southwest Japan, and historical accounts document shocks extending back as far as 684 A.D. and indicate repeat times of 100 to 200 years for major earthquakes in this region (Ando, 1975). Two great underthrust events occurred here in 1944 and 1946; the previous episode of rupturing on the Nankai Trough was in 1854.

Extensive leveling and tide gage measurements made since 1895 on Shikoku island and Kii peninsula (Figure 5) provide an unusually detailed record of the deformation associated with the 1946 Nankaido earthquake ($M=8.2$). Since

first being considered by Imamura (1929a, 1930), geodetic data from this region have been discussed and interpreted by numerous Japanese investigators (e.g. Okada and Nagata, 1953; Matuzawa, 1964; Ando, 1975), and recent studies have emphasized quantitative modeling of the observed deformation, particularly the extensive and well-documented postseismic movements (Fitch and Scholz, 1971; Nur and Mavko, 1974; Smith, 1974).

Leveling

The principal features of the deformation cycle are conveniently summarized by first order level changes on eastern Shikoku between Muroto Point and Komatsujima tide gage station. The location of this leveling route with respect to the 1944 and 1946 rupture planes obtained by Ando (1975) is shown in Figure 5, and the level changes are plotted in Figure 6. These data show that the interseismic movements (INTER, 1929-1895) are dominated by downwarping of the Muroto promontory, which subsided as much as 6 mm/yr with respect to Komatsujima during this interval. Minor uplift, at a maximum rate of about 1 mm/yr, occurred further inland. Coseismic deformation (CO, 1947-1929, Figure 8, top) shows an almost perfect reversal in the sense of movement, with a maximum uplift of ~ 800 mm at the southern-most leveling benchmark on Muroto (which subsided most during 1895-1929), and the greatest downwarping of about 400 mm occurring near where interseismic uplift was greatest. Level changes during 1947-64 (POST1) show that postseismic movements during this time involved predominantly land uplift, with maximum changes approaching 400 mm. The most recent leveling comparison available (1971-1964, POST2) suggests a pattern of deformation somewhat intermediate between the POST1 and INTER profiles.

Leveling Datum

In Figures 6 and 8 (top) we have followed Imamura (1930) and Hayashi (1970) in assuming that Komatsujima remained fixed during the interseismic and postseismic phases of the deformation. Although no continuously operating tide gage records were available during 1895-1929 to validate this assumption, leveling data from the Kii peninsula, ~ 60 km to the east of Komatsujima (see Figure 5), independently support Imamura's choice of datum level for eastern Shikoku. This leveling is tied to a continuously operating tide gage at Kushimoto (Figure 5), and with this absolute datum level as a tie, the 1929-1895 deformation profile from Kushimoto to Osaka given by Imamura (1929a) is very similar to that obtained further west on Shikoku by assuming Komatsujima was fixed during 1895-1929. Although no record exists at Komatsujima during 1947-50, continuous measurements made since 1951 show that the net change in sealevel for 1951-71 was 60 mm or less (Coastal Movements Data Center, 1975). The coseismic profile shown in Figure 8 (top) is taken from Geographical Survey Institute (1954), where it was assumed that benchmarks on the Japan Sea coast of southwest Honshu remained unmoved during 1929-47.

Short Term Postseismic Movements

Data from sealevel records and local-scale leveling in Shikoku provide further details on the short term time dependence of the postseismic movements. Uplift time histories from four tide gages and one frequently repeated leveling segment are shown in Figure 7, and both the tidal stations and the level line are located in Figure 5. Although smoothed versions of these same tidal records have been reproduced many times (e.g. Matuzawa, 1964;

Nur and Mavko, 1974), this smoothing gives a somewhat misleading impression of the precision of the observations, and Figure 7a shows the unsmoothed monthly means taken from Kawasumi (1956). As the pre-earthquake sections show, noise levels on these records are substantial, but ground movements are correspondingly large; our estimate of the total postseismic offset is shown beside each record. Although there is no truly objective way of assessing the error in these estimates, judging from the scatter in the 1944-46 data the uncertainty in total offset may be about ± 50 mm or perhaps somewhat greater.

The tilting of Muroto Point relative to Aki (a leveling benchmark 30 km further landward of the Nankai Trough, see Figure 5), is shown as a function of time in Figure 7b. The interseismic deformation is represented by a regular trenchward tilting of 7.5 mm/yr prior to 1946, and this trend is interrupted by the coseismic tilt offset of opposite sense, followed by a rapidly decaying transient. Note that the character of this transient agrees well with the postseismic uplift recorded by the URADO tide gage, located 40 km to the west of Aki. Note also that although the tilt transient appears to merge into a steady secular trend by 1950, the relative subsidence of Muroto during 1950-71 is 9.0 mm/yr, still somewhat higher than the pre-earthquake rate.

5. MODELING THE SEISMIC DEFORMATION CYCLE IN SOUTHWEST JAPAN

Computed Leveling Profile

Comparison of the observed level changes in eastern Shikoku (Figure 6) with the model calculations of Figure 1 clearly makes the point we shall advocate here, that the postseismic uplift is predominantly the result of

buried aseismic slip, while the observed interseismic downwarping is due principally to the effects of asthenospheric relaxation. Figure 8 compares the Muroto-Komatsujima leveling observations with computations made using a simple two-dimensional model of faulting in an elastic lithosphere overlying a Maxwell asthenosphere. The plate thickness assumed is 60 km, close to Kanamori's (1970) estimate of the lithospheric thickness for Japan obtained from the travel times of seismic bodywaves, and the asthenospheric viscosity is 4×10^{20} poise ($\tau = 40$ years). To fit the coseismic deformation in 1946, 3.2 m of slip occurs on a 30° -dipping fault that ruptures half the lithospheric thickness. Postseismic uplift is modeled by imposing comparable amounts of aseismic slip (2.5 m of underthrusting) on 30 km of the downdip extension of the coseismic fault, and the computed profile also includes the effects of 17 years of asthenospheric relaxation following the earthquake. The interseismic deformation, actually primarily the result of the 1854 earthquakes (INTER profile is for 1895-1929), is due to asthenospheric relaxation between $t=\tau$ and $t=2\tau$, supplemented by 30 mm/yr of steady aseismic slip on the bottom one quarter of the lithosphere-wide megathrust, which in our model does not rupture in either the coseismic or postseismic phases of the deformation cycle. The rate of slip is close to Seno's (1978) estimate of the dip-slip component of the plate convergence rate.

Two considerations suggest that the estimate of asthenospheric viscosity obtained in our model is too low by a factor of two to three. First of all, historic records of coastal deformation in 1854 (see Ando, 1975; Figure 8) suggest that relative to the 1946 shock, coseismic slip was at least 50% greater in 1854. Since the asthenospheric relaxation scales linearly with the

22

seismic slip, the same amount of downwarping between 1895 and 1929 can thus be obtained with a correspondingly higher asthenospheric viscosity. A second consistent bias results because the effects of earthquakes earlier than 1854 are not included in our model, and again these recurrent events will add to the deformation occurring in 1895-1929. For example, examination of Figure 2b indicates that the effects of the next-previous Nankai Trough event in 1707 would still be substantial 200 years later, producing 4 cm of downwarping per meter of coseismic slip between $t=5\tau$ and 6τ at $y/H = 1.0$. Approximately accounting for these biases in the way we have described here suggests that $\sim 10^{21}$ poise may be a more appropriate value for the effective viscosity of the asthenosphere beneath southwest Japan.

Detailed Postseismic Model

In order to demonstrate that postseismic buried slip can also satisfactorily explain the shorter term deformation exhibited by the tide gage and leveling data in Figure 7, we have computed the surface deformation due to a simple three-dimensional model whose geometry is suggested by Ando's (1975) coseismic model (Figure 5) and our two-dimensional postseismic model of Figure 8. This three-dimensional model consists of two fault segments each 150 km long and 30 km wide that dip 30° WNW, that have their top edge at 30 km depth, and that each have 2 m of dip-slip and 1 m of right-lateral strike-slip movement across them. The 2:1 ratio of reverse dip-slip to strike-slip motion is precisely that which Ando (1975) determined in his coseismic model, and as Figure 9 shows, the two aseismic fault planes are immediately adjacent to Ando's two coseismic faults.

Examination of Figure 9 and Table 1 demonstrates that this model, or obvious minor modifications to it, can explain all of the basic features of the observed postseismic deformation on Shikoku. The strikingly different behaviour of UWAJIMA and TOSASHIMIZU relative to URADO or Aki is seen to be due to the combination of an end effect of rupture termination and a minor asymmetry in the deformation contours that results from the small right-lateral component to the motion. Table 1 shows that, aided somewhat by the small amounts of subsidence expected from asthenospheric relaxation during 1947-50, the computed vertical offsets at UWAJIMA and TAKAMATSU provide a satisfactory match with the observations. Although the uplift at URADO is overestimated by 100 mm, the observed value represents only deformation to early 1949, and examination of the similarities between this tide gage recording and the Muroto-Aki tilt suggests the likelihood that an additional 100-150 mm of uplift occurred at URADO between 1949 and 1964. The observed ~200 mm of downwarping at TAKAMATSU is poorly accounted for by our model, and we can see no simple modification to it that will produce more than about 100 mm of subsidence so distant from the coseismic fault plane. Perhaps agreement to within a factor of two is satisfactory in any case. The general shape of the computed deformation profile between Muroto and Komatsujima agrees well with observations. Although the maximum uplift is about 150 mm too small, a short eastward extension of the western fault segment will result in a better match. The model provides an excellent fit to the Muroto-Aki postseismic tilt offset to 1964. About 0.3 m of additional slip during 1964-71 will adequately explain the continued tilting here as well as the level changes between Muroto and Komatsujima (POST2, Figure 6), although judging from the growing

similarities between post-1965 and pre-1946 deformation patterns seen in Figures 6 and 7b, the interseismic mechanisms of asthenospheric relaxation and steady buried slip are beginning to make significant contributions to the observed movements as well.

Finally, although the observed time-dependent features of the postseismic movements are not documented well enough to provide firm constraints for our model, these features can be readily accounted for by postulating appropriate time histories for the aseismic slip and its spatial growth. For example, the URADO record and the Muroto-Aki tilt history may be faithfully reflecting the slip-time function on the fault immediately beneath these locations, while the more uniform, smaller amplitude movements seen on the other three sealevel records might be attributed to the lateral and downdip growth of the aseismic slip zone on the fault, supplemented by the less impulsive effects of asthenospheric relaxation. Some evidence for downdip growth of slip comes from leveling carried out in 1950 and later in western Shikoku (see Fitch and Scholz, 1971, Figure 14; Hayashi, 1970, Figure I-10), which shows uplift of ~ 100 mm in NW Shikoku, rather than the downwarping shown by our model, although our results are consistent with 1948-50 level changes observed there.

An Alternative Model

Using results obtained in section 2 we can consider the applicability to the Nankaido data of a faulting model in which coseismic rupture breaks through the entire lithosphere. The surface displacements due to asthenospheric relaxation in one such model are shown in Figure 3, and a comparison of Figure 3a with the 1947-64 postseismic profile in Figure 6

suggests that the observed post-1946 upwarping of southern Shikoku might be identified with the local post-slip uplift shown near $y/H = 2.0$ in Figure 3a. This model is similar to one proposed by Smith (1974) and follows the spirit of Nur and Mavko's (1974) suggestion that post-Nankaido earthquake movements are due predominantly to the effects of asthenospheric relaxation.

Although in some respects appealing, this model (call it model B) has several defects that we believe make it a less likely explanation of the Nankaido data than the model we have proposed (which we shall refer to as model A). The features of this alternative model and its consistency with observations can be seen by referring to the data profile in Figure 6 and the model calculations in Figure 3. First of all, model B will have the same computed coseismic profile as that shown in Figure 8 for model A if we take $H = 30$ km for B. This follows because all lengths in each model (except fault slip) are scaled by H , and the instantaneous elastic response of the medium is independent of the plate thickness and does not depend on how much of it is ruptured coseismically. With this coseismic model, the post-slip surface deformation for model B can be obtained directly from Figure 3 and compared with data simply by multiplying the vertical axis by 3.2 m (the coseismic slip) and scaling horizontal distance y by taking $H = 30$ km. Proceeding in this way, it can be verified that in order to produce the observed uplift of Aki or URADO during 1947-48 with model B requires $\tau = 0.5$ years or $\eta = 5 \times 10^{18}$ poise. In addition, the time-dependence of the asthenospheric relaxation near $y/H = 2$ (Figure 2b) does not match the observations: uplift will continue past 1949 at rates comparable to those which obtained during the first few years after 1946, and the longer term deformation profile will

contain comparable amounts of uplift and downwarping, thus failing to satisfy the observed interseismic movement pattern. Although a non-linear rheology such as power law creep might account for the low asthenospheric viscosity, the other shortcomings of model B remain, and on balance we do not feel that lithospheric-wide rupturing and asthenospheric relaxation alone provide a viable mechanism for explaining the post-1946 Nankaido deformation.

6. OBSERVED DEFORMATION IN THE SOUTH KANTO REGION

On its northern boundary the Philippine Sea plate is being subducted beneath the Eurasian plate along the Sagami Trough (Figure 4), and near the western edge of this boundary the M=7.9 1923 Kanto earthquake occurred. Seno's (1978) analysis indicates ~ 30 mm/yr of relative plate motion here, with the preponderance of movement across the western end of the Sagami Trough being right-lateral strike slip, supplemented by a smaller amount of reverse dip-slip motion. Underthrusting is evidently quite recent in the South Kanto district, since no intermediate focal depth earthquakes (≥ 60 km) associated with the subduction of the Philippine Sea plate have been clearly identified here.

Great earthquakes apparently occur somewhat irregularly along the Sagami Trough, since historical accounts document a more extensive event in the same region in 1703 but no comparably large shock prior to this time until 818 A.D. Uplifted marine terraces record at least four 1703-type earthquakes in the previous ~ 6000 years (Matsuda et al., 1978).

Both seismic (Kanamori, 1971) and geodetic data (Ando, 1971, 1974) indicate that the 1923 earthquake involved shallow oblique angle underthrusting on a fault plane dipping 30° to 45° NE, with 6 m of right-lateral strike-slip movement and 3 m of reverse dip-slip. The location map in Figure 10 shows the surface projection of the coseismic fault plane inferred by Ando (1971).

All of the leveling and tide gage data used here has been taken from Scholz and Kato's (1978) study of the recent deformation history of the South Kanto region. We have also accepted their choices of absolute datum levels for the vertical movements, determined for the most part by ties between level lines and two continuously recording tide gages located at the tips of the Boso and Muira peninsulas.

Leveling

Figure 10 shows the locations of geodetic survey lines and tide gage stations we use here to summarize the characteristic features of the deformation cycle in the South Kanto region. The principal observations come from repeated first order leveling surveys carried out since 1895 along two routes, from Tokyo to Atami, crossing the landward extension of the Sagami Trough, and along the east coast of the Boso peninsula, extending to within ~30 km of the subduction boundary.

Examination of the CO-, POST-, and INTER-seismic level changes on the East Boso route, Figure 11, illustrates clearly the two most distinctive features of the deformation cycle in the South Kanto region: first, that the post-seismic vertical movements are very small, and second that they do not

differ substantially from the interseismic secular deformation, either in their spatial pattern or in their average rates of movement. Although differences between the POST and INTER profiles do occur near the landward extension of the Sagami Trough on the Tokyo-Atami line (Figure 12), outside this narrow ~ 20 km wide region the movement patterns seen on the two leveling routes are very similar, and indicate predominately downwarping in both the postseismic and interseismic time intervals. A comparison of Figure 11 with Figure 8 (top) shows clearly that the postseismic vertical movements in South Kanto differed very markedly from those which followed the 1946 Nankaido earthquake: on leveling segments in comparable positions with respect to the coseismic fault plane (refer to Figures 5 and 10), there are only modest amounts of downwarping in the 15 years after the Kanto earthquake (~ -50 mm), while uplift reaching almost 400 mm occurred in the 17 years following the Nankaido earthquake.

Although the CO and INTER profiles for East Boso are generally similar to those for corresponding phases of the deformation cycle in southwest Japan, several significant differences are worthwhile mentioning. First of all, distinct offsets clearly evident on the coseismic profile (e.g. near 25 and 40 km in Figure 11) are due to secondary faulting at the time of the 1923 earthquake (Imamura, 1929b; Ando, 1974; Scholz and Kato, 1978). Secondly, the INTER profile does not steepen monotonically toward the Sagami Trough, but instead flattens or perhaps even reverses its trend inside about 30 km distance from the trench. This feature is also seen, though rather less definitively, on the POST profile, and also shows up well on the West Boso leveling data presented by Scholz and Kato (1978; see their Figure 8).

Additional differences and complexities in the deformation pattern within the South Kanto district are evident in Figure 12, the leveling results from the Tokyo-Atami line. Comparison with Figure 11 shows that the coseismic uplift is considerably narrower here, a feature that Ando (1971) has attributed to the asymmetry that results from the large right-lateral component in the coseismic slip. Furthermore, the CO profile has no sharp offset near the Kozu-Matsuda fault, the landward extension of the Sagami Trough, but instead shows an irregular drop to zero level change between this fault trace and Atami. During the POST and INTER phases of the deformation cycle, the Tokyo-Atami data exhibit considerable complexity within ~ 20 km of the Kozu-Matsuda fault, showing a sharp offset near -10 km in 1924-39 and steep gradients centered near $+10$ km during 1939-64. However, outside this region both the POST and INTER movements are very similar to those which occurred on the eastern side of Boso peninsula, ~ 70 km to the SE.

Short Term Postseismic Movements

In addition to the rather coarse time resolution provided by the leveling measurements, a smaller number of additional observations, summarized in Figure 13, provide more detail on the movements that occurred immediately after 1923 and show how, in a few locations at least, the postseismic movements gradually merged into the steady interseismic phase of the deformation cycle. Figure 13a shows shear strain changes determined from repeated length measurements of a ~ 100 m aperture rhombus located at Mitaka, near Tokyo (see Figure 10) and discussed previously by Fujita (1972) and Scholz and Kato (1978). These results show that postearthquake horizontal

strain changes at Mikata were remarkably large compared with post-1923 vertical deformation observed elsewhere in the South Kanto district, amounting by 1932 to more than one third of the coseismic strain drop. The data suggest changes occurred episodically in two time intervals, during a one year period immediately following the earthquake, and spread over ~ 4 years in 1928-32. The total change to 1935 is about 15 μ strain and the plane of maximum right-lateral shear straining has an azimuth of 138° , nearly parallel to the Sagami Trough and the strike of Ando's (1971) coseismic fault. The time history of movements at Mitaka is somewhat reminiscent of the post-Nankaido tilt changes between Muroto and Aki (Figure 7b), although the timescale is somewhat longer at Mitaka. Note also that data from the nearby segments of the Tokyo-Atami leveling route, only about 10 km southeast of Mitaka, show that the large postseismic shear strain changes that occurred there were accompanied by only modest amounts of vertical deformation: 1924-1939 level changes are less than 40 mm, and corresponding tilts are about 1×10^{-6} radians, or about an order of magnitude smaller than the shear strain changes.

Tide gage recordings (Figure 13b) provide some constraints on the postseismic movements as well as giving precise estimates for the average rates of interseismic subsidence at the tips of Boso and Muira peninsulas. Sealevel measurements made since 1896 at ABURATSUBO indicate an average subsidence rate of 4 mm/yr since that time, and observations made since 1931 at MERA are also consistent with this average rate. Both tide gage records appear to show a slightly higher subsidence rate following the Kanto earthquake until ~ 1940 , roughly 5-6 mm/yr, and a rather lower rate thereafter, about 3-4 mm/yr, although the indicated rate change is small and perilously close to the noise level of the observations.

7. MODELING THE SEISMIC DEFORMATION CYCLE IN THE SOUTH KANTO REGION

Although several features of the observed deformation in the South Kanto district are distinctively different from those occurring in southwest Japan, the same basic model is capable of explaining both sets of observations. The clearest differences are seen in the postseismic movements, and the almost universal subsidence shown in both the POST- and INTER-seismic profiles for South Kanto, as well as their other general similarities, suggests that both are predominantly the result of asthenospheric relaxation. However, the Mitaka strain data provide a crucial clue that suggest buried postseismic slip nonetheless did occur after 1923, since the large and irregular movements observed at Mitaka cannot be adequately accounted for by asthenospheric relaxation, which can produce only modest, smoothly varying postseismic strain changes. The absence of comparable vertical deformation near Mitaka indicates that any buried slip that may be responsible for the movements occurring there must involve predominantly strike-slip rather than dip-slip fault motions. Such a predominance of strike-slip over dip-slip movements is also suggested by Ando's (1971) coseismic model of the Kanto earthquake, and we shall show that with this same partitioning modest amounts of postseismic slip (< 1 m) can adequately satisfy the Mitaka strain data without violating the post-1923 leveling observations.

Computed Leveling Profiles

As Figures 11 and 12 show, the general features of the South Kanto leveling data can be matched adequately using two dimensional faulting models that include the effects of asthenospheric relaxation. The model geometry is

similar to that used in fitting the Nankaido data: the lithosphere is assumed to be 60 km thick and the coseismic fault plane ruptures the free surface but does not break through the entire plate thickness. The timescale of the post- and inter-seismic movements is matched by taking an asthenospheric viscosity of 6×10^{20} poise ($\tau = 60$ years). Following Ando's (1971) coseismic model the assumed fault dip is 45° , and in order to fit both coseismic profiles, which differ somewhat, we take $d/H = 0.5$ for the Tokyo-Atami line and $d/H = 0.75$ for the east Boso profile. Slippage is 3.4 m in both cases. Because faulting in the middle of the plate between $d/H = 0.5$ and 0.75 causes very little stress change in the Maxwell half-space, the asthenospheric relaxation effects of these two models are nearly identical, so the slight differences in the coseismic models has no significant effect on the rest of the deformation cycle. Based on Seno's (1978) estimate of a 30 mm/yr convergence rate between the Asian and Philippine Sea plates in this region, and a 2:1 partitioning between strike-slip and dip-slip motion, the steady aseismic slip presumed to occur downdip of the coseismic rupture is imposed at a rate of 10 mm/yr. Because of its low rate, the surface effects of this slippage are small compared with those due to asthenospheric relaxation. For the moment we assume that any effect of a dip-slip component to the postseismic movements is small and can be safely neglected, arguing this point later in our discussion of modeling the shear strain changes for the Mitaka rhombus.

For the East Boso level profile, Figure 11, both the shape and the amplitude of the computed POST and INTER curves match the observations quite well. Although Scholz and Kato (1978) have suggested that the shallowing in the trenchward tilt of the observed profiles inside ~ 30 km from the Sagami

Trough is due to block tilting on the Boso peninsula, this feature can also be very naturally explained by our model as an effect of asthenospheric relaxation (recall Figure 1a and our earlier discussion of this feature of our model). The shape and level of the coseismic changes are successfully fitted, but details such as the secondary faulting offsets are not reproduced by the model.

The match of our model to the Tokyo-Atami data, Figure 12, is clearly not as good. Though satisfactory elsewhere, the CO model, which ruptures to the surface, completely fails to match the observations between -20 and 0 km on the profile. Although some improvement to the fit can be obtained by postulating that coseismic slip does not reach the surface (Scholz and Kato, 1978), the problem may lie elsewhere. We believe that the observed uplift in this region is due to a complex superposition of the effects of splay faulting and rupturing of semi-consolidated alluvial deposits, effects that are commonly observed in thrust faulting (e.g. Buwalda and St. Amand, 1955; U.S. Geological Survey Staff, 1971), but which are difficult to reproduce quantitatively without using many near-surface fault segments (e.g. see Jungels and Frazier, 1973). Although the mismatch between model and data is quite large, if our interpretation is correct the effects on the Maxwell half-space will be the same as those produced by surface rupturing, and accordingly we retain this model in calculating the deformation due to asthenospheric relaxation. With this model, the general shape and amplitude of the POST profile from -10 to +60 km and the INTER profile more than 20 km from the Kozu-Matsuda fault are satisfied well. However, the sharp gradients near -10 km in the POST profile, and around +10 km in the INTER phase of the

deformation cannot be matched by a source as deep as the asthenosphere, and must be due to deformation occurring at shallow depths. We do not attempt to model these complex fault zone movements, conceding that our preferred interpretation satisfies only the long wavelength features of deformation observed on the Tokyo-Atami leveling route.

Using our model derived from the leveling data, the time history of post-1923 vertical movements at the MERA and ABURATSUBO tide gages is specified, and as Figure 13b shows, the model calculations are in excellent agreement with the observations. Not only is the general shape reproduced in each case, but the indicated decrease in subsidence rate is also matched by the slightly concave downward shape of the model curves: the average rate to 1940 is 5.4 mm/yr, while since 1940 the subsidence rate is near 3.5 mm/yr. Because we have modeled the asthenospheric relaxation effects of only a single earthquake, the 1923 event, our model cannot be used to fit the 1896-1923 tide gage record for ABURATSUBO, since the movements during this time are due to the superposed effects of previous great earthquakes in the Sagami Trough.

Detailed Postseismic Model

The asthenospheric relaxation models used to fit the postseismic leveling data cannot satisfy the large post-1923 strain changes at Mitaka. However, two features of the observed movements suggest a modification to the POST models of Figures 11 and 12 that will leave the computed leveling profiles essentially unchanged yet will produce enough horizontal deformation to explain the Mitaka data. First of all, significant movements near Mitaka are largely horizontal, since nearby portions of the Tokyo-Atami leveling route

show comparatively little contemporaneous tilting. Furthermore, the maximum right-lateral shear at Mitaka is oriented nearly parallel to the Sagami Trough, so that the maximum compressive strain axis is not perpendicular to the trench but is rotated 45° counter-clockwise with respect to it. Both observations suggest effects due predominantly to right-lateral strike-slip rather than reverse dip-slip fault movements. The surface deformation effects produced by one satisfactory postseismic buried slip model that incorporates these features is shown in Figure 14. The slip ratio is 2:1 (0.6 m strike-slip, 0.3 m reverse dip-slip), as in Ando's coseismic model, and the fault plane, whose top edge is at 30 km depth, is 20 km wide, 130 km long, and dips 45° NE. The dashed rectangle is the surface projection of Ando's coseismic fault plane and the adjacent solid rectangle shows the corresponding postseismic fault.

Model results showing the orientation of the horizontal principal strains, their magnitudes, and the maximum right-lateral shear (in μ strain) are plotted at Mitaka (circled) and nearby in Figure 14. Although the 1923-35 observations are not matched exactly, minor modifications to the simple model of Figure 14 can arbitrarily improve the fit. The computed value of γ_{12} , 10 μ strain, is less than the observed, but additional post-slip shearing resulting from the asthenospheric relaxation due to the 6 m of right-lateral coseismic slip (not modeled here) is probably sufficient to make up the difference. Alternatively, increasing only the strike-slip component of the postseismic slip will accomplish the same purpose, and do so without measurably changing the vertical deformation field. The computed maximum compressive strain orientations agree with observations almost exactly ($N6^\circ E$

versus $N30^{\circ}E$), but the observed ratio of maximum to minimum compressive strain is badly misfit (6.5 versus 0.3). However, as Figure 14 reveals, locations NW of Mitaka show a favorable ratio, so this misfit can be easily remedied by shortening the postseismic fault ~ 20 km at its northwestern end.

As Figure 14 demonstrates, relatively small amounts of postseismic uplift are produced by our buried slip model. The computed uplift pattern shows no sharp gradients across either the Tokyo-Atami or East Boso leveling routes, and maximum elevation changes along these lines are less than 40 mm. Long wavelength level changes of this magnitude will not substantially alter the computed POST profiles in Figures 11 and 12 or the conclusions based on them, since any slight degradation in the model fit can be recovered by further modest changes in model parameters. Note also that the observed POST profiles on the Boso Peninsula are not tied to an absolute datum, since MERA did not begin operation until 1931. The profile in Figure 11 follows Scholz and Kato's (1978) assumption that B.M. 3934 (Figure 10) subsided at a rate of 0.3 mm/yr during 1924-31, and if this inference is in error there is an additional degree of freedom available in modeling the POST data on Boso peninsula.

Finally, it is perhaps worthwhile mentioning that the postseismic straining at Mitaka is not easily explained by the backslip model proposed by Fitch and Scholz (1971) and Scholz and Kato (1978) to account for postearthquake vertical deformation in SW Japan and in the South Kanto district. In this model, postseismic slip is postulated to act in opposite sense to the coseismic slip, implying left-lateral normal faulting after the Kanto earthquake. Clearly, such movements on the coseismic fault or its downward extension will not result in a satisfactory fit to the 1923-1935

Mitaka data, which require the maximum right-lateral shear strain direction to be parallel to the strike of the coseismic fault plane.

8. AN ELASTIC HALF-SPACE MODEL

Faulting in an elastic half-space is a simpler alternative to the model proposed here that may be appropriate if the downdip width of the coseismic fault plane is much less than the lithospheric thickness or if asthenospheric viscosity is much greater than 10^{21} poise. Considering this alternative model, in which the effects of asthenospheric relaxation are absent, also serves the purpose of highlighting distinctive features of the deformation produced by underthrusting models that do include the effects of lithosphere-asthenosphere coupling. Though not precisely identical, the elastic half-space model has many similarities with the finite element models used by Shimazaki (1974), Scholz and Kato (1978), Kato (1978), and Seno (1978) to simulate subduction and interseismic deformation in several regions of Japan.

One particular half-space model, for an infinitely long 30° inclined fault, is shown in Figure 15. Calculations have been made using the simple expressions given by Freund and Barnett (1976). The specific example that is illustrated has parameters close to those of the Nankaido model in Figure 8, and except for the unit scaling of the uplift, the CO profile is the same as the computed model profile at the bottom of Figure 8.

Since in our model the postseismic effects of the Nankaido earthquake are dominated by the purely elastic uplift due to downdip aseismic slip, the

differences between the two models is slight and in this instance the half-space formulation is satisfactory. Other documented examples of uplift following underthrust earthquakes (e.g. Kasahara, 1975; Abe, 1977; Brown et al., 1977) have been explained in the same way, suggesting that half-space models are adequate to explain postseismic movements in these regions as well. However, the widespread downwarping that followed the 1923 Kanto earthquake is not readily explained in this way, although, as our results show, the superposed effects of predominantly right-lateral buried slip and asthenospheric relaxation can account for the differing features of the South Kanto deformation quite well.

The greatest differences between the two models is shown by the interseismic deformation profiles. Steady buried slip, occurring to great depth in the half-space model (simulating slip extending to the base of a plate that overlies a completely relaxed asthenosphere), produces only uplift of the hanging wall block of the fault, and although the trenchward tilt of the profile agrees with observations, the absence of coastal downwarping does not. At first glance it appears that this difficulty can be circumvented by arbitrarily assigning most or all of the relative motion between the blocks to the footwall side of the fault (i.e. the oceanic plate), shifting the zero level of the INTER profile upward so that it shows little or no net uplift. Although agreeing with intuition (the oceanic plate must be subducted), this expedient solves one problem at the expense of creating another: imposing the same partitioning in absolute motion across the CO and POST faults will result in little or no coseismic and postseismic uplift, contrary to observations; on the other hand, unless the same partitioning of absolute motion between

oceanic and continental plates obtains across the CO and POST fault planes there will be an unphysical mismatch in slip (discontinuity in Burger's vector) between these segments and the INTER portion of the fault. Therefore, without some other mechanism to continually downwarp the INTER profile in Figure 15, models like this half-space one are not by themselves appropriate for explaining the interseismic deformation landward of trenches. These models may, however, be applicable to convergent plate boundaries where continent-continent collision is occurring, since in these regions the lithosphere is thicker, asthenospheric viscosities may be higher, and normal subduction does not occur.

The interseismic model we propose contains as one of its elements a deformation profile similar to the INTER curve in Figure 15, and although asthenospheric relaxation supplies some of the needed downwarping that modifies this uplift, our model shares one significant shortcoming with the half-space model. Within a few lithospheric thicknesses of the plate boundary, subsidence due to asthenospheric relaxation dominates uplift resulting from steady buried slip. However, at greater distances from the trench, the block offsets produced by steady slip become progressively more important. Vertical movements as large as these so far inland from the trench are not observed on the Japanese islands, and their presence is a clear defect of our model.

We believe that this unrealistic feature of the dislocation formulation that is common to both our model and the elastic half-space problem will be absent or subdued if buoyant restoring forces are accounted for in the asthenospheric relaxation model. In the absence of an exact solution

incorporating this effect, we use an order-of-magnitude calculation to argue that near the coseismic rupture buoyant effects are insignificant compared to viscous relaxation but that they become dominant further away from the plate boundary. In general, buoyant forces will oppose coseismically-imposed viscous stresses and act to reduce elevation changes resulting from fault slip in the lithosphere. Near the coseismic fault the viscous stresses in the asthenosphere are of the order of the earthquake stress drop, tens of bars, and like the coseismic deformation profile will rapidly decrease in amplitude away from the fault and become negligible beyond a few fault dimensions away from the trench. In contrast, the buoyant pressure exerted on the lithosphere due to its upward or downward displacement into the asthenosphere is much smaller but does not decay away from the coseismic fault. For a displacement ΔH of the lithosphere-asthenosphere boundary, the incremental pressure on the lithosphere is roughly

$$\Delta \rho \, g \, \Delta H$$

where $\Delta \rho$ is the density contrast and g is the acceleration due to gravity. If $\Delta \rho$ is 0.3 g/cm^3 , this pressure is 0.03 bars per meter of ΔH . A related effect is the isostatic compensation due to lateral pressure gradients in the asthenosphere beneath coseismically uplifted and downwarped portions of the lithosphere. Neglecting coseismic density perturbations, an upper bound to this effect is a pressure difference

$$\bar{\rho} \, g \, \Delta H$$

where $\bar{\rho}$ is the mean density of lithosphere and asthenosphere. If $\bar{\rho} = 3.0 \text{ g/cm}^3$, then this pressure is less than $\sim 0.3 \text{ bars/m}$ of ΔH . Due to recurrent earthquakes, the viscous stresses near the fault will never decay to

values comparable to this, and so close to the plate boundary neglect of buoyant forces and isostatic compensation effects is justified. However, further away the viscous stresses become negligible and buoyant restoring forces and isostatic effects will ensure that the large block offsets indicated by the dislocation solutions for steady buried slip do not in fact occur.

9. SUMMARY AND DISCUSSION

Our calculations have indicated that with conventionally-accepted values of lithospheric thickness and asthenospheric viscosity, the interseismic subsidence observed landward of the major subduction zones off Japan can be understood as predominantly an effect of asthenospheric relaxation following major trench earthquakes, supplemented by steady aseismic slip on the deeper parts of the plate boundary, while the sometimes spectacular deformation episodes that follow great underthrust earthquakes in Japan and elsewhere are adequately explained by buried aseismic slip that acts in the same sense as the coseismic movements and occurs on the downdip extension of the coseismic fault plane. Although the effects of asthenospheric relaxation on observed postseismic movements are in general much smaller, data from the South Kanto region suggest they can nonetheless be locally significant.

The model used here is obviously quite idealized in several respects. The effects of plate motion on subduction boundary deformation are approximated kinematically by imposing steady slip at the relative motion rate on the bottom segment of the plate bounding fault, thus evading explicit

consideration of the forces that drive the plates. The surface deformation effects of the downward pull of the subducting oceanic lithosphere are ignored in our model, and this neglect may be the most significant defect of our simplified approach. If this process is important, its effects may be very similar to asthenospheric relaxation and produce downwarping of the overlying continental lithospheric plate. However, the magnitude of this effect is very difficult to assess, since its importance depends on the magnitude of the net downward force acting on the subducted plate, the difference between the negative buoyant force of the dense sinking plate and the viscous resistance to its descent into the mantle. Forsyth and Uyeda (1975) suggest that both forces are large and of comparable size, so their difference is quite uncertain and hence the net effect on surface deformation is correspondingly in doubt. A further shortcoming of our model is its neglect in considering any modifications to asthenospheric flow that may result from the presence of a long subducted slab in the upper mantle beneath the continental plate. However, since the underthrust Philippine Sea plate has not yet penetrated below 100 km beneath either SW Japan or the South Kanto region, neglect of both subduction effects may be justified for the model calculations made here. Furthermore, in both Northern Honshu and Hokkaido (see Figure 4) the subducted Pacific plate is nearly 2000 km long and reaches depths in excess of 700 km, yet the pattern of interseismic subsidence (see Kato, 1978; Seno, 1978; Shimazaki, 1974) closely matches that occurring landward of the Nankai and Sagami Troughs. This comparison indicates that the length of underthrust slab has little effect on the surface deformation and suggests that downdrag by the subducted plate may be less important than asthenospheric relaxation in accounting for the observed interseismic movements.

Although the deformation model proposed here is certainly not unique, our analysis does indicate that in underthrust zones the deformation effects of asthenospheric relaxation are likely to be quite significant and cannot be safely ignored. While our model may not match all features of the observations significantly better than those proposed previously (Fitch and Scholz, 1971; Nur and Mavko, 1974; Smith, 1974; Bischke, 1974; Scholz and Kato, 1978), it is worthwhile emphasizing that not only does our model provide an adequate fit to the entire deformation cycle, but does so without requiring extreme or exotic properties for the lithosphere and asthenosphere or ad hoc slip laws for the plate boundary megathrust. While our choice of lithospheric thickness, 60 km is consistent with seismic estimates, and the value determined for asthenospheric viscosity, 10^{21} poise, agrees with independent determinations from studies of isostatic rebound, our conclusions are not strongly tied to these choices, and changes by as much as factors of two to three in these model parameters would not substantially alter our interpretation. Furthermore, if standard estimates of lithospheric thickness, asthenospheric viscosity, and depth of coseismic underthrusting are even approximately correct, something like the asthenospheric relaxation effects we have modeled here must occur at subduction zones. In addition, the predicted effects of asthenospheric relaxation produce deformation that agrees with observations and cannot be simply matched by dislocation models. These characteristics contrast significantly with the analogous situation at strike-slip plate boundaries, where features of the observed deformation can be suitably explained with half-space dislocation models, and where the shallow depth of coseismic faulting makes it somewhat less likely that asthenospheric relaxation contributes significantly to the surface movement pattern.

APPENDIX A

In Rundle (1978) it was shown how to compute the surface uplifts due to a finite, rectangular thrust fault. Here we specialize to the case of an infinitely long fault. From Equation (2) of Rundle (1978), the uplift can be written as

$$u_z = \int_S ds \int_0^\infty \left\{ (X_{10}(0) J_0(kr) + X_{12}(0) J_2(kr) \cos 2\varphi) \sin 2\theta + X_{11}(0) J_1(kr) \sin \varphi \cos 2\theta \right\} k dk \quad (A1)$$

where all quantities are as defined in Rundle (1978). The integral over x' reduces to an integral from $(y - y')$ to infinity. Since the kernel functions $X_{lm}(0)$ contain only the source depth h , layer thickness H and fourier wave number k , the x' integral can be performed analytically (Erdelyi et al, 1954):

$$u_z = 2 \int_0^\infty dk \int_{\xi_1}^{\xi_2} \left\{ (X_{10}(0) + X_{12}(0)) \cos k(y - y') \sin 2\theta + X_{11}(0) \sin k(y - y') \cos 2\theta \right\} d\xi \quad (A2)$$

where $y' = h \cot \theta = \xi \cos \theta$ is the horizontal source coordinate as in Rundle (1978). Note that the kernels $X_{10}(0)$, $X_{11}(0)$, $X_{12}(0)$ can be viewed as distributions on k for fixed values of h (and thus y' or ξ). If we can define an "average" $\bar{k}$ such that

$$u_z \approx 2 \int_{\xi_1}^{\xi_2} \left\{ (\bar{X}_{10}(0) + \bar{X}_{12}(0)) \cos \bar{k}(y - y') \sin 2\theta + \bar{X}_{11}(0) \sin \bar{k}(y - y') \cos 2\theta \right\} d\xi \quad (A3)$$

then we expect the solution for the infinitely long fault to be approximately periodic in y . That this in fact is the case can be verified by actual calculation. Many such infinite problems in mathematical physics

display similar "unphysical" characteristics and reflect the general conclusion that the solution is only valid near the source. In our case, we find that $\bar{k} \cong 2/(W^2 + H^2)^{1/2}$, a reasonable result.

APPENDIX B

Rundle (1978) gave the functions $\xi_n(t)$ for the case where the loading on the fault was a Heaviside step $H(t)$ in time. For the case of a Heaviside ramp $R(t)$, a reasonable approximation to steady slip, it can easily be verified that the result is:

$$\xi_n(t) = V \left\{ R(t) - \tau H(t) \left[n - e^{-t/\tau} \sum_{m=0}^{n-1} \sum_{p=0}^m \frac{1}{p!} \left(\frac{t}{\tau} \right)^p \right] \right\} \quad (B1)$$

where τ is the relaxation time defined in Rundle (1978) and V is the constant relative slip velocity. For the Maxwell rheology used in this paper, $\tau = 2\eta/\mu$ where η is the effective viscosity and μ is the rigidity. A superposition of ramp loads at different times can be used to represent a fault sliding steadily for some finite time. The corresponding functions $\xi_n(t)$ are then similar superpositions of (B1).

ACKNOWLEDGMENTS

Discussion with T. J. Fitch, T. C. Hanks, K. Kasahara, A. H. Lachenbruch, J. C. Savage, C. H. Scholz and A. T. Smith resulted in considerable improvement in the manuscript. M. Ando provided helpful comments on Japanese tide gage measurements and supplied an elusive reprint. This work was supported in part by a contract from the U.S. Department of Energy to Sandia Laboratories.

REFERENCES

- Abe, K., Tectonic implications of the large Shioya-oki earthquakes of 1938, Tectonophysics, 41, 269-289, 1978.
- Anderson, D. L., Accelerated plate tectonics, Science, 187, 1077-1079, 1975.
- Ando, M., A fault-origin model of the great Kanto earthquake of 1923 as deduced from geodetic data, Bull. Earthquake Res. Inst. Tokyo Univ., 49, 19-32, 1971.
- Ando, M., Seismotectonics of the 1923 Kanto earthquake, J. Phys. Earth, 22, 263-277, 1974.
- Ando, M., Source mechanisms and tectonic significance of historical earthquakes along the Nankai trough, Japan, Tectonophysics, 27, 119-140, 1975.
- Bischke, R. E., A model of convergent plate margins based on the recent tectonics of Shikoku, Japan, J. Geophys. Res., 79, 4845-4857, 1974.
- Bott, M. H. P., and D. S. Dean, Stress diffusion from plate boundaries, Nature, 243, 339-341, 1973.
- Brown, L. D., R. E. Reilinger, S. D. Holdahl and E. I. Balazs, Postseismic crustal uplift near Anchorage, Alaska, J. Geophys. Res., 82, 3369-3378, 1977.
- Buwalda, J. P., and P. St. Amand, Geological effects of the Arvin-Tehachapi earthquake, in Earthquakes in Kern County California during 1952, edited by G. B. Oakshott, State of California, Department of Natural Resources, Division of Mines, Bull. 171, 41-56, 1955.
- Cathles, L. M., III, The Viscosity of the Earth's Mantle, Princeton University Press, Princeton, N.J., 1975

- Chinnery, M. A., Deformation of the ground around surface faults, Bull. Seismol. Soc. Amer., 51, 355-372, 1961.
- Coastal Movements Data Center, Table and Graphs of Annual Mean Sea Level Along the Japanese Coast, 37 pp., Tokyo, 1976.
- Elsassar, W. M., Convection and stress propagation in the upper mantle, in The Application of Modern Physics to the Earth and Planetary Interiors, edited by S. K. Runcorn, 223-246, Wiley-Interscience, 1969.
- Erdelyi, A., W. Magnus, F. Oberhettinger, and F. G. Tricomi, Tables of Integral Transforms, II, McGraw Hill, 1954.
- Fitch, T. J. and C. H. Scholz, Mechanism of underthrusting in southwest Japan: A model of convergent plate interactions, J. Geophys. Res., 76, 7260-7292, 1971.
- Forsyth, D., and S. Uyeda, On the relative importance of the driving forces of plate motion, Geophys. J. R. Astr. Soc., 43, 163-200, 1975.
- Freund, L. B., and D. M. Barnett, A two-dimensional analysis of surface deformation due to dip-slip faulting, Bull. Seismol. Soc. Amer., 66, 667-676, 1976.
- Fujita, N., Deformation of the rhombus base line at Mitaka (in Japanese), J. Geod. Soc. Jap., 18, 8-16, 1972.
- Fung, Y. C., Foundations of Solid Mechanics, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1965.
- Geographical Survey Institute of Japan, Resurvey of the South-Western part of Japan after the Great Nankaido earthquake of 1946, J. Geogr. Surv. Inst. Jap., 4, 1-69, 1954.

- Hanks, T. C., The Kuril trench - Hokkaido rise system: Large shallow earthquakes and simple models of deformation, Geophys. J. Roy. Astron. Soc., 23, 173-190, 1971.
- Hayashi, T., A study of vertical movements of the earth's crust by means of the precise leveling, J. Geod. Soc. Jap., 15, 1-69, 1970.
- Hastie, L. M. and J. C. Savage, A dislocation model for the 1964 Alaska earthquake, Bull. Seismol. Soc. Amer., 1389-1392, 1970.
- Imamura, A., On the chronic and acute earth-tilting in the Kii Peninsula, Jap. J. Astron. Geophys., 7, 31-45, 1929a.
- Imamura, A., On the multiple source of origin of the great Kanto earthquake and its relation to the fault system connected with the earthquake, Proc. Imp. Acad. Jap., 5, 330-333, 1929b.
- Imamura, A., On the chronic and acute earth-tilting in the southern part of Shikoku, Jap. J. Astron. Geophys., 8, 29-37, 1930.
- Jungels, P. H., and G. A. Frazier, Finite element analysis of the residual displacements for an earthquake rupture: source parameters of the San Fernando earthquake, J. Geophys. Res., 78, 5062-5083, 1973.
- Kanamori, H., Mantle beneath the Japanese arc, Phys. Earth Planet. Interiors, 3, 475-483, 1970.
- Kanamori, H., Faulting of the great Kanto earthquake as revealed by seismological data, Bull. Earthquake Res. Inst. Tokyo Univ., 49, 13-18, 1971.
- Kasahara, K., The nature of seismic origins as inferred from seismological and geodetic observations, 1, Bull. Earthquake Res. Inst. Tokyo Univ., 35, 473-532, 1957.
- Kasahara, K., Aseismic faulting following the 1973 Nemuro-oki earthquake, Hokkaido, Japan (a possibility), Pure Appl. Geophys., 113, 127-139, 1975.

- Kato, T., Recent crustal movement in the Tohoku district for the period 1900-1975, Preprint, 1978.
- Kawasumi, H., Crustal deformation in the Shikoku district as deduced from tide gage records, second report (in Japanese), Shikoku-Chiho Sogo Kaihatsu Shingikai, __, 30-36, 1956.
- Matsuda, T., Y. Ota, M. Ando and N. Yonekura, Fault mechanism and recurrence time of major earthquakes in southern Kanto district, Japan as deduced from coastal terrace data, Bull. Geol. Soc. Amer., 89, in press, 1978.
- Matuzawa, T., Study of Earthquakes, Uno Shoten, Tokyo, 213 pp., 1964.
- McNutt, M., and H. W. Menard, Lithospheric flexure and uplifted atolls, J. Geophys. Res., 83, 1206-1212, 1978.
- Melosh, H. J., Nonlinear stress propagation in the earth's upper mantle, J. Geophys. Res., 81, 5621-5632, 1976.
- Nur, A. and G. Mavko, Postseismic viscoelastic rebound, Science, 183, 204-206, 1974.
- Okada, A., and T. Nagata, Land deformation of the neighborhood of Muroto Point after the Nankaido great earthquake in 1946, Bull. Earthquake Res. Inst., 31, 169-177, 1953.
- Plafker, G. and J. C. Savage, Mechanism of the Chilean earthquakes of May 21 and 22, 1960, Bull. Geol. Soc. Amer., 81, 1001-1030, 1970.
- Rundle, J. B., Viscoelastic crustal deformation by finite, quasistatic sources, submitted to J. Geophys. Res., 83, 1978.
- Rundle, J. B. and D. D. Jackson, A kinematic viscoelastic model of the San Francisco earthquake of 1906, Geophys. J. Roy. Astron. Soc., 50, 441-458, 1977.

- Savage, J. C. and R. O. Burford, Accumulation of tectonic strain in California, Bull. Seismol. Soc. Amer., 60, 1877-1896, 1970.
- Savage, J. C. and R. O. Burford, Geodetic determination of relative plate motion in central California, J. Geophys. Res., 78, 832-845, 1973.
- Savage, J. C., and W. H. Prescott, Comments on "Nonlinear stress propagation in the earth's upper mantle" by H. J. Melosh, J. Geophys. Res., 83, in press, 1978a.
- Savage, J. C. and W. H. Prescott, Asthenosphere readjustment and the earthquake cycle, J. Geophys. Res., 83, 3369-3376, 1978b.
- Scholz, C. H. and T. Kato, The behaviour of a convergent plate boundary: crustal deformation in the South Kanto district, Japan, J. Geophys. Res., 83, 783-797, 1978.
- Seno, T., Instantaneous rotation vector of Philippine Sea plate relative to the Eurasian plate, Tectonophysics, 42, 209-226, 1977.
- Seno, T., Intraplate seismicity in Tohoku and Hokkaido and large interplate earthquakes: a possibility of a large interplate earthquake off the southern Sanriku coast, Northern Japan, J. Phys. Earth, in press, 1978.
- Shimizaki, K., Pre-seismic crustal deformation caused by an underthrusting oceanic plate, in eastern Hokkaido, Japan, Phys. Earth Planet. Interiors, 8, 148-157, 1974.
- Smith, A. T., Time-dependent strain accumulation and release at island arcs: implications for the 1946 Nankaido earthquake, Ph.D. thesis, Massachusetts Institute of Technology, Cambridge, Massachusetts, 295 pp., 1974.

Spence, D. A. and D. L. Turcotte, Viscoelastic damping of cyclic displacements on the San Andreas fault, Proc. Roy. Soc. London A, submitted, 1978.

Thatcher, W., Strain accumulation and release mechanism of the 1906 San Francisco earthquake, J. Geophys. Res., 80, 4862-4872, 1975.

U.S. Geological Survey Staff, Surface faulting, in The San Fernando, California, Earthquake of February 9, 1971, U.S. Geol. Surv. Prof. Paper 733, 55-76, 1971.

Watts, A. B. and M. Talwani, Gravity anomalies seaward of deep-sea trenches and their tectonic implications, Geophys. J. Roy. Astron. Soc., 36, 58-90, 1974.

Table 1: Observed Versus Computed Postseismic Deformation for 1946 Nankaido earthquake

Location	Time Interval	Observed Vertical Movement (mm)	Computed (aseismic Slip + Asthenosphere Relaxation) mm
UWAJIMA	1947-50	-100	-70 (-50-20)
TOSASHIMIZU	1947-50	-60	-40 (-30-10)
URADO	1947-48	+300	+400 (+420-20)
TAKAMATSU	1947-50	-200	-50 (-40-10)
Aki-Muroto	1947-64	+350	+350 (+360-10)
Maximumum Relative Uplift, Muroto-Komatsujima Leveling	1947-64	+370	+220 (+250-30)

FIGURE CAPTIONS

Figure 1: (a) Coseismic vertical displacement ($t=0$, solid line) and subsequent viscoelastic response ($t=2\tau$, dashed line) due to slip on a fault rupturing three-quarters of an elastic plate of thickness H overlying a viscoelastic (Maxwell) half-space. Vertical uplift is normalized by the reverse fault slip, and distance perpendicular to the fault, y , is in multiples of the lithospheric thickness H . Fault dip is 30° .

(b) Similar to (a), but for buried slip extending from a depth of three-quarters of the lithospheric thickness to its base.

Figure 2: (a) Cumulative viscoelastic (post-slip) vertical displacement versus distance for successively longer times following sudden slip on a thrust fault that ruptures three-quarters of the lithospheric thickness (model of Figure 1a, top). Distance perpendicular to the fault, y , is normalized by the lithospheric thickness H , and vertical uplift is in cm per meter of fault slip. Time units are in terms of the asthenospheric relaxation time τ .

(b) Vertical displacement history at selected values of y/H for the same model as in (a) above. Offset at $t/\tau = 0$ is the coseismic displacement at each point.

Figure 3: Similar to Figure 2 but for a fault dipping 30° and rupturing the entire thickness of the lithosphere. In (b), note the break in scale of the vertical co-ordinate axis between +2 and +20 cm.

Figure 4: Location map of Japan, showing major plates and subduction boundaries. Shaded boxes outline two regions where vertical movement history through a major earthquake cycle is particularly well-documented. Larger box shows the region of southwest Japan affected by the 1944 Tonankai and 1946 Nankaido earthquakes, and the smaller shaded area outlines the South Kanto region, site of a great earthquake in 1923.

Figure 5: Area of the 1944 Tonankai ($M=8.0$) and 1946 Nankaido ($M=8.2$) earthquakes in southwest Japan. Rectangles show surface projections of inferred coseismic fault planes as determined by Ando (1975). The two western-most rectangles are for the 1946 earthquake, and the eastern one outlines the inferred 1944 fault plane. Tide gage stations (triangles, with names) are referred to in the text, and parts of their sealevel records are plotted in Figure 7. Dashed lines show leveling routes across eastern Shikoku and Kii Peninsula for which elevation change data are plotted in Figures 6-8.

Figure 6: Elevation changes in eastern Shikoku between Muroto Point and Komatsujima tide gage station. Komatsujima is assumed fixed in each time interval, and elevation changes are projected onto a profile perpendicular to the Nankai Trough. The 1929-1895 changes, typical of the interseismic secular deformation, are taken from Imamura (1930). The 1964-1947 and 1971-1964 level changes, taken from unpublished data of the Geographical Survey Institute, show postseismic uplift due to the 1946 earthquake.

Figure 7: Short-term postseismic changes related to 1946 Nankaido earthquake.

- (a) Monthly means of tide gage records for stations on Shikoku shown in Figure 5 (from Kawasumi, 1956). Estimates of total postseismic offset are indicated for each record. Note that land subsidence is measured positive upward on these sealevel records.
- (b) Elevation change of Muroto Point relative to Aki (locations in Figure 5) as determined from first-order leveling (data from Okada and Nagata (1953) and unpublished leveling, Geographical Survey Institute). Interseismic relative subsidence rate and corresponding rate during 1951-71 are shown for reference, as is the total 1947-64 elevation change between the two benchmarks.

Figure 8: Observed (top) and computed (bottom) profiles of inter-, co-, and postseismic level changes in eastern Shikoku related to the 1946 Nankaido earthquake. Two-dimensional model (middle) is drawn at the same scale as the graphs, and shows faulting in an elastic plate overlying a viscoelastic (Maxwell) halfspace. Coseismic model (Co) is a fault 60 km wide and infinitely long with 3.2 m of underthrusting. The interseismic profile (INTER) is the asthenospheric response to this sudden slip during the interval $t=\tau$ and $t=2\tau$ after the earthquake ($\tau=2\eta/\mu$ is the asthenospheric relaxation time), added to the effect of 30 mm/yr of steady aseismic slip at the bottom of the plate. Postseismic model (POST) shows the elastic response to 2.5 m of aseismic underthrusting on 30 km

of the downdip extension of the coseismic rupture, along with the asthenospheric relaxation expected to occur during 1947-64 as a result of the earthquake. See text for further details.

Figure 9: Computed postseismic uplift in southwest Japan. Solid rectangles are surface projections of faults that are downdip extensions of Ando's (1974) two coseismic fault planes (dashed rectangles). Tops of postseismic faults are at 30 km depth and faults are 30 km wide, 150 km long, dip 30° NNW and each have 2 m of reverse dip-slip and 1 m of right-lateral strike-slip movement across them. Contour interval is 100 mm except for addition of -50 mm contour. Solid triangles show tide gage stations of Figure 7a (UW = Uwajima, TS = Tosashimizu, UR = Urado, TK = Takamatsu), and solid squares denote endpoints of leveling routes (A = Aki, M = Muroto, K = Komatsujima).

Figure 10: South Kanto region, showing the coseismic fault plane of the 1923 earthquake determined by Ando (1971). Leveling routes from Tokyo to Atami and on the eastern side of the Boso Peninsula are shown dashed, and data from these lines are plotted in Figures 11 and 12. Open diamond symbol shows the location of the chained rhombus at Mitaka, for which shear strain changes are plotted in Figure 13, and triangles locate ABURATSUBO and MERA tide gage stations, whose sealevel records are also shown in Figure 13.

Figure 11: Elevation changes from a leveling route on eastern side of the Boso peninsula. Data are projected onto a profile perpendicular to the Sagami Trough, which is the zero point for the profile (data from Scholz and Kato, 1978). Solid and dashed lines show fit to the

data of a model described in the text. INTER data for 1931-65 have been normalized to a 24-year interval (1939-63) in order to make them comparable with the Tokyo-Atami level changes during this time that are plotted in Figure 12.

Figure 12: Similar to Figure 11, but for the Tokyo-Atami leveling route. Zero point on the profile is the Kozu-Matsuda fault, the presumed landward extension of the Sagami Trough. Open circles on the INTER profile denote level changes affected by non-tectonic artificially induced subsidence in the Tokyo area.

Figure 13: Short-term postseismic changes related to the 1923 Kanto earthquake.

(a) Shear strain changes versus time determined from chained length measurements of the Mitaka rhombus (from Fujita, 1972; Scholz and Kato, 1978). Arbitrary zero level is taken in 1923 immediately before the earthquake. Shear strain is then calculated as the difference between the greatest and the least principal horizontal strains e_1 and e_3 referenced to this zero level.

(b) Yearly mean sealevel at ABURATSUBO and MERA tide gage stations (Coastal Movements Data Center, 1976). Dashed lines show the predicted deformation for a model of asthenospheric relaxation used to fit the leveling data of Figures 11 and 12 and described in the text.

Figure 14: Computed postseismic uplift contours and horizontal strain changes (crosses with numbers) in the South Kanto district. Contour interval is 20 mm. Principal strain axes, with the most compressive

axis shown with arrowheads, are plotted near the location of the Mitaka rhombus, and maximum shear strain changes, in μ strain are given in each case. Circled value is at Mitaka. Solid rectangle is the surface projection of the assumed postseismic fault, which is the downdip extension of Ando's (1971) coseismic fault plane (dashed rectangle). Top of postseismic fault is at 30 km depth, and this fault is 20 km wide, 130 km long, dips 45° NE and has 0.6 m of right-lateral strike-slip and 0.3 m of reverse dip-slip movement across it. Solid squares and dashed lines show leveling routes, and solid triangles locate ABURATSUBO and MERA tide gage stations.

Figure 15: Two-dimensional elastic half-space model for the deformation cycle at convergent plate boundaries. Uplift in the CO, POST, and INTER phases of the deformation are each normalized for 1 m of fault slip. The sum of the three curves gives a block offset of 1 m (50 cm vertical displacement) across the fault. Arrows on the fault segments give the relative proportions of slip on the hanging wall (continental) and footwall (oceanic) blocks of the fault as they are prescribed by the dislocation model. Arbitrarily partitioning the slip differently between the two blocks will alter the zero levels of the model curves but leave their shapes unchanged.

Figure 1

Figure 2

Figure 3
815

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

a)

b)

Figure 13
825

Figure 15

SPACE-TIME MIGRATION OF EARTHQUAKES
ALONG THE NORTH ANATOLIAN FAULT
ZONE AND SEISMIC GAPS

M.N. Toksöz, A.F. Shakal and A.J. Michael

Department of Earth and Planetary Sciences
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Abstract

The North Anatolian fault is a well defined tectonic feature extending for 1000 km across northern Turkey. The space-time distribution of seismicity and faulting of this zone has been examined with a particular emphasis on the identification of possible seismic gaps. Results suggest several conclusions with respect to the temporal and spatial distribution of seismicity. First, the earthquake activity appears not to be stationary over time. Periods of high activity in 1850-1900 and 1940 to the present bracket a period of relatively low activity in 1910-1939. Second, there appears to have been a two-directional migration of earthquake epicenters away from a central region located at about 39°E . The migration to the west has a higher velocity than the migration to the east. The faulting associated with successive earthquakes generally starts where the previous rupture ended. Some existing gaps were filled by later earthquakes.

At present there are two possible seismic gaps along the North Anatolian fault zone. One is at the western end of the fault, from about 29 to 30°E . Unless this is a region of ongoing aseismic creep, it could be the site of a magnitude 6 or greater earthquake. The other possible gap is at the eastern end, from about 42 to 43°E , to the west of the unexpected $M = 7.3$ event of 24 November 1976.

INTRODUCTION

The North Anatolian fault zone is a major tectonic feature with a well-defined fault trace and an established history of seismicity (Allen, 1969, 1975; Ketin, 1968, 1976; Ambraseys, 1970; Seymen and Aydin, 1972; Sengor, 1978; Dewey, 1976). Most intermediate and large magnitude earthquakes occurring on this fault have produced surface breaks (Ambraseys and Zatopek, 1969; Ambraseys, 1970; Arpat et al., 1977; Ketin, 1976; Toksöz et al., 1977; Wallace, 1968). Furthermore, because of topography and water resources, the fault zone is a relatively densely populated, agricultural area dotted by many villages and small towns. Even in historic times, the fault dimensions could be estimated from geographic distribution of damage and isoseismals. In this study, we have examined the space-time distribution of earthquakes on the North Anatolian fault zone with a particular emphasis on seismicity gaps. Although we have examined historic catalogs, the primary data we analyzed in this study cover the period 1910-1977. The earthquake catalogs for this period appear to be reasonably complete and field data are more readily available.

The previous studies (Dewey, 1976; Allen, 1975; Ambraseys and Zatopek, 1969; Canitez and Toksöz, 1978) have noted the migration of the earthquake epicenters along the fault zone. Trends that were noticed included eastward migration, westward migration, and alternating occurrence of the earthquakes

between the east and the west. Dewey (1976) carried out an extensive study and identified a clear westward migration. He further sought some physical mechanism for such migration.

In this paper we look at both the eastward and westward trends of earthquake migration along the North Anatolian fault. There are indications of two possible gaps which require further study.

SEISMICITY OF TURKEY AND THE NORTH ANATOLIAN FAULT ZONE

The seismicity of Turkey is controlled by a complicated interaction of several plates. Figure 1 is a seismotectonic map showing some major fault zones and earthquake epicenters. In the southeast the Arabian plate collides against the Anatolian and Asian plates. In the southwest, the Mediterranean plate is subducting under the Anatolian and Aegean plates. In western Turkey, seismicity occurs along a series of east-west trending normal faults. This region consists of a series of horsts and grabens.

In the north, the North Anatolian fault zone is a major boundary between the Anatolian and Asian plates. The sense of the motion is right handed and the slip rate is relatively high, ranging from 3.2 to 11 cm/yr (Brune, 1968; North, 1974). Our estimate of the slip rate is 6 cm/yr averaged for the period 1910-1977, and 12 cm/yr for the period 1939-1977.

The East Anatolian fault zone appears to be a left-handed

conjugate fault to the North Anatolian fault (Arpat and Saroglu, 1972; McKenzie, 1972). It has a much lower slip rate and fewer earthquakes. Butting against this from the south is the north-south trending rift zone through Hatay and Maras. Although the tectonics of the whole region are complex, the North Anatolian fault zone can be distinguished as a separate feature, and the associated seismicity can be clearly identified. Thus, our present study is confined only to the North Anatolian fault zone.

In identifying the seismicity pattern of the region it is important to have data over as long a time period as possible. There are earthquake catalogs for Turkey dating back 2000 years (Ergin et al., 1967). However, it is not clear how complete the list of historic earthquakes is and how accurate the intensities are. Generally, population distributions, education and political climate affect the historic data sets.

In this study we examined several catalogs: Ergin et al., 1967; Karnik, 1968; Gutenberg and Richter, 1949; Alsan et al., 1975 and the U.S.G.S.-N.E.I.S. catalogs. Looking at the seismicity and average seismic energy release per year, it is clear that the historic records are very incomplete until about the year 1500. For the following four hundred years, the data base improved. In Figure 2, seismic energy release is plotted as a function of time. The energy E_s was calculated using the empirical relationship (Båth, 1973)

$$\log E_s = 12.24 + 1.44M \quad (1)$$

with intensity to magnitude conversions taken from the I.T.U. data (Ergin et al., 1967), for the pre-instrumental period. The incompleteness of data is seen from a step jump in 1840 on the energy release curve shown in Figure 2. After that period, the data base becomes more complete for western Turkey but not for eastern Turkey.

The reliable data base for the North Anatolian fault zone starts at about 1910. During this period the whole region was reasonably well populated. Both observers' descriptions of local effects and instrumental data are available for many events. In this study we limit our analysis to the period 1910-1977.

The largest event in the seismic record is the $M = 7.9$ event of 26 December 1939. The 20 years prior to this event was a period of low activity. This is apparent in the strain release graph of Figure 2 and also in Figure 3, which shows the number of events in individual magnitude ranges. Although the estimation of a threshold of data completeness is difficult, Alsan et al. (1975) suggest that the catalog is complete above $M \sim 5.5$ for the entire 1913-1970 interval, but there is no quantitative confirmation of this. The low number of $M \geq 6$ events in the 20 or 30 years prior to the sequence of large events starting in 1939 is therefore probably real and may represent a quiescent period similar to those described by Mogi (1968).

SPACE-TIME MIGRATION OF EARTHQUAKES

The North Anatolian fault zone is basically a linear feature with a small curvature in the middle. Thus the spatial distribution of seismicity can be analyzed as an east-west one-dimensional model. In this study we investigate the space-time distribution of the earthquakes along the trend of the fault in a manner similar to that of Dewey (1976). Since the record of small earthquakes is not complete, we include only earthquakes of $M_s \geq 5.9$ (intensity $\geq$ VII). These are listed in Table 1.

In Figure 4, the distribution of epicenters along the east-west trend of the fault versus time is shown. There appear to be two trends, one to the west with an average slope of 0.02 yr/km and another to the east with 0.1 yr/km. It is clear that the westward migration is faster immediately after the 1939 earthquake and slows down after 1945. The average velocities are about 50 km/yr to the west and 10 km/yr for eastward migration when averaged over a 38 year time span.

A better illustration of the migrations and their relevance to seismic gaps can be illustrated by plotting the length of observed surface faulting for each earthquake. In most cases there are surface breaks associated with earthquakes. For the larger events these have been mapped. For other events the length of the highest intensity contour gives some measure of the fault length. We have used these criteria to

plot fault length vs magnitude for the North Anatolian fault zone earthquakes (Figure 5). Fault length data are tabulated in Table 1. An empirical equation fitting these data gives

$$\log L = 0.78 M_S - 3.62 \quad (2)$$

where L = fault length, M_S = surface wave magnitude. This relationship is similar to those obtained for California earthquakes (Smith and Van de Lindt, 1969; Canitez and Toksöz, 1978). When information for actual surface breaks was not available, we used equation (2) to estimate the fault lengths.

The distribution of surface faulting is shown on the maps in Figures 6a,b,c for three different periods. In these and subsequent figures, fault breaks are plotted as they were mapped in the field, when such information was available. In other cases damage scales and isoseismals were used to fix the fault trace. (Ergin et al. (1967) give a brief description of these.) For smaller events, where neither of these data are generally available, the surface break is assumed to be parallel to the main fault zone with the epicenter in the middle. Since the fault length of these events is generally small (~ 20 km), any error that may be introduced is small.

In Figures 6a,b,c the time periods covered are 1910-1937, 1938-1977 and 1910-1977, respectively. As seen in Figure 3 and in Table 1, there were relatively few large events between 1910 and 1938. The largest earthquake in recent history

($M = 7.9$) occurred on 26 December 1939. This appears to have been the first in a series of intermediate and large earthquakes. The fault breaks associated with these events cover almost the entire length of the North Anatolian fault zone. There are very few gaps and relatively little overlap.

The same phenomenon is better illustrated in the simplified diagram in Figure 7. Fault breaks are plotted along an east-west line as a function of time. Starting with the 1939 ($M = 7.9$) event there is a very rapid propagation of earthquakes to the west. In three successive events, rupture occurs along about a 500 km segment of the fault (velocity ~ 100 km/yr). Afterward there is a continued progression westward, but at a slower rate. There is an obvious filling of a gap at 31°E longitude by an earthquake occurring 15 years later. There is a 60 km gap at about 30°E longitude. This will be discussed later and we consider it a potential site for monitoring.

The progression of the earthquakes eastward from $\lambda = 40^\circ\text{E}$ appears to be at a much slower rate (about 10 km/yr). Complete rupture had occurred between $\lambda = 40-42^\circ\text{E}$ in about 30 years. Then, there was a quiescent period until 24 November 1976, when a magnitude $M_S = 7.3$ earthquake occurred further east. The location of this event (see Figure 6) is somewhat displaced to the north from a linear extrapolation of the North Anatolian

fault zone. However, the right-handed strike-slip faulting is consistent with the North Anatolian events (Toksöz et al., 1977; Arpat et al., 1977). Thus on the eastern side, the North Anatolian fault may have been segmented northward through a set of en-echelon steps. Recent mapping has indicated the presence of geologically active strike-slip faults to the west of the 1976 earthquake. Thus the region between $\lambda = 42^{\circ}$ - 43° E longitude probably represents another seismic gap. However, more work needs to be done in these areas to understand the nature of this gap.

DISCUSSION OF POSSIBLE SEISMIC GAPS

Our study of the distribution of earthquakes and surface breaks is based primarily on local observations. However, because of the uniformity of the mechanisms and simplicity of tectonics of the North Anatolian fault zone, it is feasible to combine teleseismic information and field observations to understand the space-time distributions of earthquakes.

The general patterns of the seismicity include:

1. Earthquake occurrences are not stationary over time. There has been relatively high activity between 1850-1900, a quiet period from 1910-1939 and an active period from 1940 to the present.

2. There appears to be a two direction migration of earthquake epicenters from a central region located at about

$\lambda=39^\circ$ longitude. The westward migration is faster (50 km/yr). This is very similar to the 80 km/yr velocity obtained by Dewey (1976). Eastward migration is slower.

3. The faulting associated with successive earthquakes generally seems to start from where the previous rupture has ended. Some existing gaps have been filled by later earthquakes.

4. At present there are two possible earthquake gaps on the North Anatolian fault zone. The gap in the west between 29° and 30°E longitudes is shown in more detail in Figure 8. There have been two large earthquakes to the east and west of this gap in 1963 and 1967. Neither the local reports nor seismicity catalogs indicate earthquakes of $M \geq 5.0$ in this area. Unless this zone is deforming by creep or some other aseismic means, it should be considered a gap with a potential earthquake of magnitude 6 or greater.

The second gap is along the eastern side of the North Anatolian fault zone between 42° and 43°E longitudes. The importance of this zone really emerged after the 24 November 1976 earthquake ($M_S = 7.3$) in an unsuspected area. There is no uniformity of opinion on the eastward extension of the North Anatolian fault zone. The 1976 earthquake seems to indicate a possible northward translation of the fault zone. If this indeed is the case then the identified gap is a major one. Only recently geologically active faults have been mapped in this area. In eastern Turkey, the fault lengths

corresponding to the magnitude of the events appear to be short $L \approx 20$ km for the 1966 Varto earthquake with $M_S = 6.7$ (Wallace, 1968) and $L = 55$ km for the Van earthquake of 1976 with $M_S = 7.3$ (Toksöz et al., 1977). Thus the dimensions of the gap have the potential for a large earthquake ($M \geq 7$) in this area. However, because of the limited knowledge of geology, tectonics and seismicity of this area, this should be considered a speculative assessment.

It is necessary to undertake some detailed studies to determine whether these sites are indeed seismicity gaps. Monitoring of microseismicity, looking for evidence of creep or aseismic strain release, a very careful analysis of local earthquake catalogs and interviews with people to determine the existence or lack of earthquakes in these "gaps" are necessary.

References

- Allen, C.R., Transcurrent faults in continental areas, Phil. Trans. R. Soc. Lond., A258, 82-89, 1965.
- Allen, C.R., Active faulting in northern Turkey, Contr. No. 1577, Div. Geol. Sci., Calif. Inst. Tech., 32, 1969.
- Allen, C.R., Geological criteria for evaluating seismicity, Bull. Geol. Soc. Am., 86, 1041-1057, 1975.
- Alsan, E., L. Tezucan and M. Båth, An earthquake catalog for Turkey for the interval 1913-1970, Rept. 7-75, Kandilli Obs., Istanbul and Seismological Inst., Uppsala, 1975.
- Alsan, E., L. Tezucan and M. Båth, An earthquake catalog for Turkey for the interval 1913-1970, Tectonophysics, 31, 13-20, 1976.
- Ambraseys, N.N., Some characteristic features of the North Anatolian fault zone, Tectonophysics, 9, 143-165, 1970.
- Ambraseys, N.N. and A. Zatopek, The Mudurnu Valley, West Anatolian, Turkey, earthquake of 22 July 1967, Bull. Seism. Soc. Am., 59, 521-589, 1969.
- Arpat, E. and F. Saroglu, The East Anatolian fault system: Thoughts on its development, Bull. Mineral Res. Expl. Inst. Turkey, 78, 33-39, 1972.
- Arpat, E., F. Saroglu and H.B. Iz, 1976 Caldiran earthquake Yeryuvarive İnsan, 2, 29-41, 1977.
- Båth, M., Introduction to Seismology, John Wiley and Sons, New York, 1973.

- Brune, J.N., Seismic moment, seismicity and rate of slip along major fault zones, *J. Geophys. Res.*, 73, 777-784, 1968.
- Canitez, N. and M.N. Toksöz, Strain release and earthquake occurrences along the North Anatolian fault zone, *Tectonophysics*, in press, 1978.
- Dewey, J.W., Seismicity of Northern Anatolia, *Bull. Seism. Soc. Am.*, 66, 843-868, 1976.
- Ergin, K., U. Guolu and Z. Uz, A catalog of earthquakes for Turkey and surrounding areas, *Tech. Univ. Istanbul Mining Eng. Fac.*, Publ. No. 24, 74 pp, 1967.
- Gutenberg, B. and C.F. Richter, Seismicity of the Earth and Associated Phenomena, Princeton University Press, 1949.
- Karnik, V., Seismicity of the European Area, Part 1, Academic Praha, Publishing House of the Czechoslovak Academy of Sciences, 1968.
- Ketin, I., Relation between general tectonic features and the main earthquake region of Turkey, *Bull. Mineral Res. Expl. Inst.*, 71, 63-67, 1968.
- Ketin, I., On the North Anatolian fault, *Bull. Mineral Res. Expl. Inst.*, 72, 1-26, 1969.
- Ketin, I., A comparison between San Andreas and N. Anatolian faults, *Bull. Turkish Geol. Soc.*, 19, 149-154, (in Turkish), 1976.
- McKenzie, D.P., Active tectonics of the Mediterranean region, *Geophys. J. R. astr. Soc.*, 30, 109-185, 1972.

- Mogi, K., Migration of seismic activity, Bull. Earthquake Res. Inst., Tokyo Univ., 46, 53-74, 1968.
- North, R.C., Seismic moment, source dimension and stresses associated with earthquakes in the Mediterranean and Middle East, Geophys. J. R. astr. Soc., 48, 137-161, 1977.
- North, R.C., Seismic slip rate in the Mediterranean and Middle East, Nature, 252, 560-563, 1974.
- Richter, C.F., Elementary Seismology, W.H. Freeman and Co., San Francisco, 1958.
- Sengor, A.M.C., The North Anatolian fault: its age, offset and tectonic significance, to be published, 1978.
- Seymen, I. and A. Aydin, The Bingöl earthquake fault and its relation to the North Anatolian fault zone, Bull. Mineral Res. Expl. Inst. Turkey, 79, 1-8, 1972.
- Smith, S.W. and W. Van de Lindt, Strain adjustments associated with earthquakes in Southern California, Bull. Seism. Soc. Am., 54, 1569-1589, 1969.
- Tatar, Y., Tectonic structures along the North Anatolian fault zone, northeast of Refalize (Erzincan), Tectonophysics, 29, 401-410, 1975.
- Toksöz, M.N., E. Arpat and F. Saroglu, East Anatolian earthquake of 24 November 1976, Nature, 270, 423-425, 1977.
- Wallace, R.E., Earthquake of August 19, 1966, Varto area, Eastern Turkey, Bull. Seism. Soc. Am., 58, 11-45, 1968.

List of Figures

Figure 1 - Seismotectonic map of Turkey showing earthquake epicenters and the major faults.

Figure 2 - Seismic energy release along the Anatolian fault zone between 1500-1977, averaged over 20 year segments. Estimated energy is subject to errors in magnitude and intensity to energy conversion, and completeness of the catalogs for the historic data. The 26 December 1939 $M = 7.9$ event has been included as a 1940 event to more clearly show the low activity level of the preceding 20 years.

Figure 3 - Number of earthquakes per year with magnitudes $M \geq 5$, 6 and 7 in the 1913-1975 time period.

Figure 4 - Time-space distribution of earthquake epicenters along the North Anatolian fault zone in the period 1910-1977. The origin corresponds to year 1940, and the epicenter (39.8°N and 39.7°E) of the 26 December 1939 earthquake. Numbers identifying earthquakes are the same as those given in Table 1.

Figure 5 - Fault length vs magnitude for events of the Anatolian fault system, including fault length estimates from field studies (circles) and estimates from isoseismals (triangles). The line through these points ($\log L = .78M - 3.62$) is only slightly different from the relationship found for California earthquakes

($\log L = .59M - 2.24$) by Smith and Van de Lindt (1969).

Figure 6 - Earthquake fault breaks along the North Anatolian fault zone. Fault trace is shown as a light line, fault ruptures as heavy straight lines. Three figures correspond to three time periods:

(6a) 1910-1937

(6b) 1938-1977

(6c) 1910-1977

Figure 7 - Space-time distribution of earthquake faulting along the North Anatolian fault zone. Fault breaks are projected along an east-west line. Length of each line corresponds to the length of the observed break or the length estimated using eq. 2. The earthquakes are identified by the numbers listed in Table 1. Note the two possible gaps near the eastern and western end of the diagram.

Figure 8 - Earthquake epicenters and observed faulting along the west end of the North Anatolian fault zone. Note the possible gap to the west of the 22 July 1967 earthquake (#31) fault break. It is not known whether aseismic slip occurs in this area, or this is a locked zone and a possible site of a future earthquake.

Table 1. List of earthquakes along the North Anatolian Fault Zone used in this study.

Number	Date			Time Hr:Min	Epicenter		M	Fault Length
	Yr	Mth	Day		Lat	Lon		
1	1910	6	25	19:26	41.00	34.00	6.2	
2	1916	1	24	6:56	40.27	36.83	7.1	
3	1923	4	29	9:38	40.07	36.43	5.9	
4	1924	9	13	14:35	39.96	41.94	6.8	
5	1928	5	2	21:57	39.64	29.14	6.5	
6	1929	5	18	6:42	40.20	37.90	6.5	
7	1934	11	27		37.90	40.20	6.0	
8	1939	11	21	8:49	39.82	39.71	5.9	
9	1939	12	26	23:59	39.80	39.51	7.9	350
10	1941	11	12	10:09	39.74	39.43	5.9	
11	1942	12	11	02:40	40.76	34.83	6.1	
12	1942	12	20	14:03	40.87	36.47	7.0	50
13	1943	6	20	15:37	40.85	30.51	6.6	30
14	1943	11	26	22:24	41.05	33.72	7.4	280
15	1944	2	1	3:25	41.41	32.69	7.3	160
16	1945	10	26	13:56	40.90	33.29	6.0	45*
17	1946	5	31	3:12	39.29	41.21	5.9	
18	1949	8	17		39.54	40.57	6.8	75
19	1951	8	13	18:36	40.88	32.87	6.9	60*
20	1953	9	7	3:59	41.09	33.01	6.4	60*
21	1954	10	24		40.00	40.00	6.0	
22	1956	2	20	20:35	39.89	30.49	6.4	
23	1957	5	26	6:36	40.67	31.00	7.1	50*
24	1957	5	26	9:39	40.76	30.81	5.9	
25	1960	1	26	9:53	40.19	38.75	5.9	
26	1963	9	18	16:59	40.77	29.12	6.3	35*
27	1964	6	14	12:18	38.13	38.51	6.0	40*
28	1964	10	6	14:33	40.30	28.23	7.0	60*
29	1966	8	19	12:23	39.17	41.56	6.7	20
30	1966	8	20	12:05	39.16	40.70	6.1	
31	1967	7	22	16:56	40.67	30.69	7.2	
32	1967	7	26	18:53	39.54	40.33	6.2	
33	1968	9	3	8:24	41.81	32.39	6.5	
34	1976	11	24	12:22	39.10	44.02	7.3	50

*designates fault length estimated from isoseismals

Fig. 1

ANATOLIAN FAULT ZONE SEISMIC ENERGY RELEASE

SEISMIC ENERGY PER 20 YRS (ERGS)

10^{24}

10^{23}

10^{22}

10^{21}

10^{20}

1500

1600

1700

1800

1900

-8

-

-7

EQUIV

-6

-

20

ANATOLIAN FAULT ZONE

Fig. 3

Fig. 4

1910 - 1937

Fig. 6a

1938 - 1977

Fig. 6b

1910 - 1977

FAULT BREAKAGE ALONG NORTH ANATOLIAN FAULT

Fig. 7

PATTERNS OF FAULTING AND SEISMIC GAPS
IN THE GREAT BASIN PROVINCE

by

Robert E. Wallace

U.S. Geological Survey
345 Middlefield Road
Menlo Park, CA 94025

Abstract

Patterns and rates of faulting provide a basis for estimating long-term seismicity in the Great Basin province. A conspicuous zone of large historic earthquakes accompanied by surface faulting trends northward through central Nevada and extends beyond the province to the south. Zones of relatively high long-term prehistoric seismic activity along the eastern and western margins of the province are indicated by extensive young faulting. A third broad zone of young faulting through central Nevada brackets the narrower zone of historic activity. In much of eastern Nevada and the westernmost part of Utah young fault scarps are relatively fewer and smaller, suggesting that the area is less seismic.

Estimated rates of occurrence of earthquakes greater than magnitude 7 range from an average of one event every 22 years in the zone of historic events to an average over the last 12,000+ years of one event every 100,000 years per 1000 km² in a relatively active zone of north-central Nevada, to almost no large events in 500,000 years in the quiet zones.

Three seismic gaps exist: the Stillwater, White Mountain, and the southern Sierra areas. Estimates of the average recurrence of M7 events in these gaps range at least from one event within a few tens of years to one event within a few thousand years.

Introduction

The intent of this paper is to assess the seismic gaps in the Great Basin province. The Great Basin (see fig. 1) is a subdivision of the Basin and Range province and encompasses most of Nevada, western Utah, and eastern California, plus small parts of adjacent States.

A method based on fault scarp analysis is used to evaluate the history of seismicity over the last several hundred thousand years. The term "seismic gap" implies an orderly pattern of seismic events both spatially and temporarily. The term also carries the connotation that ruptures may be expected to propagate between two areas previously ruptured. The term "young fault scarps" is used in this paper to refer to scarps that have pronounced topographic relief preserved from one or a few displacement events, presumably accompanying earthquakes. The scarps are generally preserved in colluvium or alluvium. Data on young fault scarps are still so fragmentary that the following patterns and estimates of rates of faulting are extremely crude. But young fault scarps represent a potentially rich source of data about large prehistoric earthquakes.

Method

This attempt to identify and evaluate seismic gaps is based on fault scarps produced during seismic events in prehistoric as well as historic time. The oldest of the "young" fault scarps considered in the analysis is believed

to be several hundred thousand years old. Although the fault scarps become eroded and more obscure with age, scarps less than 12,000 years old can generally be recognized to be related to single seismic events. Older scarps commonly represent multiple seismic events, and although they provide less specific data than younger scarps, they are nevertheless evidence of seismicity and indicate the general size and distribution of earthquakes.

Relating scarps to the history of seismicity requires dating of the scarps and mapping the distribution of scarps of different ages. For this study the ages of scarps are based primarily on scarp morphology. For example, older scarps have lower angles of slope and more rounded crests (Wallace, 1977b). Bucknam and Anderson (1978) find that a scarp height-slope angle relationship provides additional evidence of the age of a scarp.

The Basin and Range province is especially favorable for this approach inasmuch as scarps are preserved in the desert climate for thousands of years, fault movement generally has a large component of vertical displacement creating topographic relief, and the faulting events are spatially discrete because they tend to occur along range-block boundaries.

After individual seismic events or clusters of events have been identified by the age and distribution of scarps produced, then the dimensions of the scarps, their heights and lengths, provide a basis for estimating the size of the earthquakes. Methods involving seismic moment and local magnitude relations (Thatcher and Hanks, 1973; Hanks, Hileman, and Thatcher, 1975) are useful in this regard.

Patterns of Faulting

Large seismic events ($>M7$) accompanied by surface faulting in the Great Basin province have been concentrated along the eastern and western margins of the province and in a north-trending belt 200+ km wide in the western half of the province. The largest historic events in the $M7-8$ range occurred in 1872, 1915, 1932, and 1954, and an event in 1952 occurred along the same trend but outside the province to the southwest. All of these earthquakes were accompanied by extensive surface faulting.

Smaller earthquakes ($M6-7$) also accompanied by minor to moderate surface faulting occurred in 1869?, 1932, 1950, and twice in 1934.

A longer term pattern can be determined from an analysis of prehistoric surface faulting. Figure 1 shows the pattern of occurrence of larger earthquakes, based both on surface faulting and seismicity.

The province is divided into three main zones: a) zones of historic surface faulting, b) zones in which large faulting events occurred during the Holocene (last 12,000+ years), and c) zones in which young fault scarps are absent or are so short that the events probably did not produce earthquakes of $M7-8$ during the last 500,000 years. Large events may have occurred earlier, but that has not been confirmed.

Both the eastern and western margins of the province have relatively high seismicity, and prominent fault scarps of Holocene age are present. A similar zone 200+ km wide trends northward through central Nevada. Although the central Nevada zone is referred to as a separate zone, it merges with the eastern Sierra zone in the southern part of the province and is separated from the eastern Sierra zone only by an indistinct zone of lower seismicity in the northern part of the province.

A narrower belt 10-30 km wide which contains the major historic events lies within the central Nevada zone, and extends southwestward, crossing the Great Basin province boundary at least to the region of the 1952 event at Arvin-Tehachapi, California. This belt has been termed the Ventura-Winnemucca zone by Ryall and others (1966).

The pattern depicted in figure 1 is based in part on maps of young faults by Howard and others (1978) and Slemmons (1967), plus consultation with J. H. Stewart, R. E. Anderson, and R. Bucknam. Unfortunately, existing data about young faults are grossly inadequate, and the boundary between areas in which large earthquakes ($>M7$) occurred in Holocene time and those in which only smaller earthquakes occurred may be incorrect in detail. In addition, scarps formed in Holocene time have been distinguished from those formed in the earlier part of late Quaternary time (last 500,000+ years) in relatively few places. The rate of faulting indicated for the entire area of Holocene faulting is estimated from the area of figure 2. Despite the inadequacy of the data, they do indicate a difference between western Nevada-eastern California and eastern Nevada-western Utah. The Rocky Mountains zone has similarities to the western Nevada-eastern California region, but present data suggest a somewhat lower abundance of larger Holocene scarps.

A more detailed study in north-central Nevada (fig. 2) provides a sample of the pattern of development of individual scarps or clusters of scarps in time and space. The pattern observed is one of a rather random occurrence of events. This is surprising in that some coupling mechanism between most of the ranges in the area appears to exist. Most ranges shown in figure 2 have been progressively tilted dominantly eastward for approximately the last 10-14 million years. This tilting might be expected to be accompanied by a more regular pattern of earthquakes. Individual clusters of scarps, such as on the west flank of the Humboldt Range or the west flank of the Tobin Range, display repeated movement during periods of time when other clusters of scarps nearby were inactive. For example, the faults on the west flank of the East Range have remained inactive for possibly several tens of thousands of years, while the faults along the Tobin and Humboldt Ranges, which bracket the East Range, were repeatedly active.

On the time scale of a few hundred thousand years, no pattern of migration of events across the whole province is yet apparent, although data are still too fragmentary to rule out such a pattern.

Rates of Faulting and Recurrence of Large Earthquakes

Various approaches can be used to estimate the rates of faulting and the recurrence of large earthquakes in the region. Ryall and others (1966) analyzed the historic record and normalized results according to events per year per 1000 km². An extension of their work was carried out by Douglas and Ryall (1975). A preliminary analysis of prehistoric fault scarps by Wallace (1977a) provides another approach. Several approaches and the values obtained are given below.

Approach A - Between the large historic earthquakes of 1872, 1915, 1932, 1952, and 1954 are four intervals of 47, 17, 22, and 2 years, or an average of 22 years. If the zone of these earthquakes is considered to be 650 x 30 km or 19,500 km², this rate can be expressed as 2.3×10^{-3} events per year per 1000 km².

Approach B - In north-central Nevada Wallace (1977a) found no more than seven large events of Holocene (<12,000+ years) in an area of 17,000 km². This represents a rate of 3.4×10^{-5} events per year per 1000 km².

Approach C - Rates of uplift of range blocks provide another approach. R. C. Speed (written commun., 1978) reports that the White Mountains scarp represents a rate of uplift of 0.8 m per 1000 years. Similarly, I estimate that the steep eastern flank of the Stillwater Range may represent an uplift rate of as much as 0.5 m per 1000 years. Considering that an M7 event would involve an area of approximately 1000 km² and 3 m uplift per event, the rates for the White Mountains and Stillwater areas are, respectively, 2.7 and 1.6×10^{-4} events per year per 1000 km².

Approach D - For the Ventura-Winnemucca zone, the historic record prior to 1962, as analyzed by Ryall and others (1966), indicates a rate of 2×10^{-3} per year per 1000 km² (Douglas and Ryall, 1975). Ryall and Douglas (written commun., 1976) added data for 1962-1968 and normalized the seismic data over different areas. By using seismic data from a variety of sources, an M of about 7.3 and an area of 150,000 km² for the western Nevada active zone, they derived a value of 2×10^{-4} per year per 1000 km². By using only their own data for 900+ earthquakes, Ryall and Douglas derived a recurrence rate as low as 7×10^{-5} events per year per 1000 km², where an event is considered to have a magnitude of 7.3.

The widely different rates derived by different approaches reflect the inadequacy of data, nevertheless, it is instructive to begin to assign tentative values to different regions of the province. Ranges of rates thus are shown in figure 1.

For the main zone of historic faulting, the values for numbers of events per year per 1000 km² range from 2.3×10^{-3} (approach A) to 2×10^{-3} (approach B) to 1.6×10^{-4} (approach C). In figure 1 this range is bracketed by 3×10^{-3} to 1×10^{-4} .

For the zones of Holocene faulting, the value of 3.4×10^{-5} events per year per 1000 km² (approach B) is the only value derived from Holocene data.

Inasmuch as the density of faults large enough to produce earthquakes $>M7$ appears to be as great in north-central Nevada as anywhere in the province, this value may be considered to be near the upper limit of values suitable for the entire region. For example, in the Rocky Mountains zone photo-geologic reconnaissance by Slemmons (1967) suggests a much lower density of large fault scarps. Values between 5×10^{-5} and 5×10^{-6} events per year per 1000 km^2 would seem to bracket the likely rates in the zones shown as Holocene faulting in figure 1.

The area of eastern Nevada and western Utah is characterized by fault scarps of Late Quaternary age (last 500,000 years) that are relatively few kilometers long, suggesting few or no large seismic events in that period of time. Whether large events occurred before 500,000 years ago cannot be stated from the evidence at hand.

Despite the wide discrepancies in rates derived by different approaches, a strong suggestion emerges that during the last century the rate was substantially greater than the long-term rate. The seismic gaps do not appear to have more young scarps than the region of Holocene faulting as a whole. Nor have scarps been recognized yet in the areas of the historic events to suggest surface faulting at a greater rate than in surrounding areas. If more were known about the history of fault scarps in the region, it might be possible to document greater rates in localized areas. The report by Thompson and Burke (1973) is suggestive. They say, "The shoreline deposits (12,000+ years old in Dixie Valley) are offset by faults at many places . . . the vertical displacement is 9 m., and it may have occurred in two or more episodes of faulting." As such data are accumulated, a much more reliable value for the rates of faulting in prehistoric time should be possible.

Seismic Gaps

Ryall and others (1966) documented the existence of gaps in the seismicity of the region for the historic period, and indicated the tendency for the gaps to be filled by successive large earthquakes. The record of fault events in prehistoric time supplements this interpretation. Some of the seismically quiet zones which might be interpreted as gaps are interpreted as relatively stable blocks in which large seismic events are considered to be unlikely.

Stillwater gap.--Between the southern end of the 1915 Pleasant Valley scarp and the northern end of the 1954 Dixie Valley scarp is a segment of fault-generated range front which has not broken in historic time. The gap is about 40 km long, and includes the precipitous east flank of the northern end of the Stillwater Range.

This precipitous segment of range front is about 20 km long and rises approximately 1000 m from the floor of Dixie Valley. A slope of between 30° and 35° is maintained over much of the scarp height. The steepness of this scarp suggests a rate of uplift much higher than the regional average and thus accelerated uplift in the last few million years. If uplift of

1000 m is assumed to have occurred in 2 million years, the rate of uplift is 0.5 m per 1000 years. If a 3-m displacement event produces an M7 event, the average interval of time between such events would be 6000 years, discounting the effect of small events and tectonic creep.

The age of the latest scarp at the base of the east flank of the Still-water Range is not well determined, but a preliminary analysis of its morphology suggests that the latest displacement is no older than $12,000 \pm$ years (Holocene). The basal part of the scarp, which possibly was produced in a single event, is approximately 8 m high and, corrected for erosion, represents displacement of approximately 5-6 m.

If the 40 km-long gap were to be filled by a single fault event having an average displacement of 5 m, an earthquake of approximately $7.4 M_L$ could be generated.

White Mountain gap.--No surface faulting has occurred in historic time between the northern end of the fracture produced in 1872 and the fractures produced in 1932 and 1934. The gap encompasses an area of complex faulting, much of it of Quaternary age. In the southern part of the gap lies the precipitous north-trending scarp on the west flank of the White Mountains. In the northern half of the gap the faults of the Caldelaria Hills strike east or northeast.

The southern part of the west flank of the White Mountains is step faulted downward toward Hamill Valley, which is underlain by unconsolidated sediments approximately 2 km thick. The northern part of the west flank of the White Mountains has relief greater than 600 m, and an average slope of 30° is maintained over much of the scarp's height. Scarps in alluvium at the range front have profiles suggestive of only a few thousand years age, and in the adjacent basin unconsolidated sediments are approximately 600 m thick (Speed, written commun., 1978; and Speed and Cogbill, written commun., 1978).

The rate of uplift of the northern part of the White Mountains is estimated by Speed from erosional and debris-accumulation rates to have been approximately 0.8 m per thousand years for the last 2.7 million years.

The eastern Sierra zone.--Although containing no certain historic earthquakes over 7M, this zone, nevertheless, is the site of relatively high seismic flux (Ryall and others, 1966) and prominent young fault scarps such as that at Genoa, Nevada (Moore, 1969; and R. C. Pease, personal commun., 1978), along the east flank of the Carson Range. Some of the scarps represent Holocene displacement and earthquakes probably in the M7-8 range. With present data, average recurrence intervals for M7 earthquakes cannot be defined with precision. The range may be from 2×10^{-3} events per year per 1000 km^2 (Ryall and others, 1966) to 3×10^{-5} events per year per 1000 km^2 (Wallace, 1977a).

Southern Sierra gap.--For the southern Sierra gap, the likelihood of an M7 event is difficult to suggest from scanty geologic evidence. The area would fall within the Ventura-Winnemucca zone of Ryall and others (1966). Fault scarps on the east flank of the Sierra Nevada Mountains appear to be

potential sites of large earthquakes. Although some seismic activity has been recorded near the Kern Canyon fault, Moore and du Bray (1978) have shown that it has not moved in the last 3.5 million years.

Gap-Filling Mechanism

In the Great Basin the seismic generating ruptures in the brittle crust (approximately upper 15 km) are discontinuous even in a single earthquake. For example, in 1915 four en echelon surface breaks developed along four separate mountain blocks. Individual breaks were separated by as much as 6.4 km. The breaks also follow old fault lines having a variety of trends and types of displacement. The model shown diagrammatically in figure 3 suggests that a previously broken brittle crust is subjected to extension by an underlying anelastic dilational zone, and that overlying old faults, whatever their trends or types of displacement, become reactivated. What localizes the anelastic dilational zone is unknown, but it is this type of zone that may underlie the whole Ventura-Winnemucca seismic zone, which extends across the southwestern boundary of the Great Basin province. If the zone is continuous, gap filling would appear likely, or conversely, if gap filling is real, the zone must have continuity.

Gaps of several kilometers between individual fault scarps probably do not constitute "gaps" in a premonitory sense; they are lower order features in comparison to the underlying extensional structure. The elongate, narrow, nature of the anelastic dilational zone also suggests that gaps in activity along it may be significant in a premonitory sense. Little argument can be given for subparallel adjacent zones to become successively active. Subparallel zones, however, may become active at random times if the pattern in figure 2 is a representative sample. To put it another way, gaps along north- and northeast-trending axes are more likely to be filled in a regular sequence than are gaps along east-west trending axes.

Areas of hundreds of square kilometers that are devoid of young fault scarps, such as can be identified in figure 2, or the Trinity Range block in figure 1, cannot be considered gaps potentially to be filled. The argument can be made that many are underlain by tectonic blocks that have retained their integrity and relative stability over long periods of time. The margins of such blocks, however, may very likely be the sites of future major ruptures.

References

- Bucknam, R. C., and Anderson, R. E., 1978, Estimation of fault scarps from a scarp height-slope angle relationship: *Geology* (in press).
- Douglas, B. M., and Ryall, Alan, 1975, Return periods for rock acceleration in western Nevada: *Seismological Society of America Bulletin*, v. 65, no. 6, p. 1599-1611.
- Hanks, T. C., Hileman, J. A., and Thatcher, Wayne, 1975, Seismic moments of the larger earthquakes of the southern California region: *Geological Society of America Bulletin*, v. 86, no. 8, p. 1131-1139.
- Howard, K. A., Aaron, J. M., Brabb, E. E., Brock, M. R., Gower, H. D., Hunt, S. J., Milton, D. J., Muehlberger, W. R., Nakata, J. K., Plafker, G., Prowell, D. C., Wallace, R. E., and Witkind, I. J., 1978, Preliminary map of young faults in the United States as a guide to possible fault activity: U.S. Geological Survey Miscellaneous Field Studies Map MF-916, scale 1:7,500,000, 2 sheets.
- Moore, J. G., 1969, Geology and mineral deposits of Lyon, Douglas, and Ormsby Counties, Nevada: *Nevada Bureau of Mines and Geology Bulletin* 75, p. 1-45.
- Moore, J. G., and du Bray, E., 1978, Mapped offset on the right-lateral Kern Canyon fault, southern Sierra Nevada, California: *Geology*, v. 6, no. 4, p. 205-208.
- Ryall, Alan, Slemmons, D. B., and Gedney, L. D., 1966, Seismicity, tectonism, and surface faulting in the western United States during historic time: *Seismological Society of America Bulletin*, v. 56, no. 5, p. 1105-1135.
- Slemmons, D. B., 1967, Pliocene and Quaternary crustal movements of the Basin-and-Range province, USA: *Journal of Geosciences, Osaka City University*, v. 10, art. 1-11, p. 91-103.
- Thatcher, Wayne, and Hanks, T. C., 1973, Source parameters of southern California earthquakes: *Journal of Geophysical Research*, v. 78, p. 8547-8576.
- Thompson, G. A., and Burke, D. B., 1973, Rate and direction of spreading in Dixie Valley, Basin and Range province, Nevada: *Geological Society of America Bulletin*, v. 84, p. 627-632.
- Wallace, R. E., 1977a, Time-history analysis of fault scarps and fault traces--A longer view of seismicity: *Sixth World Conference on Earthquake Engineering*, New Delhi, India, Proc., v. 2, p. 2-409-2-412.
- _____, 1977b, Profiles and ages of young fault scarps, north-central Nevada: *Geological Society of America Bulletin*, v. 88, no. 9, p. 1267-1281.
- _____, 1978, Geometry and rates of change of fault-generated range fronts, north-central Nevada: *U.S. Geological Survey Journal of Research* (in press).

SURFACE FAULTING IN THE GREAT BASIN PROVINCE

1915
HISTORIC FAULTING
 $3 \times 10^{-5} > \text{event} > 1 \times 10^{-6}$
per year per 1000 KM²

HOLOCENE FAULTING
(last 12,000±years)
 $5 \times 10^{-5} > \text{event} > 5 \times 10^{-6}$
per year per 1000 KM²

NO LATE QUATERNARY EVENTS
(last $5 \times 10^5 \pm$ years)

NOTE: An event is an occurrence of surface faulting large enough to have produced an earthquake $>M7$. Earthquakes of 1869, 1903, 1934 and 1950 were $<M7$. Those of 1872, 1915, 1932, 1952 and 1954 were $>M7$.

Figure 1

YOUNG FAULT SCARPS IN NORTH-CENTRAL NEVADA
SHOWING AGES OF LATEST DISPLACEMENTS IN SELECTED CLUSTERS OF SCARPS

Figure 2

DIAGRAMMATIC MODEL INDICATING A ZONE OF REACTIVATED FAULTS (HEAVY SOLID LINES) IN A FRACTURED BRITTLE CRUST OVERLYING AN ANELASTIC ZONE OF EXTENSION

Figure 3

LONG TERM PRECURSORY
SEISMICITY FLUCTUATIONS

by

M. Wyss, R.E. Habermann and A.C. Johnston

Cooperative Institute for Research in Environmental Sciences
University of Colorado/NOAA
Boulder, Colorado 80309

ABSTRACT

Seismicity patterns before major earthquakes were examined on the basis of historical felt reports, the PDE world wide locations, and local network data. All of these sources contained useful data which indicate that characteristic seismicity fluctuations do exist before major earthquakes. These patterns involve quiescence and precursory clusters. The precursor times are approximately the same as those derived from other parameters. Some properties of earthquakes in precursory clusters seem to be different from background seismicity. Much more work is needed to document and understand precursory seismicity.

INTRODUCTION

The seismicity in a given crustal volume is expected to be a function of the state of stress and the resistance offered to faulting. If the earthquake preparatory process involves changes of stress, or changes of resistance to faulting, the local seismicity should also change. It now appears that precursory seismicity patterns of quiescence accented by an earthquake swarm near the future hypocenter are detectable.

Numerous authors have reported precursory seismicity patterns (e.g., Nersesov, 1973; Kelleher and Savino, 1975; Brady, 1976; Evison, 1977; Ishida and Kanamori, 1977; Sekiya, 1977; Engdahl and Kisslinger, 1977; McNally, 1977; Khattri and Wyss, 1977; Habermann and Wyss, 1977; Caputo et al., 1978). The features of these patterns are either periods of relative quiescence, clustering of events in time and space, or a combination of the two. The fact that clusters cannot always be identified may in some cases be due to poor resolution of local seismicity; in other cases precursory clusters may not occur.

The purpose of this paper is to present data showing precursory seismicity patterns before several large earthquakes, to discuss some techniques by which these patterns may be identified, and to examine the question of what kind of seismicity resolution may be necessary to detect the patterns.

METHODS

Methods used to detect precursory seismicity variations must include examination of the number of seismic events occurring in a given region, as well as their distributions in space and time. We have used four techniques of data presentation to study these parameters. In addition to visual examination by histograms, seismicity maps and time-distance plots, we have used a computer algorithm to estimate clustering of seismicity in space and time.

Histograms such as Figures 2 and 6 are an obvious means to inspect visually whether outstanding temporal seismicity fluctuations occurred in a given crustal volume. Numbers of events per unit time are plotted for crustal volumes with horizontal dimensions of about two source radii and depths from 0 to 100 km around the epicenter of large recent earthquakes.

In order to examine possible spatial patterns it is helpful to plot sequences of seismicity maps (e.g., Figure 3) and epicentral distances from the main shock as a function of time (e.g., Figure 4). These figures bring out spatial as well as temporal clustering and quiescence if such fluctuations exist.

An examination of seismicity patterns by algorithms, however, provides more objective measures for precursory activity. A proposed algorithm is described below. For a constant volume and a moving time window of constant duration the following parameters are defined:

- (1) The number of earthquakes N within the time window
- (2) The temporal concentration of activity

$$T = \sum_{i=1}^N \frac{1}{(t_i - \bar{t})^2 + 10}$$

where t_i is the time of occurrence of the i th event, and $\bar{t}$ is the mean time of the window. T becomes large for windows centered on activity which is clustered in time.

- (3) The spatial concentration of events

$$D = \sum_{i=1}^N \frac{1}{(p_i - \bar{p})^2 + 10}$$

where p_i is the position (latitude and longitude) of the i th event, and $\bar{p}$ is the average position in the window. D becomes large when events are clustered in space.

- (4) Local aftershock activity is excluded by considering only peaks of activity which also satisfy the condition

$$m_{\max} - m_{\max - 1} \leq 0.3$$

where $m_{\max}$ is the magnitude of the largest, and $m_{\max - 1}$ the second largest, event in a cluster.

- (5) Parameters (1) through (4) were designed to detect swarm activity in a given volume. We will also add the condition that relative quiescence has to precede the swarm within an area of radius of 50 km around the swarm center. We test for quiescence with a parameter N_Q , defined as

$$N_Q = \sum_{i=M-7}^{M-1} N_i \quad r \leq 50 \text{ km}$$

where M designates the month containing the swarm. N_Q sums all events contained in the six months preceding the swarm.

The above conditions were applied to the data in the following way. For the window length we chose two months. Windows containing less than two events were excluded as temporal and spatial clustering in windows containing only one event is meaningless. The averages of N , T , and D for the entire data set were calculated, and all windows with any of the parameters N , T , or D deviating more than one standard deviation from the mean were selected for further analysis.

Since we were interested in data sets for which all three parameters N , T and D were high, we determined S , the sum of the number of standard deviations that N , T and D deviated from their respective means. The mean of S was then calculated and the sets with S exceeding its mean by more than one standard deviation were retained.

Mainshock-aftershock sequences were excluded and only swarms satisfying condition (4) were processed. If quiescence preceded such a swarm (N_Q was smaller than the average number of events in six months, $N(6)$), we initiated an alarm period.

The alarm period must be constrained in time. Caputo et al. (1978) used a fixed alarm period of six years. Because the precursor times T scale with magnitude, the uncertainties in estimates of T in months will also scale with magnitude. We therefore have scaled the duration of the alarm to the length of the stage 1 quiescence, which we denote as T_{Q1} . Habermann and Wyss (1977) found the average length of time between the precursory swarm and the main shock to be $0.8T_{Q1}$. We therefore define the alarm period as centered on the time of the swarm plus $0.08T_{Q1}$ and having a duration of T_{Q1} ; in other words, it extends from $T_s + 0.3T_{Q1}$ to $T_s + 1.3T_{Q1}$, where T_s is the time of the swarm.

DATA

The seismicity data studied were obtained from NOAA's world seismicity tapes, with the exception of the Hawaii data, which were obtained from the U.S. Geological Survey Hawaii Volcano Observatory network. Seismicity on a world scale is not resolved much below magnitude 5. Therefore we searched for precursory patterns for large earthquakes only, in the hope that events in the range $4.5 < m_b < 5.5$ can serve as a measure of background seismicity. Hawaii is an exception because its local network yields a coverage complete down to at least $M_L = 2.0$.

The minimum magnitude, $m_{\min}$, for which all occurring quakes were located was estimated for the areas studied by plotting the frequency-magnitude distributions and determining the magnitude below which the cumulative numbers began to be lower than the numbers expected from the law $\log N = a - bm$. Figure 1 shows $m_{\min}$ for the areas studied, together with the four station P-wave detection threshold published by Basham and Whitham (1971). For the areas studied with teleseismic data, $m_{\min}$ varied from 4.6 to 5.0. For Europe it could not be estimated because many epicenters without magnitudes were reported.

In most of these regions approximately 40% of the located earthquakes have magnitudes less than $m_{\min}$. The total numbers of reported epicenters are generally less than 10 background events per year in volumes of dimensions 200 by 200 by 100 km. If we remove from our data set events with $m < m_{\min}$, many of the regions cannot be studied. We therefore have not removed any quakes from our sets except for aftershock sequences.

The analysis procedure was to search the NOAA data file for all quakes (depth ≤ 100 km) around recent large main shocks. Then we plotted a histogram of the events, epicenter maps, and distance from main shock as a function of time (e.g., Figures 2, 3, 4). If background seismicity was high, we searched the data for a possible precursory swarm by algorithm (e.g., Figure 5); if it was low, we felt that the application of an algorithm was not warranted and might lead to erroneous conclusions. Many of the low seismicity areas seem to give a hint of precursory activity peaks (Habermann and Wyss, 1978), which might be identified if smaller magnitude events could be located. Periods of quiescence can still be identified for low background areas by plotting the cumulative seismicity as a function of time (e.g., Figures 9, 14). Below we will present data sets of both types.

The Cape Kamchatka Earthquake

This earthquake is discussed by Habermann and Wyss (1978). We include it here because it represents a good example of precursory seismicity fluctuations and because Fedotov et al. (1977) found significant deviations of the amplitude ratios of P- to S-waves for the precursory swarm.

An earthquake with surface wave magnitude of 7.8 occurred off Cape Kamchatka on December 15, 1971. The epicenter was located at 56.0°N and 163.3°E within an aftershock zone about 100 km long. The number of earthquakes per year in the region between 55° - 57°N and 161° - 165.5°E and in the 1971 aftershock zone are shown in Figure 2 by the light and dark histograms, respectively. It can be seen that the 1971 aftershock zone experienced few earthquakes between 1960 and 1970, the only exception being 17 events which occurred during 1969.

The locations of all epicenters in the region with depths less than 55 km are shown in Figure 3. The time between January 1960 and April 1976 has been divided into eight sequential periods of lengths between 24 and 34 months, with the exception of the aftershock period of December 1971, which is shown as a separate frame. The locations of the 17 events which occurred between January 21 and 30, 1969, are shown in Figure 3D. These were virtually the only events in the 1971 aftershock zone to occur between June 1962 and December 15, 1971. These events occurred in a circle with a diameter of 40 km, and are located within 36 km of the epicenter of the 1971 earthquake. The body wave magnitudes of the swarm events ranged from 4.3 to 5.5. The isolation of this swarm, both in time and space, is further demonstrated in Figure 4, which shows the distribution of epicenters in the region of Figure 3 as a function of time and distance from the 1971 epicenter. Between the time of this swarm and the occurrence of the main shock on December 15, 1971, no earthquakes were reported in the 1971 aftershock zone (Figure 3E).

Figure 5 shows deviations from the means of spatial and temporal clustering and from numbers of events as a function of time for the Kamchatka data set minus the aftershocks of the main event. The zeroes represent the means of each parameter and the scales are in standard deviations from the mean. This is the basic output of the algorithm discussed above. Time windows for which D , N or T are one or more standard deviation from the mean are termed class one windows. Windows for which the sum of D , N and T are more than one standard deviation from the mean of the sums for all class one windows are termed class two windows. These are windows for which spatial and temporal clustering as well as the number of events are high.

Class two windows occur in Kamchatka near the end of 1964 and near the beginning of 1969. These correspond to a volcanic swarm at volcano Sheveluch, shown in the upper left corner of Figure 3C, and the 1969 swarm centered 36 km from the 1971 aftershock zone (Figure 3D). The condition for quiescence before the 1969 cluster was fulfilled (Figure 3). According to our definitions, an alarm for a major shock is issued to start in June, 1971, and to last until March, 1976. The main shock occurred in December, 1971.

The minor peak of seismicity in early 1965 was due to an aftershock sequence, which did not qualify as a significant cluster. The cluster in mid-1965 was of the swarm type, however, the condition of quiescence was not fulfilled. Therefore, no alarm was issued here, and no large main shock occurred in the area.

The Friuli Earthquake

Two quakes of magnitude 6.4 struck the Friuli region of northern Italy on May 6 and September 11, 1976. In many respects these quakes were similar to the San Fernando shock of 1971: same magnitude, similar source dimensions, and the mechanisms were mostly thrusting with several strike-slip aftershocks. Unlike the San Fernando event, the one in Italy caused extensive damage and cost approximately 1000 lives, because it affected a more populated area.

Before 1960 no earthquakes within 80 km of the main epicenters have been located. The histogram (Figure 6) shows an outstanding peak of activity during March, 1975. The events in this peak are strongly clustered in space and they occurred within the location accuracy at the same hypocenter as the subsequent main shock (Figure 7). Biagi et al. (1976) reported additional shocks for the general area without giving the actual locations. They showed a total of seven events in March, 1975, plus thirteen events distributed throughout the remainder of 1975, with no shock during 1976, before the main event. These data suggest that better resolution would in this case have improved cluster detection. The 1975 cluster contained in the NOAA data was the only class two cluster identified by the algorithm (Figure 8).

The absence of recorded seismicity before 1960, the number of cluster events, and the precursor time are very similar to the 1971 San Fernando earthquake. This suggests that earthquakes of similar size and focal mechanism may have consistently the same seismicity precursor patterns. Also the

similarity of the patterns strengthens the suggestion that these are actually precursors, i.e., part of the earthquake preparatory process.

The Great Assam Earthquake

The area around the 1897 great earthquake has experienced significant fluctuations of seismicity even though the resolution is not good enough to detect precursory clusters if they exist. Khattri and Wyss (1977) have presented this case. We show the data here because this is the largest magnitude shock for which a precursor was reported.

Figure 9 shows the cumulative seismicity within the area I shown in the inset. The epicenter of this earthquake is at the center of the square chosen for the search. It is remarkable that not a single event was reported in that area during the 28 years before this great earthquake. For comparison, during the 28 years preceding this period of quiescence 38 earthquakes were reported. This decrease of seismicity is especially surprising since one would expect that reports become more complete closer to the present because the population, industrialization, and general news coverage is increasing. Just as outstanding is the remarkable period of quiescence during the early part of the 20th century, from 1899 to 1930, which was terminated by the sequence of a magnitude 7.6 and two 7.1 earthquakes. After these events, the background level of seismicity sharply increased.

The more recent seismicity data for the area (upper curve in Figure 9) defines very clearly two periods of relative quiescence before two magnitude 6.7 events in 1950 and 1975 and a period of four years of quiescence immediately following the 1950 earthquake. The two periods of low precursory seismicity lasted eight and six years, respectively. These fluctuations could not have been due to a change in detection capability, because starting in 1963 the World Wide Seismograph network provided uniform coverage of the area, and because world wide seismicity coverage during the 1940s was better than during the 1930s. Further discussion of these data can be found in Khattri and Wyss (1978).

The Kalapana, Hawaii, Earthquake

This earthquake occurred on November 29, 1975, had a $M_s = 7.2$, and a source length of 40 km. It was the largest earthquake in Hawaii for over a century. The seismicity on Hawaii is monitored carefully because of the active volcano Kilauea. Since 1968 the data are homogeneous with complete resolution down to $M = 2.0$.

No obvious precursory cluster could be found during the period 1968-1975. Although the seismicity of this region is high, few teleseismic events occur. A study of the local seismicity revealed a distinct pattern, as shown in Figures 10 and 11. The cumulative seismicity of an area with a radius about 12 km around the main shock epicenter (Region 1 in Figure 11) is displayed in Figure 10. A significant decrease in seismicity starting about May, 1972 is clearly shown by the change in slope. Region 2 as defined by the foreshock sequences in Figure 11 also shows a decrease in seismicity beginning near May

1972. However, in March, 1974 activity increases sharply in this zone. This may indicate precursory cluster activity in this small epicentral area; the larger Region 1 remains at its lowered seismicity level.

The above results are especially strong since they are based on a demonstrably complete and homogeneous set of data and on very large numbers of events. Care was taken to include only events on the Hilina fault system and to exclude events associated with the east rift zone of the Kilauea volcano. The Hilina fault events are tectonic (Johnston, 1978) and clearly separate from events due to magma movement.

DISCUSSION

1. Relation of Seismicity Fluctuations to Other Precursors

It is important to obtain evidence which links "seismicity precursors" to the ensuing main shock or to its preparatory process, so that one may be convinced that one is dealing with precursors rather than with fluctuations unrelated to future local tectonic developments. For issuing a warning in real time, it will be indispensable to develop means of identifying precursory clusters as such. The strongest evidence in this paper linking the clusters and periods of quiescence to the main shocks, i.e., identifying them as precursors, is the spatial proximity to the main epicenter and to the rupture area, respectively. In the case of a real time analysis, one would not have the benefit of this information. Many false alarms could be avoided if the properties of the radiated waves of precursory clusters differ from those of background events. Therefore it will be important to develop diagnostic criteria based on physical parameters for precursory cluster identification.

Below we will discuss the relation of precursory seismicity fluctuations to other precursors for several earthquakes, a connection which strengthens the suggestion that the seismicity patterns are indeed related to the preparatory process.

The Friuli precursory cluster may have been accompanied by a tilt anomaly recorded at a distance of about 17 km according to Biagi et al. (1976). These authors suggest that a tilt anomaly and the cluster earthquakes during 1974 were both part of the preparatory process. They further suggest that a southward tilting trend during 1973 was also part of the preparation process. Extensive tilt anomalies may perhaps generally accompany precursory earthquake clusters because the preparatory process may affect crustal volumes in the order of the dimensions of the future earthquake. In contrast, earthquake swarms, which are not precursors, may be more localized in their effect on tilts.

The Kalapana period of decreased seismicity coincided with a well documented period of travel time delays, as shown in Figure 12 (Johnston, 1978). The travel time delays were only observed at WHA, the station closest to the epicenter (5 km distance), and not at any of the other Hawaiian stations studied (epicentral distances larger than 15 km). The coincidence of these

two precursory parameters, and their spatial relation to the main shock epicenter, strongly suggest that these changes were not caused by chance, but by the preparatory process of the Kalapana earthquake.

A preliminary study of the average magnitude, i.e., b-values, occurring in Region 1 of Figure 11 suggests that it increases significantly from 1972-1975 compared to 1968-1972. Also it is higher than in a similar region further west on the Hilina fault. This result plus the seismic velocity change plus the seismicity decrease all are consistent with a decrease of tectonic stress for at least a portion of Region 1 of Figure 11, beginning in early 1972. Work is still in progress on the seismicity for the post-earthquake time period.

The Kamchatka precursory cluster was made up of events which had been identified by Fedotov et al. (1977) as possible precursory events on the basis of their exceedingly low ratio of P-wave to S-wave energy. This result is particularly significant because it suggests that there may be characteristic changes in the radiation properties of precursory clusters.

The Garm earthquake of 22 March 1969 was preceded by a seismicity decrease which coincided approximately with a change in focal mechanism and a decrease of the P-velocity (Neresov et al., 1973; Wyss, 1975).

The San Fernando 1971 earthquake also was preceded by a precursory cluster made up of earthquakes with signal characteristics different from the background events in the areas before and after the preparation phase (Ishida and Kanamori, 1977).

2. Precursor Times Estimated from Seismicity Changes.

The precursor time can be defined as starting at the beginning of quiescence or at the time of the cluster. Depending on the data, one or the other measurement may be available. We therefore collected both types of precursor times in Table 1, marking with Q and C those obtained from quiescence and clusters, respectively. These precursor times are plotted as a function of magnitude in Figure 13, with open circles for Q and solid dots for C data.

The precursor times derived from quiescence agree with the curve proposed by Scholz et al. (1973) approximately to within factors of two. This suggests that the seismicity decrease is associated with the beginning of the preparatory process as measured by other parameters. This is the same conclusion as the one reached in the previous paragraph for the individual cases of the Kalapana and Garm earthquakes, where changes of other parameters were observed for the same events as those for which quiescence was observed.

The precursor times estimated from clusters are always shorter than those from quiescence for the events where both were observed. The C-precursor times also fall below the curve by Scholz et al. (1973) in every instance except for one data point which seems to be way off. It appears that the clusters occur at a time after the preparatory phase has already started. The clusters evidently mark a new phase in the preparatory process,

possibly a phase marked by local reorientation of the stresses, as suggested by Brady (1976).

CONCLUSIONS

1. Seismicity patterns before the main shocks seem to be real and related to the preparatory process. This is evidenced by the following facts: Similar patterns are found to occur before different earthquakes and by several authors. The precursor times derived from seismicity patterns scale with magnitude as expected from the precursor time-magnitude relation. Statistically some of these patterns cannot be considered random fluctuations, and the locations of the anomalous seismicity is closely tied to the main shock hypocenter and to the aftershock area.

2. In general, it appears that the full pattern is one of an extended period of quiescence in and around the future rupture area, briefly interrupted by a burst of swarm activity, mostly concentrated near the future hypocenter. The period of quiescence then continues up to the time of the main shock, which can be preceded by foreshocks during a very brief period before the main rupture.

3. Some precursory cluster events appear to have radiation properties different from the background events in the respective areas. This change is partly, or possibly entirely, due to changes in the focal mechanisms.

4. The resolution of the world wide location capabilities is sufficient to identify precursory seismicity patterns before large quakes in some cases, but not in others.

5. Much more data are needed to describe and understand the precursory seismicity patterns.

REFERENCES

- Basham, P.W. and K. Whitham, Seismological detection and identification of underground nuclear explosions, Publ. Earth Phys. Branch, 41, Number 9, Department of Energy, Mines and Resources, Canada, 1971.
- Biagi, P.F., P. Caloi, M. Migani and M.C. Spadea, Tilting variations and seismicity that preceded the strong Friuli earthquake of May 6, 1976, Annali di Geofisica, 29, 137, 1976.
- Brady, B.T., Theory of earthquakes IV: General implications of earthquake prediction, Pure Appl. Geophys., 114, 1031, 1976.
- Brady, B.T., Anomalous seismicity prior to rock bursts: Implications for earthquake prediction, Pure Appl. Geophys., 115, 357, 1977.
- Caputo, M., V. Keilis-Borok, P. Gasperini, L. Marcelli and I. Rotwain, Earthquake swarms as forerunners of strong earthquakes in Italy, EOS, Trans. Amer. Geophys. Union, 59, 330, 1978.
- Engdahl, E.R., and C. Kisslinger, Seismological precursors to a magnitude 5 earthquake in the Central Aleutian Islands, J. Phys. Earth, 25, 243, 1977.
- Evison, F.F., Fluctuations of seismicity before major earthquakes, Nature, 266, 710, 1977.
- Fedotov, S.A., G.A. Sobolev, S.A. Boldyrev, A.A. Gusev, A.M. Kondratenko, O.V. Potapova, L.B. Slavina, V.D. Theophylaktov, A.A. Khramov and V.A. Shirokov, Long and short-term earthquake prediction in Kamchatka, Tectonophysics, 37, 305, 1977.
- Habermann, R.E., and M. Wyss, Seismicity patterns before five major earthquakes, EOS, Trans. Amer. Geophys. Union, 58, 1194, 1977.
- Habermann, R.E., and M. Wyss, Seismicity patterns before five major earthquakes, submitted to Bull Seism. Soc. Amer., 1978.
- Ishida, M., and H. Kanamori, The spatio-temporal variation of seismicity before the 1971 San Fernando earthquake, California, Geophys. Res. Lett., 4, 345, 1977.
- Johnston, A.C., Localized compressional velocity decrease precursory to the Kalapana, Hawaii earthquake, Science, 199, 882, 1978.
- Kelleher, J.A. and J. Savino, Distribution of seismicity before large strike-slip and thrust-type earthquakes, J. Geophys. Res., 80, 260, 1975.
- Khatti, K., and M. Wyss, Precursory variation of seismicity rate in the Assam area, India, EOS, Trans. Amer. Geophys. Union, 58, 1194, 1977.

Khatttri, I., and M. Wyss, Precursory variation of seismicity rate in the Assam area, India, Submitted to Geology, 1978.

McNally, K., Patterns of earthquake clustering preceding moderate earthquakes, central and southern California, EOS, Trans. Amer. Geophys. Union, 58, 1195, 1977.

Nersesov, I.L., A.A. Lukk, V.S. Ponomarev, T.G. Rautian, B.G. Rulev, A.N. Semonov, and I.G. Simbireva, Possibilities of earthquake prediction, exemplified by the Garm area of the Tadzlink SSR, in Earthquake precursors, Acad. Sciences, USSR, eds. M.A. Sadovsky, I.L. Nersesov and L.A. Latynina, 72-99, 1973.

Scholz, C.H., L.R. Sykes and Y.P.A. Aggarwal, Earthquake prediction: A physical basis, Science, 181, 803, 1973.

Sekiya, H., Anomalous seismic activity and earthquake prediction, J. Phys. Earth., 25, 585, 1977.

Wyss, M., Precursors to the Garm earthquake of March, 1969, J. Geophys. Res., 80, 1975.

Table 1: Precursor Times Estimated from Seismicity Fluctuations

Earthquake	Year	M	Source dimension (km)	Precursor time (months)		Source
				Q	C	
Kamchatka	1971	7.8	115	143	35	1
Friuli	1976	6.4	20	-	25	new
Hawaii	1975	7.2	40	38	-	new
Assam I	1869	7.8		276	-	2
	1897	8.7		336	-	2
	1918	7.6		228	-	2
	1922	7.1		60	-	2
	1950	6.7		96	-	2
	1975	6.7		72	-	2
Assam II	1947	7.7		204	-	2
Assam III	1950	8.7		>360	-	2
Kuriles	1975	7.0	150	46	26	1
Kuriles	1976	7.0	130	97	37	1
Mexico	1970	7.3	160	88	51	1
Peru	1970	7.6	75	66+	58	1
San Fernando	1971	6.4	20	72	22	3,4
Garm	1969	5.7	10	18	-	5
Adak	1976	5.0	-	4.5	-	6
Milford Sound	1976	7.0	-	-	98	7
San Andreas	1972	5.1	7	-	120	8
San Andreas	1972	4.0	3	-	0.3	8
Izu	1974	6.9		(120?)	120	9
Gifu	1969	6.6		-	72	9
Ebino		4.1	-	-	0.6	9

1 = Habermann and Wyss, 1978; 2 = Khattri and Wyss, 1978; 3 = Brady, 1976; 4 = Ishida and Kanamori, 1977; 5 = Nersesov et al., 1973; 6 = Engdahl and Kisslinger, 1977; 7 = Evison, 1977; 8 = McNally, 1977; 9 = Sekiya, 1977.

Figure 1. Map showing seismicity resolution $m_{\min}$ for which locations are complete are shown in circles for eight areas studied. The contours are those of the four station P-wave detection by Basham and Whitham (1971).

Figure 2. Histogram of annual number of quakes in the aftershock zone (shaded) and in the surrounding area of the 1971, $M = 7.8$, Cape Kamchatka main shock.

Figure 3. Epicenter maps of approximately equal time intervals for the Cape Kamchatka, $M = 7.8$ earthquake of 1971.

Figure 4. Epicentral distance from future main shock as a function of time for seismicity at Cape Kamchatka.

KAMCHATKA 15 DEC. 71 $M_S = 7.8$

Figure 5. Deviations from the mean of the functions D, T and N plotted versus time for the Cape Kamchatka data. Peaks in these functions indicate clustering.

Figure 6. Histogram for seismicity in the Friuli area by three-month intervals. Distances < 80 km.

Figure 7. Epicentral maps for the Friuli area.

Figure 8. Deviations from the mean of the functions D, T and N plotted versus time for the Friuli data.

Figure 9. Cumulative seismicity as a function of time in the area of the great Assam earthquake of 1897 (area I in the inset). Occurrence times for all shocks with $M > 6.6$ are indicated.

Figure 10. Seismicity, both cumulative and yearly, for Regions 1 and 2 of Figure 11. Each indicated slope is a least squares fit to monthly totals.

Figure 11. Detail of the epicentral area of the 29 Nov 1975 Kalapana earthquake. Symbols: *'s - main shock and major foreshock; x's - early (same day) aftershocks; o's - Nov 10-16 foreshock sequence; ●'s - Nov 29 foreshocks. Symbol sizes represent relative magnitudes. Region 2 is defined by the boundary which includes the maximum number of foreshocks and the minimum number of aftershocks.

Figure 12. Relative residuals versus time for (A) WHA-MLO and (B) WHA-MPR for the entire Fiji-Tonga zone; (C) relative residuals versus time for MPR-MLO, for events in the north Fiji-Tonga zone only. Note the different time scale of (C). Data begin in the year of installation of each station. Symbols: o, excellent or good quality reading (both stations); ●, fair or poor; x, other than first cycle; arrows designate the date of the Kalamana earthquake; horizontal lines mark off the bounds of ± 1 standard deviation from the average (see text). The inset is the island of Hawaii; on the scale of the inset. WHA overlies the Kalapana epicenter (from Johnston, 1978).

Figure 13. Precursor time versus magnitude data based on seismicity fluctuations. The data are those summarized in Table 1.

CLOSING COMMENTS

Keiiti Aki

I think physical models such as a stress model is very important to get at this problem but a statistical model is also very important. We have heard of statistical models in this conference, including some parameterized in x, y, z for the epicenters, origin times, and now we have magnitude and even have M_S : m_b as some kind of spectral estimate. When we increase the number of parameters as much as possible, these statistical models may become more useful. Another reason why a statistical model is necessary is clearly illustrated by the example of the Oaxaca Gap. We need some objective quantity describing the probability of the break of the gap.

K-B questioned whether introduction of more parameters would really improve the quality of statistical estimates of critical failure processes or times of occurrence.

Perhaps the most fundamental problem that we have to attack is the spatial and temporal variability of aseismic slip. The definition of aseismic gap is only simple when all slip is accompanied by seismic radiation. On the other hand, there are suggestions that aseismic slip may be substantial in some regions, and we have very little idea whether this is a local phenomenon or a general characteristic of subduction. If it is a local effect, and a rapid spatial variation in the amount of aseismic slip is possible, then seismic gaps may not be regions of strain accumulation at all.

Even less is known about the temporal variation of aseismic slip. However, more and more evidence for the non-stationarity of seismic slip is being brought forward. Temporal changes in seismic activity, perhaps on a time scale of several hundreds of years, must be explained either as an indication of episodic plate motion, or as the result of physical changes on the fracture surface. Each of these possibilities raise some interesting geophysical questions. If seismic activity does change with time, the estimation of return periods of large earthquakes is not likely to be easy.

A related question, which has almost surfaced several times during this meeting, is an old one: Are large earthquakes more effective in releasing accumulated strain than small ones? If they are, it would appear that there should be substantial differences in source characteristics (particularly ambient stress drop) between these large events and all of the rest. There is astonishingly little evidence for these differences available at present. It is important that we try to obtain more information bearing on this question.

It is very noticable, for example, that the size of the "largest possible" earthquake appears to vary substantially along the Pacific subduction zone. Even within a single zone, the sizes of the earthquakes that are assumed to represent the total slip in the zone often vary appreciably. The assumption that the aftershock zones of these earthquakes represent the total zone of slippage is a convenient one, but detailed evidence in favor of this assumption is very limited.

Progress has been good toward the assessment of the long term seismic potential of seismic gaps. Geological studies, such as the papers on marine terraces and on rates of Quaternary surface faulting presented in this conference, seem particularly effective in supplementing seismicity catalogues to provide estimates of recurrence rates of major earthquakes in present seismic gaps. In addition broad-scale hypotheses developed in a plate-tectonics framework give hope for general guidelines to estimate the maximum sizes of earthquakes in subduction zones as a function of such characteristics of the zones as the width of the seismically-active interface between the colliding plates or the impinging on the zones of aseismic oceanic ridges.

It may be several years before enough high quality data on seismic gaps are available to permit an adequate test of the hypothesis that characteristics of the moderate earthquakes associated with seismic gaps can be used to predict the large gap earthquakes. We saw two observational approaches to the prediction of large gap earthquakes. One approach develops algorithms to objectively identify precursors to major earthquakes from the large amounts of catalogued seismological data. This approach may be limited by the low precision of many catalogued data, which make it difficult to construct realistic statistical models that take into account such complexities as the variations of focal mechanisms within rather small seismic regions. The second approach concentrates on high-precision studies of particular seismic gaps. This approach suffers from the small number of major earthquakes in each gap, which are too few to permit a particular pattern of precursory phenomena to be confidently identified.

It seems a seismic gap can mean different things to different people. Classically it means a zone of quiescence along a plate boundary premonitory to a major earthquake. But, it has also been used in the context of quiet zones before less than major earthquakes. It seems that we need better definition of long term historical patterns to understand gaps and possible differences or similarities in relation to the size of the impending earthquake.

Another obvious need is to find ways to measure stress levels in seismic gaps, either directly or indirectly. Indirect measurements can perhaps be made by analysis of temporal changes in the waveforms of earthquakes occurring within a gap.

With regard to the problem of earthquake prediction, in particular, the identification of a seismic gap, by whatever criteria are adopted, is not very useful without some measure of the regional strain rate and the strain history. For example, a seismic gap may exist in a region that is either temporarily or permanently decoupled from the mechanisms that drive the accumulation and release of strain in surrounding regions.

Within the earth's seismic regions there may exist a suite of tectonically active strain concentrators each of which is related to a particular structural defect such as a magma chamber or join between different rock types. The level of such activity may be indicative of the regional strain accumulation or strain rate. For example, increased swarm activity and major volcanic eruptions have preceded great earthquakes along the Nankai through marginal to SW Japan. Similar activity could provide essential supporting evidence for strain accumulation in regions where geodetic surveys are impossible or non-existent. The statistical studies presented at the meetings are interesting in that correlations were proposed between seemingly remote swarms of earthquakes and future great earthquakes at plate boundary faults.

Hanks

By way of my own summary, I'd like to address Jack's remarks about what he wanted us to say by way of summary. I think definitions are important if only because the one we are presently working with, that a seismic gap is somewhere where earthquakes haven't occurred for some time, is so useless.

What Mike Chinnery said about aseismically slipping regions, what Bill Page said about recurrence interval problems, what Nafi Toksoz, Willie and Gary Latham said about variations in levels of Seismicity are all, terribly important in being precise about what you mean when you say seismic gap. Because of all of these uncertainties, I think anyone's estimate of a soon-to-break seismic gap is little more than a guess, even as I am aware that open-ended predictions have been made successfully on the basis of identifying seismic gaps.

But nevertheless, I enjoyed this conference a great deal. I learned a lot about what we don't know about gaps.

Jones (as transcribed from tape)

I found myself seeing that gaps really could be a useful concept for determining where to concentrate research on possibly upcoming earthquakes. But I was also struck by a lot of the problems that are going to be involved in using them. We just really don't know how large a random factor goes into earthquake occurrence and how accurate are recurrence times, how often earthquakes will repeat themselves in the same place.

I'd like to reemphasize other people's points about using geology and other means of getting a larger data base to try and determine some of these problems.

Also, I thought of the foreshock and using other means to add to the accuracy of the gap theory. They still aren't feasible for doing right now and there needs to be more work, I think especially in individual areas determining the individual geologic history of the particular area to be able to use any of the techniques for predicting.

In general, I think we need to admit that we are going to be making mistakes as we go along.

Kagan:

I think that the most important conclusion that I derived from the discussion here is that seismic gaps studies should be tied to investigation of the whole picture of earthquake occurrence. So we should take into account clustering of events, migration, stationarity (or non-stationarity) of seismic process and geometry of faulting. Seismic gap is a time-space-magnitude phenomenon, so to describe, explain and possibly predict its break, we need data which are of similar format, i.e., seismicity. So it is not coincidental that nearly all papers have used seismicity patterns as their major data. A problem with seismicity is that it is highly variable and has an extremely random character. To study effectively such data we need a statistical approach. But the problem with statistics is that we have very few strong earthquakes recorded adequately. We cannot wait a thousand years to get a sufficient number of earthquakes. Another approach is to use self-similarity of the seismic process. That means that the seismic process is auto-modelling itself, that patterns of small earthquakes are similar with a change of time-distance scale to those of large earthquakes. Another possibility is to use new data, and the most interesting new parameters in my opinion are fault-plane solutions. The information content of catalogs we have now is about one bit per earthquake. This means we can reduce uncertainty of earthquake risk prediction about two times comparing with completely random estimates. I believe that with the inclusion of fault-plane solutions in our catalog data we will have a great increase in the amount of information with corresponding reduction of uncertainty.

Although "seismic gap" has never been defined precisely, historically (Imamura, 1928; Fedotov, 1965; Mogi, 1968; Kelleher et al., 1973), it was simply defined as an area that corresponds to the known rupture zone of a previous major earthquake but has not broken for a considerable length of time. More recently, an area where seismic activity is significantly lower than in the surrounding area is often called a gap regardless of whether the previous activity is accurately known or not. The San Fernando gap, and the Kern County gap belong to this category. Seismic gaps in this category are more controversial.

The Suruga-Bay gap, Japan, belongs to the first category, although the spatial extent of the previous activity and the repeat time are still uncertain. It appears that most major (length longer than 200 km) gaps have broken during the last 25 years or so, and very few, if any, seem to remain as a ready-to-break gap. Unfortunately, the nature of most gaps which may be identified now is uncertain with respect to the previous activity, the spatial extent and the repeat time. It is, therefore, extremely difficult to identify a ready-to-break gap with any confidence. In view of the number of recent studies on seismicity, it seems most promising to first investigate the spatio-temporal pattern of seismicity in and around the suspected gaps. If any pattern diagnostic of foreshock activity, such as clustering, change in mechanism, change in spectra, is found, more intensified observations may be called for.

However, as the example for the Kern County earthquake suggests, it seems difficult to predict deterministically when the final rupture (main shock) is

finally triggered. Also, if one gap breaks, it may trigger the neighboring gaps causing a multiple shock or a doublet (e.g., large earthquakes in the Solomon Islands). Thus, deterministic prediction of the size may be difficult too.

References

- Fedotov, S. A., Regularities of the distribution of strong earthquakes in Kamchatka, the Kuril islands and northeastern Japan, Trans. Acad. Sci. U.S.S.R., Inst. Phys. Earth, 36 (203), 66-93, 1965 (in Russian).
- Imamura, A., On the seismic activity of central Japan, Jap. J. Astron. Geophys., 6, 119-137, 1928.
- Kelleher, J., L. Sykes, and J. Oliver, Possible criteria for predicting earthquake locations and their application to major plate boundaries of the Pacific and the Caribbean, J. Geophys. Res., 78, 2547-2585, 1973.
- Mogi, K., Development of aftershock areas of great earthquakes, Bull. Earthquake Res. Inst., Tokyo Univ., 46, 175-203, 1968.

My opinion is mostly similar to that described by previous speakers but let me summarize in this way (see figure). Here is a seismic event to be predicted (S), here is an event which we alarm (A). This area constitutes a successful prediction (P), so that our goal is to get the parameter $(P/S) \div (P/A)$ as close to one (1) as possible. To improve this first parameter (P/S), several items should be considered: to collect seismic geographic data, seismic historical data (especially for uncovered areas); data interpretation hopefully by standardized techniques for facilitation of communication, there being several techniques depending on the items to be subject to survey (interplate gaps, intra-plate gaps, long time and short time prediction, etc.), followed by proper evaluation depending upon the respective category.

The second item (P/A) should be improved in the following ways. The first step is to determine the seismotectonic significance as much as possible. I think there should be a level of space and time which belongs in tectonic inevitability. Simple detection of time-space gap will not be sufficient. It is needless to say that it is necessary to promote better understanding of seismo-tectonic processes. Also, it is important and necessary that intensive field work be organized. lastly, and the most undeveloped area to date, there should be development of comprehensive diagnostic of the data including statistical tests. It will always be necessary to undertake proper assessment of each technique proposed, including retrospective examination.

Goal

$$\left(\frac{P}{S}\right)\left(\frac{P}{A}\right) \rightarrow 1$$

To improve $\left(\frac{P}{S}\right)$:

seismic geography
(history)
tectonic environments } especially for uncovered areas
data interpretation by standardized techniques
proper evaluation of respective category

To improve $\left(\frac{P}{A}\right)$:

seismotectonic significances
better understanding of
physical mechanism, seismic processes
intensive field work in critical areas
comprehensive diagnostics & statistic tests

Additionally:

assessments, including
retrospective examination

V. Keilis-Borok

1. Let me first suggest an answer to the question proposed by Dr.

Evernden. I believe, that soon-to-break seismic gaps are such gaps, near which activation takes place. Activation can be diagnosed, for example, by recent occurrences of sufficiently strong earthquakes at the same fault where a gap is observed. This pattern is formally described in my paper as precursor AQ. Activation can also be diagnosed by the bursts of seismicity, formally described as precursors B, S, Σ . I think that other descriptions could be as good. Many presentations have pointed in this direction: combination of quiescence (or gap) and activation, as the premonitory seismicity pattern.

2. I believe the problem will be less difficult, if not easier, if we will separate long and short term predictions, and separate predictions of the strongest earthquakes possible in the area and the medium ones, which can also cause damage. I think that the less difficult part is the long term prediction of the strongest earthquakes. It requires the following minimum of data.

- (a) a reliable earthquake catalog for at least 30 - 40 years, for magnitudes of at least 2.5 - 3 units smaller than the earthquakes we are trying to predict;
- (b) a scheme of lineaments of at least the first three ranks;
- (c) some preliminary study of earthquake-prone segments of these lineaments.

I believe that with such a data base, about 80% of the strongest earthquakes can be predicted with a total duration of alarms about 10 - 15%. So, we are on the verge of solution of a part of the problem: prediction in the long term sense, i.e., without pinpointing the exact time and epicenter. It is a necessary step in short term prediction and it would be of some independent practical importance.

I'm still in a mist as far as short term prediction and prediction of medium earthquakes are concerned.

3. I think one of the difficulties in the search for long term precursors is the expectation to have them only very close to a future fault-break. It would certainly make the problem easier and also decrease the danger of self-deception. However, we have no real reasons to count on such luck. Generally speaking, an earthquake occurs if the difference between strength and stress is small enough, whereas the strain energy is large enough. I see no reasons to expect that this pattern would be strongly localized. Qualitative conclusions won't be unique. Facts indicate the opposite.
4. The absence of theory is a grave difficulty, but I feel that we are sometimes too scared of this. Any theory at this moment would be non-unique. Prediction is in the same state as gravity studies would be prior to formulation of Kepler's laws. We have to get more hard facts from our observations to design a theory. Otherwise,

we may be hypnotized by one particular hypothesis or by the lack of a confidence level. So far, the greatest difficulty is the state of the data base.

5. If we are really close to actual prediction, there is the difficulty of how to release it. A good public service and a scientifically valuable result would be to have one real alarm for one or two false alarms. It would be from 20 to 100 times better than random statements. So, it is informative. But, with such probability of false alarms, we need to have some administrative procedure; some written orders of how to respond to prediction, if the probability of an earthquake is 30% or 50%, but not 99%.

Latham

Well we've defined gaps here as either the space between the aftershock zones of very large earthquakes such as Bill has done, or the onset of quiescence in previously active zones and that was the thrust of the study that I reported on.

I think it's relevant that Dr. Kagan has pointed out that behavior of small earthquakes in his modeling of space-time relationships could not be distinguished from that of large earthquakes. So it gives me some hope that patterns that we might see in very large earthquakes might be revealed in those of the small ones as well, since our statistical basis is a little better for those events.

We and others have suggested a pattern in which quiescence has developed in a zone some years preceding a main shock with an eventual resumption of activity occurring prior to the shock. And of course the crucial question is "How common is that pattern?". We should not be surprised that the pattern will vary greatly from region to region, nor disappointed that our favorite pattern that developed in one region does not apply at all in another.

The details of the precursory seismic activity will almost certainly vary. A note of caution has been raised in my mind at this session that the assumption of stationarity in space time relationships over intervals of say 100 years is probably a very dangerous assumption.

Willie Lee

I have two points to make. One is that the gap theory seems to work very well when the plate boundaries are clear cut. However, in many places where we have damaging earthquakes like in China and possibly in the western United States, the plate boundaries are not clear. I don't know how to extrapolate the gap theory to those areas.

The second point I would like to make is well known and has been said by many people, i.e., we cannot bring out any information that is not in the data in the first place. I was struck by the fact that very little effort was put into collecting long-term data properly. For example, we have about 80 years of instrumental earthquake data. These data are extremely inhomogeneous and we have good earthquake data for no more than 15 to 20 years.

In order to study earthquake recurrence rate and long-term seismicity patterns, we need to pay more attention to the older instrumental records and historical documents on past earthquakes. By studying areas with long historical records (like China, Turkey, and Middle East), we may be able to figure out if the earthquake recurrence rate is 100 years or so as some seismologists believe or thousands of years as some geologists argue.

I'd like to say a few words about how to go about testing the gap hypothesis. Jefferys once said that if a certain physical effect is sufficiently strong it shouldn't require statistical testing to bring it out. If this is true, then the gap hypothesis really refers to what one might call a second order effect. It's not a really dominant effect in the sense that there can be an argument as to whether a gap does exist. In that case, we really need some kind of a model to use for testing against. What kind of model could be used?

There are physical models and statistical models and we've been talking about this with some of the people at this conference. There seems to be an agreement that the model should be a physical model, and that poses a certain number of questions. Number one: Are we going to look at plate boundaries only or are we also looking at intraplate phenomena? It seems to me that personally looking at interaction between the plates. There has not been universal agreement as to that.

There is also a question if we look at the plate boundary whether we should look only at "shallow" events or at all events that comprise the interaction between the plates. This may involve a redefinition of "shallow" as function of thickness of the lithosphere.

And finally, there is the problem of what mechanical analog we might use to simulate the interaction between the plates. There aren't really too many. There is the Burridge-Knopoff model. In conclusion what I think I got out of the meeting more definitely is that the one thing to test for is clustering, and a gap of quiescence is negative clustering. In other words, there are tests for clustering that can be applied and there seems to be a very promising possibility for doing that.

McCann (as transcribed from tape)

I just want to say a few brief words about seismic gaps. There are many regions as indicated on our map which fit some very simple criteria on the very simple plate boundaries which may have accumulated sufficient strain that would be released in large earthquakes within the next few to several decades. A few of the regions might have a higher potential than others. I'm not sure that the simple seismic gap analysis that we can do given our limited state of knowledge can sufficiently differentiate from one region to another. We can only push this so far especially since we've seen that some complicated events have occurred like the events off the Kuriles in 1970.

In order to really complete this analysis, we're probably going to need at least one full seismic cycle in each area where we are sure that each event was a thrust event, and wasn't an intraplate event such as normal faulting in the ocean floor or in the inland arc structure.

As far as using other methods to pick out which gaps might have a higher potential for being the source region for a large earthquake, there seems to be some evidence from what Dr. Keilis-Borok and Dr. Kasahara have done that swarms of events do seem to occur before some of these moderate and large earthquakes, and other people have also seen periods of quiescence occurring before these large events. I think if we can start doing some pattern recognition and start concentrating on these gaps and try to look for variations in the seismic activity, we might be getting somewhere.

I'm struck by the large differences between the recurrence intervals that seismologists and geologists talk about; seismologists discuss less than a few 100 years, geologists 1,000 to 50,000 years. Is the difference due to episodic times for earthquakes or are geologists just not seeing but one in ten. This difference pertains to the definition of a seismic gap. If a seismic gap is related to space and time, how long or short is the time between earthquakes before it is called a gap? How do you define a "soon-to-break" gap? If major earthquakes in an area have a 1,000 year recurrence interval, then scientists need to look back much more than 1,000 years. Geologists are needed to help extend the seismological framework. There are some areas of the world where the geology can be studied by ordinary techniques that we now know with a high probability of success in extending back the seismic record. It would be interesting to study the areas where known historical events in China have occurred. By examining the geomorphology of particular faults, trenching those faults, and looking for cultural materials, a comparison of recurrence based on geology with one based on the historic record could be made. Other areas could similarly be studied. Geology can also contribute to defining the differences between aseismic gaps and seismic gaps. Thatcher's type of model for uplifted coasts in seismic areas could be applied to an aseismic area to predict differences; then several plate margin coasts could be examined to test the model.

I'd like to make three points. I think the issue of understanding the occurrence and closing of gaps is important enough from several stand-points and we have come to a stage where we need to organize thinking as well as effort with respect to getting a better understanding of how gaps occur and how they are filled. I'd like to suggest that this effort should be twofold. It should certainly be pursued as a scientific inquiry but, at the same time, we should also keep in view the risk-oriented aspects of gaps since there is a lot that depends on what the effects may be when a gap-filling earthquake occurs. One particular example is offshore petroleum development. I would like to suggest that, although our knowledge at present is imperfect, we can try to improve it in two ways. One is by continuing to collect more observational data, however, what might be more purposeful is to make attempts toward modeling this process of formation and filling of gaps. Like any other situation, there are two ways of modeling a process, a physical model and a statistical model. At this stage, I feel that the basic framework of a model ought to be physical. The parameters in that model may be so chosen on a statistical basis to reflect the scatter and the uncertainty that would go into it. Such approaches have been used before and they may be worth trying.

The third point I'd like to make is that we have heard quite a few presentations regarding precursors which might serve as indicators of the imminence of a large gap-filling earthquake. It appears that it is not clear at this stage whether these precursors have any special features that go with having waited for a long time or they are like the precursors to any other earthquake and we need to pursue that further.

What was most impressive to me at this meeting is the order-of-magnitude difference in postulated return times for large and great earthquakes as determined from purely seismologic data and those obtained from geologic studies, long historic records, or by consideration of strain rates from plate tectonics theory. Worldwide seismological data such as those summarized here by Bill McCann suggest return times measured in decades whereas all the other available lines of evidence indicate that it is more likely on the order of centuries.

For example, in subduction zones the shortest known return time for large earthquakes is about 100 years in the Nakaido, Japan area. My studies of marine terraces in Alaska, and similar studies by Matsuda and others in the Kwantu, Japan area suggest return times as long as 500 to 1500 years. Furthermore, when we consider the amount of slip required for great earthquakes such as the 1960 Chile (~ 20 m), 1964 Alaska (20 m +) or 1899 Yakutat Bay (16 m+) events together with reasonable strain rates in these focal regions, we must conclude that such earthquakes cannot return in the same place in less than about 250 to 350 years even if the system is perfectly efficient.

Similarly, in transform zones recurrence times of no less than 150 years are indicated for the 1906 earthquake on the San Andreas fault and the 1976 earthquake on the Motagua, Guatemala fault. Lensen's studies of offset river terraces along the northern part of the Alpine fault system in New Zealand indicate that fault displacements recur there about once every 500 to 900 years.

Of course, when we look at intraplate zones, return times are even longer. Bob Wallace's geologic study of Great Basin Faults suggests return times of 1,000 to 10,000 years. In the same region geologic studies at Borrego Mountain by Malcolm Clark and others show returns of about 200 years and Bonilla's work at Pleasant Valley suggests roughly 700 years. The historic data for China presented by Willie Lee indicate that an event like the destructive Sian earthquake requires more than 440 years to recur and Ambrasey's analysis of the historic record in Turkey suggests that recurrence of fault displacement on the North Anatolian fault may be on the order of 1,000 years.

What all this implies, is that if we are trying to catch a really large earthquake, the wait may be longer than many of our expected lifetimes. Furthermore, the best opportunities are likely to be geologically identified active faults that have not had any known displacement for 100 years or more. The alternative is to concentrate on the more frequent moderate-sized events. However, for such events the errors in the older epicentral locations begin to approach the dimensions of the focal region as was made apparent from the discussions here by Lomnitz, Latham, and Reyes with reference to the Oaxaca "gap". In any case, if we are to make significant progress towards understanding what is going on along major fault zones where there may be a gap, it is essential to integrate all available geological, geophysical, geodetic and historical data in addition to the seismology in order to focus in on the potential seismic sources in that particular segment of the earth's crust.

CICESE

Centro de Investigación Científica y de Educación Superior de Ensenada, B. C.

AV. ESPINOZA No. 843 - TELEFONOS 8-16-60, 8-12-16, 8-10-48 - TELEX 056539 - APDO. POSTAL 2732 - ENSENADA, BAJA CALIFORNIA

August 22, 1978

Dr. Jack F. Evernden
United States
Department of the Interior
Geological Survey
Menlo Park, Ca. 94025

Dear Dr. Evernden:

First of all, I am deeply thankful for having the opportunity to participate in this conference, and to get in contact with leading scientists, which I considered a great opportunity for my own formation as a seismologist.

Thinking over the results of the conference, it appears to me that there are still some fundamental problems to be solved. For instance: What is the relationship between earthquake recurrence times of the order of 1000 years, and those with 50 years, as has been observed along the Alaskan and Peru-Chile Trench?. Why does the earthquake cycle varies even along the same tectonic environment?

Earthquake fault lengths of the order of 150 km are as important as those with fault lengths of the order of 1000 km, for public safety.

Plate tectonics provide us with a convenient Kinematic model to our understanding of earthquake potential along major plate boundaries, characterized by strike-slip faults, and thrust faults. Precise measurements of slip rates inferred from world-wide plate models, along with those determined from seismic source studies, should provide an insight into our understanding of seismic gaps as originally defined. This recalls for additional studies on relocation of major earthquakes, its aftershock distribution and associated tectonic and geological settings.

Plafker's detailed geological and geophysical studies of the Aleutian and the Peru-Chile Arc. Show that the 1964 Alaskan and 1960 Chilean Earthquake occurred along megathrust and subsidiary thrust faults.

This leaves an open possibility that megathrust faults are strong enough to resist slippage over periods of time of the order of 1000 years. While subsidiary thrust faults go through cycles of strain build up and release of the order of fifty years. In the first case, strain release occurs along most of the sector of the arc, and in the former over a few hundred kilometers of the arc.

There are additional complexities when we look at the repeat times of earthquakes with characteristic source dimensions of a few hundred kilometers, which indicate that the slippage strength along a subduction zone varies substantially and in particular areas is strong enough to resist slippage while adjacent sections go through at least two cycles of strain build up and release.

Complexities of this type call for additional studies of seismicity using local arrays to obtain better estimates of hypocentral distribution and focal mechanism and in addition to improve our teleseismic travel times, for relocation of seismic foci, using teleseismic data for previous seismicity.

Yours sincerely,

Dr. C. Alfonso Reyes Z.
Head of the Geophysics Department

Albert Smith

To understand the origin of seismic gaps and whether migration occurs, we need a data base. This coincides with Dr. Lee's idea for additional studies of historical earthquakes and Dr. Kagan's suggestion of similarity. Only then will the application of quantitative or statistical analysis determine the existence of gaps and migration. We need this analysis to provide tests for our physical models.

From the point of view of physical models, what kinds of processes are we dealing with? What controls the rates that determine the occurrence of seismic gaps and the resulting seismic migration? What is the importance of inhomogenieties? These and other questions must be answered if seismic migration and seismic gaps are to provide a predictive tool.

At this stage of the go-around, one can only reiterate and agree with what previous speakers have said.

In terms of our goals of hazard assessment and earthquake prediction, I think that seismic gaps, despite present problems, may in fact turn out to be the most useful aspect of hazard mitigation for two reasons; firstly, because we need intermediate or long term predictions so that we know where to concentrate our attention, and, secondly, because I think it's likely that our attempts to use shorter term precursors will not be infallible or may not even work at all. I suggest that we concentrate first of all on what we would term simple seismic gaps where the processes can be understood in a fairly straightforward way and from where the phenomenology can be extrapolated to more complicated regions. I think such a focus is very important. Understanding of the processes even at the simple boundaries at this point is quite imperfect and we have to begin to construct models which will account for the basic processes. We still don't really understand quite how subduction occurs! That's a very important process and until we really understand the simple situations, extrapolation, more complex boundaries is going to be very hard. I certainly agree with the points made first of all by Bill Page and later by subsequent speakers about the importance of geologic studies of recurrence. Such studies are central to assessing seismic gaps. I should also like to add a pitch for the applicability of my own special interest at the moment, geodetic measurements of crustal deformation. If strain accumulation can be detected in a suspected seismic gap, then we know that a large earthquake will occur someday. If we cannot measure detectable deformation, larger than say a few parts in 10^8 /year, then we can be pretty sure the suspected gap is either entirely aseismic or the recurrence interval is very long, 1000 years or more. Measurements such as these might settle, for example, whether or not the Marianas arc is a gap or not.

I'd like to elaborate on a few of the points that have been mentioned. Quiescent periods in seismicity in a given area seem to indicate one of three things: (1) aseismic gaps in the terms we are interested in. (2) an area of strain being released by aseismic slip, and (3) an area in which the strain was so thoroughly released by some previous earthquake that it may be many years before another event occurs. We should look at what we have called "gaps" in this framework and try to identify as many of those of the first kind as possible. Out of these we should select a manageable number, with preliminary instrumentation and studies, for more detailed instrumentation and monitoring.

Another aspect that has been brought up in this conference is that there seem to be methods that may enable us to estimate the possible time of an earthquake from pattern recognition and other studies. This "time" element may be combined with the "space element of the seismic gaps. Both time and space characteristics of identified gaps must be combined to choose those for immediate instrumentation and careful monitoring.

In order to know whether a seismic gap is ready to break, we need the ability to know the state of stress in a crustal volume and possibly also the state of strength if that can change as a function of time, which has been suggested. How to measure the in-situ stress is very costly and hard to do, so we are trying to measure crustal parameters that reflect somehow the state of stress in this volume. I found that to my taste at this conference there was somewhat of a lack of discussion of the various parameters that we could measure to try to identify the state of stress and which would reflect a preparatory phase of a crustal volume. We have discussed one such parameter, which is seismicity, almost exclusively to other parameters. Other things that I would think one should discuss at some length at some point are things like crustal deformation, velocity changes, magnetic, resistivity and other parameters. Naturally the seismicity has one advantage which is you can measure some of it teleseismically and at no further expense than the present existing worldwide network and insofar it might be the first parameter to be measured. But I think that ultimate predictions can really only be based on a combination of several changes of several parameters.