

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

TEXT TO ACCOMPANY:
COAL RESOURCE OCCURRENCE
AND
COAL DEVELOPMENT POTENTIAL
MAPS
OF THE
FATS DRAW QUADRANGLE,
CAMPBELL COUNTY, WYOMING

BY
INTRASEARCH INC.
DENVER, COLORADO

OPEN FILE REPORT 79-057
1979

This report is preliminary, and has not been
edited or reviewed for conformity with
United States Geological Survey standards or
stratigraphic nomenclature.

TABLE OF CONTENTS

	<u>PAGE</u>
I. INTRODUCTION	1
II. GEOLOGY	3
III. DATA SOURCES	8
IV. COAL BED OCCURRENCE	10
V. GEOLOGICAL AND ENGINEERING MAPPING PARAMETERS	13
VI. COAL DEVELOPMENT POTENTIAL	15
Table 1.--Strippable Coal Reserve Base Data (in short tons) for Federal Coal Lands in the Fats Draw Quadrangle, Campbell County, Wyoming.	19
Table 2.--Coal Resource Base Data (in short tons) for Underground Mining Methods for Federal Coal Lands in the Fats Draw Quadrangle, Campbell County, Wyoming.	20
Table 3.--Coal Resource Base Data (in short tons) for In-Situ Gasification for Federal Coal Lands in the Fats Draw Quadrangle, Campbell County, Wyoming.	21
SELECTED REFERENCES	22

TABLE OF CONTENTS (continued)

	<u>MAPS</u>	<u>PLATES</u>
1.	Coal Data Map	1
2.	Boundary and Coal Data Map	2
3.	Coal Data Sheet	3
4.	Isopach and Mining Ratio Map of Lower Felix Coal Bed	4
5.	Structure Contour Map of Lower Felix Coal Bed	5
6.	Isopach Map of Overburden of Lower Felix Coal Bed	6
7.	Areal Distribution of Identified Resources of Lower Felix Coal Bed	7
8.	Identified Resources of Lower Felix Coal Bed	8
9.	Isopach Map of Smith Coal Bed	9
10.	Structure Contour Map of Smith Coal Bed	10
11.	Isopach Map of Overburden of Smith Coal Bed	11
12.	Areal Distribution of Identified Resources of Smith Coal Bed	12
13.	Identified Resources of Smith Coal Bed	13
14.	Isopach Map of Wyodak Coal Zone	14
15.	Structure Contour Map of Wyodak Coal Zone	15
16.	Isopach Map of Overburden of Wyodak Coal Zone	16
17.	Areal Distribution of Identified Resources of Wyodak Coal Zone	17
18.	Identified Resources of Wyodak Coal Zone	18
19.	Isopach Map of Pawnee Coal Bed	19
20.	Structure Contour Map of Pawnee Coal Bed	20
21.	Isopach Map of Overburden of Pawnee Coal Bed	21
22.	Areal Distribution of Identified Resources of Pawnee Coal Bed	22
23.	Identified Resources of Pawnee Coal Bed	23

TABLE OF CONTENTS (continued)

<u>MAPS</u>	<u>PLATES</u>
24. Isopach Map of Wildcat-Moyer-Oedekoven Coal Zone	24
25. Structure Contour Map of Wildcat-Moyer-Oedekoven Coal Zone	25
26. Isopach Map of Overburden of Wildcat-Moyer-Oedekoven Coal Zone	26
27. Areal Distribution of Identified Resources of Wildcat-Moyer-Oedekoven Coal Zone	27
28. Identified Resources of Wildcat-Moyer-Oedekoven Coal Zone	28
29. Coal Development Potential for Surface Mining Methods	29
30. Coal Development Potential for In-Situ Gasification	30

CONVERSION TABLE

<u>TO CONVERT</u>	<u>MULTIPLY BY</u>	<u>TO OBTAIN</u>
inches	2.54	centimeters (cm)
feet	0.3048	meters (m)
miles	1.609	kilometers (km)
acres	0.40469	hectares (ha)
tons (short)	0.9072	metric tons (t)
cubic yards/ton	0.8428	cubic meters per metric tons
acre feet	0.12335	hectare-meters
Btu/lb	2.326	kilojoules/kilogram (kJ/kg)
Btu/lb	0.55556	kilocalories/kilogram (kcal/kg)
Fahrenheit	5/9 (F-32)	Celsius

I. Introduction

This report and accompanying maps set forth the Coal Resource Occurrence (CRO) and Coal Development Potential (CDP) of coal beds within the Fats Draw Quadrangle, Campbell County, Wyoming. This CRO and CDP map series (U. S. Geological Survey Open-File Report 79-057) includes 30 plates. The project is compiled by IntraSearch Inc., 5351 South Roslyn Street, Englewood, Colorado, under KRCRA Eastern Powder River Basin, Wyoming Contract Number 14-08-0001-17180. This contract is part of a program to provide an inventory of unleased federal coal in Known Recoverable Coal Resource Areas (KRCRAs) in the western United States.

The Fats Draw Quadrangle is located in Campbell County, in northeastern Wyoming. It encompasses all or parts of Townships 45, 46, and 47 North, Ranges 75 and 76 West, and covers the area: 43°52'30" to 44°00' north latitude; 105°52'30" to 106°00' west longitude.

Main access to the Fats Draw Quadrangle is provided by a maintained gravel road that traverses across the southeast corner of the study area. This gravel road intersects Wyoming State Highway 50, approximately 10 miles (16 km) northeast of the quadrangle and 24 miles (38 km) southwest of Gillette, Wyoming. Minor roads and trails that branch from this gravel road provide additional access to the more remote areas. The closest railroad is the Burlington Northern trackage, 25 miles (40 km) to the northeast at Gillette, Wyoming.

Northwestward-flowing Pumpkin Creek and its tributaries provide the major drainage for the quadrangle. The North Prong, Middle Prong, and South Prong of Pumpkin Creek drain the northern, central and southern portions of the quadrangle, respectively. Fats Draw, Hartzog Draw, and numerous intermittent streams supplement the drainage throughout the quadrangle. Pumpkin Creek flows into the Powder River approximately 10 miles

(16 km) to the northwest of the quadrangle. Elevations attain heights of 5021 feet (1530 m) above sealevel in the southeast corner of the study area, 500 to 600 feet (152 to 183 m) above the valley floors to the northwest. The somber grays, yellows, and browns of outcropping shales and siltstones, contrast strikingly with the brilliant reds, oranges, and purples of "clinker," and the deep greens of the juniper and pine tree growth.

The 13 to 14 inches (33 to 36 cm) of annual precipitation falling in this semi-arid region accrue principally in the springtime. Summer and fall precipitation usually originates from thunderstorms, and infrequent snowfalls of 6 inches (15 cm) or less generally characterize winter precipitation. Although temperatures ranging from less than -25°F (-32°C) to more than 100°F (38°C) have been recorded near Gillette, Wyoming, average wintertime minimums and summertime maximums range from +5° to +15°F (-15° to -9°C) and 75° to 90°F (24° to 32°C), respectively.

Surface ownership is divided among fee, state, and federal categories with the state and federal surface generally leased to ranchers for grazing purposes. Details of surface ownership are available at the Campbell County Courthouse in Gillette, Wyoming. Details of mineral ownership on federal lands are available from the U. S. Bureau of Land Management in Cheyenne, Wyoming. Federal coal ownership is shown on Plate 2 of the Coal Resource Occurrence maps. The non-federal coal ownership comprises both fee and state coal resources.

The Coal Resource Occurrence and Coal Development Potential program pertains to unleased federal coal and focuses upon the delineation of lignite, subbituminous coal, bituminous coal, and anthracite at the surface, and in the subsurface. In addition, the program identifies total tons of coal in place, as well as recoverable tons. These coal

tonnages are then categorized into units of measured, indicated, and inferred reserves and resources, and hypothetical resources. Finally, recommendations are made regarding the potential for surface mining, underground mining, and in-situ gasification of the coal beds. This report evaluates the coal resources of all unleased federal coal beds in the quadrangle which are 5 feet (1.5 m) or greater in thickness and occur at depths down to 3000 feet (914 m). No resources or reserves are computed for leased federal coal, state coal, fee coal, or lands encompassed by coal prospecting permits and preference-right lease applications.

Surface and subsurface geological and engineering extrapolations drawn from the current data base suggest the occurrence of approximately 7.1 billion tons (6.5 billion metric tons) of unleased federal coal resources in the Fats Draw Quadrangle.

The suite of maps that accompany this report sets forth and portrays the coal resource and reserve occurrence in considerable detail. For the most part, this report supplements the cartographically displayed information with minimum verbal duplication of the CRO-CDP map data.

II. Geology

Regional. The thick, economic coal deposits of the Powder River Basin in northeastern Wyoming occur mostly in the Tongue River Member of the Fort Union Formation, and in the lower part of the Wasatch Formation. Approximately 3000 feet (914 m) of the Fort Union Formation, including the Tongue River, Lebo, and Tullock Members of Paleocene age, are unconformably overlain by approximately 700 feet (213 m) of the Wasatch Formation of Eocene age. These Tertiary formations lie in a structural basin flanked on the east by the Black Hills uplift, on the

south by the Hartville and Casper Mountain uplifts, and on the west by the Casper Arch and the Big Horn Mountain uplift. The structural configuration of the Powder River Basin originated in Late Cretaceous time, with episodic uplift thereafter. The Cretaceous Cordillera was the dominant positive land form throughout the Rocky Mountain area at the close of Mesozoic time.

Outcrops of the Wasatch Formation and the Tongue River Member of the Fort Union Formation cover most of the areas of major coal resource occurrence in the Powder River Basin. The Tongue River Member is composed of very fine-grained sandstones, siltstones, claystones, shales, carbonaceous shales, and numerous coal beds. The Lebo Member of the Fort Union Formation consists of light- to dark-gray very fine-grained to conglomeratic sandstone with interbedded siltstone, claystone, carbonaceous shale and thin coal beds. Thin bedded calcareous ironstone concretions interbedded with massive white sandstone and slightly bentonitic shale occur throughout the unit (Denson and Horn, 1975). The Lebo Member is mapped at the surface northeast of Recluse, Wyoming. Here, the Lebo Member is east of the principal coal outcrops and associated clinkers (McKay, 1974), and it presumably projects into the subsurface beneath much of the basin. One of the principal characteristics for separating the Lebo and Tullock Members (collectively referred to as the Ludlow Member east of Miles City, Montana) from the overlying Tongue River Member is the color differential between the lighter-colored upper portion and the somewhat darker lower portion (Brown, 1958). Although geologists are trying to develop criteria for subsurface recognition of the Lebo-Tullock and Tongue River-Lebo contacts through the use of subsurface data from geophysical logs, no definitive guidelines are known to have been published. Hence, for subsurface mapping purposes,

the Fort Union Formation is not divided into its members for this study.

During the Paleocene epoch, the Powder River Basin tropical to subtropical depositional environment included broad, inland flood basins with extensive swamps, marshes, freshwater lakes, and a sluggish, but active, northeastward-discharging drainage system. These features were superimposed on an emerging sea floor, near base level. Much of the vast area where organic debris collected was within a reducing depositional environment. Localized uplifts began to disturb the near sea-level terrain of northeastern Wyoming following retreat of the Cretaceous seas. However, the extremely fine-grained characteristics of the Tongue River Member clastics suggest that areas of recurring uplift peripheral to the Powder River Basin were subdued during major coal deposit formation.

The uplift of areas surrounding the Powder River Basin created a structural basin of asymmetric character, with the steep west flank located on the eastern edge of the Big Horn Mountains. The axis of the Powder River Basin is difficult to specifically define, but it is thought to be located in the western part of the Basin and to display a north-south configuration some 15 to 20 miles (24 to 32 km) east of Sheridan, Wyoming. Thus, the sedimentary section described in this report lies on the east flank of the Powder River Basin, with gentle dips of two degrees or less disrupted by surface structure thought to relate to tectonic adjustment and differential compaction.

Some coal beds in the Powder River Basin exceed 200 feet (61 m) in thickness. Deposition of these thick, in-situ coal beds requires a delicate balance between subsidence of the earth's crust and in-filling by tremendous volumes of organic debris. These conditions in concert with a favorable ground water table, non-oxidizing clear water,

and a climate amenable to the luxuriant growth of vegetation produce a stabilized swamp critical to the deposition of coal beds.

Deposition of the unusually thick coal beds of the Powder River Basin may be partially attributable to short-distance water transportation of organic detritus into areas of crustal subsidence. Variations in coal bed thickness throughout the basin relate to changes in the depositional environment. Drill hole data that indicate either the complete absence or extreme attenuation of a thick coal bed probably relate to location of the drill holes within the ancient stream channel system draining this lowland area in Early Cenozoic time. Where thick coal beds thin rapidly from the depocenter of a favorable depositional environment, it is not unusual to encounter a synclinal structure over the maximum coal thickness due to the differential compaction between organic debris in the coal depocenter and fine-grained clastics in the adjacent areas.

The Wasatch Formation of Eocene age crops out over most of the central part of the Powder River Basin and exhibits a disconformable contact with the underlying Fort Union Formation. The contact has been placed at various horizons by different workers; however, for the purpose of this report, the contact is positioned near the top of the Roland coal bed as mapped by Olive (1957) in northwestern Campbell County, Wyoming. It is considered to descend disconformably in the stratigraphic column to the top of the Wyodak-Anderson coal bed (Roland coal bed of Taff, 1909) along the eastern boundary of the coal measures. No attempt was made to differentiate the Wasatch and Fort Union Formations on geophysical logs or in the subsurface mapping program for this project.

Although Wasatch and Fort Union lithologies are too similar to allow differentiation in some areas, most of the thicker coal beds occur in the Fort Union section on the east flank of the Powder River Basin. Furthermore, orogenic movements peripheral to the basin apparently increased in magnitude during Wasatch time causing the deposition of friable, coarse-grained to gritty, arkosic sandstones, fine- to very fine-grained sandstones, siltstones, mudstones, claystones, brown-to-black carbonaceous shales, and coal beds. These sediments are noticeably to imperceptibly coarser than the underlying Fort Union clastics.

The Fats Draw Quadrangle is located in an area where surface rocks are classified as the Wasatch Formation. Olive (1957) correlated coal beds in the Spotted Horse coal field with coal beds in the Sheridan coal field (Baker, 1929) and Gillette coal field (Dobbin and Barnett, 1927), Wyoming, and with coal beds in the Ashland coal field (Bass, 1932) in southeastern Montana. This report utilizes, where possible, the coal bed nomenclature used in previous reports. The Felix coal bed was named by Stone and Lupton (1910), and the Smith coal bed was named by Taff (1909). Baker (1929) assigned names to the Anderson, Canyon, and Wall coal beds. The Cook coal bed was named by Bass (1932), and the Pawnee coal bed was named by Warren (1959).

IntraSearch's correlation of thick coal beds from the Spotted Horse coal field to Gillette points out that the Wyodak coal bed, named the "D" coal bed by Dobbin and Barnett (1927), is equivalent to the Anderson, Canyon and all, or part of the Cook coal beds to the north and west of Gillette, Wyoming. Correlation of this suite of coal beds with the Wyodak coal bed south and southwest of Gillette suggests that the Anderson and Canyon coal beds equate with the upper 10 to 25 percent of the

thick Wyodak coal bed, and the Cook and Wall or Upper Wall coal beds are equivalent to the major part of the Wyodak coal bed. Due to problematic correlations outside of the Gillette area, the name Wyodak has been informally used by many previous authors to represent the coal beds in the area surrounding the Wyodak coal mine. The Wildcat, Moyer, and Oedekoven coal beds were informally named by IntraSearch (1978b, 1979, and 1978a).

Local. The Fats Draw Quadrangle lies on the eastern flank of the Powder River Basin, where the strata generally dip gently westward. The Wasatch Formation crops out over the entire surface of the quadrangle, and is comprised of friable, coarse-grained to gritty arkosic sandstones, fine- to very fine-grained sandstones, siltstones, mudstones, claystones, brown-to-black carbonaceous shales, and coal beds. The Fort Union Formation directly underlies the Wasatch Formation, and is composed of very fine-grained sandstones, siltstones, claystones, shales, carbonaceous shales, and numerous coal beds.

Structure contours drawn on the tops of the various coal beds mapped in the quadrangle indicate a broad westward-plunging anticline extending across the southern two-thirds of the study area. A smaller northward-plunging anticlinal high also extends into the northeast quarter of the quadrangle. Strata in the western half of the quadrangle dip gently westward.

III. Data Sources

No significant coal outcrop or associated clinker is mentioned in any publication regarding the Fats Draw Quadrangle at the time of this report. It is probable that no significant coal outcrops exist within the quadrangle boundaries.

Geophysical logs from oil and gas test bores and producing wells comprise the source of subsurface control. Some geophysical logs are not applicable to this study, for the logs relate only to the deep, potentially productive oil and gas zones. More than 80 percent of the logs include resistivity, conductivity, and self-potential curves. Occasionally the suite of geophysical logs includes gamma, density, and sonic curves. These logs are available from several commercial sources.

All geophysical logs available in the quadrangle are scanned to select those with data applicable to Coal Resource Occurrence mapping. Paper copies of the logs are obtained and interpreted, and coal intervals are annotated. Maximum accuracy of coal bed identification is accomplished where gamma, density, and resistivity curves are available. Coal bed tops and bottoms are picked on the logs at the midpoint between the minimum and maximum curve deflections. The correlation of coal beds within and between quadrangles is achieved utilizing a fence diagram to associate local correlations with regional coal occurrences.

The reliability of correlations, set forth by IntraSearch in this report, varies depending on: the density and quality of lithologic and geophysical logs; the detail, thoroughness, and accuracy of published and unpublished surface geological maps; and interpretative proficiency. There is no intent on the part of IntraSearch to refute nomenclature established in the literature or used locally by workers in the area. IntraSearch's nomenclature focuses upon the suggestion of regional coal bed names applicable throughout the eastern Powder River Basin. It is expected, and entirely reasonable, that some differences of opinion regarding correlations, as suggested by IntraSearch, exist. Additional drilling for coal, oil, gas, water, and uranium, coupled with expanded mapping of coal bed outcrops and associated clinkers will

broaden the data base for coal bed correlations and allow continued improvement in the understanding of coal bed occurrences in the eastern Powder River Basin.

The topographic map of the Fats Draw Quadrangle is published by the U. S. Geological Survey, compilation date 1953. Land network and mineral ownership data are compiled from land plats available from the U. S. Bureau of Land Management in Cheyenne, Wyoming. This information is current to October 13, 1977.

IV. Coal Bed Occurrence

Wasatch and Fort Union Formation coal beds that are present in all or part of the Fats Draw Quadrangle include, in descending stratigraphic order, an unnamed local, the Upper Felix, Lower Felix, Smith, Wyodak, Pawnee, Wildcat, Moyer, and Oedekoven coal beds. A complete suite of maps (coal isopach, mining ratio where appropriate, structure, overburden isopach, areal distribution of identified resources and identified resources) is prepared for the Lower Felix, Smith and Pawnee coal beds, and for the Wyodak and Wildcat-Moyer-Oedekoven coal zones. Insufficient data, thickness, and areal extent preclude any detailed mapping of the local and Upper Felix coal beds.

No physical and chemical analyses are known to have been published regarding the coal beds in the Fats Draw Quadrangle. However, the proximate analysis performed on an "as received" basis for Campbell County coal beds is as follows:

COAL BED NAME		ASH %	FIXED CARBON %	MOISTURE %	VOLATILES %	SULFUR %	BTU/LB
	Hole						
Felix	(U) 7314C	6.623	32.727	28.925	31.726	0.595	8076
	Hole						
Smith	(U) 7312C	6.167	33.340	29.610	30.883	1.068	8215
	Hole						
Wyodak	(U) 757	6.024	32.831	26.907	34.237	0.336	8366

(U) - U. S. Geological Survey & Montana Bureau of Mines & Geology - 1974 and 1976.

The Coal Data Sheet, Plate 3, shows the down hole identification of coal beds within the quadrangle as interpreted from geophysical logs, from oil and gas test bores, and producing sites. A datum coal bed is utilized to position columnar sections on Plate 3. This portrayal is schematic by design; hence, no structural or coal thickness implications are suggested by the dashed correlation lines projected through no record (NR) intervals. Inasmuch as the Wyodak coal zone underlies the entire quadrangle, it is designated as datum for the correlation diagram. The Wyodak coal zone shows the thickest single coal bed occurrence throughout the quadrangle. The Lower Felix, Smith, Pawnee, Wildcat, Moyer and Oedekoven coal beds are relatively thin throughout most of the area.

The Lower Felix coal bed lies approximately 140 to 540 feet (43 to 165 m) beneath the surface throughout the quadrangle. The coal bed thickness ranges from 0 to 7 feet (0 to 2.1 m) with thickest occurrences located in the northeast quarter of the study area, with thinning to the southwest. The Lower Felix coal is absent from approximately 15 percent of the quadrangle in the southwest portion. Structure contours drawn on top of the Lower Felix coal bed indicate a northward-plunging anticline extending across the eastern half of the quadrangle. The western half of the study area dips gently to the west. The Lower Felix coal bed lies less than 500 feet (152 m) in depth beneath the surface throughout approximately 98 percent of the quadrangle.

The Smith coal bed lies approximately 243 to 398 feet (74 to 121 m) below the Lower Felix coal bed. The coal bed thickness ranges from 4 to 10 feet (1.2 to 3.0 m) with the coal bed thinning to less than 5 feet (1.5 m) thick throughout the central and south-central portions of the quadrangle. The Smith coal bed dips gently to the west with a broad west-

ward-plunging anticline extending across the southern half of the quadrangle. The Smith coal bed lies approximately 550 to 1070 feet (168 to 326 m) beneath the surface throughout the entire quadrangle.

The Wyodak coal zone occurs 330 to 579 feet (101 to 177 m) beneath the Smith coal bed, and is composed of a thick, locally split coal bed overlying two-to-three thin, somewhat lenticular coal beds. The total coal zone thickness ranges from 80 to 153 feet (24 to 47 m) with maximum thicknesses occurring in the northeast and southeast quarters of the quadrangle. The clastic interval separating the various coal beds comprising the coal zone varies from 6 to 290 feet (1.8 to 88 m). Structure contours drawn on top of the Wyodak coal zone indicate a regional westward dip with a broad westward-plunging anticline extending across the southern one-fourth of the study area. The Wyodak coal zone lies approximately 950 to 1625 feet (290 to 495 m) beneath the surface throughout the entire quadrangle.

The Pawnee coal bed occurs 320 to 426 feet (98 to 130 m) below the Wyodak coal zone. It occurs primarily along the northern and western boundaries of the quadrangle, showing a minor occurrence along the southeastern edge. The coal bed thickness ranges from 0 to 17 feet (0 to 5 m) with maximum thicknesses occurring in the northwest quarter of the study area. Structure contours drawn on top of the Pawnee coal bed indicate a gentle regional dip to the west. The Pawnee coal bed lies approximately 1650 to 2200 feet (503 to 671 m) beneath the surface throughout the entire quadrangle.

The Wildcat-Moyer-Oedekoven coal zone is separated from the Pawnee coal bed by from 149 to 318 feet (45 to 97 m) of clastic debris. The coal zone is composed of the thin Upper Wildcat, Lower Wildcat, Moyer, and Oedekoven coal beds. The total coal zone thickness ranges from

4 to 30 feet (1.2 to 9 m) with coals in excess of 20 feet (6 m) thick in the northwest quadrant and less than 5 feet (1.5 m) thick over the southernmost part of the quadrangle. Individually, the Wildcat coal beds are present over the entire quadrangle and contribute the greatest coal thickness to the zone. Individual Wildcat coal bed thicknesses of over 10 feet (3 m) occur. The Moyer coal bed is present in only one location and is less than 5 feet (1.5 m) thick. The Oedekoven coal bed is present over most of the northern two-thirds of the quadrangle and averages slightly over 5 feet (1.5 m) in thickness. Total non-coal interburden within the zone ranges from 0 to 312 feet (0 to 95 m). Structure contours drawn on top of the Wildcat coal bed indicate a regional dip to the west with a broad anticline extending across the southern part of the quadrangle. The Wildcat-Moyer-Oedekoven coal zone lies approximately 1850 to 2400 feet (564 to 732 m) beneath the surface throughout the entire quadrangle.

V. Geological and Engineering Mapping Parameters

The correct horizontal location and elevation of drill holes utilized in subsurface mapping are critical to map accuracy. Intra-Search Inc., plots the horizontal location of the drill hole as described on the geophysical log heading. Occasionally this location is superimposed on or near to a drillsite shown on the topographic map, and the topographic map, horizontal location is utilized. If the ground elevation on the geophysical log does not agree with the topographic elevation of the drillsite, the geophysical log ground elevation is adjusted to conformance. If there is no indication of a drillsite on the topographic map, the "quarter, quarter, quarter" heading location is shifted within a small area until the ground elevation on the heading agrees with the topographic map elevation. If no elevation agreement

can be reached, the well heading or data sheet is rechecked for footage measurements and ground elevation accuracy. Inquiries to the companies who provided the oil and gas geophysical logs frequently reveal that corrections have been made in the original survey. If all horizontal location data sources have been checked and the information accepted as the best available data, the drillsite elevation on the geophysical log is modified to agree with the topographic map elevation. IntraSearch Inc., considers this agreement mandatory for the proper construction of most subsurface maps, but in particular, the overburden isopach, the mining ratio, and Coal Development Potential maps.

Subsurface mapping is based on geologic data within, and adjacent, to the Fats Draw Quadrangle area. Data from geophysical logs are used to correlate coal beds and control contour lines for the coal thickness, structure, and overburden maps. Structure contour maps are constructed on the tops of the main coal beds.

In preparing overburden isopach maps, no attempt is made to identify coal beds that occur in the overburden above a particular coal bed under study. Mining ratio maps for this quadrangle are constructed utilizing a 95 percent recovery factor. Contours of these maps identify the ratio of cubic yards of overburden to tons of recoverable coal. Where ratio control points are sparse, interpolated points are computed at the intersections of coal bed and overburden isopach contours using coal structure, coal isopach, and topographic control. On the Areal Distribution of Identified Resources Map (ADIR), coal bed reserves are not calculated where the coal is less than 5 feet (1.5 m) thick, where the coal occurs at a depth greater than 500 feet (152 m), where non-federal coal exists, or where federal coal leases, preference-right lease applications, and coal prospecting permits exist.

Coal tonnage calculations involve the planimetering of areas of measured, indicated, inferred reserves and resources, and hypothetical resources to determine their areal extent in acres. An Insufficient Data Line is drawn to delineate areas where surface and subsurface data are too sparse for CRO map construction. Various categories of resources are calculated in the unmapped areas by utilizing coal bed thicknesses mapped in the geologically controlled area adjacent to the insufficient data line. Acres are multiplied by the average coal bed thickness and 1750, or 1770--the number of tons of lignite A or sub-bituminous C coal per acre-foot, respectively (12,874 or 13,018 metric tons per hectare-meter, respectively), to determine total tons in place. Recoverable tonnage is calculated at 95 percent of the total tons in place. Where tonnages are computed for the CRO-CDP map series, resources and reserves are expressed in millions of tons. Frequently the planimetering of coal resources on a sectionized basis involves complexly curvilinear lines (coal bed outcrop and 500-foot stripping limit designations) in relationship with linear section boundaries and circular resource category boundaries. Where these relationships occur, generalizations of complexly curvilinear lines are discretely utilized, and resources and/or reserves are calculated within an estimated 2 to 3 percent, plus or minus, accuracy.

VI. Coal Development Potential

Strippable Coal Development Potential. Areas where coal beds are 5 feet (1.5 m) or more in thickness and are overlain by 500 feet (152 m) or less of overburden are considered to have potential for surface mining and are assigned a high, moderate, or low development potential based on the mining ratio (cubic yards of overburden per ton

of recoverable coal). The formula used to calculate mining ratios for subbituminous coal is as follows:

$$MR = \frac{to (0.911)*}{tc (rf)}$$

where MR = mining ratio
to = thickness of overburden
tc = thickness of coal
rf = recovery factor
0.911* = conversion factor (cu. yds./ton)

*A conversion factor of 0.922 is used for lignite.

A surface mining development potential map (Plate 29) was prepared utilizing the following mining ratio criteria for coal beds 5 to 40 feet (1.5 to 12 m) thick:

1. Low development potential = 15:1 and greater ratio.
2. Moderate development potential = 10:1 to 15:1 ratio.
3. High development potential = 0 to 10:1 ratio.

The following mining ratio criteria are utilized for coal beds greater than 40 feet (12 m) thick:

1. Low development potential = 7:1 and greater ratio.
2. Moderate development potential = 5:1 to 7:1 ratio.
3. High development potential = 0 to 5:1 ratio.

The surface mining potential is considered low for approximately 30 percent of the Fats Draw Quadrangle. This low potential rating is scattered throughout the eastern two-thirds of the quadrangle. It is attributed to low overburden-to-coal ratios for the Lower Felix coal bed. None of the study area qualifies for high or moderate surface mining development potential. The remaining 70 percent of the quadrangle is classified as either non-federal coal land or as no development potential for surface mining. Table 1 sets forth the estimated strippable reserve base tonnages per coal bed for the quadrangle.

Underground Mining Coal Development Potential. Subsurface coal mining potential throughout the Fats Draw Quadrangle is considered low. Inasmuch as recovery factors have not been established for the underground development of coal beds in this quadrangle, reserves are not calculated for coal beds that occur more than 500 feet (152 m) beneath the surface. Table 2 sets forth the estimated coal resources in tons per coal bed.

In-Situ Gasification Coal Development Potential. The evaluation of subsurface coal deposits for in-situ gasification potential relates to the occurrence of coal beds more than 5 feet (1.5 m) thick buried from 500 to 3000 feet (152 to 914 m) beneath the surface. This categorization is as follows:

1. Low development potential relates to: 1) a total coal section less than 100 feet (30 m) thick that lies 1000 feet (305 m) to 3000 feet (914 m) beneath the surface, or 2) a coal bed or coal zone 5 feet (1.5 m) or more in thickness which lies 500 feet (152 m) to 1000 feet (305 m) beneath the surface.
2. Moderate development potential is assigned to a total coal section from 100 to 200 feet (30 to 61 m) thick and buried from 1000 to 3000 feet (305 to 914 m) beneath the surface.
3. High development potential involves 200 feet (61 m) or more of total coal thickness buried from 1000 to 3000 feet (305 to 914 m).

The coal development potential for in-situ gasification within the Fats Draw Quadrangle is moderate to low. The moderate rating covers the entire quadrangle, except for small scattered areas classified as either low or non-federal coal land. The Low potential rating covers approximately 5 percent of the quadrangle occurring as small scattered

areas in the central portion. The remaining 10 percent is classified as non-federal coal land and not evaluated for in-situ gasification. None of the quadrangle qualifies for a high in-situ gasification potential. The coal resource tonnages for in-situ gasification with moderate and low development potential are given on Table 3.

Table 1.--Strippable Coal Reserve Base Data (in short tons) for Federal Coal Lands in the Fats Draw Quadrangle, Campbell County, Wyoming.

Development potentials are based on mining ratios (cubic yards of overburden/ton of recoverable coal).

Coal Bed	High Development Potential (0-10:1 Mining Ratio)	Moderate Development Potential (10:1-15:1 Mining Ratio)	Low Development Potential ($> 15:1$ Mining Ratio)	Total
Lower Felix	-----	-----	119,800,000	119,800,000
TOTAL	-----	-----	119,800,000	119,800,000

Table 2.--Coal Resource Base and Data (in short tons) for Underground Mining Methods for Federal Coal Lands in the Fats Draw Quadrangle, Campbell County, Wyoming.

Coal Bed Name	High Development Potential	Moderate Development Potential	Low Development Potential	Total
Lower Felix	-----	-----	340,000	340,000
Smith	-----	-----	186,500,000	186,500,000
Wyodak	-----	-----	6,042,660,000	6,042,660,000
Pawnee	-----	-----	136,050,000	136,050,000
Wildcat-Moyer- Oedekoven	-----	-----	619,670,000	619,670,000
TOTAL	-----	-----	6,985,220,000	6,985,220,000

Table 3.--Coal Resource Base Data (in short tons) for In-Situ Gasification
for Federal Coal Lands in the Fats Draw Quadrangle, Campbell
County, Wyoming.

Coal Bed Name	High Development Potential	Moderate Development Potential	Low Development Potential	Total
	-----	6,542,870,000	442,350,000	6,985,220,000
TOTAL	-----	6,542,870,000	442,350,000	6,985,220,000

SELECTED REFERENCES

- Baker, A. A., 1929, The northward extension of the Sheridan coal field, Big Horn and Rosebud Counties, Montana: U. S. Geological Survey Bull. 806-B, p. 15-67.
- Bass, N. W., 1932, The Ashland coal field, Rosebud, Powder River, and Custer Counties, Montana: U. S. Geological Survey Bull. 831-B, p. 19-105.
- Brown, R. W., 1958, Fort Union Formation in the Powder River Basin, Wyoming: Wyoming Geological Association Guidebook, Thirteenth Annual Field Conf., p. 111-113.
- Denson, N. M., and Horn, G. H., 1975, Geologic and structure map of the southern part of the Powder River Basin, Converse, Niobrara, and Natrona Counties, Wyoming: U. S. Geological Survey Miscellaneous Investigations Series Map I-877, scale 1:125,000.
- Dobbin, C. E., and Barnett, V. H., 1927 (1928), The Gillette coal field, north-eastern Wyoming: U. S. Geological Survey Bull. 796-A, p. 1-50.
- Glass, G. B., 1975, Review of Wyoming coal fields, 1975: Wyoming Geological Survey Public Information Circ. 4, p. 10.
- IntraSearch Inc., 1978a, Coal resource occurrence and coal development potential of the Cabin Creek Northeast Quadrangle, Sheridan and Campbell Counties, Wyoming, and Powder River County, Montana: U. S. Geological Survey Open-File Report 78-064, 21 p.
- _____, 1978b, Coal resource occurrence and coal development potential of the Rocky Butte Quadrangle, Campbell County, Wyoming: U. S. Geological Survey Open-File Report 78-830, 22 p.
- _____, 1979, Coal resource occurrence and coal development potential of the Larey Draw Quadrangle, Campbell County, Wyoming: U. S. Geological Survey Open-File Report 79-023, 29 p.

- Jacob, A. F., 1973, Depositional environments of Paleocene Tongue River Formation: Am. Assoc. of Petroleum Geologists Bull., vol. 56, no. 6, p. 1038-1052.
- McKay, E. J., 1974, Preliminary geologic map of the Bertha 2 NW (Rocky Butte) Quadrangle, Campbell County, Wyoming: U. S. Geological Survey Open-File Report 74-173, scale 1:24,000.
- Olive, W. W., 1957, The Spotted Horse coal field, Sheridan and Campbell Counties, Wyoming: U. S. Geological Survey Bull. 1050, 83 p.
- Schell, E. M., and Mowat, G. D., 1972, Reconnaissance map showing some coal and clinker beds in the Fort Union and Wasatch Formations in the eastern Powder River Basin, Campbell and Converse Counties, Wyoming: U. S. Geological Survey Open-File Report, scale 1:63,360.
- Stone, R. W., and Lupton, C. T., 1910, The Powder River coal field, Wyoming, adjacent to the Burlington Railroad: U. S. Geological Survey Bull. 381-B, p. 115-136.
- Taff, J. A., 1909, The Sheridan coal field, Wyoming: U. S. Geological Survey Bull. 341-B, p. 123-150.
- U. S. Bureau of Mines and U. S. Geological Survey, 1976, Coal Resource classification system of the U. S. Bureau of Mines and U. S. Geological Survey: U. S. Geological Survey Bull. 1450-B, 7 p.
- U. S. Geological Survey and Montana Bureau of Mines and Geology, 1976, Preliminary report of coal drill hole data and chemical analyses of coal beds in Campbell and Sheridan Counties, Wyoming; Custer, Prairie, and Garfield Counties, Montana; and Mercer County, North Dakota: U. S. Geological Survey Open-File Report 76-319, 377 p.

- _____, 1977, Preliminary report on 1976 drilling of coal in Campbell and Sheridan Counties, Wyoming: and Big Horn, Dawson, McCone, Richland, Roosevelt, Rosebud, Sheridan and Wibaux Counties, Montana: U. S. Geological Survey Open-File Report 77-283, 403 p.
- Warren, W. C., 1959, Reconnaissance geology of the Birney-Broadus coal field, Rosebud and Powder River Counties, Montana: U. S. Geological Survey Bull. 1072-J, p. 561-585.
- Weimer, R. J., 1977, Stratigraphy and tectonics of western coals, in Geology of Rocky Mountain Coal, A Symposium, 1976: Colorado Geological Survey Resource Series 1, p. 9-27.