

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

Text to accompany:
OPEN-FILE REPORT 79-633

1985

FEDERAL COAL RESOURCE OCCURRENCE MAPS
OF THE MESITA AMERICANA 7 1/2-MINUTE QUADRANGLE,
McKINLEY COUNTY, NEW MEXICO

[Report includes 6 plates]

Prepared by Berge Exploration, Inc.

This report was prepared under contract to the U.S. Geological Survey, and has not been edited for conformity with Geological Survey editorial standards or stratigraphic nomenclature. Opinions expressed herein do not necessarily represent those of the Geological Survey.

MESITA AMERICANA QUADRANGLE CONTENTS

	Page
Introduction	1
Purpose	1
Location	4
Accessibility	4
Physiography	4
Climate	5
Land status	5
General geology	6
Previous work	6
Stratigraphy	6
Depositional environments	7
Structure	9
Coal geology	9
Menefee Cleary No. 1 coal bed	10
Coal resources	10
Coal development potential	12
Selected references	14
Glossary	15

ILLUSTRATIONS

Plates 1-6. Coal resource occurrence maps:

1. Coal data map
2. Boundary and coal data map
3. Coal data sheet
4. Isopach map of the Menefee Cleary No. 1 coal bed.
5. Structure contour map of the Menefee Cleary No. 1 coal bed.
6. Isopach map of overburden of the Menefee Cleary No. 1 coal bed.

	Page
Figure 1. Location of project area	2
2. Index to USGS 7 1/2-minute quadrangles and coal resource occurrence/coal development potential maps in the southern San Juan Basin area, New Mexico	3

TABLE

	Page
Table 1. Analyses of a coal sample from the Cleary Coal Member of the Menefee Formation	11

INTRODUCTION


Purpose

This text complements the Coal Resource Occurrence (CRO) maps of the Mesita Americana 7½ minute quadrangle, McKinley County, New Mexico. These maps and report are part of an evaluation of fifty-six 7½ minute quadrangles in northwestern New Mexico which were completed under U. S. Geological Survey Contract No. 14-08-0001-17459 (see figs. 1 and 2).

The purpose of this Coal Resource Occurrence-Coal Development Potential program, which was conceived by Congress as part of its Federal Coal Leasing Amendments Act of 1976, is to obtain coal resource information and to determine the geographical extent of Federal coal deposits. In addition, the program is intended to provide information on the amount of coal recoverable by various mining methods and to serve as a guide for land-use planning.

The U. S. Geological Survey initiated the program by identifying areas underlain by coal resources. These areas were designated Known Recoverable Coal Resource Areas based on the presence of minable coal thicknesses, adequate areal extent of these coal deposits, and the potential for developing commercial quantities of coal at minable depths.

This report is limited to coal resources which are 3,000 ft (914 m) or less below ground surface. Published and unpublished public information was used as the data base for this study. No new drilling or field mapping was performed as part of this study, nor were any confidential data used.




LOCATION OF PROJECT AREA

FIGURE 1

FIGURE 2.--Index to USGS 7 1/2-minute quadrangles and coal resource occurrence/
coal development potential maps for the southern San Juan Basin area, New Mexico

Map No.	Quadrangle	Open-file report	Map No.	Quadrangle	Open-file report
1	Cuba	79- 623	31	Nose Rock	79- 641
2	San Pablo	79- 624	32	Becenti Lake	79-1124
3	La Ventana	79-1038	33	Heart Rock	79- 642
4	Headcut Reservoir	79-1043	34	Crownpoint	79-1125
5	San Luis	79-1044	35	Antelope Lookout Mesa	79-1376
6	Arroyo Empedrado	79-1045	36	Milk Lake	79-1377
7	Wolf Stand	79-1046	37	La Vida Mission	79-1378
8	Tinian	79- 625	38	The Pillar 3 SE	79-1379
9	Canada Calladita	79- 626	39	Red Lake Well	79-1380
10	Cerro Parido	79- 627	40	Standing Rock	79-1381
11	El Dado Mesa	79- 628	41	Dalton Pass	80- 026
12	Mesa Cortada	79- 629	42	Oak Spring	80- 027
13	Mesita del Gavilan	79- 630	43	Hard Ground Flats	80- 028
14	Rincon Marquez	79- 631	44	Big Rock Hill	80- 029
15	Whitehorse Rincon	79- 632	45	Twin Lakes	80- 030
16	Mesita Americana	79- 633	46	Tse Bonita School	80- 031
17	El Dado	79- 634	47	Samson Lake	80- 032
18	Cerro Alesna	79- 635	48	Gallup West	80- 033
19	San Lucas Dam	79- 636	49	Gallup East	80- 034
20	Piedra de la Aguila	79-1039	50	Bread Springs	80- 035
21	Hospah	79- 637	51	Manuelito	80- 036
22	Whitehorse	79-1040	52	Borrego Pass	80- 037
23	Seven Lakes NE	79- 638	53	Casamero Lake	80- 038
24	Kin Nahzin Ruins	79- 639	54	Twin Buttes	80- 039
25	Orphan Annie Rock	79-1041	55	Pinehaven	80- 040
26	Mesa de los Toros	79-1122	56	Upper Nutria	80- 041
27	Laguna Castillo	79- 640			
28	Seven Lakes	79-1042			
29	Seven Lakes NW	79-1123			
30	Kin Klizhin Ruins	79-1047			



Location

The Mesita Americana 7½ minute quadrangle includes acreage in Tps. 16, 17, and 18 N., Rs. 6 and 7 W. of the New Mexico Principal Meridian, McKinley County, northwestern New Mexico (see figs. 1 and 2).

Accessibility

No paved roads pass through the Mesita Americana quadrangle. A light duty maintained road provides access to the town of San Mateo, 22 mi (35 km) southwest of the quadrangle. Dirt roads and jeep trails traverse most parts of the area. The Atchison, Topeka, and Santa Fe Railroad line passes about 41 mi (66 km) due south of the quadrangle (see fig. 1).

Physiography

The Mesita Americana quadrangle is in the Navajo section of the southernmost part of the Colorado Plateau physiographic province (U. S. Geological Survey, 1965). The topography of the quadrangle is mesa-and-canyon. Mesita Americana is a prominent land form in the northwestern part of the quadrangle.

No perennial streams are present in the quadrangle. Local drainage is provided by three intermittent arroyos: North Fork Arroyo Chico, Arroyo Lucero, and Arroyo de Alfredo Padilla. Elevations within the quadrangle range from less than 6,440 ft (1,963 m) at the eastern edge along Arroyo Lucero to 6,969 ft (2,124 m) on Mesita Americana.

Climate

The climate of this area is semiarid to arid. The following temperature and precipitation data were reported by the National Oceanic and Atmospheric Administration for the Torreon Navajo Mission Station. The Mesita Americana quadrangle is about 16 mi (26 km) WSW. of the Torreon Navajo Mission Station. Average total annual precipitation for thirteen of the previous fifteen years is 9.94 in. (25.25 cm). Intense thunderstorms in July, August, and September account for the majority of precipitation. The area is susceptible to flash flooding associated with these thunderstorms. Mean annual temperature for eleven of the previous fifteen years is 49.5°F (9.7°C). The average daily temperatures in January and July are 27.4°F (-2.6°C) and 72.7°F (22.6°C), respectively.

Land status

The Federal Government holds coal rights to approximately 15 percent of the Mesita Americana quadrangle. For the specific coal ownership boundaries, see plate 2. It is not within the scope of this report to provide detailed land-surface ownership. All but about 7,200 acres (2,914 ha) in the northern and eastern portions of the quadrangle are within the Hospah Known Recoverable Coal Resource Area. As of October 26, 1978, there were no Federal coal leases, coal preference right lease applications or coal exploration licenses within the Mesita Americana quadrangle.

GENERAL GEOLOGY

Previous work

Early reports on the area include that of Hunt (1936) who mapped a Menefee Cleary coal outcrop in the quadrangle, noting that no coal beds exceeding 14 in. (36 cm) were identified. Shomaker, Beaumont, and Kottlowski (1971) noted Hunt's outcrop, and also reported "coal beds apparently exist in the south half of sec. 19 and the north half of sec. 20 as evidenced by interpretation of photographs in T. 17 N., R. 7 W." Shomaker and Whyte (1977) estimated 19.8 million short tons (18.0 million t) of Menefee Formation coal resources in the upper 500 ft (152 m) of T. 17 N., R. 7 W. About 60 percent of the Mesita Americana quadrangle is within T. 17 N., R. 7 W.

Stratigraphy

Within the San Juan Basin, the shoreline positions of the Cretaceous seaways changed innumerable times. The overall regional alignment of the shorelines trended N. 60° W. - S. 60° E. (Sears, Hunt, and Hendricks, 1941). The transgressive and regressive shoreline migrations are evidenced by the intertonguing relationships of continental and marine facies. Rates of trough (geosynclinal) subsidence and the availability of sediment supplies are the major factors that controlled the transgressive-regressive shoreline sequences.

Exposed rock units in the Mesita Americana quadrangle include some of the sedimentary units of Upper Cretaceous age. There is Quaternary alluvium along drainages in the area. The Menefee Formation is exposed at the surface over the entire Mesita Americana quadrangle.

The Point Lookout Sandstone is a prominent sandstone marker in most of the San Juan Basin and represents nearshore or littoral deposits which formed during the most extensive northeastward retreat prior to the final withdrawal of the Cretaceous seaways in the San Juan Basin (Sears, Hunt, and Hendricks, 1941). Light gray to reddish-brown, fine-to medium-grained sandstone with interbedded shales comprises the lithologies of the unit, which ranges from 110 to 190 ft (33 to 58 m) thick locally. The continental sediments deposited inland from the beach area during deposition of the Point Lookout Sandstone compose the overlying Menefee Formation.

The Menefee Formation consists of dark gray to brown carbonaceous to noncarbonaceous shales, light gray sandstones, and coal beds, and is divisible into the basal Cleary Coal Member and the upper Allison Member. A massive channel sandstone sequence, which crops out in the northern part of this quadrangle, defines the boundary between the two members. The Allison Member was defined as the Allison Barren Member (Sears, 1925) as containing only thin, noncommercial coal beds, although the upper part of the Allison Member commonly contains commercially important coal beds in areas nearby.

All of the identified coal beds in this quadrangle are within the Cleary Coal Member, which ranges from 350 to 410 ft (107 to 125 m) thick locally. Erosion has reduced the Allison Member to about 210 ft (64 m) thick locally.

Depositional environments

The Cretaceous System sedimentary units in the quadrangle represent transgressive and regressive depositional conditions. There were

innumerable minor cycles of widely varying duration and extent within the major sedimentary sequences. The paucity of data in this quadrangle and the intended scope of this report permit only general interpretations of the depositional environments.

The Cretaceous coal deposits of the San Juan Basin are products of former coastal swamps and marshes. These swamps and marshes were supported by heavy precipitation and a climate conducive to rapid vegetal growth in moderately fresh water. Due to the relatively low sulfur contents of the San Juan Basin coals, Shomaker and Whyte (1977) suggest the coals formed in fresh water environments.

Most of the coal-bearing units were deposited in coastal plain environments. The majority of the peat deposits formed in a transition zone between lower and upper deltaic sediments during periods of relative shoreline stability. Coals also formed in lake margin swamps inland from the coastal area. Shoreline oscillations and the subsequent influx of continental or marine debris upon the peat accumulations produced the vertical buildup or "stacking" of peat deposits. This sediment debris represents variable ash contents, rock partings, and splits within the coal seams.

The peat accumulated in lenses or pods which were generally parallel to the ancient shorelines. The coals in the lower portions of the coal-bearing units represent regressive depositional conditions (Sears, Hunt, and Hendricks, 1941). The coals in the upper portions of these units are relatively sporadic in occurrence.

Structure

The Mesita Americana quadrangle is in the Chaco Slope structural division in the southern portion of the structural depression known as the San Juan Basin (Kelley, 1950). The quadrangle is northwest of the uplifted and eroded San Miguel Creek Dome. The rock units dip from 1° to 2° E. to NE. Hunt (1936) mapped an east-west trending fault in the southeast corner of the quadrangle. No localized folding has been mapped in the area.

COAL GEOLOGY

In this quadrangle, the authors identified one coal bed and one coal zone in oil and gas well logs and Hunt's (1936) surface mapping. The bed and zone are here informally called the Menefee Cleary No. 1 coal bed and the Menefee Cleary coal zone.

The Menefee Cleary No. 1 bed is the first persistent coal bed above the Point Lookout Sandstone. It occurs up to 8 ft (2 m) above the Point Lookout Sandstone in this quadrangle, although in nearby quadrangles it is up to 15 ft (5 m) above the Point Lookout Sandstone. The bed is inferred to be continuous, although it may be several individual beds that are stratigraphically equivalent.

The Menefee Cleary coal zone contains up to three beds which occur from 25 to 165 ft (8 to 50 m) above the Point Lookout Sandstone. These beds may be correlated for limited distances in portions of the area but they lack sufficient continuity with poorly defined stratigraphic position and cannot be designated as persistent coal beds.

There are no published coal quality analyses for coal beds from the Mesita Americana quadrangle. An analysis of a core sample of Cleary Coal Member beds sampled about 12 mi (19 km) west of the Mesita Americana quadrangle, has been reported by Shomaker, Beaumont, and Kottlowski (1971) and is shown in table 1. The Cleary Coal Member beds analyzed are probably similar in quality to the Cleary Coal Member beds in this quadrangle. Rank of the Cleary Coal Member seams is probably subbituminous A in this area.

Menefee Cleary No. 1 coal bed

The Menefee Cleary No. 1 bed was identified in all four drill hole logs and does not exceed 3.0 ft (0.9 m) in thickness. It is shown (plate 4) to increase to a thickness of 5.0 ft (1.5 m), based on Menefee Cleary No. 1 coal data from the northern adjacent Whitehorse Rincon quadrangle. Existence and character of the bed are unknown in the southern and eastern parts of the quadrangle because of insufficient data.

COAL RESOURCES

Because there are no identified correlative coal beds on Federal coal land with thicknesses of 3.0 ft (0.9 m) or greater, no reserve base or reserves have been calculated for the Mesita Americana quadrangle. The U. S. Geological Survey specified that only coal beds 3.0 ft (0.9 m) or greater in thickness be included in reserve base and reserve data, rather than the 28 in. (71 cm) minimum prescribed in U. S. Geological Survey Bulletin 1450-B.

Table 1.- Analysis of a coal sample from the Cleary Coal Member of the Menefee Formation.

(Core sample from sec. 36, T. 17 N., R. 10 W.)

[Form of analysis: A, as received; B, moisture free; C, moisture and ash free]
from Shomaker, Beaumont, and Kottlowski, 1971

Form of analysis	Proximate analysis (percent)			Sulfur	Heating Value (Btu/lb)
	Moisture	Volatiles Matter	Fixed Carbon		
A	16.5	33.4	40.4	9.7	0.6
B	-----	40.0	48.3	11.7	0.7
C	-----	45.3	54.7	-----	0.8
					10,070
					12,060
					13,650

Remarks:

A moist, mineral-matter-free (MMMF) calculation using the Parr formula (American Society for Testing and Materials, 1973) yields a heating value of 11,256 Btu/lb (26,181 kJ/kg). The free-swelling index of the analysis shows the sample to be nonagglomerating.

COAL DEVELOPMENT POTENTIAL

The factors used to determine the development potential are the presence of a potentially coal-bearing formation, and the thickness and overburden of correlative coal beds. The U. S. Geological Survey supplied the criteria to evaluate the coal development potential for Federal lands in this quadrangle. These criteria are based on current industry practice, U. S. Geological Survey Bulletin 1450-B, and anticipated technological advances.

Any area underlain by a potentially coal-bearing formation with 200 ft (61 m) or less of overburden has potential for surface mining. The U. S. Geological Survey designated the 200 ft (61 m) maximum depth as the strip-ping limit. Areas where a potentially coal-bearing formation is overlain by more than 200 ft (61 m) of overburden have no potential for surface mining. Areas with no correlative coal bed or a correlative coal bed less than 3.0 ft (0.9 m) in thickness and overlain by 200 ft (61 m) or less of overburden have unknown surface mining potential.

Any area underlain by a potentially coal-bearing formation with 200 to 3,000 ft (61 to 914 m) of overburden has potential for subsurface mining. Areas where a potentially coal-bearing formation is overlain by more than 3,000 ft (914 m) of overburden have no subsurface mining potential. Development potential for subsurface mining is unknown where a potentially coal-bearing formation within 200 to 3,000 ft (61 to 914 m) of the surface contains no identified correlative coal bed or a correlative coal bed less than 3.0 ft (0.9 m) thick.

Based on all presently available data, the Menefee Cleary No. 1 coal bed

does not underlie Federal coal land with thicknesses of 3.0 ft (0.9 m) or greater in the Mesita Americana quadrangle. In addition, the Menefee Formation is potentially coal-bearing in this area. Based on coal development potential criteria, all Federal coal land has unknown development potential for surface and subsurface mining methods. In situ gasification of coal has not been done on a commercial scale in the United States and criteria for rating the development potential of this method are unknown.

The coal development potential of this quadrangle is subject to revision. As further coal information becomes available, it is possible that correlative coal beds with sufficient thicknesses may be identified. These coal data will likely define areas of Federal coal lands with development potentials other than unknown.

SELECTED REFERENCES
(MESITA AMERICANA QUADRANGLE)

- American Society for Testing and Materials, 1973, Standard specification for classification of coals by rank, in American Society for Testing and Materials Standards for coal and coke: Designation D388-66, p. 54-57.
- Baltz, E. H., 1967, Stratigraphy and regional tectonic implications of part of Upper Cretaceous and Tertiary rocks, east-central San Juan Basin, New Mexico: U.S. Geological Survey Professional Paper 552, 101 p.
- Dane, C. H., 1936, The La Ventana-Chacra Mesa coal field, part 3 of Geology and fuel resources of the southern part of the San Juan Basin, New Mexico: U.S. Geological Survey Bulletin 860-C, p. 81-161.
- Hunt, C. B., 1936, The Mount Taylor coal field, part 2 of Geology and fuel resources of the southern part of the San Juan Basin, New Mexico: U.S. Geological Survey Bulletin 860-B, p. 31-80.
- Kelley, V. C., 1950, Regional structure of the San Juan Basin, in New Mexico Geological Society Guidebook of the San Juan Basin, New Mexico and Colorado, 1st Field Conference, 1950: p. 101-108.
- Keroher, G. C., and others, 1966, Lexicon of geologic names of the United States for 1936-60: U.S. Geological Survey Bulletin 1200, 4341 p.
- National Oceanic and Atmospheric Administration, 1964-78, Climatological data, New Mexico: National Climatic Center, Asheville, N.C., v. 68-82.
- Petroleum Information Well Log Library: Denver, Colo.
- Rocky Mountain Well Log Service, 1974, Catalog of electrical, radioactivity and hydrocarbon surveys: Electrical Log Services, 1974, 819 p.
- Sears, J. D., 1925, Geology and coal resources of the Gallup-Zuni Basin, New Mexico: U.S. Geological Survey Bulletin 767, 54 p.
- Sears, J. D., Hunt, C. B., and Hendricks, T. A., 1941, Transgressive and regressive Cretaceous deposits in southern San Juan Basin, New Mexico: U.S. Geological Survey Professional Paper 193-F, p. 101-121.
- Shomaker, J. W., Beaumont, E. C., and Kottowski, F. E., 1971, Strippable low-sulfur coal resources of the San Juan Basin in New Mexico and Colorado: New Mexico Bureau of Mines and Mineral Resources Memoir 25, 189p.
- Shomaker, J. W., and Whyte, M. R., 1977, Geologic appraisal of deep coals, San Juan Basin, New Mexico: New Mexico Bureau of Mines and Mineral Resources Circular 155, 39 p.
- U.S. Bureau of Mines, 1936, Analyses of New Mexico coals: U.S. Bureau of Mines Technical Paper 569, 112 p.
- U.S. Bureau of Mines and U.S. Geological Survey, 1976, Coal resource classification system of the U.S. Bureau of Mines and U.S. Geological Survey: U.S. Geological Survey Bulletin 1450-B, 7 p.
- U.S. Geological Survey, 1965, Mineral and water resources of New Mexico: New Mexico Bureau of Mines and Mineral Resources Bulletin 87, 437 p.

GLOSSARY

- coal bed--A stratified sequence of coal, composed of relatively homogeneous material, exhibiting some degree of lithologic unity and separated from the rocks above and below by physically rather well defined boundary planes.
- coal bed separation line--A line on a map plate separating areas where different coal beds or zones are mapped.
- coal bench--One of two or more divisions of a coal bed separated by rock.
- coal conversion factor--A factor used to convert acre-feet of coal into short tons of coal; bituminous coal is 1800 tons/acre-ft; subbituminous coal is 1770 tons/acre-ft.
- coal development potential--A subjective determination of the comparative potential of Federal coal lands for development of a commercially viable coal mining operation.
- coal exploration license--An area of Federal coal lands in which the licensee is granted the right, after outlining the area and the probable methods of exploration, to investigate the coal resources. An exploration license has a term not to exceed 2 years and does not confer rights to a lease.
- coal lease--An area of Federal coal lands in which the Federal Government has entered into a contractual agreement for development of the coal deposits.
- coal split--A coal bed resulting from the occurrence of a noncoal parting within the parent coal bed which divides the single coal bed into two or more coal beds.
- coal zone--A distinctive stratigraphic interval containing a sequence of alternating coal and noncoal layers in which the coal beds may so lack lateral persistence that correlating individual beds in the zone is not feasible.
- Federal coal land--Land for which the Federal Government holds title to the coal mineral rights, without regard to surface ownership.
- hypothetical resources--Undiscovered coal resources in beds that may reasonably be expected to exist in known mining districts under known geologic conditions. In general, hypothetical resources are in broad areas of coal fields where points of observation are absent and evidence is from distant outcrops, drill holes or wells. Exploration that confirms their presence and reveals quantity and quality will permit their reclassification as a Reserve or Identified Subeconomic Resource.
- identified resources--Specific bodies of coal whose location, rank, quality, and quantity are known from geologic evidence supported by engineering measurements.
- indicated--Coal for which estimates for the rank, quality, and quantity have been computed partly from sample analyses and measurements and partly from reasonable geologic projections.
- inferred--Coal in unexplored extensions of demonstrated resources for which estimates of the quality and quantity are based on geologic evidence and projections.
- isopach--A line joining points of equal bed thickness.
- Known Recoverable Coal Resource Area (KRCRA)--Formerly called Known Coal Leasing Area (KCLA). Area in which the Federal coal land is classified (1) as subject to the coal leasing provisions of the Mineral Leasing Act of 1920, as amended, and (2) by virtue of the available data being sufficient to permit evaluation as to extent, location, and potential for developing commercial quantities of coal.
- measured--Coal for which estimates for rank, quality, and quantity can be computed, within a margin of error of less than 20 percent, from sample analyses and measurements from closely spaced and geologically well known sample sites.
- mining ratio--A numerical ratio equating the in-place volumes, in cubic yards, of rocks that must be removed in order to recover 1 short ton of coal by surface mining.
- overburden--A stratigraphic interval (composed of noncoal beds and coal beds) lying between the ground surface and the top of a coal bed. For coal zones, overburden is the stratigraphic interval lying between the ground surface and the structural datum used to map the zone.
- parting--A noncoal layer occurring along a bedding plane within a coal bed.
- Preference Right Lease Application (PRLA)--An area of Federal coal lands for which an application for a noncompetitive coal lease has been made as a result of exploration done under a coal prospecting permit. PRLA's are no longer obtainable.
- quality or grade--Refers to measurements such as heat value; fixed carbon; moisture; ash; sulfur; phosphorus; major, minor, and trace elements; coking properties; petrologic properties; and particular organic constituents.
- rank--The classification of coal relative to other coals, according to degree of metamorphism, or progressive alteration, in the natural series from lignite to anthracite (Classification of coals by rank, 1973, American Society for Testing and Materials, ASTM Designation D-388-66).
- recovery factor--The percentage of total tons of coal estimated to be recoverable from a given area in relation to the total tonnage estimated to be in the Reserve Base in the ground.
- reserve--That part of identified coal resource that can be economically mined at the time of determination. The reserve is derived by applying a recovery factor to that component of the identified coal resource designated as the reserve base.
- reserve base--That part of identified coal resource from which Reserves are calculated.
- stripping limit--A vertical depth, in feet, measured from the surface, reflecting the probable maximum, practical depth to which surface mining may be technologically feasible in the foreseeable future. The rock interval, expressed in feet, above the stripping limit is the "strippable interval."
- structure contour--A line joining points of equal elevation on a stratum or bed.