




Base from U.S.G.S. Wenatchee Sheet, 1957
Limited revision 1963

Plate 2 Map showing generalized outcrop patterns and sample localities for Swauk (S), Manastash (M), Chumstick (C), Roslyn (R), Naches (N), and Wenatchee (W) Formations, Puget Group (P), isolated arkosic bodies (CS), Swauk (C) near Wenatchee (SW), and Raging River (RR) and Tokwila (T) Formations. Outcrop patterns from Taber and others (1978), and Taber and Frizzell, in prep.

This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards and nomenclature.




Plate 1 Map showing generalized outcrop patterns and sample localities for Chumstick (CH) and Huntington (H) Formations, rocks of Bolson Creek (BC), and rocks in isolated arkosic bodies (CS), Washington. Also shown: Devils Mountain Fault and a line separating rocks with very low hylt values from other Chumstick samples. Generalized geology after Huntington and others, 1961, and Eastbrook, 1976. Base from U.S.G.S. Victorville Sheet (U.S. revised 1974).

This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards and nomenclature.




Plate 3 Map showing sample localities for Puget Group (P), Blalakey (B), and Tokwila (T) Formations. Base: USGS Seattle Sheet, 1961. This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards and nomenclature.