

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

Chemical analyses and norms of 81 volcanic rocks  
from part of the Mogollon-Datil volcanic field,  
southwestern New Mexico

By  
James C. Ratté and Terry Grotbo

Open-File Report 79-1435

1979

This report is preliminary and has not been edited  
or reviewed for conformity with U.S. Geological Survey standards.

Chemical analyses and norms of 81 volcanic rocks  
from parts of the Mogollon-Datil volcanic field,  
southwestern New Mexico

By James C. Ratté and Terry Grotbo

The analyzed rocks (tables 1-3) were collected during mineral resource studies of the Gila Wilderness study area (Ratté and Gaskill, 1975) and subsequent geologic mapping, which is continuing in the Mogollon-Datil volcanic field. For this preliminary release of data, the rocks are divided into three groups:

Group I - Rocks of the early andesitic complexes that largely predate major ash-flow eruptions and related cauldron subsidence in the area considered here. These rocks range generally from 30 to 35 million years in age.

Group II - Quartz latitic or dacitic to rhyolitic ash-flow tuffs and associated intrusive and extrusive rocks. These rocks range mainly from 25 to 30 million years in age; they overlap in time with the early andesitic complexes on the one end and with the post ash-flow tuff andesites and associated rocks on the other.

Group III - Post ash-flow tuff andesites and associated rocks. Most of these rocks are between 20 and 25 million years in age, but rocks as young as 5-6 million years are included in this group.

For general characterization and classification, the rocks have been plotted on an alkali-silica diagram (fig. 1), an AFM diagram (fig. 2), and a normative-color-index vs. normative-plagioclase-composition diagram (fig. 3). The alkali-silica plot shows most of the rocks to be in the subalkaline field. Three ash-flow tuff analyses fall in the alkaline field, but are probably beyond the main discriminating part of the divider. Otherwise, a number of the andesitic rocks plot in the alkaline field close to the dividing line, but only two analyses, nos. 80 and 81, are clearly separated from the others. These analyses represent alkali olivine basalt flows interlayered in Gila Conglomerate.

The AFM diagram (fig. 2) shows nearly all of the subalkaline rocks to be in the calc-alkaline series as opposed to the tholeiitic series, and there is no clear separation of rocks of tholeiitic affinity.

Rock names applied to the analyzed rocks on the location and correlation chart were derived from figure 3.

#### References cited

- Elston, W. E., 1968, Terminology and distribution of ash flows of the Mogollon-Silver City-Lordsburg region, New Mexico: Arizona Geological Society Guidebook III to Southern Arizona, p. 231-240.
- Irvine, T. N., and Baragar, W. R. A., 1971, A guide to the chemical classification of the common volcanic rocks: Canadian Journal of Earth Sciences, v. 8, p. 523-548.
- Ratté, J. C., 1977, Geologic map of the Mogollon quadrangle, Catron County, New Mexico: U.S. Geological Survey Open-File Report 77-324.
- Ratté, J. C., and Finnell, T. L., 1978, Road log from Silver City to Reserve via Glenwood and the Mogollon mining district: New Mexico Geological Society Special Publication no. 7, p. 49-63.
- Ratté, J. C., and Gaskill, D. L., 1975, Reconnaissance geologic map of the Gila Wilderness study area, southwestern New Mexico: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-886, 2 sheets [1976].
- Rhodes, R. C., 1976, Volcanic geology of the Mogollon Range and adjacent areas, Catron and Grant Counties, New Mexico: New Mexico Geological Society Special Publication no. 5, p. 45-46.
- Shapiro, Leonard, 1967, Rapid analysis of rocks and minerals by a single solution method: U.S. Geological Survey Professional Paper 575-B, p. B187-191.
- \_\_\_\_\_, 1975, Rapid analysis of silicate, carbonate, and phosphate rocks, revised edition: U.S. Geological Survey Bulletin 1401, 76 p.
- Shapiro, Leonard, and Brannock, W. W., 1962, Rapid analysis of silicate, carbonate, and phosphate rocks: U.S. Geological Survey Bulletin 1144-A, 56 p.

## Analytical methods and analysts

### Analytical methods:

#### Code

1. Two-solution method (Shapiro and Brannock, 1962). Supplemented by atomic absorption.
2. Single-solution method (Shapiro, 1975)
3. Single-solution method (Shapiro, 1967)

### Analysts:

Artis, L.; Botts, S., Budinsky, J.; Chloe, G.; Elmore, P.;  
Glenn, J.; Kelsey, J.; Moore, R.; and Smith, H.

Table 1.--Rapid rock chemical analyses, adjusted oxides and CIPW norms for 81 volcanic rocks from part of Mogollon-Datil volcanic field, southwestern New Mexico.

EARLY ANDESITIC ROCKS

SYMBOL	1	2	3	4	5	6	7	8	9	10	11	12
S102	56.50	62.90	61.00	63.70	69.30	63.80	60.90	63.20	66.20	65.30	55.40	71.40
AL2O3	17.50	16.90	16.40	18.20	16.20	16.80	16.90	15.50	14.70	15.50	16.50	14.70
FE2O3	5.10	3.50	3.40	3.90	1.50	3.30	3.90	3.00	2.30	3.90	6.00	1.40
FeO	1.40	1.50	2.60	0.48	0.24	0.92	0.80	1.70	1.10	0.40	2.70	0.10
MgO	2.10	1.70	3.00	0.49	0.20	0.78	0.80	2.20	1.60	1.80	4.90	0.27
CaO	5.90	4.00	5.30	3.00	1.30	2.70	1.00	4.00	2.70	1.20	6.60	0.14
Na2O	4.40	3.80	3.80	4.10	4.40	4.30	4.40	4.40	3.10	3.30	3.50	3.20
K2O	2.90	2.40	2.30	3.20	3.60	3.80	4.20	2.20	4.20	5.30	1.70	5.50
H2O	1.50	1.82	2.05	1.80	1.90	2.14	3.20	2.49	3.26	2.25	1.60	2.41
TiO2	1.00	0.56	0.87	0.54	0.41	0.82	0.73	0.92	0.68	0.85	1.20	0.54
P2O5	0.42	0.21	0.23	0.34	0.16	0.28	0.23	0.22	0.16	0.18	0.32	0.08
MnO	0.08	0.08	0.09	0.08	0.02	0.06	0.08	0.07	0.05	0.06	0.11	0.13
CO2	0.35	0.04	0.01	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.06	0.05
CL	0.06	0.02	0.01	0.05	0.01	0.01	0.09	0.04	0.04	0.06	0.04	0.06
TOTAL(-O)	99.18	99.22	101.06	99.91	99.89	99.80	98.44	99.98	100.18	100.12	100.08	99.95

ADJUSTED OXIDES - H2O FREE

S102	57.84	64.58	61.61	64.93	70.72	65.33	63.94	64.83	68.31	66.72	55.91	75.20
AL2O3	17.91	17.35	16.56	18.55	16.53	17.20	17.74	15.90	15.17	15.84	16.65	15.07
FE2O3	5.22	3.39	3.43	3.98	1.53	3.38	4.09	3.08	2.37	3.98	6.06	1.44
FeO	1.43	1.54	2.63	0.49	0.24	0.94	0.84	1.74	1.13	0.41	2.73	0.10
MgO	2.15	1.75	3.03	0.50	0.82	0.80	2.10	2.26	1.65	1.84	4.95	0.28
CaO	6.04	4.11	5.35	3.06	1.33	2.76	1.05	4.10	2.79	1.23	6.66	0.14
Na2O	4.50	3.90	3.84	4.18	4.49	4.39	4.41	4.51	3.20	3.37	3.53	3.28
K2O	2.97	2.46	2.32	3.26	3.67	3.89	4.26	2.26	4.33	5.42	1.72	5.64
TiO2	1.02	0.57	0.88	0.55	0.42	0.84	0.77	0.94	0.70	0.87	1.21	0.55
P2O5	0.43	0.22	0.23	0.35	0.16	0.29	0.24	0.23	0.17	0.18	0.32	0.08
MnO	0.09	0.08	0.09	0.08	0.02	0.06	0.08	0.07	0.05	0.06	0.11	0.13
CO2	0.36	0.04	0.01	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.06	0.05
CL	0.06	0.02	0.01	0.05	0.01	0.01	0.09	0.04	0.04	0.06	0.04	0.06

NORMATIVE MINERALS - H2O FREE

O	7.438	21.779	15.607	22.021	27.135	13.891	15.870	19.090	25.717	21.815	9.939	32.195
C		1.414		3.588	3.281	1.692	4.558		0.906	2.879		1.570
OR	17.543	14.561	13.727	19.274	21.709	22.992	26.059	13.335	25.608	31.999	10.139	33.310
AB	38.114	33.012	32.476	35.362	37.918	37.180	39.091	38.114	26.760	28.530	29.218	27.759
AN	19.896	18.696	21.108	12.437	5.192	11.106	2.765	16.498	11.920	4.229	24.872	
HL					0.017	0.017			0.017		0.150	
W0	2.015		1.615					0.795			2.337	
EN	5.354	4.347	7.546	1.244	2.033	1.989	5.230	5.620	4.112	4.580	12.317	0.572
FS			0.703									
MT	1.919	3.565	4.979	0.248		0.803	0.760	3.119	1.793	5.635		
HM	3.897	0.929		3.804	1.531	2.825	3.571	0.926	1.137	3.985	2.169	1.235
IL	1.944	1.092	1.669	1.045	0.561	1.595	1.456	1.792	1.333	0.994	2.500	0.502
QU					0.123	0.679	0.572	0.535	0.391	0.345	0.765	0.289
AP	1.018	0.511	0.550	0.821	0.387	0.116	0.150	0.043	0.139	0.436	0.194	
FR	0.047	0.003		0.041		0.116	0.119	0.117	0.117	0.092	0.024	0.049
CC	0.815	0.093	0.023	0.116	0.116	0.116				0.116	0.138	
MG												0.094
TOTAL	100.001	100.001	100.001	100.001	100.003	100.001	100.001	100.001	100.001	100.001	100.001	99.994
SALIC	22.591	89.461	82.917	92.682	95.252	91.878	88.143	87.054	90.979	89.452	74.318	96.644
FEMIC	17.010	10.540	17.085	7.319	4.751	8.173	11.857	12.947	9.021	10.548	25.684	3.141

# EARLY ANDESITIC ROCKS

SYMBOL	13	14	15	16	17
SiO2	63.70	58.60	62.70	60.60	69.40
Al2O3	15.50	16.00	14.90	15.80	13.00
Fe2O3	4.90	5.00	3.40	3.70	0.73
FeO	0.24	2.00	2.00	2.20	0.88
MgO	1.20	3.60	2.70	2.80	0.68
CaO	3.30	5.20	4.40	4.50	2.10
Na2O	3.50	3.50	3.20	3.20	3.90
K2O	4.10	3.00	3.50	3.60	2.40
H2O	2.50	1.40	1.92	2.50	6.60
TiO2	0.76	0.94	0.76	0.83	0.24
P2O5	0.19	0.29	0.22	0.20	0.06
MnO	0.02	0.12	0.08	0.09	0.04
CO2	0.05	0.28	0.08	0.05	0.05
CL	0.01	0.01	0.05		
F	0.11	0.07	0.08	0.09	0.03
TOTAL (-O)	100.03	99.98	99.94	100.12	100.10

## ADJUSTED OXIDES - H2O FREE

SiO2	65.31	59.45	63.96	62.08	74.23
Al2O3	15.89	16.23	15.20	16.18	13.90
Fe2O3	5.02	5.07	3.47	3.79	0.78
FeO	0.25	2.03	2.04	2.25	0.94
MgO	1.23	3.65	2.75	2.87	0.73
CaO	3.38	5.27	4.49	4.61	2.25
Na2O	3.59	3.55	3.26	3.28	4.17
K2O	4.20	3.04	3.57	3.69	2.57
TiO2	0.78	0.95	0.78	0.85	0.26
P2O5	0.19	0.29	0.22	0.20	0.06
MnO	0.02	0.12	0.08	0.09	0.04
CO2	0.05	0.28	0.08	0.05	0.05
CL	0.01	0.01	0.05		
F	0.11	0.07	0.08	0.09	0.03

## NORMATIVE MINERALS - H2O FREE

Q	20.311	12.787	19.623	16.151	34.318
C	0.143				0.528
OR	24.841	17.983	21.099	21.792	15.169
AB	30.290	29.968	27.246	27.737	35.296
AN	14.490	19.402	16.477	18.557	10.192
HL	0.017	0.017	0.084		
WO	1.137	1.137	1.402	0.880	1.811
FN	3.064	9.095	6.860	7.143	0.739
FS					1.132
NT		4.172	4.595	5.100	
HM	5.024	2.194	0.299	0.273	
TL	0.564	1.811	1.472	1.615	0.488
RU	0.682				
AP	0.461	0.697	0.532	0.485	0.152
FR	0.196	0.092	0.127	0.152	0.034
CC	0.117	0.646	0.186	0.116	0.122
MG					
TOTAL	100.000	100.002	100.001	100.001	100.000
SALIC	90.092	80.157	84.529	84.237	95.503
FEMIC	9.908	19.844	15.473	15.764	4.498

ASH-FLOW TUFFS AND ASSOCIATED ROCKS

SYMBOL	18	19	20	21	22	23	24	25	26	27	28	29
S102	64.20	73.80	63.50	75.80	74.80	76.00	78.70	71.40	69.80	72.90	55.90	63.30
AL203	16.80	13.30	16.70	12.70	12.70	12.50	10.60	14.70	14.90	13.50	17.30	16.60
FE203	3.90	1.30	2.40	0.48	1.60	1.00	0.76	1.30	2.00	1.90	4.30	4.80
FE0	0.24	0.04	1.60	0.84	0.32	0.44	0.04	0.36	0.32	0.08	2.10	0.24
CaO	1.30	0.45	1.60	0.24	0.32	0.10	0.42	0.27	0.39	0.36	4.90	0.84
MgO	2.80	0.44	3.20	0.44	0.39	0.34	1.70	0.98	1.90	1.20	7.50	2.80
Na2O	4.90	2.40	4.60	3.00	2.90	3.30	5.90	4.50	4.00	3.60	3.20	4.50
K2O	2.80	5.80	3.70	5.10	4.50	5.10	5.90	5.10	4.80	4.60	1.80	3.60
H2O	1.00	1.50	1.30	1.10	1.30	0.93	0.90	0.85	1.10	0.85	1.50	1.80
TiO2	0.74	0.21	0.72	0.08	0.13	0.11	0.08	0.37	0.43	0.37	0.84	0.81
P2O5	0.20	0.08	0.17	0.13	0.07	0.10	0.08	0.06	0.04	0.10	0.22	0.30
MnO	0.04	0.02	0.09	0.13	0.07	0.10	0.08	0.06	0.04	0.04	0.10	0.13
Co2	0.05	0.06	0.36	0.05	0.07	0.05	0.02	0.05	0.05	0.01	0.05	0.12
CL	0.07	0.02	0.03	0.03	0.03	0.02	0.01	0.08	0.03	0.02	0.08	0.07
F	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
TOTAL (-0)	99.01	99.41	99.94	99.98	98.73	99.98	99.13	100.06	99.93	99.52	99.76	99.97
ADJUSTED OXIDES - H2O FREE												
S102	65.50	75.37	64.36	76.66	76.78	76.73	80.12	71.97	70.63	73.88	56.89	64.51
AL203	17.14	13.58	16.93	12.84	13.04	12.62	10.79	14.82	15.08	13.68	17.61	16.92
FE203	3.98	1.33	2.43	0.49	1.64	1.01	0.77	1.31	2.02	1.93	4.38	4.99
FE0	0.24	0.04	1.62	0.85	0.33	0.44	0.04	0.36	0.32	0.08	2.14	0.24
MgO	1.33	0.46	1.62	0.24	0.33	0.10	0.43	0.27	0.39	0.36	4.99	0.84
CaO	2.86	0.45	3.24	0.44	0.40	0.34	1.73	0.99	1.92	1.22	7.63	2.85
Na2O	5.00	2.45	4.66	3.03	2.98	3.33	5.93	4.54	4.05	3.65	3.26	4.59
K2O	2.86	5.92	3.75	5.16	4.62	5.15	6.01	5.14	4.86	4.66	1.83	3.67
TiO2	0.76	0.21	0.73	0.08	0.13	0.11	0.08	0.37	0.44	0.37	0.85	0.83
P2O5	0.20	0.08	0.17	0.13	0.07	0.10	0.08	0.06	0.04	0.10	0.22	0.31
MnO	0.04	0.02	0.09	0.13	0.07	0.10	0.08	0.06	0.04	0.04	0.10	0.13
Co2	0.05	0.06	0.36	0.05	0.07	0.05	0.02	0.05	0.05	0.01	0.05	0.12
CL	0.07	0.02	0.03	0.03	0.03	0.02	0.01	0.08	0.03	0.02	0.08	0.07
F	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
NORMATIVE MINERALS - H2O FREE												
Q	18.275	37.238	15.054	37.614	41.071	36.961	46.446	23.963	24.487	32.018	9.759	17.749
C	1.380	2.696	0.599	1.660	2.948	1.113	0.916	0.480	0.257	0.719	0.438	1.348
OR	16.882	35.005	22.162	30.479	27.294	30.426	35.494	30.378	28.701	27.549	10.825	21.690
AB	42.304	20.741	39.454	25.673	25.187	28.191	14.645	38.382	34.248	30.872	27.558	38.823
AN	12.127	1.212	12.549	1.666	0.522	1.236	1.439	3.577	7.925	5.225	28.013	11.069
HL												
W0	3.303	1.145	4.039	0.605	0.818	0.251	0.193	0.678	0.983	0.909	3.182	0.404
EN				0.704		1.439	0.131	0.286	2.024	1.926	4.742	0.530
FS			3.409	0.704	1.642	0.017	0.683	1.113	0.770	0.258	1.105	0.724
MT	3.979	1.328	0.081	0.154		0.211		0.708			1.624	0.894
HM	0.604	0.130	1.386									0.900
IL												
TN	0.437	0.146	0.408	0.062	0.316	0.041	0.006	0.167	0.029	0.239	0.530	0.404
PU	0.483	0.194	0.031	0.062	0.039	0.041	0.006	0.153	0.029	0.023	0.126	0.091
FR	0.109	0.027	0.830	0.115	0.163	0.115	0.046	0.115	0.115	0.023	0.116	0.278
CC	0.116	0.139										
MG												
TOTAL	100.000	100.000	100.001	100.002	100.000	100.001	100.000	100.001	100.000	100.000	100.001	100.002
SALIC	90.968	96.892	89.817	97.092	97.021	97.926	98.940	96.781	95.619	96.383	76.155	90.677



FEMIC 9.032 3.108 10.185 2.909 2.979 2.075 1.060 3.220 4.382 3.618 23.846 9.325

ASH-FLOW TUFFS AND ASSOCIATED ROCKS

SYMBOL	30	31	32	33	34	35	36	37	38	39	40	41
SI02	74.90	76.50	76.10	75.70	75.50	65.00	68.70	64.30	66.30	75.20	71.80	71.90
AL2O3	13.00	12.10	11.70	12.90	12.60	17.10	14.70	17.20	16.70	12.70	13.80	13.80
FE2O3	1.00	0.74	1.30	0.90	0.87	2.80	2.20	2.60	2.10	1.00	2.00	2.00
FeO	0.20	0.24	0.02	0.36	0.12	0.36	0.48	0.28	0.24	0.28	0.28	0.28
MgO	0.23	0.11	0.24	0.10	0.07	0.59	0.57	0.62	0.47	0.16	0.18	0.42
CaO	0.48	0.30	0.32	0.29	0.20	1.20	1.70	1.50	2.00	0.18	0.40	0.71
Na2O	3.30	3.50	1.30	3.00	1.30	4.20	3.80	5.20	4.60	3.80	2.60	3.20
K2O	5.10	4.60	6.00	5.20	6.90	5.60	4.80	6.00	5.50	5.10	7.40	5.70
H2O	0.70	0.69	2.60	0.73	1.10	2.07	2.30	0.54	0.98	0.42	0.64	0.95
TiO2	0.22	0.14	0.21	0.27	0.07	0.75	0.57	0.73	0.49	0.13	0.48	0.49
P2O5	0.05	0.03	0.02	0.02	0.07	0.14	0.16	0.19	0.10	0.02	0.13	0.11
MnO	0.06	0.05	0.07	0.03	0.07	0.07	0.04	0.09	0.12	0.05	0.03	0.06
CO2	0.05	0.05	0.05	0.05	0.02	0.05	0.05	0.05	0.35	0.05	0.05	0.10
CL	0.01	0.01	0.05	0.05	0.02	0.01	0.03	0.01	0.07	0.03	0.08	0.08
F	0.25	0.13	0.05	0.03	0.02	0.13	0.09	0.13	0.07	0.03	0.08	0.08
TOTAL (-O)	99.44	99.13	99.76	99.57	98.76	100.01	100.15	99.34	99.99	99.31	99.84	99.77
ADJUSTED OXIDES - H2O FREE												
SI02	75.85	77.71	78.16	76.59	77.31	66.37	70.21	65.08	66.96	76.05	72.38	72.76
AL2O3	13.17	12.29	12.02	13.05	12.90	17.46	15.02	17.41	16.87	12.84	13.91	13.97
FE2O3	1.01	0.75	1.34	0.91	0.89	2.86	2.25	2.63	2.12	1.01	2.02	2.02
FeO	0.20	0.24	0.02	0.36	0.12	0.37	0.49	0.24	0.24	0.28	0.28	0.28
MgO	0.23	0.11	0.25	0.10	0.07	0.60	0.58	0.63	0.47	0.16	0.18	0.43
CaO	0.49	0.30	0.33	0.29	0.20	1.23	1.74	1.52	2.02	0.18	0.40	0.72
Na2O	3.34	3.56	1.34	3.04	1.33	4.29	3.88	5.26	4.65	3.94	2.62	3.24
K2O	5.16	4.67	6.16	5.26	7.07	5.72	4.91	6.07	5.55	5.36	7.46	5.77
TiO2	0.22	0.14	0.22	0.27	0.07	0.77	0.58	0.74	0.49	0.13	0.48	0.50
P2O5	0.05	0.03	0.02	0.02	0.07	0.14	0.16	0.19	0.10	0.02	0.13	0.11
MnO	0.06	0.05	0.07	0.03	0.07	0.07	0.04	0.09	0.12	0.05	0.03	0.06
CO2	0.05	0.05	0.05	0.05	0.02	0.05	0.05	0.05	0.35	0.05	0.05	0.10
CL	0.01	0.01	0.05	0.05	0.02	0.01	0.03	0.01	0.07	0.03	0.08	0.08
F	0.25	0.13	0.05	0.03	0.02	0.13	0.09	0.13	0.07	0.03	0.08	0.08
NORMATIVE MINERALS - H2O FREE												
Q	36.366	39.013	46.096	38.307	42.284	16.992	25.259	8.132	15.090	32.837	28.178	30.492
C	2.092	1.383	2.854	2.073	2.949	2.786	0.928	0.316	0.765	0.631	1.408	1.780
OR	30.521	27.613	36.417	31.090	41.750	33.787	28.989	35.885	32.826	31.672	44.085	34.086
AN	28.204	30.009	11.299	23.684	11.264	36.210	32.636	44.459	39.313	32.516	22.179	27.402
HL	0.017	0.045	0.809	0.795	0.315	3.943	6.661	5.118	6.675	0.242	0.321	1.677
W0		0.017				0.017	0.051	0.017				
EN	0.558	0.278	0.614	0.252	0.179	1.500	1.451	1.563	1.182	0.403	0.452	1.039
FS												
MT	0.205	0.539		0.481	0.396		0.026			0.697		
HM	0.871	0.380	1.335	0.579	0.618	2.859	2.230	2.631	2.121	0.531	2.016	2.024
IL	0.423	0.270	0.197	0.519		0.929	1.106	0.708	0.771	0.250	0.661	0.778
TH												
RU			0.112			0.276		0.366	0.089		0.136	0.112
AP	0.120	0.072	0.049	0.048	0.170	0.339	0.387	0.455	0.239	0.048	0.310	0.264
FP	0.511	0.266	0.102	0.059	0.029	0.247	0.159	0.235	0.127	0.059	0.142	0.146
CC	0.094	0.116	0.117	0.115	0.047	0.116	0.116	0.115	0.804	0.115	0.115	0.230
MG	0.018											
TOTAL	100.001	100.001	100.001	100.000	100.000	100.001	100.001	100.001	100.001	100.001	100.000	100.001
SALIC	97.200	98.080	97.475	97.948	98.562	93.735	94.524	93.927	94.668	97.899	96.169	95.458

FEMIC	2.801	1.921	2.525	2.052	1.437	6.266	5.476	6.074	5.333	2.102	3.832	4.563
-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

ASH-FLOW TUFFS AND ASSOCIATED ROCKS

SYMBOL	42	43	44	45	46	47	48	49	50	51	52	53
	*	*	*	*	*	*	*	*	*	*	*	*
ST02	65.00	72.10	74.40	76.60	77.20	75.90	77.60	73.40	75.70	73.90	75.50	76.50
AL203	15.30	12.70	12.40	12.60	12.10	12.30	11.80	11.20	13.00	12.00	12.80	11.50
FE2O3	3.90	2.00	1.30	0.75	0.77	0.63	0.77	1.70	0.62	1.10	0.27	0.82
FE0	0.32	0.48	0.04	0.32	0.48	0.52	0.08	0.12	0.20	0.60	0.44	0.04
MgO	1.20	0.41	0.08	0.09	0.08	0.15	0.14	0.50	0.11	0.77	0.06	0.22
CaO	1.70	0.38	0.49	0.18	0.23	0.21	0.42	0.85	0.24	1.70	0.47	0.67
Na2O	3.10	2.20	3.20	3.40	3.40	3.70	2.70	3.10	4.00	1.30	3.50	1.70
K2O	6.00	7.70	5.00	4.80	4.90	5.20	5.20	6.00	4.40	4.30	4.80	7.80
H2O	1.29	1.24	0.79	0.83	0.49	0.95	0.90	0.57	0.82	3.80	0.89	0.76
TiO2	1.10	0.51	2.10	0.18	0.11	0.21	0.16	0.07	0.10	0.18	0.11	0.17
P2O5	0.30	0.13								0.16		
MnO	0.11	0.04	0.07	0.06	0.09	0.07		0.04	0.08	0.04	0.03	0.04
CO2	0.62	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
CL												
F	0.10	0.03	0.10	0.01	0.02	0.03	0.02	0.04	0.03	0.03	0.03	0.01
TOTAL(-G)	100.00	99.96	99.98	99.87	99.91	99.91	100.03	99.96	99.36	99.92	99.07	100.08

ADJUSTED OXIDES - H2O FREE

ST02	65.85	73.04	75.01	77.35	77.65	76.70	78.48	73.85	76.82	76.89	76.90	76.83
AL203	15.50	12.87	12.50	12.72	12.17	12.43	11.90	13.28	13.19	12.48	13.04	11.58
FE2O3	3.95	2.03	1.31	0.76	0.77	0.64	0.78	1.71	0.63	1.14	0.27	0.81
FE0	0.32	0.49	0.04	0.32	0.48	0.53	0.08	0.12	0.20	0.62	0.45	0.04
MgO	1.22	0.42	0.08	0.09	0.08	0.15	0.14	0.50	0.11	0.80	0.06	0.22
CaO	1.72	0.38	0.49	0.18	0.23	0.21	0.42	0.86	0.24	1.77	0.48	0.67
Na2O	3.14	2.23	3.23	3.43	3.42	3.74	2.72	3.12	4.06	1.35	3.67	1.71
K2O	6.08	7.80	5.04	4.85	4.93	5.25	5.25	6.04	4.47	4.47	4.89	7.85
TiO2	1.11	0.52	2.12	0.18	0.11	0.21	0.16	0.34	0.10	0.19	0.11	0.17
P2O5	0.30	0.13						0.07	0.07	0.17		
MnO	0.11	0.04	0.07	0.06	0.09	0.07		0.04	0.08	0.04	0.03	0.04
CO2	0.63	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
CL												
F	0.10	0.03	0.10	0.01	0.02	0.03	0.02	0.04	0.03	0.03	0.01	0.01

NORMATIVE MINERALS - H2O FREE

O	21.626	29.349	36.230	38.473	38.376	34.295	41.648	30.466	35.824	47.577	36.013	35.690
C	3.005	0.539	1.227	1.643	0.960	0.404	1.146	0.437	1.482	2.765	1.076	1.076
OR	35.920	46.093	29.788	28.641	29.124	31.052	30.997	35.672	26.386	26.436	28.890	46.410
AN	26.575	18.858	27.299	29.050	28.937	31.638	23.047	26.391	34.123	11.445	30.574	16.484
AR	2.045	0.593	1.394	0.509	0.682	0.511	1.635	3.216	0.665	7.238	1.978	0.717
HL									0.050		0.101	
W0												
EN	3.028	1.034	0.201	0.226	0.200	0.378	0.352	1.253	0.278	1.995	0.152	0.817
FS					0.232	0.220					0.468	0.552
MT	0.202			0.712	1.123	0.923			0.625	1.605	0.399	
HM	3.951	1.887	1.311	0.266			0.777	1.710	0.198	0.038		0.826
IL	0.923	0.981	0.236	0.345	0.210	0.403	0.170	0.341	0.193	0.356	0.213	0.199
TN												
RU	0.628		1.993				0.072	0.162		0.394		
AP	0.720							0.167				
FR	0.152	0.038	0.207	0.021	0.041	0.062	0.041	0.070	0.063	0.034	0.021	0.021
CC	1.428	0.115	0.115	0.115	0.114	0.115	0.115	0.114	0.115	0.118	0.116	0.114
M6												
TOTAL	100.001	100.000	100.001	100.001	100.001	100.001	100.000	100.000	100.001	100.000	100.001	100.000
SALIC	89.170	95.431	95.936	98.315	98.080	97.900	98.473	96.183	98.529	95.461	98.533	97.301

FEMIC	10.831	4.570	4.062	1.685	1.922	2.101	1.527	3.818	1.472	4.539	1.308	2.699
-------	--------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

ASH-FLOW TUFFS AND ASSOCIATED ROCKS

SYMBOL 54  
\*

SiO2 77.80  
Al2O3 12.30  
Fe2O3 0.42  
FeO 0.16  
MgO 0.24  
CaO 0.31  
Na2O 2.40  
K2O 4.30  
H2O 1.57  
TiO2 0.13  
P2O5  
MnO  
CO2 0.05  
CL  
F 0.02  
TOTAL (-O) 99.69

ADJUSTED OXIDS - H2O FREE

SiO2 79.29  
Al2O3 12.54  
Fe2O3 0.43  
FeO 0.16  
MgO 0.24  
CaO 0.32  
Na2O 2.45  
K2O 4.38  
TiO2 0.13  
P2O5  
MnO  
CO2 0.05  
CL  
F 0.02

NORMATIVE MINERALS - H2O FREE

C 47.453  
C 3.367  
OR 25.896  
AB 20.697  
AN 1.096  
HL  
WO  
EN 0.609  
FS  
MT 0.142  
HM 0.330  
IL 0.252  
TN  
RU  
AP  
FR 0.042  
CC 0.116  
MG  
TOTAL 100.000  
SALIC 98.509

FEMIC 1.491

# POST ASH-FLOW TUFF ANDESITES AND ASSOCIATED ROCKS

SYMBOL	55	56	57	58	59	60	61	62	63	64	65	66
SiO2	57.10	52.90	57.20	57.30	59.80	52.10	54.10	49.30	58.30	51.90	50.00	48.10
Al2O3	13.70	16.30	15.80	16.40	15.60	16.60	16.10	17.60	16.80	17.60	14.90	14.90
Fe2O3	5.50	6.30	3.20	5.80	3.00	8.30	9.00	7.40	2.70	9.30	8.70	10.00
FeO	0.88	1.90	4.20	1.60	3.30	0.80	0.92	3.50	3.80	1.00	1.70	1.50
MnO	4.70	4.90	3.60	2.90	2.80	4.50	4.10	4.50	3.30	4.10	6.40	6.00
CaO	5.60	7.00	5.90	5.80	5.10	6.60	6.70	9.50	5.50	6.70	7.90	7.90
Na2O	2.60	3.60	3.60	3.60	3.50	3.70	3.20	2.80	3.70	3.60	2.70	2.90
K2O	5.40	2.60	2.60	2.90	1.90	2.40	2.00	1.10	2.40	2.30	1.80	2.00
H2O	2.40	2.40	1.07	1.96	3.30	3.00	2.60	1.40	1.61	3.50	3.80	4.50
TiO2	1.70	1.00	0.89	1.00	0.72	1.70	1.80	2.00	1.30	1.80	1.40	1.60
P2O5	0.69	0.74	0.54	0.65	0.42	0.81	0.74	0.41	0.43	0.69	0.47	0.50
MgO	0.10					0.10	0.09	0.15	0.09	0.12	0.16	0.14
CO2	0.05	0.06	0.07	0.01	0.01	0.02	0.01	0.22	0.05	0.05	0.05	0.05
CL									0.01			
F	0.12	0.06	0.04	0.07	0.04	0.03	0.03	0.02	0.05	0.13	0.09	0.07
TOTAL(-O)	99.89	99.73	98.69	99.96	99.47	100.65	101.38	99.89	100.02	99.84	100.04	100.13
ADJUSTED OXIDES - H2O FREE												
SiO2	58.57	54.35	58.59	58.47	62.18	53.36	54.77	50.06	59.24	53.87	51.95	50.30
Al2O3	14.05	16.75	16.18	16.73	16.22	17.00	16.30	17.87	17.07	15.26	15.48	15.58
Fe2O3	5.64	6.47	3.28	5.92	3.12	8.50	9.11	7.51	2.74	9.65	9.04	10.46
FeO	0.90	1.95	4.30	1.43	3.43	0.82	0.93	3.55	3.86	1.04	1.77	1.57
MgO	4.82	5.03	3.69	2.96	2.91	4.61	4.15	4.57	3.35	4.26	6.65	6.27
CaO	5.74	7.19	6.04	5.92	5.30	6.76	6.78	9.65	5.59	6.95	8.21	8.26
Na2O	2.67	3.70	3.69	3.67	3.64	3.79	3.24	2.84	3.76	3.74	2.81	3.03
K2O	5.54	2.67	2.66	2.96	1.98	2.46	2.02	1.12	2.44	2.39	1.87	2.09
TiO2	1.13	1.03	0.91	1.02	0.75	1.74	1.82	2.03	1.32	1.87	1.45	1.67
P2O5	0.71	0.76	0.55	0.66	0.44	0.83	0.75	0.42	0.44	0.72	0.49	0.52
MnO	0.10					0.10	0.09	0.15	0.09	0.12	0.17	0.15
CO2	0.05	0.06	0.07	0.01	0.01	0.02	0.01	0.22	0.05	0.05	0.05	0.05
CL									0.01		0.01	
F	0.12	0.06	0.04	0.07	0.04	0.03	0.03	0.02	0.05	0.13	0.09	0.07
NORMATIVE MINERALS - H2O FREE												
Q	7.521	3.982	9.914	11.582	18.161	4.220	10.621	5.931	11.661	6.553	5.443	2.396
C												
OR	32.732	15.785	15.738	17.487	11.674	16.524	11.965	6.600	14.612	14.108	11.052	12.359
AB	22.567	31.296	31.204	31.084	30.794	32.065	27.613	26.056	31.740	31.621	23.662	23.660
AN	10.014	21.203	19.743	20.433	22.089	22.119	23.952	32.699	22.542	17.811	24.168	22.725
NE												
HL									0.017		0.017	
MO	5.455	3.807	2.577	1.854	0.535	1.410	0.985	4.601	0.803	3.827	5.290	5.978
FN	12.007	12.538	9.184	7.370	7.251	11.477	10.338	11.379	8.352	10.600	16.562	15.626
FS			3.687		2.488				2.813			
FO												
FA												
MT		3.314	4.753	2.304	4.523			6.065	3.978		2.020	0.684
HM	5.642	4.187	4.329	4.329		8.500	9.111	3.331		9.654	7.647	9.985
IL	2.126	1.951	1.938	1.938	1.422	1.949	2.162	3.857	2.509	2.459	2.763	3.178
TN	0.022					1.753	1.678			1.408		
PF												
RU												
AP	1.676	1.801	1.310	1.571	1.034	1.965	1.774	0.986	1.035	1.696	1.157	1.238
FR	0.123			0.025	0.005				0.024	0.146	0.103	0.055



CC	0.117	0.140	0.163	0.023	0.024	0.047	0.023	0.508	0.116	0.118	0.119
TOTAL	100.001	100.004	100.005	100.000	100.000	100.026	100.022	100.012	100.001	100.002	100.002
SALIC	72.834	72.266	76.599	80.585	82.719	72.925	73.951	69.285	80.372	64.343	63.139
FEMIC	27.167	27.738	23.406	19.415	17.281	27.101	26.071	30.726	19.629	35.659	36.862

# POST ASH-FLOW TUFF ANDESITES AND ASSOCIATED ROCKS

SYMBOL	67	68	69	70	71	72	73	74	75	76	77	78
SI02	48.00	52.30	55.10	58.20	55.10	62.80	48.80	53.60	53.40	59.40	49.80	62.90
AL2O3	16.50	15.00	16.90	16.10	17.40	17.70	17.70	15.80	16.00	17.40	15.30	15.40
FE2O3	11.70	9.00	4.20	2.90	5.00	2.10	5.20	6.90	3.90	4.40	5.70	4.90
FE0	0.84	1.50	2.70	3.90	0.60	2.40	6.00	1.60	4.80	0.86	7.50	0.44
MGO	4.90	3.70	3.80	3.40	3.50	1.80	4.90	3.90	5.20	2.60	3.40	2.30
CAO	7.30	6.90	6.00	3.50	6.20	5.10	9.20	3.20	6.90	5.90	7.10	4.60
MA2O	2.90	3.70	4.20	3.80	2.80	3.70	2.70	3.00	3.50	4.40	3.80	3.70
K2O	1.70	2.00	2.60	3.50	1.10	2.40	0.86	2.70	2.20	2.00	1.60	2.80
H2O	3.20	2.20	1.60	2.64	7.00	1.29	1.88	2.70	1.30	1.40	1.20	1.86
TI02	2.10	1.90	1.30	0.89	0.72	0.65	2.00	1.50	1.40	0.58	2.60	0.80
P2O5	0.63	0.67	0.55	0.24	0.30	0.19	0.35	0.62	0.58	0.30	0.66	0.42
MNO	0.16	0.13	0.11	0.16	0.04	0.06	0.14	0.09	0.11	0.08	0.12	0.10
CO2	0.05	0.05	0.05	0.74	0.01	0.02	0.05	0.08	0.35	0.01	0.10	0.01
CL		0.02										
F	0.06	0.12	0.06	0.03		0.09	0.02	0.04	0.06	0.03	0.04	0.04
TOTAL (-O)	100.01	100.13	99.14	99.99	99.77	100.26	99.79	98.71	99.67	99.35	98.90	100.25
ADJUSTED OXIDES - H2O FREE												
SI02	49.58	53.40	56.49	59.79	59.39	63.45	49.84	55.83	54.28	60.64	50.97	63.93
AL2O3	17.04	16.34	17.33	16.54	18.76	17.88	18.08	16.46	16.26	17.76	15.66	15.65
FE2O3	12.08	9.19	4.31	2.98	5.39	2.12	5.31	7.19	3.96	4.49	5.83	4.98
FE0	0.87	1.53	2.77	4.01	0.65	2.42	6.13	1.67	4.88	0.88	7.68	0.45
MGO	5.06	3.78	3.90	3.49	3.77	1.82	5.00	4.06	5.29	2.65	3.48	2.34
CAO	7.54	7.05	6.15	3.60	6.68	5.15	9.40	6.46	7.01	6.02	7.27	4.68
MA2O	3.00	3.78	4.31	3.90	3.02	3.74	2.76	3.12	3.56	4.49	3.89	3.76
K2O	1.76	2.04	2.67	3.60	1.19	2.42	0.88	2.81	2.24	2.04	1.64	2.55
TI02	2.17	1.94	1.33	0.91	0.78	0.66	2.04	1.56	1.42	0.59	2.66	0.81
P2O5	0.65	0.68	0.56	0.25	0.32	0.19	0.36	0.65	0.59	0.31	0.68	0.43
MNO	0.17	0.13	0.11	0.16	0.04	0.06	0.14	0.09	0.11	0.08	0.12	0.10
CO2	0.05	0.05	0.05	0.76	0.01	0.02	0.05	0.08	0.36	0.01	0.10	0.01
CL		0.02										
F	0.06	0.12	0.06	0.03		0.09	0.02	0.04	0.06	0.03	0.04	0.04
NORMATIVE MINERALS - H2O FREE												
Q	4.789	6.999	5.393	11.529	18.298	18.718	4.356	10.067	4.644	11.931	3.460	19.528
C				2.064	1.156	0.445						
OR	10.376	12.068	15.751	21.246	7.007	14.330	5.190	16.618	13.215	12.066	9.677	16.816
A9	25.346	31.817	36.434	33.031	25.559	31.634	23.334	26.439	30.105	38.012	32.910	31.820
AN	27.871	21.668	20.075	11.356	30.974	23.642	34.334	22.571	21.804	22.277	20.434	17.422
NE												
HL	0.034											
MO	0.578	3.231	2.652				4.009	1.970	2.852	2.303	4.403	0.918
EN	12.605	9.409	9.702	8.698	9.396	4.529	12.464	10.116	13.165	6.611	8.667	5.822
FS				3.692		1.728	3.758		3.543		5.110	
FO												
FA												
MT			5.426	4.319		3.076	7.700	1.149	5.748	1.380	8.459	
HM	12.085	9.190	0.563		5.390			6.394		3.540		4.980
IL	2.186	3.519	2.531	1.736	1.458	1.247	3.879	2.967	2.703	1.125	5.054	
TN	2.498	0.214										0.494
PF												
RU					0.008							
AP	1.541	1.620	1.336	0.584	0.766	0.455	0.847	1.530	1.396	0.725	1.600	1.011
FR	0.009	0.126	0.023	0.018		0.152			0.017	0.007		0.005

CC	0.117	0.116	0.117	1.729	0.025	0.046	0.116	0.189	0.809	0.023	0.233	0.023
TOTAL	100.002	100.002	100.001	100.002	100.017	100.001	100.008	100.010	100.001	100.001	100.013	100.001
SALIC	68.383	72.576	77.652	79.225	82.974	88.767	67.234	75.695	69.768	84.286	66.482	85.586
FEMIC	31.619	27.426	22.350	20.777	17.043	11.234	32.774	24.316	30.233	15.715	33.531	14.415

# POST ASH-FLOW TUFF ANDESITES AND ASSOCIATED ROCKS

SYMBOL	79	80	81
	•	•	•
SiO <sub>2</sub>	51.20	47.50	47.40
Al <sub>2</sub> O <sub>3</sub>	15.50	14.60	15.00
Fe <sub>2</sub> O <sub>3</sub>	2.70	6.50	12.10
FeO	8.20	6.10	1.20
MgO	7.00	5.90	6.00
CaO	8.20	8.20	7.50
Na <sub>2</sub> O	3.50	4.40	4.20
K <sub>2</sub> O	1.00	2.20	2.10
H <sub>2</sub> O	0.79	1.00	1.30
TiO <sub>2</sub>	1.40	2.00	2.10
P <sub>2</sub> O <sub>5</sub>	0.38	0.69	0.75
MnO	0.10	0.20	0.21
CO <sub>2</sub>	0.02	0.32	0.05
CL		0.09	
F	0.03	0.07	
TOTAL(-O)	99.81	99.72	99.91
ADJUSTED OXIDES - H <sub>2</sub> O FREE			
SiO <sub>2</sub>	51.71	48.12	48.07
Al <sub>2</sub> O <sub>3</sub>	15.65	14.79	15.21
Fe <sub>2</sub> O <sub>3</sub>	2.73	6.58	12.27
FeO	8.28	6.12	1.22
MgO	7.07	5.98	6.08
CaO	8.28	8.31	7.61
Na <sub>2</sub> O	3.33	4.46	4.26
K <sub>2</sub> O	1.01	2.23	2.13
TiO <sub>2</sub>	1.41	2.03	2.13
P <sub>2</sub> O <sub>5</sub>	0.38	0.70	0.76
MnO	0.10	0.20	0.21
CO <sub>2</sub>	0.02	0.32	0.05
CL		0.09	
F	0.03	0.07	
NORMATIVE MINERALS - H <sub>2</sub> O FREE			
Q			
C			
OR	5.968	13.169	12.584
AB	28.201	28.948	33.785
AN	24.770	14.123	16.098
HE		4.384	1.222
HL		0.150	
W0	5.711	8.520	6.045
EN	16.115	6.402	5.224
FS	9.892	1.264	
FO	1.045	5.944	6.958
FA	0.707	1.293	
MT	3.954	9.547	
HM			12.271
IL	2.685	3.848	3.026
TN			
PF			0.913
RU			
AP	0.909	1.656	1.801
FR		0.018	

CC	0.046	0.737	0.115
TOTAL	100.004	100.003	100.042
MALIC	58.939	60.774	63.689
FEMIC	41.065	39.228	36.353

Table 2: Location and correlation of analyzed samples and key to analytical methods; ~~Informal~~ Geologic names used here are largely from Ratte and Gaskill, 1975; Ratte, 1977; or are related to geologic mapping in progress, except as otherwise noted.

Sample No.	Field No.	Quadrangle	Latitude	Longitude	Rock Name	Correlation	Analytical methods code
1/	SP-33-75	Saliz Pass	33° 33' 55" N	108° 53' 50" W	andesite	Older andesitic complex north and west of Mogollon	1
2/	SP-7E-76	Saliz Pass	33° 34' 06" N	108° 53' 05" W	andesite	"	1
3/	DM-3	Dillon Mtn.	33° 51' 12" N	108° 48' 41" W	andesite	"	1
4	GR-88A	Canyon Hill	33° 03' 36" N	108° 29' 28" W	dacite	Volcanic complex of Brock Canyon	2
5	GR-79C	Canteen Canyon	33° 04' 06" N	108° 31' 05" W	rhyolite	"	2
6	GR-76A	Canteen Canyon	33° 05' 07" N	108° 31' 37" W	dacite	Andesitic flows of Murtocks Hole	2
7	GR-95A	Canteen Canyon	33° 03' 17" N	108° 50' 12" W	rhyolite	Volcanic complex of Brock Canyon	2
8	GR-47	Copperas Pk.	33° 02' 33" N	108° 13' 27" W	andesite	"	2
9	GR-41A	Granny Mtn.	33° 02' 05" N	108° 16' 56" W	dacite	Volcanic complex of Alum Mountain	2
10	GR-35B	Copperas Pk.	33° 04' 30" N	108° 12' 27" W	dacite	"	2
11	GR-33A	Copperas Pk.	33° 05' 29" N	108° 07' 48" W	basalt	"	2
12	GR-21D	Copperas Pk.	33° 03' 01" N	108° 12' 44" W	rhyolite	"	2
13	GR-20H	Copperas Pk.	33° 03' 04" N	108° 13' 09" W	dacite	Andesitic flows of Gila Flat	2
14	GR-18F	Copperas Pk.	33° 04' 13" N	108° 13' 03" W	andesite	Volcanic complex of Alum Mountain	2
15	GR-14	Gila Hot Springs	33° 08' 38" N	108° 11' 32" W	andesite	"	2
16	GR-13B	Gila Hot Springs	33° 08' 12" N	108° 12' 41" W	andesite	"	2
17	GR-128C	L. Turkey Park	33° 07' 42" N	108° 15' 13" W	dacite	Andesitic lava flows of Gila Flat	2
18	GR-50C	Holt Mtn.	33° 19' 45" N	108° 48' 58" W	dacite	Cooney Tuff	2
19	MR-4A-76	Mogollon	33° 22' 85" N	108° 49' 29" W	rhyolite	"	1
20	GR-59A	Rice Ranch	33° 10' 55" N	108° 40' 06" W	dacite	"	2
21	GR-216A	Shelley Peak	33° 30' 25" N	108° 34' 32" W	rhyolite	Tuff of Fall Canyon	2
22	MR-116-74	Mogollon	33° 23' 35" N	108° 49' 38" W	rhyolite	"	3
23	GR-157H	Canyon Hill	33° 01' 17" N	108° 26' 20" W	rhyolite	Tuff of Davis Canyon	2
24	MR-31-74	Mogollon	33° 24' 12" N	108° 49' 15" W	rhyolite	"	2
25	GR-70B	Shelley Peak	33° 11' 06" N	108° 33' 57" W	rhyolite	Tuff of Shelley Peak	2
26	GR-142C	Canyon Hill	33° 04' 18" N	108° 24' 44" W	dacite	"	2
27	SP-5B-75	Saliz Pass	33° 35' 02" N	108° 55' 59" W	rhyolite	"	1
28	GR-70H	Shelley Peak	33° 10' 24" N	108° 35' 08" W	basalt	Andesitic flows of Gila Flat	2
29	GR-180J	Canteen Canyon	33° 05' 46" N	108° 32' 58" W	dacite	Andesitic flows of Murtocks Hole	2

Sample No.	Field No.	Quadrangle	Latitude	Longitude	Rock Name	Correlation	Analytical Method & Code
30	DG-155	Woodland Park	33° 22' 22" N	108° 18' 55" W	rhyolite	Bloodgood Canyon Rhyolite of Elston (1968)	2
31	DG-344	Lilley Mtn.	33° 16' 10" N	108° 22' 53" W	rhyolite	"	2
32	GR-88	Gila Hot Spring	33° 11' 41" N	108° 10' 47" W	rhyolite	"	2
33	GR-129A	Granny Mtns.	33° 06' 00" N	108° 18' 58" W	rhyolite	"	2
34	MR-41-74	Mogollon	33° 26' 36" N	108° 48' 22" W	rhyolite	"	3
35	DG-176	Grouse Mtn.	33° 22' 12" N	108° 43' 33" W	rhyolite	Tuff of Apache Springs Post Apache Springs	2
36	DG-347	Negrito Mtn.	33° 28' 30" N	108° 36' 14" W	dacite	Quartzite of Elston (1968)	2
37	DG-342	Lilley Mtn.	33° 16' 18" N	108° 25' 00" W	rhyolite	"	2
38	GR-99G	Grouse Mtn.	33° 18' 13" N	108° 40' 05" W	rhyolite	"	2
39	GR-107C	Grouse Mtn.	33° 16' 41" N	108° 38' 29" W	rhyolite	"	2
40	GR-71A	Shelley Peak	33° 12' 41" N	108° 37' 08" W	rhyolite	Rhyolite of Sacaton Mtn.	2
41	GR-84B	Shelley Peak	33° 14' 06" N	108° 36' 46" W	rhyolite	"	2
42	GR-62F	Grouse Mtn.	33° 15' 25" N	108° 43' 38" W	rhyolite	"	2
43	GR-101C	Grouse Mtn.	33° 21' 04" N	108° 44' 49" W	rhyolite	"	2
44	GR-34A	North Star Mesa	33° 06' 55" N	108° 00' 18" W	rhyolite	Rhyolite of Rocky Canyon	2
45	GR-74B	Canteen Canyon	33° 07' 18" N	108° 03' 07" W	rhyolite	Rhyolite of Diablo Range	2
46	GR-200B	Wall Lake	33° 21' 54" N	108° 06' 39" W	rhyolite	Rhyolite of Beaver Creek	2
47	GR-125C	L. Turkey Park	33° 09' 25" N	108° 21' 41" W	rhyolite	Rhyolite of Diablo Range	2
48	GR-59B	Rice Ranch	33° 11' 02" N	108° 40' 12" W	rhyolite	"	2
49	GR-56B	Holt Mtn.	33° 19' 31" N	108° 47' 52" W	rhyolite	Quartzite of Nabours Mountain (Rhodes, 1976)	2
50	DG-161	Canyon Ck. Mtns.	33° 25' 11" N	108° 17' 00" W	rhyolite	Rhyolite of Indian Creek	2
51	GR-113B	Grouse Mtn.	33° 21' 58" N	108° 44' 55" W	? dacite	Deadwood Gulch Member of Fanny Rhyolite (Rettig, 1978)	2
52	DG-175	Grouse Mtn.	33° 22' 20" N	108° 41' 09" W	rhyolite	Fanny Rhyolite	2
53	GR-54G	Holt Mtn.	33° 21' 29" N	108° 45' 37" W	rhyolite	"	2
54	GR-66B	Holt Mtn.	33° 15' 13" N	108° 47' 28" W	rhyolite	"	2
55	DG-10	Gila Hot Springs	33° 13' 40" N	108° 13' 46" W	Tholeiitic andesite	Post-ash flow tuff Andesitic flows	2
56	MR-7-74	Mogollon	33° 27' 44" N	108° 45' 12" W	andesite	"	2
57	MR-6-74	Mogollon	33° 28' 28" N	108° 45' 32" W	andesite	"	3
							3

SAMPLE NO.	FIELD NO.	Quadrangle	Latitude	Longitude	Rock Name	Correlation	Analytical methods code
58	MR-43A-74	Mogollon	33° 29' 07" N	108° 46' 34" W	andesite	Post-ash-flow tuff andesitic flows	3
59	MR-51A-74	Mogollon	33° 25' 37" N	108° 46' 19" W	andesite	"	3
60	MR-81B-75	Mogollon	33° 25' 13" N	108° 47' 14" W	andesite	"	1
61	MR-81A-75	Mogollon	33° 25' 08" N	108° 47' 23" W	andesite	"	1
62	OB-4E-76	O-Block Canyon	33° 33' 52" N	108° 50' 35" W	basalt	"	1
63	GR-54A	Holt Mtn.	33° 22' 04" N	108° 45' 29" W	andesite	Last Chance Andesite	2
64	GR-6B	Gila Hot Springs	33° 11' 40" N	108° 09' 20" W	andesite	Post-ash-flow tuff andesitic flows	2
65	GR-6A	Gila Hot Springs	33° 11' 40" N	108° 09' 20" W	basalt	"	2
66	DG-6	Gila Hot Springs	33° 11' 49" N	108° 11' 37" W	basalt	"	2
67	DG-5	Gila Hot Springs	33° 11' 48" N	108° 12' 12" W	basalt	"	2
68	DG-3	Gila Hot Springs	33° 11' 43" N	108° 10' 35" W	andesite	"	2
69	GR-110A	Grouse Mtn.	33° 19' 19" N	108° 40' 47" W	andesite	"	2
70	GR-176A	Holt Mtn.	33° 17' 35" N	108° 49' 17" W	Tholeiitic andesite	Dacitic intrusive rock of Holt Gulch	2
71	MR-48D-74	Mogollon	33° 29' 36" N	108° 45' 31" W	basalt	Hypabyssal andesitic intrusive rock (Ratté, 1977)	3
72	EP-3-75	Eagle Peak	33° 42' 18" N	108° 34' 30" W	andesite	Andesite of Eagle Peak	1
73	SP-50A-75	Saliz Pass	33° 06' 35" N	108° 56' 52" W	basalt	Basalt of Pueblo Park	1
74	OB-7E-76	O-Block Canyon	33° 32' 50" N	108° 52' 17" W	andesite	Post-ash-flow tuff andesitic flows	1
75	SP-42-75	Saliz Pass	33° 36' 18" N	108° 53' 30" W	basalt	Basalt of Saliz Hill	1
76	SP-27B-75	Saliz Pass	33° 34' 19" N	108° 53' 13" W	andesite	Hypabyssal andesite intrusive of Saliz Canyon	1
77	BR-10-74	Bearwallow Mtn.	33° 24' 55" N	108° 44' 45" W	basalt	Basalt of Coney Peak	3
78	BB-16E-76	Bull Basin	33° 42' 00" N	108° 53' 40" W	andesite	Hypabyssal intrusive of Legett Spring	1
79	GL-2-74	Glenwood	33° 21' 10" N	108° 54' 13" W	basalt	Basalt flow interlayered in Gila Conglomerate	3
80	GR-30C	North Star Mesa	33° 02' 39" N	108° 03' 55" W	olivine basalt	Basalt of North Star Mesa	2
81	GR-26D	North Star Mesa	33° 03' 22" N	108° 02' 38" W	olivine basalt	"	2



Table 3.--Description of analyzed rocks

Group I - Early andesitic rocks

Sample no.	Field no.	
1	SP-33-75	Gray, porphyritic lava flow in upper part of early andesitic complex. Contains microphenocrysts of olivine and pyroxene as well as plagioclase.
2	SP-7E-76	Gray, porphyritic, glassy monomictic vent(?) breccia with abundant plagioclase (andesine) and biotite phenocrysts, and amphibole and orthopyroxene microphenocrysts.
3	DM-3-76	Highly porphyritic, dull black glassy plug rock with abundant phenocrysts of plagioclase, pyroxene, and opaque oxide.
4	GR-88A	Blue-gray, porphyritic flow rock with about 15 percent 1-3 mm plagioclase and oxidized mafic phenocrysts.
5	GR-79C	Red, porphyritic flow rock with about 25 percent 1-4 mm plagioclase and minor biotite crystals from area of steep flow layering that may mark a plug.
6	GR-76A	Red, porphyritic flow rock with about 10-15 percent plagioclase and sparse green pyroxene phenocrysts.
7	GR-95A	Light-gray porphyritic flow rock with 1-3 mm altered feldspar phenocrysts in a pyritized and argillically altered matrix. Mafic phenocrysts, probably biotite, almost completely sericitized.
8	GR-47	Black rock with vitreous appearance has scattered pink feldspar crystals and glomeroporphyritic groups of plagioclase, pyroxene, and opaque oxide crystals in an aphanitic but holocrystalline matrix of flow-aligned plagioclase microlites.
9	GR-41A	Gray vitrophyre with about 20-25 percent 1-5 mm white plagioclase (andesine) and 1 mm biotite books in a colorless glass. Minor pyroxene, xenocrystic(?) quartz, opaque oxide, and zircon. Another sample has K-Ar ages of $29.6 \pm 1.0$ (sanidine) and $29.3 \pm 1.0$ (biotite) m.y.
10	GR-35B	Fine-grained, light-gray, granitic rock with some 1-4 mm plagioclase phenocrysts, sparse altered biotite, and disseminated pyrite. The potassic feldspar has a K-Ar age of $29.7 \pm 1.0$ m.y.

Table 3.--cont.

Sample no.	Field no.	
11	GR-33A	Fine-grained, dark-gray, vesicular flow rock with scattered olivine and pyroxene grains in a microlitic matrix of plagioclase (labradorite). Flows may correlate with andesitic flows that are younger than the volcanic complex of Alum Mountain.
12	GR-21D	Light-gray, fluidal, aphanitic rhyolite dike with sparse sodic plagioclase phenocrysts <1 mm long. Veinlets of quartz and magnetite commonly parallel the fluidal structure.
13	GR-20H	Light-gray, porphyritic flow rock with 10-15 percent 1-4 mm white plagioclase phenocrysts, partly sericitized biotite, greenish-yellow pyroxene, and rare quartz grains in an aphanitic matrix.
14	GR-18F	Dark-greenish-gray, porphyritic flow rock with abundant plagioclase (andesine) phenocrysts and biotite, altered pyroxene, and sparse amphibole microphenocrysts. Some granophyric quartz and feldspar in matrix.
15	GR-14	Dark-gray to black flow with with 4-5 mm plagioclase phenocrysts (andesine) in an aphanitic matrix of microphenocrysts and microlitic plagioclase. Microphenocrysts are biotite, pyroxene, and brown hornblende; rock also contains sparse rounded quartz xenocrysts.
16	GR-13B	Dark-gray, sparsely porphyritic rock from small stock or dike. Phenocrysts (10-15 percent) include plagioclase (oligoclase-andesine), biotite, and pyroxene. Rare quartz and sanidine xenocrysts(?).
17	GR-128C	Black vitrophyre with about 5 percent plagioclase (oligoclase-andesine), biotite, and pyroxene phenocrysts. Perlitic glass matrix is crowded with flow-aligned microlites.
Group II - Ash-flow tuffs and associated rocks		
18	GR-50C	Reddish-brown, densely welded tuff with about 40 percent small phenocrysts, mainly plagioclase (andesine), minor biotite and opaque oxide, and rare pyroxene, zircon, and brown hornblende; from upper part of Cooney Tuff sequence of ash flows.
19	MR-4A-76	Pink to red, densely welded phenocryst-poor ash-flow tuff with 5-10 percent sanidine, partly argillized sodic plagioclase, and rare quartz phenocrysts from lithophyssal zone in bottom of Whitewater Canyon.

Table 3.--cont.

Sample no.	Field no.	
20	GR-59A	Greenish-gray, propylitically altered, densely welded ash-flow tuff with 30-40 percent altered plagioclase, biotite, and pyroxene phenocrysts.
21	GR-216A	Light-lavender-gray, densely welded ash-flow tuff with 25-30 percent 1-3 mm quartz and sanidine and minor plagioclase phenocrysts; practically no mafic minerals other than rare oxidized biotite and opaque oxide.
22	MR-11B-74	Light-lavender-gray, moderately welded, ash-flow tuff with 20-30 percent quartz, plagioclase, and sanidine phenocrysts and accessory biotite, opaque oxide, zircon, and apatite in a devitrified vitroclastic matrix. Irregular, angular quartz phenocrysts as large as 1/2 cm.
23	GR-157H	Light brown, pumice-rich, moderately welded ash-flow tuff with less than 5 percent tiny quartz, plagioclase, and sanidine (moonstone) phenocrysts. Compacted dark-brown to creamy-white pumice fragments are as much as several centimeters long.
24	MR-31-74	Light gray to nearly white, partially welded, porous vaporphase ash-flow tuff with rare tiny quartz, sanidine, and plagioclase phenocrysts, and small gray eutaxitic pumice generally <1 cm long in a devitrified microcrystalline matrix.
25	GR-70B	Pinkish-brown, finely eutaxitic to fluidal, phenocryst-poor, densely welded ash-flow tuff near base of unit. Five to ten percent phenocrysts are mostly sanidine and plagioclase plus accessory biotite, brown hornblende, pyroxene, sphene, zircon, and apatite.
26	GR-142C	Pinkish-gray, densely welded tuff with abundant small white feldspar phenocrysts and scattered bronze biotite 1-2 mm in diameter, from near top of ash-flow tuff sheet.
27	SP-5B-75	Red, densely welded ash-flow tuff with 20-25 percent phenocrysts of aligned glassy plagioclase, and rare sanidine, scattered brown biotite, and sparse green pyroxene in a devitrified matrix. Sample is from 10-20 meters below top of ash-flow tuff sheet.
28	GR-70H	Light- to medium-gray, fine-grained flow rock with small plagioclase laths and dark-green pyroxene grains 1 mm or less in a fairly coarse microlitic matrix.

Table 3.--cont.

Sample no.	Field no.	
29	GR-180J	Reddish-brown, porphyritic flow rock with about 10 percent 1-4 mm plagioclase (andesine) laths and sparse pyroxene and opaque oxide microphenocrysts in a felted microlitic matrix. Mafics are highly oxidized.
30	DG-155	Light-gray, densely welded ash-flow tuff with abundant quartz and sanidine phenocrysts and accessory brown hornblende, opaque oxides, sphene, biotite, and zircon in a microeutaxitic to granophyric matrix.
31	DG-344	Light-gray, densely welded ash-flow tuff with abundant quartz and sanidine phenocrysts and minor accessory biotite, sphene, and opaque oxides in a devitrified matrix.
32	GR-8B	Light-gray, moderately to densely welded ash-flow tuff with 10-20 percent quartz and sanidine phenocrysts and eutaxitic pumice several cm long in a granophyric to cryptocrystalline matrix.
33	GR-129A	Light-gray, densely welded ash-flow tuff with eutaxitic white pumice several cm long and 10-15 percent 1-3 mm phenocrysts of sanidine and amoebic quartz grains, and smaller, yellow, euhedral sphene grains, opaque oxide, and rare brown hornblende and biotite in an aphanitic devitrified matrix.
34	MR-41-74	Pink, densely welded ash-flow tuff with 20-25 percent 1-3 mm quartz and sanidine phenocrysts in a devitrified eutaxitic matrix. Rare accessory minerals include sodic plagioclase, opaque oxide, sphene, biotite, and zircon.
35	DG-176	Pink, densely welded ash-flow tuff with abundant quartz and sanidine phenocrysts in a devitrified cryptocrystalline matrix.
36	DG-347	Reddish-brown, densely welded ash-flow tuff with 25-30 percent feldspar and scattered biotite phenocrysts. Quartz, sphene, and opaque oxides common in a devitrified matrix.
37	DG-342	Reddish-gray, densely welded ash-flow tuff with abundant feldspar, quartz, and biotite phenocrysts in an aphanitic, devitrified matrix.
38	GR-99G	Reddish-gray, densely welded ash-flow tuff with 55 percent 1-3 mm phenocrysts, which include white, somewhat skeletal sanidine, glassy plagioclase, quartz, rare black biotite and accessory opaque oxide, sphene, zircon, and apatite in a devitrified microeutaxitic, cryptocrystalline to granophyric matrix.

Table 3.--cont.

Sample no.	Field no.	
39	GR-107C	Reddish-brown, densely welded ash-flow tuff that contains abundant phenocrysts of sanidine with a sericitic sheen, and quartz in a devitrified cryptocrystalline matrix with granophyric eutaxitic pumice. Rock looks similar to tuff of Apache Spring, but rare plagioclase and foreign lithic fragments support its chemical similarity to Bloodgood Canyon Tuff, and thus sample is apparently from a block of Bloodgood Canyon Tuff enclosed in the younger tuff of Apache Spring.
40	GR-71A	Gray, porphyritic flow rock with 15-20 percent 1-3 mm quartz and feldspar phenocrysts in an aphanitic granophyric to micropegmatitic matrix. Rare biotite is largely altered to sericite.
41	GR-84B	Pink, porphyritic flow rock with 1-4 mm perthitically altered pink sanidine and quartz phenocrysts. Sparse plagioclase and accessory biotite, opaque oxide, apatite, sphene, and zircon are present in a micropegmatitic matrix.
42	GR-62F	Reddish-gray, porphyritic flow rock with 10-15 percent argillized feldspar and altered mafic phenocrysts in an aphanitic granophyric to micropegmatitic matrix.
43	GR-101C	Gray, porphyritic flow rock with about 30 percent quartz and feldspar phenocrysts in an aphanitic matrix that contains sparse opaque oxide, biotite, apatite, and zircon, and has a granophyric to micropegmatitic texture.
44	GR-34A	Light-gray, aphanitic rhyolite with <1 percent very sparsely scattered 1-2 mm quartz and sanidine phenocrysts and a rare flake of biotite. White lenticular lithophysae commonly have a medial lens of tridymite.
45	GR-74B	Light-gray, rhyolite porphyry with <5 percent inconspicuous quartz and sanidine phenocrysts in an aphanitic micropegmatitic to aplitic matrix containing numerous plumose spherulites.
46	GR-200B	White, friable, lithophysal rhyolite with 5-10 percent sanidine and amethystine quartz phenocrysts as much as 1/2 cm in diameter. Crystalline hematite or other metallic oxides project into some lithophysal cavities.
47	GR-125C	Light-gray to nearly white, strongly flow-banded porphyritic rhyolite with about 5 percent quartz and sanidine phenocrysts as much as 1/2 cm in diameter in an aphanitic granophyric to micropegmatitic matrix. Wisps of fine muscovite, rare oxidized biotite, and sphene in thin sections. Vugs 1-2 cm long are aligned with amethystine quartz.

- | no.   | no.       |   |
|---|-----------|---|
| 48  | GR-59B    | Light-pinkish-gray rhyolite porphyry dike with about 1-2 percent, 1-2 mm quartz phenocrysts and rare 1-3 mm pink sanidine phenocrysts, and a rare flake of biotite in an aphanitic granophyric to spherulitic matrix.   |
| 49  | GR-56B    | Light-pinkish-gray, porphyritic flow rock with about 20 percent 1-4 mm phenocrysts and microphenocrysts that are predominantly sodic plagioclase, and some quartz plus biotite, brown amphibole, pyroxene and opaque oxide, sphene, and zircon in a spherulitic to glassy matrix. |
| 50  | DG-161    | Light-gray rhyolite flow rock with about one or two percent scattered quartz phenocrysts in an aphanitic matrix.  |
| 51  | GR-113B   | Light-pink, pumiceous, poorly welded ash-flow tuff with about 20 percent quartz, feldspar, and biotite phenocrysts mostly <1 mm, and a few lithic lapilli in a devitrified cryptocrystalline matrix.  |
| 52  | DG-175    | Light-brown to white flowbanded rhyolite with about 1 percent quartz, sanidine, and plagioclase microphenocrysts in a spherulitic to cryptocrystalline matrix.  |
| 53  | GR-54G    | Light-lavender-gray, irregularly fluidal rhyolite with about 1 percent 1-2 mm pinkish-white plagioclase crystals in an aphanitic cryptocrystalline to granophyric matrix.   |
| 54  | GR-66B    | Massive, white, aphanitic rhyolite with less than 1 percent tiny quartz and feldspar crystals in a microgranular matrix much clouded with fine clay.  |
| Group III Post-ash-flow tuff andesites and associated rocks |           |   |
| 55  | DG-10     | Gray, amygdaloidal, aphanitic flow rock with rare yellow-green pyroxene crystals <1 mm long and sparse but conspicuous small quartz xenocrysts.   |
| 56  | MR-7-74   | Dark-gray, aphanitic flow rock with small white amygdules, and scattered brown, oxidized mafic phenocrysts generally <1 mm long. Rock is about 98 percent flow-aligned plagioclase microlites with scattered red-orange iddingsite after olivine.                                 |
| 57  | MR-6-74   | Gray-black, subvitreous, amygdaloidal flow rock with sparse tiny glassy-brown olivine grains in an aphanitic matrix. Microphenocrysts (<5 percent) of plagioclase, pyroxene, and olivine are in a flow-aligned microlitic plagioclase matrix.                                     |
| 58  | MR-43A-74 | Zebra-striped (flow-banded) purplish-gray rock with sparse tiny oxidized mafic crystals in a matrix of microlitic plagioclase.  |

Table 3.--cont.

Sample no.	Field no.	
59	MR-51A-74	Black, glassy flow rock with small white amygdules and sparse glomeroporphyritic clots. About 1 percent plagioclase (andesine-labradorite) and pyroxene microphenocrysts in a matrix of fine-grained plagioclase and mafic granules with an intersertal texture.
60	MR-81B-75	Gray, aphanitic, vesicular flow rock with a flow-aligned microlitic texture of andesine-labradorite microlites and opaque oxide and pyroxene granules. Sample is from flows above 20-meter purplish-gray sandstone lens within flow sequence.
61	MR-81A-75	Reddish-brown, aphanitic, vesicular flow rock with a dictytaxitic, microlitic matrix which contains abundant small subhedral opaque grains and local areas of ophitic-textured pyroxene and plagioclase.
62	OB-4E-76	Gray, massive, aphanitic flow rock with tiny (<1 mm) rare olivine phenocrysts. Flow-aligned microlitic plagioclase matrix contains sparse iddingsitic olivine and pyroxene microphenocrysts.
63	GR-54A	Black, subvitreous, massive flow rock with a few scattered 1-2 mm plagioclase phenocrysts. Sparse andesine and both clinopyroxene and orthopyroxene in a microlitic plagioclase and cryptocrystalline matrix. Rare but conspicuous quartz xenocrysts have reaction borders of pyroxene.
64	GR-6B	Gray, finely vesicular flow rock with andesine and clinopyroxene in dictytaxitic, microlitic texture with sparse scattered opaque oxide and pyroxene microphenocrysts.
65	GR-6A	Dark-gray, blotchy, massive flow rock with a diabasic texture of clinopyroxene, altered olivine, andesine, and opaque oxide.
66	DG-6	Brownish-gray, dictytaxitic to diabasic, amygdaloidal flow rock with common opaque oxide and minor plagioclase and clinopyroxene phenocrysts.
67	DG-5	Dark-gray, aphanitic flow rock with clinopyroxene and opaque oxide grains in a dictytaxitic microlitic plagioclase matrix.
68	DG-3	Brownish-gray, aphanitic flow rock of clinopyroxene grains and trachytoid plagioclase.
69	GR-110A	Gray, aphanitic flow rock with a few small vesicles and microphenocrysts of plagioclase, and rare pyroxene in a flow-aligned matrix of microlitic plagioclase.

Table 3.--cont.

Sample no.	Field no.	
70	GR-176A	Greenish-gray, fine-grained, intrusive rock consisting largely of altered subhedral plagioclase crystals with interstitial microlitic to micropegmatitic quartz and feldspar.
71	MR-48D-74	Gray, fine-grained, hypabyssal intrusive rock with small 1-4 mm plagioclase (andesine) and hornblende phenocrysts in a fine-grained matrix.
72	EP-3-75	Gray, fine-grained hypabyssal intrusive rock with abundant 1 mm and smaller plagioclase and pyroxene phenocrysts in a cryptocrystalline to glassy matrix. Orthopyroxene is more abundant than augitic pyroxene.
73	SP-50A-75	Mottled gray flow rock with labradorite phenocrysts several cm long locally, and augitic pyroxene phenocrysts as much as 1-2 cm long. Plagioclase, olivine, and pyroxene intergrown in a subophitic texture.
74	OB-7E-76	Dark-gray, fine-grained amygdaloidal flow rock. Felted microlitic labradorite forms polygonal to radial mosaic pattern with interstitial olivine, pyroxene, and opaque oxide grains.
75	SP-42-75	Black, vesicular, subvitreous flow rock with sparse small (1 mm) olivine and plagioclase phenocrysts in a very fine grained microlitic matrix of flow-aligned andesine and interstitial pyroxene.
76	SP-27B-75	Gray, fine-grained hypabyssal intrusive rock with abundant andesine, hornblende, and pyroxene phenocrysts 1-2 mm or smaller in a microgranular matrix.
77	BR-10-74	Dark-gray, aphanitic flow rock with about 10 percent small phenocrysts and microphenocrysts of andesine-labradorite, olivine, and opaque oxide in a microlitic plagioclase matrix.
78	BB-16E-76	Gray, porphyritic hypabyssal intrusive rock with about 15 percent phenocrysts including pink plagioclase and amethystine quartz as much as 1 cm long, and smaller mafic phenocrysts of biotite, greenish-yellow hornblende, sphene, and pyroxene. Some of quartz phenocrysts (or xenocrysts) have reaction rims with pyroxene; others show only corroded borders.
79	GL-2-74	Dark-gray, finely vesicular aphanitic flow rock with sparse green olivine crystals. Microscopically, rock has a diabasic texture of calcic plagioclase laths, olivine, and pyroxene.



Table 3.--cont.

Sample no.	Field no.	
80	GR-30C	Dark-gray to black, massive to vesicular, aphanitic flow rock with sparsely scattered olivine and pyroxene microphenocrysts in a poorly aligned microlitic plagioclase matrix.
81	GR-26D	Dark-gray, flow-banded, aphanitic flow rock with rare small (<1 cm) ultramafic inclusions. Less than 1 percent olivine phenocrysts in a well-aligned microlitic plagioclase matrix with interstitial olivine and pyroxene and opaque oxide grains.

EARLY ANDESITIC ROCKS

ASH-FLOW TUFTS AND ASSOCIATED ROCKS

POST ASH-FLOW TUFT ANDESITES AND ASSOCIATED ROCKS

$\text{Na}_2\text{O} + \text{K}_2\text{O}$  (weight percent)

40

50

60

70

80

90

100

110


120

130

ALKALINE FIELD

SUB-ALKALINE FIELD

Figure 1.--Alkali-Silica plot of some volcanic rocks of the Hogston-Dat II field. Divider between alkaline and subalkaline rocks from Irvine and Barnet, 1971.



$F = \text{FeO} + 0.8988\text{Fe}_2\text{O}_3$

Figure 2.--AFM plot of volcanic rocks from part of the Mogollon-Datil volcanic field. All values in weight percent

1. EARLY ANDESITIC ROCKS ..
2. ASH-FLOW TUFFS AND ASSOCIATED ROCKS ▲▲
3. POST ASH-FLOW TUFF ANDESITES AND ASSOCIATED ROCKS ++




Figure 3.--Classification diagram for subalkaline volcanic rocks,  
from Irvine and Baragar, 1971.

