DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

OPEN-FILE REPORT
80-2084

SIERRA ANCHA WILDERNESS, ARIZONA

CORRELATION OF MAP UNITS

QTu		QUATERNARY AND TERTIARY
Cs		CAMBRIAN
Ydb		PROTEROZOIC Y
Yt		
Ym		
Yds		
Yu	Yp	Ypr
	Yr	

LIST OF MAP UNITS

QTu	Alluvial and colluvial deposits, undivided (Quaternary and Tertiary)
Cs	Sandstone (Cambrian)
Ydb	Diabase (Proterozoic Y)
Yt	Troy Quartzite (Proterozoic Y)
Ym	Mescal Limestone (Proterozoic Y)
Yds	Dripping Spring Quartzite (Proterozoic Y)
Yp	Pioneer Formation (Proterozoic Y)
Yr	Ruin Granite (Proterozoic Y)
Ypr	Pioneer Formation and Ruin Granite, undivided (Proterozoic Y)
Yu	Dripping Spring Quartzite, Pioneer Formation, and Ruin Granite, undivided (Proterozoic Y)

CONTACT

FAULT--Dotted where concealed, dashed where inferred

X CASTLE PK — PROMINENT TOPOGRAPHIC FEATURE

LOCATION OF KNOWN MINERALIZATION--From Granger and Raup (1969a, 1969b), Light (1980), and Shride (1969)

- A — Asbestos
- F — Fluorine
- C — Copper
- Fe — Iron
- U — Uranium

APPROXIMATE BOUNDARY OF SIERRA ANCHA WILDERNESS AREA

APPROXIMATE BOUNDARY OF SALOME STUDY AREA

0.4 — RADIOMETRIC CONTOUR--Contour interval 0.4 and 0.8. Hachures indicate closed area of lower values. Values expressed in arbitrary units

U.S. GEOLOGICAL SURVEY DENVER
DEC 8 1980
LIBRARY
GEOLOGIC DIVISION

PLEASE RETURN TO
IN BACK OF BOUND VOLUME

Base from U.S. Geological Survey Mc Fadden Peak and Rockinstraw Mtn, 1949, 1:62,500; Armer Mountain, Copper Mountain, Greenback Creek, Picture Mtn, Theodore Roosevelt Dam, and Windy Hill, 1964, 1:24,000

SCALE 1:48 000

1 ½ 0 1 2 3 MILES

1 .5 0 1 2 3 KILOMETERS

CONTOUR INTERVAL FEET
DATUM IS MEAN SEA LEVEL

Geology modified from Bergquist and others (1980) and from A. F. Shride (unpub. mapping, 1967)

PLATE 5. RADIOMETRIC MAP OF RATIO OF APPARENT SURFACE DISTRIBUTION OF eU/eTh