Base from U.S. Geological Survey 1:250,000 Mt. Fairweather, 1961; Sitka, 1951 OPEN FILE REPORT 81-0027-H

STUDIES RELATED TO WILDERNESS

The Wilderness Act (Public Law 88-577, September 3, 1964) and related acts require the U.S. Geological Survey and the U.S. Bureau of Mines to survey certain areas on Federal lands to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a geochemical survey of the Western Chichagof-Yakobi Islands Wilderness Study Area in the Tongass National Forest, Alaska. About 65 percent of the study area was established as a wilderness on December 2, 1980, under the Alaska National Interest Lands Conservation Act (P.L. 96-487).

CORRELATION OF MAP UNITS

Tf Tm

Kd

Ks

Kkb

KJf KJm

Tag

MzPzu

Teg

Boundary of study area

Geochemical sample site

Lead (AA) = 30-50 ppm

Lead (AA) > 50 ppm

QUATERNARY

TERTIARY (?)

CRETACEOUS (?)

CRETACEOUS

TRIASSIC(?)

LIST OF MAP UNITS

Alluvial deposits, undivided

Diorite sill, extensively altered

Undivided metasedimentary, metavolcanic,

and metaplutonic rocks

SYMBOLS

Contact, approximately located, dotted where concealed

Felsic plutonic rocks

Mafic plutonic rocks

Felsic plutonic rocks

Mafic plutonic rocks
Whitestripe Marhle

Goon Dip Greenstone

Sitka Graywacke

Kelp Bay Group

CRETACEOUS AND JURASSIC

MESOZOIC AND PALEOZOIC

Forest, Alaska. About 65 percent of the study area was established as a wilderness on December 2, 1980, under the Alaska National Interest Lands Conservation Act (P.L. 96-487).

In the course of the U.S. Geological Survey investigations of the Western Chichagof-Yakobi Islands Wilderness Study Area, 2,230 bedrock geochemical samples were collected. Samples were analyzed for 31 elements by a 6-step, semiquantitative spectrographic method (Grimes and Marranzino, 1968) and for 4 elements by atomic absorption spectrophotometry (Ward and others, 1969). Complete analytical data, station coordinates, and a station location map are available in two reports: Johnson, 1982, and Johnson and Elliott, 1984. A map and discussion of the mineral

resource potential of the study area is also available (Johnson, Kimball, and Still, 1982). Background levels for each element vary for different lithologies in the study area. Because of this and variability introduced from other sources such as sampling technique, analytical variance, and chemical weathering, it is impossible to select a specific analytical level above which values indicate mineralization. Higher values may indicate a greater likelihood of bedrock mineralization, but confidence levels are low for single element high values and results which are not supported by neighboring values. This map shows the distibution of high analytical values for the element lead by two analytical techniques, as well as the locations of all 2,230 samples. Multiple symbols for a single analytical technique at one sample site represent

REFERENCES CITED

multiple samples at that site.

Grimes, D. J., and Marranzino, A. P., 1968, Directcurrent arc and alternating-current spark emission spectrographic field methods for the semiquantitative analysis of geologic materials: U.S. Geological Survey Circular 591,

Johnson, B. R., 1982, Magnetic tape containing trace element data for bedrock geochemical samples from the West Chichagof-Yakobi Islands Wilderness Study Area, southeastern Alaska: National Technical Information Service Report No. USGS-GD-82-005, computer tape, 1 reel.


Johnson, B. R., and Elliott, G. S., 1984, Map showing bedrock geochemical station locations, Western Chichagof-Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Open-File Report 81-0027-A, scale 1:125,000.

Johnson, B. R., and Karl, S. M., 1982, Reconnaissance geologic map of the Western Chichagof and Yakobi Islands Wilderness Study Area, southeastern Alaska: U.S. Geological Survey Miscellaneous


Field Studies Map MF-1476-A, scale 1:125,000.

Johnson, B. R., Kimball, A. L., and Still, Jan, 1982,
Mineral resource potential map of the Western
Chichagof and Yakobi Islands Wilderness Study
Area, southeastern Alaska: U.S. Geological
Survey Miscellaneous Field Studies Map MF-1476-B,
scale 1:125,000.

Ward, F. N., Nakagawa, H. M., Harms, T. F., and Van Sickle, G. H., 1969, Atomic absorption methods of analysis useful in geochemical exploration: U.S. Geological Survey Bulletin 1289, 45 p.



Index Map Showing Location of Study Area



2 0 2 4 6 8 KILOMETERS

MAP SHOWING THE DISTRIBUTION AND ABUNDANCE OF LEAD IN BEDROCK SAMPLES, WESTERN CHICHAGOF AND YAKOBI ISLANDS WILDERNESS STUDY AREA, SOUTHEASTERN ALASKA

Geology generalized from Johnson and Karl (1982)