

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

MINERAL RESOURCES OF THE PHIPPS-DEATH HOLLOW
INSTANT STUDY AREA, GARFIELD COUNTY, UTAH

By

G. W. Weir, U. S. Geological Survey

and

M. E. Lane, U.S. Bureau of Mines

Open-File Report 81-558

1981

This report is preliminary and has not been
edited or reviewed for conformity with U.S.
Geological Survey standards.

Mineral Surveys

Wilderness Studies Related to Bureau of Land Management

The Federal Land Policy and Management Act (Public Law 94-579, October 21, 1976), requires the U. S. Geological Survey and the U.S. Bureau of Mines to conduct mineral surveys on certain areas to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a mineral survey of the Phipps-Death Hollow Instant Study Area, Utah.

Contents

	Page
Summary.....	1
Introduction.....	2
Geology and geochemistry.....	4
Mining districts and mineralized areas.....	10
Oil and gas exploration.....	11
Mineral resource potential.....	11
References.....	15

Illustrations

Figure 1. Index map of Utah showing location of Phipps-Death Hollow Instant Study Area.....	3
2. Mineral resource potential and generalized geologic map of the Phipps-Death Hollow Instant Study Area showing locations of manganese prospect and dry wells.....	9

Tables

Table 1. Generalized stratigraphic section of the Phipps-Death Hollow Instant Study Area.....	5
2. Record of wildcat wells drilled in the Phipps-Death Hollow Instant Study Area.....	12

Mineral resource potential of the Phipps-Death Hollow
Instant Study Area, Utah

SUMMARY

A geologic and geochemical investigation and an examination of the only existing prospect have been conducted to determine the mineral resource potential of the Phipps-Death Hollow Instant Study Area, Garfield County, Utah. The study area encompasses about 80 sq mi (200 sq km) of mesas and canyon between the towns of Escalante and Boulder. Mesozoic and Paleozoic strata known from subsurface tests are overlain by outcropping rocks of Triassic and Jurassic age. The Navajo Sandstone forms the most extensive outcrops. The Boulder-Collett Canyon anticline is the major fold, but rocks dip gently south or southeast in most of the area. The mineral and energy resource potential of the Phipps-Death Hollow Instant Study Area is low.

The Van Hamet manganese prospect, in the southwestern corner of the area, consists of manganese minerals in concretions in sandstone. The deposit is too small to yield ore in commercial quantities. Triassic rocks underlying the study area may be locally uranium and copper bearing, but no evidence suggests that they contain economic deposits. Spectrographic analyses of stream sediments and rocks in and near the study area do not suggest derivation from mineralized terranes.

The Boulder-Collett Canyon anticline was tested for oil and gas near the northeast edge of the area. The well was dry, as were six other tests on this anticline outside the study area. Although one well 6 mi (10 km) to the south had a show of oil in Permian strata, the oil and gas potential of the area appears low.

INTRODUCTION

During 1979 and 1980 the U. S. Geological Survey and the U. S. Bureau of Mines conducted field studies to evaluate the mineral-resource potential of the Phipps-Death Hollow Instant Study Area, Garfield County, Utah. Field studies included geological mapping (Weir and Beard, 1981), geochemical sampling, and a search for mines, prospects and mineralized areas.

The Phipps-Death Hollow Instant Study Area includes about 80 sq mi (200 sq km) of mesas and canyons between the towns of Escalante and Boulder in central Garfield County, Utah (fig. 1). The area is bounded approximately on the south and east by a paved highway, on the west by an improved road leading to points north, from Escalante and on the north by the Dixie National Forest. Access to most of the area is difficult even by foot.

Except on the outskirts of the town of Escalante no permanent residents are in or near the area. Escalante had a population of 638, and the total population of Garfield County was 3,157 in 1970.

The Phipps-Death Hollow Instant Study Area lies within the western part of the Canyon Lands section of the Colorado Plateau physiographic province (Thornbury, 1965, p. 426-434). Steep-walled canyons, mesas, and plateaus cut into Paleozoic and Mesozoic sedimentary rocks are the major landforms in this section. The dominant structures are gently dipping homoclines associated with broad upwarps and basins. The study area lies southwest of the Circle Cliffs upwarp and northeast of the Kaiparowits basin. It is south of the basalt-capped, flat-lying rocks of the Aquarius Plateau.

Exploration within the Phipps-Death Hollow Instant Study Area has been limited to drilling for oil and gas and prospecting for manganese. In 1979 and 1980 no wells were drilled for oil and gas and the only known manganese prospect was dormant.




Figure 1.--Index showing location of the Phipps-Death Hollow Instant Study Area, Utah

GEOLOGY AND GEOCHEMISTRY

Rocks of Triassic and Jurassic age, totalling about 2,000 ft (600 m) in thickness, and thin deposits of Quaternary age crop out in the study area (Weir and Beard, 1981). Underlying Mesozoic and Paleozoic rocks known from subsurface tests in the region are more than 4,000 ft (1,200 m) thick. Characteristics of the rock units are listed in table 1. Grayish-orange, crossbedded sandstone of the Navajo Sandstone forms the most extensive outcrops (fig. 2). Younger units are exposed along the northern, western, and southern fringes of the area. The base of the Navajo and the upper part of the underlying Kayenta Formation are exposed only in canyons near the east edge of the study area.

In most of the area the rocks dip gently to the south or southeast (Weir and Beard, 1981). The major folds are the Boulder-Collett Canyon anticline, whose southwestern flank is the steeply dipping Escalante monocline. The rocks are complexly jointed but are not displaced by faults.

Geochemical investigation of the study area included sampling stream sediments (38 samples), bedrock (8 samples), and water from streams (3 samples). The stream-sediment and rock samples were analyzed for 30 elements using semiquantitative spectrographic methods and the water samples were analyzed for uranium using atomic-absorption and liquid-chromatograph techniques.

The analyses of the stream-sediment and rock samples do not suggest derivation from mineralized terranes, but rather appear characteristic of the country rock, mostly Jurassic sandstone and shale (Weir and Beard, 1981). A few analytical values are high relative to the whole set of analyses but probably are related to contamination of fine debris derived from volcanic rocks that crop out north of the study area (McFall and Peterson, 1971). None

Table 1.—Generalized stratigraphic section of the Phipps-Death Hollow Instant Study Area, Utah

[Data on units below Navajo Sandstone from Zeller (1973) and Peterson (1973) based on wells in and near the Upper Valley oilfield about 12 mi (20 km) west of study area]

System	Series	Formation	Member	Thickness		Description
				Feet	(Meters)	
Quaternary	Holocene and Pleistocene	Alluvial, eolian, and colluvial deposits		0-40	(0-12)	Clay, silt, sand, and gravel along streams; silt and sand on mesas; pediment gravels consisting chiefly of pebbles to boulders of basalt. Colluvium consists chiefly of slope-wash aprons and lobate masses of basalt gravel.

Unconformity

Middle		Entrada Sandstone		150+	(60+)	Reddish-brown silty sandstone. Only lower part of formation present in area; as much as 1,000 ft (300 m) thick in adjoining Dave Canyon quadrangle (Zeller, 1973).
Middle		Camel Formation	Upper member	400-600	(120-180)	Midstone and minor fine-grained sandstone, chiefly reddish-brown; light-gray gypsum in irregular lenses; thin bedded yellowish-gray limestone near base.
Jurassic	Middle	Page Sandstone	Thousand Pockets Tongue	10-60	(3-18)	Sandstone, chiefly yellowish-gray, fine-grained; crossbedded, locally contorted. Contains layer of reddish-brown mudstone near middle of unit.
Middle		Camel Formation	Judd Hollow Tongue	10-40	(3-12)	Midstone, reddish-brown; sandstone, limonite-stained, fine-grained, 2 to 10 ft (0.6-3 m) thick at base.

Table 1.—Generalized stratigraphic section of the Phipps-Death Hollow Instant Study Area, Utah—Continued

System	Series	Formation	Member	Thickness		Description
				Feet	(Meters)	
	Middle	Page Sandstone	Harris Wash Tongue	0-40	(0-12)	Sandstone, grayish-orange, crossbedded; contains chert pebbles at base.
Unconformity						
Jurassic and Triassic(?)		Navajo Sandstone		1,300-1,500	(400-455)	Sandstone, grayish-orange, crossbedded in part contorted; contains sparse thin lenses of limestone and reddish-brown siltstone, mostly near base.
Triassic(?)	Upper(?)	Kayenta Formation		200-320	(60-100)	Sandstone interbedded with siltstone, reddish brown. Base not exposed; thickness of exposed rock about 180 ft (55 m).
Triassic	Upper	Wingate Sandstone		200-260	(60-80)	Sandstone, light-reddish orange to reddish-brown, fine-grained, cross-bedded.
Unconformity						
Triassic	Upper	Chinle Formation	Church Rock Member Owl Rock Member Petrified Forest Member Monitor Butte Member	0-25 150-250 150-350 100-200	(0-8) (45-75) (45-105) (30-60)	Sandstone, brown, fine-to-medium grained. Sandstone and mudstone, red, brown, and greenish-gray; limestone, greenish-gray, in thin lenses. Mudstone, bentonitic, variegated, and sandstone; contains silicified wood. Mudstone, bentonitic, green and grayish-red; sandstone, light-brown, micaceous, ripple-laminated.


Table 1.—Generalized stratigraphic section of the Phipps-Death Hollow Instant Study Area, Utah—Continued

System	Series	Formation	Member	Thickness		Description
				Feet	(Meters)	
			Shinarump Member	0-60	(0-18)	Sandstone, conglomeratic, yellowish-gray, crossbedded; mudstone, greenish-gray, in thin lenses. In same general stratigraphic position as mottled unit.
			or			
			Mottled siltstone unit	0-50	(0-15)	Siltstone and minor sandstone, mottled red and white.
Unconformity						
	Lower and Middle(?)	Moenkopi Formation		800-880	(245-270)	Sandstone, shale, and siltstone, reddish-brown; minor limestone.
			Timpowap Member	60-70	(18-21)	Dolomite, light-brown to light-yellowish-red.
Unconformity						
Permian	Lower	Torowap Formation		360-420	(110-130)	Dolomite, light-brown dense; interbedded light-gray sandstone and anhydrite.
	Lower					
	Lower	Coconino Sandstone		50	(15)	Sandstone, light-gray.
		Outler Formation	Organ Rock Member	140-160	(45-50)	Siltstone, light-gray to light-red, some brown sandstone
	Lower	Outler Formation	Cedar Mesa Sandstone Member	1,350-1,400	(410-425)	Sandstone, light-gray to light-yellowish-red, fine- to medium-grained; lower one-third contains interbedded light-brown to light-yellowish-red limestone and dolomite.

Table 1.—Generalized stratigraphic section of the Phipps-Death Hollow Instant Study Area, Utah—Continued

System	Series	Formation	Member	Thickness		Description
				Feet	(Meters)	
Unconformity						
Pennsylvanian		Hemosa Formation		340	(105)	Sandstone, light-gray; interbedded light-yellowish-red dolomite and reddish- and purplish gray siltstone.
Pennsylvanian		Molas Formation		40-70	(12-21)	Shale, reddish-gray, some limestone concretions and sandstone.
Unconformity						
Mississippian	Upper and Lower	Redwall Limestone		900	(270)	Limestone and dolomite, light-gray to light-yellowish-red; extensive karst surface at top; in part cavernous.
Devonian	Upper	Ouray Limestone		160	(50)	Limestone, light-yellowish-red, dense.
Devonian		Unnamed		230	(70)	Dolomite, light-yellowish-red, some interbedded sandstone and greenish-gray shale.
Unconformity						
Cambrian		Unnamed		100	(30)	Dolomite, light-yellowish-red and interbedded greenish- and reddish-gray shale.

¹Subsurface terminology.



EXPLANATION

Je	Entrada Sandstone (Middle Jurassic)	1. Gulf Oil Co. No. 1 Garfield - X well
Jcp	Carmel Formation and Page Sandstone (Middle Jurassic)	2. Van Hamet manganese prospect
Jrn	Navajo Sandstone (Jurassic & Triassic ?)	3. Mountain Fuel Supply Co. No. 1 - Collett Well
Rk	Kayenta Formation (Triassic ?)	
—	Contact	x Prospect
		⊙ Oil well, dry hole

Figure 2. Mineral resource potential and generalized geologic map of the Phipps-Death Hollow Instant Study Area showing locations of manganese prospect and dry wells.

of the anomalies in the sediments or rocks are extremely high nor do they form a pattern suggesting a need for additional sampling. None of the water samples showed a uranium content of more than 1.1 parts per billion.

MINING DISTRICTS AND MINERALIZED AREAS

The Phipps-Death Hollow Instant Study Area does not lie within an organized mining district. Mining claim records were examined at the office of the Garfield County Recorder in Utah. No mining claims are known to have been located within the study area.

Development and exploration of mineral resources in and near the Phipps-Death Hollow Instant Study Area have been limited to the prospecting for manganese and the quarrying of road material.

The Van Hamet prospect in the southwestern corner of the area is on a small manganese deposit less than 200 ft (60 m) in diameter. Purplish-black manganese minerals are in irregular nodules, from a fraction of an inch to 6 in. (15 cm) across, scattered through layers less than a ft (30 cm) thick in a lensing sandstone about 6 ft (2 m) thick. The host rock is reddish-brown, fine-grained sandstone in the Judd Hollow Tongue, the basal unit of the Carmel Formation of Middle Jurassic age. Samples of the mineralized material collected by Doelling (1975, p. 138) ranged from 16 to 27 percent manganese and from 45 to 54 percent silica. Such mineralized material, however, is estimated to make up less than 5 percent of the host sandstone.

Road material has been quarried in and near the study area from Quaternary colluvial deposits, consisting mostly of pebbles to boulders of basalt, and from siltstone, shale, and minor sandstone of the lower part of the Carmel Formation of Middle Jurassic age. None of this quarried material has been trucked more than a few miles.

OIL AND GAS EXPLORATION

Two exploratory wells have been drilled in the area for oil and gas in Triassic and Permian rocks. The Gulf Oil Co. No. 1 Garfield-X well tested the Escalante anticline. The Mountain Fuel Supply Co. No. 1-Collett well tested the Boulder-Collett Canyon anticline. Both wells were dry and have been plugged and abandoned. Summary data for these wells are given on table 2.

MINERAL RESOURCE POTENTIAL

The mineral resource potential of the Phipps-Death Hollow Instant Study Area is very low. A single metallic mineral deposit is present (fig. 2). The spectrographic analyses of 46 samples of stream sediments and rocks in and near the study area do not suggest derivation from mineralized terranes. Test wells for oil and gas were dry.

The only known mineral deposit is at the Van Hamet manganese prospect in the southwestern corner of the area. The deposit consists of purplish-black manganese minerals in irregular nodular concretions, as much as 6 in. (15 cm) across, in reddish-brown, fine-grained sandstone in the Judd Hollow Tongue of the Carmel Formation. The mineralized material is scattered through layers less than a foot (30 cm) thick in a lensing sandstone about 6 ft (2 m) thick and makes up less than 5 percent of the host sandstone. It lies within an area less than 200 ft (60 m) in diameter. Samples of the mineralized rock collected by Doelling (1975, p. 138) ranged from about 16 to 27 percent manganese and from 45 to 54 percent silica. The deposit is too small to yield ore in commercial quantities. Furthermore, geologically similar deposits elsewhere in western Utah have not proved economic (Crittenden, 1951, p. 14). Thus, the manganese potential of the area is negligible.

Uranium-copper deposits in Triassic rocks are exposed in the Circle Cliffs area about 10 to 15 mi (16-24 km) east of the study area. The Circle

Table 2.—Record of wildcat wells drilled in the Phipps-Death Hollow Instant Study Area, Utah

[Data from records of the Conservation Division of the U. S. Geological Survey]

Operator	Well	Location	Year completed	Total depth (ft)	Formation at surface	Oldest formation penetrated	Remarks
Gulf Oil Co.	No. 1 Carfield-X	Sec 10, T. 35 S., R. 3 E.	1972	4,399	Navajo Ss. (Jurassic and Triassic?)	Cedar Mesa Ss. Mbr. of Outler Fm. (Permian)	Dry hole.
Mountain Fuel	No. 1 Collett	Sec 14, T. 34 S., R. 2 E.	1969	3,225	Pediment deposits (Quaternary) and Navajo Ss. (Jurassic and Triassic?)	White Rim Ss. Mbr. of Outler Fm. (Permian)	Dry hole.

Cliffs deposits are relatively small and weakly mineralized (Davidson, 1965, p. 65-91; Doelling, 1975, p. 107-109, 131-135). The Triassic rocks underlying the Phipps-Death Hollow area may also be locally uranium- and copper-bearing, but no evidence suggests that they contain economic deposits.

The oil and gas potential of the area appears low, although oil is produced from a fold similar to the Escalante anticline in the Upper Valley field about 12 mi (20 km) southwest of the study area. The productive strata are in the Timpoweap Member of the Moenkopi Formation (Triassic) and the Kaibab Limestone (Permian). Shows of oil have also been noted in the upper part of the Moenkopi, in the Cedar Mesa Sandstone Member of the Cutler Formation (Permian) and in the Redwall Limestone of Mississippian age (Peterson, 1973; Doelling, 1975, p. 91-96). The Escalante anticline was tested in the area by the Gulf Oil Co. No. 1 Garfield-X well in 1972. According to records of the Conservation Division of the U. S. Geological Survey, the well penetrated the Moenkopi and Kaibab and bottomed in the Cedar Mesa Sandstone Member of the Cutler Formation (Permian) at a depth of 4,399 ft (1,340 m) with no recorded shows of oil or gas. Three other tests of the Moenkopi and Kaibab drilled on the Escalante anticline, 8 to 10 mi (13-16 km) north of the area, were also dry wells. The Boulder-Collett Canyon anticline was tested in 1969 by the Mountain Fuel Supply Co. No. 1-Collett well near the northeast edge of the area. The well, which was dry, bottomed below the Kaibab in the White Rim Member of the Cutler Formation (Permian) at a depth of 3,225 ft (973 m). Six other tests on this anticline outside the study area, 6 to 12 mi. (10-20 km) south of the Mountain Fuel Supply Co. well, were also unsuccessful though one well 6 mi (10 km) to the south had a show of oil in Permian strata.

Potential construction materials within the area include gravel in

Quaternary alluvial and colluvial deposits, and gypsum and limestone in the Carmel Formation. These materials are, however, not economically significant, because ample supplies are readily available at nearby localities outside the study area.

References

- Crittenden, M.D., 1951, Manganese deposits of western Utah: U. S. Geological Survey Bulletin 979-A, 62 p.
- Davidson, E. S., 1965, Geology of the Circle Cliffs area, Garfield and Kane Counties, Utah: U. S. Geological Survey Bulletin 1229, 140 p.
- Doelling, H. H., 1975, Geology and mineral resources of Garfield County, Utah: Utah Geological Mineral Survey Bulletin 107, 175 p.
- McFall, C. C., and Peterson, P. R., 1971, Geology of the Escalante-Boulder area, Garfield County, Utah: Utah Geological and Mineralogical Survey Map 31, scale 1:62,500.
- Peterson, P. R., 1973, Upper Valley Field: Utah Geological and Mineralogical Survey Oil and Gas Field Studies 7.
- Thornbury, E. F., 1965, Regional geomorphology of the United States: New York, John Wiley, 609 p.
- Weir, G. W., and Beard, L. S., (1981), Geologic map of the Phipps-Death Hollow Instant Study Area, Garfield County, Utah: U. S. Geological Survey Miscellaneous Field Studies Map MF-__A, scale 1:48,000.
- Zeller, H. D., 1973, Geologic map and coal resources of the Dave Canyon quadrangle, Garfield County, Utah: U. S. Geological Survey Coal Investigations Map C-59, scale 1:24,000.