

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

PRELIMINARY GEOLOGIC MAP OF THE GRAND GULCH BENCH QUADRANGLE,  
MOHAVE COUNTY, ARIZONA

By

I. Lucchitta and S. Beard

Open-File Report 81-1321

1981

This report is preliminary and has not  
been edited or reviewed for conformity  
with U.S. Geological Survey standards  
and nomenclature.

## GEOLOGIC SETTING

This map is part of a regional study of the Grand Wash Fault. Geologic mapping was carried out by Lucchitta in 1973, and by Lucchitta and Beard in 1981.

The Grand Wash fault forms the boundary between the Colorado Plateau and the Basin and Range Province to the west. The fault is a high-angle normal fault that trends approximately north and is down to the west. Along most of its trace, the fault is buried by the Tertiary Muddy Creek Formation and other upper Cenozoic deposits. In this quadrangle, however, the fault is composed of a complex system of subparallel breaks, many of which are excellently exposed along the Grand Wash Cliffs. Typically, these breaks are of small displacement and up to the west. Where the fault system is widest and best developed, the up-to-the-west faults are associated with westerly tilts of 15-30° in the Paleozoic beds. The faulting and tilting combine to maintain structural elevation as one goes west. Substantial net down-to-the-west displacement is attained only at the main breaks. The first of these is composed of en-echelon segments located at or near the base of the Grand Wash Cliffs, and typically drops the uppermost Paleozoic units (Kaibab, Toroweap and Hermit Formations, and possibly part of the Esplanade Sandstone) against Mississippian rocks. One or more additional breaks are located west of the Grand Wash Cliffs and are buried by basin fill and other upper Cenozoic deposits. Most of the displacement that formed the basin occurred along these buried faults.

The map area is in a region where stratigraphic units change and thicken rapidly westward from the platform sequence of the Grand Canyon to the shelf sequence of the eastern Great Basin. The changes in stratigraphy, which are matched by changes in nomenclature, are not yet studied adequately. To avoid potential conflicts and to facilitate structural mapping, which is the main purpose of this study, we have simplified parts of the stratigraphic column into map units with distinctive lithologic and outcrop characteristics.


In the classic Grand Canyon section, a thick sequence of redbeds occupies the interval between the Redwall Limestone and the Coconino Sandstone. The redbeds are subdivided into the Supai Group of McKee (1975) and the overlying Hermit Shale. Both are composed of hematitic sandstone, siltstone and shale in various proportions. To the west, the Supai Group becomes increasingly calcareous. At the Grand Wash Cliffs, in rocks equivalent to the Supai Group of McKee, 1975 are composed of well bedded limestone with interbedded reddish cross-bedded sandstone and red shale. Only the upper part of the uppermost unit of the Supai, the Esplanade Sandstone remains carbonate-free McNair (1951) referred to the underlying calcareous units as the Pennsylvanian Callville Limestone and the Permian Pakoon Limestone. Although disconformities separate many of these units, the main break in lithology and weathering characteristics occurs between the cliff-forming limestones and the overlying slope-forming redbeds of the upper Esplanade and Hermit. We therefore have mapped the former as the Permian and Pennsylvanian Pakoon and Callville undivided, and the latter as the Permian Hermit and Esplanade Formations undivided. The Callville as mapped thus includes rocks equivalent to all the Supai Group of the Grand Canyon except the upper part of the Esplanade Sandstone. A more extensive discussion of the nomenclature and correlations of these rocks is given by G. H. Billingsley (1978).

The Devonian as mapped the Grand Gulch Bench quadrangle includes the interval between the unclassified dolomites of McKee and Resser (1945), also called undifferentiated dolomites by Billingsley (1978), and the Redwall Limestone. These rocks are equivalent to the Middle Devonian Temple Butte Formation of the Grand Canyon and the Muddy Peak Formation of the Lake Mead area.

The Muddy Creek Formation is a term applied by Longwell (1936) and Lucchitta (1966) to Miocene interior-basin deposits filling the Grand Wash trough, even though these rocks are not physically continuous with beds in the type locality (Muddy Valley of Nevada) which were called "Muddy Creek beds" by Stock (1921).

Several inactive copper prospects and mines occur within the quadrangle. In the surface workings, mineralization occurs in cross bedded sandstone beds of the Callville Formation (*sensu lato*). Structural control of mineralization appears likely. The presence of basaltic dikes in the vicinity also suggests the possibility of hydrothermal mineralization.

#### CORRELATION OF MAP UNITS



## DESCRIPTION OF MAP UNITS

[Areas marked with queried unit symbols have not been field checked]

- Qa1 ALLUVIUM (HOLOCENE)--Silt, sand, gravel, cobbles and boulders in active washes. Consists of angular to subangular, poorly sorted, unconsolidated material of local derivation. Thickness unknown; probably less than 2 m. Grades into, and locally includes pediment gravels
- PIEDMONT-SLOPE DEPOSITS (HOLOCENE AND PLEISTOCENE?)
- Qpf Pediment and fan gravels (Holocene)--Silt, sand gravel, cobbles and boulders on active pediments and fans. Consists of angular, poorly sorted, unconsolidated material of local derivation. Thickness unknown; probably less than a few meters. Pediment and fan gravels grade laterally into each other. Both grade laterally into, and locally include recent alluvium
- Qopf Older pediment and fan gravels (Pleistocene?)--Silt, sand, gravel, cobbles and boulders on gently sloping surfaces that are at various elevations above modern washes, by which they are being dissected. Consists of angular, poorly sorted, unconsolidated material of local derivation. Thickness generally less than 10 m.
- Tb BASALTIC FLOWS (PLEISTOCENE? OR MIOCENE?)--Fresh to moderately weathered, fine- to medium-grained, vesicular to dense, olivine basalt. Groundmass locally diktytaxitic. Groundmass plagioclase and olivine visible in hand specimen. Fresh to moderately altered olivine present as 0.5 to 3 mm phenocrysts that form less than 1 percent of rock
- Tmc MUDDY CREEK FORMATION (MIOCENE)--Claystone, siltstone, sandstone, pebble to boulder fanglomerate, freshwater limestone, dolomite and gypsum, deposited under conditions of interior drainage in the Grand Wash basin, which was formed by basin-range faulting. The various facies grade into each other both laterally and vertically. In quadrangle, unit consists chiefly of well- to poorly bedded, moderately consolidated sandstone and fanglomerate containing subangular to subrounded pebbles to boulders of Paleozoic rocks derived from the east. The fanglomerate locally fills steep and narrow canyons and cuts into the Grand Wash Cliffs. Unit is at least 600 m, and probably several thousand meters thick
- uPz PERMIAN ROCKS--Includes all or part of the Kaibab Limestone, Toroweap Formation, Coconino Sandstone, Hermit Shale, and Esplanade Sandstone. Present as intricately broken downfaulted wedges

- Ph s HERMIT SHALE, AND UPPER PART OF ESPLANADE SANDSTONE OF SUPAI GROUP UNDIVIDED (PERMIAN)--Redbed sequence that includes the Hermit Shale and that part of the Esplanade Sandstone Member of Supai Formation of White (1929) that overlies the highest carbonate beds. The Hermit comprises moderately to poorly indurated deep-red siltstone and light-red thin-bedded sandstone. The upper part of the Esplanade includes moderately indurated, brick-red to white, fine-grained, cross stratified, cliff-forming sandstone as well as poorly indurated deep-red gypsiferous shale. Map unit thickness 450-500 m
- PPc PAKOON LIMESTONE OF MCNAIR, 1951, (PERMIAN), AND CALLVILLE LIMESTONE (PENNSYLVANIAN) UNDIVIDED UNDIFFERENTIATED--Predominantly light-gray, fossiliferous, locally cherty, well-bedded limestone that forms ledges and ledgy slopes, with interbeds of purplish-red cliff-forming cross bedded sandstone and red-slope forming shale. Thickness about 300 m
- Mr REDWALL LIMESTONE (MISSISSIPPIAN)--light-gray, aphanitic to crystalline, fossiliferous, mostly thick bedded limestone, cherty in lower part. Forms massive, conspicuous cliff. Thickness 180-200 m
- D DEVONIAN ROCKS--Medium- to dark-gray, medium-grained, medium- to thick-bedded, typically fetid dolomite, limestone, and dolomitic sandstone. Upper part forms alternating ledge and slope; lower part forms color banded cliff. Thickness 140-200 m


#### REFERENCES

- Billingsley, G. H., 1978, A synopsis of stratigraphy in the western Grand Canyon: Museum of Northern Arizona Research Paper 16, 27 p.
- Longwell, C. R., 1936 Geology of the Boulder Reservoir flow, Arizona-Nevada: Geological Society of America Bulletin, v. 47, p. 1393-1476.
- Lucchitta, Ivo, 1936, Cenozoic geology of the Upper Lake Mead area adjacent to the Grand Wash Cliffs, Arizona: Unpublished Ph. D thesis, The Pennsylvania State University, 218 p.
- McKee, E. D., 1975, The Supai Group--Subdivision and nomenclature: U.S. Geological Survey Bulletin 1395-J, 11 p.
- McKee, E. D., and Resser, C. E., 1945, Cambrian history of the Grand Canyon region: Carnegie Institute of Washington Publication 563, pt. II, p. 171-232.
- McNair, A. H., 1951, Paleozoic stratigraphy of northwestern Arizona: American Association of Petroleum Geologists Bulletin v. 35, no. 3, March.
- Stock, Chester, 1921, Later Cenozoic mammalian remains from the Meadow Valley region, southeastern Nevada: American Journal of Science, 5th Series, v. 2, p. 250-264.

White, D., 1929, Flora of the Hermit Shale, Grand Canyon, Arizona: Carnegie Institute of Washington Publication 405, 219 p.

EXPLANATION OF MAP SYMBOLS

———— CONTACT--Dashed where approximate

 32° 60' ---? FAULT--Showing dip, direction and plunge of slickensides, or rack of slickensides. Bar and ball on downthrown side. Dashed where approximately located or probable, queried where doubtful, dotted where concealed.

STRIKE AND DIP OF BEDS

 30  


Inclined  
Horizontal