

Map and tables showing preliminary results of K-Ar age studies in the Ugashik quadrangle, Alaska Peninsula
by
Frederic H. Wilson

Reported here are 15 new potassium-argon age determinations on igneous and altered igneous rocks from the Ugashik quadrangle. These determinations are part of an ongoing petrographic and K-Ar geochronologic study of igneous rocks and mineralized areas in the Ugashik and Karluk quadrangles. The new age determinations reported here provide support data for preliminary conclusions in Wilson and Shew (1981). They are also in close agreement with results from studies in the Chignik and Sutvik Island quadrangles to the south (Wilson and others, 1981; Wilson, 1980). The dates reported here include three ages determined on rocks from the Meahik Formation (Detterman and others, 1981; Knappen, 1929). These determinations are in the same range as ages from the Meahik Formation in the Chignik and Sutvik Island quadrangles to the south (Wilson and others, 1981).

Also reported here are four age determinations on three Pleistocene igneous intrusions. Two of these intrusions are from magnetic centers that may have been active during Holocene time. The age of 1.78 m.y. from a sample that may be related to Yantarni Volcano contrasts with an age of 0.62 m.y. reported previously (Wilson and others, 1981) and suggests an eruptive history spanning at least 1.1 m.y. The two Yantarni(?) samples are very similar petrographically. The dated sample 2 suggests a mixing process was important in an unusual rock, in which thin section study (Table 2) discards results on plagioclase and impure hornblende separates. The isopropy (< 10%) in the hornblende is plagioclase and no explanation for the discordance can be advanced at this time. Blue Mountain was indicated to be Jurassic in age by Betkman (1980), however the dates reported here indicate that the volcanic center is Pleistocene. Nork (1965) had reported Blue Mountain to be a Tertiary intrusion.

Three samples from the Rex Prospect yielded early Oligocene ages on both mineralized and unmineralized rocks. Samples R-41358A and R-41360 were collected, described, and analyzed by M.L. Silberman. Reconnaissance mapping of the Rex prospect by Silberman, D.P. Cox, and this author indicates a multi-event history and further dating work is in progress in an attempt to clearly define this history.

One sample from the Mike prospect, a enphysematous system, yielded concordant ages on biotite and plagioclase of 3.48 and 3.42 m.y., respectively. These ages are interpreted to be hydrothermal alteration ages.

A concordant Late Jurassic age on a granite cobble collected from conglomerates in the Upper Jurassic Naknek Formation is reported here. This sample was analyzed to place a maximum age on the Naknek Formation at this locality.

Potassium was determined by flame photometry using a lithium molybdate fusion technique (Engels and Igenells, 1970). Potassium analyzers were Byron Lal, B. Vivit, and Paul Klack. Argon extraction and measurement was accomplished using standard technique described by Dalrymple and Lanphere (1965), essentially as described by Dalrymple and Lanphere (1965). The analytical error precision using the method of Cox and Dalrymple (1967) together with the standard deviation of analytical estimates of uncertainties in the concentration of ³⁹K tracer and potassium measurements. Sample preparation, argon extraction and data reduction was by the author with assistance from Nora Shew, Rita Taylor, Brian Ho, and Leda Beth Gray, except for the two samples (R-41358A and R-41360) by M.L. Silberman.

Analytical data is listed in Table 1, rock sample descriptions in Table 2 and sample locations are plotted on the map.

References

Betkman, H.M., 1980, Geologic map of Alaska: U.S. Geol. Survey, scale 1:2,500,000, 2 sheets.

Burk, C.A., 1985, Geology of the Alaska Peninsula - island arc and continental margin: Geol. Society of America Memoir 99, 250p., 2 maps, scales 1:250,000 and 1:500,000.

Cox, Allan, and Dalrymple, G.B., 1967, Statistical analysis of geochronologic reversal data and the resolution of potassium-argon dating: Journal of Geophysical Research, v. 72, p. 2603-2614.

Dalrymple, G.B., and Lanphere, P.M., 1969, Potassium-argon dating: principles, techniques, and applications in geochronology: W.H. Freeman and Co., San Francisco, California, 258p.

Detterman, R.C., Miller, T.H., Young, M.L., and Wilson, F.H., 1981, Geologic map of the Chignik and Sutvik Island quadrangles, Alaska: U.S. Geol. Survey Misc. Geologic Investigations Map 1-129, scale 1:250,000.

Engels, J.C., and Igenells, C.O., 1970, Effect of sample inhomogeneity in K-Ar dating: Geochimica et Cosmochimica Acta, v. 34, p. 1007-1017.

Knappen, B.S., 1929, Geology and mineral resources of the Aniakchak National Monument, in Smith, P.S., Mineral resources of Alaska, report on progress of investigations in 1924: U.S. Geol. Survey Bull. 797, p. 161-227.

Wilson, F.H., 1980, Late Mesozoic and Cenozoic tectonics and the age of porphyry copper prospects: Chignik and Sutvik Island quadrangles, Alaska Peninsula: U.S. Geol. Survey Open-File Report 80-343, 94p., 5 plates, scales 1:250,000 and 1:2,500,000.

Wilson, F.H., and Shew, Nora, 1981, Preliminary results of potassium-argon age determinations from the Ugashik quadrangle, Alaska Peninsula: in Gordon, W.L., ed., The U.S. Geological Survey in Alaska, Accomplishments during 1980: U.S. Geol. Survey Circular 843, in press.

Wilson, F.H., Dunn, K.C., and Herson, P.L., 1981, Map and tables showing geochronology and whole-rock geochemistry of the Chignik and Sutvik Island quadrangles, Alaska: U.S. Geological Survey Misc. Field Studies Map MF-103-M, 3 sheets, scale 1:250,000.

Table 1. Potassium-argon age determinations from the Ugashik quadrangle.

Sample No. and name	Location Latitude Longitude (quadrangle)	Rock type	Mineral or component dated*	$\pm K_2O$	$^{40}Ar_{rad}$	$\pm K_2O$	$^{40}Ar_{rad}$	Age and $\pm 1\sigma$ (m.y.)**
				wt %	mmol/gm	wt %	mmol/gm	
79A02 Meahik fm.	57°08.2'N 157°30.2'W Ugashik A-5	Andesite	HR	1.564	6.2488	86.0	27.85 ± .25	27.85 ± .25
				1.567	1.547	6.2912	79.0	28.03 ± .45
				1.534				$\bar{x} = 27.96 \pm .36$
79A04a Yantarni Volcano Ugashik A-5	57°02.2'N 157°26.2'W	Andesite or basalt	HR	1.04	0.2647	7.2	1.77 ± .024	1.77 ± .024
				1.03	1.04	2.876	9.3	1.79 ± .019
				1.04				$\bar{x} = 1.78 \pm .034$
79A0617 Kialagvik Volcano Ugashik A-3	57°15.6'N 156°49.4'W	Andesite	Plag	0.429	0.0589	3.3	0.80 ± .109	0.80 ± .109
				0.430	0.429	0.551	7.6	.89 ± .029
				0.429				$\bar{x} = .90 \pm .113$
79A0618a Clast in Naknek fm. Ugashik B-3	57°23.0'N 156°54.8'W	Granite	Hbl	.860	.862	20.35	78.6	161 ± .65
				.843		19.54	71.3	155 ± .21
				.843				$\bar{x} = 158 \pm 4.6$
79A0623 Rex prospect Ugashik A-3	57°12.1'N 156°36.0'W	Altered andesite porphyry	Hbl	8.73	202.3	84.2	354 ± .59	354 ± .59
				8.72	8.72	198.4	90.1	352 ± .41
				8.72				$\bar{x} = 353 \pm 2.2$
79A0623 Rex prospect Ugashik A-3	57°12.1'N 156°36.0'W	Altered andesite porphyry	Hbl	.502	2.475	44.1	33.9 ± .17	33.9 ± .17
				.503	.503	2.563	36.1	35.0 ± .16
				.504				$\bar{x} = 34.5 \pm .27$
				.504				
79A0635 Meahik fm. Ugashik A-3	57°15.9'N 156°54.8'W	Andesite porphyry	Hbl	.732	3.670	48.6	34.5 ± .29	34.5 ± .29
				.738	.732	3.811	44.1	35.4 ± .29
				.725				$\bar{x} = 35.2 \pm 1.01$
				.732				
79A0620 Meahik fm. Ugashik A-6	57°05.9'N 157°51.2'W	Basalt	HR	.563	2.007	52.5	26.6 ± .15	26.6 ± .15
				.565	.563	2.060	46.0	25.2 ± .13
				.561				$\bar{x} = 24.9 \pm .49$
801358A Rex prospect Ugashik A-4	57°14.2'N 157°03.0'W	Quartz diorite	Hbl	1.056	5.298	52.	34.8 ± .65	34.8 ± .65
				1.050	1.048	5.267	57.	35.2 ± .45
				1.037				$\bar{x} = 35.0 \pm 0.27$
				8.67	8.70	39.67	77.	31.4 ± 0.9
				8.67				
801360 Rex prospect Ugashik A-5	57°14.2'N 157°03.0'W	Altered diorite porphyry	Rio (secondary)	7.20	7.21	38.84	68.	37.0 ± 1.1
				7.22				
79AY90 Blue Mountain Ugashik C-3	57°51.5'N 156°50.0'W	Andesite	Plag	.418	.1014	9.9	1.68 ± .048	1.68 ± .048
				.421	.420	.0992	10.2	1.66 ± .048
				.420				$\bar{x} = 1.66 \pm .072$
				.420				
			Hbl with plag	.95	.0736	2.3	.53 ± .021	.53 ± .021
				.97	.0797	2.4	.57 ± .020	.57 ± .020
				.96				$\bar{x} = .95 \pm .042$
				.98				
79AY101 Mike prospect Ugashik A-3	57°03.8'N 157°13.3'W	Dacite	Plag	1.008	.474	12.7	3.25 ± .09	3.25 ± .09
				1.016	1.014	.526	15.0	3.59 ± .09
				1.008				$\bar{x} = 3.42 \pm .28$
				1.002				
			Hbl	8.82	4.815	11.7	3.79 ± .08	3.79 ± .08
				8.79	8.83	4.024	14.3	3.16 ± .03
				8.67				$\bar{x} = 3.48 \pm .45$

*HR = whole rock, Plag = plagioclase and plagioclase are HF treated, see Wilson (1980), Hbl = hornblende, Rio = biotite.

** $k_1 = 5.72 \times 10^{-11} \text{yr}^{-1}$, $k_2 = 8.78 \times 10^{-11} \text{yr}^{-1}$, $k_3 = 4.963 \times 10^{-10} \text{yr}^{-1}$, $k_4 = 1.167 \times 10^{-4} \text{mol/mol}$.




Table 2. Rock Descriptions

79AWs 2, Meahik Formation. Gray weathering, very fine-grained basalt or andesite flow. Contains sparsely distributed phenocrysts of ortho- and clinopyroxene and rare plagioclase (anorthite An 50) in a flow-textured groundmass of very fine-grained plagioclase, amphibole or augite, and glass.

79AWs 6a, Yantarni Volcano. Very fine-grained porphyritic andesite or basalt sill or dike. Clinopyroxene phenocrysts of augite and clinopyroxene are more transparent than ortho-; plagioclase, opaque minerals, and devitrified glass. Some phenocrysts of augite are altered to calcite and urtite. Weak development of flow texture indicated by plagioclase laths in groundmass.

79AWs 17, Kialagvik Volcano. Dark-green porphyritic two-pyroxene andesite with abundant phenocrysts of plagioclase, hornblende, and augite in a very fine-grained to aphanitic groundmass of plagioclase, glass, and opaque minerals. Parts of groundmass are more transparent than others; these more transparent parts containing fewer opaque minerals are where the augite is concentrated. Plagioclase phenocrysts are of two types, some of which are poikilitic and have overgrown edges. Other plagioclase phenocrysts are unaltered, free of inclusions, and may have obscured edges. Most plagioclase phenocrysts of both types have compositions between An 60 to An 70 though some may have compositions nearer to An 45.

79AWs 24a, Naknek Formation. Clast of granite from conglomerate in Naknek Formation. Medium to coarse-grained hornblende-biotite granite. Estimated mode is: 30% plagioclase (An 47), 30% potassium feldspar, 15% quartz, 15% biotite and chlorite, and 10% hornblende. Hornblende typically has prograde cores, biotite grains may be chloritized. Hypoclinopyroxene granular texture.

79AD1 95, Rex prospect. Porphyritically altered hornblende andesite porphyry. Closely packed phenocrysts of hornblende and plagioclase (An 54) in a groundmass of devitrified glass, plagioclase and sericite. Most plagioclase phenocrysts are partially altered to epidote, many hornblende phenocrysts are partially altered to chlorite and calcite. Abundant epidote.

79AD1 23, Meahik Formation. Ductile- to irregularly-shaped hornblende andesite porphyry. Phenocrysts of green pleochroic hornblende in a groundmass of fine-grained anhedral plagioclase crystals (An 66?) and devitrified glass. A small percentage (< 5%) quartz crystals, some of which are subhedral or euhedral.

79ACe 20, Meahik Formation. Porphyritic flow-textured olivine basalt. Phenocrysts of olivine and unaltered pyroxene(?) in a fine-grained groundmass of plagioclase (anorthite An 62), opaque minerals, olivine and possibly augite. Olivine phenocrysts are altered to idiomorphic boundaries and some plagioclase crystals have cores that appear to be chlorite or urtite. Fresh plagioclase, poikilitic hornblende, and quartz in a sparse groundmass of quartz, plagioclase, and minor potassium feldspar. Minor biotite is present and appears to be primary or formed in reaction relationship with hornblende; it is not hydrothermal. Hornblende has inclusions of plagioclase and quartz, and shows minor replacement by chlorite. Pyrite is present and is concentrated around and within hornblende and biotite grains.

R-41358A, Rex prospect. Biotite-hornblende quartz diorite. Abundant phenocrysts of relatively fresh plagioclase, poikilitic hornblende, and quartz in a sparse groundmass of quartz, plagioclase, and minor potassium feldspar. Minor biotite is present and appears to be primary or formed in reaction relationship with hornblende; it is not hydrothermal. Hornblende has inclusions of plagioclase and quartz, and shows minor replacement by chlorite. Pyrite is present and is concentrated around and within hornblende and biotite grains.

R-41360, Rex prospect. Hydrothermally altered diorite porphyry. Alterated, saussuritized plagioclase phenocrysts in a fine-grained, saccharine-textured groundmass of quartz, plagioclase, and minor potassium feldspar(?). All mafic phenocrysts are altered to hydrothermal biotite; biotite also occurs in veinlets. Sulfides concentrated around biotite.

79AYb 101, Mike prospect. Porphyritic hornblende-plagioclase andesite. Hornblende phenocrysts are poikilitic, pleochroic in brown and green, and generally have sharp, euhedral grain boundaries, with no evidence of resorption. Plagioclase phenocrysts are unaltered; though boundaries are rounded and have lines of inclusions in a thin zone parallel to the grain boundaries suggesting resorption and recrystallization. The plagioclase grains include weakly zoned and strongly oscillatory zoned grains which have an average composition of An 54 or greater. The very fine-grained groundmass is primarily glass and devitrified glass with a small proportion of microcrystalline plagioclase and hornblende(?).

79AWs 6a, Blue Mountain. Porphyritic hornblende-biotite dacite porphyry. Coarse phenocrysts of plagioclase (An 46), biotite, quartz, and altered hornblende in a fine-grained groundmass of feldspar, secondary biotite, and quartz. Pseudomorphs after hornblende of fine-grained biotite and chlorite. Plagioclase phenocrysts are generally unaltered though grain boundaries show resorption into the groundmass. The plagioclase grain boundaries are and may be partially saussuritized. Quartz phenocrysts have rounded outlines and grain boundaries show resorption. Biotite phenocrysts show minor alteration to chlorite; chlorite is more abundant in pseudomorphs after hornblende.

Map and tables showing preliminary results of K-Ar age studies in the Ugashik quadrangle, Alaska Peninsula

by
Frederic H. Wilson
1982

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature.