



DESCRIPTION OF MAP UNITS		
LANDFORMS	GEOMORPHOLOGY	SYMBOLS
EOLIAN SAND DEPOSITS	ACTIVE DUNES	Contact between geomorphic/landform units, plotted from aerial photographs
	YOUNGER STABILIZED DUNES	Drainage divide
	OLDER STABILIZED DUNES	Stream channel
	RELATIVELY FLAT LOWLANDS PARTLY SAND COVERED	Dune crests
	UNDULATING UPLANDS AND LOWLANDS, PARTLY SAND COVERED WITH LONGITUDINAL DUNES	Area with linear arrangement of blowouts
FLUVIAL DEPOSITS	MODERN RIVER BARS	Stabilized longitudinal dunes
	YOUNGEST POINT BAR SYSTEM	Thaw lake, including coalescing lakes
	YOUNGER POINT BAR SYSTEM	In floodplain channel scrotts
	OLDER POINT BAR SYSTEM	Drained thaw lake basins
	VALLEYS	Abandoned channel
UNDIFFERENTIATED SURFICIAL DEPOSITS	UNDULATING LOWLANDS	Point bar system with arrow pointing
	SLOPES OF TRANSPORT	In direction of build up
	UNDULATING UPLANDS	Polygonal ground cracks
	FOOTSLOPES	Swamp
	BEDROCK	Terrace scarp, hachures point downslope
		Inferred fault scarp, teeth on downthrow side
		Village with roads
		Elevation in meters above sea level



**INTRODUCTION**

The Noghahabara Sand Dunes and part of the Koyukuk Lowland are located in the Kotzeb River (1250,000) quadrangle between the northeastern spurs of the Hulo Hills and the Koyukuk River near the village of Huslia. The Noghahabara Sand Dunes lie approximately 45 km west from the village of Huslia in the northern and northwest corners of the Kotzeb River C-3 and C-4 (1:250,000) quadrangles. The dunes lie within an area of about 300 km<sup>2</sup> of dunes in the lower Koyukuk Valley (Péwé, 1975, p. 36), in the zone of discontinuous permafrost and the area has a mean annual temperature of approximately -4°C. Geologic maps of the area, scale 1:250,000, (Coss, 1957; Patton, 1964) show three eolian units and two flood-plain units. Some of the surficial geologic units in the map of the Yukon-Koyukuk Lowland (Weber and Péwé, 1970) are useful guides for photo-interpretation in our study area, even though it is 10 km to the north.

This map (scale 1:60,000) is based solely on the interpretation of thirteen false-color infrared and seven black and white aerial photographs (A). Four major map units, bedrock, undifferentiated surficial deposits, eolian and fluvial, form the basis for our map. Both eolian and fluvial landforms have been listed in order of relative age. Dunes and sand sheets are recognized, on the photographs, by: 1) color differences, 2) topography and geomorphic form, 3) presence or absence of drainage patterns, 4) distribution of flow lines, and 5) relative topographic position. Fluvial landforms, point bars and abandoned channels, are separated on the basis of: 1) color differences, 2) meander development, 3) degree of modification by thermokarst activity, 4) presence or absence of an-ice lakes or abandoned channels, and 5) cross-cutting relationships. Undifferentiated surficial deposits are separated on the basis of: 1) color differences, 2) degree of modification by thermokarst activity, and 3) geomorphologic criteria like relative height and slope angles. Landforms in which the deposits are undifferentiated were mapped and described without consideration of age. Bedrock was recognized on the photographs by: 1) color differences and 2) topographic expression.

**EOLIAN SAND DEPOSITS**

Based on topographic expression five eolian sand units have been recognized: active dunes (Q<sub>5</sub>), younger stabilized dunes (Q<sub>4</sub>), older stabilized dunes (Q<sub>3</sub>), relatively flat lowlands partly covered with sand (Q<sub>2</sub>), and undulating uplands and lowlands partly covered with sand and longitudinal dunes (Q<sub>1</sub>). The largest active dune field forms a circular dune of sand having a diameter of approximately 7 km. The edge of the field is sharply defined all around by a distinct escarpment. Lying on an elevation of 100 to 170 meters above sea level the active dune fields cover an area of 44.5 km<sup>2</sup>. This unit (Q<sub>5</sub>) consists of actively drifting sand, free of vegetation, which appears white to light-blue on the photographs. Dunes with relatively low height and even spacing are clearly visible on the photographs and appear to be transverse or bar-shaped in form. The low dune height is probably influenced by shallow subsurface permafrost as seen from the many areas of deflation located in the central part of the dune field. Deflation areas, red in color on the photographs, have a linear arrangement of W-E to NW-SE. This direction slightly deviates from the inferred transport direction of the dunes. We have delineated the dune height, trough to crest, to be generally less than 10 meters with local spacing being 50 to 200 meters. Dune forms are surprisingly uniform in spacing, height and crest line orientation. Most dune crests can be followed for a few kilometers. Crest lines are parallel in orientation NNE-SSW. There is a slight deviation of the orientation along the edge of the dune field. Evidence of present-day surficial drainage and flow lines is absent in the active dune field. However, former drainage lines are still visible. Coss (1957) described the dune field as consisting of quartz sand, coarse grained on the western edge decreasing in grain size towards the eastern edge. Results of a mechanical analysis of a sand sample collected from the Noghahabara Sand Dunes was reported by Weber and Péwé (1970). The midpoint on the cumulative frequency curve lies at 2.15 phi, fine sand. Mechanical analysis of sand samples from another active dune field (The

Great Kobuk Sand Dunes) show a slightly coarser sand with the midpoint on the cumulative frequency curve ranging from 1.75 phi, medium sand, to 3.0 phi, fine sand (Galloway and Koster, n.d.). We do not agree with Patton (1964) and Coss (1957) that the active dunes are moving northward and eastward or southward. The major transport direction appears to be to the west and northwest with a possible secondary (secondary) direction to the east.

Younger stabilized dunes (Q<sub>4</sub>) lie to the west of the active dune field. This unit appears white to green-grey on the photographs, suggesting that stabilization is still incomplete. Dune crests are clearly visible and are approximately parallel to the crest lines in the active dune field. Dunes have climbed up the eastern lower slopes of the Hulo Hills. As in the Noghahabara Sand Dunes no flow lines are present.

A third eolian sand unit (Q<sub>3</sub>) consists of dunes stabilized by sparse vegetation (color reddish-grey). Based on topographic position these dunes appear to be older than the dunes previously described. These rose-like dunes lie just east and south of the largest active dune field. They form circular to elliptical bodies and on the photographs appear medium to dark grey in color. Trunks of the dunes, transverse in form, vary strongly in orientation. Development of the dunes may have occurred under periglacial conditions. Downwind direction before stabilization was variable originating from the southeast, east and northeast. The presence of several flow lines and lack of a white line on the photographs suggest the dunes have been stable for some time. Similar rose-like dunes have been identified along the Koyukuk River south of Noghahabara, and in various other Alaskan stabilized dune fields: Snow Creek Sand Dunes, Tetlin Sand Dunes, and Karibuk Sand Dunes. Although the dune fields (Q<sub>4</sub>, Q<sub>3</sub>, Q<sub>2</sub>) cover an area of more than 80 km<sup>2</sup>.

Large areas, having a relief of less than 15 m, surrounding the dune fields and in the area south of the Dagliit River have a partial cover of eolian sand (Q<sub>2</sub>). These areas are dark green to reddish green in color on the photographs. The eolian sand appears to be relatively thin and discontinuous. Formation of this sand cover might be related to an early phase of dune-building, however in this unit no dune forms have been recognized. These areas are conspicuous for their absence of flow lines.

**UNDIFFERENTIATED SURFICIAL DEPOSITS**

The largest eolian map unit (Q<sub>1</sub>) occupies the Koyukuk Flats just west and east of the flood-plain. Patton (1964) called this area modified eolian sand deposits. "Dunes much modified and locally dissected by stream erosion but some longitudinal and parabolic forms recognizable in aerial photographs. Deposits have strong NE-SW trend. However accumulations on northeast-facing slopes and orientation of parabolic dunes convex toward the southwest suggests deposition by prevailing northerly winds." Apart from an incipient, probably discontinuous eolian sand cover, longitudinal dunes are readily recognizable on the photographs. The longitudinal dunes have been indicated on the map only schematically and have a strong NE-SW trend indicating former transport directions to the southwest. This unit probably rests on the older Dagliit terrace deposits as described for the Yukon-Koyukuk Lowland by Weber and Péwé (1970). The large number of flow lines, dissected flow lines and the terrace scarp of unit Q<sub>1</sub> cutting into the dune forms suggest a long period of stabilization. These dunes lie on an elevation of approximately 80 meters and vary in area from 0.1 to 3.5 km<sup>2</sup>. Dune forms are strongly modified by flow lines and dissected by stream erosion.

Although the dunes have been stabilized by vegetation massive subsidence of a sand sheet to the east has occurred due to recent tectonic activity. "Associated with the Huslia earthquake of April 1, 1958, were several thousand sand flows up to one-quarter square mile in area. This material appears to have flowed upwards from old lake and stream beds currently covered by vegetated sand dunes. Local collapse of the sand dune layer accompanied the sand flowage" (Davis, 1958).

**FLUVIAL DEPOSITS**

As noted by Weber and Péwé (1970) and Péwé (1973) the Koyukuk River flood-plain is a classic example of a flood-plain developed by strongly meandering rivers under periglacial conditions. Our four fluvial units are basically equivalent to the units developed by Weber and Péwé (1970) and have been differentiated and arranged according to relative age. Point bars and abandoned channels are located in a large valley where the river is locally capable of building point bars by channel migration. Near Huslia present-day lateral erosion of the river bank still occurs.

Modern river bars, Unit Q<sub>4</sub>, appear bluish-white in color on the photos and are free of vegetation. The youngest point bar system (Q<sub>3</sub>) appears red on false color infrared photographs suggesting the presence of a dense and healthy vegetation. Point bars appear fresh showing little or no evidence of erosion by thermokarst activity. Drainage is not integrated and flow lines are absent.

Apparently bluish-grey to red in color the youngest point bar system (Q<sub>2</sub>) consists of modified point bars and abandoned channels. This unit represents several generations of point bars suggesting a complex history. Just below Huslia the width of the meander belt is approximately 10 km with a radius of 2.5 km. Although drainage is not integrated, linear flow lines parallel to drainage trends are common in abandoned channels.

The older point bar system (Q<sub>1</sub>) consists of oxbow lakes, flow lines with no preferred orientation, abandoned channels and remnants of point bars. Thermal erosion around the edges of flow lines, integration of drainage between flow lines, and the degree of modification of the point bars suggest this unit was abandoned a relatively long time ago. The absence of red color on the photographs indicates trends such as willows, larch and birch are absent. Colors to purple colors suggest the development of a typical tundra mat consisting of peat moss and sedges (Weber and Péwé, 1970). This unit cuts into Q<sub>2</sub>, producing a well defined terrace scarp of 10 to 30 meters in height. The total width of the Koyukuk River flood-plain in this area is from 13 to 22 km.

**UNDIFFERENTIATED SURFICIAL DEPOSITS**

As previously mentioned, all other landforms, developed in deposits of unknown origin were mapped without consideration for their relative age. Unit Q<sub>1</sub>, (red) is found along tributaries and the Dagliit and Hulo Hills rivers which are strongly meandering streams having a gradient of less than 1:200. Mainly reworked alluvium is present along these stream valleys. Presently the Hulo Hills and Dagliit Rivers and their tributaries are incapable of building point bars. Polygonal ground cracks thought to be an expression of ice-wedges in this unit are shown where recognized. Patterned ground elsewhere in the valleys is not visible on aerial photographs. Weber and Péwé (1970) have described large rounded ground-ice masses as horizontal and vertical sheets in the Koyukuk Lowland which lies south of our study area.

Undulating lowlands (Q<sub>2</sub>) are characterized by large numbers of flow lines and an intricate drainage pattern. Evidence of eolian morphology is completely absent. Surficial deposits probably consist of reworked terrace deposits and colluvium. South of Huslia is an area of undifferentiated surficial deposits (Q<sub>3</sub>), approximately 15 meters above river level, which might be equivalent to the younger (low) terrace deposits as mapped by Weber and Péwé (1970) for the central part of the Yukon-Koyukuk Lowland.

Relative height differences, slope angles, the frequent occurrence of deflation stripes and surface runoff features form the basis for the distinction of unit Q<sub>1</sub> (various colors on the photographs), undulating uplands from unit Q<sub>2</sub> (slopes of transport reddish-green color). Undulating uplands and slopes of transport vary in elevation from approximately 80 to 215 meters and have a surficial cover of colluvium. Unit Q<sub>2</sub> consists of sediments which have been transported from the undulating uplands to the lowlands and valleys. Unit Q<sub>1</sub> is probably partly underlain at very shallow depth by bedrock, consisting of volcanic rocks (Coss, 1957).

**BEDROCK**

The hilly region west of the Noghahabara Sand Dunes forms part of the Hulo Hills. With the exception of some crests, where a white-bluish color on the photographs indicates outcrops of bare rock, the base of unit K<sub>2</sub> is covered by forests (red in color). Footcliffs, unit K<sub>2</sub>, are less densely vegetated (grey-red color) and covered by deflation stripes. These two units (K<sub>1</sub> and K<sub>2</sub>) vary in height from approximately 170 to 500 meters above sea level and slope angles vary between 10 and 30 degrees. The prominent scarp visible along the northern side of the Dagliit River is probably of structural origin. According to Patton (1964) most of the andesitic volcanic rocks in this area are of mid-Cretaceous (Eocene) age.

**DISCUSSION**

The Yukon-Koyukuk Lowland was not subjected to glaciation during the Quaternary, but was influenced by glacial activity to the north. The glacial periods provided conditions in the Koyukuk Flood-plain and Upper Kuskokwim Region most favorable to the formation of dunes (Fernald, 1968). Aggradation of streams produced a large source of sediment. Sand moved by wind was derived from fluvial or glacial/fluvial deposits.

Based on degree of dune modification, topographic position, and the presence or absence of flow lines we recognize at least three periods of eolian activity. Active dunes and younger stabilized dunes (Q<sub>4</sub> and Q<sub>3</sub>) represent the most recent period of eolian activity where the major eolian transport direction is westerly to northwesterly. Our second period, older stabilized dunes, rose-like in form, (Q<sub>2</sub>) are stabilized by a sparse vegetation cover. Downwinds for both periods can be variable originating from the southeast, east and northeast.

A sharp contrast, between clearly recognizable dune forms (Q<sub>5</sub>, Q<sub>4</sub>, Q<sub>3</sub>) and sand sheets (Q<sub>2</sub>, Q<sub>1</sub>) suggest a considerable time difference between the formation of parabolic, transverse, and rose-shaped dunes and the sand sheets with longitudinal dunes. No inferred parabolic dunes is given for unit Q<sub>2</sub> because the eolian sand cover is relatively thin and discontinuous and has no clearly recognizable dune forms. Based on topographic position unit Q<sub>2</sub> appear to be older than units Q<sub>3</sub>, Q<sub>4</sub>, Q<sub>5</sub>, yet younger than unit Q<sub>1</sub>.

For the Upper Kuskokwim Region, which lies approximately 325 km south-southwest of Huslia, Fernald (1968) has noted two periods of eolian activity based on the sharp degree of dune modification. Whether our period of eolian activity on comparable to Fernald's (1968) remains to be seen. The position of the dune fields and the geomorphology of individual dunes indicate general easterly winds as the effective present and past dune-building winds. This is also the case for comparable dune fields in the Kotzeb River Valley (Fernald, 1968), which lie approximately 150 km to the northwest of Huslia. We feel the clearly recognizable dune forms (units Q<sub>5</sub>, Q<sub>4</sub>, and Q<sub>3</sub>) are Holocene in age and the sand sheets (unit Q<sub>2</sub>) are probably Late Wisconsin in age. The very conspicuous Noghahabara Sand Dunes merit further attention, as the recent nature of the dunes and the processes leading to their actively drifting character yet remain unknown.

**REFERENCES CITED**

Coss, I. T., 1957, Reconnaissance Geologic Map of the Kotzeb River Quadrangle, Alaska. U.S. Geological Survey Miscellaneous Geologic Investigation Map 1-263, 1 sheet, scale 1:250,000.

Lewis, T. H., 1956, Yakutat and Huslia Earthquakes of 1956. Science in Alaska 1956. Proceedings Ninth Alaskan Science Conference, p. 94-95 (eds.).

Fernald, A. T., 1968, Geomorphology of the Upper Kuskokwim Region, Alaska. U.S. Geological Survey Bulletin 1181-A, 31 p.

Galloway, J. P., and Koster, E. A., in press, Comparison of grain-size statistics from two northern Alaskan dune fields. In: Coss, W. and Elliott, R., eds., The United States Geological Survey in Alaska Accomplishments during 1981. U.S. Geological Survey Circular.

Gary, M., Jacobson, L., Jr., and Wolf, C. L., eds., 1972, Glossary of Geology, American Geological Institute, Washington, D.C., 802 p.

Patton, W. W., Jr., 1966, Regional geology of the Kotzeb River Quadrangle, Alaska. U.S. Geological Survey Miscellaneous Geologic Investigations Map 1-431, 1 sheet, scale 1:250,000.

Péwé, J. L., 1975, Quaternary Geology of Alaska. U.S. Geological Survey Professional Paper 835, 145 p.

Weber, J. H., and Péwé, J. L., 1970, Surficial and engineering geology of the central part of the Yukon-Koyukuk Lowland, Alaska. U.S. Geological Survey Miscellaneous Geologic Investigations Map 1-390, 2 sheets, scale 1:125,000.

**ACKNOWLEDGMENTS**

We wish to thank Dr. H. van Zuiden (R.T.C.), Drs. A.C.J.M. Hession and Drs. J.L.Y. Castel (University of Amsterdam) for assistance in preparing this map.

**FOOTNOTES**

(1) UNIVERSITY OF AMSTERDAM, Physical Geography and Soil Science Laboratory, The Netherlands.

(2) U.S. GEOLOGICAL SURVEY, Heron Park, CA, U.S.A., 94025.

(3) INTERNATIONAL INSTITUTE FOR AERIAL SURVEY AND EARTH SCIENCES, Enschede, The Netherlands.

(4) aerial photograph used for this study: false-color infrared - 1:60,000, July 70 Alaska C-3, 60 red IR, photographs 143 to 155 (black and white - NASA, SC 386, June 70 Alaska SW 120 red IR, photographs 079 to 085).

(5) divide (arrow) the line of separation, or ridge, apex, or narrow crest of high ground, marking the boundary between two adjacent drainage basins or dividing the surface waters that flow laterally in one direction from those that flow in the opposite direction (U.S. drainage divide) (Gary and others, 1972).

Photo-interpretation Map of Surficial Deposits and Landforms of the Noghahabara Sand Dunes and part of the Koyukuk Lowland, Alaska

by  
Eduard A. Koster (1), John P. Galloway (2), and Toon Pronk (3)  
1984

This report is preliminary and has not been edited for conformity with Geological Survey standards and nomenclature.