

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

**Analytical results and sample locality map
of stream-sediment and heavy-mineral-concentrate samples
from the Moses and Dennison Peak Roadless Areas,
Tulare County, California**

By

J. A. Domenico, R. J. Goldfarb, D. L. Leach,
and S. M. Smaglik

Open-File Report 84-283

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature. Any use of trade names is for descriptive purposes only and does not imply endorsement by the USGS.

1984

CONTENTS

	Page
Studies related to wilderness	1
Introduction.....	1
Methods of study.....	1
Sample collection.....	1
Stream-sediment samples.....	3
Heavy-mineral-concentrate samples.....	3
Sample preparation.....	3
Sample analysis.....	3
Spectrographic method.....	3
RASS.....	4
References cited.....	4

TABLES

TABLE 1. Limits of determination for spectrographic analysis of rocks and stream sediments.....	5
TABLE A1. Data for stream-sediment samples from the Moses and Dennison Peak Roadless Areas.....	6
TABLE A2. Data from heavy-mineral, nonmagnetic concentrate samples from the Moses and Dennison Peak Roadless Areas.....	12

ILLUSTRATIONS

FIGURE 1. Location map of the Moses and Dennison Peak Roadless Areas, Tulare County, California.....	2
PLATE 1. Sample localities and generalized geology of the Moses and Dennison Peak Roadless Areas, California.....	In pocket

STUDIES RELATED TO WILDERNESS

The Wilderness Act (Public Law 88-577, September 3, 1964) and related acts require the U.S. Geological Survey and the U.S. Bureau of Mines to survey certain areas on Federal lands to determine their mineral resource potential. Results must be made available to the public and be submitted to the President and the Congress. This report presents the results of a geochemical survey of the Moses and Dennison Peak Roadless Areas in the Sequoia National Forest, Tulare County, California. Moses and Dennison Peak Roadless Areas were classified as further planning areas during the Second Roadless Area Review and Evaluation (RARE II) by the U.S. Forest Service, January 1979.

INTRODUCTION

From 1979 to 1982, we conducted a reconnaissance geochemical survey of the Moses and Dennison Peak Roadless Areas, Tulare County, California.

The Moses and Dennison Peak Roadless Areas comprise about 45 mi² in the Sequoia National Forest, of Tulare County, California, and lie immediately southwest of Sequoia National Park (see plate 1). Springville lies 4 miles to the southwest and Visalia is 30 miles to the northwest. Access to the study area is provided by California Highway 190, Gulch Park Road and Bear Creek Road.

The exposed rocks within Moses and Dennison Peak Roadless Areas are primarily Jurassic or Cretaceous plutons of the Sierra Nevada Batholith. Some of these plutons are separated by septa of metamorphic roof pendants. These septa consist of quartzite, schist, marble, and calc-silicate rocks of the Tule River pendants within the Jurassic-Triassic Kings Sequence (Saleeby and others, 1978). The geology of the study areas is discussed in detail by Sawlan and Ort (in press).

Both roadless areas are characterized by extreme relief and dense vegetation. Dennison Peak Roadless Area ranges in elevation from 8400 ft near Dennison Mountain to 3600 ft on Backbone Creek. The North Fork of the Tule River and Dillonwood Grove State Forest, separate Dennison Peak Roadless Area from the northern part of Moses Roadless Area to the southeast. In the northern part of Moses Roadless Area, elevations range from 9300 ft on top of Moses Mountain to 3700 ft along the North Fork of the Tule River. Mountain Home State Forest divides this northern part from the southern part of the Moses Roadless Area, which is centered about the North Fork of the Middle Fork of the Tule River.

METHODS OF STUDY

Sample Collection

We collected stream-sediment samples at 85 sites (plate 1). At many of those sites, we also collected a heavy-mineral concentrate. We analyzed 85 stream-sediment samples, and 58 panned-concentrate samples, for a sampling density of about 1 sample per 0.5 mi² for the stream sediment.




Figure 1. Location map of the Moses and Dennison Peak Roadless Areas, Tulare County, California.

Stream-sediment samples

Analyses of the stream-sediment samples represent the chemistry of the rock material eroded from the drainage basin upstream from each sample site. Such information is useful in identifying those basins which contain concentrations of elements that may be related to mineral deposits.

The stream-sediment samples consisted of active alluvium collected primarily from first-order (unbranched) and second-order (below the junction of two first-order) streams as shown on USGS topographic maps (scale = 1:62,500). Each sample was composited from several localities within an area that may extend as much as 200 ft from the site plotted on the map.

Heavy-mineral-concentrate samples

We panned heavy-mineral-concentrate samples from the same active alluvium as the stream-sediment samples. Each bulk sample was passed through a 2.0-mm (10-mesh) screen to remove the coarse material. The sediment passing through the screen was panned until most of the quartz, feldspar, organic material, and clay-sized material was removed. The sample was air dried.

Sample Preparation

We sieved the stream-sediment samples at the collection site through a 2-mm screen and the minus 2 mm material was retained. The samples were air dried and sieved to minus-80-mesh using stainless steel sieves. The portion of the sediment passing through the sieve was split and a representative fraction was saved for analysis.

After panning the sediment, we used bromoform to separate and remove the remaining quartz and feldspar from the heavy-mineral concentrate. The heavy minerals (specific gravity >2.8) were separated into three fractions using a large electromagnet (in this case a modified Frantz Isodynamic Separator). The most magnetic material (largely magnetite) was discarded. The second fraction (largely ferromagnesian silicates and iron oxides) was saved for analysis/archival storage. The third fraction (the least magnetic material including nonmagnetic ore and ore related minerals) was hand ground for spectrographic analysis.

The magnetic separates discussed are the same separates that would be produced by removing the magnetite with a hand magnet and then using a Frantz Isodynamic Separator set at a slope of 15° and a tilt of 10° with a current of 0.1 ampere to remove the ilmenite, and a current of 1.0 ampere to split the remainder of the sample into magnetic and nonmagnetic fractions.

Sample Analysis

Spectrographic method

We analyzed the stream-sediment and heavy-mineral-concentrate samples for 31 elements using a semiquantitative, direct-current arc emission spectrographic method (Grimes and Marranzino, 1968) (tables A1 and A2). Spectrographic results were obtained by visual comparison of spectra derived from the sample against spectra obtained from standards made from pure oxides and carbonates. Standard concentrations are geometrically spaced over any given order of magnitude of concentration as follows: 100, 50, 20, 10, and so forth. Samples whose concentrations are estimated to fall between those

values are assigned values of 70, 30, 15, and so forth. The precision of the analytical method is approximately plus or minus one reporting unit at the 83 percent confidence level and plus or minus two reporting units at the 96 percent confidence level (Motooka and Grimes, 1976). Values determined for the major elements (iron, magnesium, calcium, and titanium) are given in weight percent; all others are given in parts per million (micrograms/gram) (table 1).

ROCK ANALYSIS STORAGE SYSTEM

Upon completion of all analytical work, the analytical results were entered into a computer-based file called RASS (Rock Analysis Storage System). This RASS file contains both descriptive geological information and analytical data. Any or all of this information may be retrieved and converted to a standard form (STATPAC) for computerized statistical analysis or publication (VanTrump and Miesch, 1976).

REFERENCES CITED

- Grimes, D. J., and Marranzino, A. P., 1968, Direct-current arc and alternating-current spark emission spectrographic field methods for the semiquantitative analysis of geologic materials: U.S. Geological Survey Circular 591, 6 p.
- Motooka, J. M., and Grimes, D. J., 1976, Analytical precision of one-sixth order semiquantitative spectrographic analyses: U.S. Geological Survey Circular 738, 25 p.
- Saleeby, J. B., Goodwin, S. E., Sharp, W. D., and Busby, C. J., 1978, Early Mesozoic paleotectonic reconstruction of the southern Sierra Nevada region, in Mesozoic Paleogeography of the Western United States: Society of Economic Paleontologists and Mineralogists Pacific Section, p. 311-336.
- Sawlan, M. G., and Ort, K. M., in press, Geologic map of the Moses and Dennison Peak Roadless Areas, Tulare County, California,: U.S. Geological Survey MF Map 1651-C.
- VanTrump, George, Jr., and Miesch, A. T., 1976, The U.S. Geological Survey RASS-STATPAC system for management and statistical reduction of geochemical data: Computers and Geosciences, v. 3, p. 475-488.

TABLE 1.--Limits of determination for the spectrographic analysis of stream sediments, based on a 10-mg sample

[The spectrographic limits of determination for heavy-mineral-concentrate samples are two reporting units higher than the limits given for stream sediments]

Elements	Lower determination limit	Upper determination limit
Percent		
Iron (Fe)	0.05	20
Magnesium (Mg)	.02	10
Calcium (Ca)	.05	20
Titanium (Ti)	.002	1
Parts per million		
Manganese (Mn)	10	5,000
Silver (Ag)	0.5	5,000
Arsenic (As)	200	10,000
Gold (Au)	10	500
Boron (B)	10	2,000
Barium (Ba)	20	5,000
Beryllium (Be)	1	1,000
Bismuth (Bi)	10	1,000
Cadmium (Cd)	20	500
Cobalt (Co)	5	2,000
Chromium (Cr)	10	5,000
Copper (Cu)	5	20,000
Lanthanum (La)	20	1,000
Molybdenum (Mo)	5	2,000
Niobium (Nb)	20	2,000
Nickel (Ni)	5	5,000
Lead (Pb)	10	20,000
Antimony (Sb)	100	10,000
Scandium (Sc)	5	100
Tin (Sn)	10	1,000
Strontium (Sr)	100	5,000
Vanadium (V)	10	10,000
Tungsten (W)	50	10,000
Yttrium (Y)	10	2,000
Zinc (Zn)	200	10,000
Zirconium (Zr)	10	1,000
Thorium (Th)	100	2,000

Table A1--Data for stream sediment samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.

Sample	LATITUDE	LONGITUDE	S-FEX	S-MGX	S-CAZX	S-TIX	S-MN	S-AG	S-AS	S-AU	S-B	S-BA
01	36 18 12	118 38 46	3.0	1.5	3.0	.70	1,500	N	N	N	15	500
02	36 18 19	118 39 44	3.0	.7	.7	.50	1,000	N	N	N	70	300
03	36 18 8	118 39 41	3.0	.5	1.0	.50	1,500	N	N	N	10	300
04	36 18 3	118 39 59	3.0	1.0	.7	.70	1,000	N	N	N	70	300
05	36 17 19	118 39 49	2.0	.7	.7	.30	1,000	N	N	N	20	200
06	36 17 5	118 39 52	7.0	1.5	1.0	.70	1,000	N	N	N	70	300
07	36 16 32	118 39 52	5.0	1.0	.3	.50	700	N	N	N	70	300
08	36 16 13	118 39 59	7.0	1.5	1.0	.70	1,500	N	N	N	70	700
09	36 14 30	118 37 4	7.0	3.0	2.0	1.00	1,000	N	N	N	70	300
10	36 14 27	118 37 1	10.0	3.0	2.0	.70	1,000	N	N	N	50	200
11	36 14 1	118 37 50	7.0	2.0	1.5	.50	700	N	N	N	100	300
12	36 13 57	118 37 49	15.0	3.0	3.0	1.00	1,500	N	N	N	100	300
13	36 13 28	118 38 25	7.0	3.0	2.0	.70	1,000	N	N	N	70	200
14	36 13 5	118 39 6	3.0	1.5	.2	.50	1,000	N	N	N	150	500
15	36 12 11	118 38 44	2.0	.5	>20.0	.10	3,000	N	N	N	10	100
16	36 11 27	118 40 8	7.0	2.0	5.0	.30	1,500	N	N	N	10	300
17	36 10 11	118 42 23	5.0	1.5	2.0	.30	1,000	N	N	N	10	200
18	36 9 34	118 43 49	3.0	.3	.5	.20	500	N	N	N	15	500
19	36 11 25	118 44 51	5.0	1.5	5.0	.30	1,000	N	N	N	15	300
20	36 17 8	118 46 5	1.0	.2	.3	.20	1,000	N	N	N	30	300
21	36 17 52	118 46 30	2.0	.2	.3	.30	700	N	N	N	30	500
22	36 18 19	118 46 48	1.0	.2	.2	.20	700	.5	N	N	20	500
23	36 18 8	118 47 25	3.0	1.0	.7	.50	500	N	N	N	20	200
24	36 18 8	118 47 17	3.0	.3	.2	.20	1,000	N	N	N	20	700
25	36 18 28	118 48 28	7.0	3.0	2.0	1.00	1,500	N	N	N	30	700
26	36 17 26	118 42 11	3.0	.5	.3	.30	1,500	N	N	N	20	1,000
27	36 17 11	118 42 36	3.0	.2	.2	.20	700	N	N	N	15	500
28	36 13 24	118 36 59	7.0	3.0	3.0	1.00	1,500	N	N	N	20	500
29	36 12 38	118 36 51	3.0	2.0	3.0	.50	700	.5	N	N	150	300
30	36 11 59	118 38 8	7.0	2.0	1.5	1.00	1,000	N	N	N	100	300
31	36 11 39	118 41 33	5.0	.5	1.0	.20	500	N	N	N	N	200
32	36 11 30	118 36 49	7.0	3.0	2.0	.70	1,500	.5	N	N	70	500
33	36 10 59	118 41 34	5.0	.7	1.0	.20	700	N	N	N	10	200
34	36 9 16	118 45 6	3.0	.5	2.0	.20	700	N	N	N	10	700
35	36 11 23	118 46 7	3.0	1.5	3.0	.30	1,000	N	N	N	20	700
36	36 18 45	118 41 37	2.0	.2	.5	.15	500	N	N	N	15	300
37	36 18 27	118 41 8	1.5	.7	.7	.20	300	N	N	N	20	200
38	36 18 30	118 42 59	1.5	.5	.7	.20	300	N	N	N	N	300
39	36 18 30	118 43 5	1.5	.7	.5	.20	300	N	N	N	N	200
40	36 17 8	118 44 1	2.0	.3	.3	.20	700	N	N	N	20	500
41	36 16 35	118 44 51	3.0	.7	.7	.30	1,000	N	N	N	70	700
42	36 15 40	118 44 10	7.0	1.5	1.0	.30	700	N	N	N	10	300
43	36 19 7	118 47 40	3.0	1.5	1.0	.50	1,000	N	N	N	50	200
44	36 18 18	118 47 2	3.0	.7	.5	.20	500	N	N	N	50	700
45	36 18 7	118 47 18	3.0	.7	.7	.20	700	N	N	N	30	700

Table A1--Data for stream sediment samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.

Sample	S-BE	S-BI	S-CD	S-CO	S-CR	S-CU	S-LA	S-MO	S-NB	S-NI	S-PB
01	3.0	N	N	7	20	10	100	<5	N	10	70
02	2.0	N	N	10	70	15	50	N	N	30	20
03	2.0	N	N	7	20	10	50	N	N	<5	30
04	2.0	N	N	15	70	15	70	N	20	30	20
05	10.0	N	N	5	50	7	N	N	N	20	70
06	3.0	N	N	20	100	20	50	N	20	30	50
07	5.0	N	N	15	70	20	70	7	20	50	70
08	3.0	N	N	20	100	20	50	N	20	30	50
09	1.0	N	N	30	100	20	50	N	N	20	30
10	1.0	N	N	30	150	20	30	N	<20	20	15
11	1.0	N	N	30	150	20	50	N	N	30	20
12	1.0	N	N	50	300	30	100	N	<20	70	20
13	1.0	N	N	30	150	20	30	N	N	50	10
14	2.0	N	N	20	70	20	30	N	N	70	70
15	1.5	N	N	5	20	7	N	20	N	20	20
16	2.0	N	N	15	100	10	N	N	N	10	50
17	1.5	N	N	15	30	5	N	N	N	5	20
18	2.0	N	N	<5	N	20	50	N	N	<5	30
19	2.0	N	N	10	15	15	20	N	N	5	30
20	5.0	N	N	5	10	15	50	N	N	15	50
21	5.0	N	N	7	15	30	100	5	N	15	100
22	3.0	N	N	N	30	10	70	N	N	<5	50
23	1.0	N	N	20	70	20	N	N	N	15	20
24	2.0	N	N	7	20	20	70	<5	N	10	70
25	1.0	N	N	30	100	50	300	N	N	20	30
26	5.0	N	N	7	20	20	200	7	<20	15	150
27	3.0	N	N	<5	<10	5	70	<5	N	5	30
28	1.0	N	N	50	150	30	50	N	N	30	30
29	1.0	N	N	50	150	100	30	N	N	50	30
30	1.0	N	N	30	100	30	30	N	N	50	30
31	2.0	N	N	10	20	5	20	N	N	5	30
32	1.0	N	N	30	100	50	50	N	N	50	50
33	2.0	N	N	7	30	<5	20	N	N	5	20
34	1.5	N	N	5	<10	20	30	N	N	5	50
35	1.5	N	N	10	30	15	100	N	N	20	50
36	2.0	N	N	5	N	<5	50	N	N	<5	30
37	2.0	N	N	5	15	7	N	N	N	10	30
38	1.5	N	N	5	N	<5	30	N	N	<5	30
39	2.0	N	N	5	N	<5	20	N	N	<5	30
40	3.0	N	N	5	10	10	70	N	N	5	50
41	2.0	N	N	7	15	15	N	5	N	5	30
42	2.0	N	N	15	20	10	70	N	N	5	30
43	1.5	N	N	15	20	15	N	N	N	15	20
44	2.0	N	N	10	30	20	70	N	20	15	70
45	2.0	N	N	10	30	20	70	N	20	20	70

Table A1--Data for stream sediment samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.

Sample	S-SB	S-SC	S-SN	S-SR	S-SV	S-SW	S-Y	S-ZN	S-ZR	S-TH
01	N	20	N	300	150	N	70	N	300	N
02	N	15	N	150	150	N	50	N	300	N
03	N	10	N	150	100	N	50	N	500	N
04	N	15	N	100	100	N	50	N	500	N
05	N	5	N	100	50	N	70	N	200	N
06	N	20	N	100	150	N	70	N	300	N
07	N	15	N	N	100	N	70	N	200	N
08	N	20	N	200	150	N	70	N	500	N
09	N	50	N	300	300	N	70	N	300	N
10	N	50	N	200	300	N	70	N	700	N
11	N	30	N	100	200	N	50	N	200	N
12	N	30	N	200	300	N	70	N	300	N
13	N	30	N	200	300	N	70	N	500	N
14	N	15	N	N	200	N	30	N	300	N
15	N	5	N	500	50	N	20	N	30	N
16	N	30	N	300	200	N	70	N	500	N
17	N	20	N	200	150	N	50	N	500	N
18	N	7	N	200	70	N	50	N	300	N
19	N	20	N	300	100	N	50	N	300	N
20	N	5	N	<100	30	N	70	N	200	N
21	N	7	N	N	70	N	100	N	500	N
22	N	5	N	N	20	N	70	N	300	N
23	N	20	N	150	200	N	50	N	1,000	N
24	N	5	N	N	100	N	100	N	500	N
25	N	50	N	300	300	N	100	N	1,000	N
26	N	10	N	N	100	N	150	200	500	N
27	N	5	N	N	50	N	70	N	700	N
28	N	50	N	300	300	N	100	<200	500	N
29	N	20	N	300	200	N	50	<200	300	N
30	N	30	N	150	300	N	50	N	200	N
31	N	15	N	150	100	N	30	N	300	N
32	N	30	N	200	200	N	70	N	300	N
33	N	15	N	150	150	N	50	N	700	N
34	N	7	N	150	100	N	50	N	300	N
35	N	15	N	200	100	N	50	N	300	N
36	N	5	N	200	50	N	20	N	300	N
37	N	5	N	150	50	N	15	N	100	N
38	N	5	N	200	50	N	10	N	50	N
39	N	5	N	100	50	N	15	N	200	N
40	N	5	N	N	30	N	70	N	300	N
41	N	7	N	100	70	N	50	N	1,000	N
42	N	20	N	150	150	N	70	N	500	N
43	N	15	N	150	100	N	20	N	200	N
44	N	10	N	150	100	N	50	N	500	N
45	N	7	N	100	100	N	50	N	300	N

Table A1--Data for stream sediment samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.--continued

Sample	LATITUDE	LONGITUDE	S-FEX	S-MGX	S-CAZ	S-TIX	S-MN	S-AG	S-AS	S-AU	S-B	S-BA
46	36 17 57	118 47 11	5.0	1.0	.7	.30	700	N	N	N	70	700
47	36 17 56	118 47 23	5.0	1.0	1.0	.70	1,000	30.0	N	N	100	500
48	36 17 52	118 47 17	5.0	1.0	.7	1.00	700	N	N	N	70	700
49	36 17 45	118 47 13	5.0	1.5	1.5	.70	700	N	N	N	50	700
50	36 17 21	118 47 3	3.0	1.0	1.0	.50	700	N	N	N	70	700
51	36 14 32	118 39 14	2.0	1.0	.5	.20	700	N	N	N	20	100
52	36 14 22	118 38 56	5.0	1.5	2.0	.50	1,000	.5	N	N	100	200
53	36 17 59	118 42 4	2.0	.7	1.0	.30	300	N	N	N	N	300
54	36 17 30	118 42 7	1.5	.3	.3	.20	500	N	N	N	15	700
55	36 15 27	118 42 41	1.5	.5	.7	.20	200	N	N	N	N	700
56	36 12 10	118 39 11	3.0	.7	.5	.30	300	N	N	N	100	300
57	36 9 37	118 44 34	15.0	.5	2.0	.30	1,000	N	N	N	10	1,000
58	36 10 41	118 41 56	15.0	3.0	5.0	.20	1,500	7.0	N	N	15	300
59	36 18 32	118 47 26	7.0	.7	.7	.50	1,500	1.0	N	N	30	1,000
60	36 15 50	118 41 45	7.0	2.0	.7	.50	1,500	<.5	N	N	20	1,000
61	36 16 3	118 42 8	5.0	1.0	.5	.30	1,000	N	N	N	10	1,500
62	36 16 50	118 41 32	7.0	.5	.5	.30	1,500	N	N	N	10	1,000
63	36 17 14	118 41 33	5.0	.3	.5	.30	2,000	N	N	N	30	1,000
64	36 17 40	118 41 44	7.0	.5	1.0	.50	1,000	N	N	N	50	700
65	36 18 10	118 41 36	5.0	.7	2.0	.50	1,000	20.0	N	N	10	1,000
66	36 16 45	118 43 23	3.0	.7	.7	.20	700	N	N	N	20	1,000
67	36 16 47	118 42 23	5.0	1.0	.7	.50	1,500	N	N	N	20	1,000
68	36 15 55	118 42 41	5.0	1.0	.5	.50	1,000	N	N	N	20	1,000
69	36 15 57	118 43 29	5.0	.7	.5	.30	1,500	N	N	N	20	1,000
70	36 16 5	118 44 44	10.0	1.5	1.5	.20	1,000	N	N	N	10	700
71	36 10 10	118 43 30	15.0	3.0	10.0	1.00	2,000	N	N	N	10	500
72	36 10 51	118 42 56	7.0	1.0	.7	.50	1,000	N	N	N	20	1,000
73	36 18 46	118 43 36	7.0	1.0	.5	.70	1,000	N	N	N	20	1,000
74	36 17 55	118 44 15	2.0	.5	.5	.30	1,000	N	N	N	20	700
75	36 17 20	118 44 39	7.0	2.0	2.0	.50	2,000	N	N	N	150	700
76	36 16 0	118 40 16	7.0	2.0	5.0	.50	1,500	N	N	N	10	500
77	36 10 40	118 44 7	20.0	5.0	7.0	1.00	2,000	N	N	N	10	700
78	36 18 40	118 44 24	7.0	1.5	.7	.50	1,000	N	N	N	20	1,500
79	36 18 0	118 46 0	5.0	.7	.5	.50	1,000	N	N	N	20	1,500
80	36 18 35	118 47 14	10.0	2.0	5.0	>1.00	1,500	N	N	N	100	700
81	36 15 21	118 40 45	10.0	.3	.7	1.00	3,000	N	N	N	20	2,000
82	36 16 51	118 41 56	2.0	.5	.1	.20	700	N	N	N	30	700
83	36 14 50	118 42 44	2.0	1.0	1.0	.30	500	N	N	N	15	1,000
84	36 14 10	118 41 33	2.0	1.5	2.0	.30	500	N	N	N	<10	700
85	36 14 0	118 39 1	3.0	2.0	1.0	.50	700	N	N	N	100	200

Table A1--Data for stream sediment samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.--continued

Sample	S-BE	S-BI	S-CD	S-CO	S-CR	S-CU	S-LA	S-MO	S-NB	S-NI	S-PB
46	1.0	N	N	15	50	15	N	N	N	10	20
47	1.0	N	N	15	50	20	30	N	N	15	15
48	1.0	N	N	15	70	30	N	N	N	20	30
49	1.5	N	N	20	70	30	20	N	N	20	50
50	1.5	N	N	15	70	20	30	N	N	15	50
51	5.0	<10	N	10	30	10	30	70	N	20	200
52	3.0	N	N	20	150	20	100	N	N	100	50
53	1.0	N	N	7	20	<5	70	<5	N	5	20
54	3.0	N	N	<5	15	15	70	5	<20	5	30
55	1.5	N	N	5	10	10	50	<5	N	5	30
56	1.5	N	N	50	100	20	70	<5	N	70	30
57	N	N	N	5	20	<5	500	N	N	N	20
58	N	N	N	15	100	5	N	N	N	10	20
59	<5.0	N	N	5	30	30	100	5	30	10	150
60	<5.0	N	N	5	20	50	100	5	20	5	70
61	<5.0	N	N	5	10	10	50	N	<20	N	30
62	<5.0	N	N	N	30	20	200	10	20	10	100
63	5.0	N	N	N	10	10	150	N	20	5	70
64	N	N	N	5	20	5	100	N	<20	5	50
65	N	N	N	5	20	N	100	N	N	N	30
66	N	N	N	5	15	7	50	N	N	10	50
67	5.0	N	N	10	30	10	150	N	20	10	70
68	<5.0	N	N	10	20	10	70	<5	<20	10	50
69	<5.0	N	N	5	20	15	100	N	<20	10	70
70	N	N	N	10	30	<5	N	N	N	5	30
71	N	N	N	20	100	7	N	N	N	10	20
72	<5.0	N	N	7	20	10	100	N	20	7	50
73	<5.0	N	N	5	20	10	100	N	<20	7	50
74	5.0	N	N	5	30	30	100	N	<20	10	50
75	7.0	N	N	20	70	30	200	<5	20	20	100
76	N	N	N	20	50	5	70	N	N	5	30
77	N	N	N	30	100	7	N	N	N	15	30
78	<5.0	N	N	5	10	20	100	N	30	5	50
79	N	N	N	5	15	20	50	N	<20	7	70
80	N	N	N	20	100	20	50	N	N	15	30
81	7.0	N	N	N	30	10	200	5	50	10	100
82	5.0	N	N	7	20	20	150	7	20	15	50
83	2.0	N	N	7	20	10	100	N	N	10	20
84	1.0	N	N	15	30	7	N	N	N	7	30
85	2.0	N	N	50	150	30	50	N	N	150	30

Table A1--Data for stream sediment samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.--continued

Sample	S-SB	S-SC	S-SN	S-SR	S-V	S-W	S-Y	S-ZN	S-ZR	S-TH
46	N	15	N	150	150	N	20	N	500	N
47	N	30	N	150	200	N	50	N	1,000	N
48	N	20	N	150	200	N	30	N	500	N
49	N	30	N	200	200	N	30	N	700	N
50	N	30	N	150	200	N	30	N	1,000	N
51	N	10	N	N	50	N	70	N	150	N
52	N	20	N	200	150	N	70	<200	200	N
53	N	10	N	200	150	N	30	N	200	N
54	N	5	<10	N	30	N	100	N	500	N
55	N	7	N	200	50	N	30	N	200	N
56	N	15	N	100	200	N	30	N	200	N
57	N	--	N	200	150	N	70	N	--	N
58	N	--	N	300	200	N	50	N	--	N
59	N	--	N	N	50	N	100	N	--	N
60	N	--	20	100	30	N	100	<200	--	N
61	N	--	30	N	20	N	50	N	--	N
62	N	--	N	N	20	N	150	N	--	N
63	N	--	N	N	15	N	100	N	--	N
64	N	--	N	300	50	N	70	N	--	N
65	N	--	N	500	100	N	20	N	--	N
66	N	--	N	100	50	N	50	N	--	N
67	N	--	20	N	50	N	100	N	--	N
68	N	--	10	N	50	N	70	N	--	N
69	N	--	N	100	50	N	70	N	--	N
70	N	--	N	300	200	N	50	N	--	N
71	N	--	N	300	300	N	70	N	--	N
72	N	--	N	100	70	N	70	N	--	N
73	N	--	N	100	50	N	70	N	--	N
74	N	--	N	N	50	N	70	N	--	N
75	N	--	30	200	30	N	100	500	--	N
76	N	--	N	500	200	N	50	N	--	N
77	N	--	N	500	300	N	100	N	--	N
78	N	--	20	100	50	N	70	N	--	N
79	N	--	N	100	30	N	50	N	--	N
80	N	--	N	300	200	N	50	N	--	N
81	N	--	10	N	30	N	150	N	--	N
82	N	10	10	N	70	N	150	N	300	N
83	N	10	N	200	150	N	50	N	700	N
84	N	20	N	200	150	N	30	N	300	N
85	N	15	N	150	150	N	50	N	300	N

Table A2--Data for heavy-mineral, non-magnetic concentrate samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.

Sample	S-BE	S-BI	S-CD	S-CO	S-CR	S-CU	S-LA	S-MO	S-NB	S-NI	S-PB
02	1	N	N	N	100	15	300	<5	30	N	15
03	1	N	N	10	70	15	150	<5	30	N	30
09	1	200	N	N	50	7	1,000	N	20	N	30
10	1	N	N	N	30	5	>1,000	N	<20	N	30
11	1	300	N	70	70	5	200	10	20	N	150
12	1	70	N	50	50	5	300	N	<20	N	50
13	1	50	N	100	70	20	700	7	<20	N	20
16	1	N	N	15	N	5	300	N	<20	N	100
17	1	20	N	N	20	10	200	N	<20	N	300
19	1	N	N	10	20	7	300	N	30	N	70
21	1	>1,000	N	N	<10	<5	>1,000	7	50	N	500
23	1	N	N	N	<10	N	100	N	N	N	30
24	1	>1,000	N	200	15	50	>1,000	50	20	N	3,000
25	1	70	N	N	N	N	>1,000	N	N	N	N
28	1	500	N	20	100	5	200	N	<20	N	15
30	1	100	N	100	150	10	500	N	<20	N	150
31	1	1,000	N	10	30	5	1,000	N	30	N	100
33	1	N	N	N	20	<5	1,000	N	20	N	100
34	1	N	N	N	N	7	>1,000	10	1,000	N	50
35	1	50	N	N	70	N	>1,000	<5	1,000	N	70
36	1	500	N	N	30	<5	>1,000	N	150	N	70
38	1	N	N	N	20	N	>1,000	N	50	N	20
42	1	>1,000	N	N	10	7	>1,000	20	20	N	N
43	1	50	N	N	N	7	100	N	N	N	15
44	2	>1,000	N	5	50	10	200	20	200	N	500
47	N	200	N	15	70	10	150	<5	N	N	200
48	<1	20	N	10	50	5	50	<5	N	N	50
49	1	N	N	5	70	<5	70	N	N	N	300
50	2	N	N	5	50	N	50	N	N	15	20
53	N	N	N	<10	70	10	500	30	100	10	50
54	2	N	N	N	20	<10	200	N	<50	20	70
56	3	N	N	<10	200	20	70	N	N	30	30
57	N	N	N	N	N	N	700	N	N	N	30
58	N	N	N	10	<20	<10	150	N	N	N	70
59	<2	>2,000	N	<10	N	10	N	70	50	N	700
62	2	30	N	N	N	10	70	10	N	N	50
63	3	<20	N	N	N	N	200	70	50	N	30
64	2	50	N	N	20	N	700	70	50	N	100
65	2	100	N	N	N	N	500	10	70	N	50
66	3	70	N	N	N	N	N	N	N	N	20
67	2	50	N	N	N	N	200	N	<50	N	30
68	3	30	N	N	N	10	N	100	N	N	50
69	3	200	N	N	<20	N	500	10	N	N	30
70	3	200	N	N	N	N	50	N	N	N	30
71	<2	150	N	N	<20	10	N	N	N	N	20

Table A2--Data for heavy-mineral, non-magnetic concentrate samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.

Sample	LATITUDE	LONGITUDE	S-FEZ	S-MGX	S-CAZ	S-TIX	S-MN	S-AG	S-AS	S-AU	S-B	S-BA
02	36 18 19	118 39 44	.50	.30	7.00	1.00	700	N	N	N	70	100
03	36 18 8	118 39 41	.50	2.00	5.00	>1.00	700	N	N	N	50	70
09	36 14 30	118 37 4	.30	.70	5.00	>1.00	700	2.0	N	N	1,000	100
10	36 14 27	118 37 1	.50	.30	3.00	>1.00	700	3.0	N	N	500	150
11	36 14 1	118 37 50	.50	.30	3.00	>1.00	500	2.0	N	N	300	100
12	36 13 57	118 37 49	.70	.70	3.00	>1.00	700	2.0	2,000	N	500	70
13	36 13 28	118 38 25	2.00	.70	5.00	>1.00	700	N	N	N	300	70
16	36 11 27	118 40 8	.50	.10	3.00	>1.00	500	N	N	N	20	150
17	36 10 11	118 42 23	.20	.05	3.00	>1.00	300	N	N	N	20	100
19	36 11 25	118 44 51	.20	.20	5.00	>1.00	700	N	N	N	30	150
21	36 17 52	118 46 30	.50	.03	.50	>1.00	200	N	300	N	20	70
23	36 13 8	118 47 25	.10	.03	.50	1.00	100	N	N	N	30	100
24	36 18 7	118 47 18	1.00	.05	3.00	>1.00	700	2.0	10,000	10	1,000	150
25	36 18 28	118 48 28	.05	.02	.07	.30	70	N	N	N	N	100
28	36 13 24	118 36 59	.20	.70	3.00	>1.00	200	N	N	N	150	70
30	36 11 59	118 38 8	1.00	1.50	3.00	>1.00	500	N	500	N	1,500	500
31	36 11 39	118 41 33	.20	.15	5.00	>1.00	300	N	N	N	30	100
33	36 10 59	118 41 34	.20	.10	3.00	>1.00	200	N	N	N	N	150
34	36 9 16	118 45 6	.05	.05	.20	.02	70	N	N	N	N	70
35	36 11 23	118 46 7	.15	.10	.20	.10	200	N	N	N	500	500
36	36 18 45	118 41 37	.10	.05	5.00	>1.00	200	N	N	N	N	100
38	36 12 30	118 42 59	.20	.05	2.00	.30	200	N	N	N	700	20
42	36 15 40	118 44 10	.50	.03	2.00	>1.00	200	N	500	N	N	70
43	36 19 7	118 47 40	.10	.02	.10	.70	100	N	N	N	100	100
44	36 18 18	118 47 2	.70	.10	2.00	>1.00	300	30.0	N	N	20	300
47	36 17 56	118 47 23	1.00	.50	3.00	>1.00	500	N	N	N	2,000	300
48	36 17 52	118 47 17	1.00	.30	3.00	1.00	500	N	N	N	200	500
49	36 17 45	118 47 13	1.50	.30	5.00	1.00	700	N	N	N	300	150
50	36 17 21	118 47 3	.30	.05	.70	1.00	100	N	N	N	150	300
53	36 17 59	118 42 4	.50	.20	5.00	>2.00	500	N	N	N	100	<50
54	36 17 30	118 42 7	.70	.10	1.00	1.50	300	N	N	N	100	50
56	36 12 10	118 39 11	1.00	.50	1.00	.50	300	N	N	N	200	150
57	36 9 37	118 44 34	.30	.05	.50	.20	70	N	N	N	N	200
58	36 10 41	118 41 56	.30	<.05	2.00	>2.00	100	N	N	N	N	150
59	36 18 32	118 47 26	1.00	.30	.20	2.00	200	50.0	N	30	N	300
62	36 16 50	118 41 32	1.50	.70	.70	.50	300	N	N	N	<20	200
63	36 17 14	118 41 33	.70	<.05	2.00	1.00	150	N	N	N	N	150
64	36 17 40	118 41 44	.50	.05	1.00	2.00	200	N	N	N	100	100
65	36 13 10	118 41 36	.30	<.05	2.00	>2.00	200	N	N	N	N	50
66	36 16 45	118 43 23	.70	.05	.10	.70	150	N	N	N	N	300
67	36 16 47	118 42 23	1.00	.07	.20	2.00	300	N	N	N	N	50
68	36 16 3	118 42 8	1.50	.07	.10	1.00	150	N	N	N	N	200
69	36 15 57	118 43 29	1.00	.07	.15	1.00	200	N	N	N	20	100
70	36 16 5	118 44 44	.50	.05	1.00	2.00	150	N	N	N	30	100
71	36 10 10	118 43 30	.50	.05	1.50	>2.00	150	N	N	N	N	70

Table A2--Data for heavy-mineral, non-magnetic concentrate samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.

Sample	S-SB	S-SC	S-SN	S-SR	S-V	S-W	S-Y	S-ZN	S-ZR	S-TH
C2	N	10	30	N	100	<50	200	N	1,000	200
C3	N	10	50	N	100	<50	300	N	>1,000	<100
C9	N	15	70	N	100	N	700	N	>1,000	N
10	N	15	30	N	100	N	300	N	>1,000	N
11	N	15	50	N	100	200	500	N	>1,000	N
12	N	15	30	100	100	N	300	N	>1,000	N
13	N	15	50	200	100	150	500	N	>1,000	N
16	N	15	30	N	100	N	700	N	>1,000	1,000
17	N	15	30	N	100	N	500	N	>1,000	1,000
19	N	15	70	N	100	N	500	N	>1,000	700
21	100	15	300	N	100	N	700	N	>1,000	300
23	N	15	<10	N	100	<50	500	N	>1,000	1,000
24	100	15	500	N	100	100	2,000	N	>1,000	300
25	N	15	N	N	100	N	300	N	>1,000	300
28	N	15	50	N	100	N	500	N	>1,000	N
30	<100	15	30	N	100	<50	300	N	>1,000	1,000
31	N	15	100	N	100	N	500	N	>1,000	1,000
33	N	15	50	N	100	N	700	N	>1,000	1,000
34	<100	15	10	N	100	N	1,500	N	>1,000	1,500
35	<100	15	<10	N	100	N	1,500	N	>1,000	700
30	N	15	50	N	100	<50	1,500	N	>1,000	500
38	N	15	50	N	100	50	200	N	>1,000	1,000
42	N	15	30	N	100	200	200	N	>1,000	1,000
43	N	15	N	N	100	N	300	N	>1,000	700
44	N	10	100	N	100	200	700	N	>1,000	N
47	N	15	20	N	300	N	200	N	>1,000	200
48	N	10	20	100	150	N	150	N	>1,000	500
49	N	10	N	N	150	N	300	N	>1,000	200
50	N	10	N	N	20	N	700	N	>1,000	1,500
53	N	15	200	N	300	N	500	N	2,000	700
54	N	15	50	N	20	<100	700	N	>2,000	<200
56	N	15	N	<200	300	N	20	N	500	N
57	N	30	20	N	N	<100	200	N	>2,000	500
58	N	30	70	N	300	N	200	N	>2,000	5,000
59	N	50	50	N	70	N	150	N	2,000	300
62	N	30	<20	N	100	N	500	N	>2,000	<200
63	N	30	N	N	20	<100	500	N	>2,000	300
64	N	30	50	N	100	150	500	N	>2,000	1,000
65	N	30	100	N	200	N	300	N	>2,000	1,000
66	200	30	20	N	20	<100	200	N	>2,000	N
67	N	30	150	N	100	N	500	N	>2,000	200
68	N	30	50	N	50	200	300	N	>2,000	300
69	N	30	30	N	50	200	500	N	>2,000	<200
70	N	30	50	N	70	<100	300	N	>2,000	200
71	N	30	70	N	300	<100	200	N	>2,000	200

Table A2--Data for heavy-mineral, non-magnetic concentrate samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.--continued

Sample	LATITUDE	LONGITUDE	S-FEX	S-MGZ	S-CAZ	S-TIX	S-MN	S-AG	S-AS	S-AU	S-B	S-BA
72	36 10 51	118 42 56	.70	.05	2.00	>2.00	150	N	N	N	N	200
73	36 18 46	118 43 36	.70	.10	1.00	2.00	200	N	N	N	N	200
74	36 17 55	118 44 15	.70	.05	1.00	>2.00	200	N	N	N	N	70
75	36 17 20	118 44 39	2.00	.50	.70	2.00	300	N	N	N	N	70
76	36 16 0	118 40 16	1.00	.50	7.00	1.50	1,000	N	N	N	500	150
77	36 10 40	118 44 7	.20	<.05	1.00	>2.00	150	N	N	N	N	100
78	36 18 40	118 44 24	1.50	.05	.50	2.00	700	N	N	N	N	200
79	36 13 0	118 46 0	.70	.05	.50	>2.00	200	N	N	N	N	150
80	36 18 35	118 47 14	1.00	.20	.50	1.50	300	N	N	N	200	200
81	36 15 21	118 40 45	2.00	<.05	<.10	2.00	300	N	N	N	N	150
83	36 14 50	118 42 44	1.50	.10	.20	2.00	300	N	N	N	70	70
84	36 14 10	118 41 33	.50	.10	2.00	2.00	300	N	N	N	50	100
85	36 14 0	118 39 1	1.00	1.00	1.00	2.00	300	N	N	N	500	150

Table A2--Data for heavy-mineral, non-magnetic concentrate samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.--continued

Sample	S-BE	S-BI	S-CD	S-CO	S-CR	S-CU	S-LA	S-MO	S-NB	S-NI	S-PB
72	<2	N	N	10	<20	10	300	N	50	N	100
73	2	N	N	N	<20	<10	N	30	70	N	20
74	N	N	N	N	<20	10	N	N	<50	N	30
75	2	1,000	N	10	70	30	N	10	N	50	50
76	<2	<20	N	15	20	<10	N	N	50	N	30
77	3	N	N	N	N	<10	N	N	N	N	20
78	2	20	N	N	N	10	N	15	50	N	30
79	3	50	N	N	N	<10	N	N	50	N	30
80	5	N	N	N	N	N	N	20	N	N	30
81	2	N	N	N	N	N	N	<10	100	N	30
83	N	50	N	20	20	30	2,000	N	50	50	1,000
84	N	N	N	N	20	15	100	N	50	50	1,000
85	2	N	N	15	150	70	200	N	70	30	200

Table A2--Data for heavy-mineral, non-magnetic concentrate samples from Moses Mountain Wilderness and Dennison Peak Wilderness, California.--continued

Sample	S-SB	S-SC	S-SN	S-SR	S-V	S-W	S-Y	S-ZN	S-ZR	S-TH
72	N	30	100	N	300	N	300	N	>2,000	>5,000
73	N	30	70	N	150	N	300	N	>2,000	700
74	N	30	70	N	20	<100	1,000	N	>2,000	500
75	N	30	30	N	50	N	1,000	N	>2,000	<200
76	N	30	30	N	30	100	150	N	>2,000	N
77	N	30	30	N	100	100	200	N	>2,000	<200
78	N	30	30	N	30	N	300	N	>2,000	200
79	N	30	50	N	20	N	300	N	>2,000	700
80	N	30	N	N	50	100	200	N	>2,000	1,000
81	N	30	20	N	<20	N	150	N	>2,000	N
83	N	15	50	N	50	N	700	N	>2,000	500
84	500	15	500	N	150	N	500	N	>2,000	200
85	N	15	70	<200	300	N	150	N	>2,000	N