

**STUDIES OF GEOLOGY AND HYDROLOGY IN THE
BASIN AND RANGE PROVINCE, SOUTHWESTERN UNITED STATES,
FOR ISOLATION OF HIGH-LEVEL RADIOACTIVE WASTE**

CHARACTERIZATION OF THE SONORAN REGION, ARIZONA

Edited by M.S. Bedinger, K.A. Sargent, and William H. Langer

U.S. Geological Survey Open-File Report 84-741



In press as Professional Paper 1370-D

This is Chapter D of an eight-chapter Professional Paper series being prepared by the U.S. Geological Survey in cooperation with States in the Basin and Range province.

UNITED STATES
DEPARTMENT OF THE INTERIOR
DONALD PAUL HODEL, Secretary
GEOLOGICAL SURVEY
Dallas L. Peck , Director



Copies of this report can be purchased from:

Open-File Services Section
Western Distribution Branch
P.O. Box 25425, Denver Federal Center
Denver, Colorado 80225

BASIN AND RANGE PROVINCE WORKING GROUP

U.S. Geological Survey Members

Chairman of the Province Working Group:

M.S. Bedinger
Hydrologist
U.S. Geological Survey
Denver, CO

Member:

K.A. Sargent
Geologist
U.S. Geological Survey
Denver, CO

State Members and Alternates

ARIZONA

Member:

Larry D. Fellows
State Geologist and Assistant Director
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

Alternate:

H. Wesley Peirce
Principal Geologist
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

CALIFORNIA

Member:

Robert Streitz
Geologist
California Division of Mines and Geology
Sacramento, CA

IDAHO

Member:

Frank B. Sherman
Chief, Ground-Water Section
Idaho Department of Water Resources
Boise, ID

Alternate:

Darrel Clapp
Chief, Technical Services Bureau
Idaho Department of Water Resources
Boise, ID

NEVADA

Member:

John Schilling
State Geologist
Nevada Bureau of Mines and Geology
University of Nevada, Reno
Reno, NV

Alternate:

Susan L. Tingley
Deputy to the State Geologist
Nevada Bureau of Mines and Geology
University of Nevada, Reno
Reno, NV

NEW MEXICO

Member:

James M. Hill
Chief, Bureau of Geology
New Mexico Energy and Minerals Department
Santa Fe, NM

Alternate:

Frank E. Kottowski
Director
New Mexico Bureau of Mines and Mineral Resources
Socorro, NM

TEXAS

Member:

Christopher D. Henry
Geologist
Texas Bureau of Economic Geology
University of Texas at Austin
Austin, TX

Alternate:

Douglas Ratcliff
Associate Director
Texas Bureau of Economic Geology
University of Texas at Austin
Austin, TX

UTAH

Member:

Genevieve Atwood
State Geologist
Utah Geological and Mineral Survey
Salt Lake City, UT

Alternate:

Don R. Mabey
Senior Geologist
Utah Geological and Mineral Survey
Salt Lake City, UT

CONTENTS

	Page
Abstract-----	1
Introduction, by M.S. Bedinger, K.A. Sargent, and Robert B. Scarborough-----	4
Background and purpose-----	4
Geographic setting-----	7
Acknowledgments-----	10
References cited-----	11
Geology, by Robert B. Scarborough-----	13
Stratigraphy-----	13
Precambrian rocks-----	13
Paleozoic rocks-----	15
Mesozoic rocks-----	19
Cenozoic rocks-----	21
Summary of geologic events-----	27
Precambrian tectonics-----	27
Phanerozoic tectonics and volcanism-----	29
Paleozoic-----	29
Mesozoic and Cenozoic denudation-----	30
Pre-Laramide Mesozoic tectonics-----	32
Laramide orogeny-----	38
Metamorphic-core complexes and middle Tertiary detachments-----	41
Middle Tertiary volcanism-----	52
Basin-and-range faulting and volcanism-----	55
Miocene, Pliocene and Quaternary history---	62

	Page
References cited-----	65
Potential host media for radioactive waste, by	
K.A. Sargent-----	82
Intrusive rocks-----	83
Tuffaceous rocks-----	84
Basaltic rocks-----	85
Salt-----	87
Other rocks-----	88
Unsaturated zone-----	88
References cited-----	89
Quaternary tectonism, by K.A. Sargent and Robert B.	
Scarborough-----	90
Seismicity-----	92
Heat flow-----	93
Quaternary faulting-----	93
Late Cenozoic volcanics-----	94
Vertical crustal movement-----	95
References cited-----	96
Ground-water hydrology, by M.S. Bedinger, William H.	
Langer, and J.E. Reed-----	98
Major hydrogeologic units-----	98
Ground-water flow regime-----	101
Ground-water flow analysis-----	103
Areal ground-water flow-----	103
Cross-sectional models-----	105

	Page
Quality of ground water-----	108
Pleistocene hydrologic conditions-----	111
References cited-----	112
Mineral and energy resources, by B.T. Brady and	
Robert B. Scarborough-----	115
Metallic-mineral resources-----	116
Nonmetallic and industrial mineral resources-----	130
Geothermal resources-----	133
Coal, oil, and gas resources-----	134
References cited-----	135

ILLUSTRATIONS

[Plates in pocket]

Plate 1.--Map showing potential host rocks and areas of thick unsaturated zones in the Sonoran region, Arizona.

2.--Map showing hydrogeologic units and relative velocity of ground water at the water table in the Sonoran region, Arizona.

3.--Map showing ground-water traveltimes, flow paths, and natural discharge areas in the Sonoran region, Arizona.

4.--Hydrogeologic sections A-A', B-B', and C-C' showing ground-water flow paths and relative traveltime, Sonoran region, Arizona.

5.--Map showing location of the metallic-mineral districts and mineralized areas in the Sonoran region, Arizona.

	Page
Figure 1.--Map showing physiographic features-----	8
2.--Geographic index map of the Sonoran region and vicinity-----	9
3.--Photograph of overturned Paleozoic section in the Martin Peak area of the Little Harquahala Mountains-----	17
4.--Hypothetical geologic section transverse to main drainage across a typical basin-----	26
5.--Photograph of the Hercules thrust fault, exposed near "S" Mountain, southeast of Salome, La Paz County, Arizona-----	34
6.--Hypothetical geologic section through the southern Plomosa and Harcurvar Mountains-----	43
7.--Photograph of the southern Cabeza Prieta Mountains (Yuma County) looking north- northwest-----	47
8.--Photograph of the southern Growler Mountains capped by volcanic strata-----	53
9.--Photograph of Hualapai Valley and Red Lake Playa-----	58
10.--Map showing deep basins and areas of Pleistocene and Holocene erosion-----	59

	Page
11.--Map showing seismicity, heat flow, Quaternary faults, late Cenozoic volcanic rocks and vertical crustal movement-----	91
12.--Map showing dissolved-solids concentration in ground water-----	109
13.--Map showing distribution of chemical types of ground water-----	110

Table 1.--Hydraulic properties of hydrogeologic units and hydraulic gradients used in estimating relative ground-water velocities at the water table-----	104
2.--Hydraulic properties of hydrogeologic units used in cross-sectional models-----	106
3.--Production summary of base and precious metals for metallic-mineral districts with total production in excess of 100,000 short tons by county (including estimates for years data not available)-----	117

CONVERSION FACTORS

For use of readers who prefer to use inch-pound units, conversion factors for terms used in this report are listed below.

U.S. Customary units and troy weights are used in the "Mineral and Energy Resources" section of this report where such terms are widely accepted and utilized by the mineral and energy fuel industries.

<u>Multiply</u>	<u>By</u>	<u>To obtain</u>
<u>Length</u>		
millimeter (mm)	0.03937	inch (in.)
meter (m)	3.281	foot (ft)
kilometer (km)	0.6214	mile (mi)
<u>Area</u>		
square kilometer (km ²)	3.861	square mile (mi ²)
<u>Acceleration</u>		
milligal (mGal)	No Conversion	
<u>Volume</u>		
liter (L)	0.2642	gallon (gal)
cubic kilometer (km ³)	0.2399	cubic mile (mi ³)
<u>Flow</u>		
liter per minute (L/min)	0.2642	gallon per minute (gal/min)
meter per day (m/d)	3.281	foot per day (ft/d)

Temperature

degree Celsius (° C)

$$F = \frac{9}{5} C + 32$$

degree Fahrenheit (° F)

Mass

megagram (Mg) or
metric ton

1.102

short ton (2,000 lb)

Chemical concentrations

milligram
per liter (mg/L)

About 1

part per million

**STUDIES OF GEOLOGY AND HYDROLOGY IN THE
BASIN AND RANGE PROVINCE, SOUTHWESTERN UNITED STATES,
FOR ISOLATION OF HIGH-LEVEL RADIOACTIVE WASTE**

**CHARACTERIZATION OF THE
SONORAN REGION, ARIZONA**

Edited by M.S. Bedinger, K.A. Sargent, and William H. Langer,
U.S. Geological Survey

ABSTRACT

The Sonoran region, southwestern Arizona, of the Basin and Range province, is south and east of the Colorado River. The structural basins typically are filled with 200 to 1,500 meters of clastic material with thick evaporite sections present locally. Basins of the region generally trend north-northwest. Relief between the valleys and adjacent mountains generally is 300 to 1,300 meters. Surface drainage is to the Colorado River, major tributaries being the Bill Williams and Gila Rivers. A few basins have interior or poorly integrated drainage.

Bedrock is exposed in about one-fourth of the region. Bedrock consists of Precambrian metamorphic and plutonic rocks, Paleozoic sedimentary rocks, Mesozoic volcanic and sedimentary rocks, and middle to late Tertiary volcanic and plutonic rocks. In addition to igneous tectonism, the rocks have been repeatedly deformed and moved along slight and steep-angle faults. Present topography was largely shaped by Tertiary basin-and-range faulting, erosion, and deposition.

Potential host media for isolation of high-level radioactive waste in the Sonoran region, Arizona, include intrusive rocks, tuffaceous rocks, basaltic rocks, laharic breccias, and salt. Basin-fill deposits, and possibly other rock types, have potential as host media in the unsaturated zone.

Quaternary tectonism in the region is indicated in the northwest and southern part of the region by relatively substantial strain release, epicenters of few earthquakes greater than magnitude 4 (Richter scale, surface wave), and vertical crustal movement. Quaternary volcanic rocks may be present in the late Cenozoic volcanic field in the southeastern part of the region.

The Sonoran region is arid with annual precipitation less than 200 millimeters throughout most of the region and annual potential evaporation greater than 2.5 meters. Recharge of ground water occurs principally in areas of higher altitude where bedrock is permeable and by infiltration of mountain runoff in the basin areas. Discharge of ground water occurs by seepage to gaining streams, evapotranspiration, springs, and by withdrawal from wells. In most of the region, the concentration of dissolved solids in ground water is less than 500 milligrams per liter. Ground water near the Gila River and the Colorado River contains 1,000 milligrams per liter or more dissolved solids. The ground water is mostly of the calcium magnesium or sodium bicarbonate type. Sulfate and chloride type waters occur in and near large discharge areas and in some playas.

Mineral development in the 19th century was confined to precious-metal lodes, base-metal vein and replacement ores, and placer deposits. More recent development has focused on large scale open-pit copper mines, which produce substantial quantities of precious metals as byproducts.

INTRODUCTION

By M.S. Bedinger and K.A. Sargent, U.S. Geological Survey
and Robert B. Scarborough,
Arizona Bureau of Geology and Mineral Technology

Background and Purpose

A study by the U.S. Geological Survey to evaluate potential hydrogeologic environments for isolation of high-level radioactive waste in the Basin and Range physiographic province of the southwestern United States was begun in May, 1981 with the introduction of the study to the Governors of eight Basin and Range States--Arizona, California, Idaho, Nevada, New Mexico, Oregon, Texas, and Utah, and respective Indian tribes in those States. Accordingly, these States were invited to participate in the study by designating an Earth scientist to serve on a Province Working Group with the Survey--membership of the working group follows the title page. State representatives have provided consultation in selecting guidelines, assembling geologic and hydrologic data, and assessing such information to identify environments that meet the guidelines for further study.

The guidelines for evaluation of the regions and the rationale for the treatment, and the basis for hydrogeologic characterization of the regions are given in Chapter A of this Professional Paper (Bedinger, Sargent, and others, 1985). The evaluation of the region is given in Chapter H (Bedinger, Sargent, and Langer, 1985). The titles of chapters in this series are as follows:

- A - Basis of characterization and evaluation
- B - Characterization of the Trans-Pecos region, Texas
- C - Characterization of the Rio Grande region, New Mexico and Texas
- D - Characterization of the Sonoran region, Arizona
- E - Characterization of the Sonoran region, California
- F - Characterization of the Death Valley region, Nevada and California
- G - Characterization of the Bonneville region, Utah and Nevada
- H - Evaluation of the regions

These chapters are closely integrated and contain a minimum of repetition. The reader needs to consult Chapters A and H, and the appropriate regional Chapters B through G in order to achieve a complete understanding of the characterization and evaluation of an individual region.

Additional background information on this study is given in reports on the province phase of characterization and evaluation by Bedinger, Sargent, and Reed (1984); Sargent and Bedinger (1985); and Bedinger, Sargent, and Brady (1985).

This report, characterization of the Sonoran region, Arizona, Chapter D, is one of six reports characterizing the geology and hydrology of regions in the Basin and Range province. Chapter D is divided into six separately authored sections: (1) Introduction; (2) geology; (3) potential host media; (4) Quaternary tectonism; (5) ground-water hydrology; and (6) mineral and energy resources. Although this report was prepared under the general guidelines set up by the Province Working Group, the scope of individual sections is established by their respective authors.

This chapter provides the geologic and hydrologic framework necessary to evaluate the region for relative potential for isolation of high-level radioactive waste. Because of the limited and specific goals of the project, emphasis is placed on the characteristics of the region that relate to waste isolation.

The results of this study are not based on original data; no new field work was conducted specifically for this project. It is not intended to be a definitive report on the geologic and hydrologic aspects of the region, but it provides a general summary of published and unpublished data that are available. In parts of the region, inadequate data exists to characterize the area. In these places it was necessary to discuss the geologic or hydrologic characteristics in the vicinity of the region, and then project that data into the region.

Geographic Setting

The Sonoran region, Arizona, is south and east of the Colorado River (fig. 1). The structural basins typically are filled with 200 to 1,500 m of clastic material with thick evaporite sections present locally. Bedrock pediments or shoulders of variable width are adjacent to the mountain blocks. Within the region, most basins trend north-northwest, except for the Butler and McMullen Valleys in the central part of the area that trend northeasterly. Regional drainage-outlet areas are through the south- and west-flowing Big Sandy, Santa Maria, and Bill Williams Rivers in the north and the Gila River system in the south; they collectively drain into the south-flowing lower Colorado River. Several basins have interior or poorly integrated drainages. Most basins are 150 to 300 m above sea level. Relief between valleys and adjacent mountains generally is 300 to 1,300 m. Bedrock probably is exposed in less than one-fourth of the total area. The location of geographic features is shown in figure 2.





Figure 1.--Physiographic features of the Sonoran region, Arizona and vicinity.



EXPLANATION

1. Ajo Range
2. Aquarius Mountains
3. Artillery Mountains
5. Baker Peaks
6. Black Mesa
7. Black Mountains
8. Buck Mountains
9. Buckskin Mountains
10. Cabeza Peak
11. Cabeza Prieta Mountains
12. Castaneda Hills
13. Castle Dome Mountains
14. Cerbat Mountains
15. Chocolate Mountains
16. Crater Range
17. Dome Rock Mountains
18. Eagletail Mountains
19. Gila Bend Mountains
20. Grand Wash Cliffs
21. Granite Wash Mountains
22. Growler Mountains
23. Harcuvar Mountains
24. Harquahala Mountains
25. Hualapai Mountains
26. Kofa Mountains
27. Little Harquahala Mountains
28. Martin Peak
30. Mohave Mountains
31. Mohawk Mountains
32. Mohon Mountains
33. New Water Mountains
34. North Camobabi Mountains
35. Plomosa Mountains
36. Quijotoa Mountains
37. Rawhide Mountains
38. "S" Mountain
39. Sauceda Mountains
40. South Camobabi Mountains
41. Vulture Mountains
42. Butler Valley
43. Dextrital Valley
44. Dutch Flat
45. Growler Wash
46. Hualapai Valley
47. McMullen Valley
48. Mohawk Valley
49. Ranegras Plain
50. Sacramento Valley
51. San Cristobal Wash
52. Valley of the Ajo
53. Big Sandy River
54. Bill Williams River
55. Burro Creek
56. Date Creek
57. Imperial Reservoir
58. Radium Hot Spring
59. Red Lake Playa
60. Salt River
61. Santa Maria River
62. Whipple Mountains
63. Mule Mountains
64. Baboquiviri Mountains
65. Big Maria Mountains

Figure 2.--Geographic index of the Sonoran region and vicinity.

Acknowledgments

This chapter and the other chapters in this series were prepared in cooperation with the States of Arizona, California, Idaho, Nevada, New Mexico, Texas, and Utah. Each of these States were represented by members of the Basin and Range Province Working Group. The cooperating agencies in each State and members and alternates of the Province Working Group are given following the title page. Alternate member of the Province Working Group, H. Wesley Peirce of Arizona, contributed significantly to the regional phase of the study. The following individuals provided continued advice and assistance to the Basin and Range Province Working Group and in overall planning and execution of the work in preparation of this series of chapters: John W. Hawley and William J. Stone of the New Mexico Bureau of Mines and Mineral Resources; Robert B. Scarborough of the Arizona Bureau of Geology and Mineral Technology; T.L.T. Grose of the Nevada Bureau of Mines and Geology and the Colorado School of Mines; George Dinwiddie and George I. Smith of the U.S. Geological Survey.

References Cited

- Bedinger, M. S., Sargent, K. A., and Brady, B. T., 1985, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part III, Geologic and hydrologic evaluation: U.S. Geological Survey Circular 904-C, 27 p.
- Bedinger, M. S., Sargent, K. A., and Langer, W. H., 1985, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Evaluation of the regions: U.S. Geological Survey Professional Paper 1370-H, [in press].
- Bedinger, M. S., Sargent, M. S., and others, 1985, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Basis of geohydrologic characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].
- Bedinger, M. S., Sargent, K. A., and Reed, J. E., 1984, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part I, Introduction and guidelines: U.S. Geological Survey Circular 904-A, 16 p.

Sargent, K. A., and Bedinger, M. S., 1985, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part II, Geologic and hydrologic characterization: U.S. Geological Survey Circular 904-B, 30 P.

GEOLOGY

By

Robert B. Scarborough,

Arizona Bureau of Geology and Mineral Technology

Stratigraphy

PRECAMBRIAN ROCKS

Precambrian rocks are exposed locally beneath Cenozoic volcanic rocks in many parts of the Sonoran region, and extensively in the Cerbat and Hualapai Mountains in the north. Precambrian rocks consist of plutonic rocks and metamorphic rocks of greenschist to amphibolite grade, 1.6 to 1.7 billion years old, and porphyritic granite, 1.4 billion years old (Wasserburg and Lanphere, 1965; Livingston, 1969; Anderson and Silver, 1971), all intruded by pegmatite and diabase dikes. Most crystalline terranes range from granodiorite to granite, with minor diorite. A nonpervasive regional Precambrian foliation ascribed to the Mazatzal Revolution (Wilson, 1939) strikes north to east-northeast and generally is steeply dipping. Sedimentary rocks of late Precambrian age that occur in adjacent regions, for example, the Apache Group to the east, the Grand Canyon Supergroup to the northeast, and the Pahrump Group to the northwest in California, are not recognized in the Sonoran region of Arizona.

Bedrock geology consists of Precambrian crystalline and metamorphic rocks overlain by a sequence of Paleozoic cratonic-shelf sedimentary rocks, Mesozoic volcanic rocks and clastic sedimentary rocks, and Oligocene and Miocene volcanic and sedimentary rocks. In all but the extreme northeast part of the Sonoran region, this assemblage of rock has been repeatedly deformed and moved along slight- and steep-angle faults, and intruded by Mesozoic and Cenozoic plutons. Regionally, most of the Cenozoic section and some crystalline rocks beneath have been displaced along northeast or southwest directions for unknown distances along middle Tertiary detachment faults. Reconstruction of major tectonic events is difficult because of extensive burial of all older rocks by upper Cenozoic basin-fill deposits.

Precambrian crystalline rocks exposed in the Hualapai and Cerbat Mountains are both foliated and unfoliated. Elsewhere, as in the northern Black Mountains and in various ranges of Yuma, La Paz, and Maricopa Counties, many of the crystalline rocks are gneissic. Metamorphosed crystalline terranes that contain variably foliated and mylonitized Proterozoic gneisses and granites lie beneath regional detachment fault in the Buckskin, Rawhide, and Harcuvar Mountains, and other ranges in the central part of the Sonoran region. Descriptions of Precambrian rock of the region are found in Gilluly (1946), Dings (1951), Thomas (1953), Kessler (1976), Anderson and Silver (1976), Stensrud and More (1980), and Howard and others (1982).

PALEOZOIC ROCKS

Small exposures of Paleozoic strata are known in some ranges of the region. These rocks, composed of limestone, dolomite, quartzite, and calcareous sandstone, have been described or mapped in the Plomosa, Harquahala, Little Harquahala, Rawhide, and the Black Mountains and south of Ajo, respectively, by Miller (1970), Reynolds and others (1980), Richard (1983), Shackelford (1976), and Wilson and others (1969). A comprehensive description of Paleozoic rocks in Arizona by Peirce (1976a) includes thicknesses in areas adjacent to the Sonoran region.

Paleozoic strata in the Sonoran region are correlated with:

(1) The Bolsa Quartzite and Abrigo Formation of Cambrian age, the Martin Formation of Devonian age, and Escabrosa Limestone of Mississippian age, which are cratonic in southeast Arizona, and (2) the Supai Formation of Pennsylvanian and Permian age, and Coconino Sandstone and Kaibab Limestone of Permian age, which are cratonic on the Colorado Plateau. Some workers correlate the Mississippian carbonate rocks with the Redwall Limestone of the Colorado Plateau. Only two complete and relatively unmetamorphosed Paleozoic sections are known in the Sonoran region southwest of the Colorado Plateau. These are in the southern Plomosa Mountains (Miller, 1970), and the Little Harquahala Mountains (Richard, 1983) as seen in figure 3. Virtually all other sections are tectonically attenuated and stratigraphically incomplete. Miller (1970) reports the following lithology and thickness: Bolsa Quartzite--maroon, wavy bedded quartz sandstone, 21 m; Abrigo Formation--dark sandy shale with minor quartz sandstone beds, 50 m; Martin Formation--dolomitic limestone, dolomite, and sandy dolomite, 110 m; Escabrosa Limestone--dolomite, limestone, and cherty limestone, fossiliferous, 136 m; Supai Formation--quartzite, limestone, sandy limestone, thin conglomerate, 180 m; Coconino Sandstone--white vitreous quartzite, eolian cross-bedding, 215 m; and the Kaibab Limestone--chert-bearing limestone of various colors and minor upper gypsum-anhydrite, fossiliferous, 220 m. Total




Figure 3.--Overturned Paleozoic section in the Martin Peak area of the Little Harquahala Mountains. View is to the northeast. Dark layered rocks on the right (A) are quartzite and calcareous quartzite correlated to the Supai Formation (Pennsylvanian and Permian). Light-colored rocks in the center (B) are correlated to the Redwall or Escabrosa Limestone of Mississippian age, and the darker units near the skyline on the left (C) may be Redwall, or Martin Formation of Devonian age. This section is near the southwest end of a major northeast-trending belt of Paleozoic rocks overturned to the southeast, that extends for more than 35 kilometers through the Little Harquahala and adjacent Harquahala Mountains. Photograph by Stephen J. Reynolds, Arizona Bureau of Geology and Mineral Technology, 1981.

aggregate thickness of 930 m is considered a minimum. As in other parts of Arizona, the Permian rocks are thicker than the rest of the Paleozoic combined (Peirce, 1976a).

Small, Paleozoic masses, some of which are tectonically encased within crystalline rocks, are found in the Rawhide and Buckskin Mountains (Shackelford, 1980), the Plomosa Mountains (Miller, 1970; Scarborough and Meader, 1983), the southern Grouler Mountains, the hills east of Parker, and at the extreme northern end of the Black Mountains (Wilson and others, 1969). Metamorphosed and deformed, but probably complete, Paleozoic sections are preserved beneath major thrust faults in the Harquahala and Granite Wash Mountains (Keith and others, 1982; Reynolds and others, 1980; Stephen J. Reynolds, Arizona Bureau of Geology and Mineral Technology, oral commun., 1984).

MESOZOIC ROCKS

Rocks of Mesozoic age in the Sonoran region of Arizona are represented by moderately sorted clastic sedimentary rocks with minor evaporites, volcanic rocks, Jurassic and Laramide plutons, and quartzose medium-grade metasedimentary rocks, all of which are variably metamorphosed. Jurassic volcanic rocks, part of the middle Mesozoic volcanic-arc complex of southwestern North America (Rogers and others, 1974; Coney, 1978) are recognized as intermediate to silicic flows and ash-flow tuff beneath Jurassic and Cretaceous sedimentary rocks in scattered localities in and around the Sonoran region (Damon and others, 1981, fig. 16). Jurassic granitic rocks are recognized in the northern Dome Rock Mountains (Reynolds, 1980) and in the Quijotoa, Camobabi, and Baboquivari Mountains (Haxel and others, 1980). Clastic and volcanic sequences, locally as thick as 7 km, are found in the Dome Rock, southern Plomosa, Buckskin, Granite Wash, Little Harquahala, and Harquahala Mountains (Robison, 1980; Reynolds, 1980; Harding, 1980). They are interpreted in part as having been deposited in a variety of terrestrial and lacustrine environments in a possible east-trending graben structure, and then deformed and overridden by thrusts by Late Jurassic time (Harding, 1982; Harding and others, 1983).

Laramide plutons occur throughout Arizona Basin and Range country and are the source rocks for the region's important porphyry copper deposits in the Cerbat Mountains of Mohave County at Mineral Park (Wilkinson and others, 1982), and at Ajo in Pima County (Dixon, 1966). Sizable Laramide plutons also are present in the Little Harquahala and Granite Wash Mountains, around Wickenburg, and as a batholithic mass underlying most of the region from Ajo to Yuma. This batholith, termed the Gunnery Range batholith, is dominated by leucocratic granite and has an area of about 15,000 km² in Arizona. It has K/Ar ages of 53 million years (Shafiqullah and others, 1980). Age dates on Laramide plutons in western Arizona generally cluster between 85 to 50 million years old.

A unit, probably correlative with the Orocopia Schist of Mesozoic(?) age of southern California, is recognized in three localities north of Yuma (Haxel and Dillon, 1978). The rocks are considered a regional allochthon composed of graywacke, chert, marble, and ultramafic rocks metamorphosed to greenschist or amphibolite grade in Late Cretaceous and Paleocene time. Their tectonic setting at this time is unknown. No Cretaceous volcanic rocks are recognized in the Sonoran region (Reynolds, 1980; Shafiqullah and others, 1980). However, many undated Mesozoic rocks are known in the region.

CENOZOIC ROCKS

Older Cenozoic rocks in western Arizona are characterized by a diverse assemblage of middle Tertiary volcanic rocks, presumed buried subvolcanic plutons, and volcanoclastic sedimentary rocks, 30 to 13 million years in age. This assemblage is unconformably overlain by locally thick basin-fill sediments and sparse intercalated basalt flows, formed in response to rift tectonics (younger than 13 million years) of the Basin and Range disturbance (Shafiqullah and others, 1980).

The middle Tertiary assemblage in the Sonoran region is characterized by a tripartite succession (Eberly and Stanley, 1978; Wilt and Scarborough, 1981) of: (1) Older fluvial and eolian redbeds with subordinate ash flows and tuff, commonly dominated by debris flows and mud flows; (2) an overlying locally thick volcanic sequence with ash flows, andesite, and pyroclastic sediments grading upwards into commonly less deformed basaltic andesite; and (3) a capping sequence of fluvial and lacustrine rocks that contain coarse redbeds and uraniumiferous organic-enriched lakebeds. The middle Tertiary volcanic-sedimentary succession may attain an aggregate thickness of as much as 2,000 m (Eberly and Stanley, 1978, fig. 8), but is non-uniform in thickness throughout the region. The upper part of the sequence contains significant low-grade uranium deposits in some middle Miocene carbonaceous lacustrine sediments in the Date Creek basin that is at the north end of the Harcuvar Mountains (Sherborne and others, 1979). Other nearby valleys also contain middle Miocene, uraniumiferous, fine-grained sediments buried under basin fill (Wilt and Scarborough, 1981).


The boundary between the middle Tertiary tripartite sequence (Eberly and Stanley, 1978) (Unit I of fig. 3) and the younger post-Basin and Range basin-fill sediments and basalts (Unit II of fig. 3) is a "... widespread unconformity surface resulting from an important period of subsidence, block-faulting, and erosion that began in Miocene time (13 to 12 m.y. ago)" (Eberly and Stanley, 1978). Basin-fill sediments are as much as 1,700 m thick where drilling has penetrated pre-basin-fill rocks (Eberly and Stanley, 1978). Peirce (1976b) suggests that as much as 3,300 m of basin fill occurs in the Hualapai Valley south of Red Lake, based on geometry of tilting of the Cerbat Mountains and gravity studies. The Bouse Formation is an assemblage of limestone, sandstone, and claystone deposited in a Pliocene estuary environment along most of the lower Colorado River.

Thick accumulations of middle Tertiary volcanic rocks that survived the subsequent Basin and Range block-faulting and that remain part of exposed range blocks are found in the Black, Aquarius, and Mohon Mountains of Mohave and Yavapai Counties, the Chocolate, Castle Dome, Kofa, New Water, Eagletail, and Gila Bend Mountains of La Paz and Yuma Counties, and the Crater Range and Saucedo Mountains of Maricopa County. Prominent late-stage, middle Tertiary, potassic, basaltic andesite fields are found surrounding the Mohon Mountains of Mohave County, in the Chocolate, Kofa, and Eagletail Mountains of Yuma County, in the Gila Bend Mountains of Maricopa County, and as a series of mesa-capping units east of Ajo, Pima County.

Basin and Range basalts, less than 13 million years old, are found as a few relatively thin caps on small mesas in the northern Black Mountains of Mohave County, a 130 km² mesa capping in northernmost La Paz County, the Sentinel volcanic field (600 km²) in western Maricopa County, and the northernmost part of the Pinacate volcanic field of Yuma County. The youngest basalts in the Sonoran region of Arizona are between 2 and 6 million years old in the Sentinel field; however, Pliocene and Pleistocene flows are found in the Pinacate field of Sonora and southernmost Yuma County, just south of the study area (Shafiqullah and others, 1980).

A thick sequence of predominantly halite in the Sonoran region has been penetrated in the subsurface south of Red Lake playa in Hualapai Valley (Peirce, 1976b). Geophysical modeling indicates a mass with a width of 8 km, length of 18 km, and bottom depth of 3.2 km (1,220 m of which has been drilled). Possible volume is 250 to 500 km³. The salt probably represents closed-basin accumulation under nonmarine conditions during the Miocene (Peirce, 1976b).

Basin-fill geology is a major factor in characterization of regional hydrology. Sand and gravel facies within basin fill are the most permeable of all near-surface units in the Basin and Range province, and hence their spatial distribution is important in formulating ground-water travel times. A hypothetical geologic section that indicates most facies relationships recognized in a typical southern Arizona basin is given in figure 4. Basin and Range faults that commonly are transmissive near the surface may be much less permeable at depth because of overburden pressure and silicification or alteration of fault zones related to fossil-geothermal systems.



EXPLANATION

GENERAL DESCRIPTION OF ROCK UNITS

1. Pleistocene piedmont alluvial fans, capping a beveled pediment that was eroded into basin-fill sediments. Fans thicken and contain coarser debris mountainward.
2. Late Pleistocene and Holocene strath-terrace gravel, sand, and overband silt, related to erosion-backfill cycles of the valley's trunk stream. These landforms and deposits better developed in valleys nearer the Colorado and Gila Rivers.
3. Basin fill, generally interpreted as post-13 million years in age, and assigned as Unit II by Eberly and Stanley (1978).
 - 3a. Proximal alluvial fan facies (poorly sorted sand and gravel)
 - 3b. Distal-fan facies (gravelly sand, sand, silt, and air-fall ash)
 - 3c. Lacustrine facies (bedded clay, silt, gypsum stringers, air-fall ash, and diatomites as pods)
 - 3d. Evaporite facies (halite, anhydrite, and clay stringers)
 - 3e. Stripped pediment exposed at the mountain fronts within those valleys undergoing stream erosion. Basin-fill deposits thin and coarsen mountainward, and rest on the pediment that is eroded into older Tertiary deposits and bedrock.
4. Mid-Tertiary volcanic rocks, and sediments, generally 28 to 13 million years in age, and assigned as Unit I by Eberly and Stanley (1978). Individual units generally lack lateral continuity. Volcanic rocks consist of welded and unwelded ash-flows, andesite and basalt flows, agglomerate, tuff, and intrusive domes. Sediments are heterogeneous assemblage of redbeds, fluvial sand, lacustrine clay, silt, and limestone.
5. Pre-Cenozoic bedrock, including Precambrian crystalline and metamorphic rocks, Paleozoic limestone and sandstone, and Mesozoic volcanic rocks, volcanoclastic rocks, redbeds, and sandstone.

CONTACT

FAULT.---Dashed where approximately located. B, Basin and Range faults, steep-angle normal, active during block faulting; L, listric (curvilinear) faults, active during denudational tectonics

Figure 4.--Hypothetical geologic section transverse to main drainage across a typical basin.

Summary of Geologic Events

PRECAMBRIAN TECTONICS

The Precambrian basement that extends throughout the Sonoran section records an older Precambrian (1.7 to 1.6 billion years ago) island-arc volcanic and submarine sedimentary terrane (including banded iron formations, Gilmour and Still, 1968), that was intruded by a series of granite and subjected to a non-pervasive metamorphism (Wilson, 1939; Anderson, 1951; Anderson and others, 1971; Conway, 1976; Shafiqullah and others, 1980). Regional schistosity produced by this metamorphism generally is northeast trending, and central-Arizona fold belts and ore-deposit clusters trend northeast as well. This orogeny, termed the Mazatzal Revolution by Wilson (1939), is represented by the Bamori, Vishnu Schist, Yavapai Schist, and Pinal Schist terranes throughout the State, and is now attributed to regional accretionary tectonics by Shafiqullah and others (1980). Attendant greenstone belts and volcanogenic massive sulfide deposits in central Arizona are discussed by Stensrud and More (1980), Donnelly and Hahn (1981), and Anderson and Silver (1976).

This metamorphosed terrane was intruded by regionally extensive batholiths dated at 1.4 to 1.45 billion years (Silver, 1968). Plutonic rocks of the batholiths are composed of moderately alkalic granite and quartz monzonite characterized by the presence of rapakivi-type orthoclase phenocrysts ("megacrysts"), and, in one spot in the Aquarius Mountains, pods of high-grade radioactive carbonatite material. This intrusive episode is termed "anorogenic" in reference to the apparent absence of related volcanic rocks or evidence of associated deformation (Silver and others, 1977; Anderson, 1983).

Based on data from areas north and east of the Sonoran region, the region was eroded to low relief after the 1.4-billion-year granite metamorphism and younger Precambrian sedimentary rocks of the Apache Group, Grand Canyon Supergroup, and Pahrump Group were deposited in a nearshore sedimentary environment, perhaps related to a North America-Siberia continental rifting (Stewart, 1976a, 1976b; Sears and Price, 1978; Condie, 1982). Main lithologies of the Apache Group include quartz sandstone, tuffaceous siltstone, maroon shale, black carbonaceous shale, and minor conglomerate. Late-stage, rift-related sills and dikes of diabase and minor syenite intruded the sedimentary rocks throughout central Arizona at about 1,100 to 1,200 million years ago (Silver, 1960; Livingston, 1969). Sedimentation occurred between 1,420 and 1,100 million years ago. Some diabase dikes, presumably of this age, intrude the crystalline terrane of the Hualapai Mountains (Kessler, 1976) and the Mohave Mountains (Nakata, 1982).

PHANEROZOIC TECTONICS AND VOLCANISM

Paleozoic

The Arizona Paleozoic cratonic assemblage is hypothesized to have been deposited throughout the Sonoran region, although most Paleozoic rock occurrences there are now mostly limited to a few sparse generally deformed, ductilely attenuated masses. Peirce (1976a) and Schumacher and others (1976) suggest that Paleozoic crustal movements in Arizona primarily were epeirogenic, consisting of mild tilting induced by broad regional arching and sagging. Stone and others (1983) conclude that the cratonic Paleozoic section appears to have been deposited throughout the Sonoran region westward to the central Mojave Desert in California about 300 km west of the region.

Mesozoic and Cenozoic denudation

The Sonoran region is characterized by Oligocene and younger volcanic and sedimentary rocks deposited on Precambrian and lower Tertiary crystalline and metamorphic basement and local areas of Paleozoic through Eocene strata. On the adjacent Colorado Plateau, a northeast-dipping basal Tertiary erosion surface that has been incised progressively downward through the Paleozoic section to the southwest, is capped with southwest-derived Tertiary gravels, the "rim gravels", along the plateau edge. Farther east along the plateau edge, a northeast-sloping Cretaceous (pre-Turonian) surface also truncates progressively older Paleozoic strata to the southwest. These conditions indicate that the part of the Basin and Range province bordering the plateau was topographically higher than the plateau and was largely stripped of its Mesozoic and Paleozoic sedimentary cover by later Mesozoic and early Tertiary erosion that preceded the intense middle Tertiary volcanic pulse (Peirce and others, 1979). This central Arizona Mesozoic highland was termed the Mogollon Highland by Harshbarger and others (1957) and Cooley and Davidson (1963). It evidently was a southwest source for Mesozoic and lower Tertiary sediments on the southern Colorado Plateau. The Mogollon Highland then was lowered in altitude relative to the plateau as a consequence of some combination of further erosion and tectonic effects, mostly during Eocene and Oligocene(?) time, with subsequent regional drainage disruption and creation of new Basin and Range drainage trends. This newly created, post "rim gravel" erosion cycle and the earlier Mesozoic erosion of the Highland are jointly responsible for the removal of nearly all

the Phanerozoic cover from the area extending some unknown distance southwest of the plateau edge. Cretaceous erosion largely preceded Laramide tectonism and magmatism in southern Arizona; the middle Tertiary loss of altitude was contemporaneous with, and probably the consequence of, major crustal extension and associated volcanism in the Basin and Range province.

The modern Mogollon Rim, the southwestern edge of the Colorado Plateau, was erosionally carved out in crystalline rocks by the newly created Oligocene(?) Basin and Range drainages as they eroded northeastward in an area of southwestward pinchout of resistant Paleozoic and Jurassic strata under the Cretaceous surface (Young and Brennan, 1974; Peirce and others, 1979; Shafiqullah and others, 1980).

It is not known how far southwest this erosion occurred; the complex tectonics southwest of the metamorphic-core complexes disturbs the record so that it is not now clear concerning the relative effects in that region carried by erosion and tectonics.

Pre-Laramide Mesozoic tectonics

The lowermost Mesozoic deposits known in the Sonoran region are middle Jurassic felsic volcanic rocks found in a zone 200 km wide, extending southwest from Parker. These are typically overlain by a series of locally thick Jurassic or Cretaceous clastic sedimentary rocks or both. One of the more regionally recognized sequences, the McCoy Mountains Formation, has been described as a west-northwest-trending graben (Harding, 1982).

Regional thrust-faulting occurred after the deposition of Jurassic and Cretaceous volcanic and sedimentary rocks, and then the intrusion of some early Laramide (85 million years) plutons occurred. Rocks juxtaposed by thrust faults are found from the Rawhide Mountains southward to near the Yuma area and southeastward, east of Ajo. These thrust sequences consist of stacked tectonic plates of Precambrian metamorphic and plutonic rocks, Paleozoic sedimentary rocks, and Mesozoic sedimentary and volcanic rocks. The bounding faults are at times nearly flat, as in the Harquahala Mountains (Reynolds and others, 1980) and in the Little Harquahala Mountains (Richard, 1983), shown in figure 5, and at times moderately to steeply dipping, as in the southern Plomosa Mountains (Miller, 1970) and the northern Dome Rock Mountains (Harding, 1982). Plates composed of crystalline rocks appear to have had little internal deformation in comparison to structurally confined plates composed of Paleozoic and Mesozoic rocks that commonly contain tight to isoclinal folds, and in some places recumbent folds, that indicate attenuation and extensive ductile deformation. Kinematic indicators of this thrust deformation in the Sonoran region of Arizona indicate an older southeast-vergent transport and a younger south-southwest or north-northeast transport (Reynolds and others, 1980; Stephen J. Reynolds, oral commun., 1984). Shackelford (1976) indicates the presence of probable Cretaceous northeast-thrusting, thermal metamorphism, and penetrative deformation of Precambrian and Paleozoic rocks in the Rawhide




Figure 5.--Hercules thrust fault, exposed near "S" Mountain, southeast of Salome, La Paz County. The ledge-forming unit consists of quartzite and stretched pebble conglomerate of Mesozoic age, correlated with the McCoy Mountains Formation (Harding, 1982). The unit is capped by the slightly dipping Hercules thrust, and overlain by sheets of light-colored leucocratic granite and gray-colored quartz diorite of Precambrian or Mesozoic age. Beneath the ledge-forming units is quartz-feldspar schist derived from Mesozoic volcanic rocks and volcanoclastic sediments. The Hercules thrust fault has been mapped in the Harquahala, Little Harquahala, and Granite Wash Mountains. This thrust terrane is intruded elsewhere by undeformed granite (85 million years old) of the Tank Pass pluton. Photograph by Stephen J. Reynolds, Arizona Bureau of Geology and Mineral Technology, 1981.

Mountains. Hamilton (1982) suggests that a pre-80- to 90-million-year-old northeast overfolding with attendant metamorphism affected the rocks of the Big Maria Mountains in nearby southeastern California. Miller and McKee (1971) noted west-vergent thrusts in the southern Plomosa Mountains that juxtaposed a complex array of Precambrian, Paleozoic, and Mesozoic rocks in an hourglass-shaped, east-dipping thrust pile. Directly to the east is a nonmetamorphosed terrane that contains one of the two relatively complete fossiliferous Paleozoic sections in western Arizona. The other relatively complete Paleozoic section is illustrated in figure 3. Scarborough and Meader (1983) mapped a thrust stack in the northern Plomosa Mountains that records a ductile, knappe-like deformation that produced an attenuated, imbricately faulted, south-dipping Paleozoic and Mesozoic plate overlain and underlain by Precambrian crystalline plates. East- or northeast-vergent minor folds occur in the plate. The plate is intruded by a post-kinematic pluton dated at 85 million years by K/Ar (biotite). Outcrops of Mesozoic(?) Orocochia Schist mapped by Haxel and Dillon (1978) north of Yuma, may be exposed in windows through a regionally extensive thrust plate that may include all of the southern part of the Sonoran region.

This late Mesozoic thrust terrane extends from at least as far south as the southern Dome Rock Mountains, and perhaps southward to near Yuma, northward to the Rawhide and Buckskin Mountains, westward into the eastern Mojave Desert, and eastward at least to the Harquahala Mountains. The terrane is thought by some to be a continuation of the Sevier orogenic belt of western Utah and southern Nevada (Armstrong, 1968); but unlike the Sevier belt, thrusts involve surficial crystalline rocks and ductilely attenuated Phanerozoic sedimentary rocks which are both southeast and northwest vergent (Burchfiel and Davis, 1975; 1981; Dickinson and Snyder, 1978; Carr, 1983).

The east-trending belt containing the 4-km-thick Jurassic(?) McCoy Mountains Formation trends through the Dome Rock and southern Plomosa Mountains of the Sonoran region of Arizona (Harding, 1982). This belt has been deformed into an oppositely dipping pair of folds partly cored with McCoy Mountains sedimentary rocks that are surrounded by Jurassic volcanics rocks. Moderate north- and south-dipping faults juxtapose this belt into a lower-plate position with respect to the terranes to the north and south that do not contain thick Mesozoic clastics. Harding (1982) calls the northern terrane, Cratonic North America; the central terrane, the McCoy terrane; and the southern region, the Mojave-Sonora composite terrane. The deformation of the belt is suggested to have taken place in Jurassic time, based on paleomagnetic evidence (Harding and others, 1983). Preliminary structural analysis from the northern fault complex, the Boyer Gap thrust, indicates south-vergent movement, placing cratonic rocks over McCoy terrane rocks (Yeats, 1984). Tosdal (1982) notes a possible continuation of the southernmost of the two bounding faults, the Mule Mountains thrust, from the Mule Mountains of California to the southern Dome Rock Mountains of Arizona, based on similar juxtapositions along a south-dipping reverse fault.

Possible pre-Sevier strike-slip faulting is hypothesized by Silver and Anderson (1974) and Anderson and Silver (1981). This fault or series of faults is termed the Mojave-Sonora megashear, and is interpreted to have a northwest trend. About 800 km of left-lateral offset has been indicated based on: (1) An offset pattern of radiometrically dated Precambrian rocks in Arizona, California, and Sonora, Mexico; (2) plate-tectonic modeling of the opening of the Caribbean Sea (Anderson and Schmidt, 1983); and (3) Mesozoic pinchout patterns in Sonora. The zone has been drawn through an area just west of Ajo and the lower Colorado River near Ehrenberg, Yuma County. If it exists, dissimilar Precambrian through Middle Jurassic rocks should be juxtaposed across the line.

Laramide orogeny

The Laramide orogeny in southern and western Arizona is represented by a period of intense subduction-related magmatism and volcanism, significant copper-molybdenum-lead-zinc-silver-gold metallogenesis, and late-stage northeast-directed compressional tectonics (Coney, 1976 and 1978; Keith and Barrett, 1976; Drewes, 1978; Damon and others, 1981; Titley, 1981 and 1982; and Keith, 1982).

Laramide plutons with related porphyry-copper mineralization appear to have been emplaced into structurally high crust, and localized along structural intersections related to Precambrian fabric (Titley, 1981). Some evidence exists to indicate that Laramide lead-zinc-silver vein-type mineralization and copper-molybdenum porphyry-type deposits were associated with late Mesozoic shallowing subduction (Coney and Reynolds, 1977; Keith, 1978 and 1981). Some workers now believe that a subsequent Eocene magmatic-tectonic event occurred that was related to, or possibly the culmination of, the Laramide orogeny because of a direct relationship between magmatic-metallogenic space-time patterns and inferred paleosubduction geometries (Keith, 1978, 1981, and 1982; Shafiqullah and others, 1980). The possibility has been suggested (Keith, 1981) that the subduction zone reached highest crustal levels (slightest dip angles) during the Eocene at which time crustal anatexis and high-level shearing produced plutonized tectonite basement that eventually became locally arched, and formed the core of the exposed southern Arizona metamorphic-core complexes. Alternatively, some workers view the ductile basement shearing as due to regional distension during the middle Tertiary, unrelated to the Laramide (Reynolds, 1980).

Laramide dikes in the Sonoran region strike generally northeast to east, indicative of an overall regime of northeast compression (Rehrig and Heidrick, 1976). Andesite dikes at Ajo strike nearly eastward. In the Vulture Mountains, dikes probably related to the 68 million-year-old Wickenburg batholith, itself elongate in a northeastern direction, generally trend northeast as well. At Bagdad, dioritic dikes that transect the 71 million-year-old mineralizing stock trend east-northeast.



Metamorphic-core complexes and middle Tertiary detachments

In the Sonoran region, metamorphic core complexes, as defined by Rehrig and Reynolds (1980), Davis (1979), and Davis and others (1980), are found in the Buckskin, Rawhide, and Harcuvar Mountains, near the western terminus of a northwest-trending zone, which is 600 km long in Arizona and California, that contains a number of such complexes. The core-complex rocks contain a distinct rock assemblage, and are separated from "normal" geology by a slight-angle fault (Coney, 1980, fig. 2). Beneath the fault, the lower-plate assemblage consists of mixed crystalline rocks subjected to Late Cretaceous(?) to middle Tertiary mylonitization, heating, and deformation by ductile shearing within a subhorizontal shear couple. The intensity of lower-plate mylonitization increases upward through a thin zone that is characterized by a chloritized breccia and capped by a ledge-forming microbreccia and overlying fault surface, commonly referred to as a detachment fault. The mylonitic zone usually continues structurally downward through the lower-plate crystalline rocks for some fraction of a kilometer. The mylonitic zone contains a mineral lineation in the foliation plane that is interpreted as being parallel to and produced by upper-plate transport during mylonitization. This lineation direction is usually at right angles to the strike direction of upper-plate tilted Cenozoic strata, an observation that leads some workers (Rehrig and Reynolds, 1980; Rehrig and others, 1980) to infer that mylonitization and detachment represent ductile and brittle aspects, respectively, of the same phenomenon, namely a crustal-scale shear zone that accommodated

major extension across the Basin and Range province.

Detached (upper-plate) rocks structurally above the core complexes range in age from Precambrian to Miocene. Upper-plate Cenozoic stratified rocks near the core complexes usually consist of Oligocene sequences and younger early to middle Miocene strata that typically have a consistent regional direction of strike (northwest) and dip (northeast or southwest). The Miocene rocks usually are less deformed than the older upper-plate rocks, indicating episodic deformation. Regionally, upper-plate volcanic rocks as young as 15 million years were involved in rotation and detachment faulting (Scarborough and Wilt, 1979; Rehrig, 1982; Rehrig and others, 1980; Suneson and Lucchitta, 1979). The thermal history of the lower plate is typically one of thermal changing of radiometric ages to early or middle Cenozoic age for all mylonitized rocks. Upper-plate rocks have radiometric ages that have not been affected by Tertiary thermal effects, unlike lower-plate rocks. This history indicates that the mylonitic zone has been subjected to a substantial thermal flux. A hypothesized geologic section through the northeast end of the Harcuvar metamorphic-core complex (fig. 6) and shows the lower- and upper-plate relationship and the detachment fault. The vertical exaggeration of 10 times makes the fault, which dips 20°, appear nearly vertical. This geologic section is a part of one of 32 constructed for the Sonoran Desert region for this study, on open file at the Arizona Bureau of Geology and Mineral Technology.

SW
MILLIGALS
60
0
-60
Complete residual Bouguer gravity anomaly profile
from Lysonski and others (1981)



0 10 20 30 40 50 60 60 KILOMETERS
VERTICAL EXAGGERATION X 10

	EXPLANATION
QTS	QUATERNARY AND TERTIARY SEDIMENTS.--Mostly post-13 million years in age
Tb	OLIGOCENE AND MIOCENE BASALTIC ROCKS
Tv	OLIGOCENE AND MIOCENE VOLCANIC ROCKS.--Ash flows, tuff, and andesite
Tvs	OLIGOCENE AND MIOCENE VOLCANIC AND SEDIMENTARY ROCKS
KTM	MYLONITIC CRYSTALLINE ROCKS OF THE METAMORPHIC CORE COMPLEXES
Ki	CRETACEOUS CRYSTALLINE ROCKS.--Quartz monzonite, granodiorite, and so forth
Kpcc	CRETACEOUS AND PRECAMBRIAN CRYSTALLINE ROCKS
Mzs	MESOZOIC CLASTIC SEDIMENTS.--Mostly sandstones; in places with minor andesite flows
Mzm	METAMORPHOSED MESOZOIC CLASTIC SEDIMENT.--Phyllites, argillites, and so forth
Pzs	PALEOZOIC LIMESTONE, DOLOMITE, AND QUARTZITE
Mzpz	MESOZOIC AND PALEOZOIC SEDIMENTARY AND METAMORPHIC ROCKS
pCg	PRECAMBRIAN GRANITIC ROCKS
pCm	PRECAMBRIAN METAMORPHIC ROCKS

The section crosses two detachment faults that separate lower plate thrust-juxtaposed rocks or mylonitic crystalline rocks with post-kinematic plutons from upper-plate allochthonous, but thermally intact, Precambrian crystalline and Tertiary volcanic rocks. Northeast transport of upper-plate rocks is indicated by antithetic southwest rotation of upper-plate volcanic rocks throughout the region, as seen in the section, and hence, a post-detachment arching, of the detachment faults and some mountain ranges also is indicated. It is not known whether the master detachments continue to dip at moderate angles to great depths, or if they are simply flat faults, now warped so that they intersect the land surface at different places.

The complete residual Bouguer gravity profile above the section, from Lysonski and others (1981), indicates a .10 mGal negative anomaly in the Ranegras Plain and a nearly 30-mGal negative anomaly in this part of the Date Creek basin, indicative of the depths to bedrock shown. Dense evaporite facies in basin fill and thick, dense, buried Tertiary volcanic rocks make interpretation of depth to crystalline bedrock difficult without subsurface control.

Detachment-fault outcrops and lower-plate mylonitic fabrics together define two sets of large-scale, fold-like irregularities in the Sonoran region. In the Harcuvar, Buckskin, Rawhide and possibly the Harquahala Mountains, one of these fold-sets is well defined and has axes trending east-northeast. Synforms and antiforms are represented respectively by sediment-filled valleys and arch-like ranges. The sinusoidal, wave-like form of the detachment fault and mylonitic fabrics is readily discernible on geologic maps of the area (Rehrig and Reynolds, 1980). The arches are thought to be somehow related to mylonitization and detachment faulting because their axes are parallel to the movement direction on the detachment fault and to mineral lineation in the mylonites, but their precise origin is unknown.

A second, broader warp has a northwest-trending axis that is orthogonal to the east-northeast arches and perpendicular to detachment-fault movement direction. It appears to have formed as a result of isostatic rebound after tectonic denudation (Rehrig and Reynolds, 1980; Spencer, 1984). The plunging axes at the ends of some of the arches in the Harcuvar complex reflect this warp, as might well the overall regional northwest trend of the Arizona-southeast California line of core complexes. In the Lincoln Ranch basin area of the Buckskin Mountains, a northwest-trending, southwest-vergent reverse fault places Precambrian gneisses over middle Miocene redbeds (Wilson, 1962, pl. 24). It is possible to relate this thrusting to the formation of the northwest arch or warp.

Rotated Cenozoic strata also are found well away from the core complexes throughout all of western Arizona and eastern California, and consistently strike northwest and dip either southwest or northeast. Tilted volcanic strata in the Cabeza Prieta Mountains are shown in figure 7. It is now assumed, but not proven, that these terranes were subjected to detachment movement contemporaneously with movement of upper-plate rocks around the core complexes.



Figure 7.--Southern Cabeza Prieta Mountains (Yuma County)

looking north-northeast nearly parallel to the strike of tilted dark-colored, bedded volcanic rocks and pyroclastic sediments. Underlying rocks include Mesozoic(?) schist and granite of the 53-million-year-old Gunnery Range batholith. The uniform 25° northeast dip on the strata may be explained by southwest transport on a series of northwest-striking, listric, normal faults that affected the volcanic rocks and underlying crystalline bedrock. The high point on the skyline is Cabeza Peak (A). Beyond the mountains lies Mohawk Valley. View is taken from about 1.6 kilometers south of Cabeza Peak (B). Photograph by Peter Kresan, University of Arizona, 1980.

Strata with uniform northeast or southwest dip direction, as seen in figure 6, can be grouped into elongate belts that extend through western Arizona in a northwest direction parallel to the Colorado Plateau margin (Rehrig and Heidrick, 1976; Scarborough and Wilt, 1979; Rehrig and others, 1980; Stewart, 1980). The belts, 100 km or more wide, are interpreted by Rehrig and others (1980) as brittle upper-plate expansive regimes between narrow northwest-trending zones (belt boundaries) of concentrated lower-plate ductile expansion and magmatic ingress into high crustal levels. Similarly Stewart (1980) suggests the belts are related to stress relief extending outward from boundaries of zones that are interpreted as initial sites of rupture during late Cenozoic extension. Klein (1982) suggests the existence of a zone, 30 km wide, containing a positive residual aeromagnetic anomaly that parallels the tilt domain boundaries that extend through the Parker area. He suggests a crustal, magnetic-source region at least 6 km below the land surface, and notes that the magnetic zone separates the terrane to the northeast with a relatively positive residual-gravity signature from the terrane to the southwest with a negative residual gravity. Perhaps this magnetic anomaly signifies the presence of one of these zones of concentrated lower-plate ductile expansion of Rehrig and others (1980).

The depths to which the basal detachment faults extend are not known. In some areas, such as the detached terrane including the Whipple Mountains of east-central California, available evidence based on reconstructing upper-plate thicknesses indicates that detachment faults dipped gently to depths of 10 to 15 km at the time of formation (Howard and others, 1982). Wernicke (1981) has proposed that basalt detachments continue at moderate dip angles through the entire lithosphere. It is not known how many different detached terranes are present in the Sonoran region. Because there are at least four different dip domains present in the Sonoran region (Scarborough and Wilt, 1979, fig. 28), each presumably having moved in response to distention along an individual master detachment fault, there may be several, or numerous large-scale, low-angle faults in the region.

The Gila trough is adjacent to and parallels an anomalous east-northeast regional photolinear trend occupied by reaches of the Salt and lower Gila Rivers that extends from Yuma through Phoenix and on to at least as far as the Mazatzal Mountains (80 km northeast of Phoenix). The origin of the trough between Yuma and Phoenix is unknown although it appears to be an elongate series of depocenters partly filled with relatively thick middle Tertiary sedimentary and volcanic rocks (Eberly and Stanley, 1978), and hence is inferred to have been forming during that time. Its form has been described as a graben or half-graben (Eberly and Stanley, 1978), and as a synform or downwarp (Pridmore and Craig, 1982). Along the southern flank of the trough, detached middle Tertiary redbeds at Baker Peaks, that probably were deposited in the trough, contain indications of northward sediment transport from the volcanic-free Gunnery Range batholith terrane to the south. These rocks have a northeast trend of detachment movement as indicated by their general southwest dips, a trend that approximately parallels the long axis of the trough (Pridmore and Craig, 1982). The east-northeast trend of the trough and the trend of upper-plate detachment movement along its flanks appears geometrically analogous to that of the line of northeast-trending, arched-core complexes and intervening synformal valleys in the Buckskin-Harquahala region to the north, that also have indicated northeast tectonic transport (southwest dips) of upper-plate rocks.

Miller and McKee (1971) discuss a west-northwest-trending zone of strike-slip faults in the southern Plomosa Mountains that disrupt rocks as young as some middle Tertiary volcanic rocks that contain a flow 20 million years old. Apparent left-lateral separation of 6 km is dispersed across a zone 8 km wide that contains at least 3 large-offset faults. It is not clear whether this faulting is related more to detachment faulting or to later Basin and Range pull-apart rhombochasm basin development (Davis, 1980; Anderson and others, 1972). Curiously, this strike-slip trend is parallel to the strike of upper-plate rotated strata in the immediate region that have been deformed by west northwest-striking, listric, normal faults of about the same age.

Middle Tertiary volcanism

The middle Tertiary volcanic episode in the Sonoran region is believed as being subduction related (Coney and Reynolds, 1977). It consisted of earlier andesite-rhyodacite, ash-flow tuff volcanism and penecontemporaneous sedimentation in local basins, then by a voluminous mass of dacite-latitude-rhyodacite ash-flow tuff, and finally a series dominated by potassic basaltic andesite with local ultrapotassic trachyte (Shafiqullah and others, 1980). Later andesite and ash flows of the Growler Mountains are illustrated in figure 8. Local variations of this sequence are known, such pre-ash flow basaltic volcanism in the Castaneda Hills (Suneson and Lucchitta, 1983). The middle Tertiary rock suites in the Sonoran region range in age from about 28 to 15 million years, at which time an abrupt change to basaltic volcanism took place (Eberly and Stanley, 1978; Shafiqullah and others, 1980; Suneson and Lucchitta, 1983). Local volcanic-tectonic processes consisted of differential block subsidence with local landsliding and sedimentation as noted in the southern Black Mountains (Thorson, 1971) and in the Mohave Mountains (Pike and Hansen, 1982). Pike and Hansen (1982) suggest that most volcanic and sedimentary rocks lack lateral continuity because of numerous unconformities produced by coeval volcanic-tectonic processes. Thick successions of ash-flow tuff and concentrated faulting, possible indicators of calderas, are found in the southern Black mountains (Thorson, 1971), and in the



Figure 8.--View northwestward towards the southern Growler Mountains (A) of western Pima County in the distance, which are capped by dark-colored, slightly eastward dipping middle Tertiary volcanic strata. All dark-colored rocks are Oligocene and Miocene andesite, basalt, and some ash flows. Basement rock of Mesozoic(?) granite is exposed in the small irregularly shaped gray-colored hill (B). Growler Wash (C) drains nearly 700 square kilometers of the Valley of the Ajo, out of view to the right. This drainage joins San Cristobal Wash downstream, and together they empty into the Gila River at the Mohawk Mountains 100 kilometers downstream. Most of the scene is in T14S, R7W. Photograph by Peter Kresan, University of Arizona, 1981.

Kofa and Castle Dome Mountains, and the Ajo Range. Detailed studies in the latter three areas are lacking.

As summarized in Zoback and others (1981), the beginning time of middle Tertiary volcanism progressed westward from New Mexico (40 million years ago) to east-central California (25 million years ago) in keeping with the post-Eocene steepening Benioff-zone concept of Coney and Reynolds (1977). Middle Tertiary volcanism lasted about 10 to 14 million years in any given area.

Middle Tertiary dikes in the Sonoran region generally trend northwest and are nearly orthogonal to Laramide dikes in the region (Rehrig and Heidrick, 1976; Nakata, 1982; Howard and others, 1982). Quartz-monzonite-porphyry dikes at Bagdad strike north-northwest. In the Vulture Mountains, extensive rhyodacite-to latite-porphyry dikes, 15 to 18 million years old, trend north-northwest. The Tertiary-dike trend in both the Castle Dome and Mohave Mountains is northwest. In the southern Black Mountains, rhyolite-porphyry dikes and associated banded quartz-calcite epithermal gold-silver veins trend northwest to west-northwest. The general north-northwest dike trend in middle Tertiary time (25 to 15 million years ago) generally is thought to be perpendicular to the direction of tectonic extension during the middle Tertiary magmatic episode.

Basin-and-range faulting and volcanism

Basin-and-range faulting, as characterized by Eberly and Stanley (1978) and Zoback and others (1981), has affected the southwestern one-half of Arizona. Basin and Range mountain blocks in the Sonoran region trend north to north-northwest. In contrast, the Buckskin, Rawhide, Harquahala, and Harcuvar Mountains of the Sonoran region trend east-northeast, but appear to be a manifestation of undulatory northeast-trending Tertiary (Miocene?) arching that has profoundly affected the form and surface distribution of mylonitic rock of the metamorphic-core complexes. Hence, these ranges with their different trend are not believed to be Basin and Range fault blocks.

Most of the steep-angle, non-rotational horst-and-graben faulting generally has been bracketed in time as younger than 13 million years and older than 5 million years throughout most of the southern part of the Basin and Range province (Stewart, 1971; Anderson and others, 1972; Scarborough and Peirce, 1978; Eberly and Stanley, 1978; Zoback and others, 1981). Where investigated by seismic methods and drilling, most basins morphologically resemble non-rotated, wedge-shaped sags downdropped along one or more steep, planar, normal faults, or prism-shapes above tilted-bedrock ramps that are displaced by moderately to deeply penetrating listric(?) normal faults (Stewart, 1978; Eberly and Stanley, 1978; Zoback and others, 1981; Anderson and others, 1983), and filled with from 600 to perhaps 1,500 m of basin-fill sediments (fig. 4). However, it is not clear to what degree the present-day structural relief between range tops and adjacent basin bottoms is the result of steep-angle block faulting or of other events such as local isostatic rebound of denuded terrane after detachment faulting (Spencer, 1984). New evidence indicates that many western Arizona range blocks, including some outside the line of metamorphic-core complexes, are bounded just at the mountain fronts by detachment faults that now appear to have steeper dips than at the time of detachment, indicating post-detachment relative uplift or arching of the range blocks. The northeast-trending arches that affected the metamorphic-core complexes of the Sonoran region produced minimum, vertical structural relief on the mylonites of 1,200 m, which is a significant fraction of that postulated for southeast Arizona range top to basin-bottom relief, 1,800 to 3,300 m (Scarborough and Peirce, 1978).

Depth to bedrock within the basins has been modeled using the residual Bouguer-gravity anomalies by Oppenheimer and Sumner (1980). Peirce (1976b) suggests the presence of 3,000 m of basin fill, including a significant fraction of halite thickness, for Hualapai Valley south of Red Lake (fig. 9). Large residual anomalies that indicate depth to bedrock possibly as great as 2,500 m (Scarborough and others, 1983) are found in the valley west of the Cerbat Mountains, west of the southern Black Mountains, east of the Mohave and Buck Mountains, east of the southern Castle Dome Mountains, and between Baker Peaks and the Mohawk Mountains, as shown in figure 10.



Figure 9.--Red Lake playa (A) of the Hualapai Valley in Mohave County. View is to the northeast towards the Grand Wash Cliffs (B), composed of Precambrian crystalline rocks overlain by Paleozoic strata as young as Devonian. In the low foreground are hills at the base of the Cerbat Mountains (C). The farthest skyline, just visible, consists of cliffs of Permian sandstone and limestone that form the Mogollon Rim (D), the physiographic edge of the Colorado Plateau. Although Red Lake retains water for short intervals after storms, no Pleistocene pluvial record of the lake is known. A large body of halite (at least 1,200 meters thick, and possibly 150 cubic kilometers in volume) is known in the subsurface south (to the right) of the playa. Throw on the Grand Wash fault, parallel to the valley edge on the far side of the playa, may be as great as 4,900 meters. Photograph by Peter Kresan, University of Arizona, 1980.




Figure 10.--Deep basins and areas of Pleistocene and Holocene erosion.

The northern block of the Hualapai Mountains has been down-thrown 350 m since 18 million years ago with respect to the Colorado Plateau edge, based on present altitudes of the Miocene Peach Springs Tuff (Young and Brennan, 1974) on both sides of the Hualapai Valley, and the hypothesis that the ash-flow sheet flowed northeasterly from a now-buried source west of Kingman down into canyons already incised into the edge of the Colorado Plateau (Young and Brennan, 1974). However, Precambrian crystalline rocks in the Cerbat Mountains north of Kingman are now found at least 760 m higher than corresponding rocks beneath the Paleozoic strata at the Plateau edge. These observations indicate that, while Basin and Range grabens formed by subsidence, adjacent range blocks need not have been vertically shifted by great distances during Basin and Range faulting. Peirce (1976b) suggests a total vertical throw of 4,900 m on the Grand Wash fault just north of the Peach Springs Tuff outcrops at the western edge of the Colorado Plateau. The throw is distributed in unknown proportions between earlier rotational faulting and later basin-and-range faulting, but still left the Precambrian rocks of the Cerbats 760 m higher than their position at the plateau edge.

Zoback and others (1981) suggest that total Cenozoic Basin and Range crustal extension may be to 100 to 300 percent, but the extension during the late steep-angle block faulting may be only 10 to 30 percent; the rest may be assumed to have occurred during middle Tertiary time.

Coinciding with the change of tectonic style from low-angle detachment to steep-angle block faulting is a change of volcanic-rock chemistry (Rehrig and others, 1980; Shafiqullah and others, 1980). Volcanism older than 15 million years is subduction related; is dominated by andesite; silicic ash flows and tuff, and some basaltic andesite; and is characterized by strontium-isotope initial ratios in the 0.707 to 0.709 range. Volcanism younger than 15 million years throughout Arizona is dominated by alkali basalts that have large characteristic titanium, and strontium initial ratios of 0.704 to 0.705, indicative of mantle-derived, relatively uncontaminated magmas. The statewide distribution of basalt-dominated volcanic rocks younger than 15 million years is depicted on a map by Scarborough and others (1983).

Miocene, Pliocene and Quaternary history

An incursion of the Gulf of California into the lower Colorado River trough in response to the opening of the Gulf of California in late Miocene, and Pliocene time (Karig and Jansky, 1972) is postulated to explain the deposition of the estuarine-fluvial Bouse Formation from near Yuma to as far north as the Lake Mohave area (Smith, 1970; Eberly and Stanley, 1978; Lucchitta, 1979). The almost impermeable basal limestone of the Bouse Formation is a subsurface marker horizon in the region, and presumably affects ground-water flow characteristics.

Thickness of the Bouse Formation ranges from several meters along valley flanks to a few hundred meters in the southern part of the region around Yuma (Metzger and others, 1973). Present altitudes of the top of the formation indicate post-Bouse crustal warping of about 2,400 m between the southwest corner of Arizona and the Lake Mohave area (Lucchitta, 1979; Blair, 1978). The downwarping of the Bouse Formation south of Yuma probably is related to continued Gulf of California-rift tectonics; the postulated Bouse uplift in the Lake Mohave region has an unknown tectonic source.

A series of lacustrine, gypsiferous, red-brown clays that interfinger with or predate the Bouse Formation are penetrated in the subsurface in most, if not all, of the valleys along the lower Colorado and lower Gila-Salt Rivers (Ross, 1923; Eberly and Stanley, 1978). As shown in figure 4, Eberly and Stanley (1978) suggest that the clays were products of the same period of Basin and Range interior drainage that produced the massive evaporites in the basins farther east. The clays usually are overlain by Pleistocene sand and gravel along valley axes, and piedmont gravel along mountain fronts, indicating a progressive development of the modern Colorado-Gila-drainage system in post-Bouse time. Although aspects of the modern drainage pattern have undoubtedly been evolving through the Tertiary, the through-flowing Colorado and Gila River systems apparently began to form some time between 10 and 5 million years ago, based on evidence of closed-basin, nonmarine-halite accumulation older than 10.5 million years in the Luke basin near Phoenix (Eberly and Stanley, 1978) and interior drainages older than 5.9 million years in the Lake Mead area (Lucchitta, 1979; Eberly and Stanley, 1978).

Several cycles of intermittent river incision and stabilization have occurred through the last part of the Pleistocene. Areas where river incision, terrace formation, and pediment stripping have been the most pronounced are shown in figure 10. Several valleys in the southeastern part of the Sonoran region are still topographically closed as is most of Hualapai Valley in Mohave County. Even so, no Pleistocene pluvial lakes are recognized in the internally drained areas (Williams and Bedinger, 1984) either because they did not form, or possibly because of the easily eroded and abraded character of the Pleistocene surficial silt and sand in the valleys that may have recorded the shorelines.

References Cited

- Anderson, C. A., 1951, Older Precambrian structure in Arizona: Geological Society of America Bulletin, v. 62, no. 11., p. 1331-1346.
- Anderson, C. A., Blacet, P. M., Silver, L. T., and Stern, T. W., 1971, Revision of the Precambrian stratigraphy in the Prescott-Jerome area, Yavapai County, Arizona: U.S. Geological Survey Bulletin 1324-C, p. C1-C16.
- Anderson, C. A., and Silver, L. T., 1976, Yavapai series--A greenstone belt: Arizona Geological Society Digest, v. 10, p. 12-26.
- Anderson, J. L., 1983, Proterozoic anorogenic granite plutonism of North America, in Medaris, L. G., and others, eds., International Proterozoic symposium: Geological Society of America Memoir 161, p. 133-154.
- Anderson, R. E., Longwell, C. R., Armstrong, R. L., and Marvin, R. F., 1972, Significance of K-Ar ages of Tertiary rocks from the Lake Mead region, Nevada-Arizona: Geological Society of America Bulletin, v. 83, no. 2, p. 273-288.
- Anderson, R. E., Zoback, M. L., and Thompson, G. A., 1983, Implications of selected subsurface data on the structural form and evolution of some basins in the northern Basin and Range province, Nevada and Utah: Geological Society of America Bulletin, v. 94, no. 9, p. 1055-1072.
- Anderson, T. H., and Schmidt, V. A., 1983, The evolution of middle America and the Gulf of Mexico-Caribbean Sea region during Mesozoic time: Geological Society of America Bulletin, v. 94, no. 8, p. 941-966.

- Anderson, T. H., and Silver, L. T., 1971, Preliminary history for Precambrian rocks, Bamori region, Sonora, Mexico (abs.): Geological Society of America Abstracts with Programs, v. 3, no. 2, p. 72-73.
- 1981, The role of the Mojave-Sonosa megashear on the tectonic evolution of Sonora Mexico (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 2, p. 42.
- Armstrong, R. L., 1968, Sevier orogenic belt in Nevada and Utah: Geological Society of America Bulletin, v. 79, no. 4, p. 429-458.
- Blair, W. N., 1978, Gulf of California in Lake Mead area of Arizona and Nevada during late Miocene time: American Association of Petroleum Geologists Bulletin, v. 62, no. 7, p. 1159-1170.
- Burchfiel, B. C., and Davis, G. A., 1975, Nature and controls of Cordilleran orogenesis, western United States--Extensions of an earlier synthesis: American Journal of Science, v. 275-A, p. 363-396.
- 1981, Mojave Desert and environs, in Ernst, W. G., ed., The geotectonic development of California: Englewood Cliffs, New Jersey, Prentice-Hall, p. 217-252.
- Carr, M. D., 1983, Geometry and structural history of the Mesozoic thrust belt in the Goodspring district, southern Spring Mountains, Nevada: Geological Society of America Bulletin, v. 94, no. 10, p. 1185-1198.
- Condie, K. C., 1982, Plate tectonic model for Proterozoic continental accretion in southwestern United States: Geology, v. 10, no. 1, p. 37-42.

- Coney, P. J., 1976, Plate tectonics and the Laramide orogeny, in Woodward, L. A., and Northrup, S. A., eds., Tectonics and mineral resources of southwestern North America: New Mexico Geological Society Special Publication 6, p. 5-10.
- 1978, Magmatism and sedimentation in western North America during Mesozoic and Cenozoic time (abs.), in Najera, J. G., and Ortlieb, Lucien, eds., Resumes del Primer Simposio Sobre la Geologia y Potential Minera del Estado de Sonora, Hermosillo, Sonora, Mexico: Universidad de Norte America, Instituto de Geologia, p. 127-128.
- 1980, Cordilleran metamorphic core complexes--An overview, in Crittenden, M. D., Coney, J. P., and Davis, G. H., eds., Cordilleran core complexes: Geological Society of America Memoir 153, p. 7-31.
- Coney, P. J., and Reynolds, S. J., 1977, Cordilleran Benioff zones: *Nature*, v. 270, no. 5636, p. 403-405.
- Conway, C. M., 1976, Petrology, structure, and evolution of a Precambrian volcanic and plutonic complex, Tonto Basin, Gila County, Arizona: Pasadena, California Institute of Technology, unpublished Ph.D. dissertation, 460 p.
- Cooley, M. E., and Davidson, E. S., 1963, Mogollon Highlands-- Their influence on Mesozoic and Cenozoic erosion and sedimentation: *Arizona Geological Society Digest*, v. 6, p. 7-36.
- Damon, P. E., Shafiqullah, Muhammad, and Clark, K. F., 1981, Age trends of igneous activity in relation of metallogenesis in the southern Cordillera: *Arizona Geological Society Digest*, v. 14, p. 137-154.

- Davis, G. A., 1979, Problems of intraplate extensional tectonics, western United States, with special emphasis on their Great Basin, *in* Newman, G. W., and Goode, H. D., eds., 1979 Basin and Range symposium and Great Basin field conference: Denver, Colorado, Rocky Mountain Association of Geologists, p. 41-54.
- Davis, G. A., Anderson, J. L., Frost, E. G., and Shackelford, T. J., 1980, Mylonitization and detachment faulting in the Whipple-Buckskin-Rawhide Mountains terrane, southeastern California and western Arizona, *in* Crittenden, M. D., Jr., Coney, J. P., and Davis, G. H., eds., Cordilleran core complexes: Geological Society of America Memoir 153, p. 79-130.
- Dickinson, W. R., and Snyder, W. S., 1978, Plate tectonics of the Laramide orogeny, *in* Matthews, V., ed., Laramide folding associated with basement block faulting in the western United States: Geological Society of America Memoir 151, p. 355-366.
- Dings, M. G., 1951, The Wallapai mining district, Cerbat Mountains, Mohave County, Arizona: U.S. Geological Survey Bulletin 978-E, p. 123-162.
- Dixon, D. W., 1966, Geology of the New Cornelia mine, Ajo, Arizona, *in* Titley, S. R., and Hicks, C. L., eds., Geology of the porphyry copper deposits, southwestern North America: Tucson, University of Arizona Press, p. 123-132.
- Donnelly, M. E., and Hahn, G. A., 1981, A review of the Precambrian volcanogenic massive sulfide deposits in central Arizona and the relationship to their depositional environment: Arizona Geological Society Digest, v. 14, p. 11-22.

- Drewes, Harald, 1978, The Cordilleran orogenic belt between Nevada and Chihuahua: Geological Society of America Bulletin, v. 89, no 5, p. 641-657.
- Eberly, L. D., and Stanley, T. B., Jr., 1978, Cenozoic stratigraphy and geologic history of southwestern Arizona: Geological Society of America Bulletin, v. 89, no. 6, p. 921-940.
- Gilluly, James, 1946, The Ajo mining district, Arizona: U.S. Geological Survey Professional Paper 209, 112 p.
- Gilmour, Paul, and Still, A. R., 1968, The geology of the Iron King Mine, in Ridge, J. E., ed., Ore deposits of the United States, 1933-1967; The Graton-Sales volume: New York, American Institute of Mining and Metallurgical Engineers, v. 2, p. 1238-1257.
- Hamilton, Warren, 1982, Structural evolution of the Big Maria Mountains, southeastern California, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 1-27.
- Harding, L. E., 1980, Petrology and tectonic setting of the Livingston Hills Formation, Yuma County, Arizona: Arizona Geological Society Digest, v. 12, p. 135-146.
- 1982, Tectonic significance of the McCoy Mountains Formation, southeastern California and southwestern Arizona: Tucson, University of Arizona, unpublished Ph.D. dissertation, 197 p.

- Harding, L. E., Butler, R. F., and Coney, P. J., 1983,
Paleomagnetic evidence for Jurassic deformation of the McCoy
Mountains Formation, southeast California and southwest Arizona:
Earth and Planetary Science Letters, v. 62, no. 1, p. 104-114.
- Harshbarger, J. W., Repenning, C. A., and Irwin, J. H., 1957,
Stratigraphy of the uppermost Triassic and Jurassic rocks of
the Navajo Country: U.S. Geological Survey Professional
Paper 291, 74 p.
- Haxel, Gordon, and Dillon, John, 1978, The Pelona-Orocopia Schist
and Vincent-Chocolate Mountain thrust system, southern
California, in Howell, D. G., and McDougall, K. A., eds.,
Mesozoic paleogeography of the western United States--
Pacific Coast Paleogeography symposium 2: Los Angeles,
California, Society of Economic Paleontologists and
Mineralogists, p. 453-469.
- Haxel, Gordon, Wright, J. E., May, D. J., and Tosdal, R. M.,
1980, Reconnaissance geology of the Mesozoic and lower
Cenozoic rocks of the southern Papago Indian Reservation,
Arizona--A preliminary report: Arizona Geological Society
Digest, v. 12, p. 17-29.
- Howard, K. A., Goodge, J. W., and John, B. E., 1982, Detached
crystalline rocks of the Mohave, Buck, and Bill Williams
Mountains, western Arizona, in Frost, E. G., and Martin,
D., L., eds., Mesozoic-Cenozoic tectonic evolution of the
Colorado River region, California, Arizona, and Nevada: San
Diego, California, Cordilleran Publishers, p. 377-392.
- Karig, D. E., and Jensky, W., 1972, The proto-Gulf of California:
Earth and Planetary Science Letters, v. 17, no. 1, p. 169-174.

- Keith, S. B., 1978, Paleosubduction geometries inferred from Cretaceous and Tertiary magmatic patterns in southwestern North America: *Geology*, v. 6, no. 9, p. 516-521.
- 1981, Laramide tectonic and metallogenic evolution within the porphyry copper cluster of the southwest (abs.): *Arizona Geological Society Bulletin* 14, p. 287.
- 1982, Paleoconvergence rates determined from K_2O/SiO_2 ratios in magmatic rocks and their application to Cretaceous and Tertiary tectonic patterns in southwestern North America: *Geological Society of America Bulletin*, v. 93, no. 6, p. 524-532.
- Keith, S. B., and Barrett, L. F., 1976, Tectonics of the central Dragoon Mountains--A new look: *Arizona Geological Society Digest*, v. 10, p. 169-204.
- Keith, S. B., Reynolds, S. J., and Richard, S. M., 1982, Preliminary geologic map of the western Harquahala Mountains, west-central Arizona: Arizona Bureau of Geology and Mineral Technology Open-File Report 82-6, scale 1:1,000,000, 1 sheet.
- Kessler, E. J., 1976, Rubidium-strontium geochronology and trace element geochemistry of precambrian rocks in the northern Hualapai Mountains, Mohave County, Arizona: Tucson, University of Arizona, unpublished M.S. thesis, 193 p.
- Klein, D. P., 1982, Regional gravity and magnetic evidence for a tectonic weakness across southwestern Arizona, *in* Frost, E. G., and Martin, D. L., eds., *Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada*: San Diego, California, Cordilleran Publishers, p. 61-68.

- Livingston, D. E., 1969, Geochronology of older Precambrian rocks in Gila County, Arizona: Tucson, University of Arizona, unpublished Ph.D. dissertation, 224 p.
- Lucchitta, Ivo, 1979, Late Cenozoic uplift of the southwestern Colorado Plateau and adjacent lower Colorado River region, in McGretchin, T. R., ed., Plateau uplift; mode and mechanism: Tectonophysics, v. 61, no. 1-3, p. 63-95.
- Lysonski, J. C., Aiken, C. L. Z., and Sumner, J. S., 1981, Complete residual Bouguer gravity anomaly maps of Arizona: Arizona Bureau of Geology and Mineral Technology Open-File Report 81-24, scale 1:250,000, 23 maps.
- Menges, C. M., 1983, Neotectonic framework of Arizona-- Implications for the regional character of Basin-Range tectonism: Arizona Bureau of Geology and Mineral Technology Open-File Report 83-19, 101 p.
- Metzger, D. G., Loeltz, O. J., and Irelan, Burdge, 1973, Geohydrology of the Parker-Blythe-Cibola area, Arizona and California: U.S. Geological Survey Professional Paper 486-G, 130 p.
- Miller, F. K., 1970, Geologic map of the Quartzsite Quadrangle, Yuma County, Arizona: U.S. Geological Survey Geologic Quadrangle Map GQ-841, scale 1:62,500.
- Miller, F. K., and McKee, E. H., 1971, Thrust and strike-slip faulting in the Plomosa Mountains, southwestern Arizona: Geological Society of America Bulletin, v. 82, no. 3, p. 717-722.

- Nakata, J. K., 1982, Preliminary report on diking events in the Mohave Mountains, Arizona, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 85-90.
- Oppenheimer, J. M., and Sumner, J. S., 1980, Depth-to-bedrock map, Basin and Range province, Arizona: Tucson, University of Arizona Laboratory of Geophysics, scale 1:1,000,000.
- Peirce, H. W., 1976a, Elements of Paleozoic tectonics in Arizona: Arizona Geological Society Digest, v. 10, p. 37-57.
- 1976b, Tectonic significance of Basin and Range thick evaporite deposits: Arizona Geological Society Digest, v. 10, p. 325-339.
- Peirce, H. W., Damon, P. E., and Shafiqullah, Muhammad, 1979, An Oligocene(?) Colorado Plateau edge in Arizona, in McGretchin, T. R., ed., Plateau uplift; mode and mechanism: Tectonophysics, v. 61, no. 1-3, p. 1-24.
- Pike, J. E. N., and Hansen, V. L., 1982, Complex Tertiary stratigraphy and structure, Mohave Mountains, Arizona--A preliminary report, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 91-96.

- Pridmore, C. L., and Craig, C., 1982, Upper plate structure and sedimentation of the Baker Peaks area, Yuma County, Arizona, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 356-376.
- Rehrig, W. A., 1982, Metamorphic core complexes of the southwestern United States--An update, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 551-559.
- Rehrig, W. A., and Heidrick, T. L., 1976, Regional tectonic stress during the Laramide and late Tertiary intrusive periods, Basin and Range province, Arizona: Arizona Geological Society Digest, v. 10, p. 205-228.
- Rehrig, W. A., and Reynolds, S. J., 1980, Geologic and geochronologic reconnaissance of a northwest-trending zone of metamorphic core complexes in southern and western Arizona, in Crittenden, M. D., Jr., Coney, J. P., and Davis, G. H., eds., Cordilleran core complexes: Geological Society of America Memoir 153, p. 131-158.
- Rehrig, W. A., Shafiqullah, Muhammad, and Damon, P. E., 1980, Geochronology, geology, and listric normal faulting of the Vulture Mountains, Maricopa County, Arizona: Arizona Geological Society Digest, v. 12, p. 89-110.
- Reynolds, S. J., 1980, Geologic framework of west-central Arizona: Arizona Geological Society Digest, v. 12, p. 1-16.

- Reynolds, S. J., Keith, S. B., and Coney, P. J., 1980, Stacked overthrusts of Precambrian crystalline basement and inverted Paleozoic sections emplaced over Mesozoic strata, west-central Arizona: Arizona Geological Society Digest, v. 12, p. 45-52.
- Richard, S. M., 1983, Structure and stratigraphy of the southern Little Harquahala Mountains, La Paz County, Arizona: Tucson, University of Arizona unpublished M.S. thesis, 154 p.
- Robison, B. A., 1980, Description and analysis of Mesozoic "Red Beds", western Arizona and southeastern California: Arizona Geological Society Digest, v. 12, p. 147-154.
- Rogers, J. J. W., Burchfiel, B. C., Abbott, E. W., Anepohl, J. K., Ewing, A. H., Koehnken, P. J., Novitsky-Evans, J. M., and Talukdar, S. C., 1974, Paleozoic and lower Mesozoic volcanism and continental growth in the western United States: Geological Society of America Bulletin, v. 85, no. 12, p. 1913-1924.
- Ross, C. P., 1923, Lower Gila region, Arizona--A geographic, geologic, and hydrologic reconnaissance, with a guide to desert watering places: U.S. Geological Survey Water-Supply Paper 498, 237 p.
- Scarborough, R. B., and Meader, Norman, 1983, Reconnaissance geology of the northern Plomosa Mountains, La Paz County, Arizona: Arizona Geological Society Field Guidebook for Fall 1983 Trip, p. 1-35.

- Scarborough, R. B., Menges, C. M., and Pearthree, P. A., 1983, Map of Basin and Range (post 15 m.y.o.) exposed faults, grabens, and basalt-dominated volcanism in Arizona: Arizona Bureau of Geology and Mineral Technology Open-File Report 83-21, scale 1:500,000.
- Scarborough, R. B., and Peirce, H. W., 1978, Late Cenozoic basins of Arizona, in Callender, J. F., Wilt, J., Clemons, R. E., and James, H. L., eds., Land of Cochise: New Mexico Geological Society 29th Annual Field Conference Guidebook, p. 253-259.
- Scarborough, R. B., and Wilt, J. C., 1979, A study of uranium favorability of Cenozoic sedimentary rocks, Basin and Range province, Arizona: U.S. Geological Survey Open-File Report 79-1429, 101 p.
- Schumacher, Dietmer, Witter, D. P., Meader, S. J., and Keith, S. B., 1976, Late Devonian tectonism in southeastern Arizona: Arizona Geological Society Digest, v. 10, p. 59-70.
- Sears, J. W., and Price, R. A., 1978, The Siberian connection--A case for Precambrian separation of the North American and Siberian cratons: *Geology*, v. 6, no. 5, p. 267-270.
- Shackelford, T. J., 1976, Structural geology of the Rawhide Mountains, Mohave County, Arizona: Los Angeles, University of Southern California, unpublished Ph.D. dissertation, 166 p.
- 1980, Tertiary tectonic denudation of a Mesozoic-early Tertiary(?) gneiss complex, Rawhide Mountains, western Arizona: *Geology*, v. 8, no. 4, p. 190-194.

- Shafiqullah, Muhammad, Damon, P. E., Lynch, D. J., Reynolds, S. J., Rehrig, W. A., and Raymond, R. H., 1980, K-Ar geochronology and geologic history of southwestern Arizona and adjacent areas: *Arizona Geological Society Digest*, v. 12, p. 210-260.
- Sherborne, J. E., Jr., Buckovic, W. A., DeWitt, D. G., Hellinger, T. S., Pavlak, S. J., 1979, Major uranium discovery in volcaniclastic sediments, Basin and Range province, Yavapai County, Arizona: *American Association of Petroleum Geologists Bulletin*, v. 63, no. 4, p. 621-646.
- Silver, L. T., 1960, Age determinations on Precambrian diabase differentiates in the Sierra Ancha, Gila County, Arizona (abs.): *Geological Society of America Bulletin*, v. 71, no. 12, pt. 2, p. 1973-1974.
- 1968, Precambrian batholiths of Arizona (abs.): *Geological Society of America Special Paper 121*, p. 558-559.
- Silver, L. T., and Anderson, T. H., 1974, Possible left-lateral early to middle Mesozoic disruption of the southwestern North American craton margin (abs.): *Geological Society of America Abstracts with Programs*, v. 6, no. 7, p. 955-956.
- Silver, L. T., Bickford, M. E., and Van Schmus, W. R., 1977, The 1.4-1.5 B.Y. transcontinental anorogenic plutonic deformation of North America (abs.): *Geological Society of America Abstracts with Programs*, v. 9, no. 7, p. 1176-1177.
- Smith, P. B., 1970, New evidence for a Pliocene marine embayment along the lower Colorado River area, California and Arizona: *Geological Society of America Bulletin*, v. 81, no. 5, p. 1411-1420.

Spencer, J. E., 1984, The role of tectonic denudation in warping and uplift of low-angle normal faults: *Geology*, v. 12, no. 2, p. 95-99.

Stensrud, H. L., and More, Syver, 1980, Precambrian geology and massive sulfide environments of the west-central Hualupai Mountains, Mohave County, Arizona: *Arizona Geological Society Digest*, v. 12, p. 155-166.

Stewart, J. H., 1971, Basin and Range structure--A system of horsts and grabens produced by deep-seated extension: *Geological Society of America Bulletin*, v. 82, no. 4, p. 1019-1044.

——— 1976a, Late Precambrian evolution of North America--Plate tectonic implications: *Geology*, v. 4, no. 1, p. 11-15.

——— 1976b, Late Precambrian evolution of North America--Plate tectonic implications, discussion: *Geology*, v. 4, no. 6, p. 325-326.

——— 1978, Basin and Range structure in western North America--A review, in Smith, R. B., and Eaton, G. P., eds., *Cenozoic tectonics and regional geophysics of the western Cordilleran*: *Geological Society of America Memoir* 152, p. 1-31.

——— 1980, Regional tilt patterns of late Cenozoic basin-range fault blocks, western United States: *Geological Society of America Bulletin*, pt. 1, v. 91, no. 8, p. 460-464.

- Stone, Paul, Howard, K. A., and Hamilton, Warren, 1983,
Correlation of metamorphosed Paleozoic strata of the
southeastern Mojave Desert region, California and Arizona:
Geological Society of America Bulletin, v. 94, no. 10,
p. 1135-1147.
- Suneson, N. H., and Lucchitta, Ivo, 1979, K/Ar ages of Cenozoic
volcanic rocks, west-central Arizona: IsoChron/West, no. 24,
p. 25-29.
- 1983, Origin of bimodal volcanism, southern Basin and Range
province, west-central Arizona: Geological Society of
America Bulletin, v. 94, no. 8, p. 1005-1019.
- Thomas, B. E., 1953, Geology of the Chloride Quadrangle, Arizona:
Geological Society of America Bulletin, v. 64, no. 4, p. 391-420.
- Thorson, J. P., 1971, Igneous petrology of the Oatman district,
Mohave County, Arizona: University of California at Santa
Barbara, unpublished Ph.D dissertation, 233 p.
- Titley, S. R., 1981, Geologic and geotectonic setting of porphyry
copper deposits in the southern Cordillera: Arizona
Geological Society Digest, v. 14, p. 79-97.
- 1982, ed., Advances in geology of the porphyry copper
deposits of southwestern North America: Tucson, University of
Arizona Press, 560 p.

- Tosdal, R. M., 1982, The Mule Mountains thrust in the Mule Mountains, California and its probable extension in the southern Dome Rock Mountains, Arizona, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonic evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 56-60.
- Wasserburg, G. L., and Lanphere, M. A., 1965, Age determinations in the Precambrian of Arizona and Nevada: Geological Society of America Bulletin, v. 76, no., 7, p. 735-758.
- Wernicke, B., 1981, Low angle normal faults in the Basin and Range province--nappe tectonics in an extending orogen: Nature, v. 291, no. 5817, p. 645-648.
- Wilkinson, W. H., Vega, L. A., and Titley, S. R., 1982, Geology and ore deposits at Mineral Park, in Titley, S. R., ed., Advances in geology of the porphyry copper deposits: Tucson, University of Arizona Press, p. 523-542.
- Williams, T. R., and Bedinger, M. S., 1984, Selected geologic and hydrologic characteristics of Basin and Range province--Pleistocene lakes and marshes: U.S. Geological Survey Miscellaneous Investigations Series Map I-1522-D, scale 1:2,500,000.
- Wilson, E. D., 1939, Pre-Cambrian Mazatzal revolution in central Arizona: Geological Society of America Bulletin, v. 50, no. 7, p. 1113-1164.
- 1962, Resume of the geology of Arizona: Arizona Bureau of Mines Bulletin 171, 140 p.

- Wilson, E. D., Moore, R. T., and Cooper, J. R., 1969, Geologic map of Arizona: Arizona Bureau of Mines and U.S. Geological Survey, scale 1:500,000.
- Wilt, J. C., and Scarborough, R. B., 1981, Cenozoic sediments, volcanics, and related uranium in the Basin and Range province of Arizona, in Goodell, P. C., and Waters, A. C., eds., Uranium in volcanic and volcanoclastic rocks: American Association of Petroleum Geologists Studies in Geology 13, p. 123-144.
- Yeats, K., 1984, Nature of a major tectonic boundary which juxtaposes cratonic North America an Mesozoic sedimentary terrane, northern Dome Rock Mountains, Western Arizona: University of Arizona Geoscience Department, Geoscience Daze Abstract, Master's thesis, Research Progress Report, March 1984, p. 47.
- Young, R. A., and Brennan, W. J., 1974, Peach Springs Tuff--Its bearing on structural evolution of the Colorado Plateau and development of Cenozoic drainage in Mohave County, Arizona: Geological Society of America Bulletin, v. 85, no. 1, p. 83-90.
- Zoback, M. L., Anderson, R. E., and Thompson, G. A., 1981, Cainozoic evolution of the state of stress and style of tectonism of the Basin and Range province of the western United States, in Vine, F. J., convener, Extensional tectonics associated with convergent plate boundaries: Royal Society of London, Philosophical Transactions, Series A., v. 300, no. 1454, p. 407-434.

POTENTIAL HOST MEDIA FOR RADIOACTIVE WASTE

By

K. A. Sargent, U.S. Geological Survey

Host media considered to have potential for isolation of high-level radioactive waste in the Sonoran region, Arizona, include intrusive rocks, such as granite and other coarse-grained plutonic rocks; ash-flow tuff, especially where densely welded and having a thickness greater than 100 m; basalt and basaltic andesite lava flows, where greater than 100 m thick; and salt. Other, less abundant rock types that have potential may occur in the region. These include laharic and mudflow breccia and certain shallow intrusive bodies such as rhyolite domes. The occurrence and thickness of argillaceous sedimentary and metasedimentary rocks is summarized by Johnson (1984). Argillaceous rocks of the Sonoran region, Arizona, are folded, faulted, and deformed; they appear to have little or no potential as host rock in the region. In addition to the above mentioned rock types, basin-fill deposits and possibly other rock types have potential as host media in the unsaturated zone. The outcrop areas of potential host media and outcrop areas of potential host media and areas believed to have thick unsaturated zones in the Sonoran region, Arizona, are shown in plate 1.

Intrusive Rocks

Granitic rocks are widespread in the Sonoran region of Arizona. In some areas, granitic rocks form the entire core of mountain ranges, elsewhere the granitic rocks are sheared and have overridden younger rocks in upper plates of detachment faults. The distribution, age, and lithology of the granitic rocks of the region are summarized by Johnson and Scarborough (1984a).

Ground-water units SA-01, SA-02, and western ground-water unit SA-03 are underlain by the extensive Tertiary and Cretaceous age Gunnery Range batholith. Elsewhere in ground-water unit SA-03 are late Mesozoic to early Tertiary and Precambrian granitic rocks. The granitic rocks crop out in mountain ranges and probably underlie Quaternary and Tertiary basin fill at relatively shallow depths. The granitic rocks are mostly leucogranite and quartz monzonite, which are locally extensively fractured and foliated.

Very few granitic rocks crop out in ground-water units SA-04 and SA-05. These are Mesozoic to Tertiary granite and quartz diorite. Granite and quartz monzonite are widespread in the mountain ranges of ground-water unit SA-06. Granitic rocks are of Precambrian, Mesozoic and early Tertiary age. Granitic stocks that occur in the Harquahala Mountains core complexes in the eastern part of ground-water unit SA-06 are believed to be underlain by shallow thrust faults.

In ground-water unit SA-07 the intrusive rocks are granite, quartz monzonite and quartz diorite mostly of Precambrian age with some of Mesozoic and Tertiary age. Granite occurs as younger Precambrian and Mesozoic stocks intruding older Precambrian granite. Of special interest is a large body of Precambrian gabbro and anorthosite in the vicinity of Bagdad.

Ground-water units SA-08, SA-09, and SA-10 contain Precambrian granite, granodiorite, and quartz monzonite, that have intruded older Precambrian granitic rocks. A few Cretaceous and Tertiary granitic stocks intrude Precambrian gneissic and granitic rocks.

Tuffaceous Rocks

Tuffs of widespread occurrence, include ash-flow tuff, some interbedded tuffaceous sedimentary rocks, ash-fall tuff and minor rhyolitic lava flows. Welded tuffs generally are thin and mixed with other volcanic rocks. For details on distribution and description of individual occurrences the reader is referred to Jenness and others (1984).

In ground-water units SA-01 to SA-05 are mixed volcanic rocks consisting of ash-flow tuff, volcanic sedimentary rocks, rhyolitic and andesitic lava flows and breccia. Ash flows of middle Tertiary age are as much as 100 m thick in ground-water units SA-04 and SA-05. Tuffs and lavas in the northeastern part of ground-water unit SA-03 of Cretaceous age are deformed and fractured.

In ground-water unit SA-06, mixed volcanoclastic rocks of middle Tertiary age occur in the central part of the unit. The thickness of these rocks generally is unknown; however, thick tuffs are reported in the Eagletail Mountains in the easternmost part of ground-water unit SA-06.

Ground-water unit SA-07 contains mostly thin ash-flow deposits of rhyolitic composition. However, southwest of Bagdad, the Grayback Mountain Rhyolite Tuff of Late Cretaceous(?) or early Tertiary(?) age is as much as 150 m thick. Ground-water units SA-08, SA-09, and SA-10 contain outcrops of ash-flow tuffs complexly mixed with agglomerate, and andesitic, rhyolitic, and basaltic lava flows. Dated ash flows are 17 to 18 million years old, about the age of inception of the basin-and-range faulting. These, as well as other ash flows in the area, are thin, generally less than 50 m.

Basaltic Rocks

Basalt flows are widespread and of middle to late Tertiary age, contemporaneous with basin-and-range faulting. For details on distribution and descriptions of individual occurrences, the reader is referred to Johnson and Scarborough (1984b).

In ground-water units SA-01 to SA-06, most of the basalts are 10 to 20 million years old (Miocene) and are olivine basalts and basaltic andesites. Maximum aggregate thickness is 400 m in the Lomosa Mountains and 450 m near Ajo; individual flows are no more than 15 to 20 m thick. In ground-water unit SA-07 upper Tertiary olivine basalt, basaltic andesite, and interbedded clastic rocks are no more than 100 m thick. Ground-water units SA-08, SA-09, and SA-10 have scattered upper Tertiary basalt and andesite flows. Individual flows generally are 3 to 10 m thick, but some flows in the Miocene Mount Davis Volcanics are greater than 75 m thick. Flows commonly are interbedded with tuffaceous sedimentary rocks.

Salt

Large salt deposits occur in Hualapai and Detrital Valleys, in the northern part of the Sonoran region of Arizona (Peirce, 1981a; Ege, 1984). The Red Lake deposit, beneath Hualapai Valley, is buried by more than 450 m of sedimentary cover. Geophysical modeling indicates the halite body to be 8 km wide, 18 km long, and probably 3.2 km thick. A drill hole has penetrated 1,220 m of salt. Peirce (1981b) reports that a natural-gas storage facility is planned for the Red Lake salt. The Detrital Valley salt deposit occurs south of Lake Mead. It is buried by as much as 600 m of sedimentary fill and is as much as 335 m thick. The Detrital Valley salt is associated with gypsum, anhydrite, and clastic sediments belonging to the Muddy Creek Formation of Miocene and Pliocene age. Other deep valleys in western Arizona may contain thick salt or other evaporitic bodies that are unknown at this time.

Other Rocks

Metamorphosed and unmetamorphosed Mesozoic sedimentary rocks, Precambrian metamorphic rocks, Quaternary basalts, and basin-fill material also are present in the Sonoran region of Arizona. Generally these rocks are not considered potential host rocks and were not discussed separately. However, some metamorphic rocks, mainly granite gneiss, were mapped with the granite because of their close spatial relationship to, and similar lithologic properties with, granitic rocks. Quaternary basaltic rocks locally are extensive, but because of their relatively recent age they are not considered host rocks. Basin fill is considered as potential host rock in the next section on the unsaturated zone.

Unsaturated Zone

With the exception of ground-water unit SA-07, the ridges in the region are believed to contain unsaturated sections as great as 150 m thick. Basin fill contains as much as 150 m of unsaturated Quaternary and Tertiary sediment in the Sacramento Valley (ground-water unit SA-08), Hualapai Valley (SA-09), Detrital Valley (SA-10), Dutch Flat (SA-07 and SA-08), and Date Creek (SA-07).

Volcanic rocks generally are above the water table and because they are relatively permeable, they are considered satisfactory for potential host rocks in the unsaturated zone. However, their thickness probably is great enough only locally to provide a site for a repository in the unsaturated zone.

References Cited

- Ege, J. R., 1985, Maps showing distribution, thickness, and depth of salt deposits of the United States: U.S. Geological Survey Open-File Report 85-28, 4 plates, scale 1:2,500,000.
- Jenness, J. E., Lopez, D. A., and LaFortune, J. R., 1984, Map showing outcrops of pre-Quaternary ash-flow tuffs and volcanoclastic rocks, Basin and Range province, Arizona: U.S. Geological Survey Water-Resources Investigations Map 83-4114-F, scale 1:500,000.
- Johnson, W. D., Jr., 1984, Map showing outcrops of thick, dominantly argillaceous sedimentary and metasedimentary rocks, Basin and Range province, Arizona: U.S. Geological Survey Water-Resources Investigations Map 83-4114-E, scale 1:500,000.
- Johnson, W. D., Jr., and Scarborough, R. B., 1984a, Map showing outcrops of granitic rocks, Basin and Range province, Arizona: U.S. Geological Survey Water-Resources Investigations Map 83-4114-D, scale 1:500,000.
- 1984b, Map showing outcrops of pre-Quaternary basaltic rocks, Basin and Range province, Arizona: U.S. Geological Survey Water-Resources Investigations Map 83-4114-G, scale 1:500,000.
- Peirce, H. W., 1981a, Major Arizona salt deposits: Arizona Bureau of Geology and Mineral Technology Fieldnotes, v. 11, no. 4, p. 1-5.
- 1981b, Natural gas storage in Arizona salt: Arizona Bureau of Geology and Mineral Technology Fieldnotes, v. 11, no. 3, p. 8.


QUATERNARY TECTONISM

By K. A. Sargent, U.S. Geological Survey

and Robert B. Scarborough,

Arizona Bureau of Geology and Mineral Technology

Quaternary tectonism of the Sonoran region, Arizona is indicated by the seismicity, heat flow, Quaternary faulting, late Cenozoic volcanic activity and vertical movement of the region. Each of these features is depicted in figure 11.



- EXPLANATION
- EPICENTER OF EARTHQUAKE OF MAGNITUDE 4-5 (RICHTER SCALE)
 - EPICENTER OF EARTHQUAKE OF MAGNITUDE 5-6 (RICHTER SCALE) AS DEFINED BY ASKEW AND ALGERMISSEN, 1983
 - △ EPICENTER OF EARTHQUAKE OF MAGNITUDE 6-7 (RICHTER SCALE) AS DEFINED BY ASKEW AND ALGERMISSEN, 1983
 - ||||| LINE OF EQUAL STRAIN RELEASE IN EQUIVALENT MAGNITUDE 4.0 (RICHTER SCALE) EARTHQUAKES PER 823 SQUARE KILOMETERS.--Hachured on side of lesser strain release
 - QUATERNARY FAULT.--Dashed where approximately located
 - HEAT-FLOW MEASUREMENT IN EXCESS OF 2.5 HEAT FLOW UNITS
 - ||||| LINE OF EQUAL UPWARD CRUSTAL MOVEMENT, IN METERS PER 10,000 YEARS.--Hachured on side of lesser upward crustal movement
 - ||||| LINE OF EQUAL UPWARD CRUSTAL MOVEMENT, IN MILLI-METERS PER YEAR.--Hachured on side of lesser upward crustal movement
 - OUTCROP OF UPPER CENOZOIC VOLCANIC ROCK

Figure 11.--Seismicity, heat flow, Quaternary faults, late Cenozoic volcanic rocks, and vertical crustal movement.

Seismicity

Less than 30 earthquakes are reported for the Sonoran region of Arizona in a regional compilation by Algermissen and others, (1983). Two earthquakes have Richter magnitude (surface wave) 5 to 6 and two have magnitude 4 to 5 (fig. 11), the rest are less than magnitude 4. Of the two largest earthquakes, one is located near Lake Mead and may be related to its filling, the other is near the lower Gila River. Two additional large earthquakes, magnitude 5 to 6 and 6 to 7, occurred north of Imperial Reservoir, just outside the region to the west, but affect the region as shown by the strain-release lines. Areas showing greatest strain release are in the southwest, where the two large earthquakes north of Imperial Reservoir occurred, and near Lake Mead in the northwest part of the region. A similar compilation of earthquake data has been made for Arizona by DuBois and others (1982). Within the Sonoran region they report 11 instrumentally recorded and 5 historically recorded epicenters of Richter magnitude less than 5.0 between 1830 and 1980. Menges and Pearthree (1983, fig. 5) suggest a correlation between historical seismicity and late Pliocene and Pleistocene faulting within a 200-km-wide zone that extends from the western Colorado Plateau diagonally southeastward into southwestern New Mexico. Menges also indicates a 60-km-wide zone of historical seismicity (11 recorded epicenters) in the southernmost Sonoran region that trends west-northwest through the Yuma area, parallel to the Mexican border, that has only minimal prehistoric surface-rupture expression. The Algodones fault with obvious surface rupture lies in this zone. It is located near the Mexican border 35 km

southeast of Yuma. Menges' 60-km-wide seismicity zone may merge to the west with the San Andreas fault system in California, although the two trends are divergent by 30^o.

Heat Flow

About 40 heat-flow measurements are reported for the Sonoran region of Arizona (J.H. Sass, U.S. Geological Survey, written commun., 1982). Of these measurements, only two exceed 2.5 heat-flow units (HFU) (fig. 11). The maximum measurement is 2.82 HFU. The entire Sonoran region of Arizona has a heat flow ranging from 1.5 to 2.5 HFU (J.H. Sass, written commun., 1982).

Quaternary Faulting

Mapped Quaternary faults are exceedingly few in the Sonoran region of Arizona (Nakata and others, 1982; Menges and Pearthree, 1983). In addition to those faults mentioned in the previous section on seismicity, a few short segments occur around the periphery of the region; none have been mapped in the interior (fig. 11). Quaternary faults that have been studied show displacements older than 10,000 years.

Late Cenozoic Volcanics

The Sentinel Plain-Arlington volcanic field occurs north and south of the Gila River near Gila Bend. Only the Sentinel Plain part of the field is in the region (fig. 11). It contains fine-grained olivine basalt flows and cinder cones, have late Cenozoic dates ranging from 1.72 to 3.19 million years (Aldrich and Laughlin, 1981; Eberly and Stanley, 1978; Shafiqullah and others, 1980). Judging on weathering characteristics of the flows and the K/Ar dates, only a small part of the Sentinel Plain is believed to be Quaternary in age (R.B. Scarborough, Arizona Bureau of Geology, unpublished data, 1984). Most of the Quaternary radiometric dates come from the Arlington and Gillespie cones farther to the northeast.

Aldrich and Laughlin (1981) show basalt no older than 3 million years cropping out north of Imperial Reservoir in the southwestern part of the region. Garner and others (1982) suggest that the flow, shown on the State geologic map as Quaternary basalt (Qb) (Wilson and others, 1969), is actually late Miocene or Pliocene in age.

Vertical Crustal Movement

Gable and Hatton (1983) show an area of uplift in southern California and southwestern Arizona. Based on geodetic-leveling, the southwestern Sonoran region of Arizona is rising at a rate of 4 mm/yr. Gable and Hatton (1983) show historic subsidence in the Lake Mead area of as much as 2 m based on leveling data from 1935 to 1950, although the general area is rising at a rate of 10 mm/yr. The subsidence related to the filling of Lake Mead is summarized by Anderson and Laney (1975).

References Cited

- Aldrich, M. J., and Laughlin, A. W., 1981, Age and location of volcanic centers <3.0 m.y. old in Arizona, New Mexico, and the Trans-Pecos area west of Texas: Los Alamos, New Mexico, Los Alamos National Laboratory Map LA-8812-MAP, revised, UC-11, scale 1:1,000,000.
- Algermissen, S. T., Askew, B. L., Thenhaus, P. C., Perkins, D. M., Hanson, S., and Bender, B. L., 1983, Seismic energy release and hazard estimation in the Basin and Range province: U.S. Geological Survey Open-File Report 83-358, 13 p.
- Anderson, R. E., and Laney, R. L., 1975, The influence of late Cenozoic stratigraphy on distribution of impoundment-related seismicity at Lake Mead, Nevada-Arizona: U.S. Geological Survey Journal of Research, v. 3, no. 3, p. 337-343.
- Askew, Bonny, and Algermissen, S. T., 1983, An earthquake catalog for the Basin and Range province, 1803-1977: U.S. Geological Survey Open-File Report 83-86, 21 p.
- DuBois, S. M., Smith, A. W., Nye, N. K., and Nowak, T. A., Jr., 1982, Arizona earthquakes, 1776-1980: Arizona Bureau of Geology and Mineral Technology Bulletin 193, 456 p.
- Eberly, L. D., and Stanley, T. B., Jr., 1978, Cenozoic stratigraphy and geologic history of southwestern Arizona: Geological Society of America Bulletin, v. 89, no. 6, p. 921-940.

- Gable, D. J., and Hatton, Tom, 1983, Maps of vertical crustal movements in the conterminous United States over the last 10 million years: U.S. Geological Survey Miscellaneous Investigations Series Map I-1315, scale 1:5,000,000, 2 sheets.
- Garner, W. E., Frost, E. G., Tanges, S. E., and Germinario, M. P., 1982, Mid-Tertiary detachment faulting and mineralization in the Trigo Mountains, Yuma County, Arizona, in Frost, E. G., and Martin, D. L., eds., Mesozoic-Cenozoic tectonics evolution of the Colorado River region, California, Arizona, and Nevada: San Diego, California, Cordilleran Publishers, p. 158-172.
- Menges, C. M., and Pearthree, P. A., 1983, Map of Neotectonic (latest Pliocene-Quaternary) deformation in Arizona: Arizona Bureau of Geology and Mineral Technology Open-File Report 83-22, scale 1:500,000, 4 sheets.
- Nakata, J. K., Wentworth, C. M., and Machette, M. N., 1982, Quaternary fault map of the Basin and Range and Rio Grande rift provinces, western United States: U.S. Geological Survey Open-File Report 82-579, scale 1:2,500,000, 2 sheets.
- Shafiqullah, Muhammad, Damon, P. E., Lynch, D. J., Reynolds, S. J., Rehrig, W. A., and Raymond, R. H., 1980, K-Ar geochronology and geologic history of southwestern Arizona and adjacent areas, in Jenney, J. P., and Stone, Claudia, eds., Studies in western Arizona: Arizona Geological Society Digest, v. 12, p. 201-260.
- Wilson, E. D., Moore, R. T., and Cooper, J. R., 1969, Geologic map of Arizona: Arizona Bureau of Mines and U.S. Geological Survey, scale 1:500,000.

GROUND-WATER HYDROLOGY

By

M. S. Bedinger, William H. Langer, and J. E. Reed,
U.S. Geological Survey

The Sonoran region of Arizona is arid for the most part, with annual precipitation less than 200 mm throughout most of the region, and potential evaporation greater than 2.5 m. Annual precipitation is greatest in ground-water unit SA-07 reaching a maximum of about 500 mm. Surface drainage of the structural basins is open, except for part of the Hualapai basin (ground-water unit SA-09).

Major Hydrogeologic Units

The hydrogeologic units in the region include basin fill, volcanic rocks, clastic indurated sedimentary rocks, carbonate rocks, and crystalline rocks, including intrusive and metamorphic rocks.

Basin fill (including alluvial material of stream valleys) occurs largely in structural basins and is as thick as 3,000 m. The basin fill crops out at the land surface throughout a large part of the region, however, pediment surfaces underlie the fill at shallow depths beneath one-half to two-thirds of the basin-fill area.

The basin fill consists mostly of nonindurated to semi-indurated sedimentary terrestrial and marine deposits. The age of the deposits ranges from middle Tertiary to Holocene. The terrestrial deposits consist largely of poorly sorted to moderately sorted mixtures of gravel, sand, silt, and clay that were derived largely from the consolidated rocks in the nearby mountains. The fill also contains volcanic flows and ash falls from episodic volcanic activity during the Tertiary and Quaternary Periods. Fine-grained lake deposits, silt and clay, and evaporitic deposits of salt, gypsum, and limestone, have been found in many of the basins. The stratigraphic sequence of basin-fill deposits in the Ranegras Plain, the Wellton-Mohawk area, and the Gila Bend basin have been described from drill cuttings by Metzger (1952a, 1952b). The basin fill in most basins has not been fully penetrated by wells and the lithology is variable areally. However, the lithologic sequence described may be similar to sequences in other basins. The older fill is described as semi-indurated fine-grained deposits, identified as lake-bed deposits in some areas, with minor interbedded sand, ranging in thickness from 60 to 240 m. Overlying the fine-grained material is a coarser unit composed of sand and gravel described as being about 100 m thick. A thickness of more than 300 m of salt, and interbedded fine-grained material, presumably of lake deposition, has been reported near Red Lake in the Hualapai Valley (Gillespie and others, 1966).

Marine estuarine deposits in Tertiary (Miocene and Pliocene) extensions of the Gulf of California have been described in the Yuma area (Olmsted and others, 1973), and in the Parker-Blythe-Cibola area (Metzger and others, 1973). The marine deposits predominantly are thin-bedded silt and clay with some calcareous material, and may be as thick as 600 m in the Yuma area. Marine deposits are present in other basins in the region but have not been positively identified. Metzger (1952a) noted a fine-grained sequence in the Gila River valley, possibly equivalent to the marine deposits of the Yuma area. The maximum thickness of fill in individual basins ranges from about 250 m to 3,000 m with most basins having a maximum thickness of 1,000 to 2,000 m (Oppenheimer and Sumner, 1980).

Volcanic rocks are grouped hydrologically as undifferentiated volcanic rocks, which include flows and pyroclastic materials of Mesozoic and Tertiary age, and tuff of Tertiary age. Volcanic rocks overlie older consolidated rocks and locally compose a part of the basin filling. Coarse-clastic sedimentary rocks of Mesozoic age are as thick as 4,000 m locally, and are scattered throughout the region. Carbonate rocks of Paleozoic age occur in a limited area in the northern part of the region.

Crystalline rocks are widespread and underlie the entire region at depth. The rocks in this hydrogeologic unit include metamorphic rocks and intrusive-igneous rocks and range in age from Precambrian to Tertiary.

Ground-Water Flow Regime

Ground-water recharge occurs by infiltration of precipitation. The greatest opportunity for recharge is in areas of permeable surface rock during such periods when precipitation is in excess of evapotranspiration. Lacking direct evidence of the location and timing of recharge, it generally is assumed that much of the recharge occurs during surface runoff to basin-fill deposits near the mountain fronts. Some recharge occurs within the mountain ranges, but the minimal permeability of the rocks and steep slopes promote rapid runoff of rainfall and snowmelt. The basin fill, being more permeable, is more receptive to infiltration of runoff from the mountain ranges. The central parts of the basins receive some recharge, but probably minor quantities because of generally minimal precipitation and substantial potential evapotranspiration. Recharge to the central parts of the basin fill may largely occur at infrequent times during rare storm runoff.

Discharge of ground water occurs by seepage to gaining streams, evapotranspiration, springs, and by withdrawal from wells. Gaining streams include reaches of the Colorado River, the Gila River, and the Bill Williams River and its tributaries. Ground-water unit SA-07 contains numerous gaining stream reaches of the Bill Williams River and its tributaries, and contains a dozen or more noteworthy springs.

Springs having a discharge temperature of 31 °C or greater, and a discharge of 200 L/min or greater are shown in plate 3. The largest concentration of springs is in ground-water unit SA-07. These include two springs in the Big Sandy River drainage having temperatures of 37 °C. Two springs with temperature less than 30 °C, having known discharge of greater than 200 L/min also occur in ground-water unit SA-07.

Radium Hot Spring in the Gila River valley is now reported as dry (Berry and others, 1980). The spring was at one time developed as a health resort and had a temperature of 60 °C. The original flow was small and has ceased, possibly because of ground-water withdrawal by wells in the vicinity.

Four springs with temperature less than 30 °C flowing greater than 200 L/min are known to occur in the region outside ground-water unit SA-07. Two are in the eastern arm of ground-water unit SA-09 and two are in ground-water unit SA-08.

Ground water is withdrawn through wells, largely for irrigation, in localized areas scattered throughout the region. Greatest withdrawals are along the Gila and Colorado Rivers, and in the Sacramento Valley, Ranegras Plain, Butler Valley, and Haulapai Valley.

Ground-Water Flow Analysis

AREAL GROUND-WATER FLOW

Ground-water traveltime near the water table was analyzed using the procedure described in Chapter A (Bedinger, Sargent, and others, 1984). The relative velocity in the hydrogeologic units at the water table is shown in plate 2. Relative velocities are reported because site-specific data are not available for the region. Hydraulic properties of the hydrogeologic units were estimated using values in Chapter A and modified from the lithologic and hydrologic description of the units in the literature, and further modified as appropriate during verification of the cross-sectional models (Bedinger, Langer, and Reed, 1984). The values of hydraulic properties of the units and hydraulic gradients used in estimating relative ground-water velocities are given in table 1.

The hydraulic gradients for the hydrogeologic units are representative gradients obtained from the water-level contour map of the region (Langer and others, 1984). Relative ground-water traveltimes are shown in plate 3. Flow paths along which the relative ground-water traveltimes were calculated and major discharge areas also are shown in plate 3. Traveltimes are computed from ground-water divides to the gaining reaches of streams and major pumping centers.

Table 1.--Hydraulic properties of hydrogeologic units and hydraulic gradients used in estimating relative ground-water velocities at the water table.
 [K, hydraulic conductivity, in meters per day;
 \emptyset , effective porosity]

Hydrogeologic unit	Map symbol (pl. 2)	K/ \emptyset (meters per day)	Hydraulic gradient
Basin fill	a	6×10^1	0.003
Volcanic rocks			
Undifferentiated	v	1×10^{-1}	.03
Lava flows	b	3×10^0	.03
Ash-flow tuff	t	1×10^{-1}	.03
Granitic rocks	g	2×10^{-1}	.03
Metamorphic rocks	m	2×10^{-1}	.03
Coarse clastic rocks	s	2×10^{-1}	.03
Carbonate rocks	c	1×10^1	.003

The longest relative traveltimes, 10 to 20, occur from a few small areas of undifferentiated volcanic rocks and intrusive and metamorphic rocks near ground-water divides to discharge areas. Generally, the relative flow time from points in basin fill to discharge areas is less than 1.

The relative traveltimes indicated in the map from divide areas to discharge areas are extremely conservative because actual flow paths in recharge areas dip below the water table and take much longer flow paths and much longer traveltimes. The relative traveltimes are useful for comparing relative velocities near the water table and for calculating relative traveltimes at shallow depths between nearby points. A more realistic estimate of relative traveltime between widely spaced points, such as from near a water-table divide to a discharge area, is given in the discussion of cross-sectional models.

CROSS-SECTIONAL MODELS

Cross-sectional models were used to analyze ground-water flow along selected flow paths. The mathematical model used in modeling flow in section is given in Chapter A of this report (Reed, 1984). The map location of the sections and the model parameters and results are shown in plate 4. The values of hydraulic properties of the rock units in the hydrogeologic sections used in analysis of the ground-water flow are given in table 2.

Table 2.--Hydraulic properties of hydrogeologic units modeled in hydrogeologic sections.
 [K, hydraulic conductivity, in meters per day; ϕ , effective porosity]

Hydrologic unit	Symbol (plate 4)	Hydrogeologic sections in plate 4					
		A-A'		B-B'		C-C'	
		K	ϕ	K	ϕ	K	ϕ
Ash-flow tuff	t	4 X 10 ⁻⁴	3.5 X 10 ⁻¹	---	---	---	---
Crystalline rock, upper part of section	G	5 X 10 ⁻⁴	3 X 10 ⁻³	5 X 10 ⁻⁴	3 X 10 ⁻³	5 X 10 ⁻⁴	3 X 10 ⁻³
Crystalline rock, lower part of section	g	3 X 10 ⁻⁷	1 X 10 ⁻⁴	3 X 10 ⁻⁷	1 X 10 ⁻⁴	3 X 10 ⁻⁷	1 X 10 ⁻⁴
Undifferentiated volcanic rocks	v	4 X 10 ⁻⁴	4 X 10 ⁻³	4 X 10 ⁻⁴	4 X 10 ⁻³	4 X 10 ⁻⁴	4 X 10 ⁻³
Coarse-grained basin fill, upper part of section	a	5 X 10 ⁰	1.8 X 10 ⁻¹	5 X 10 ⁰	1.8 X 10 ⁻¹	1 X 10 ¹	1.8 X 10 ¹
Coarse-grained basin fill, lower part of section	B	1 X 10 ⁻¹	1.2 X 10 ⁻¹	1 X 10 ⁻¹	1.2 X 10 ⁻¹	1 X 10 ⁻¹	1.2 X 10 ⁻¹
Fine-grained basin fill	A	---	---	2 X 10 ⁻⁴	3.2 X 10 ⁻¹	---	---

Distribution of rock units, relative traveltime, and stream functions, are given in plate 4. Relative traveltimes are given in intervals of one order of magnitude from 10^1 and longer. Relative traveltimes indicate the relative time of travel from points on the flow line to the discharge area. Stream functions show the directions of ground-water movement and relative quantity of flow in the section below the flow line.

The sections give a more realistic concept of the traveltime between widely spaced points in the region, for example, between the water-table divide areas and the discharge areas. As shown in plate 4, the flow paths in the divide areas of the flow system dip steeply into the flow system and take the longest flow paths to the discharge areas. Relative traveltimes from the divide areas to discharge areas are as great as 10^6 to 10^8 . Commonly, these longest relative traveltimes are of restricted surface area at the water table. The areas of longer relative traveltime enlarge with depth. One would be more confident in locating an area of long traveltime at depth beneath the water table than near or above the water table where the area of long traveltime is small.

Broad areas of relative traveltime of 10^5 or greater exist at the water table in all sections and areas with traveltime from 10^6 to 10^8 exist within 1,000 m of the water table in all sections.

Quality of Ground Water

The quality of ground water in the Sonoran region, Arizona, is characterized by the areal distribution of dissolved solids (fig. 12) and predominant chemical constituents in solution (fig. 13). These maps are generalized from those compiled by Thompson and others (1984). The maps of Thompson and others (1984) were compiled from data in the water-quality files of the U.S. Geological Survey (WATSTORE) and in published reports. The data are mostly from non-geothermal springs and wells less than 150 m deep completed in alluvial and basin-fill deposits. In areas where data are not available, the water-quality characteristics were estimated from the position in the ground-water flow system and the lithology of the local bedrock.

In most of the region, the concentration of dissolved solids in ground water is less than 500 mg/L. Ground water near the Gila River contains 3,000 to 5,000 mg/L dissolved solids with concentrations as great as 15,000 mg/L. The dissolved-solids concentration near Lake Mead and the Colorado River is between 1,000 and 3,000 mg/L. Concentrations of dissolved solids are between 500 and 1,000 mg/L in a large part of the Ranegras Plain. The ground water is mostly of the calcium magnesium or sodium bicarbonate type. Waters in which the dominant anion is sulfate or chloride occur in and near large discharge areas and in some playas.




Figure 12.--Dissolved-solids concentration in ground water.




Figure 13.--Distribution of chemical types of ground water.

Pleistocene Hydrologic Conditions

There is no evidence of Pleistocene lakes in the Sonoran region of Arizona and the drainage in the Pleistocene Epoch is presumed to have been integrated as it is today, except for the closed basin occupied by the dry Red Lake. During glacial maxima, sea level may have been as much as 80 to 140 m lower (Bloom, 1978, p. 406) and entrenchment of the Colorado and Gila Rivers may have resulted in a lower base level for ground-water discharge. Glacial climate was probably characterized by increased precipitation and decreased temperature and evaporation as inferred from maximum Pleistocene lake levels in the Basin and Range province (Leopold, 1951; Reeves, 1966; and Snyder and Langbein, 1962) and from plant macrofossils (Spaulding, 1983). The dry Red Lake bed occupies part of a closed basin in ground-water unit SA-09, but contains no reported evidence of having been occupied by a lake during the Pleistocene. Peirce (1976) believes that the basin was occupied by a closed-basin lake during the Miocene during which time several hundred meters of evaporites with interbedded silt and clay accumulated. If the closed basin existed in the Pleistocene, as it probably did, the precipitation in the basin was lost by evaporation and infiltration to the subsurface and discharge as ground water to the Colorado River to the north. The entrenchment of the Colorado River probably in Pliocene (subsequent to 5.9 million years ago, Shafiqullah and others, 1980) and consequent development of a base level for ground-water discharge 625 m below the surface of the playa may have prevented the basin from developing a lake during the Pleistocene.

References Cited

- Bedinger, M. S., Langer, W. H., and Reed, J. E., 1985, Hydraulic properties of rocks in Bedinger, M. S., Sargent, K. A., and others, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Basis of characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].
- Bedinger, M. S. Sargent, K. A., and others, 1985, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Basis of characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].
- Berry, G. W., Grim, P. J., and Ikelman, J. A., 1980, Thermal springs list for the United States: National Oceanic and Atmospheric Administration Key to Geophysical Records Document 12, 59 p.
- Bloom, A. L. 1978, Geomorphology--A systematic analysis of late Cenozoic landforms: Englewood Cliffs, New Jersey, Prentice-Hall, Inc., 510 p.
- Gillespie, J. G., Bentley, C. B., and Kam, William, 1966, Basic hydrologic data of the Hualapai, Sacramento, and Big Sandy Valleys, Mohave County, Arizona: Arizona Land Department Water-Resources Report 26, 39 p.

- Langer, W. H., Mulvihill, D. A., and Anderson, T. W., 1984, Maps showing ground-water levels, springs, and depth to ground water, Basin and Range province, Arizona: U.S. Geological Survey Water-Resources Investigations WRI 83-4114-B, scale 1:500,000.
- Leopold, L. B., 1951, Pleistocene climate in New Mexico: American Journal of Science, v. 249, no. 2, p. 152-168.
- Metzger, D. G., 1952a, Welton-Mohawk area, Yuma County, in Halpenny, L. C., and others, Ground water in the Gila River basin and adjacent areas, Arizona--A summary, Part II, Ground water in individual areas: U.S. Geological Survey open-file report, p. 165-170.
- 1952b, Ranegras Plain area, Yuma County, in Halpenny, L. C., and others, Ground water in the Gila River basin and adjacent areas, Arizona--A summary, Part II, Ground water in individual areas: U.S. Geological Survey open-file report, p. 171-176.
- Metzger, D. G., Loeltz, O. J., and Irelan, Burdge, 1973, Geohydrology of the Parker-Blythe-Cibola area, Arizona and California: U.S. Geological Survey Professional Paper 486-G, p. G1-G130.
- Olmsted, F. H., Loeltz, O. J., and Irelan, Burdge, 1973, Geohydrology of the Yuma area, Arizona and California: U.S. Geological Survey Professional Paper 468-H, p. H1-H227.
- Oppenheimer, J. M., and Sumner, J. S., 1980, Depth-to-bedrock map of southern Arizona: University of Arizona Laboratory of Geophysics Department of Geosciences Map 6, scale 1:500,000.
- Peirce, H. W., 1976, Elements of Paleozoic tectonics in Arizona: Arizona Geological Society Digest, v. 10, p. 37-57.

- Reed, J. E., 1985, Mathematical model for simulating ground-water and heat flow, *in* Bedinger, M. S., Sargent, K. A., and others, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of radioactive waste--Basis of characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].
- Reeves, C. C., Jr., 1966, Pleistocene climate of the Llano Estacado, Part 2: *Journal of Geology*, v. 74, no. 1, pt., 1, p. 642-647.
- Shafiqullah, Muhammad, Damon, P. E., Lynch, D. J., Reynolds, S. J., Rehrig, W. A., and Raymond, R. H., 1980, K-Ar geochronology and geologic history of southwestern Arizona and adjacent areas, *in* Jenney, J. P., and Stone, Claudia, eds., Studies in western Arizona: Arizona Geological Society Digest, v. 12, p. 201-260.
- Snyder, C. T., and Langbein, W. B., 1962, Pleistocene lake in Spring Valley, Nevada, and its climatic implications: *Journal of Geophysical Research*, v. 67, no. 6, p. 2385-2394.
- Spaulding, W. G., 1983, Late Wisconsin macrofossil records of desert vegetation in the American southwest: *Quaternary Research*, v. 19, no. 2, p. 256-264.
- Thompson, T. H., Nuter, J. A., and Anderson, T. W., 1984, Maps showing distribution of dissolved solids and dominant chemical type in ground water, Basin and Range province, Arizona: U.S. Geological Survey Water-Resources Investigations WRI-83-4114-C, scale 1:500,000.

MINERAL AND ENERGY RESOURCES

By

B. T. Brady, U.S. Geological Survey

and

Robert B. Scarborough,

Arizona Bureau of Geology and Mineral Technology

The Sonoran region of Arizona contains mineral deposits which occur in diverse geologic environments and are of several ages. Historically, base- and precious-metal ores have been the principal sources of metals in the region. Of these commodities copper, molybdenum, silver and gold are currently the most important to the State's mineral economy. Early mineral development in the Sonoran region of Arizona was confined to precious-metal lodes, base-metal vein and replacement ores, and placer deposits. These deposits with few exceptions were extensively developed by the close of the 19th century. More recent development has been focused on large scale open-pit copper mines which produce substantial quantities of precious metals as coproducts or byproducts.

Metallic-Mineral Resources

The mineralized areas discussed in this report have been delineated by Keith, Gest, and DeWitt (1983), and Keith, Schnabel, DeWitt, Gest, and Wilt (1983). These areas are characterized as unique metallogenic systems, which have been defined on the basis of mineralization age and style. These areas are not all inclusive, however, they generally comprise areas of identified resources with documented production. This report is not meant to contain an assessment of areas of mineral potential. Commodities of primary importance in the mineralized areas are metallic minerals that contain any of the following elements: copper, lead, zinc, molybdenum, iron, manganese, barium, gold, silver, mercury, platinum, lithium, beryllium, tin, tungsten, and uranium.

The Sonoran region of Arizona contains more than 115 metallic-mineral districts (Keith, Gest, and DeWitt, 1983; Keith, Schnabel, DeWitt, Gest, and Wilt, 1983), which have individually and collectively contributed substantially to the total State non-fuel mineral production. Boundaries of these metallic-mining districts (plate 5) represent the general extent of known productive workings and prospects, and do not indicate limits of mineralized rock. These predominantly base- and precious-metal ores have been mined from deposits of various forms and ages. The principal production in the area to date has been from disseminated and stockwork porphyry deposits, vein and replacement deposits, and to a lesser extent, from stratabound deposits. The principal base- and precious-metal production for the metallic-mineral districts that have produced more than 90,780 Mg of ore are summarized in table 3.

Table 3.--Production summary of base and precious metals for metallic-mineral districts with total production in excess of 100,000 short tons, by county (including estimates for years data not available). Commodities listed in other commodity column are abbreviated as follows: Mn, manganese; Mo, molybdenum; W, tungsten [Data from Keith, Gest, DeWitt, Toll, and Everson (1983) unless otherwise noted].

Metallie-mineral district	Production period	Ore (thousands of short tons)	Copper (thousands of lbs)	Lead (thousands of lbs)	Zinc (thousands of lbs)	Gold (ozs)	Silver (hundreds of ozs)	Other commodity, (thousands of lbs) unless noted
LaPaz County								
Little Harquahala	1888-1963	159	50	156	-	143,000	900	-
Silver	1880-1951	103	2	2,456	-	<100	13,110	-
Mohave County								
Artillery Peak	1946-59	-	-	-	-	-	-	95,108(Mn) ⁽¹⁾
Boriana	1919-57	239	408	-	-	100	125	>121,324(W) ⁽²⁾
Emerald Isle	1917-73	1,418	22,166	-	-	<100	4	-
Hualapai	1918-70	161	7,247	897	11,370	700	990	-
Mc Cracken	1911-81	173	10	3,031	43	100	6,990	100(Mo) ⁽³⁾
Oatman	1870-1980	4,073	60	-	-	1,966,000	11,470	-

Table 3.--Production summary of base and precious metals for metallic-mineral districts with total production in excess of 100,000 short tons, by county (including estimates for years data not available)--Continued

Metallie-mineral district	Production period	Ore (thousands of short tons)	Copper (thousands of lbs)	Lead (thousands of lbs)	Zinc (thousands of lbs)	Gold (ozs)	Silver (hundreds of ozs)	Other (commodity, thousands of lbs) unless noted
Mohave County Continued								
Pilgrim	1929-45	281	-	-	-	48,000	720	-
Planet	1862-1970	1,010	19,520	-	-	400	3	-
Swansea	1909-62	545	26,457	-	-	500	330	-
Union Pass	1868-1943	704	-	-	-	128,000	3,130	-
Wallapai	1901-81	96,920	666,137	80,096	126,487	151,000	115,350	53,180 (Mo)
Pima County								
Ajo	1899-1979	430,038	6,026,380	30	-	1,558,000	196,720	450 (Mo)

Table 3.--Production summary of base and precious metals for metallic-mineral districts with total production in excess of 100,000 short tons, by county (including estimates for years data not available)--Continued

Metallic-mineral district	Production period	Ore (thousands of short tons)	Copper (thousands of lbs)	Lead (thousands of lbs)	Zinc (thousands of lbs)	Gold (ozs)	Silver (hundreds of ozs)	Other commodity, (thousands of lbs) unless noted
Yavapai County								
Copper Basin	1901-68	357	19,631	508	1,400	500	457	-
Eureka (includes 1880 estimate from Hillside mine)	1890-1981	101,417	1,306,111	7,866	3,624	67,000	46,910	16,535(Mo) 46(Mn) 116(U ₃ O ₈), (3) 13(V ₂ O ₅), (3)
Martinez (Congress mine)	1887-1950	990	201	2	-	432,500	4,660	-
Old Dick	1917-77	1,683	106,396	3,041	306,584	3,500	6,520	-
Rich Hill	1895-1970	745	124	882	-	203,000	1,890	-

Table 3.--Production summary of base and precious metals for metallic-mineral districts with total production in excess of 100,000 short tons, by county (including estimates for years data not available)--Continued

Metallie-mineral district	Production period	Ore (thousands of short tons)	Copper (thousands of lbs)	Lead (thousands of lbs)	Zinc (thousands of lbs)	Gold (ozs)	Silver (hundreds of ozs)	Other (commodity, thousands of lbs) unless noted
Yuma County								
Castle Dome	1870-1900	4		7,350			2,480	
	1884-1900				3,617			
	1901-81	120	53	11,582	29	3,000	2,540	-
Kofa	1897-1957	675	7	2	-	237,000	1,070	-

(1) Area extends southward into La Paz County

(2) Production in short-ton units. A short-ton unit of WO₃ is 1 percent of a short ton or 20 pounds of WO₃, and contains 15.86 pounds of tungsten metal

(3) Production in pounds

(4) Placer production estimated from Tenney, J. B., in Wilson (1933)

Copper has been noted in the majority of the mineralized areas within the Sonoran region of Arizona (Anderson, 1969; Keith, Gest, and DeWitt, 1983; Keith, Schnabel, DeWitt, Gest, and Wilt, 1983), and the principal sources of this metal in the study area are the mines in the Eureka, Ajo, and Wallapai mining districts. Eighteen of the nations 25 leading copper-producing mines are located in Arizona. During 1981 the Eureka, Ajo, and Wallapai districts ranked 9th, 17th, and 23rd, respectively, in total domestic copper production (Butterman, 1983).

Production of copper from the disseminated and stockwork deposits in the Eureka (Bagdad) district during 1981 was 64,869 Mg (Burgin, 1983), and large quantities of molybdenum and gold are recovered annually as byproducts. Base- and precious-metal veins and massive replacement deposits also have been important metal sources in the district. Mineralization at Eureka was related to the intrusion of a Laramide quartz-monzonite stock. The geologic history and character of the mineral deposits of the Eureka district is complex and has been described in detail by Butler (1938), Anderson (1948, 1950a, 1950b), and Anderson and others (1955).

The New Cornelia disseminated porphyry copper ore body in the Ajo district occurs in the brecciated Laramide Cornelia quartz monzonite, and to a smaller extent in the surrounding quartz diorite facies and Cretaceous(?) Concentrator volcanics. Copper, substantial quantities of byproduct silver and gold, and lesser quantities of molybdenum constitute the primary metals produced since the inception of large-scale mining about 1917. The cumulative base- and precious-metal production from the New Cornelia open-pit mine exceeded \$1.5 billion by 1972 (Keith, 1974). Discussions of the geology and mineral deposits near Ajo are contained in articles by Joralemon (1914), Bryan (1925), Ingham and Barr (1932), Gilluly (1937, 1938, 1942, 1946, 1952), and Dixon (1966).

The Wallapai mining district is located in the north-trending Cerbat Mountains in northern Mohave County. Isoclinally folded schist, gneiss, and amphibolite of the Precambrian Cerbat complex are locally intruded by Precambrian granite and a compositionally and texturally zoned Laramide quartz-monzonite stock. Mineralization in the district is of multiple ages and types. An early stage of stockwork-copper deposits formed mainly within predominately northwest-trending fractures in the brecciated (Mesozoic(?) Ithaca Peak Granite (Eidel and others, 1968). Northwest-trending, lead-zinc vein deposits overlie and are peripheral to the porphyry-copper deposits (Thomas, 1949). Tertiary supergene turquoise deposits that occur superimposed above the stockwork deposits, and Quaternary supergene(?) - chrysocolla deposits which formed at Emerald Isle, are the results of late-stage oxidation and mineralization in the Wallapai district. Initially the principal production was from vein deposits, whereas more recent development has been confined to the porphyry-copper deposits. The cumulative total ore production from the Wallapai district through 1981 has been estimated to exceed 87,148 Mg (Keith, Gest, DeWitt, Toll, and Everson, 1983).

Eleven of the 46 mineral districts in Arizona that have produced more than 10,000 oz of gold occur within the Sonoran region of Arizona study area. These include the Ajo, Cienega, Eureka, Kofa, Little Harquahala, Martinez, Oatman, Pilgrim, Rich Hill, Union Pass and Wallapai districts. Collectively they have produced nearly 5 million oz of gold (Koschman and Bergendahl, 1968; Keith, Gest, Dewitt, Toll, and Everson, 1983). The majority of the gold produced in Arizona prior to 1900 came from lode deposits of various mineralization ages and styles, and placer deposits, whereas the majority of the State's gold mined during the 20th century has been produced as a byproduct of large-scale, low-grade copper mining. Arizona ranked fourth among the Nation's leading gold producers during 1981 (Lucas, 1982).

Lead and zinc minerals occur primarily in vein and replacement deposits in the Sonoran region of Arizona. Lead minerals commonly contain substantial quantities of silver, and historically lead-bearing ores have been mined as a source of silver. The most significant production of lead has come from deposits in the Wallapai district, Mohave County, and the Eureka area in Yavapai County. Smaller, although locally important tonnages of lead, were mined from vein deposits in the Castle Dome district, Yuma County, and in the Mc Cracken district, Mohave County. The Mc Cracken district deposits were reworked substantially in 1981 (Keith, Gest, DeWitt, Toll, and Everson, 1983). More than one-half of the lead produced in Arizona during 1981 came from large copper mines (Burgin, 1983). The Mc Cracken district and Mineral Park area in the Wallapai district ranked 1st and 6th, respectively in Arizona lead output during 1981.

The Old Dick and Copper Basin districts in Yavapai County, and the Wallapai and Hualapai districts in Mohave County, have been the principal sources of zinc in the Arizona Sonoran region. Zinc was recovered at the Mineral Park area in the Wallapai district, and from a few base- and precious-metal mines in the study area during 1981 (Burgin, 1983).

Silver and molybdenum have been recovered in significant quantities as byproducts of open-pit copper mining at the Mineral Park area in the Wallapai district and in the Eureka district. The New Cornelia mine in the Ajo district also has produced substantial silver and small quantities of molybdenum. Furthermore, silver has been mined from important precious-metal lode and placer deposits, and base-metal fissure veins in the study area. The most notable of these areas, which have produced more than 500,000 oz of silver, include the Silver, Oatman, Mc Cracken, Old Dick, Hackberry, and Castle Dome districts. Arizona mines accounted for 20 percent of the 1981 domestic silver production, and 30 percent of the Nation's total molybdenum output (Burgin, 1983).

A large domestically important manganese resource occurs in the Artillery Peak district. This syngenetic deposit occurs in the upper few hundred meters of clastic sediments in the Miocene Chapin Wash Formation. Two horizons of mineralization have been noted, and the upper zone has been the principal development target to date. An estimated 337,702 Mg of ore were mined from the shallow manganese-bearing horizon for government stockpiles during 1951-56 (Farnham and Stewart, 1958). This area contains more than 159 million Mg of manganese ore which averages 3.5 to 4 percent manganese (Lasky and Webber, 1949). A smaller deposit of manganese oxide similar to the syngenetic ores in the Artillery Mountains occurs at the Doyle mine in the Alamo district. Several million pounds of manganese also have been produced from deposits in the Lincoln Ranch and Bouse districts in LaPaz County, and the Trigo Mountains district in Yuma County (Keith, Gest, DeWitt, Toll and Everson, 1983). Numerous other scattered occurrences of manganese occur in western Arizona as epigenetic fissure fillings and irregular replacement deposits. These deposits commonly are of low grade and usually are associated with base metals, which decrease the commercial value of the ore.

Tungsten has been mined from deposits in 10 metallic-mineral districts in the Sonoran region of Arizona (Wilson, 1941b).

(A short-ton unit of tungsten equals 1 percent per ton or 20 pounds WO_3 , which contains 15.8614 pounds of tungsten metal.)

Although tungsten occurs in several modes, irregular vein and replacement deposits in silicic plutonic and metamorphic rocks are the principal sources of this metal in Arizona. Wolframite- and scheelite-bearing quartz veins in the Borianna district have yielded more than 133,735 Mg units of tungsten (Keith, Gest, DeWitt, Toll and Everson, 1983). A few thousand megagram units of tungsten were mined in the Tungstona, Wheeler Wash and Aquarius Mountains districts, and smaller tonnages were produced from mines in the Ellsworth, Kofa, Fluorescent, Zannaropolis, Ophir, and Lost districts.

A few hundred flasks of mercury were produced from a narrow brecciated quartz vein in the La Cholla district in the Dome Rock Mountains prior to 1914 (Lausen and Gardner, 1927; Wilson, 1941a; and Bailey, 1969).

Within the region, three areas have been intensively explored and drilled for uranium. The Date Creek basin contains reserves of 18 million pounds of U_3O_8 in a Miocene carbonaceous lacustrine sequence within 200 m of the surface (Sherborne and others, 1979), and very large potential resources exist in deeper parts of the basin (Otton, 1981; May and others, 1982). Similiar aged sediments in the Big Sandy River valley 10 to 30 km north of Wikieup contains several low-grade uraniferous zones, buried beneath shallow less-deformed gravels throughout an area of several square kilometers (Scarborough, 1981). Numerous vein and fissure uranium occurrences, some with thorium, are known in the Cerbat Mountains north of Kingman, and are related to base- and precious-metal mineralization in that area (Scarborough, 1981). Of these three areas, minor uranium production is recorded only from the Date Creek basin.

Although many other metallic minerals have been reported to occur sporadically in the mineralized areas of the Sonoran region of Arizona, they are chiefly of mineralogical importance, and will not be discussed herein.

Nonmetallic and Industrial Mineral Resources

Nonmetallic and industrial minerals are of widespread occurrence in the Sonoran region of Arizona. However, these materials have been produced intermittently primarily for local use. More than 28 varieties of nonmetallic industrial minerals or rocks are reported to occur within the study area. The distribution and nature of these minerals and rocks have been shown or described in publications by McCrory and O'Haire (1961), Wilson and Roseveare (1949), U.S. Geological Survey, Arizona Bureau of Mines, and U.S. Bureau of Reclamation (1969), and Elevatorski (1975). The occurrences of semi-precious or precious gemstones, ornamental and building stone, and natural aggregates will not be described in this report. The nonmetallic mineral commodities, which have generally small production, include: barite, fluorspar, bentonite, kaolinite, feldspar, mica, gypsum, halite, brucite, and magnesite.

Approximately 2,723 Mg of barite worth nearly \$27,000 were produced collectively from veins in faulted Precambrian granite, schist, and gneiss, and to a lesser extent from fissure fillings in Tertiary volcanic rocks in the Harcuvar, Plomosa Pass, and Mohawk districts (Keith, 1978). Small quantities of barite also have been mined from deposits north of the Bouse and Three-In-One districts in Mohave County (Elevatorski, 1975). Barite also occurs as a common gangue mineral in many of the vein and replacement deposits in the Sonoran region of Arizona.

Argentiferous galena-fluorite veins in the Castle Dome district yielded nearly 2,723 Mg of metallurgical-grade fluorspar during 1902-17 and in 1953 (Wilson, 1950; and Van Alstine and Moore, 1969). These veins comprise north-to northwest-trending fissure filling deposits that crosscut Mesozoic shales and diorite and rhyolite porphyry dikes (Wilson, 1933). Fluorite localities are common in the study area. The mineral usually is associated with barite, calcite or quartz, manganese oxides, and to a lesser extent with beryllium, tungsten, and uranium minerals in several precious- and base-metal vein deposits.

A few hundred megagrams of bentonite were mined for drilling mud near Bouse (Wilson and Roseveare, 1949), and small quantities of lithium-bearing montmorillonite have been produced for a sealer from the Lyles clay deposit 14.5 km northwest of Kirkland (Norton, 1965). An estimated 1 million Mg of bentonite was reported to occur within the smectite deposit along Burro Creek 12.8 km west of Bagdad (Funnell and Wolfe, 1964). The majority of these deposits of bentonite probably formed by alteration of Cenozoic volcanic ash, or tuffaceous or lacustrine sediments. However, the genesis of these occurrences has not been fully documented. Kaolinite associated with quartz and cristobalite occurs in altered volcanic rocks at the Klaner and Doolin prospect near the southwest end of Black Mesa, Mohave County (Elevatorski, 1975). Although kaolinite has been intermittently mined as a refractory material from a few small deposits, no production data are currently available.

Pegmatites at several localities in the region have been mined sporadically to acquire materials for abrasives, ceramics, glass, enamels, drilling mud, and local smelter flux (Elevatorski, 1975).

Gypsum has been extracted for agricultural use from small deposits in Permian limestone at the Blue Moon and Townsend properties near Bouse (Elevatorski, 1975).

The principal occurrences of halite in the Arizona Sonoran region are located in the Hualapai Valley and in the Detrital Valley in central Mohave County. The Detrital Valley bedded-salt deposit attains a maximum thickness of 218 m and is associated with gypsum- and anhydrite-bearing clastic sediments of the Miocene Muddy Creek Formation (Peirce, 1981b). The massive body of halite at Red Lake in the Hualapai Valley is more than 1,000 m thick. Southwest Gas Co., currently plans to utilize this salt deposit for storage of natural gas (Peirce, 1981a). Subsidence due to solution of salt has not been recognized to date in the vicinity of either salt body.

A potential brucite resource in the Tertiary volcanic rocks in the Oatman district reportedly contains more than 9,078 Mg of low-grade magnesium oxide (Gildersleeve, 1962; Ericksen, 1969). The brucite-bearing horizons, which contain abundant calcite, dolomite, and lesser quantities of magnesite and hydromagnesite, are parallel to the enclosing volcanic rocks, and locally attain a maximum composite thickness of 9 m. Although a small quantity of brucite was mined in 1953, the low grade of the material prevented sustained shipments.

Nonmetallic and industrial rocks and minerals occur throughout the region. However, these commodities have been developed primarily for local markets, and there has been no sustained production for any of these commodities described herein.

Geothermal Resources

Four areas with potential for discovery or development of low-temperature geothermal waters (less than 100 °C) have been identified in the Sonoran region of Arizona (Witcher and others, 1982). These areas, about 104 to 777 km², are located in the southern Sacramento Valley near Yucca, west of the Aquarius Mountains in the Big Sandy River valley, along the Santa Maria River east of Alamo Crossing, and in the Ranegras Plain west of Vicksburg. Water in three thermal wells in the study area has a temperature that exceeds 50 °C. Only 1 well with a water temperature of 89 °C is in an area with a gradient greater than 15 °C/km. The majority of the geothermal wells are shallow and produce water with a temperature less than 50 °C. There are few thermal springs in the region, none of which have temperatures in excess of 50 °C. There are no known Geothermal Resources Areas in the Sonoran region of Arizona (Witcher and others, 1982).

Coal, Oil, and Gas Resources

No coal occurrences have been identified in the Sonoran region of Arizona. A few wildcat boreholes have been drilled in the study area (Conley and Stacey, 1981; Brady, 1983). However, no production has been recorded to date. The potential for oil and gas discoveries in southwest Arizona has been described by McCarthy (1961) and Peirce and others (1970), and for the recently publicized overthrust oil and gas reserves, by Anschutz (1980) and Reif and Robinson (1981). Oil and gas potential is unknown except perhaps for basin-fill sand along the lower Colorado River trough, as in such basins as the Yuma basin, where leasing is ongoing after a commercial-gas well drilled by Pemex in the northwest part of Sonora, Mexico, near the mouth of the Colorado River (World Oil, 1981; Davis, 1981). There, Pemex announced a pre-production flow of 5.7 million m³/d of gas from a depth of 4,115 m in Miocene and Pliocene sands. These sands may extend northward, thinning toward the Yuma area.

References Cited

- Anschutz, P. F., 1980, The overthrust belt--Will it double U.S. gas reserves?: World Oil, January, p. 111-116.
- Anderson, C. A., 1948, Structural control of copper mineralization, Bagdad, Arizona: American Institute of Mining and Metallurgical Engineering, Technical Publication 2352, Mining Technology, v. 12, no. 2, 11 p.
- 1950a, Lead-zinc deposits, Bagdad area, Yavapai County, Arizona, in Arizona zinc and lead deposits, Part 1: Arizona Bureau of Mines Bulletin 156, p. 122-138.
- 1950b, Alteration and metallization in the Bagdad porphyry copper deposit, Arizona: Economic Geology, v. 45, no. 7, p. 609-628.
- 1969, Copper, in Mineral and water resources of Arizona: Arizona Bureau of Mines Bulletin 180, p. 117-156.
- Anderson, C. A., Scholz, E. A., and Strobell, J. D., Jr., 1955, Geology and ore deposits of the Bagdad area, Yavapai County, Arizona: U.S. Geological Survey Professional Paper 278, 103 p.
- Bailey, E. H., 1969, Mercury, in Mineral and water resources of Arizona: Arizona Bureau of Mines Bulletin 180, p. 226-230.
- Brady, B. T., 1983, Map showing coal, oil and gas wells, seeps, and tar sandstone occurrences in the Basin and Range province: U.S. Geological Survey Open-File Report 83-549, 104 p.
- Bryan, Kirk, 1925, The Papago country, Arizona: U.S. Geological Survey Water-Supply Paper 499, 436 p.

- Burgin, L. B., 1983, The mineral industry of Arizona: U.S. Bureau of Mines Minerals Yearbook, v. 2, p. 55-73.
- Butler, B. S., 1938, Bagdad mine, Eureka district: Arizona Bureau of Mines Bulletin 145, p. 98-103.
- Butterman, W. C., 1983, Copper: U.S. Bureau of Mines Minerals Yearbook, v. 1, p. 279-307.
- Conley, J. N., and Stacey, D. A., 1981, Well location map for State of Arizona: Arizona Oil and Gas Conservation Commission, sheet 1, scale 1:500,000; sheet 2, scale 1:126,720; supplement, 23 p.
- Davis, Ben, 1981, Arizona discovery race reaching home stretch: Western Oil Reporter, v. 38, no. 11, p. 67.
- Dixon, D. W., 1966, Geology of the New Cornelia mine, Ajo, Arizona, in Titley, S. R., and Hicks, C. L., eds., Geology of the porphyry copper deposits southwestern North America: Tucson, University of Arizona Press, p. 123-132.
- Eidel, J. J., Frost, J. E., and Clippinger, D. M., 1968, Copper-molybdenum mineralization at Mineral Park, Mohave County, Arizona, in Ridge, J. D., ed., Ore deposits of the United States, 1933-1967; The Graton-Sales volume: New York, American Institute of Mining, Metallurgical, and Petroleum Engineers, v. 2, p. 1258-1281.
- Elevatorski, E. A., 1975, Arizona industrial minerals: Santa Ana, California, MINOBRAS, 73 p.
- Ericksen, G. E., 1969, Brucite and magnesite, in Mineral and water resources of Arizona: Arizona Bureau of Mines Bulletin 180, p. 320-324.

- Farnham, L. L., and Stewart, L. A., 1958, Manganese deposits of western Arizona: U.S. Bureau of Mines Information Circular 7843, 87 p.
- Funnell, J. E., and Wolfe, E. J., 1964, Compendium on nonmetallic minerals of Arizona: San Antonio, Texas, Southwest Research Institute, 374 p.
- Gildersleeve, Benjamin, 1962, Magnesite and brucite in the United States: U.S. Geological Survey Minerals Investigations Resource Map MR-27, scale 1:3,168,000.
- Gilluly, James, 1937, Geology and ore deposits of the Ajo Quadrangle, Arizona: Arizona Bureau of Mines Geological Series 9, Bulletin 141, 83 p.
- 1938, Ajo District: Arizona Bureau of Mines Geological Series 12, Bulletin 145, p. 86-90.
- 1942, The mineralization of the Ajo copper district, Arizona: Economic Geology, v. 37, no. 4, p. 257-309.
- 1946, The Ajo mining district, Arizona: U.S. Geological Survey Professional Paper 209, 112 p.
- 1952, Geology of the New Cornelia Mine, Ajo, Arizona, in Economic geology--Ajo porphyry copper, Trip 4, of the Arizona Geological Society guide book for field trip excursions in southern Arizona: Geological Society of America, Cordilleran Section, Meeting, April 10-14, 1952, Tucson, Arizona, University of Arizona, p. 57-61.
- Ingham, G. R., and Barr, A. T., 1932, Mining methods and costs at the New Cornelia Branch, Phelps Dodge Corporation, Ajo, Arizona: U.S. Bureau of Mines Information Circular 6666, 17 p.

- Joralemon, I. B., 1914, The Ajo copper mining district: American Institute of Mining Engineering Bulletin, no. 92, p. 2011-2028.
- Keith, S. B., 1974, Index of mining properties in Pima County, Arizona: Arizona Bureau of Mines Bulletin 189, 156 p.
- 1978, Index of mining properties in Yuma County, Arizona: Arizona Bureau of Geology and Mineral Technology Bulletin 192, 185 p.
- Keith, S. B., Gest, D. E., and DeWitt, Ed, 1983, Metallic mineral districts of Arizona: Arizona Bureau of Geology and Mineral Technology map, scale 1:1,000,000.
- Keith, S. B., Gest, D. E., DeWitt, Ed, Toll, Netta, and Everson, B. A., 1983, Metallic mineral districts and production in Arizona: Arizona Bureau of Geology and Mineral Technology Bulletin 194, map scale 1:1,000,000, 58 p.
- Keith, S. B., Schnabel, Lorraine, DeWitt, Ed, Gest, D. E., and Wilt, J. C., 1983, Map and bibliography of the mineralized areas of the Basin and Range province in Arizona: U.S. Geological Survey Open-File Report 84-86, map scale 1:500,000, 81 p.
- Koschman, A. H., and Bergendahl, M. H., 1968, Principal gold producing districts of the United States: U.S. Geological Survey Professional Paper 610, 283 p.
- Lasky, A. G., and Webber, B. N., 1949, Manganese resources of the Artillery Mountains region, Mohave County, Arizona: U.S. Geological Survey Bulletin 936-R, p. 417-448.
- Lausen, Carl, and Gardner, E. D., 1927, Quicksilver resources of Arizona: Arizona Bureau of Mines Bulletin 182, 112 p.
- Lucas, J. M., 1982, Gold, in Minerals yearbook, 1981: U.S. Bureau of Mines, v. 1, p. 365-391.

- May, R. T., White, D. L., and Nystrom, R. J., 1982, National uranium resource evaluation, Prescott Quadrangle, Arizona: U.S. Department of Energy Report GJQ-015(82), 62 p.
- McCarthy, D. P., 1961, Geological evaluation of oil and gas possibilities in southwestern Arizona, in Oil, gas, and helium in Arizona: Arizona Development Board, Arizona Oil and Gas Conservation Commission, p. 30-34.
- McCrary, F. J., and O'Haire, R. T., 1961, Map of known nonmetallic mineral occurrences of Arizona: Arizona Bureau of Mines, scale 1:1,000,000.
- Norton, J. J., 1965, Lithium-bearing bentonite deposit, Yavapai County, Arizona: U.S. Geological Survey Professional Paper 525-D, p. D163-D166.
- Otton, J. K., 1981, Geology and genesis of the Anderson Mine, a carbonaceous lacustrine uranium deposit, western Arizona-- A summary report: U.S. Geological Survey Open-File Report 81-780, 24 p.
- Peirce, H. W., 1981a, Natural gas storage in Arizona salt: Arizona Bureau of Geology and Mineral Technology Fieldnotes, v. 11, no. 3, p. 8.
- 1981b, Major Arizona salt deposits: Arizona Bureau of Geology and Mineral Technology Fieldnotes, v. 11, no. 4, p. 1-4.
- Peirce, H. W., Keith, Stanley B., and Wilt, J. C., 1970, Coal, oil, natural gas, helium, and uranium in Arizona: Arizona Bureau of Mines Bulletin 182, 289 p.

- Reif, D. M., and Robinson, J. P., 1981, Geophysical, geochemical, and petrographic data and regional correlation from the Arizona State A-1 Well, Pinal County, Arizona: Arizona Geological Society Digest, v. 13, p. 99-110.
- Scarborough, R. B., 1981, Radioactive occurrences and uranium production in Arizona: U.S. Department of Energy Report GJBX-143(81), 297 p.
- Sherborne, J. E., Buckovic, W. A., and Dewitt, D. B., 1979, Major uranium discovery in volcanoclastic sediments, Basin and Range province, in Yavapai County, Arizona: American Association of Petroleum Geologists Bulletin, v. 63, no. 4, p. 621-646.
- Thomas, B. E., 1949, Ore deposits of the Wallapai district, Arizona: Economic Geology, v. 44, no. 8, p. 663-705.
- U.S. Geological Survey, Arizona Bureau of Mines, and U.S. Bureau of Reclamation, 1969, Mineral and water resources of Arizona: Arizona Bureau of Mines Bulletin 180, 638 p.
- Van Alstine, R. E., and Moore, R. T., 1969, Fluorspar, in Mineral and water resources of Arizona: Arizona Bureau of Mines Bulletin 180, p. 348-357.
- Wilson, E. D., 1933, Geology and mineral deposits of southern Yuma County, Arizona: Arizona Bureau of Mines Bulletin 134, Geological Series 7, v. 4, no. 2, 236 p.
- 1941a, Quicksilver (mercury): Arizona Bureau of Mines Circular 9, 13 p.
- 1941b, Tungsten deposits of Arizona: Arizona Bureau of Mines Bulletin 148, Geological Series 14, v. 12, no. 2, 54 p.
- 1950, Fluorspar in Arizona: Arizona Bureau of Mines Circular 15, 13 p.

Wilson, E. D., and Roseveare, G. H., 1949, Arizona nonmetallics,
(2d ed.): Arizona Bureau of Mines Bulletin 155, 60 p.

Witcher, J. C., Stone, Claudia, and Hahman, R. W., Sr., 1982,
Geothermal resources of Arizona: Arizona Bureau of Geology
and Mineral Technology, scale 1:500,000.

World Oil, 1981, Mexico, v. 193, no. 1, p. 84.