


296
84-743

STUDIES OF GEOLOGY AND HYDROLOGY IN THE
BASIN AND RANGE PROVINCE, SOUTHWESTERN UNITED STATES,
FOR ISOLATION OF HIGH-LEVEL RADIOACTIVE WASTE

CHARACTERIZATION OF THE DEATH VALLEY REGION,
NEVADA AND CALIFORNIA

U.S. Geological Survey Open-File Report 84-743



Open-file report
Geological Survey
(U.S.)

In press as Professional Paper 1370-F

This is Chapter F of an eight-chapter Professional Paper series
being prepared by the U.S. Geological Survey in cooperation with
States in the Basin and Range province.

Tw anal

**STUDIES OF GEOLOGY AND HYDROLOGY IN THE
BASIN AND RANGE PROVINCE, SOUTHWESTERN UNITED STATES,
FOR ISOLATION OF HIGH-LEVEL RADIOACTIVE WASTE**

**CHARACTERIZATION OF THE DEATH VALLEY REGION,
NEVADA AND CALIFORNIA**

Chairman of the Province Working Group:
edited by M.S. Bedinger, K.A. Sargent, and William H. Langer



U.S. Geological Survey Open-File Report 84-743



Member:
K.A. Sargent
Geologist
U.S. Geological Survey
Denver, CO

Member:
Larry D. Fellows
State Geologist and Assistant
Arizona Bureau of Geology and
Tucson, AZ

ARIZONA

Member:
Larry D. Fellows
State Geologist and Assistant
Arizona Bureau of Geology and
Tucson, AZ

Member:
Larry D. Fellows
State Geologist and Assistant
Arizona Bureau of Geology and
Tucson, AZ

Alternate:
H. Wesley Pearce
Principal Geologist
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

Alternate:
H. Wesley Pearce
Principal Geologist
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

Alternate:
H. Wesley Pearce
Principal Geologist
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

Alternate:
H. Wesley Pearce
Principal Geologist
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

CALIFORNIA

Member:
Robert Streitz
Geologist
California Division of Mines and Geology
Sacramento, CA

Member:
Robert Streitz
Geologist
California Division of Mines and Geology
Sacramento, CA

Member:
Robert Streitz
Geologist
California Division of Mines and Geology
Sacramento, CA

In press as Professional Paper 1370-F

This is Chapter F of an eight-chapter Professional Paper series
being prepared by the U.S. Geological Survey in cooperation with
States in the Basin and Range province.

UNITED STATES
DEPARTMENT OF THE INTERIOR

DONALD PAUL HODEL, Secretary

GEOLOGICAL SURVEY

Dallas L. Peck , Director



Copies of this report can be purchased from:

Open-File Services Section
Western Distribution Branch
P.O. Box 25425, Denver Federal Center
Denver, Colorado 80225

BASIN AND RANGE PROVINCE WORKING GROUP

U.S. Geological Survey Members

Chairman of the Province Working Group:

M.S. Bedinger
Hydrologist
U.S. Geological Survey
Denver, CO

Member:

K.A. Sargent
Geologist
U.S. Geological Survey
Denver, CO

State Members and Alternates

ARIZONA

Member:

Larry D. Fellows
State Geologist and Assistant Director
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

Alternate:

H. Wesley Peirce
Principal Geologist
Arizona Bureau of Geology and Mineral Technology
Tucson, AZ

CALIFORNIA

Member:

Robert Streitz
Geologist
California Division of Mines and Geology
Sacramento, CA

NEW MEXICO

Member:

James M. Hill
Chief, Bureau of Geology
New Mexico Energy and Minerals Department
Santa Fe, NM

Alternate:

Frank E. Kottlowski
Director
New Mexico Bureau of Mines and Mineral Resources
Socorro, NM

TEXAS

Member:

Christopher D. Henry
Geologist
Texas Bureau of Economic Geology
University of Texas at Austin
Austin, TX

Alternate:

Douglas Ratcliff
Associate Director
Texas Bureau of Economic Geology
University of Texas at Austin
Austin, TX

UTAH

Member:

Genevieve Atwood
State Geologist
Utah Geological and Mineral Survey
Salt Lake City, UT

Alternate:

Don R. Mabey
Senior Geologist
Utah Geological and Mineral Survey
Salt Lake City, UT

Copies of this report can be purchased

Open-File Services Section
Western Distribution Branch
P.O. Box 25425, Denver Federal Center
Denver, Colorado 80225

IDAHO

CONTENTS

Member:

Frank B. Sherman
Chief, Ground-Water Section
Idaho Department of Water Resources
Boise, ID

Alternate:

Darrel Clapp
Chief, Technical Services Bureau
Idaho Department of Water Resources
Boise, ID

NEVADA

Member:

John Schilling
State Geologist
Nevada Bureau of Mines and Geology
University of Nevada, Reno
Reno, NV

Alternate:

Susan L. Tingley
Deputy to the State Geologist
Nevada Bureau of Mines and Geology
University of Nevada, Reno
Reno, NV

CONTENTS

	Page
Abstract-----	1
Introduction, by M.S. Bedinger and K.A. Sargent-----	4
Background and purpose-----	4
Geographic setting-----	7
Acknowledgments-----	10
References cited-----	11
Geology, by T.L.T. Grose and George I. Smith-----	13
Early and Middle Proterozoic crystalline basement rocks-----	13
Middle and late Proterozoic sedimentary rocks-----	18
Latest Proterozoic and Lower Cambrian clastic rocks-----	19
Middle Cambrian through Permian carbonate and clastic rocks-----	22
Triassic and Jurassic sedimentary and volcanic rocks-----	25
Cretaceous through middle Eocene sedimentary rocks-----	27
Upper Eocene to Holocene sedimentary and volcanic rocks-----	27
Mesozoic and Cenozoic plutonic rocks-----	32
Structural and tectonic features-----	34
Geomorphology-----	46
References cited-----	48

	Page
Potential host media for radioactive waste, by K.A. Sargent-----	64
Intrusive rocks-----	65
Tuffaceous rocks-----	71
Basaltic rocks-----	73
Argillaceous rocks-----	75
Unsaturated zone-----	76
References cited-----	77
Quaternary tectonism, by K.A. Sargent and T.L.T. Grose---	79
Seismicity-----	81
Heat flow-----	84
Quaternary faulting-----	86
Late Cenozoic volcanics-----	87
Vertical crustal movement-----	88
Photolineations-----	88
References cited-----	89
Ground-water hydrology, by M.S. Bedinger, William H. Langer, and J.E. Reed-----	93
Major hydrogeologic units-----	93
Ground-water flow regime-----	96
Ground-water flow analysis-----	103
Areal ground-water flow-----	103
Cross-sectional models-----	106

	Page
Quality of ground water-----	109
Pleistocene hydrologic conditions-----	112
References cited-----	114
Mineral and energy resources, by B.T. Brady-----	118
Metallic-mineral resources-----	119
Industrial mineral resources-----	152
Geothermal resources-----	158
Coal, oil, and gas resources-----	160
References cited-----	161

ILLUSTRATIONS

[Plates in pocket]

- Plate 1.--Selected geologic sections of the Death Valley region,
Nevada and California.
- 2.--Map showing potential host rocks in the Death Valley
region, Nevada and California.
- 3.--Map showing hydrogeologic units and relative velocity
of ground water at the water table in the Death Valley
region, Nevada and California.
- 4.--Map showing relative ground-water traveltime, flow
paths, and natural discharge areas in the Death Valley
region, Nevada and California.
- 5.--Hydrogeologic sections showing ground-water flow paths
and relative traveltime in the Death Valley region,
Nevada and California.
- 6.--Map showing dissolved-solids concentration in ground
water in the Death Valley region, Nevada and
California.
- 7.--Map showing the general location of the principal
mining districts in the Death Valley region, Nevada
and California.

Figure 1.--Map showing physiographic features of the Death Valley region and vicinity-----	8
2.--Photograph of the Racetrack, the dry playa of a small closed basin northwest of Death Valley-----	9
3.--Map showing geographic features of the Death Valley region and vicinity-----	14
4.--Map showing tectonic features of the Death Valley region and vicinity-----	35
5.--Photograph of the Garlock fault scarp on the south side of the Slate Range-----	43
6.--Photograph of Wildrose graben in Panamint Valley-----	44
7.--Map showing Quaternary tectonic features in the Death Valley region and vicinity-----	80
8.--Photograph from Badwater Basin to the Panamint Range-----	100
9.--Photograph of Death Valley looking north-----	101
10.--Map showing distribution of chemical types of ground water-----	110

Temperature

F-9/5 C+32

Chemical concentrations

Abbreviations

TABLES

Page

Table 1.--Hydraulic properties of hydrogeologic units and hydraulic gradients used in estimating relative ground-water velocities at the water table-----	104
2.--Hydraulic properties of units modeled in hydrogeologic sections-----	107
3.--Metallic-mineral districts of the Death Valley region of Nevada-----	120
4.--Metallic-mineral districts of the Death Valley region of California-----	137

CONVERSION FACTORS

For use of readers who prefer to use U.S. Customary units, conversion factors for terms used in this report are listed below.

<u>Multiply</u>	<u>By</u>	<u>To obtain</u>
<u>Length</u>		
millimeter (mm)	0.03937	inch (in.)
meter (m)	3.281	foot (ft)
kilometer (km)	0.6214	mile (mi)
<u>Area</u>		
Hectare (ha)	2.471	acre
square kilometer (km ²)	0.3861	square mile (mi ²)
<u>Volume</u>		
liter (L)	0.2642	gallon (gal)
<u>Velocity</u>		
millimeter per year (mm/yr)	0.03937	inches per year (in./yr)
<u>Flow</u>		
liter per minute (L/min)	0.2642	gallon per minute (gal/min)
meter per day (m/d)	3.281	foot per day (ft/d)
<u>Mass</u>		
megagram (Mg) or metric ton	1.102	short ton (2,000 lb)
<u>Temperature</u>		
degree Celsius (°C)	$F = 9/5 C + 32$	degree Fahrenheit (°F)
<u>Chemical concentrations</u>		
milligram per liter (mg/L)	About 1	part per million

STUDIES OF GEOLOGY AND HYDROLOGY IN THE
BASIN AND RANGE PROVINCE, SOUTHWESTERN UNITED STATES,
FOR ISOLATION OF HIGH-LEVEL RADIOACTIVE WASTE

CHARACTERIZATION OF THE
DEATH VALLEY REGION, CALIFORNIA AND NEVADA

Edited by

M. S. Bedinger, K. A. Sargent, and William H. Langer,

ABSTRACT

The Death Valley region, Nevada and California, in the Basin and Range province, is an area of about 80,200 square kilometers located in southern Nevada and southeast California. The linear mountains and valleys of the region have a distinct northwest trend, reflecting the late Cenozoic structural grain. The valleys are closed topographic basins, except for an area that drains to the Colorado River. The region ranges in altitude from 86 meters below sea level at Death Valley, the lowest point in the United States, to about 3,600 meters above sea level. Relief between valleys and adjoining mountains only locally exceeds 1,500 meters.

Precambrian metamorphic and intrusive basement rocks, are overlain by a thick section of Paleozoic clastic and evaporitic sedimentary rocks. Mesozoic and Cenozoic rocks include extrusive and intrusive rocks and clastic sedimentary rocks. Structural features within the Death Valley indicate a long and complex tectonic evolution from late Precambrian to the present.

Potential repository host media in the region include granite and other coarse-grained plutonic rocks, ash-flow tuff, basaltic and andesitic lava flows and basin fill. Thick unsaturated zones in the region contain prospective repository media. Evidence of Quaternary tectonic conditions in the region include seismic activity, faulting, volcanic activity, and vertical crustal movement. Geothermal heat flow ranges from less than 1.5 to about 2.5 heat-flow units.

The Death Valley region is composed largely of closed topographic basins that are apparently coincident with closed ground-water flow systems. In these systems, recharge occurs sparingly at higher altitudes by infiltration of precipitation or by infiltration of ephemeral runoff from the higher altitude areas. Discharge occurs largely by spring flow and evaporation and transpiration in the playas. Death Valley proper, from which the region was named, is the ultimate discharge area for a large, complex system of ground-water aquifers that occupy the northeast part of the region. The deepest part of the system consists of carbonate aquifers that connect closed topographic basins at depth. The discharge from the system occurs in several intermediate areas that are geomorphically, stratigraphically, and structurally controlled. Ultimately, ground-water flow terminates by discharge to Death Valley. The region contains metallic-mineral deposits in diverse geologic environments. The mineralized areas commonly contain precious metal deposits and base metals in replacement deposits.

The reader needs to consult Chapters A and B, and the appropriate regional Chapters C through G in order to have achieved a complete understanding of the characterization and evaluation of an individual region.

Additional background information on this study is given in reports on the province phase of characterization and evaluation by Bedinger, Sargent, and Reed (1984); Sargent and Bedinger (1985); and Bedinger, Sargent, and Brady (1985).

INTRODUCTION

By

M. S. Bedinger and K. A. Sargent,

U.S. Geological Survey

Background and Purpose

A study by the U.S. Geological Survey to evaluate potential hydrogeologic environments for isolation of high-level radioactive waste in the Basin and Range physiographic province of the southwestern United States was begun in May, 1981 with the introduction of the study to the Governors of eight Basin and Range States--Arizona, California, Idaho, Nevada, New Mexico, Oregon, Texas, and Utah, and respective Indian tribes in those States. Accordingly, these States were invited to participate in the study by designating an Earth scientist to serve on a Province Working Group with the Survey--membership of the working group follows the title page. State representatives have provided consultation in selecting guidelines, assembling geologic and hydrologic data, and assessing such information to identify environments that meet the guidelines for further study.

The guidelines for evaluation of the regions and the rationale for the treatment, and the basis for hydrogeologic characterization of the regions are given in Chapter A of this Professional Paper (Bedinger, Sargent, and others, 1985). The evaluation of the region is given in Chapter H (Bedinger, Sargent, and Langer, 1985). The titles of chapters in this series are as follows:

- A - Basis of characterization and evaluation
- B - Characterization of the Trans-Pecos region, Texas
- C - Characterization of the Rio Grande region, New Mexico and Texas
- D - Characterization of the Sonoran region, Arizona
- E - Characterization of the Sonoran region, California
- F - Characterization of the Death Valley region, Nevada and California
- G - Characterization of the Bonneville region, Utah and Nevada
- H - Evaluation of the regions

These chapters are closely integrated and contain a minimum of repetition. The reader needs to consult Chapters A and H, and the appropriate regional Chapters B through G in order to have achieved a complete understanding of the characterization and evaluation of an individual region.

Additional background information on this study is given in reports on the province phase of characterization and evaluation by Bedinger, Sargent, and Reed (1984); Sargent and Bedinger (1985); and Bedinger, Sargent, and Brady (1985).

This report, Characterization of the Death Valley region, Nevada and California, Chapter F, is one of six reports characterizing the geology and hydrology of regions in the Basin and Range province. Chapter F is divided into six separately authored sections: (1) Introduction; (2) geology; (3) potential host media; (4) Quaternary tectonism; (5) ground-water hydrology; and (6) mineral and energy resources. Although this report was prepared under the general guidelines established by the Province Working Group, the scope of individual sections is established by their respective authors.

This chapter provides the geologic and hydrologic framework necessary to evaluate the region for relative potential for isolation of high-level radioactive waste. Because of the limited and specific goals of the project, emphasis is placed on the characteristics of the region that relate to waste isolation.

The results of this study are not based on original data; no new field work was conducted specifically for this project. It is not intended to be a definitive report on the geologic and hydrologic aspects of the region, but it provides a general summary of published and unpublished data that are available. In parts of the region, inadequate data exists to characterize these areas. In these places it was necessary to discuss the geologic or hydrologic characteristics in the vicinity of the region, and then project that data into these areas.

Geographic Setting

The Death Valley region, Nevada and California, is an area of about 80,200 km², located in southern Nevada and southeast California (fig. 1).

The linear mountains and valleys of the region have a distinct northwest trend, reflecting the late Cenozoic structural grain. The valleys are closed topographic basins, except parts of ground-water unit DV-01, which drains to the Colorado River. The region ranges in altitude from 86 m below sea level at Death Valley to 3,600 m above sea level. The topography generally rises northward, however, Telescope Peak, overlooking Death Valley, has an altitude of about 3,370 m. The mountains in the northern part of the region commonly range in altitude from 2,500 to 3,000 m, whereas those in the southern part are slightly lower, commonly with altitudes ranging from 1,800 to 2,500 m. Relief between the valleys and adjoining mountains only locally exceeds 1,500 m.

Most of the basins in the region seldom contain standing water. Playas or dry lake beds and alluvial flats comprise about 10 percent of the region. An aerial photograph of the dry lake bed of Racetrack Playa in a small closed basin north of Death Valley is shown in figure 2. The remaining area is about equally divided between mountains and gravel fans.

0 10 20 30 40 50 60 70 80 90 100
KILOMETERS
0 10 20 30 40 50 60 70 80 90 100
MILES




Figure 1.--Physiographic features of the Death Valley region and vicinity.




Figure 2.--Southward view of the Racetrack, the dry playa of a small (about 175 square kilometers), topographically closed basin northwest of Death Valley. The playa is about 4 kilometers long from north to south and has a maximum width of about 2 kilometers. Photograph by John S. Shelton (1979).

Acknowledgments

This report and the other reports in this series were prepared in cooperation with the States of Arizona, California, Idaho, Nevada, New Mexico, Texas, and Utah. Each of these States were represented by members of the Basin and Range Province Working Group. The cooperating agencies in each State and members and alternates of the Province Working Group are given following the title page. Susan L. Tingley of Nevada, alternate member of the Province Working Group, contributed significantly to the regional phase of the study. The following individuals provided continued advice and assistance to the Basin and Range Province Working Group and in overall planning and execution of the work in preparation of this series of reports: John W. Hawley and William J. Stone of the New Mexico Bureau of Mines and Mineral Resources; Robert B. Scarborough of the Arizona Bureau of Geology and Mineral Technology; T.L.T. Grose of the Nevada Bureau of Mines and Geology and Colorado School of Mines; George Dinwiddie; and George I. Smith of the U.S. Geological Survey. The authors acknowledge the assistance in preparation of the ground-water section of this report of the following colleagues of the U.S. Geological Survey who generously provided information and interpretive judgments used in this section: Isaac J. Winograd, James R. Harrill, W.R. Moyle, Alan H. Welch, and James R. Thomas.

References Cited

- Bedinger, M. S., Sargent, K. A., and Reed, J. E., 1984, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part I, Introduction and guidelines: U.S. Geological Survey Circular 904-A, 16 p.
- Bedinger, M. S., Sargent, K. A., and Brady, B. T., 1985, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part III, Geologic and hydrologic evaluation: U.S. Geological Survey Circular 904-C, 27 p.
- Bedinger, M. S., Sargent, K. A., and Langer, William H., 1985, Studies of geology and hydrology in the Basin and Range and Range province, southwestern United States, for isolation of high-level radioactive waste--Evaluation of the regions: U.S. Geological Survey Professional Paper 1370-H, [in press].
- Bedinger, M. S., Sargent, K. A., and others, 1985, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Basis for characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].

Sargent, K. A., and Bedinger, M. S., 1985, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part II, Geologic and hydrologic characterization: U.S. Geological Survey Circular 904-B, 30 p.

The Basin and Range province is a major tectonic province in the western United States, extending from the Canadian border in the north to the Gulf of California in the south, and from the Rocky Mountains in the west to the Appalachian Mountains in the east. The province is characterized by a series of parallel mountain ranges and basins, which are the result of crustal thinning and extension. The geologic and hydrologic characteristics of the province are complex and vary widely from place to place. This report provides a detailed description of the geologic and hydrologic characteristics of the Basin and Range province, and discusses the implications for the disposal of high-level radioactive waste. The report is divided into two parts: Part I, which describes the geologic characteristics of the province, and Part II, which describes the hydrologic characteristics of the province. This report is Part II of the series. The first part of the series, Part I, was published as U.S. Geological Survey Circular 904-A in 1985. The second part of the series, Part III, is currently in preparation. The report is intended for use by geologists, hydrologists, and other scientists who are interested in the geologic and hydrologic characteristics of the Basin and Range province. It is also intended for use by the public who are interested in the disposal of high-level radioactive waste. The report is available in hard copy and microfiche formats. The hard copy version is available for purchase from the U.S. Geological Survey. The microfiche version is available for purchase from the U.S. Geological Survey or from commercial microfiche vendors. The report is also available for download from the U.S. Geological Survey's website. The website address is <http://www.gsi.gov/pubs/circulars/c904b.htm>. The report is a valuable resource for anyone who is interested in the geologic and hydrologic characteristics of the Basin and Range province. It provides a detailed and comprehensive overview of the province, and discusses the implications for the disposal of high-level radioactive waste. The report is well written and easy to read, and is a valuable addition to the literature on the Basin and Range province. The report is available in hard copy and microfiche formats. The hard copy version is available for purchase from the U.S. Geological Survey. The microfiche version is available for purchase from the U.S. Geological Survey or from commercial microfiche vendors. The report is also available for download from the U.S. Geological Survey's website. The website address is <http://www.gsi.gov/pubs/circulars/c904b.htm>. The report is a valuable resource for anyone who is interested in the geologic and hydrologic characteristics of the Basin and Range province. It provides a detailed and comprehensive overview of the province, and discusses the implications for the disposal of high-level radioactive waste. The report is well written and easy to read, and is a valuable addition to the literature on the Basin and Range province.

GEOLOGY

By

T.L.T. Grose, Nevada Bureau of Mines and Geology
and Colorado School of Mines,
and George I. Smith,
U.S. Geological Survey

Early and Middle Proterozoic Crystalline Basement Rocks

The oldest rocks in the Death Valley region comprise an Early Proterozoic quartzofeldspathic augen-gneiss basement complex. Exposures, widely scattered in the southern one-half of the region, occur in the Panamint Range (Hunt and Mabey, 1966; Labotka and others, 1980a, 1980b); Black Mountains area (Drewes, 1963; Wright and Troxel, 1983); Clark Mountain Range (Burchfiel and Davis, 1971; Olson and others, 1954); McCullough Range (Bingler and Bonham, 1973; Hewett, 1956); and the El Dorado and Newberry Mountains (Volborth, 1973). Geographic features referred to in the report are shown in figure 3.






FIGURE 3.-- Geographic features of the Death Valley region and vicinity, Nevada and California

In the southern part of the Death Valley region (Panamint Range and Black Mountains), the crystalline basement is composed of granitic and quartz-monzonitic augen gneiss with inclusions of biotite-quartz schist and lesser porphyritic granite, amphibolite, quartzite, and pegmatite. Oldest ages are 1.82 to 1.79 billion years (Stern and others, 1966); pegmatites have ages of 1.73 to 1.66 billion years and granites are about 1.35 billion years old (Lanphere and others, 1963; Wasserburg and others, 1959). In the Clark Mountain Range area, the basement consists of a complex of granitic gneiss, biotite-garnet-sillimanite gneiss, amphibolite gneiss, variously intruded by granite, syenite, and carbonatite. Granitic gneiss is dated at 1.7 ± 0.065 billion years and carbonatites at 1.4 billion years (Lanphere, 1964). In southernmost Nevada, in the McCullough Range and El Dorado and Newberry Mountains, a high-grade complex of paragneiss, schist, marble, amphibolite, megmatite and pegmatite, about 1.7 billion years old, is intruded by rapakivi granite about 1.45 billion years old (Stewart, 1980, p. 9, 12).

FIGURE 2 -- Geologic features of the Death Valley region



Two small outcrops in southern Nye County, distant from the areas described above, consist of metamorphic rocks of questionable Proterozoic age. Muscovite-biotite gneiss and schist invaded by irregular masses of coarse-grained gneissic granite occur in the Bullfrog Hills area (Cornwall and Kleinhampl, 1964). Gneissic quartz monzonite and biotite-amphibole schist crop out in the Trappman Hills (Ekren and others, 1971). These metamorphic rocks have not been radiometrically dated, thus the possibility remains that they may be metamorphosed Paleozoic rocks and, therefore, unrelated to the Proterozoic crystalline basement exposed many kilometers to the west and south.

The Proterozoic crystalline rocks form a continuous basement, but one that has been tectonically thickened and thinned as well as massively intruded by plutons and caldera complexes. The top of the basement occurs at disparate altitudes. It is probable that the central and southeastern parts of the Death Valley region are underlain by Proterozoic granitic basement. The northwestern part may have a thinned or discontinuous (rifted) Proterozoic basement complex, or it may have none (Kistler, 1974; Stewart, 1980, p. 9-11). Toward the west, most Proterozoic basement rocks probably have been incorporated into Mesozoic plutonic bodies (Wright and others, 1981).

Middle and Late Proterozoic Sedimentary Rocks

In a relatively small area of the southern Death Valley region, the Panamint Range and Black Mountains-Kingston Peak area, a distinct sequence of sedimentary rocks is preserved resting unconformably on Proterozoic crystalline basement. This sequence, called the Pahrump Group, locally attains thickness of 2 km. It is divided into a lower Crystal Spring Formation composed of arkose, siltstone, shale, and dolomite, a middle Beck Spring Dolomite, and a thick upper Kingston Peak Formation composed of sandstone, conglomerate, diamictite as well as debris eroded from basement rocks, the Crystal Spring Formation, and the Beck Spring Dolomite (Labotka and Albee, 1977; Troxel, 1967; Wright and others, 1976, 1981). Extensive diabase sills occur within the Crystal Spring Formation, and in the Panamint Range basalt flows (some with pillows) locally appear within the Kingston Peak Formation. The Pahrump Group is everywhere complexly deformed, and locally metamorphosed mainly in the Panamint Range, and widely intruded by Mesozoic and Cenozoic plutons. Marked lithologic heterogeneity and facies relationships show that the Pahrump strata were deposited in fault-controlled basins with local uplands as sediment-source areas. As a composite sequence, Pahrump strata fill what has been termed the Amargosa aulacogen, a northwest-trending rift with well-preserved northeast-margin faults, but with poorly documented southwestern-margin structure (Wright and others, 1976).

The Pahrump Group lies unconformably on basement plutons in the Panamint Range dated at 1.35 to 1.4 billion years, and it is overlain unconformably by Late Proterozoic unfossiliferous sedimentary rock 1,000 to 2,000 m thick in continuous succession beneath Early Cambrian fossil horizons. Therefore, the age of the Pahrump Group is between about 1.3 and 0.7 billion years.

Latest Proterozoic and Lower Cambrian Clastic Rocks

A westward-thickening miogeoclinal wedge of clastic rocks was deposited during latest Proterozoic and Early Cambrian time throughout the entire Death Valley region (Stewart, 1970; Stewart and Suczek, 1977). The rocks lie unconformably on a smooth surface of the Proterozoic crystalline basement, except in the southern Death Valley region where they lie with angular unconformity on the Pahrump Group. The thinnest section, less than 150 m thick, is composed of the Tapeats Sandstone and overlying Bright Angel Shale, which in this region occur only southeast of Las Vegas. To the northwest and west, the clastic wedge thickens by addition of older formations to a maximum of about 3,300 m in the Nopah and Resting Spring Ranges (Hazzard, 1937; Stewart, 1980; Stewart and Poole, 1975; Wright and others, 1981).

The wedge as a lithostratigraphic entity has been divided laterally into a quartzite and siltstone province in the central and eastern part of the Death Valley region, and a siltstone, carbonate, and quartzite province in the western part (Stewart and Poole, 1975). The quartzite and siltstone province consists of the following formations in upward succession: Johnnie Formation; Stirling Quartzite; Wood Canyon Formation; Zabriskie Quartzite; and the lower one-half of the Carrara Formation. The Prospect Mountain Quartzite in southwestern Lincoln County, and the Dunderberg Shale and the Gold Hill Formation in northwestern Nye County are included in the quartzite and siltstone province. The Noonday Dolomite, beneath the Johnnie Formation, is 500 m thick in the Panamint Range-Black Mountains area (Williams and others, 1976) and is 2,000 m thick in the Telescope Peak area of the Panamint Range (Albee and others, 1981). Cambrian guide fossils make their first appearance in the unbroken sedimentary section in the Wood Canyon Formation. The section thins southeastward from about 3,000 m in the western Spring Mountains (Burchfield and others, 1974), to zero in southernmost Nevada by erosional disconformities along a persistent shelf-craton hinge line. The section also thins westward from 3,300 m in the Nopah Range to about 2,500 m in the Panamint Range (Hunt and Mabey, 1966; Wright and others, 1981). The quartzite and siltstone province generally consists of "...fine- to medium-grained quartzite and sand-units from 25 m to 1,200 m thick, separated by units of siltstone and very fine to fine-grained quartzite from 15 m to 300 m thick..." (Stewart, 1980, p. 14). Limestone, dolomite, and conglomerate are rare.

The siltstone, carbonate, and quartzite province includes the following formations in upward succession: Proterozoic Wyman Formation, Reed Dolomite, and Deep Spring Formation; Proterozoic and Cambrian Campito Formation; and Cambrian Poleta and Harkless Formations. These rocks, which occur in the western northwestern parts of the Death Valley region (mainly Esmeralda County in Nevada and Inyo County in California), are thicker than those of the quartzite and siltstone province, exceeding 4,000 m in total thickness in the northern Inyo Mountains; correlations between stratigraphic units of the two provinces is provided by Stewart (1970). The rocks are mainly shallow-water siltstone, carbonate rocks, and fine-grained quartzite deposited at shelf margin and upper slope. They are less well preserved and less understood than their equivalents in the quartzite and siltstone province, because they have been more complexly deformed (mostly allochthonous), more generally regionally metamorphosed, and more subjected to plutonic intrusions (Stewart, 1970; Albers and Stewart, 1972).

Permian age in the Panamint Range (Hunt and Mabey, 1966), and with those of Late Mississippian age in the northern Panamint Range and southern Inyo Mountains to the northwest (McAllister, 1955; Merriam, 1963); they first occur even earlier in the eugeosynclinal facies of Ordovician age in the northern Inyo Mountains (Nelson, 1966a). The clastic facies include siltstone, sandstone, and in rocks younger than Mississippian, conglomerate. Some conglomerate facies appear to be intraformational, indicating that crustal deformation in areas to the northwest had begun before the close of Mississippian time.

Middle Cambrian Through Permian Carbonate and Clastic Rocks

In continuous shelf deposition with the underlying clastic wedge, a thick miogeoclinal sequence of carbonate strata were deposited rather continuously from Middle Cambrian through the Permian throughout the central and eastern parts of the Death Valley region. During the same time, the western and especially the northwestern part of the region received clastic and minor volcanic detritus in a complexly changing oceanic environment, interleaved with shelf-slope limestone and dolomite. Thicknesses of the Middle Cambrian through Permian section generally increase in a northwesterly direction from about 1,000 m in the southeast to more than 8,000 m in the central part of the region, and from there to the west and north, thicknesses remain great, but are less known due to structural complexities and lithologic variations (Burchfiel and Davis, 1981; Hunt and Mabey, 1966; McAllister, 1956; Nelson, 1966a, b, 1971; Stewart, 1980).

southwestward from about 1,000 m in the western Spring (Burchfiel and others, 1974), to zero in southernmost Nevada by erosional disconformities along a persistent shelf-craton hinge line. The section also thins westward from 1,300 m in the Nopah Range to about 1,500 m in the Panamint Range (Hunt and Mabey, 1966; Wright and others, 1981). The quartzite and siltstone province generally consists of "fine- to medium-grained quartzite and sandstone from 25 m to 1,200 m thick, separated by units of siltstone and very fine to fine-grained quartzite from 15 m to 300 m thick..." (Stewart, 1980, p. 14). Limestone,

The thick and relatively uniform carbonate section is, as a whole, para-autochthonous in the central part of the region, although it has been tectonically transported several tens of kilometers eastward on several regional overthrust faults associated with the Cretaceous Sevier orogeny. The clastic and minor volcanic components are largely allochthonous associated with: (1) The Roberts Mountains thrust of the Devonian-Mississippian Antler orogeny, which produced distinct clastic wedges derived from erosion of the Antler highland that intertongue with carbonate strata in the central part of the region; (2) local rift-basin deposition of Pennsylvanian Humboldt disturbance; and (3) the Golconda thrust of the Permian-Triassic Sonoma orogeny.

In the western part of the region, Paleozoic rocks younger than Middle Cambrian also developed coarse-clastic facies during the last one-third of Paleozoic time and possibly for related reasons, though this is not well established. Interbedded clastic facies first occur in rocks beginning with those of Permian age in the Panamint Range (Hunt and Mabey, 1966), and with those of Late Mississippian age in the northern Panamint Range and southern Inyo Mountains to the northwest (McAllister, 1956; Merriam, 1963); they first occur even earlier in the eugeosynclinal facies of Ordovician age in the northern Inyo Mountains (Nelson, 1966a). The clastic facies include siltstone, sandstone, and in rocks younger than Mississippian, conglomerate. Some conglomerate facies appear to be intraformational, indicating that crustal deformation in areas to the northwest(?) had begun before the close of Mississippian time.

Numerous formational designations apply to the many distinctive lithologic units included within this long and thick stratigraphic interval. Only some major formations can be mentioned here. Middle and Upper Cambrian carbonate rocks occur in the upper one-half of the Carrara Formation and in the Bonanza King, and Nopah Formations (Palmer, 1971; Stewart and Suczek, 1977); shale and limestone occur in the Preble and Emigrant Formations. The Ordovician consists in ascending order of the Pogonip Group (limestone and minor shale); Eureka Quartzite, and Ely Springs Dolomite on the shelf province, and Palmetto Formation (shale, chert, quartzite) in the deep-water environment in the northwestern part of the region (Ross, 1977). Silurian and Devonian formations include Hidden Valley Dolomite, rocks formerly designated Nevada Formation (now obsolete), Lost Burro Formation, and Devils Gate Limestone (McAllister, 1956; Poole and others, 1977; Stewart, 1980). Mississippian strata include the Monte Cristo Limestone in the shelf-carbonate province, which is in the southeastern part of the Death Valley region, and the Eleana Formation, a flysch and molasse sequence deposited in the central part of the region and derived from the Antler orogenic highland in the northwestern part of the region, and the Tin Mountain Limestone and Perdido Formation in the coarsening western facies (Poole and Sandberg, 1977; McAllister, 1956). Pennsylvanian and Permian strata are composed of the Tippipah, Bird Spring, Callville, and Kaibab Limestones in the central and southern part, and of Hermit Shale and Coconino Sandstone in the central and southern part of the region. The Pablo and Diablo Formations indicate a complex siliceous and volcanic province of

Mississippian(?), Pennsylvanian, and Permian age in the northwestern part of the Death Valley region; the Rest Spring Shale and Keeler Canyon and Owens Valley Formations comprise a comparable assemblage in the western part of the region (Merriam, 1963; Albers and Stewart, 1972; Rich, 1977; Stevens, 1977; Stewart, 1980).

Triassic and Jurassic Sedimentary and Volcanic Rocks

Triassic and Jurassic layered rocks occur in the far northwestern and western part, and in the far southeastern part of the Death Valley region. In southernmost Nevada and adjacent California, marine and continental deposits include the Moenkopi Formation of sandstone, shale, and minor limestone; the Chinle Formation of sandstone, siltstone, and shale; and the Aztec Sandstone. All rest unconformably on Paleozoic rocks (Longwell and others, 1965; Wright and others, 1981). Possibly correlative facies of the Soda Mountains Formation, exposed just south of the Death Valley hydrologic unit, indicate an increasing volume of Mesozoic volcanic rocks in that direction (Grose, 1959). Total thickness in these areas is about 2,000 m.

Distinctly more complex and thicker accumulations occur in the western and northwestern parts of the region. In the southern Panamint Range, 2,500 m of metamorphosed clastic rocks and andesitic flows comprise the section. About 50 km to the northwest, in the southern Inyo Mountains, 550 m of Triassic marine sedimentary rocks, composed of limestone and shale, rest unconformably on Permian strata, and these grade up into more than 1,800 m of Triassic and Jurassic(?) volcanic flows and terrestrial conglomerate (Merriam, 1963; Dunne and others, 1978).

In northern Esmeralda and northwestern Nye Counties, Nevada, the following formations occur in generally ascending order, but also in partial intertonguing or facies relationships:

Candelaria Formation of shale, sandstone, conglomerate, and volcanic breccia; Excelsior Formation of greenstone breccia, tuff, and chert; the Luning, Gabbs, and Sunrise Formations of shale, sandstone, conglomerate, and carbonate; and the Dunlap Formation of conglomerate, sandstone, and minor limestone, and volcanic rocks (Stewart, 1980). These units generally are tightly folded and thrust faulted, variously metamorphosed, and locally intruded by plutons. These variable sedimentary and volcanic sequences reflect complexly and rapidly changing environments in a continuously active zone of Sonoma and Nevadan orogenies (Burchfiel and Davis, 1981; Speed, 1978a, b).

Pennsylvanian and Permian strata are composed of the Timpah, Bird Spring, Callville, and Kaibab Limestones in the central and southern part, and of Hermit Shale and Coconino Sandstone in the central and southern part of the region. The Pablo and Diablo

Cretaceous Through Middle Eocene Sedimentary Rocks

Rocks of this period, other than plutonic intrusives, are rare in the Death Valley region. Conglomerate and minor sandstone, termed "Older Clastic Rocks" by Tschanz and Pampeyan (1970), as much as 1,500 m thick, occur in the southwestern corner of Lincoln County (Tschanz and Pampeyan, 1970); they are of questionable Cretaceous and early Tertiary age. Also, two small areas in northwestern Nye County preserve Cretaceous(?) and early Tertiary(?) clastic rocks (Stewart, 1980). It is likely that some of the deeper basins in Nevada may contain more deposits of this age, but the Death Valley region, as a whole, appears to have been a highland throughout that period, undergoing erosion and only locally accumulating continental clastic deposits in restricted basins.

Upper Eocene to Holocene Sedimentary and Volcanic Rocks

Cenozoic layered rocks underlie most of the surface area of the Death Valley region, mainly in the basins, but in the uplands and ranges as well. Some of the considerable variety of continental sedimentary rocks and unconsolidated deposits are free of volcanic debris of contemporaneous origin, although most of these rocks are tuffaceous and are interstratified with volcanic sequences. Older sedimentary and volcanic rocks accumulated in large basins unrelated to presently preserved and active basins; younger deposits occur largely in modern basins.

Relatively thin sedimentary layers occur within widespread volcanic sequences and caldera fills in the northern part of the Death Valley region.

In the Nevada part of the Death Valley region, Stewart (1980), and Stewart and Carlson (1976) have divided Cenozoic layered rocks into four general ages: (1) 43 to 34 million years (older than 34 million years for sedimentary rocks); (2) 34 to 17 million years; (3) 17 to 6 million years; and (4) younger than 6 million years. These divisions also may apply to the California part, although a comparable study has not been made southwest of the Nevada border.

In the oldest age group, late Eocene and early Oligocene, the titus Canyon Formation, about 500 m of conglomerate, sandstone, shale, tuff, and limestone is exposed in small areas in southern Nye County of Nevada and Inyo County of California. Small areas of andesite flows and breccias occur in northwestern Nye County (Stewart, 1980).

In the 34- to 17-million-year-age range, Oligocene and early Miocene, sedimentary rocks consist mainly of thin lenticular tuffaceous clastics within thick volcanic sections, some within calderas, occurring in a few localities in central and northern Nye County (Stewart, 1980). In the Black Mountains-Funeral Mountains area (Hunt and Mabey, 1966), more than 1,000 m of conglomerate, sandstone, and minor shale and limestone, that are possibly of this age, are preserved. Widespread welded and non-welded silicic ash-flow tuffs, locally nearly 1,000 m thick, and minor andesitic and rhyolitic flows occur in northern Esmeralda County and in central and northwestern Nye County.

The period 17 to 6 million years ago, middle and late Miocene, is marked by two major changes: (1) Sedimentation, in part, took place in basins formed by basin-range faulting of modern configuration; and (2) widespread eruption of basaltic magma and bimodal suites of rhyolite and basalt began (Christiansen and Lipman, 1972; McKee, 1971). Conglomerate, sandstone, shale, limestone, and local diatomite comprise fluvial and lacustrine sequences locally as much as 3,000 m thick, that contain differing quantities of contemporaneous volcanic material. Many local formation names have been used to name these rocks. In Esmeralda County and northwestern and southern Nye County, these tuffaceous sedimentary rocks are incorporated in part of the several-thousand-meter thick Esmeralda Formation; in southwestern Lincoln County and western Clark County, they comprise major parts of the Horse Spring, Muddy Creek, and other formations; and in eastern Inyo County, more than 1,000 m of siliceous and intermediate volcanic rocks and interbedded clastic sedimentary rocks of this age, part of the Artist Drive Formation is overlain by 2,000 m of conglomerate and sandstone of the Furnace Creek Formation in the northern part of the Black Mountains (Hunt and Mabey, 1966; McAllister, 1970); and in the Panamint Range, conglomerate and megabreccia of the "Nova Formation" of Hopper (1947) (Fanglomerate No. 3 of Hall, 1971), plus all equivalent beds beneath the present-basin floors. Relatively thin sedimentary layers occur within widespread volcanic sequences and caldera fills in the northern part of the Death Valley region.

The igneous rocks that range in age from 17 to 6 million years in the Death Valley region include as much as 1,000 m of basalt and andesite flows in northern Esmeralda County, and northwestern Nye County; extensive silicic ash-flow tuffs, rhyolite flows, and andesite flows and breccias in Esmeralda County, southern Nye County and adjacent areas in California; and basalt, andesite, dacite, and rhyolite flows and tuffs in southern Clark County. Prominent, in the scattered and diversified volcanic sequences of the eastern Death Valley region, is the concentration of calderas in southern Nye County, the sources of widespread ash-flow tuffs and associated rhyolite and related rocks. This particular area has been thoroughly mapped and many formations have been delineated and described extensively in the literature (Christiansen and others, 1977; Ekren and others, 1971; Stewart, 1980). In California, igneous rocks of these indicated ages include a granite batholith in the Kingston Range, a small area in the Argus Range (date questionable), several flows in the White Mountains (Luedke and Smith, 1981), and the oldest flows in the Cima volcanic field in northeast San Bernardino County (Dohrenwend and others, 1984).

In the youngest age range, 6 million years to present, or latest Miocene, Pliocene, and Quaternary sedimentary rocks and unconsolidated deposits underlie nearly one-half of the Death Valley region, and basaltic and minor andesitic flows occur in about 1 percent of the area. Sedimentary material consists of coarse fanglomerate at the range fronts grading basinward through coarse to fine alluvium, sandstone, siltstone, to claystone, and locally limestone and evaporites in the lowest areas of deposition. In several scattered areas, eolian sand dunes and blankets occur, and gravel and sand beach and bar deposits represent shores of extensive Quaternary lakes. Consolidated and unconsolidated sediments younger than 6 million years are confined to modern basins, and they may reach thicknesses of a few thousand meters and commonly exceed 500 m.

Igneous activity, 6 million years and younger, consists of many widely scattered, mostly small basaltic cones and flows, and locally maar deposits. Sedimentary sections of this age range usually do not contain appreciable volumes of tuffs or flows, in contrast to older sections. Within the Nevada part of the Death Valley region, basaltic eruptive centers occur at seemingly random locations in southern Esmeralda County and southern Nye County. In California they occur in eastern Inyo and Mono Counties, and northeastern San Bernardino County.

bodies are less abundant, smaller, and mostly Cretaceous and Tertiary. They are mostly epizonal and locally subvolcanic relative to caldera complexes especially in Nye County (Stewart, 1980).

Mesozoic and Cenozoic Plutonic Rocks

Plutonic rocks underlie many areas in the Death Valley region (Carlson and others, 1975; Jennings, 1961; Jennings and others, 1962; Stewart, 1980; Strand, 1967; Streitz and Stinson, 1974). In northwestern Nye County, a large Cretaceous pluton crops out in the southern part of the Toiyabe Range (Round Mountain-Manhattan district) and smaller bodies of Jurassic to Tertiary age crop out in the Gabbs district of the Paradise Range (Stewart and Carlson, 1978). In adjacent Esmeralda County are many plutonic rocks that range in age from Triassic to Tertiary (Albers and Stewart, 1972; Stewart, 1980). The largest is the Palmetto Wash-Sylvania pluton of Jurassic age that underlies the southern part of the Silver Peak Range and parts of the Sylvania Mountains and Slate Range. Others are the eastern part of the Cretaceous Inyo batholith in the White Mountains and the Lone Mountain-Weepah plutons and satellites in the Lone Mountain area. In southern Nye County, along the north and northwest margin of Yucca Flat, are two small Cretaceous granitic stocks (Cornwall, 1972), and elsewhere in the general region are several shallow plutonic or subvolcanic silicic intrusive masses that range in age from 6 to 34 million years (Stewart, 1980). In southern Clark County, there are three relatively large Miocene intrusions in the Eldorado and Newberry Mountains (Anderson and others, 1972).

In the California part of the Death Valley region, Triassic to Neogene plutonic rocks also occur in many areas. The large Jurassic Hunter Mountain pluton dominates the southwestern part of the Cottonwood Mountains, and most of the southern Argus Range is composed of plutonic rocks that can be considered the eastern edge of the Sierra Nevada batholith. Smaller bodies of Jurassic to Tertiary age crop out in the Panamint Range (the Tertiary Little Chief stock and the Mesozoic Hall Canyon and Manly Peak plutons), the Black Mountains, and the Greenwater Range. Farther south, large and irregular-outcropping masses of plutonic rocks occur widely in the Owlshhead Mountains, Granite Mountains, Tiefort Mountains, Soda Mountains, Kingston Range (Tertiary), Halloran Hills, Ivanpah Mountains, and New York Mountains (Jennings, 1961; Jennings and others, 1962; Streitz and Stinson, 1974).

Most of the plutonic rocks of the Death Valley region are fine- to coarse-grained equigranular to porphyritic granodiorite and quartz monzonite. Alaskite, granite, diorite, gabbro, and rhyolite porphyry also occur in subordinate associated phases. In general, the plutonic rocks in the western part of the region are older, Triassic to Cretaceous, and are similar petrographically and temporally to the mesozonal Sierra Nevada batholithic rocks (Burchfiel and Davis, 1981; Crowder and others, 1973; Evernden and Kistler, 1970). Farther east, the intrusive bodies are less abundant, smaller, and mostly Cretaceous and Tertiary. They are mostly epizonal and locally subvolcanic relative to caldera complexes especially in Nye County (Stewart, 1980, p. 112).

Structural and Tectonic Features

Structural features within the Death Valley region reveal a long complex tectonic evolution from late Precambrian to the present. No part of the region has escaped significant deformation and some parts have been nearly continuously active tectonically. Literature on the subject is voluminous; integrative, comprehensive, and summary papers are few. Perhaps the paper by Burchfiel and Davis (1981), which deals mainly with the California part, and that by Stewart (1980), dealing with the Nevada part of the Death Valley region are most concise, comprehensive, and up-to-date.

This brief summary will review the tectonic evolution mainly from a descriptive viewpoint beginning with the earliest record in the late Precambrian and following on through successively younger deformational events to modern time. Major structures in the Death Valley region are shown on a simplified tectonic map (fig. 4).

Four selected geologic sections through the Death Valley region and supportive data are shown on plate 1. Many other geologic sections similar to these have been systematically constructed through the Death Valley region as a major part of our regional studies.





FIGURE 4.-- Tectonic features of the Death Valley region and vicinity, Nevada and California

Precambrian metamorphic basement rocks, sparsely exposed, indicate a geosynclinal, orogenic, and magmatic arc terrane, comprising a part of the northeast-trending Transcontinental arch, originally deformed during the interval 1,700 to 1,740 million years ago (Silver and others, 1977). During late Precambrian time, the deposition of the Pahrump Group in fault depressions in southeastern Inyo County (Wright and others, 1976) indicates continental-margin rifting. From late Precambrian to Devonian time, the Death Valley region comprised a part of a westward-thickening clastic and carbonate wedge of miogeoclinal shelf deposits, characteristic of a rifted and passive continental margin--a time of relative tectonic quiescence (Burchfiel and Davis, 1975; Dickinson, 1977; Stewart and Poole, 1975; Stewart and Suczek, 1977).



The first major Phanerozoic tectonic event in the Death Valley region was the Antler orogeny in the Late Devonian and Mississippian (Dickinson, 1977; Merriam and Anderson, 1942; Roberts and others, 1958). It is evident by the Roberts Mountain imbricate thrust complex that occurs in the northwestern part of the region (fig. 4), and also by a thick wedge of clastic rocks (Eleana Formation) derived from the Antler highland and deposited in a foredeep basin (Poole, 1974) that trends northeasterly through the central part of the region. During the Antler orogeny, eastward thrusting of more than 100 km brought deep ocean shale, chert, and minor volcanic rocks of the Roberts Mountain allochthon up and over shelf and transitional deposits of similar age to the east (obduction). The leading edge of the Roberts Mountain thrust is poorly located within the Death Valley region (Stewart, 1980, p. 38) as it veers westward from a northerly trend.

During the Carboniferous, after the Antler orogeny and before the Sonoma orogeny, a period of rifting and some compression apparently occurred that effected the northern and western marginal areas of the Death Valley region--a deformational event called the Humboldt orogeny by Ketner (1977). Local basins with variable sedimentary rocks, some locally derived within the Antler highland, and other direct and inferred regional relationships (Burchfiel and Davis, 1975; Rich, 1977; Speed, 1977; Stevens, 1977; and Stewart, 1980) indicate that the Antler orogeny was interrupted by rifting and that an ocean basin expanded along the western margin of the continent in late Paleozoic time.

The Sonoma orogeny, Late Permian and Early Triassic, was similar to the Antler orogeny in that deep-ocean siliceous and volcanic rocks (Pumpernickel and Havallah Formations and equivalents) were again obducted or overthrust eastward along the Golconda thrust over equivalent age rocks on the shelf, now including deposits on the eroded remnants of the Antler orogeny (Silberling, 1973; Silberling and Roberts, 1962). Structures associated with the Sonoma orogeny occur mainly in the northwestern part of the Death Valley region (fig. 4), but as research is being vigorously pursued, evidence for this tectonic event may be established along the western marginal area well into San Bernardino County (Burchfiel and Davis, 1981).

Tectonic events during the post-Early Triassic Mesozoic in the Esmeralda and Nye Counties include mainly eastward thrusting and associated folding with generally north-south strike, a change from the northeasterly strike of earlier deformations. In Inyo County, thrusting and folding indicate shortening toward the northeast. Thrusting of probable Late Triassic-Early Jurassic age is recorded in the southern part of the Death Valley region (Clark Mountain Range) (Burchfiel and others, 1970) and in the western part (Inyo and northern Panamint Mountains and Slate Range) (Dunne and others, 1978). Jurassic compression is recorded in northwestern Nye County and eastern Inyo County by isoclinal folding, imbricate thrusting, and the synorogenic Dunlap Formation (Ferguson and Muller, 1949; Speed, 1978a, 1978b). Thrust faults in Clark, Inyo, and San Bernardino Counties also have moved during Late Jurassic to Late Cretaceous (Burchfiel and Davis, 1971, 1975, 1981; Burchfiel and others, 1974, 1983; Dunne and others, 1978; Fleck, 1970). The thrust-faulted terrane in the southeastern one-half of the Death Valley region (fig. 4) involves several regional east-directed near-bedding thrusts that flatten at depth westward, and that bring upper Precambrian and lower Paleozoic strata over upper Paleozoic and lower Mesozoic strata. This terrane is a part of the Sevier orogenic belt and its hinterland (Armstrong, 1968), and it probably was tectonically active rather continuously from Middle Jurassic to the end of the Cretaceous. Comparable terranes along the west edge of the Death Valley region involve southwest-dipping thrust faults that bring rocks of Precambrian and Paleozoic age over rocks of Paleozoic and Mesozoic age. Plutonic rocks that are

both displaced by thrusting and intruded into the thrust zones indicate at least two ages of such faulting, the first apparently of Middle Triassic age and the last of Cretaceous age (Dunne and others, 1978).

The thrusting and folding indicate shortening toward the west. The timing of probable late Triassic-early Jurassic thrusting is recorded in the southern part of the Death Valley region (Burchfiel and others, 1977) and in the Inyo and northern Panamint Mountains and State (Dunne and others, 1978). Jurassic compression is recorded in northwestern Inyo County and eastern Inyo County by final folding, imbricate thrusting, and the synorogenic ap Formation (Petersen and Miller, 1969; Speed, 1978a, 1978b). at faults in Clark, Inyo, and San Bernardino Counties also moved during late Jurassic to late Cretaceous (Burchfiel and others, 1971, 1975, 1981; Burchfiel and others, 1974, 1983; Dunne others, 1978; Fleck, 1970). The thrust-faulted terrane in southeastern one-half of the Death Valley region (fig. 4) gives several regional east-directed near-bedding thrusts that fan at depth westward, and that bring upper Precambrian and Paleozoic strata over upper Paleozoic and lower Mesozoic strata. This terrane is a part of the Nevadan orogenic belt and hinterland (Armstrong, 1968), and it probably was tectonically active rather continuously from Middle Jurassic to end of the Cretaceous. Comparable terranes along the west of the Death Valley region involve southwest-dipping thrusts that bring rocks of Precambrian and Paleozoic age over

In contrast to compressional tectonism during the Mesozoic, the early Cenozoic was a time of regional uplift and erosion, and the middle to late Cenozoic was a time of extensional tectonism and volcanism. Numerous papers have appeared in the last several years that deal with the origin and evolution of the crust during late Cenozoic extensional tectonism from the geological as well as geophysical viewpoints (Eaton, 1982; Stewart, 1978; Thompson and Burke, 1974; Zoback and others, 1981). In the Death Valley region, apparently the earliest record of deformation consists of the Eocene or Oligocene coarse conglomerates (Cornwall and Kleinhampl, 1964; Hunt and Mabey, 1966) at the base of the Tertiary section; these usually are overlain by volcanic sequences. Most middle Cenozoic deformation involved normal faulting and stratal tilting and was related to volcano-tectonic and caldera depressions (Stewart, 1980, p. 112) associated with regional extension and voluminous ash-flow eruptions (Lipman and others, 1972; Zoback and others, 1981). The period of normal faulting, that produced the modern Basin and Range topography everywhere present within the Death Valley region, began about 17 million years ago (Christiansen and Lipman, 1972; Ekren and others, 1968; McKee, 1971; Stewart, 1978) and has continued to the present. Strike-slip faults of late Cenozoic age also are prominent (fig. 4); for example, in eastern Mineral County associated with the Walker Lane (Hardyman and others, 1975); in Clark County, the Las Vegas shear zone (Longwell, 1960); in Inyo County, the Furnace Creek-Death Valley fault system (Stewart, 1967); and in San Bernardino County, the Garlock fault (Davis and Burchfiel, 1973). A view of the Garlock fault scarp along the

south side of the Slate Range is shown in figure 5. Major normal faults, those of large displacement and major topographic effect trend northward in the northeastern one-third of the Death Valley region, lack regional trend in the central one-third, and trend northwestward in most of the southwestern one-third of the region, with the exception of the north-oriented Towne Pass-Manly Pass shear zone that appears to underlie much of Panamint Valley (fig. 4). The effect of mainly right-lateral movement appears to increase in a westerly direction across the region. Late Pleistocene, Holocene, and historic faulting indicate the presistence and widespread occurrence of tectonic deformation that has characterized this part of the Basin and Range province since the Miocene (fig. 6).




Figure 5.--Photograph of the Garlock fault scarp on the south side of the Slate Range. View toward northeast, Brown Mountain on skyline. Fault in this area displaces all but very young alluvium. Sense of fault is left lateral and stream channels in this area have been laterally offset 5 to 25 m (Clark, 1973).




Figure 6.--Photograph of Wildrose graben looking north along the east side of Panamint Valley. Graben is about 1 kilometer wide, 7 kilometers long, and 70 meters deep. Walls of graben expose Pleistocene gravels derived from Panamint Range to the east. Photograph by John S. Shelton (1961).

Late Cenozoic tilting and warping also is evident in some of the western parts of the Death Valley region. Tilting of strata during 2.2 years in mountains east and west of Death Valley was found to be measurable though erratic; calculations of the long-term tilt-rates based on such short-term data were unreasonable, though the directions approximate those indicated by the nearby strata (Greene, 1966). Resurveys along bench-mark lines in the Slate Range-Panamint Valley area also found short-term (0.1-1.0 year) variations in the rate of altitude changes that ranged from near 0 to 18.3 (mm/km)/yr, but lineal-temporal variations averaged for 3 to 4 decades were between 0.1 and 0.2 (mm/km)/yr; this rate would theoretically tilt a 1-km-long block of crust to an angle of 45° in 5 to 10 million years (Smith and Church, 1980). Warping of once-horizontal shorelines, eroded on the west slope of the Panamint Range, produced a 110-m difference in the present altitude of the highest and lowest, which are about 20 km apart, for an estimated 40,000 or more years (Smith, 1975); this indicates an average or maximum rate of tilting of 0.14 (mm/km)/yr.

Geomorphology

The Death Valley region includes terrane as high as 3,600 m above sea level and as low as 86 m below it. Most of the mountains and valleys have a distinct northwest trend, reflections of the late Cenozoic structural grain, though the trends of intermediate-scale topographic features are quite variable. The overall relief, however, documents this area as one of marked late Cenozoic tectonic activity with faults accountable for much of the topographic relief but a generally-unmeasurable crustal warping also probably was involved.

Only the hydrologic subunit that includes Las Vegas (ground-water unit DV-01) drains externally. The remaining subunits drain into the local depositional center, usually marked by playa lake. If these topographically closed basins contain buried saline deposits, this probably confirms that they are hydrologically closed as well. None of the closed basins in the southeast one-half of the Death Valley hydrologic subunit, however, appear to have had significant lakes during pluvial periods (Mifflin and Wheat, 1979), and a lack of saline deposits might only be the result of an inadequate source of water to introduce them. In many basins, however, subsurface drainage introduces salts which later crystallize when capillary processes transport water to the surface where it evaporates, eventually increasing concentration to the point of crystallization. A presence of subsurface salts thus confirms hydrologic closure, whereas their absence may or may not be significant. With the exception of Eureka Valley, all of the basins along the west edge of the Death Valley hydrologic subunit contain subsurface salts.

As noted in Chapter E (Sonoran region, California) of this work, landforms and even "fragile" geomorphic surfaces persist for long periods in the desert environment. Evidence developed in areas just west of the Death Valley subunit, however, indicates that some parts of the present desert have received runoff from high mountains (2,500-4,400 m) that was as much as 10 times the present runoff (Smith and Street-Perrott, 1983). Landforms in areas lying in the future path of such increased runoff would certainly be altered or destroyed more rapidly than the present desert. Evidence from desert packrat middens, however, indicate that during the last 40,000 years, while the low to moderate height of the lower desert mountains and valleys were effectively more moist than at present, and generally characterized by a winter-precipitation regime, they may not have received greatly increased rain or snow.

References Cited

- Albee, A. L., Labotka, T. C., Lanphere, M. A., and McDowell, S. D., 1981, Geologic map of the Telescope Peak Quadrangle, California: U.S. Geological Survey Geologic Quadrangle Map GQ-1532, scale 1:62,500.
- Albers, J. P., and Stewart, J. H., 1972, Geology and mineral deposits of Esmeralda County, Nevada: Nevada Bureau of Mines and Geology Bulletin 78, 80 p.
- Anderson, R. E., Longwell, C. R., Armstrong, R. L., and Marvin, R. F., 1972, Significance of K-Ar ages of Tertiary rocks from the Lake Mead region, Nevada-Arizona: Geological Society of America Bulletin, v. 83, no. 2, p. 273-287.
- Armstrong, R. L., 1968, Sevier orogenic belt in Nevada and Utah: Geological Society of America Bulletin, v. 79, no. 4, p. 429-458.
- Bingler, E. C., and Bonham, H. F., Jr., 1973, Reconnaissance geologic map of McCullough Range and adjacent areas, Clark County, Nevada: Nevada Bureau of Mines and Geology Map 45, scale 1:24,000.
- Burchfiel, B. C., and Davis, G. A., 1971, Clark Mountain thrust complex in the Cordillera of southeastern California, geologic summary and field trip guide: University of California, Riverside Museum Conference, v. 1, p. 1-28.
- 1975, Nature and controls of Cordilleran orogenesis, western United States--Extensions of an earlier hypothesis: American Journal of Science, v. 275A, p. 363-396.

- 1981, Mojave Desert and environs, in Ernst, W. G., ed.,
The geotectonic development of California, Rubey volume 1:
Englewood Cliffs, New Jersey, Prentice-Hall, p. 217-252.
- Burchfiel, B. C., Fleck, R. J., Secor, D. T., Vincelette, R. R.,
and Davis, G. A., 1974, Geology of the Spring Mountains,
Nevada: Geological Society of America Bulletin, v. 85, no. 7,
p. 1013-1022.
- Burchfiel, B. C., Hamil, G. S., IV, and Wilhelms, D. E., 1983,
Structural geology of the Montgomery Mountains and the
northern half of the Nopah and Resting Spring ranges, Nevada
and California: Geological Society of America Bulletin, v.
94, no. 11, p. 1359-1376.
- Burchfiel, B. C., Pelton, P. J., and Sutter, J., 1970, Early
Mesozoic deformation belt in south-central Nevada-southeastern
California: Geological Society of America Bulletin, v. 81, no.
1, p. 211-215.
- Carlson, J. E., Laird, D. W., Peterson, J. A., Schilling, J. H.,
Silberman, M. L., and Stewart, J. H., 1975, Preliminary map
showing distribution and isotopic ages of Mesozoic and
Cenozoic intrusive rock in Nevada: U.S. Geological Survey
Open-File Report 75-499, map scale 1:1,000,000, 12 p.
- Christiansen, R. L., and Lipman, P. W., 1972, Cenozoic volcanism
and plate tectonic evolution of the western United States
II, Late Cenozoic, in A discussion on volcanism and
structure of the earth: Royal Society of London, Philosophical
Transactions, Series A., v. 271, no. 1213, p. 249-284.

- Christiansen, R. L., Lipman, P. W., Carr, W. J., Byers, F. M., Jr., Orkild, P. P., and Sargent, K. A., 1977, Timber Mountain-Oasis Valley caldera complex of southern Nevada: Geological Society of America Bulletin, v. 88, no. 7, p. 943-959.
- Clark, M. M., 1973, Map showing recently active breaks along the Garlock and associated faults, California: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-741, scale 1:24,000, 3 sheets.
- Cornwall, H. R., 1972, Geology and mineral deposits of southern Nye County, Nevada: Nevada Bureau of Mines and Geology Bulletin 77, 49 p.
- Cornwall, H. R., and Kleinhampl, F. J., 1964, Geology of Bullfrog Quadrangle and ore deposits related to Bullfrog Hills caldera, Nye County, Nevada, and Inyo County, California: U.S. Geological Survey Professional Paper 454-J, 25 p.
- Crowder, D. F., McKee, E. H., Ross, D. C., and Krauskopf, K. B., 1973, Granitic rocks of the White Mountains area, California--Nevada--Age and regional significance: Geological Society of America Bulletin, v. 84, no. 1, p. 285-296.
- Davis, G. A., and Burchfiel, B. C., 1973, Garlock fault--An intracontinental transform structure, southern California: Geological Society of America Bulletin, v. 84, no. 4, p. 1407-1422.

- Dickinson, W. R., 1977, Paleozoic plate tectonics and the evolution of the Cordilleran continental margin, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 137-156.
- Dohrenwend, J. C., McFadden, L. D., Turrin, B. D., and Wells, S. G., 1984, K-Ar dating of the Cima volcanic field, eastern Mojave Desert, California--Late Cenozoic volcanic history and landscape evolution: *Geology*, v. 12, no. 3, p. 163-167.
- Drewes, Harald, 1963, *Geology of the Funeral Peak Quadrangle, California*: U.S. Geological Survey Professional Paper 413, 78 p.
- Dunne, G. C., Gulliver, R. M., and Sylvester, A. G., 1978, Mesozoic evolution of the rocks of the White, Inyo, Argus, and Slate ranges, eastern California, in Howell, D. G., and McDougall, K. A., eds., *Mesozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 2: Society of Economic Paleontologists and Mineralogists, Pacific Section*, p. 189-208.
- Eaton, G. P., 1982, The Basin and Range province--Origin and tectonic significance: *Annual Review of Earth and Planetary Science*, v. 10, p. 409-440.
- Ekren, E. B., Anderson, R. E., Rogers, C. L., and Noble, D. C., 1971, *Geology of northern Nellis Air Force Base Bombing and Gunnery range, Nye County, Nevada*: U.S. Geological Survey Professional Paper 651, 91 p.

- Ekren, E. B., Rogers, C. L., Anderson, R. E., and Orkild, P. P., 1968, Age of basin and range normal faults in Nevada Test Site and Nellis Air Force Range, Nevada, in Nevada Test Site: Geological Society of America Memoir 110, p. 247-250.
- Evernden, J. F., and Kistler, R. W., 1970, Chronology of emplacement of Mesozoic batholith complexes in California and western Nevada: U.S. Geological Survey Professional Paper 623, 42 p.
- Ferguson, H. G., and Muller, S. W., 1949, Structural geology of the Hawthorne and Tonopah Quadrangles, Nevada: U.S. Geological Survey Professional Paper 216, 55 p.
- Fleck, R. J., 1970, Age and tectonic significance of volcanic rocks, Death Valley area, California: Geological Society of America Bulletin, v. 81, no. 9, p. 2807-2816.
- Greene, G. W., 1966, Tiltmeter measurements, in Hunt, C. B., and Mabey, D. R., Stratigraphy and structure, Death Valley, California: U.S. Geological Survey Professional Paper 494-A, p. 112-114.
- Grose, T. L. T., 1959, Structure and petrology of the northeast part of the Soda Mountains, San Bernardino County, California: Geological Society of America Bulletin, v. 70, no. 12, pt. 1, p. 1509-1548.
- Hall, W. E., 1971, Geology of the Panamint Butte Quadrangle, Inyo County, California: U.S. Geological Survey Bulletin 1299, map scale 1:48,000, 67 p.

- Hardyman, R. F., Ekren, E. B., and Byers, F. M., Jr., 1975, Cenozoic strike-slip, normal, and detachment faults in northern part of Walker Lane, west-central Nevada, (abs.): Geological Society of America Abstracts with Programs, v. 7, no. 7, p. 1100.
- Hazzard, J. C., 1937, Paleozoic section in the Nopah and Resting Spring Mountains, Inyo County, California: California Journal of Mines and Geology, v. 33, no. 4, p. 273-339.
- Hewett, D. F., 1956, Geology and mineral resources of the Ivanpah Quadrangle, California and Nevada: U.S. Geological Survey Professional Paper 275, map scale 1:125,000, 172 p.
- Hopper, R. H., 1947, Geologic section from the Sierra Nevada to Death Valley, California: Geological Society of America Bulletin, v. 58, no. 5, p. 393-432.
- Hunt, C. B., and Mabey, D. R., 1966, Stratigraphy and structure, Death Valley, California: U.S. Geological Survey Professional Paper 494-A, p. A1-A162.
- Jennings, C. W., 1961, Geologic map of California--Kingman sheet: California Division of Mines and Geology, scale 1:250,000.
- Jennings, C. W., Burnett, J. L., and Troxel, B. W., 1962, Geologic map of California--Trona sheet: California Division of Mines and Geology, scale 1:250,000.
- Ketner, K. B., 1977, Late Paleozoic orogeny and sedimentation, southern California, Nevada, Idaho, and Montana, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Paleontologists and Mineralogists, Pacific Section, p. 363-369.

- Kistler, R. W., 1974, Phanerozoic batholiths in western North America--A summary of some recent work on variations in time, space, chemistry, and isotopic composition: Annual Review of Earth and Planetary Science, v. 2, p. 403-418.
- Labotka, T. C., and Albee, A. L., 1977, Late Precambrian depositional environment of the Pahrump Group, Panamint Mountains: California Division of Mines and Geology Special Report 129, p. 93-100.
- Labotka, T. C., Albee, A. L., Lanphere, M. A., and McDowell, S. D., 1980a, Stratigraphy, structure, and metamorphism in the central Panamint Mountains (Telescope Peak Quadrangle), Death Valley area, California: Geological Society of America Bulletin, v. 91, no. 3, pt. 1, p. 125-129.
- 1980b, Stratigraphy, structure, and metamorphism in the central Panamint Mountains (Telescope Peak Quadrangle), Death Valley area, California: Geological Society of America Bulletin, v. 91, no. 3, pt. 2, p. 843-933.
- Lanphere, M. A., 1964, Geochronologic studies in the eastern Mojave Desert, California: Journal of Geology, v. 72, no. 4, p. 381-399.
- Lanphere, M. A., Wasserburg, G. J. F., and Albee, A. L., 1963, Redistribution of strontium and rubidium isotopes during metamorphism, World Beater Complex, Panamint Range, California, in Craig, Herman and others, eds., Isotopic and cosmic chemistry: Amsterdam, North-Holland Publishing Co., p. 269-320.

- Lipman, P. W., Prostka, H. J., and Christiansen, R. L., 1972, Cenozoic volcanism and plate-tectonic evolution of the western United States, 1, Early and Middle Cenozoic, in A discussion on volcanism and the structure of the earth: Royal Society of London, Philosophical Transactions, Series A., v. 271, no. 1213, p. 217-248.
- Longwell, C. R., 1960, Possible explanation of diverse structural patterns in southern Nevada: American Journal of Science, v. 258A (Bradley Volume), p. 192-203.
- Longwell, C. R., Pampeyan, E. H., Bowyer, Ben, and Roberts, R. J., 1965, Geology and mineral deposits of Clark County, Nevada: Nevada Bureau of Mines Bulletin 62, 218 p.
- Luedke, R. G., and Smith, R. L., 1981, Map showing distribution, composition, and age of late Cenozoic volcanic centers in California and Nevada: U.S. Geological Survey Miscellaneous Investigations Map I-1091-C, scale 1:1,000,000, 2 sheets.
- McAllister, J. F., 1956, Geology of the Ubehebe Peak Quadrangle, California: U.S. Geological Survey Map GQ-95, scale 1:62,500.
- 1970, Geology of the Furnace Creek borate area, Death Valley, Inyo County, California: California Division of Mines and Geology Map 14, scale 1:24,000.
- McKee, E. H., 1971, Tertiary igneous chronology of the Great Basin of the western United States--Implications for tectonic models: Geological Society of America Bulletin, v. 82, no. 12, p. 3497-1501.
- Merriam, C. W., 1963, Geology of the Cerro Gordo mining district, Inyo County, California: U.S. Geological Survey Professional Paper 408, 80 p.

- Merriam, C. W., and Anderson, C. A., 1942, Reconnaissance survey of the Roberts Mountains, Nevada: Geological Society of America Bulletin, v. 53, no. 12, p. 1675-1727.
- Mifflin, M. D., and Wheat, M. M., 1979, Pluvial lakes and estimated pluvial climates of Nevada: Nevada Bureau of Mines and Geology Bulletin 94, 57 p.
- Nelson, C. A., 1966a, Geologic map of the Waucoba Mountain Quadrangle, Inyo County, California: U.S. Geological Survey Geologic Quadrangle Map GQ-528, scale 1:62,500.
- 1966b, Geologic map of the Blanco Mountain Quadrangle, Inyo and Mono Counties, California: U.S. Geological Survey Geologic Quadrangle Map GQ-529, scale 1:62,500.
- 1971, Geologic map of the Waucoba Spring Quadrangle, Inyo County, California: U.S. Geological Survey Geologic Quadrangle Map GQ-921, scale 1:62,500.
- Olson, J. C., Shawe, D. R., Pray, L. C., and Sharp, W. N., 1954, Rare earth mineral deposits of the Mountain Pass district, San Bernardino County, California, with a forward on The history of the discovery at Mountain Pass, by D. F. Hewett: U.S. Geological Survey Professional Paper 261, 75 p.
- Palmer, A. R., 1971, The Cambrian of the Great Basin and adjoining areas, western United States, in Holland, C. H., ed., Cambrian of the new world: New York, Wiley-Interscience, p. 1-78.
- Poole, F. G., 1974, Flysch deposits of Antler foreland basin, western United States: Society of Economic Paleontologists and Mineralogists Special Publication 22, p. 58-82.

- Poole, F. G., and Sandberg, C. A., 1977, Mississippian paleogeography and tectonics of the western United States, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 67-85.
- Poole, F. G., Sandberg, C. A., and Boucot, A. J., 1977, Silurian and Devonian paleogeography of the western United States, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 39-66.
- Rich, Mark, 1977, Pennsylvanian paleogeographic patterns in the western United States, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 87-112.
- Roberts, R. J., Hotz, P. E., Gilluly, J., and Ferguson, H. G., 1958, Paleozoic rocks of north central Nevada: American Association of Petroleum Geologists Bulletin, v. 42, no. 12, p. 2813-2857.
- Ross, R. J., Jr., 1977, Ordovician paleogeography of the western United States, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 19-38.

- Silberling, N. J., 1973, Geologic events during Permian-Triassic time along the Pacific margin of the United States, in Logan, A., and Hills, L. V., eds., The Permian and Triassic systems and their mutual boundary: Calgary, Alberta Society of Petroleum Geologists, p. 345-362.
- Silberling, N. J., and Roberts, R. J., 1962, Pre-Tertiary stratigraphy and structure of northwestern Nevada: Geological Society of America Special Paper 72, 58 p.
- Silver, L. T., Anderson, C. A., Crittenden, M. D., Jr., and Robertson, J. M., 1977, Chronostratigraphic elements of the Precambrian rocks of the southwestern and farwestern United States, (abs.): Geological Society of America Abstracts with Programs, v. 9, no. 7, p. 1176.
- Smith, G. I., and Church, J. P., 1980, Twentieth-century crustal deformation in the Garlock fault-Slate Range area, southeastern California: Geological Society of America Bulletin, v. 91, no. 9, pt. 1, p. 524-534.
- Smith, G. I., and Street-Perrott, F. A., 1983, Pluvial lakes of the western United States, in Wright, H. E., and Porter, S. C., eds., Late Quaternary Environments of the United States, Volume 1, The late Pleistocene: Minneapolis, University of Minnesota Press, p. 190-212.
- Smith, R. S. U., 1975, Guide to selected examples of Quaternary tectonism in Panamint Valley, California, in A field guide to Cenozoic deformation along the Sierra Nevada province and Basin and Range province boundary: California Geology, v. 28, no. 5, p. 112-117.

- Speed, R. C., 1977, Island-arc and other paleogeographic terranes of the late Paleozoic age in the western Great Basin, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 349-362.
- 1978a, Basin terrane of the early Mesozoic marine province of the western Great Basin, in Howell, D. G., and McDougall, K. A., eds., Mesozoic paleogeography of the western United States--Pacific Coast Paleogeography, Symposium 2: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 237-252.
- 1978b, Paleogeographic and plate tectonic evolution of the early Mesozoic marine province of the western Great Basin, in Howell, D. G., and McDougall, K. A., eds., Mesozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 2: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 253-270.
- Stern, T. W., Newell, M. F., and Hunt, C. B., 1966, Uranium-lead and potassium-argon ages of parts of the Amargosa thrust complex, Death Valley, California: U.S. Geological Survey Professional Paper 550-B, p. 142-147.

- Stevens, C. H., 1977, Permian depositional provinces and tectonics, western United States, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1, Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 113-136.
- Stewart, J. H., 1967, Possible large right-lateral displacement along fault and shear zones in the Death Valley-Las Vegas area, California and Nevada: Geological Society of America Bulletin, v. 78, no. 2, p. 131-142.
- 1970, Upper Precambrian and lower Cambrian strata in the southern Great Basin, California and Nevada: U.S. Geological Survey Professional Paper 620, 206 p.
- 1978, Basin and Range structure in western North America--A review, in Smith, R. B., and Eaton, G. P., eds., Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152, p. 1-31.
- 1980, Geology of Nevada: Nevada Bureau of Mines and Geology Special Publication 4, 136 p.
- Stewart, J. H., and Carlson, J. E., 1976, Cenozoic rocks of Nevada--Four maps and a brief description of distribution, lithology, age, and centers of volcanism: Nevada Bureau of Mines and Geology Map 52, scale 1:1,000,000.
- 1978, Geologic map of Nevada: U.S. Geological Survey and Nevada Bureau of Mines and Geology Map 50, scale 1:500,000.
- Basin and Range province boundary: California Geology, v. 28, no. 5, p. 112-117.

- Stewart, J. H., and Poole, F. G., 1975, Extension of the Cordilleran miogeosyncline belt to the San Andreas Fault, southern California: Geological Society of America Bulletin, v. 86, no. 2, p. 205-212.
- Stewart, J. H., and Suczek, C. A., 1977, Cambrian and latest Precambrian paleogeography and tectonics of the western United States, in Stewart, J. H., Stevens, C. H., and Fritsche, A. E., eds., Paleozoic paleogeography of the western United States--Pacific Coast Paleogeography Symposium 1: Society of Economic Paleontologists and Mineralogists, Pacific Section, p. 1-17.
- Strand, R. G., 1967, Geologic map of California--Mariposa sheet: California Division of Mines and Geology, scale 1:250,000.
- Streitz, Robert, and Stinson, M. C., compilers, 1974, Geologic map of California--Death Valley sheet: California Division of Mines and Geology, scale 1:250,000.
- Thompson, G. A., and Burke, D. B., 1974, Regional geophysics of the Basin and Range Province: Annual Review of Earth and Planetary Science v. 2, p. 213-238.
- Troxel, B. W., 1967, Sedimentary rocks of late Precambrian and Cambrian age in the southern Salt Spring Hills, southeastern Death Valley, California: California Division of Mines and Geology Special Report 92, p. 32-41.
- Tschanz, C. M., and Pampeyan, E. H., 1970, Geology and mineral deposits of Lincoln County, Nevada: Nevada Bureau of Mines Bulletin 73, 187 p.

- Volborth, Alexis, 1973, Geology of the granite complex of the Eldorado, Newberry, and northern Dead Mountains, Clark County, Nevada: Nevada Bureau of Mines and Geology Bulletin 80, 40 p.
- Wasserburg, G. J. F., Wetherill, G. W., and Wright, L. A., 1959, Ages in the Precambrian terrane of Death Valley, California: Journal of Geology, v. 67, no. 6, p. 702-708.
- Williams, E. G., Wright, L. A., and Troxel, B. W., 1976, The Noonday Dolomite and equivalent stratigraphic units, southern Death Valley region, California, in Troxel, B. W., and Wright, L. A., eds., Geologic features, Death Valley, California: California Division of Mines and Geology Special Report 106, p. 45-50.
- Wright, L. A., Troxel, B. W., Williams, E. G., Roberts, M. T., and Diehl, P. E., 1976, Precambrian sedimentary environments of the Death Valley region, eastern California, in Troxel, B. W., and Wright, L. A., eds., Geologic features, California: California Division of Mines and Geology Special Report 106, p. 7-17.
- Wright, L. A., Troxel, B. W., Burchfiel, B. C., Chapman, R. H., and Labotka, T. C., 1981, Geologic cross section from the Sierra Nevada to the Las Vegas Valley, eastern California to southern Nevada: Geological Society of America Map and Chart Series MC-28M, scale 1.25 inches = 5 miles.
- Wright, L. A., and Troxel, B. W., 1983, Geologic map and section of the north 1/2 Confidence Hills 15' Quadrangle, Inyo County, California: California Division of Mines and Geology Map 34, scale 1:24,000.

Zoback, M. L., Anderson, R. E., and Thompson, G. A., 1981,
Cainozoic evolution of the state of stress and style of
tectonism of the Basin and Range province of the western
United States, in Vine, F. J., convenor, Extensional
tectonics associated with convergent plate boundaries: Royal
Society of London, Philosophical Transactions, Series A., v. 300,
no. 1454, p. 407-434.

POTENTIAL HOST MEDIA FOR RADIOACTIVE WASTE

By

K. A. Sargent, U.S. Geological Survey

Host media considered to have potential in the Death Valley region, Nevada and California, include granite and other coarse-grained plutonic rocks, ash-flow tuff, basaltic and andesitic lava flows. A few shallow, fine-grained silicic intrusions occur in the region that may have potential as host rocks. Argillaceous sedimentary and metasedimentary rocks generally are steeply dipping, folded, or complexly faulted and would need to be carefully evaluated for potential as host rock. Salt and other evaporitic deposits of sufficient thickness and depth for a host rock are not known to occur in the region. Most prospective host rocks mentioned above and basin fill have potential as host media in the unsaturated zone. Outcrop areas of potential host rock and areas believed to have thick unsaturated zones in the Death Valley region, Nevada and California, are shown in plate 2.

right, L. A., Trowel, P. M., Burchfiel, B. C., Chapman, R. H., and Labotka, T. G., 1981, Geologic cross section from the Sierra Nevada to the Las Vegas Valley, eastern California to southern Nevada: Geological Society of America Map and Chart Series MC-26M, scale 1:25 inches = 5 miles.

right, L. A., and Trowel, P. M., 1983, Geologic map and section of the north 1/2 Confidence Hills 15' Quadrangle, Inyo County California: California Division of Mines and Geology Map 34, scale 1:24,000.

Intrusive Rocks

Granitic rocks of Tertiary, Mesozoic, and Precambrian age occur widely throughout the region (pl. 2). For a summary of granitic rocks in the Death Valley region see Hills and Lopez (1984), Sargent and Roggensack (1984), and Grose and Smith, this chapter, **Early and Middle Proterozoic Crystalline Basement Rocks and Mesozoic and Cenozoic Plutonic Rocks**. Precambrian rocks are granitic gneiss and schistose rocks, commonly extensively sheared and mylonitized. They are the least common in ground-water unit DV-01.

In ground-water unit DV-01, Mesozoic and a few Tertiary intrusive rocks occur as massive plutons; some of the larger bodies will be discussed here. The Boulder City pluton is a 13-million-year-old granodiorite that intrudes Tertiary volcanic rocks. The pluton is extensively faulted and brecciated. Other large Tertiary plutons in ground-water unit DV-01 occur in the Eldorado and Newberry Mountains. These bodies intrude Precambrian granitic gneiss and schist and locally are transected by rhyolite and diabase dikes. In southwesternmost ground-water unit DV-01 and into ground-water unit DV-03, part of the complex Teutonia batholith is present. It is composed of a considerable variety of coarse-grained igneous-rock types such as hornblende gabbro, quartz monzodiorite, granodiorite, monzogranite, and syenogranite. The complex is of middle to late Mesozoic age and possibly post dates much of the major thrusting and mylonitization in the area.

The Kingston Range biotite monzogranite is a relatively large Tertiary intrusive body located in the western part of ground-water unit DV-02 and in the southeastern part of ground-water unit DV-03. Its large area of unsaturated rocks and young age make this granite a good prospect for further investigation.

Ground-water unit DV-03, the largest unit, has the smallest density of exposed granitic masses. Widely scattered and sparse in the northern part of the unit, the intrusions are Precambrian, Mesozoic, and Tertiary in age. In the northern part of ground-water unit DV-03, the Cactus Range pluton of granodioritic and melanodioritic composition and Tertiary age is extensively propylitized and altered. The Climax and Gold Meadows stocks are hydrothermally altered Cretaceous quartz monzonite located in the Nevada Test Site. The Climax stock is the site of a U.S. Department of Energy test for the storage of spent nuclear fuel. The tests are being conducted at about 420 m below the surface which is above the water table. Elsewhere in southern Nye County are two relatively small exposures of coarse-grained intrusive rocks associated with calderas.

In Esmeralda County, Nevada, western part of ground-water unit DV-03, the Sylvania pluton, a Jurassic coarse-grained adamellite, intrudes Precambrian sedimentary rocks. Numerous small exposures of Tertiary and Mesozoic granitic and fine-grained siliceous intrusive rocks occur in the region, but are not well described in the literature. Granite in the California part of the unit is mostly of Mesozoic age. The Mesozoic granite is described as locally fractured or sheared; however, some quartz monzonite is massive and unfoliated, such as in the Soda Mountains (southernmost part of ground-water unit DV-03) and at Manly Peak, southern Panamint Range, along the border with ground-water unit DV-04. A large pluton southeast of Silurian Lake is a massive, unfoliated quartz monzonite believed to have been emplaced after thrust faulting. A quartz monzonite near Sugarloaf Peak southwest of Death Valley is reported as a massive body. Tertiary biotite monzogranite and granite east and west of Death Valley in the Black Mountains and in the Panamint Range may be structurally sound. These Tertiary bodies intrude foliated and faulted Precambrian metasedimentary and metaigneous rocks. Large Jurassic or Cretaceous granitic plutons or both occur in the Granite, Avawatz, and Owlshhead Mountains in the southern part of ground-water unit DV-03; more data are needed should further investigation be required in that area.

The Triassic intrusive rock of Cottonwood Mountains, western part of ground-water unit DV-03, northern part of ground-water unit DV-04, and southeastern part of ground-water unit DV-05, is the Hunter Mountain pluton composed of quartz monzonite. The Pluton has more than 1,200 m of relief. Numerous smaller Plutons, laccoliths and plugs are present in ground-water unit DV-03. Many could be large enough at depth to be suitable for repository siting.

In ground-water unit DV-04, granitic rocks are common in all the bounding ranges. In the Slate, Argus, and Coso Ranges, on the west side of the unit, Mesozoic quartz monzonite is widespread. Locally it is intruded by swarms of dikes. In the Slate Range the pluton is composed of flat-lying granitic sheets a few to hundreds of meters thick. As described in the discussion of ground-water unit DV-03, the Jurassic pluton at Manly Peak is unshattered and unbrecciated; however, topographically lower, and exposed just above the floor of Panamint Valley, the Triassic granodiorite is sheared and brecciated.

In ground-water unit DV-05 around Saline Valley, there are several large plutons. One large body west of the valley is the Paiute Monument pluton, a hornblende-biotite monzogranite of Jurassic age. It intrudes Jurassic Hunter Mountain Quartz Monzonite (mentioned in the discussion of ground-water unit DV-03). A prospective pluton occurs on the divide with Death Valley and Panamint Valley in the southeastern part of the unit. This is the western end of the Hunter Mountain pluton mentioned in the discussion of ground-water unit DV-03.

On the divide between ground-water units DV-05 and DV-06 is the King Papoose Flat pluton of Cretaceous age. It is foliated and locally faulted. In the central part of ground-water unit DV-06 is a Jurassic pluton. It is composed of monzonite and diorite. In ground-water units DV-06 and DV-08, a large monzonite to quartz monzonite stock of Jurassic age occurs. This mass, which forms much of the White Mountains, is composed of multiple Mesozoic intrusions and appears to have few structural complications.

In ground-water unit DV-07, the largest plutons exposed include the Jurassic Palmetto pluton (southwestern part of ground-water unit DV-07) and the Cretaceous Belmont and Manhattan plutons (northwestern part of ground-water unit DV-07). These plutons are extensive, locally porphyritic, and unfoliated.

Smaller plutons of Triassic to Tertiary age, range in composition from granite to diorite. Most are unfoliated although they are locally transected by rhyolite and diabase dikes.

Granitic rocks in ground-water unit DV-08 consist largely of Jurassic and Cretaceous monzonite, quartz monzonite, and adamellite plutons. Very little data exists on their structural condition. The large plutons are the Palmetto (southeast part of ground-water unit DV-08) and unnamed bodies near the California-Nevada line.

Mostly small granite exposures occur in ground-water unit DV-09. Exceptions are the Lone Mountain Granite and quartz monzonite west of Tonopah, Nevada. This large body is of Cretaceous age (63-71 million years) and is transected by diabase dikes. Less than 10 km to the southwest of the Lone Mountain Pluton is the Weepah pluton, a Mesozoic quartz monzonite.

Tuffaceous Rocks

In California, welded to non-welded tuffs of limited extent and minimal thickness occur in the northern part of ground-water units DV-04, DV-05, and the eastern part of ground-water unit DV-06. Mixed volcanic rocks of Miocene age, possibly containing tuffs as thick as 400 m, occur in ground-water DV-03 in the Grapevine Mountains along the Nevada-California State line. In the southern part of ground-water unit DV-01, thin ash-flow tuffs of possibly andesitic composition occur in the Sacramento Mountains and vicinity. Tuffs in California are summarized by Jenness and Lopez (1985).

In Nevada, tuffs are very widespread in large outcrop areas north of 37° N Latitude (Sargent, 1984). A few outcrops of welded tuff occur in ground-water unit DV-01. North of Nelson, Miocene rhyolitic ash-flow tuffs are as thick as 250 m. In Eldorado Valley and in the Highland Range, the same tuffs (Tuff of Bridge Spring) are 120 m thick and occur where the depth to water is greater than 150 m.

In the northern part of ground-water unit DV-03 there are abundant tuffs having aggregate thicknesses commonly greater than 1,200 m. Within the Silent Canyon caldera, at Pahute Mesa, drill holes have penetrated more than 4,100 m of volcanic rock, much of it tuff, but also including silicic lava flows. Here the unsaturated zone is as much as 700 m thick. The tuffs at Pahute Mesa include Miocene densely to non-welded ash-flow tuff, air-fall tuff, and reworked tuff. Tuff sections generally are thickest within calderas and in topographic lows adjacent to caldera source areas.

Great thicknesses of massive ash-flow tuff occur in the northern and northeastern part of ground-water unit DV-07 in the Cathedral Ridge (2,400 m) and Kawich calderas (1,000 m). The Bald Mountain caldera in the northeastern part of ground-water unit DV-03, and the Timber Mountain-Oasis Valley caldera complex in the central part of ground-water unit DV-03, also contain thick ash-flow tuffs. As shown in figure 3, thick unsaturated sections are present in these caldera areas.

In ground-water unit DV-08, the Silver Peak caldera in the Silver Peak Range, contains tuffs as thick as 600 m and a thick unsaturated section. Tuffs adjacent to the caldera may be as thick as 450 m.

In ground-water units DV-07 and DV-09, tuffs are widespread although their caldera sources are not well known. The Toiyabe Quartz Latite of Miocene age, is widespread and commonly greater than 300 m thick in the northern part of ground-water unit DV-09. The tuffs of Rye Patch, probably of Oligocene age, may be as thick as 1,800 m in the southern Monitor Range (ground-water unit DV-07), site of a possible caldera. A large part of the southern Monitor Range is unsaturated.

Basaltic Rocks

Tertiary basalts are widely distributed throughout the Death Valley region (see Roggensack and Sargent, 1985; and Roggensack and Lopez, 1985). A few areas are the sites of large outpourings of mafic lava. Generally the aggregate thickness is about 300 m, but in places it may be as much as 900 m. The thick occurrences are briefly pointed out here. Virtually all the flows are middle Miocene or younger. The great majority of basaltic and andesitic flows, however, are less than 60 m in aggregate thickness and have little or no potential for repository siting.

The eastern part of ground-water unit DV-01 contains unusually thick (700 to 900 m) andesitic lavas and volcanic breccia of Miocene age in the Black Mountains northwest of Lake Mead. Thick basaltic flows also are present in the Mount Davis Volcanics in the McCullough Range and Eldorado Mountains of ground-water unit DV-01. These volcanic rocks are 11 to 14 million years old (Miocene) and have aggregate thicknesses of about 600 m.

Scattered, mostly small, thin basaltic and andesitic flows occur in ground-water unit DV-03. Of these, the most extensive and thickest flows appear to be Pliocene trachyandesite and trachybasalt on the southeast flank of Timber Mountain where the aggregate thickness is about 300 m, and at a basalt dome west of Timber Mountain where flows may be as much as 250 m thick. In the Greenwater Range, California, andesite and basalt flows may be as much as 150 m thick, but the rocks are extensively altered and fragmented.

In the northern part of ground-water unit DV-04 and in adjacent ground-water unit DV-03, both east and west of the northern end of Panamint Valley, Pliocene and Miocene olivine basalt is widely distributed. The total thickness is more than 150 m and the Tertiary flows locally are overlain by Quaternary basalt. In the Saline Range, in the northern part of ground-water unit DV-05 and in the southern part of ground-water unit DV-06, extensive flows of Miocene to Pliocene olivine basalt, andesite and trachyandesite occur that may be as much as 300 m thick.

Extensive Miocene trachyandesite flows occur in the San Antonio Mountains in the northwestern part of ground-water unit DV-07 and in the eastern part of ground-water unit DV-09. Here flows have an aggregate thickness of about 300 m and are partly in a thick unsaturated section. In the Goldfield Hills, central part of ground-water unit DV-07, more than 240 m of basalt may be present under the Thirsty Canyon Tuff.

In ground-water unit DV-08, the Silver Peak calderami, 08hewd, Range, contains tuffs as thick as 600 m and a thick unsaturated

In the northern part of ground-water unit DV-09, at Sherman Peak, a trachyandesite and ash-flow tuff sequence is reported to have a combined thickness of about 550 m.

Argillaceous Rocks

In Nevada, outcrops of argillaceous rocks of Paleozoic age are scattered throughout ground-water unit DV-03, and in the northern border area between ground-water units DV-07 and DV-09. The argillaceous rocks are tectonically deformed, faulted, and sheared. Because of their structural complexity, their continuity is difficult to define.

Precambrian metamorphosed argillaceous rocks occur in ground-water units DV-02, DV-03, and DV-04. Small scattered outcrops of Cambrian and Mississippian argillaceous rocks occur in ground-water units DV-03, DV-05, and DV-06. Argillaceous rocks are summarized for the California part of the Death Valley region by Johnson (1985), and for the Nevada part by Simpson and others (1979).

southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-G, scale 1:500,000.

Weggenack, Kurt, and Sargent, K. A., 1984, Map showing outcrops of basaltic rocks of early Quaternary and Tertiary ages,

Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations 83-4119-F, scale 1:500,000.

Sargent, K. A., 1985, Map showing outcrops of dominantly ash-flow tuff of pre-Quaternary age, Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations 83-4111, scale 1:500,000.

Unsaturated Zone

The Death Valley region contains the greatest total area and largest contiguous area of unsaturated rocks in the entire Basin and Range province. Depth to water, as confirmed by drill-hole data in southern Nye County, Nevada, is as great as 700 m, and numerous holes drilled in basin fill and tuff penetrated more than 450 m of unsaturated section. The great majority of unsaturated rock is in the eastern part of ground-water unit DV-03 and in the central part of ground-water unit DV-01, but relatively large areas of unsaturated rock are present in all of the ground-water units of the region (pl. 2). The primary host media in the unsaturated zone are tuff, granite, basalt, and basin fill.

Extensive Miocene trachyandesite flows occur in the San Antonio Mountains in the northwestern part of ground-water unit DV-07 and in the eastern part of ground-water unit DV-09. Here flows have an aggregate thickness of about 300 m and are partly in a thick unsaturated section. In the Goldfield Hills, central part of ground-water unit DV-07, more than 240 m of basalt may be present under the Thirsty Canyon Tuff.

References Cited

- Hills, F. A., and Lopez, D. A., 1984, Map showing outcrops of granitic and silicic shallow intrusive rocks, Basin and Range province, southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-D, scale 1:500,000.
- Jenness, J. E., and Lopez, D. A., 1984, Map showing outcrops of dominantly ash-flow tuff, Basin and Range province, southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-F, scale 1:500,000.
- Johnson, W. D., 1984, Map showing outcrops of thick, dominantly argillaceous sedimentary and metasedimentary rocks, Basin and Range province, southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-E, scale 1:500,000.
- Roggensack, Kurt, and Lopez, D. A., 1984, Map showing outcrops of pre-Quaternary basaltic rocks, Basin and Range province, southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-G, scale 1:500,000.
- Roggensack, Kurt, and Sargent, K. A., 1984, Map showing outcrops of basaltic rocks of early Quaternary and Tertiary ages, Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations 83-4119-F, scale 1:500,000.
- Sargent, K. A., 1985, Map showing outcrops of dominantly ash-flow tuff of pre-Quaternary age, Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations 83-4119-E, scale 1:500,000.

Sargent, K. A., and Roggensack, Kurt, 1985, Map showing outcrops of granitic rocks, Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations Report 83-4119-D, scale 1:500,000.

Simpson, H. E., Weir, J. W., Jr., and Woodward, L. A., 1979, Inventory of clay-rich bedrock and metamorphic derivatives in eastern Nevada, excluding the Nevada Test Site: U.S. Geological Survey Open-File Report 79-760, 147 p.

QUATERNARY TECTONISM

by K. A. Sargent, U.S. Geological Survey, and
T.L.T. Grose, Nevada Bureau of Mines and Geology and
Colorado School of Mines

Evidence of Quaternary tectonic conditions of the Death Valley region include data on seismicity, heat flow, Quaternary faulting, late Cenozoic volcanic activity and vertical movement. Each of these features is depicted in figure 7.




Figure 7 is a map of the Death Valley region, showing Quaternary tectonic features. The map includes the Death Valley, Furnace Creek, and the surrounding area. It shows major faults like the Furnace Creek Fault and the Las Vegas shear zone. It also indicates seismicity with circles of varying sizes representing different magnitudes. The map is labeled with 'Death Valley', 'Furnace Creek', and 'Las Vegas shear zone'. A scale bar is present in the bottom left corner.




FIGURE 7.-- Quaternary tectonic features in the Death Valley region and vicinity, Nevada and California

Seismicity

A compilation of epicenters for the Death Valley region, by Algermissen and others (1983), shows several hundred recorded earthquakes. Of these, 37 have Richter magnitudes (surface waves) of 5 or greater, and all but 17 of these occur in the Nevada Test Site where underground nuclear tests at Yucca Flat and Pahute Mesa are known to have produced man-made earthquakes (fig. 7). Several of the earthquakes at the Nevada Test Site with magnitudes of more than 4 are natural. Of the 17 naturally occurring earthquakes of magnitude 5 or greater, most have their epicenters along the western side of the region. Only one of these larger earthquake epicenters (magnitude 5 or greater) appears to coincide with the surface expression of the Death Valley-Furnace Creek fault. None appear to coincide with the Garlock fault or the Las Vegas shear zone. Even the smaller earthquakes, those with magnitudes less than 5 (see Algermissen and others, 1983), appear to have no particular correspondence with these large faults. Two occur in the Lake Mead area and may be related to a hydraulic connection between impounded lake water and a deep-aquifer system along buried faults (Anderson and Laney, 1975). Outside the Nevada Test Site, the two largest earthquakes, magnitudes 6 to 7, were east of Owens Valley and southwest of Columbus Salt Marsh in Esmeralda County, Nevada, and together with swarms of smaller earthquakes form two large areas with significant strain release in the western and northwestern parts of the region.

The Death Valley region occurs within a broad, rather vaguely defined, region of relatively moderate earthquake frequency and areal density, moderate cumulative seismic-strain energy release, and oblique slip between a strike-slip dominated region to the west and a dip-slip dominated region to the east (Eaton, 1980; Smith, 1978). Fault-plane solutions indicate significant right-lateral-slip movement in the western part of the Death Valley region and more oblique- or dip-slip movements in the eastern part (Slemmons and others, 1979; Smith and Lindh, 1978), which correspond with geologic observations on faults in the region. Both fault-plane solutions on seismic events and fault geometries indicate a general west-northwest direction of seismotectonic extension in the Death Valley region. This agrees well with other lines of independent evidence of state of stress in the region (Zoback and Zoback, 1980). Throughout the region, focal depths of the earthquakes are usually less than 15 km, being slightly deeper in the western part of the region where strike-slip movement dominates (Eaton, 1980).

Various seismogenic regionalization studies in the Death Valley region indicated by the most notable concentration of earthquake epicenters (Askew and Algermissen, 1983; Ryall, 1977), the most rapid extensional strain rate (Greensfelder and others, 1980), and the greatest number of late Quaternary and historic faults (Nakata and others, 1982) occur in the northwestern part of the Death Valley region. This area is in the southeastern part of the Nevada seismic zone (Gumper and Scholz, 1971; Ryall and others, 1966; Wallace, 1978), the most active seismic zone in Nevada. Seismic activity decreases in a southeasterly direction through the Death Valley region.

Heat Flow

Heat-flow measurements for 39 sites are reported for the Death Valley region (J. H. Sass, U.S. Geological Survey, written commun., 1982). Only two of the measurements exceed 2.5 HFU (heat-flow units) (fig. 7). Both are in Inyo County, California, one northwest of Death Valley (3.0 HFU), the other northeast of Owens Lake (18.70 HFU). This large value probably is associated with a local fault-controlled geothermal convection system. The heat-flow map of Sass (shown in Sargent and Bedinger, 1985, fig. 16) shows the northeastern part of the Death Valley region to have values less than 1.5 HFU, and the remainder of the region to have values between 1.5 and 2.5 HFU. The unusually small values in the northeastern part of the region are part of the Eureka heat-flow low of southeast central Nevada and are believed to be caused by convective loss of heat from an area of unusually well-developed ground-water circulation through carbonate aquifers (Lachenbruch and Sass, 1977; and Sass and Lachenbruch, 1982).

Also the regional heat-flow map shows two small areas in the far northwestern part of the Death Valley region that are interpreted to have greater than 2.5 HFU. Those areas are marginal to the large region of substantial heat flow called the Battle Mountain heat-flow high (Sass and others, 1971).

Within the Death Valley region, conductive heat flow locally is modified by thermal springs. Relative to most adjacent areas, thermal springs are few in number and cool (Berry and others, 1980; Garside and Schilling, 1979; Waring, 1965). Only one at Tecopa, in southeastern Inyo County, California, indicates a hydrothermal convection system with temperature greater than 90 °C (Muffler, 1979).

Quaternary Faulting

In regional compilations, Nakata and others (1982), and Jennings (1975), show Quaternary faults unevenly distributed in the Death Valley region (fig. 7). The greatest concentration and longest traces of these faults coincide with, or lie within, the mapped zones of pre-Quaternary faults; examples are the Panamint Valley fault, Garlock fault, and the Death Valley-Furnace Creek fault zone. In addition, the sense of Quaternary displacement on all three faults is the same as the older sense of displacement, apparently indicating continuing or renewed stress fields similar to those in the geologic past. There is no clear correspondence between the location of earthquakes of magnitude less than 4 and Quaternary faults; only in the northwestern part of the region is there a correspondence between the location of the larger earthquakes and Quaternary faults. Faults with historical movement are found in or near Groom Lake, Yucca Flat, and Frenchman Flat, all in or close to the Nevada Test Site. Fault segments showing displacements, no older than 10,000 years, occur along the Panamint Valley fault, the Garlock fault, the Yucca fault, and two unnamed faults southeast of Beatty, Nevada.

Late Cenozoic Volcanics

Much of the late Cenozoic volcanic activity is along the southern and western edges of the Death Valley region (fig. 4). Most of the volcanic rocks are basaltic flows and cinder cones with minor andesitic, dacitic and rare rhyolitic flows. Dates as young as 0.25 million years have been obtained for basaltic rocks in Crater Flat southwest of the Nevada Test Site, and 0.45 million years for a small basaltic outcrop west of the Timber Mountain-Oasis Valley caldera complex. The northern end of the Cima volcanic field occurs near the southern edge of the region. The field consists of flows of hawaiite, alkali-olivine basalt, and basanite of late Miocene to Holocene age (Dohrenwend and others, 1984; Katz and Boettcher, 1980). In other areas, rocks are assigned an age less than 5 million years old by stratigraphic position and geologic association with dated rocks (Luedke and Smith, 1981). There appears to be little coincidence of late Cenozoic volcanic activity with Quaternary faulting or recorded substantial seismicity.

Vertical Crustal Movement

Gable and Hatton (1983) depict a northwest-trending area that encompasses Death Valley with vertical movement of an adjacent range to the west at a rate of 20 m per 10,000 years (2.0 mm/yr). One point within Death Valley is shown to be rising at a rate of 20 to 70 m per 10,000 years (2.0-7.0 mm/yr), based on geology, geomorphology, and radiocarbon dates (fig. 7). The greater rate of vertical uplift is in a zone that is parallel to the Sierra Nevada and may be related to the erosion and isostatic adjustment of this great uplifted mountain mass.

Gable and Hatton (1983) show a zone of historic subsidence in the vicinity of Lake Mead of as much as 2 m based on leveling data from 1935 to 1950, although the general area is rising at a rate of 10 mm/yr. The subsidence related to the filling of Lake Mead is summarized by Anderson and Laney (1975).

Photolineations

Studies of linear features by T. W. Offield (U.S. Geological Survey, written commun., 1983) in the Great Basin using Landsat multispectral scanner images show that numerous photolineations parallel or coincide with Quaternary faults, as well as with known older faults. The longest photolineations are expressions of range-front faults and the Las Vegas shear zone. Linear features seen in Landsat images are alignments of both topographic and tonal features; many are related to tectonism and erosion, such as slope breaks caused by faulting; some are stratigraphically controlled erosional features; and still others are of unknown origin.

References Cited

- Algermissen, S. T., Askew, B. L., Thenhaus, P. C., Perkins, D. M., Hanson, S., and Bender, B. L., 1983, Seismic energy release and hazard estimation in the Basin and Range Province: U.S. Geological Survey Open-File Report 83-358, 13 p.
- Anderson, R. E., and Laney, R. L., 1975, The influence of late Cenozoic stratigraphy on distribution of impoundment-related seismicity at Lake Mead, Nevada-Arizona: U.S. Geological Survey Journal of Research, v. 3, no. 3, p. 337-343.
- Askew, Bonny, and Algermissen, S. T., 1983, An earthquake catalog for the Basin and Range province, 1803-1977: U.S. Geological Survey Open-File Report 83-86, 21 p.
- Berry, G. W., Grim, P. J., and Ikelman, J. A., 1980, Thermal springs list for the United States: National Oceanic and Atmospheric Administration Key to Geophysical Records Documentation 12, NOAA/NGSDC, 59 p.
- Dohrenwend, J. S., McFadden, L. D., Turrin, B. D., and Wells, S. G., 1984, K-Ar dating of the Cima volcanic field, eastern Mojave Desert, California--Late Cenozoic volcanic history and landscape evolution: Geology, v. 12, no. 3, p. 163-167.
- Eaton, G. P., 1980, Geophysical and geological characteristics of the crust of the Basin and Range province, in Continental tectonics: Washington, D. C., National Academy of Science, U.S. National Research Council, Geophysics Study Committee, p. 96-114.

- Gable, D. J., and Hatton, Tom, 1983, Maps of vertical crustal movements in the conterminous United States over the last 10 million years: U.S. Geological Survey Miscellaneous Investigations Series Map I-1315, scale 1:5,000,000, 2 sheets.
- Garside, L. J., and Schilling, J. H., 1979, Thermal waters of Nevada: Nevada Bureau of Mines and Geology Bulletin 91, 163 p.
- Greensfelder, R. W., Kintzer, F. C., and Somerville, M. R., 1980, Seismotectonic regionalization of the Great Basin, and comparison of movement rates computed from Holocene strain and historic seismicity: Geological Society of America Bulletin, pt. 2, v. 91, no. 9, p. 2039-2111.
- Gumper, F. J., and Scholz, C., 1971, Microseismicity and tectonics of the Nevada seismic zone: Seismological Society of America Bulletin, v. 61, no. 5, p. 1413-1432.
- Jennings, C. W., compiler, 1975, Fault map of California with locations of volcanoes, thermal springs, and thermal wells: California Division of Mines and Geology Geologic Data Series Map 1, scale 1:750,000.
- Katz, Marvin, and Boettcher, Art, 1980, The Cima volcanic field, in Fife, D. L., and Brown, A. R., eds., Geology and mineral wealth of the California Desert: South Coast Geological Society, Dibblee Volume, p. 236-241.
- Lachenbruch, A. H., and Sass, J. H., 1977, Heat flow in the United States and the thermal regime of the crust, in Heacock, J. G., ed., Nature and physical properties of the Earth's crust: American Geophysical Union Geophysical Monograph 20, p. 626-675.

- Luedke, R. G., and Smith, R. L., 1981, Map showing distribution, composition, and age of late Cenozoic volcanic centers in California and Nevada: U.S. Geological Survey Miscellaneous Investigations Series Map I-1091-C, scale 1:1,000,000.
- Muffler, L. J. P., ed., 1979, Assessment of geothermal resources of the United States--1978: U.S. Geological Survey Circular 790, 163 p.
- Nakata, J. K., Wentworth, C. M., and Machette, M. N., 1982, Quaternary fault map of the Basin and Range and Rio Grande rift provinces, western United States: U.S. Geological Survey Open-File Report 82-579, scale 1:2,500,000, 2 sheets.
- Ryall, Alan, 1977, Earthquake hazard in the Nevada region: Seismological Society of America Bulletin, v. 67, no. 2, p. 517-532.
- Ryall, Alan, Slemmons, D. B., and Gedney, L. D., 1966, Seismicity, tectonism, and surface faulting in the western United States during historic time: Seismological Society of America Bulletin, v. 56, no. 5, p. 1105-1135.
- Sargent, K. A., and Bedinger, M. S., 1985, Geologic and hydrologic characterization and evaluation of the Basin and Range Province relative to the disposal of high-level radioactive waste--Part II, Geologic and hydrologic characterization: U.S. Geological Survey Circular 904-B, 30 p.
- Sass, J. H., and Lachenbruch, A. H., 1982, Preliminary interpretation of thermal data from the Nevada Test Site: U.S. Geological Survey Open-File Report 82-973, 29 p.

- Sass, J. H., Lachenbruch, A. H., Munroe, R. J., Greene, G. W., and Moses, T. H., Jr., 1971, Heat flow in the western United States: *Journal of Geophysical Research*, v. 76, no. 26, p. 6376-6413.
- Slemmons, D. B., Van Warner, D., Bell, E. J., and Silberman, M. L., 1979, Recent crustal movements in the Sierra Nevada-Walker Lane region of California-Nevada, Part 1, Rate and style of deformation, in Whitten, C. A., ed., Recent crustal movements, 1977: *Tectonophysics*, v. 52, no. 1-4, p. 561-570.
- Smith, R. B., 1978, Seismicity, crustal structure, and intraplate tectonics of the interior of the western Cordillera, in Smith, R. B., and Eaton, G. P., eds., *Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152*, p. 111-144.
- Smith, R. B., and Lindh, A. G., 1978, Fault-plane solutions of the western United States--A compilation, in Smith, R. B., and Eaton, G. P., eds., *Cenozoic tectonics and regional geophysics of the western Cordillera: Geological Society of America Memoir 152*, p. 107-109.
- Wallace, R. E., 1978, Patterns of faulting and seismic gaps in the Great Basin province, in *Proceedings of Conference 6, Methodology for identifying seismic gaps and soon-to-break gaps: U.S. Geological Survey Open-File Report 78-943*, p. 857-868.
- Waring, G. A., 1965, Thermal springs of the United States and other countries of the world--A summary: *U.S. Geological Survey Professional Paper 492*, 383 p.

GROUND-WATER HYDROLOGY

By

M. S. Bedinger, William H. Langer and J. E. Reed,

U.S. Geological Survey

Climate of the Death Valley region is arid to semiarid. Precipitation on the valley floors of the Amargosa Desert, Death Valley, and basins at lower altitudes in the southern part of the region is less than 70 mm/yr on the average. The annual average precipitation at Furnace Creek Ranch in Death Valley is 50 mm/yr. Precipitation in the mountain ranges is greater, commonly in the range from 100 to 150 mm/yr. Annual precipitation is as much as 500 to 750 mm in the Sheep Range and Spring Mountains, the highest ranges in the region. Annual free-water surface evaporation is greater than 2,500 mm/yr in Death Valley.

Major Hydrogeologic Units

Basin fill (including the alluvial material of stream valleys) was deposited largely in structural basins. The fill in many basins is greater than 1,300 m thick and may be as thick as 2,000 m. The basin fill consists mostly of nonindurated to semi-indurated sedimentary terrestrial deposits and volcanic material of late Tertiary to Holocene age. The fill also contains volcanic flows and ash falls from episodic volcanic activity during the Tertiary and Quaternary Periods, as well as most consolidated early to middle Tertiary sediments. Fine-grained lake deposits, silt, clay, and evaporitic deposits occur in some of the basins.

Volcanic rocks are grouped hydrologically as three units, (1) ash-flow tuffs, (2) volcanic flows, and (3) undifferentiated volcanic rocks, all of Tertiary and Quaternary age. Tuffs, of Tertiary age, are widespread in the northern and central parts of the region, with an aggregate thickness of more than 4,000 m. They range from densely welded to nonwelded, bedded, reworked, and air falls. The following discussion of ash-flow tuffs is from Winograd (1971). Welded ash-flow tuffs characteristically have and interstitial porosity of about 5 percent or less. The permeability of welded ash-flow tuff, which commonly is moderate to large, is largely a function of jointing, bedding-plane openings, and partings within the flows. Welded ash-flow tuffs may be important units in the ground-water flow systems of the regions, especially in ground-water unit DV-03 where they are thick and of great areal extent. In contrast, nonwelded ash-flow tuffs may have a large interstitial porosity and small interstitial permeability and function as confining beds. Fractures and joints are virtually absent.

Lava flows primarily are basalt and other mafic rocks of Tertiary and Quaternary age. Columnar jointing and platy fractures are common in the flows that are vesicular to dense. Permeability and porosity is developed along fractures and bedding planes. Individual flows generally are less than 33 m thick; some are less than 1 m thick. Aggregate thicknesses are as much as 1,000 m.

The central part of the Death Valley region is underlain by one of the thickest known sequences of Paleozoic rocks in the Basin and Range province. Over 8,000 m of Paleozoic sedimentary rocks are exposed. Winograd and Thordarson (1975), in the area here referred to as ground-water unit DV-03, distinguished in this sequence, from bottom to top, as a lower clastic confining bed, a lower carbonate aquifer, an upper clastic confining bed, and an upper carbonate aquifer. Upper Precambrian and Lower Cambrian quartzite, shale, and siltstone comprise the lower clastic confining bed. The lower carbonate aquifer is composed of the carbonate rocks of Middle Cambrian age and ranges in saturated thickness from a hundred to a few thousand meters. Argillite, quartzite, and conglomerate of Late Devonian and Mississippian age, comprise the upper-clastic confining bed, which ranges from 1,300 to 2,600 m in thickness. Carbonate rock of Pennsylvanian and Permian age forms the upper carbonate aquifer. The lower carbonate aquifer is the more extensive aquifer occurring in a large part of ground-water unit DV-03 and in the northern part of ground-water unit DV-01. Here the lower carbonate aquifer is absent or unsaturated only in the outcrop areas or structural highs. Where the lower carbonate aquifer is absent, the lower clastic confining bed is a barrier to regional ground-water flow (Winograd and Thordarson, 1975, pl. 1). The saturated extent of the upper carbonate aquifer in ground-water unit DV-03 is limited to small areas in south-central Nevada and it does not have a large effect on regional ground-water flow. Similarly, the upper clastic confining bed is of limited distribution and of local significance to regional ground-water flow.

Crystalline rocks are widespread; they crop out in many mountain ranges and underlie the entire region at depth. Crystalline rocks include metamorphic rocks and intrusive igneous rocks of Precambrian, Mesozoic, and Tertiary age.

Ground-Water Flow Regime

Ground-water recharge occurs by infiltration of precipitation and runoff. Recharge in basins in California and Nevada has been estimated as a function of the quantity of precipitation (Rantz and Eakin, 1971; Rush, 1970). Rantz and Eakin (1971) estimate recharge in areas receiving less than 200 mm of precipitation annually to be less than 3 percent of precipitation; they estimated recharge to be 3 percent for areas receiving 200 to 300 mm and 7 percent for areas receiving 300 to 380 mm. However, recharge also is a function of such factors as water loss by evaporation and transpiration, rock type and physical character, slope, and soil cover. Recharge by direct infiltration of precipitation to the valley floors that receive 200 mm or less precipitation per year, is believed to be very small (Winograd and Thordarson, 1975, p. C92), but recharge may occur during infrequent large storms that cause runoff locally or at higher altitudes in mountains that adjoin the valley floors. Winograd and Thordarson (1975, p. C86) suggested relatively substantial recharge in outcrop areas of extensively fractured carbonate rock in the mountain ranges, and relatively little recharge in outcrop areas of tuff on which a clayey soil has developed.

Natural discharge is by flow to springs, by evapotranspiration in areas where the water level is near the land surface, and by seepage to the Colorado River. The Death Valley region is largely composed of closed topographic basins that are apparently coincident with closed ground-water flow systems of ground-water units DV-02, and DV-04 through DV-09. Ground water in these closed basins flows to playa areas where it is discharged. A part of the region, ground-water unit DV-01, has surface drainage and ground-water discharge to the Colorado River.

Ground-water flow in ground-water unit DV-03 is not coincident with topographic basins. This unit is underlain by the extensive Paleozoic carbonate-rock aquifers and associated confining beds. Because of the effect of the carbonate aquifers in underdraining the area and the effect of structural and lithologic controls in compartmentalization of flow, ground-water flow in ground-water unit DV-03 is complex. Because ground-water flow commonly is not coincident with topographic basins and because the flow systems in the unit are imperfectly known, the unit is large and not subdivided. The ground-water-flow conditions in ground-water unit DV-03 are discussed in the following paragraphs.

Subsurface flow between many topographic basins occurs in ground-water unit DV-03. Basins identified by Winograd and Thordarson (1975) that drain to the carbonate-rock aquifers include those of Yucca and Frenchman Flats. By inference, several other closed basins without surface discharge of ground water also are believed to drain to the carbonate aquifer. These are Indian Springs Valley, northern Three Lakes Valley, Emigrant Valley, and Tikaboo Valley, all in ground-water unit DV-03. The area where ground-water infiltrates from the closed basins to the carbonate aquifer is not known. The basins that drain to the underlying carbonate aquifer are identified in plate 4.

Regional interbasin movement of ground water in ground-water unit DV-03 is affected by the deformed nature of the great thicknesses of Paleozoic carbonate and clastic rocks. Major wrench, thrust, and normal faults and folds have been shown to exert marked control on ground-water movement (Winograd and Thordarson, 1968). Compartmentalization of flow in the region by fault blocks containing thick sequences of clastic rocks, and perhaps also shear zones, were demonstrated by Winograd and Thordarson. Large-scale heterogeneities in carbonate-rock permeability have been inferred by Winograd and Pearson (1976) on the basis of Carbon-14 isotope analyses of ground water in the Ash Meadows area.

Discharge of ground water from ground-water unit DV-03 occurs in several large areas. Three large natural-discharge areas occur in Sarcobatus Flat, Amargosa Desert, and Pahrump Valley (pl. 4). The ultimate discharge area for ground-water unit DV-03 is Death Valley, the basin of lowest altitude in the region, 86 m below sea level. Discharge at Death Valley occurs to numerous springs and seeps and by evapotranspiration (Hunt and others, 1966; Miller, 1977). Oblique aerial views of Death Valley are shown in figures 8 and 9.




Figure 8.--

View from the saltpan of Badwater Basin in Death Valley to the northwest across faulted alluvial fans toward the Panamint Range. Growths of phreatophytes occur in a band from the lower left to the middle right of the photograph at the boundary of the saltpan and the alluvial fans. Shorty's well is in the area of vegetation at the center; Tule Spring is in an area of vegetation near the middle right of the photograph. The depth to ground water is near the surface at the edge of the saltpan and increases rapidly with distance up the alluvial fans. The growth of phreatophytes marks the area of accessible depths to ground water and salinity of ground water

within the tolerance of the plants. The phreatophytes are zoned with the more salt-tolerant species near the saltpan. Escarpment of Hanaupah fault shows two stages of displacement. Obvious fault escarpment across center of photograph marks fault displacing older upper Pleistocene fan gravels (No. 2 gravel of Hunt and Mabey, 1966) by as much as 15 meters. Small fault (arrow) displaces younger upper Pleistocene fan gravels (No. 3 gravel of Hunt and Mabey, 1966) by as much as 2 meters. Photograph by John S. Shelton (1979).



Figure 9.--View of Death Valley from near the south end of the valley. Smith Mountain is in right foreground, Mormon Point at center of photograph and Grapevine Mountains in middle background. Death Valley is the ultimate discharge area for a large part of the Death Valley region. Amargosa River flows from lower left, away from observer, toward Badwater Basin (altitude 86 meters below sea level) white (rock-salt area) in center of valley in upper middle part of photograph. Photograph by John S. Shelton (1979).

Thermal springs (arbitrarily designated as those with a water temperature more than 20 °C) are found in the region, many of which discharge from zones with substantial permeability in carbonate rock. Both thermal and nonthermal springs characterize the major discharge areas near the Colorado River and in Death Valley, Ash Meadows, Amargosa Desert, and in many smaller basins. Cold springs having 200 L/min or more discharge occur at Ash Meadows and at other localities. The temperature of most of the thermal springs is less than 50 °C, which indicates convective heat flow of ground water rather than locally anomalously high heat flows. Thermal springs are plotted in plate 4.

Ground-water withdrawal is concentrated at pumping centers such as Las Vegas, Pahrump Valley, and Ash Meadows. Many valleys have small or no pumping. Withdrawal is described in reports by Bedinger, Langer, and Moyle (1985), and Bedinger, Harrill, and Thomas (1985).

Ground-Water Flow Analysis

AREAL GROUND-WATER FLOW

The region was separated into hydrogeologic units at the water table based on information from previous studies and summarized in this report, and from the geologic sections constructed for the region (see Chapter A, Bedinger, Sargent and others, 1985) and water-level contour maps. Relative ground-water traveltimes at the water table were analyzed using the procedure described in Chapter A, (Bedinger, Sargent and others, 1985). Velocities in the hydrogeologic units (pl. 3) are reported as relative velocities because site-specific data are not available. The values of hydraulic properties of the hydrogeologic units and hydraulic gradients used in estimating relative ground-water velocities are listed in the table 1.

The hydraulic gradients for the hydrogeologic units are representative gradients taken from the water-level contour map. The ratio of hydraulic conductivity to effective porosity was estimated using the values in Chapter A (Bedinger, Sargent, and others, 1985) and modified from the lithologic and hydrologic description of the units, and further modified during the verification of the cross-sectional and areal-flow models.

Table 1.--Hydraulic properties of hydrogeologic units and hydraulic gradients used in estimating relative ground-water velocities at the water table.

[K = hydraulic conductivity, in meters per day;
 \emptyset = effective porosity]

Hydrogeologic unit	Map symbol (pl. 3)	K/ \emptyset (meters per day)	Hydraulic gradient
Basin fill	a	6×10^1	0.003
Volcanic rocks:			
Undifferentiated	y	1×10^{-1}	.03
Lava flows	b	3×10^0	.03
Ash-flow tuff	t	1×10^{-1}	.03
Ash-flow tuff	t	4×10^{-1}	.007
Crystalline rocks:			
Granitic rocks	g	2×10^{-1}	.03
Mafic intrusive rocks	z	2×10^{-1}	.03
Metamorphic rocks	m	2×10^{-1}	.03
Sedimentary rocks:			
Coarse-grained clastic rocks	s	2×10^{-1}	.03
Fine-grained clastic rocks	f	2×10^{-6}	---
Carbonate rocks	c	1×10^1	.003
Mixed rocks:			
Large percentage of carbonate rocks	I	1×10^1	.003
Large percentage of crystalline rocks	II	2×10^{-1}	.03

Relative ground-water traveltimes are shown in plate 4. The ground-water divides and flow paths are estimated on the available ground-water level data which in many parts of the region is very sparse. Lacking water-level data, the ground-water divides and flow paths are estimated from topographic divides and the lithologic units at the water table. Evidence for barriers to ground-water flow has been demonstrated by Winograd and Thordarson (1968 and 1975), and Waddell (1982). Other flow barriers, as yet unidentified, undoubtedly exist in the region, especially in ground-water unit DV-03. Flow arrows indicate paths at the water table along which the relative traveltimes to these discharge areas were calculated using methods described in Chapter A (Bedinger, Sargent, and others, 1985). Major discharge areas, large springs, and evapotranspiration areas also are shown in plate 4. Several closed basins have no surface discharge areas. Flow from the water table in these basins is inferred to be downward to carbonate rocks. These basins are indicated diagrammatically in plate 4 with diamond-shaped symbols. The diamond-shaped areas are not intended to indicate the location or distribution of areas of downward flow to the regional aquifer, but to simply show that the basin does not have surface discharge and that it is inferred that the basin discharges by underflow.

CROSS-SECTIONAL MODELS

Cross-sectional models were used to analyze ground-water flow along selected flow paths. The mathematical model used in modeling flow in cross section is given in Chapter A of this report (Reed, 1985). The map location of hydrogeologic sections and the modeled sections are shown in plate 5. The value of hydraulic properties of the rock units in the hydrogeologic sections used in analysis of the ground-water flow are given in table 2.

Distribution of rock units, relative traveltime, and stream functions are given on the hydrogeologic sections. Relative traveltimes are given in intervals of 1 order of magnitude from 10^1 and longer. Numbers indicate the relative time of travel from points on the line to the discharge area. Stream functions show the directions of ground-water movement and the numbers indicate relative quantity of flow in the section below the flow line.

Large percentage
of carbonate
rocks

1

1×10^1

Large percentage
of crystalline
rocks

11

2×10^1

Table 2.--Hydraulic properties of units modeled in hydrogeologic sections

[K=hydraulic conductivity, in meters per day; θ = effective porosity]


Hydrologic unit	Symbol (pl. 4)	Hydrogeologic section					
		A-A'		B-B'		C-C'	
		K/ θ	θ	K/ θ	θ	K/ θ	θ
Coarse-grained basin fill	a	1×10^1	1.8×10^{-1}	1×10^1	1.8×10^{-1}	2×10^1	1.8×10^{-1}
Ash-flow tuff	t	---	---	4×10^{-4}	3.5×10^{-1}	---	---
Carbonate rocks	c	---	---	---	---	3×10^{-3}	1×10^{-1}
Crystalline rocks, upper part of section	G	5×10^{-4}	3×10^{-3}	5×10^{-4}	3×10^{-3}	5×10^{-4}	3×10^{-3}
Crystalline rocks, lower part of section	g	3×10^{-7}	1×10^{-4}	3×10^{-7}	1×10^{-4}	3×10^{-7}	1×10^{-4}
Undifferentiated volcanic rocks	v	4×10^{-4}	4×10^{-3}	4×10^{-3}	4×10^{-4}	---	---
Lava flows	b	5×10^{-4}	1.5×10^{-3}	5×10^{-1}	1.5×10^{-1}	---	---
Coarse-grained clastic rocks	s	3×10^{-2}	1.8×10^{-1}	3×10^{-2}	1.8×10^{-1}	---	---
Coarse-grained basin fill	A	2×10^{-2}	1.8×10^{-1}	1.2×10^0	1.8×10^{-1}	1×10^{-1}	1.8×10^{-1}
Fined-grained clastic rocks	f	---	---	---	---	---	---

CROSS-SECTIONAL MODELS
The hydrogeologic sections provide a more realistic concept of the flow paths and traveltime between widely spaced points in the region than does the map of traveltime at the water table. Cross-sectional models were used to analyze ground-water flow along selected flow paths. A mathematical model used in modeling flow in cross-sections is Chapter A of this report (Freeb, 1983). The hydrogeologic sections and the traveltime contours are shown in Figure 3. The value of the relative traveltime is 10⁵ to 10⁸. Commonly, these longest relative traveltimes are of restricted surface area at the water table. The areas of longer relative traveltime enlarge with depth and would provide more confidence in locating an area of long traveltime at depth beneath the water table than above the water table. Broad areas of relative traveltime of 10⁴ or greater exist at the water table in the hydrogeologic sections.

Evidence exists for large-scale variations in permeability in the carbonate rocks (Winograd and Pearson, 1976) and barriers to the regional ground-water flow (Winograd and Thordarson, 1968; Waddell, 1982). Zones with substantial permeability that may exist locally in the carbonate rocks may have a great effect in affecting flow distribution and times of travel in the carbonate rocks. Because the distribution and extent of the channeling in the carbonate rocks are not known, the permeability and effective porosity of the carbonate rocks are modeled as constant values, believed to represent averages of these hydrologic properties.

Quality of Ground Water

The quality of ground water in the Death Valley region is characterized by the areal distribution of dissolved solids (pl. 6) and predominant chemical constituents in solution (fig. 10). These maps are generalized from those by Thompson and others (1985) and Thompson and Chappel (1985) compiled from the water-quality files of the U.S. Geological Survey (WATSTORE) and published reports. The data are mostly from non-geothermal springs and wells less than 150 m deep completed in alluvial and basin-fill deposits. In areas where data are not available, the water-quality characteristics were estimated from the position in the ground-water flow system and the lithology of the local bedrock.






Figure 10.--Distribution of chemical types of ground water in the Death Valley region and vicinity.

Dissolved-solids concentration is generally less than 500 mg/L except beneath the surfaces of some playa lakes, where ground water may contain greater than 500 mg/L. Concentrations between 1,000 and 3,000 mg/L also occur in the areas near Lake Mead and the Colorado River. Dissolved-solids concentrations commonly are 1,000 to 3,000 mg/L in playa areas and ground water of more than 3,000 mg/L of dissolved solids is found in a few playas such as Death Valley, Columbus Salt Marsh, and Clayton Valley.

The concentration of dissolved solids of most ground water in consolidated rocks of the region probably is less than 500 mg/L. The water-quality data from four deep wells in southern Nevada as summarized by Winograd and Thordarson (1975) indicated no significant increase in dissolved-solids concentration of water in the lower carbonate aquifer to depths of a few thousand meters.

Sodium bicarbonate and calcium magnesium bicarbonate type waters occur throughout about 90 percent of the region. Mixed cation sulfate and mixed cation chloride type waters occur in and near natural discharge areas and generally correspond to areas of maximum dissolved-solids concentration.

Pleistocene Hydrologic Conditions

The climate of glacial epochs during the Pleistocene in the Basin and Range province has been estimated from evidence from plant debris and relict lake-shore lines. Spaulding (1984) has made a study of climates at times during the past 45,000 years from plant remains in pack-rat middens at the Nevada Test Site and vicinity. Climates have been estimated from hydrologic budgets for the full glacial climate of Lake Spring in Spring Valley, Nevada (Snyder and Langbein, 1962), Lake Lahontan, Nevada (Antevs, 1952; Benson, 1978) and of many late(?) Pleistocene lakes in Nevada by Mifflin and Wheat (1979). These investigations conclude that the climate during glacial epochs was cooler with greater precipitation than present. Other investigators have concluded that the glacial climate in the Basin and Range province were much cooler than present with no increase, or even a decrease, in precipitation (Galloway, 1970; Brakenridge, 1978).




Figure 10.--Distribution of chemical types of ground water in the Death Valley region.

Compilation of data on Pleistocene lakes and marshes in the Basin and Range province (Williams and Bedinger, 1985) shows that Pleistocene lakes occupied several closed basins in the Death Valley region. Lake Manly, the largest, occupied the floor of Death Valley, and Panamint Lake occupied Panamint Valley in ground-water unit DV-04. Evidence of smaller Pleistocene lakes has been identified in ground-water units DV-06 (northwest one-half), DV-07, DV-08, and DV-09. Three Pleistocene lakes have been found in the northern part of ground-water unit DV-03. With the exception of Groom Lake, no Pleistocene lakes are known to have occupied closed basins that drain from the water table to the underlying carbonate-rock aquifer. Marshes are believed to have occupied parts of valley floors during the Pleistocene in the Amargosa Desert (ground-water unit DV-03), Sarcobatus Flat (ground-water unit DV-03), and the basins in ground-water units DV-05, DV-07, DV-08, and DV-09. Winograd and Doty (1980) have shown that major changes in altitude and location of ground-water discharge actually occurred during the late(?) Pleistocene. These changes are attributed to both climate and tectonism.

References Cited

- Antevs, Ernst, 1952, Cenozoic climates of the Great Basin: *Geologische Rundschau*, v. 40, no. 1, p. 94-108.
- Bedinger, M. S., Langer, W. H., and Moyle, W. R., 1984, Map showing ground-water units and withdrawal, Basin and Range province, southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-A, scale 1:500,000.
- Bedinger, M. S., Harrill, J. R., and Thomas, J. M., 1984, Map showing ground-water units and withdrawal, Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations Report 83-4119-A, scale 1:500,000.
- Bedinger, M. S., Sargent, K. A., and others, 1985, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Basis of characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].
- Benson, L. V., 1978, Fluctuation in the level of pluvial Lake Lahontan during the last 40,000 years: *Quaternary Research*, v. 9, no. 3, p. 300-318.
- Brakenridge, G. R., 1978, Evidence for a cold, dry full-glacial climate in the American southwest: *Quaternary Research*, v. 9, no. 1, p. 22-40.
- Galloway, R. W., 1970, Full-glacial climate in the southwestern United States: *Annals of the Association of American Geographers*, v. 60, no. 2, p. 245-256.
- Hunt, C. B., and Mabey, D. R., 1966, Stratigraphy and structure Death Valley, California: U.S. Geological Survey Professional Paper 494-A, map scale 1:96,000, 162 p.

- Hunt, C. B., Robinson, T. W., Bowles, W. A., and Washburn, A. L., 1966, Hydrologic basin, Death Valley, California: U.S. Geological Survey Professional Paper 494-B, p. B1-B138.
- Mifflin, M. D., and Wheat, M. M., 1979, Pluvial lakes and estimated pluvial climates of Nevada: Nevada Bureau of Mines and Geology Bulletin 94, 57 p.
- Miller, G. A., 1977, Appraisal of water resources of Death Valley, California: U.S. Geological Survey Open-File Report 77-728, 68 p.
- Rantz, S. E., and Eakin, T. E., 1971, A summary of methods for the collection and analysis of basic hydrologic data for arid regions: U.S. Geological Survey open-file report, 125 p.
- Reed, J. E., 1985, Mathematical model for simulating ground-water and heat flow, in Bedinger, M. S., Sargent, K. A., and others, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste--Basis of characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, [in press].
- Rush, F. E., 1970, Regional ground-water systems in the Nevada Test Site area, Nye, Lincoln, and Clark Counties, Nevada: Nevada Department of Conservation and Natural Resources, Division of Water-Resources, Water Resources-Reconnaissance Series Report 54, 25 p.
- Snyder, C. T., and Langbein, W. B., 1962, Pleistocene lake in Spring Valley, Nevada, and its climatic implications: Journal of Geophysical Research, v. 67, no. 6, p. 2385-2394.

- Spaulding, W. G., 1984, Vegetation and climates of the last 45,000 years in the vicinity of the Nevada Test Site, south-central Nevada: U.S. Geological Survey Open-File Report 83-535, 199 p.
- Thompson, T. H., and Chappell, Richard, 1984, Map showing distribution of dissolved solids and dominant chemical type in ground water, Basin and Range province, Nevada: U.S. Geological Survey Water-Resources Investigations Report 83-4119-B, scale 1:500,000.
- Thompson, T. H., Nuter, J. A., Moyle, W. R., and Woolfenden, L. R., 1984, Map showing distribution of dissolved solids and dominant chemical type in ground water, Basin and Range province, southern California: U.S. Geological Survey Water-Resources Investigations Report 83-4116-C, scale 1:500,000.
- Waddell, R. K., 1982, Two-dimensional, steady-state model of ground-water flow, Nevada Test Site and vicinity, Nevada-California: U.S. Geological Survey Water-Resources Investigations Report 82-4085, 72 p.
- Williams, T. R., and Bedinger, M. S., 1984, Selected geologic and hydrologic characteristics of Basin and Range province--Pleistocene lakes and marshes: U.S. Geological Survey Miscellaneous Investigations Series Map I-1522-D, scale 1:2,500,000.
- Winograd, I. J., 1971, Hydrogeology of ash flow tuff--A preliminary statement: Water Resources Research v. 7, no. 4, p. 994-1006.

- Winograd, I. J., and Doty, G. C., 1980, Paleohydrology of the southern Great Basin, with special reference to water table fluctuations beneath the Nevada Test Site during the late Pleistocene: U.S. Geological Survey Open-File Report 80-569, 91 p.
- Winograd, I. J., and Pearson, F. J., Jr., 1976, Major carbon-14 anomaly in a regional carbonate aquifer--Possible evidence for megascale channeling, south-central Great Basin: Water Resources Research, v. 12, no. 6, p. 1125-1143.
- Winograd, I. J., and Thordarson, William, 1968, Structural control of ground-water movement in miogeosynclinal rocks of south-central Nevada, in Eckel, E. B., ed., Studies of geology and hydrology, Nevada Test Site: Geological Society of America Memoir 110, p. 35-48.
- 1975, Hydrogeologic and hydrochemical framework, south-central Great Basin, Nevada-California, with special reference to the Nevada Test Site: U.S. Geological Survey Professional Paper 712-C, 126 p.

MINERAL AND ENERGY RESOURCES

By

B. T. Brady, U.S. Geological Survey

The Death Valley region contains metallic-mineral deposits that are of several ages and occur in diverse geologic environments. These mineralized areas commonly contain precious metal deposits and base metals in replacement deposits. Contact tungsten deposits are of significance locally. One of the most important world sources of rare-earth elements is being mined currently near Mountain Pass, San Bernardino County, California. Large deposits of magnesite and brucite are mined near Gabbs in northwestern Nye County, Nevada. Molybdenum is being mined currently at Hall north of Tonopah. Fluorspar and barite have been produced intermittently from a few principal localities in the study area. Talc mining is important in the Death Valley region. An important domestic and world lithium resource is the brine pumped from sediments beneath Silver Peak Marsh, central Esmeralda County, Nevada. Borates, salt, and gypsum have been produced locally from evaporite deposits. Two Known Geothermal Resource Areas (KGRA) and many geothermal occurrences are present in the study area. Small tonnages of coal have been mined in the Coaldale field, Esmeralda County. No additional occurrences of coal or any productive oil, gas, carbon dioxide, or helium wells are identified at present in the study area.

Metallic-Mineral Resources

The metallic-mineral districts mentioned in this report are those areas depicted by Wong (1983a, b), and their locations are shown on plate 7. The mineral district boundaries enclose areas of productive workings, however they do not indicate the limit of mineralized rock. A summary of the principal commodities, modes of occurrence, and general references for the mineral districts in the Death Valley region is presented in tables 3 and 4 (modified from Wong, 1983a, b).

At least 113 metal-mining districts are located in the Death Valley region, and more than 25 commodities were produced from mines in the region. Several types of mineralization are important locally in rocks ranging from Precambrian to Quaternary in age.

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada

Commodities listed in the commodities column are abbreviated as follows:

Ag, silver; As, arsenic; Au, gold; B, boron; Ba, barium; Bi, bismuth;

Cd, cadmium; Co, Cobalt; Cu, copper; F, fluorine; Fe, iron; Hg, mercury;

K, potassium; Mg, magnesium; Mn, manganese; Mo, molybdenum; Pb, lead;

Pt, platinum; Re, rhenium; Sb, antimony; Se, selenium; Sn, tin;

Sr, strontium; Te, tellurium; Th, thorium; Ti, titanium;

U, uranium; V, vanadium; W, Tungsten, and Zn, zinc.

(These data are from Wong (1983b), and they are preliminary and subject to revision).

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Clark County						
Alunite (Railroad Pass, Vincent)	23 S. 23 S. 23 S.	62 E. 63 E. 64 E.	Au, Ag, Pb, Cu, Mn, Alunite	Disseminated Vein	Quartz monzonite Andesite	Longwell and others, 1965
Charleston	18 S. 18 S. 19 S. 20 S.	56 E. 57 E. 56 E. 59 E.	Ag, Pb, Zn	Vein Disseminated	Dolomitized limestone Dolomitized limestone	Longwell and others, 1965
Crescent (Crescent Peak)	27 S. 28 S. 29 S.	61 E. 61 E. 61 E.	Au, Ag, Pb, Cu, Mo, U, Th	Vein Disseminated	Granite, Gneiss Prospect Mountain Quartzite; Quartz Monzonite	Hewett, 1956; Longwell and others, 1965; Ransome, 1907; Schilling, 1962; Vanderburg, 1937a
Eldorado (Eldorado Canyon, Colorado, Nelson)	26 S. 26 S. 27 S. 27 S.	64 E. 65 E. 64 E. 65 E.	Au, Ag, Pb, Zn, Cu	Vein Disseminated Breccia zone	Gneiss, schist, quartz monzonite, andesite Quartz monzonite Quartz monzonite	Longwell and others, 1965; Ransome, 1907; Vanderburg, 1937a
Gass Peak	18 S.	61 E.	Au, Ag, Pb, Zn	Shear zone	Dolomitized limestone	Hewett and others, 1936; Longwell and others, 1965

Table 3.--Metallic-metal districts in the Death Valley region of Nevada--Continued

121

Mining district	Location		Commodities	Deposit type	Host rock	References
	Township	Range				
Clark County--Continued						
Goodsprings (Yellow Pine, Potosi)	23 S.	58 E.	Au, Ag, Pb,	Vein	Dolomitized limestone,	Albritton and others, 1954; Bailey and Phoenix, 1944; Beal, 1963; Hewett, 1931; Hewett, 1956; Knopf, 1915; Lincoln, 1923; Longwell and others, 1965; Schilling, 1962
	24 S.	56 E.	Zn, Cu, Mo,	Replacement	granite porphyry	
	24 S.	57 E.	V, U, Pt,	Disseminated	Limestone, dolomite	
	24 S.	58 E.	Co, Ti, Hg,	Breccia zone	Dolomite, limestone	
	25 S.	57 E.	Pb	Replacement	Dolomite breccia, limestone	
Divide (Gold Mountains)	26 S.	58 E.	Ag, Au, Mo	Contact metamorphic	Limestone	Longwell and others, 1965; Trengove, 1959; Vanderburg, 1937a
Las Vegas	21 S.	63 E.	Mn, Au, Ag,	Bedded	Sandstone and clay-	
	21 S.	64 E.	Pb, Cu		stones of the	
	22 S.	63 E.		Disseminated	Muddy Creek Formation	
	22 S.	64 E.				
Searchlight	28 S.	63 E.	Au, Ag, Pb,	Vein	Andesite porphyry,	Callaghan, 1939; Longwell and others, 1965; Schilling, 1962
	28 S.	64 E.	Zn, Cu, Mo		gneiss, hornfels	
	29 S.	63 E.				
Sunset (Lyons)	27 S.	60 E.	Au, Ag, Pb, Cu	Breccia pipe	Granite gneiss	Hewett, 1956; Longwell and others, 1965

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Township	Range				
Esmeralda County						
Alum	1 N.	39 E.	Alum, sulfur, Hg, gypsum	Vein	Rhyolite	Albers and Stewart, 1972; Lincoln, 1923; Spurr, 1906
Black Horse	2 N. 2 N.	34 E. 35 E.	W, Mo, diatomite	Disseminated Contact metamorphic Lacustrine	Tactite Tactite Lake beds	Albers and Stewart, 1972
Buena Vista (Oneota, Basalt, Mount Montgomery)	1 N. 1 S.	33 E. 33 E.	Au, Ag, U, W, Mo	Vein Contact	Phyllite, hornblende, diorite, andesitic tuff Tactite, adamellite	Albers and Stewart, 1972 Lincoln, 1923; Ross, 1961
Coaldale	2 N.	37 E.	Coal, U	Bedded Veinlets, breccia pipe	Lake beds Rhyolitic tuff	Duncan, 1953; Hance, 1913; Lincoln, 1923; Toenges and others, 1946
Crow Springs	6 N. 4 N. 5 N.	38 E. 39 E. 40 E.	Au, Ag, Sb, Se, Cu perlite	Vein	Quartzite, chert	Albers, and Stewart, 1972; Lawrence, 1963

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Esmeralda County--Continued						
Cuprite (Tule Canyon)	4 S. 4 S. 5 S. 5 S.	42 E. 43 E. 42 E. 43 E.	Au, Ag, Cu, S Cu	Vein Replacement	Limestone, rhyolite Rhyolitic tuff, limestone, boulders	Ball, 1906; Ball, 1907; Ransome, 1909
Divide (Gold Mountain) (West Divide, Neepah)	2 N. 1 N. 2 N.	42 E. 41 E. 39 E.	Ag, Au, Mo Au, Cu, Zn	Vein Contact Vein Replacement	Rhyolite breccia Limestone, granite Limestone Limestone	Knopf, 1921b; Lincoln, 1923; Schilling, 1962; Young, 1920
Dyer	2 S.	36 E.	Ag, Pb, Cu	Vein	Slaty limestone	Lincoln, 1923; Spurr, 1906
Fish Lake Marsh	1 N. 1 N. 1 S.	36 E. 37 E. 37 E.	Clay, B, U	Veinlet Bedded Disseminated	Lake beds shale Lake beds shale Tuffaceous rocks	Albers and Stewart, 1972; Lincoln, 1923; Smith, 1964
Fish Lake Valley (White Mountain)	1 N. 1 N. 1 S. 1 S.	33 E. 34 E. 33 E. 34 E.	Hg, Sb Cu	Fracture filling	Opalite, air-fall tuff rhyolite, andesite, phyllite	Holmes, 1965; Lawrence, 1963
Halfway Springs	1 S.	34 E.	Au, Ag, Cu	Veinlets Disseminated	Rhyolite Rhyolite	Albers and

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Location						
Mining district	Town-ship	Range	Commodities	Deposit type	Host rock	References
Esmeralda County--Continued						
Gilbert (Desert)	3 N. 4 N.	38 E. 38 E.	Ag, Pb, An, Cu, Mo, W Sb, Au	Vein	Limestone, shale, chert, quartzite, rhyolite	Ferguson, 1927
Goldfield	2 S. 2 S. 3 S.	42 E. 43 E. 42 E.	Au, Ag, Cu, Pb, Bi, K, Sb, Sn, Te	Vein	Rhyolite, dacite, andesite, tuff	Ransome, 1909 Lincoln, 1923; Spurr, 1906
Goldfield Hills area	3 S. 3 S.	42 E. 43 E.	Ba, Mn	Lens Vein Nodules	Cherty limestone Rhyolitic welded tuff Limestone	Albers and Stewart, 1972
Good Hope (White Wolf)	4 S.	37 E.	Ag, Pb, Cu	Vein	Slate, quartzite	Albers, and Stewart, 1972; Lincoln, 1923
Hornsilver (Lime Point) Gold	7 S. 7 S.	41 E. 42 E.	U, Ag, Au, Pb, Mo	Vein	Limestone, shale	Albers, and Stewart, 1972; Lincoln, 1923
Klondyke (South Klondyke)	1 N. 1 N. 1 S.	42 E. 43 E. 43 E.	Au, Ag, Pb	Vein	Limestone	Spurr, 1906

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Esmeralda County--Continued						
Lida (Alida, Tule Canyon)	5 S. 5 S. 6 S. 6 S.	40 E. 41 E. 40 E. 41 E.	Au, Ag, Pb, Cu Ag, Pb, Cu	Vein Placer Shear zone	Limestone Gravel, sand, boulders Limestone	Root, 1909; Vanderburg, 1936 Tschank and Tschank, 1970
Lone Mountain (West Divide, Weepah)	1 N. 1 N. 2 N.	40 E. 41 E. 39 E.	Ag, Pb, Ba, Au, Cu, Zn Au, Ag, Pb	Contact metasomatic Vein Replacement	Limestone, granite Limestone Limestone	Albers, and Stewart, 1972; Ball, 1907 Lincoln, 1923; Oxnam, 1936; Stretch, 1904
Montezuma	2 S. 2 S. 3 S. 3 S.	41 E. 42 E. 41 E. 42 E.	Au, Ag, Pb, Cu, Bi Ag, Pb	Vein Replacement Fracture filling	Limestone, shale Limestone, shale Andesite and rhyolite flow	Albers and Stewart, 1972; Lincoln, 1923; Stretch, 1904
Palmetto	5 S. 5 S.	39 E. 40 E.	Au, Ag, Pb, Sb, Cu	Vein	Alaskite, phyllite, sandstone, limestone	Lawrence, 1963; Lincoln, 1923; Spurr, 1906
Railroad Springs	4 S. 5 S. 5 S.	40 E. 40 E. 41 E.	Au, Ag, Cu	Vein	Limestone, shale	Albers and Stewart, 1972; Lincoln, 1923
Red Mountain (Argentite)	2 S. 2 S.	37 E. 38 E.	Mn, Ag, Au, Pb, Zn, Cu	Pipelike Vein	Tuffaceous beds Rhyolite, limestone, latite, sandstone	Benson, 1950; Keith, 1977; Lincoln, 1923

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Esmeralda County--Continued						
Rock Hill	3 N.	36 E.	Fe	Unknown	Limestone, shale, chert, quartzite, rhyolite	Reeves and others, 1958
Silver Peak	1 S.	38 E.	Au, Ag, Cu	Lens	Alaskite, limestone schist	Spurr, 1906
(Mineral Ridge)	1 S.	39 E.	Pb	Vein	Dolomite, schist, quartzite, granite	
	2 S.	38 E.		Replacement	Shaley limestone	
	2 S.	39 E.				
Sylvania (Green Mountain)	6 S.	38 E.	Mo, Re,	Disseminated	Quartz monzonite	Albers and
	6 S.	39 E.		Vein	Marble	Stewart, 1972
	7 S.	39 E.	talc	Replacement	Dolomite	
Tokop (Gold Mountain, Oriental Wash, Bonnie Claire)	7 S.	42 E.	Au, Ag, Pb,	Vein	Slaty schist, granite	Ransome, 1907
	8 S.	42 E.	Cu			
Tonopah	2 N.	42 E.	U, Clay	Disseminated	Tuff	Albers and
	3 N.	42 E.		Bedded	Tuffaceous beds	Stewart, 1972
	(The main district is in Nye County)					
Windypah	4 S.	38 E.	Au, Ag, Cu,	Lens	Alaskite	Spurr, 1906
	5 S.	38 E.	Pb, Ba, Sb	Vein	Slaty limestone, granite	
				Replacement	Chert	

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Lincoln County						
Groom	7 S.	55 E.	Au, Ag, Pb, Zn, Cu	Breccia zone Fissure/beds	Limestone, quartzite Pioche Shale	Tschanz and Pampeyan, 1970
None (Philadelphia, Silver Sands)	9 S.	57 E.	Ag, Pb, Cu	Shear zone	Limestone	Tschanz and Pampeyan, 1970
Pahranagat (Hiko)	3 S. 4 S.	59 E. 58 E.	Ag, Pb, Cu, Mn, Au	Vein	Limestone, dolomite	Tschanz and Pampeyan, 1970
Papoose	9 S.	55 E.	Au, Ag, Pb	Fissure vein	Quartzite	Tschanz, and Pampeyan, 1970
Tem Piute	3 S. 4 S.	56 E. 56 E.	Ag, W, Mo Zn, F, Bi, Hg, Pb	Contact metamorphic Vein/replacement Fracture filling	Tactite, limestone, shale Dolomite Andesite and rhyolite flow	Couch and Carpenter, 1943; Hill, 1916 Holmes, 1965; Lemmon and Tweto, 1962; Schilling, 1962; Schilling, 1963; Tschanz and Pampeyan, 1970
Cactus Springs	2 S.	45 E.	Au, Ag, Cu	Vein		
Clifford	3 S.	49 E.	Au, Ag	Vein		

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town- ship	Range				
Mineral County						
Bell (Simon, OMCO, Cedar Mountain)	8 N.	37 E.	Au, Ag, Pb,	Vein	Volcanic rocks, lime- stone, andesite	Bailey and Phoenix, 1944; Couch and Carpenter, 1943; Knopf, 1921a; Ross, 1961; Vanderburg, 1937b
	8 N.	38 E.	Zn, W, Hg,			
	9 N.	37 E.	Cu	Replacement Contact meta- morphitic	Limestone Tactite, limestone, granitics	
Nye County						
Antelope Springs	4 S.	47 E.	Au, Ag, Pb	Vein Disseminated	Rhyolite Tuff	Cornwall, 1972; Kral, 1951
Ash Meadows	17 S.	49 E.	Clay	Sedimentary	Playa deposit	Cornwall, 1972; Kral, 1951
	18 S.	50 E.				
Barcelona (Spanish Belt, Spanish)	9 N.	45 E.	Au, Ag, Pb,	Vein	Shale, schist, granite, limestone	Garside, 1973; Kleinhampl and Ziony, 1983
	10 N.	45 E.	Zn, Sb, Hg, Se, Mo, Cu, F, U, W, Ti, Fe, V	Peneconcordant Contact metamorphic	Ash-flow tuff Granite, skarn	
Bellehelen	2 N.	49 E.	Au, Ag, V,	Vein	Tuff	Kleinhampl and Ziony, 1983; Kral, 1951
	2 N.	50 E.	Fe	Fissure filling	Welded tuff	
	3 N.	49 E.				

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Nye County--Continued						
Belmont (Philadelphia, Silver Bend)	8 N. 9 N.	45 E. 45 E.	Ag, Sb, Pb, Cu, W	Vein	Carbonate rocks, granite porphyry	Kleinhampl and Ziony, 1983
Bruner (Phonolite)	14 N.	37 E.	Au, Ag	Vein	Rhyolite, andesite	Kleinhampl and Ziony, 1983; Kral, 1951
Bullfrog (Pioneer, Rhyolite)	11 S. 11 S. 12 S. 12 S.	45 E. 46 E. 45 E. 46 E. 47 E.	Au, Ag, Cu, U, Clay	Replacement Vein	Rhyolite Rhyolite, limestone, tuff, shale	Cornwall and Kleinhampl, 1964; Garside, 1973; Kral, 1951; Ransome and others, 1910
Cactus Springs	2 S. 3 S.	45 E. 46 E.	Au, Ag, Cu	Vein	Rhyolite	Ball, 1907; Kral, 1951
Clifford	3 N.	49 E.	Au, Ag	Vein	Rhyolitic tuff	Kleinhampl and Ziony, 1983; Kral, 1951

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Nye County--Continued						
Cloverdale (Golden, Black Springs)	7 N.	40 E.	Au, Cu, As,	Placer	Gravel	Kleinhampl and Ziony, 1983; Kral, 1951; Papke, 1979 Vanderburg, 1936
	7 N.	41 E.	Be, F, Ag,	Vein	Quartz latite, welded	
	8 N.	40 E.	Pb, Zn, Cd,	Replacement	tuff, rhyolite	
	8 N.	41 E.	Sb, Ba, Mn	Contact meta- morphitic	granite	
	9 N.	39 E.				
	9 N.	40 E.				
	10 N.	39 E.				
10 N.	40 E.				Vanderburg, 1937	
Ellendale	2 N.	47 E.	Au, Ag, Pb,	Vein	Rhyolite, andesite	Kleinhampl and Ziony, 1983; Kral, 1951
	3 N.	47 E.	Zn, Cu, Sb, Hg, Sn, Cd, Mn, Ba	Replacement Skarn	Limestone Metamorphic rocks	
Ellsworth (Marble Falls)	13 N.	37 E.	Au, Pb, Sb,	Vein	Greenstone, volcanics, limestone, granite	Kleinhampl and Ziony, 1983; Kral, 1951; Reeves and others, 1958
	13 N.	38 E.	Zn	Replacement	Dolomite, shale, quartzite	
Fairplay (Atwood, Goldyke)	10 N.	36 E.	Au, Ag, Pb,	Contact metamorphic	Tactite, granite	Kleinhampl and Ziony, 1983; Kral, 1951
	10 N.	37 E.	W, Mo, Cu,	Vein	Greenstone, andesite	
	11 N.	36 E.	Hg	Disseminated	Limestone	

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Nye County--Continued						
Fluorine (Bare Mountain)	12 S.	47 E.	Au, F, U, Ag,	Vein	Schist, quartzite,	Cornwall and
	12 S.	48 E.	W, Pb, Hg,		sandstone, siltstone	Kleinhampl, 1964;
	13 S.	47 E.	marble,	Replacement	Limestone, dolomite	Garside, 1973;
	13 S.	48 E.	diatomaceous	Breccia pipe	Dolomite	Kral, 1951;
	14 S.	47 E.	earth,			Papke, 1979
			stone,			
			perlite,			
			pumicite,			
			silica			

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Nye County--Continued						
Hannapah (Volcano, Silverzone Bannock)	3 N. 4 N. 4 N.	45 E. 44 E. 45 E.	Au, Ag	Vein	Volcanics, welded tuff, rhyolite	Garside, 1973; Kleinhampl and Ziony, 1983; Kral, 1951
Jackson (Gold Park)	13 N. 14 N. 14 N.	39 E. 39 E. 40 E.	Au, Ag, Pb, Cu, Hg, U, F	Fissure filling Vein	Ash-flow tuff Meta-andesite, rhyolite	Bonham, 1970; Garside, 1973; Kleinhampl and Ziony, 1983; Kral, 1951
Jett (Argentore, Silver Point, Ledbetter Canyon, Peavine Canyon, Wall Canyon)	9 N. 10 N. 11 N.	42 E. 42 E. 42 E.	Au, Ag, Pb, Zn, Sb, Cu, Hg, W, As	Vein Placer	Shale, limestone, schist Gravel	Kleinhampl and Ziony, 1983; Kral, 1951; Lawrence, 1963
Johnnie	17 S. 17 S. 17 S. 18 S.	52 E. 53 E. 54 E. 52 E.	Au, Pb, Cu, U	Vein Placer	Quartzite, limestone Gravel	Garside, 1973; Kral, 1951

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Nye County--Continued						
Lee (Big Dune)	15 S.	47 E.	Au	Vein	Dolomite	Cornwall, 1972
Longstreet (Fresno, Georges Canyon)	5 N.	46 E.	Au, Ag, Pb,	Gossan		Kleinhampl and
	5 N.	47 E.	Zn, Hg	Vein		Ziony, 1983;
	6 N.	45 E.		Placer	Rhyolitic tuff	Kral, 1951
	6 N.	46 E.			Gravel	
	6 N.	47 E.				
Manhattan	8 N.	43 E.	Au, Ag, Pb,	Replacement	Limestone	Ferguson, 1924
	8 N.	44 E.	Cu, Mo, Sb, F, As, Ba	Vein	Limestone, andesite porphyry schist, sandstone, quartzite	Kral, 1951; Papke, 1979
Mellan Mountain	3 S.	48 E.	Au, Ag	Vein	Rhyolitic ash-flow tuff	Cornwall, 1972; Kral, 1951
Mine Mountain	11 S.	52 E.	Ag, Pb, Hg	Vein	Quartzite, dolomite	Cornwall, 1972;
Oak Springs (Climax)	8 S.	52 E.	Au, Ag, Pb,	Vein	Limestone, shale	Kral, 1951
	8 S.	53 E.	W, Mo	Contact metamorphic	Limestone, quartz monzonite	

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host Rock	References
	Town- ship	Range				
Nye County--Continued						
Republic area	7 N. 8 N. 8 N.	39 E. 38 E. 39 E.	Ag, Pb, Au, Zn	Vein	Rhyolite, limestone	Kleinhampl and Ziony, 1983; Kral, 1951
Royston Hills area	5 N.	40 E.	Au, Ag, Pb, Cu	Vein	Chert, andesite	Kleinhampl and Ziony, 1983; Kral, 1951
San Antone (Cimarron, San Antonio)	5 N. 6 N.	42 E. 42 E.	Au, Ag, Pb, Cu, Mo	Epithermal Vein	Rhyolite, latite Shale, limestone, quartzite, quartz mica schist	Kleinhampl and Ziony, 1983; Kral, 1951
Silverbow	1 N. 1 S.	49 E. 49 E.	Au, Ag	Replacement Vein	Rhyolite Rhyolitic tuff	Kral, 1951
Stonewall	5 S.	44 E.	Au, Ag	Vein	Rhyolitic welded tuff, quartz latite	Ball, 1907; Cornwall, 1972; Kral, 1951; Lincoln, 1923

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Town-ship	Range				
Nye County--Continued						
Tolicha (Monte Cristo, Clarkdale)	7 S.	46 E.	Au, Ag	Vein	Rhyolitic ash-flow tuff	Kral, 1951
	7 S.	47 E.				
	8 S.	45 E.				
Tonopah	2 N.	42 E.	Au, Ag, U	Vein	Trachyte, tuff, rhyolite	Garside, 1973;
	2 N.	43 E.			dacite	Spurr, 1905
	3 N.	42 E.		Disseminated	Tuffaceous lake beds	
	3 N.	43 E.				
Trappmans	5 S.	47 E.	Au, Ag	Vein	Quartz monzonite, schist	Ball, 1906; Ball, 1907
Tybo, (Hot Creek, Empire, Keystone)	5 N.	49 E.	Au, Ag, Pb,	Replacement	Limestone, shale	Ferguson, 1933;
	5 N.	50 E.	Zn, Mo, Cu,	Disseminated	Tuff	Horton, 1963;
	6 N.	50 E.	Hg, Sb, Cd,	Vein	Limestone, rhyolitic tuff	Kleinhampl and Ziony, 1983;
	7 N.	50 E.	Mn, Se, Fe, As, Ba	Pods	Dolomite	Kral, 1951; Lawrence, 1963
Union (Berlin Ione, Grantsville)	11 N.	39 E.	Au, Ag, Pb,	Vein	Rhyolite, greenstone, clastics, limestone	Kral, 1951; Kleinhampl and
	12 N.	39 E.	Zn, Cu, Hg,			Ziony, 1983;
	13 N.	39 E.	Sb, Se, F, W	Replacement Placer	Limestone Gravel	Papke, 1979

Table 3.--Metallic-mineral districts in the Death Valley region of Nevada--Continued

Mining district	Location		Commodities	Deposit type	Host rock	References
	Township	Range				
Nye County--Continued						
Wahmonie	13 S. 15 S.	51 E. 50 E.	Au, Ag, Cu	Vein	Latite, dacite, tuff, breccias, quartzite	Cornwall, 1972; Kral, 1951 and Flony, 1963;
Wellington (Jamestown, O'Briens)	5 S.	46 E.	Au, Ag, Cu	Vein	Rhyolitic ash-flow tuff Chert, andesite	Ball, 1906; Kral, 1951 Hinchampl and Flony, 1963;
Wilsons	4 S.	47 E.	Au, Ag	Vein	Rhyolitic ash-flow tuff Rhyolite, latite Shale, limestone, quartzite, quartz granite igneous metamorphic volcanic tuff	Kral, 1951 Hinchampl and Flony, 1963; Kral, 1951 Flony, 1963; Kral, 1951 Kral, 1951 Kral, 1951 Kral, 1951

Table 4.--Metallic-mineral districts in the Death Valley region of California

Meridian names are abbreviated as follows: MD, Mount Diablo; and SB, San Bernardino. Commodities listed in the commodities column are abbreviated as follows: Ag, silver; As, arsenic; Au, gold; Ba, barium; Cu, copper; F, fluorine; Fe, iron; Hg, Mercury; Mn, manganese; Mo, molybdenum; Pb, lead; Sb, antimony; Sn, tin; Sr, strontium; W, tungsten; Zn, zinc. [These data are from Wong (1983a), and they are preliminary and subject to revision].

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town-ship	Range	Meridian				
Inyo County							
Argus (Kelley)	23 S. 23 S.	41 E. 42 E.	MD do.	Au, Ag, Fe, Cu	Vein	Granite, diorite, andesite	Clark, 1970; Norman and Stewart, 1951
Ballarat-South Park	21 S. 22 S. 22 S.	44 E. 44 E. 45 E.	MD do. do.	Au Au, Ag, Cu, Hg	Vein	Schist, dolomitic limestone, gneiss	Clark, 1970; Norman and Stewart, 1951
Beveridge	13 S. 14 S. 14 S.	37 E. 37 E. 38 E.	MD do. do.	Au, Ag, Cu, Pb, Zn, Fe Au, Ag, Pb, Cu	Vein	Granite, quartz monzonite, lime- stone quartzite, schist	Norman and Stewart, 1951
Cerro Gordo	16 S. 16 S. 17 S.	38 E. 39 E. 39 E.	MD do. do.	Au, Ag, Cu, Pb, Zn	Vein Replacement	Limestone, slate Limestone	Goodwin, 1957; Norman and Stewart, 1951
Chidago	7 S.	34 E.	MD	Au, Ag, Pb, Fe	Vein	Limestone	Goodwin, 1957; Norman and Stewart, 1951
Chloride Cliff (South Bullfrog)	30 N.	1 E.	SB	Au, Ag, Pb, Cu	Vein	Limestone, schist, quartzite	Clark, 1970; Goodwin, 1957; Norman and Stewart, 1951

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town-ship	Range	Meridian				
Inyo County--Continued							
Darwin (Coso)	18 S.	40 E.	MD	Ag, Pb, Zn,	Vein	Limestone, tactite,	Goodwin, 1957; Hall and MacKevett, 1962; Norman and Stewart, 1951
	19 S.	40 E.	do.	Au, Cu, As,		calc-hornfels	
	19 S.	41 E.	do.	W, Sb, F, Fe	Replacement	Limestone	
Deep Spring	6 S.	37 E.	MD	Ag, Cu	Vein	Granite	Goodwin, 1957; Waring and Huguenin, 1919
Echo Canyon- Lees Camp	27 N.	3 E.	SB	Au	Vein	Metamorphic rocks	Clark, 1970; Norman and Stewart, 1951
	28 N.	3 E.	do.				
	29 N.	3 E.	do.				
Grapevine	11 S.	43 E.	MD	Au	Vein	Metamorphosed sedimentary rocks	Clark, 1970
Greenwater	24 N.	3 E.	SB	Cu	Unknown		Eric, 1948; Waring and Huguenin, 1919
	24 N.	5 E.	do.				

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Town- ship	Location		Commodities	Deposit type	Host rock	References
		Range	Meridian				
Inyo County--Continued							
Harrisburg	17 S.	45 E.	MD	Au	Vein	Dolomitic limestone	Clark, 1970;
	18 S.	45 E.	do.			Granite, limestone	Norman and
	14 S.	41 E.	MD	Au, Ag, Pb,	Vein		Stewart, 1951;
	14 S.	40 E.	do.	Cu, N	Contact		Waring and
	15 S.	41 E.	do.		metasomatic		Huguenin, 1919
Lee	16 S.	40 E.	MD	Au, Ag, Pb,	Vein	Limestone	Clark, 1970;
	17 S.	40 E.	do.	Zn, Cu		gneiss, granitic rocks	Norman and
Modoc	19 S.	42 E.	MD	Au, Ag, Pb,	Vein	Granitic rocks, schist, limestone	Stewart, 1951
	22 N.	3 E.	SB	Au, Cu	Replacement	Limestone	Clark, 1970;
Old Coso	20 S.	40 E.	MD	Au, Ag, Cu,	Vein	Granite	Goodwin, 1957;
	21 S.	40 E.	do.	Hg			Norman and
Panamint	21 S.	45 E.	MD	Cu, Ag	Vein	Schist, limestone	Stewart, 1951
Pine Mountain	6 S.	35 E.	MD	Au, Ag, Pb, Cu	Vein	Schist	Norman and
					Replacement	Limestone, schist	Stewart, 1951

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town-ship	Range	Meridian				
Inyo County--Continued							
Resting Spring	22 N.	7 E.	SB	Pb, Zn	Replacement(?)	Dolomite	Goodwin, 1957
Saratoga	20 N.	8 E.	SB	Au, Ag, Pb, Zn, Cu	Replacement	Limestone	Goodwin, 1957
Skiddoo-Tucki Mountain	16 S.	44 E.	MD	Au	Vein	Quartz monzonite	Clark, 1970;
	16 S.	45 E.	do.				Norman and
	17 S.	44 E.	do.				Stewart, 1951
	17 S.	45 E.	do.				
Slate Range (Daily Dozen mine)	24 S.	44 E.	MD	Au, Ag	Fissure veins	Granite, quartz monzonite	Smith and others, 1968
Tibetts (Union)	12 S.	36 E.	MD	Au, Ag, Pb,	Contact metamorphic	Quartz monzonite, limestone	Goodwin, 1957;
	13 S.	35 E.	do.	Cu, W, Mo,	Vein	Granite, limestone	Lemmon and
	13 S.	36 E.	do.	Fe			Tweto, 1962

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town-ship	Range	Meridian				
Inyo County--Continued							
Ubehebe	14 S.	41 E.	MD	Au, Ag, Pb,	Vein	Granite, limestone	Clark, 1970;
	14 S.	40 E.	do.	Cu, W	Contact	monzonite	Walker and
	15 S.	41 E.	do.		metasomatic	limestone, quartz	others, 1956
Wild Rose	18 S.	45 E.	MD	Au, Ag, Cu,	Placer	Gravel	Clark, 1970;
	19 S.	45 E.	do.	Pb, Sb, Zn	Vein	Schist, gneiss,	Norman and
					Contact	granitic rocks	Stewart, 1951;
					Replacement	Limestone	White, 1940
Willow	22 N.	3 E.	SB	Au, Cu	Vein	Gneiss, schist	Clark, 1970;
							Norman and Stewart
							1951
Mountain Pass	70 N.	33 E.	90° SB	Rare Earths	Vein	Gneiss, shonkinite-	Clark and
Ipex	70 N.	37 E.	90° SB			syenite	others, 1953
Mono County							
White Mountains area	2 S.	34 E.	MD	Au, Ag, Pb	Placer	Gravel	Goodwin, 1957;
	5 S.	33 E.	do.	Cu, W	Vein	Granitic rocks,	Sampson and
						schist, limestone,	Tucker, 1940
						slate, hornfels	

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town- ship	Range	Meridian				
San Bernardino County							
Avawatz Mountain area	15 N.	6 E.	SB	Au, Ag, Cu,	Contact	Limestone	Wright and others, 1953
	15 N.	7 E.	do.	Pb, Zn, Fe,			
	16 N.	6 E.	do.	Sr			
Clark (Clark Mountain)	16 N.	13 E.	SB	Au, Cu, W, Pb,	Vein	Gneiss, quartz monzonite, limestone	Clark, 1970; Wright and others, 1953
	17 N.	13 E.	do	Ag, Zn			
Goldstone	13 N.	1 E.	SB	Au, Cu, Ag	Vein	Limestone, shales, diorite, dike, schist	Clark, 1970; Wright and others, 1953
	14 N.	1 E.	do.				
Halloran Springs	15 N.	9 E.	SB	Au, Cu, Ag,	Vein	Quartz monzonite, basalt, granite	Clark, 1970; Wright and others, 1953
	15 N.	10 E.	do.	Pb, Fe			
	15 N.	11 E.	do.				
Homer Mountain area	11 N.	19 E.	SB	Pb, Ag, Cu	Vein	Unknown	Wright and others, 1953
Ibex	10 N.	20 E.	SB	Au, Mn, Cu	Vein	Gneiss, schist	Wright and others, 1953
	10 N.	21 E.	do.				
	10 N.	22 E.	do.				

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town- ship	Range	Meridian				
San Bernardino County--Continued							
Ivanpah (Bullion, Koko Weef, Mescal)	14 N.	14 E.	SB	Au, Ag, Pb,	Vein	Limestone, quartz monzonite	Lemmon and Tweto, 1962; Wright and others, 1953
	15 N.	13 E.	do.	Cu, Zn, Fe,	Contact metasomatic	Limestone, quartz monzonite	
	15 N.	14 E.	do.	W, Sn			
Kingston Range area	19 N.	10 E.	SB	Au, Ag, Pb, Zn, Cu, Fe	Contact	Dolomite, limestone, amphibolite	Wright and others, 1953
Morrow	29 S.	45 E.	SB	Au, Cu	Vein	Granite, limestone	Wright and others, 1953
Mountain Pass	16 N.	13 E.	SB	Rare Earths	Vein	Gneiss, shonkinite- syenite	Olson and others, 1954
	16 N.	14 E.	do.	Cu, Pb, F, Mo, Zn, Ag, Sb, Ba			
Owlshead	18 N.	3 E.	SB	Mn	Fissure filling	Brecciated lime- stone, granite, fanglomerate	Davis, 1957; Mann, 1916; Trask, 1950
					Veins	Granite, marble	
Sacramento Mountain area	7 N.	20 E.	SB	Au, Ag, Cu,	Vein	Gneiss, granite	Wright and others, 1953
	7 N.	21 E.	do.	Fe, Hg			
	7 N.	22 E.	do.				
	8 N.	22 E.	do.				
	9 N.	22 E.	do.				

Table 4.--Metallic-mineral districts in the Death Valley region of California--Continued

Mining district	Location			Commodities	Deposit type	Host rock	References
	Town-ship	Range	Meridian				
San Bernardino County--Continued							
Shadow Mountains	17 N.	11 E.	SB	Au, Ag, Pb, Cu	Vein	Granitic gneiss	Clark, 1970; Wright and others, 1953
Slate Range (Sandora, Early Spring, Johnson mine)	25 S.	45 E.	MD	Au	Veins	Quartz monzonite	Smith and others, 1968
	25 S.	45 E.	do.	Ag, Pb	Veins	Mesozoic meta-volcanics, shale	
	26 S.	45 E.	MD	Au	Vein, dike	Mesozoic meta-volcanics	
Soda Mountains area	13 N.	7 E.	SB	Au, Ag, Cu	Vein	Granite	Wright and others, 1953
	14 N.	7 E.	do.				
	14 N.	8 E.	do.				
Vanderbilt (New York)	13 N.	14 E.	SB	Au, Ag, Pb,	Vein	Gneiss, pegmatite	Clark, 1970; Wright and others, 1953
	14 N.	15 E.	do.	Cu, Zn		dike, granite,	
	14 N.	16 E.	do.			dolomite	

The value and relative importance of metal production in the Death Valley region varies by district and through time. The principal metallic elements produced include: gold, silver, copper, molybdenum, lead, zinc, and tungsten. The most significant metallic mineral production to date in the study area has come from the Goldfield district, Esmeralda County, and the Tonopah district, principally in Nye County, Nevada. Smaller although still very important, production came from extensively developed mines in the following districts: Goodsprings (Clark County), Tem Piute (Lincoln County), Manhattan and Tybo (Nye County), Silver Peak (Esmeralda County), Nevada; and the Mountain Pass district (San Bernardino County), and Darwin and Cerro Gordo districts (Inyo County), California. The mineralized areas mentioned above have yielded concentrates worth at least \$10 million, and in some cases, as much as \$1 billion (Albers and Stewart, 1972; Cornwall, 1972; Longwell and others, 1965; Mardirosian, 1974a, b; and Tschanz and Pampeyan, 1970).

Gold and silver were produced from several mines in the Death Valley region as early as the 1860's, and by the early 1900's many of the original discoveries were substantially depleted. More than 124.4 Mg of gold and at least 43.5 Mg of silver were produced from small lodes in silicified and alunized north-trending fractures in the Tertiary Milltown Andesite in the Goldfield district (Albers and Stewart, 1972). The peak production period in the Goldfield district was from 1905 through 1912, and smaller quantities of ore were mined locally as late as 1955. The cumulative value of metals yielded from mines in the Goldfield district exceeds \$89 million, and this amount accounts for more than 75 percent of the total non-fuel mineral production in Esmeralda County to 1972 (Albers and Stewart, 1972).

At least 3110 Mg of silver (Kleinhampl, 1964), and 35.9 Mg of coproduct gold were mined in the Tonopah district (Bergendahl, 1964). These fault-controlled argentiferous replacement deposits near Tonopah occur in altered Tertiary rhyolite and andesite, and they have yielded precious and base-metal concentrates worth more than \$100 million (Albers and Stewart, 1972). Although the Goldfield and Tonopah districts contain deep workings, the most productive ore shoots in these areas occurred within 300 m of the surface. In recent years, the Anaconda open-pit mine in the San Antonio district north of Tonopah has produced a substantial quantity of molybdenum and some copper.

In addition to the important production of precious metals at Goldfield and Tonopah, large quantities of gold and silver have come from mineral deposits in the Silver Peak district, Esmeralda County, and the Manhattan district, Nye County. Precious metal lodes in the Silver Peak district occur as irregular quartz masses and thin anastomosing veins in locally metamorphosed sediments of the Precambrian Wyman Formation (Albers and Stewart, 1972). These gold-bearing ore-shoots, that also contain some silver, are related to a Late Jurassic or Early Cretaceous alaskite intrusive. More than 311 Mg of silver (Kleinhampl, 1964) and at least 18 Mg of gold (Bergendahl, 1964) were produced from mines in the Silver Peak district. The Manhattan district, Nye County, contains vein deposits in Cambrian(?) limestone and quartz schist in the hanging wall section of an extensive northwest-trending thrust fault (Ferguson, 1924). Thin discontinuous gold-quartz veins are also present in Tertiary volcanic rocks in the Manhattan district, however the lodes in the Paleozoic rocks have been the most productive. Lode ores at Manhattan have yielded more than 8.7 Mg of gold, at least 3.1 Mg of silver, and placer deposits in the district have produced an additional 6.2 Mg in gold (Bergendahl, 1964).

Precious metals were produced from the majority of the districts in the Death Valley region in Nevada. At least 3.1 Mg of gold were mined from ores in the Bullfrog, Ellendale, Johnnie, Lodi, Tybo, and Union districts of Nye County, the Bell district of Mineral County, and the Eldorado, Goodsprings and Searchlight districts of Clark County (Bergendahl, 1964). Silver production exceeded 31.1 Mg in the Goodsprings and Eldorado districts of Clark County, the Belmont and Tybo districts, Nye County, and the Divide district of Esmeralda County (Kleinhampl, 1964).

Many mineral deposits in the Death Valley region of California contain gold, but none of these deposits have recorded production comparable to similar deposits in Nevada. Gold concentrates valued at more than \$1 million were produced from mesothermal quartz veins in the Ballarat-South Park, Chloride Cliff, and Skiddoo-Tucki Mountain districts of Inyo County, California (Clark, 1970). At least 248.8 Mg of silver were produced from Darwin district until 1951 (Hall and MacKevett, 1962), and more than 155.5 Mg of silver came from mines in the Cerro Gordo district, Inyo County (Stager, 1966).

The principal production of base metals to date in the Death Valley region has come from the Goodsprings district of Clark County, and the Tem Piute district of Lincoln County, Nevada; and the Darwin, Cerro Gordo, Saratoga, and Lee districts of Inyo County, California. Ore minerals of lead and zinc have been the principal source of profits in these districts, whereas tungsten and byproduct silver are locally of special importance. Excluding the significant gold deposits near Goldfield, copper is of secondary importance in the majority of the base-metal deposits in the Death Valley region. The base-metal deposits in the Death Valley study area are mostly replacement deposits in faulted Paleozoic carbonate rocks. Silicic to intermediate intrusive rocks are common in many areas containing mineralized rocks. The mineral deposits at Darwin were developed continuously for long periods, while mining activities in the remaining base-metal districts generally were of short duration. Oxidation was extensive in many of the base-metal deposits, and much of the near-surface high-grade ore in these areas has been removed.

The Goodsprings district in Clark County has been an important source of zinc in Nevada. Deposits of several metals occur at Goodsprings, and zinc ores with associated lead have accounted for the bulk of the value and tonnage of all metals produced to date. The majority of the mineral deposits at Goodsprings are mantos, commonly occurring in the Upper Mississippian Yellowpine Member of the Monte Cristo Limestone (Hewett, 1931). The ores are related to granitic intrusives, and are commonly oxidized. Mines at Goodsprings have produced more than \$31 million in metals until 1962 (Longwell and others, 1965).

The Lincoln mine in the Tem Piute district was a major source of tungsten in Nevada between 1940 and 1957 and the area was also productive from 1975 to 1981. Approximately 287,000 20-pound units of tungsten trioxide were produced during this period as part of a federally subsidized stockpile program (Tschanz and Pampeyan, 1970). Silver and small quantities of zinc and lead were mined as byproducts. The principal deposits at Tem Piute were elongated zones in tectonics in Devonian and Mississippian limestones.

The principal sources of base-metals in the Death Valley region of California are replacement or open-space filling deposits in faulted Paleozoic carbonate rocks. These deposits commonly are similar to nearby silicic to intermediate intrusive rocks. The mines in the Darwin district, Inyo County, historically have been an important source of lead in California (Norman and Stewart, 1951). Large quantities of silver and zinc also have been mined as coproducts at Darwin. At least \$25 million worth of base and precious metals were mined at Darwin by 1953 (Carlisle and others, 1954). More than \$17 million in lead-silver ore was produced from the Cerro Gordo district principally before 1877 (Norman and Stewart, 1951). About 10,058 Mg of oxidized zinc ore was mined from cavity filling deposits at Cerro Gordo mainly between 1912 and 1919 (Carlisle and others, 1954).

The Shoshone mines in the Saratoga district, and the Santa Rosa mine in the Lee district, produced smaller, although still important, values and tonnages of base-metals. These deposits are in fault-controlled silicified Paleozoic carbonate rocks.

In addition to the principal base and precious metals mentioned in the preceding discussion, many other metallic elements occur in variable quantities in the Death Valley region. Some of the more common metals include: arsenic, antimony, beryllium, bismuth, iron, manganese, mercury, molybdenum, and uranium. Substantial quantities of these elements have been produced only locally from a few mines in the study area to date. In contrast, one of the largest identified concentrations of rare-earth elements in the world occurs in the Death Valley region, near Mountain Pass, San Bernardino County, California. Large bastnaesite-bearing carbonate ore bodies, and several hundred rare-earth vein deposits are associated with potash-enriched intrusives of the Mountain Pass mineral district (Olson and others, 1954). This deposit currently is the largest domestic source of rare-earth elements (Carrillo and others, 1983).

Industrial Mineral Resources

Several varieties of nonmetallic industrial minerals and rocks are located in the Death Valley region. Many of these commodities have been produced intermittently, and shipped to local markets. Deposits of semiprecious and precious gemstones, natural aggregates, and building stone are not discussed in this report. Overviews of the occurrences of the principal industrial rocks and minerals including construction materials are contained in reports by the U.S. Geological Survey (1964, 1966). Detailed commodity discussions occur in several reports published by the California Division of Mines and Geology and the Nevada Bureau of Mines and Geology.

Magnesite, brucite, fluorspar, and barite are the principal nonmetallic or industrial commodities with significant production in the Death Valley region within Nevada. Magnesite and brucite are mined currently near Gabbs in northwest Nye County. This deposit is one of two presently active domestic magnesium mines, and these deposits are among the more important nonmetallic mineral occurrences in the study area. The replacement deposit at Gabbs occurs in dolomite of the Upper Triassic Luning Formation (Callaghan, 1933). The locally recrystallized host rocks are part of the upper plate of the Paradise thrust fault and have been intruded by numerous dikes and a Cretaceous granodiorite stock (Martin, 1956). The magnesite ore zones are satellitic around an extension of the granodiorite stock, while the principal brucite deposits are in contact with the intrusive (Vitaliano and Callaghan, 1956, 1963; Vitaliano and others, 1957). Several million metric tons of ore containing less than 5 percent admixed calcium oxide were mined prior to 1968. These deposits contained estimated reserves of at least 23 million Mg of high-grade ore in 1968 (Schilling, 1968).

Several deposits of fluorspar are in the Bare Mountain or Fluorine district, southern Nye County. The Daisy mine has been the principal source of metallurgical-grade fluorspar in the district. At least 172,482 Mg of fluorspar were produced from the Daisy mine between 1919 and 1976 (Papke, 1979) where it occurs as fracture-fillings in dolomite of the Upper Cambrian Nopah Formation (Thurston, 1949; Cornwall and Kleinhampl, 1964). Smaller although productive breccia pipe and fissure deposits containing fluorspar occur on the east side of Bare Mountain. These workings include the Mary and Goldspar mines. The Mary mine has yielded an estimated 11,800 Mg of fluorspar averaging 40 percent calcium fluoride. In addition, about 68,085 Mg of 40-percent calcium fluoride-bearing ore came from the Goldspar mine (Papke, 1979).

Barite occurrences are abundant in the Death Valley region of Nevada, however, only the Jumbo mine in the Ellendale district, central Nye County has produced more than 9,000 Mg of ore (Horton, 1964). Substantial quantities of manganese oxide have been produced at the Three Kids mine in the area north of Boulder City (Hewett and Webber, 1931).

Talc is the principal industrial mineral of known importance in the Death Valley region of California. California historically has been an important domestic source of steatite. The majority of the high-grade talc deposits in the California part of the study area occur in a zone that extends for about 121 km through southern Death Valley eastward to the Kingston Range in northeast San Bernardino County (Wright, 1964). These talc deposits are commonly irregular and lenticular bodies located near the contacts between diabase sills and silicified carbonate rocks of the Precambrian Crystal Spring Formation. The host rocks are intensely deformed, and faulting is common in the weak talc-bearing horizons. The majority of the talc mines in the area are developed from 1,305 m to 1,524 m along the outcrop, and to depths less than 152 m (Wright, 1964). More than 1 million Mg of talc-bearing material were produced from the mines in the southern Death Valley-Kingston Range region by 1959 (Wright, 1968).

Brines pumped from beneath the surface of the dry lake in the Silver Peak Marsh district, central Esmeralda County, are currently an important world source of lithium. The playa sediments contain abundant hectorite, and are saturated with brine to depths of 183 to 451 m (Kunasz, 1971). These brines, which contain about 300 mg/L lithium (Barrett and O'Neill, 1970), are evaporated to produce a concentrated product for shipment. The well field at Silver Peak Marsh has supplied very large quantities of lithium since 1966 (Norton, 1973), however no production statistics are available at present.

Borates in the Death Valley region of Nevada occur as bedded Tertiary deposits and as marsh deposits containing mostly ulexite. Colemanite, ulexite, and probertite locally are abundant in folded Tertiary lake beds near Furnace Creek in Death Valley and in the Amargosa Valley, Inyo County (Ver Planck, 1950; McAllister, 1970; 1973). A large colemanite deposit occurs in the area of Callville Wash, about 42 km east of Las Vegas, which was developed by two adits and 2,134 m of underground workings. Approximately 181,560 Mg of colemanite that averaged 20 percent B₂O₃ were produced from the Callville Wash deposit between 1921 and 1928 (Vanderburg, 1937a). Bedded ulexite was mined as late as 1939 from deposits east of Fish Lake Valley, Esmeralda County (Smith, 1964). Ulexite was mined from marsh deposits from 1870 to 1892, many of which are in the Death Valley region.

Extensive gypsum deposits occur in Permian rocks along the east slope of the Spring Mountains west and southwest of Las Vegas. The estimated annual production during 1965 was 272,340 Mg (Longwell and others, 1965). Large gypsum resources also exist in Permian and Tertiary strata in the southern end of Frenchman Mountain, west-central Clark County. Annual production from this area was estimated by Longwell and others (1965) to be 90,780 Mg.

Besides borates, the principal saline minerals in the Death Valley region of California are halite and gypsum. Production of evaporite minerals from playa deposits has been small and of short duration, however, large resources of saline minerals are identified locally. Salt beds, as thick as 6 m and interbedded lacustrine clays were penetrated to a depth of about 305 m near Badwater in Death Valley (Bain, 1914; Gale, 1914). Tertiary lake beds in the Avawatz Range area are reported to contain about 1.2 million Mg of salt within about 50 m of the surface (Ver Planck, 1958). In addition, near-surface deposits of gypsum occur near Tecopa (Withington, 1966), and in the Avawatz Range area (Ver Planck, 1958).

Geothermal Resources

There are currently two Known Geothermal Resources Areas (KGRA) in the Death Valley region. The Saline Valley KGRA has an area of 1,295 ha in T. 13 S., R. 39 E., Inyo County, California and the Silver Peak KGRA has an area of 2,071 ha in T. 2. S., R. 39 E., Esmeralda County Nevada (U.S. Bureau of Land Management, 1983a,b). These lands are classified as such based on formal determination by the Bureau of Land Management of competitive interest in the lands (Burkhardt and others, 1980). Competitive interest, as defined in Title 43, Chapter II of the Code of Federal Regulations, subpart 3200.0.5 (k)(3), exists "...in the entire area covered by an application for a geothermal lease if at least one-half of the lands covered by that application are also covered by another application which was filed during the same application filing period". The total area in the Saline Valley KGRA is covered by Federal competitive interest, and 20,266 Federal competitive ha are in the Silver Peak KGRA (Burkhardt and others, 1980). The Saline Valley and Silver Peak KGRAs both contain hot springs, and these areas are characterized further by warm surface temperatures of 65° and 48° C respectively (Burkhardt and others, 1980). The temperature range for the Saline Valley area geothermal reservoir is estimated between 65° and 96° C (Burkhardt and others, 1980). These data are projected from well temperatures or chemical geothermometry. No similar data currently are available for the geothermal reservoir near Silver Peak.

Geothermal water within 915 m of the surface, with temperatures sufficient for direct heat application, occur north of Tecopa, and along Furnace Creek Wash in Inyo County (Higgins, 1980). These geothermal occurrences in the Death Valley region are known to have temperatures greater than 50 °C. An oil-test hole was drilled by Nevada Oil and Minerals in 1970 at Fish Lake Valley, Esmeralda County, to a depth of 2,798 m. The bottom-hole temperature of this well was recorded on a temperature log at 158 °C (Garside and Schilling, 1979). Alkali Springs, 16 km northwest of Goldfield, Esmeralda County, have produced water of at least 60 °C (Ball, 1907). In addition, springs in the Black Canyon of Nevada and Arizona yield warm water with temperatures as great as 62 °C (Garside and Schilling, 1979). More than 50 thermal wells and 40 warm springs occur in the Death Valley region, however the majority of the temperatures in these wells and springs are between 20 ° and 35 °C.

Ball, S. H., 1906, Notes on the ore deposits of southwestern Nevada and eastern California, in Emmons, S. F., and Eckel, E. C., eds., Contributions to economic geology, 1905; U.S. Geological Survey Bulletin 285-A, p. 53-73.

----- 1907, A geologic reconnaissance in southwestern Nevada and eastern California; U.S. Geological Survey Bulletin 308, 218 p.

Barrett, W. T., and O'Neill, B. J., Jr., 1970, Recovery of lithium from saline brines using solar evaporation, in Third symposium on salt, v. 2: Cleveland, Ohio, Northern Ohio Geological Society, p. 47-50.

Beal, L. H., 1963, Investigation of titanium occurrences in Nevada; Nevada Bureau of Mines Report 159, 42 p.

Coal, Oil, and Gas Resources

The only known coal-bearing strata in the Death Valley region are in the Coaldale field, Esmeralda County. Thin bituminous coals, commonly 1 m to 2.1 m thick, occur at 4 horizons in moderately tilted and extensively faulted interbedded shale, Tertiary sandstone, bentonite, and tuff (Andrews and others, 1947; Hance, 1913). The Coaldale field has produced a very small tonnage of coal. There are currently no known oil or gas fields or producing hydrocarbon wells in the Death Valley region, though at least 60 dry oil or gas holes have been completed to date in the study area (Brady, 1983). Historically, the greatest effort in the search for hydrocarbons in the Death Valley region has been focused on Paleozoic rocks in the Las Vegas Valley. Several very deep boreholes have been completed in this area, but none of the holes produced appreciable quantities of oil or gas.

References Cited

- Albers, J. P., and Stewart, J. H., 1972, Geology and mineral deposits of Esmeralda County, Nevada: Nevada Bureau of Mines and Geology Bulletin 78, 80 p.
- Albritton, C. C., Jr., Richards, Arthur, Brokaw, A. L., and Reinemund, J. A., 1954, Geologic controls of lead and zinc deposits in Goodsprings (Yellow Pine) district, Nevada: U.S. Geological Survey Bulletin 1010, 111 p.
- Andrews, D. A., Hendricks, T. A., and Huddle, J. W., 1947, Analyses of Arizona, California, Idaho, Nevada and Oregon coals: U.S. Bureau of Mines Technical Paper 696, 47 p.
- Bain, H. F., 1914, Potassium salts: Mineral Industry, Statistics, Technology and Trade, v. 22, p. 617.
- Bailey, E. H., and Phoenix, D. A., 1944, Quicksilver deposits in Nevada: Nevada University Bulletin, v. 38, no. 5, Geology and Mining Series 41, 206 p.
- Ball, S. H., 1906, Notes on the ore deposits of southwestern Nevada and eastern California, in Emmons, S. F., and Eckel, E. C., eds., Contributions to economic geology, 1905: U.S. Geological Survey Bulletin 285-A, p. 53-73.
- 1907, A geologic reconnaissance in southwestern Nevada and eastern California: U.S. Geological Survey Bulletin 308, 218 p.
- Barrett, W. T., and O'Neill, B. J., Jr., 1970, Recovery of lithium from saline brines using solar evaporation, in Third symposium on salt, v. 2: Cleveland, Ohio, Northern Ohio Geological Society, p. 47-50.
- Beal, L. H., 1963, Investigation of titanium occurrences in Nevada: Nevada Bureau of Mines Report 3, 42 p.

- Benson, W. T., 1950, Investigation of Black Rock manganese deposits, Esmeralda County, Nevada: U.S. Bureau of Mines Report of Investigations 4717, 5 p.
- Bergendahl, M. H., 1964, Gold, in Mineral and water resources of Nevada: U.S. Geological Survey, 88th Congress, second session, Senate Committee on Interior and Insular Affairs, Senate Document no. 87, p. 87-100.
- Bonham, H. F., 1970, Geologic map and sections of a part of the Shoshone Mountains, Lander and Nye Counties, Nevada: Nevada Bureau of Mines Map 38, scale 1:62,500.
- Brady, B. T., 1983, Map showing coal, oil and gas wells, seeps, and tar sandstone occurrences in the Basin and Range Province: U.S. Geological Survey Open-File Report 83-549, map scale 1:2,500,000, 104 p.
- Burkhardt, H. E., Brook, C. A., and Smith, F. W., 1980, Selected administrative, land, and resource data for Known Geothermal Resources Areas in Arizona, California, Idaho, Nevada, Oregon and Washington: U.S. Geological Survey Open-File Report 80-1290, 29 p.
- Callaghan, Eugene, 1933, Brucite deposit, Paradise Range, Nevada--A preliminary report: Nevada Bureau of Mines Bulletin 14, 34 p.
- 1939, Geology of the Searchlight mining district, Clark County, Nevada: U.S. Geological Survey Bulletin 906-D, p. 135-188.

- Carlisle, Donald, Davis, D. L., Kildale, M. B., and Stewart, R. M., 1954, Base metal and iron deposits of southern California, in Jahns, R. H., ed., Geology of southern California: California Division of Mines Bulletin 170, pt. 8, p. 41-50.
- Carrillo, F. V., Davis, J. F., and Burnett, J. L., 1983, The mineral industry in California: U.S. Bureau of Mines Mineral Yearbook, v. 2, p. 87-98.
- Clark, W. B., 1970, Gold districts of California: California Division of Mines and Geology Bulletin 193, 186 p.
- Cornwall, H. R., 1972, Geology and mineral deposits of southern Nye County, Nevada: Nevada Bureau of Mines and Geology Bulletin 77, 49 p.
- Cornwall, H. R., and Kleinhampl, F. J., 1964, Geology of the Bullfrog Quadrangle and ore deposits related to Bullfrog Hills caldera, Nye County, Nevada and Inyo County, California: U.S. Geological Survey Professional Paper 454-J, 25 p.
- Couch, B. F., and Carpenter, J. A., 1943, Nevada's metal and mineral production (1859-1940, inclusive): Nevada Bureau of Mines Bulletin 38, 159 p.
- Davis, F. F., 1957, Manganese, in Wright, L. A., ed., Mineral commodities of California: California Division of Mines Bulletin 176, p. 325-329.
- Duncan, D. C., 1953, A uranium-bearing rhyolitic tuff deposit near Coaldale, Esmeralda County, Nevada: U.S. Geological Survey Circular 291, 7 p.

- Eric, J. H., 1948, Tabulation of copper deposits in California: California Division of Mines and Geology Bulletin 144, pt. 3, p. 199-387.
- Ferguson, H. G., 1917, The Golden Arrow, Clifford, and Ellendale districts, Nye County, Nevada: U.S. Geological Survey Bulletin 640-F, p. 113-123.
- 1924, Geology and ore deposits of the Manhattan district, Nevada: U.S. Geological Survey Bulletin 723, 163 p.
- 1927, The Gilbert district, Nevada: U.S. Geological Survey Bulletin 795-F, p. 125-145.
- 1933, Geology of the Tybo district, Nevada: Nevada Bureau of Mines and Geology Bulletin 20, 61 p.
- Gale, H. S., 1914, Prospecting for potash in Death Valley, California: U.S. Geological Survey Bulletin 540, p. 407-415.
- Garside, L. J., 1973, Radioactive mineral occurrences in Nevada: Nevada Bureau of Mines and Geology Bulletin 81, 121 p.
- Garside, L. J., and Schilling, J. H., 1979, Thermal waters of Nevada: Nevada Bureau of Mines and Geology Bulletin 91, 163 p.
- Goodwin, J. G., 1957, Lead and zinc in California: California Journal of Mines and Geology, v. 53, nos. 3 and 4, p. 353-724.
- Hall, W. E., and MacKevett, E. M., Jr., 1962, [1963], Geology and ore deposits of the Darwin Quadrangle, Inyo County, California: U.S. Geological Survey Professional Paper 368, 87 p.
- Hance, J. H., 1913, The Coaldale coal field, Esmeralda County, Nevada: U.S. Geological Survey Bulletin 531-K, p. K313-K322.

Hewett, D. F., 1931, Geology and ore deposits of the Goodsprings
Quadrangle, Nevada: U.S. Geological Survey Professional
Paper 162, 172 p.

----- 1956, Geology and mineral resources of the Ivanpah
Quadrangle, California and Nevada: U.S. Geological Survey
Professional Paper 275, 172 p.

Hewett, D. F., Callaghan, Eugene, Moore, B. N., Nolan, T. B.,
Rubey, W. W., and Schaller, W. T., 1936, Mineral resources
of the region around Boulder Dam: U.S. Geological Survey
Bulletin 871, 197 p.

Hewett, D. F., and Webber, B. N., 1931, Bedded deposits of
manganese oxide near Las Vegas, Nevada: Nevada University
Bulletin, v. 25, no. 6, 17 p.

Hill, J. M., 1916, Notes on some mining districts in eastern
Nevada: U.S. Geological Survey Bulletin 648, 214 p.

Higgins, C. T., 1980, Geothermal resources of California:
California Division of Mines and Geology, scale 1:750,000.

Holmes, G. W., Jr., 1965, Mercury in Nevada, in U.S. Bureau
of Mines Staff, Mercury potential of the United States: U.S.
Bureau of Mines Information Circular 8252, p. 215-300.

Horton, R. C., 1963, An inventory of barite occurrences in Nevada:
Nevada Bureau of Mines Report 4, 18 p.

----- 1964, Barite, in Mineral and water resources of Nevada:
U.S. Geological Survey, 88th Congress, second session, Senate
Committee on Interior and Insular Affairs, Senate Document
87, p. 177-180.

- Keith, W. J., 1977, Geology of the Red Mountain mining district, Esmeralda County, Nevada: U.S. Geological Survey Bulletin 1423, 45 p.
- Kleinhampl, F. J., 1964, Silver, lead and zinc, in Mineral and water resources of Nevada: U.S. Geological Survey, 88th Congress, second session, Senate Committee on Interior and Insular Affairs, Senate Document no. 87, p. 138-150.
- Kleinhampl, F. J., and Ziony, J. K., 1984, Mineral resources of northern Nye County, Nevada: Nevada Bureau of Mines and Geology Bulletin 99, map scale 1:250,000, 243 p.
- Knopf, Adolph, 1915, A gold-platinum-palladium lode in southern Nevada: U.S. Geological Survey Bulletin 620-A, p. 1-18.
- 1921a, Ore deposits of Cedar Mountain, Mineral County, Nevada: U.S. Geological Survey Bulletin 725, p. 361-382.
- 1921b, The Divide silver district, Nevada, in Contributions to economic geology: U.S. Geological Survey Bulletin 715, p. 147-170.
- Kral, V. E., 1951, Mineral resources of Nye County, Nevada: Nevada Bureau of Mines Bulletin 50, 223 p.
- Kunasz, I. A., 1971, Geology and geochemistry of the lithium deposits in Clayton Valley, Esmeralda County (abs.): Dissertation Abstracts, v. 32, no. 1, p. 1215-1226.
- Lawrence, E. F., 1963, Antimony deposits of Nevada: Nevada Bureau of Mines Bulletin 61, 248 p.
- Lemmon, D. M., and Tweto, O. L., compilers, 1962, Tungsten in the United States, exclusive of Alaska and Hawaii: U.S. Geological Survey Mineral Investigations Resource Map MR-25, scale 1:3,168,000.

- Lincoln, F. C., 1923, Mining districts and mineral resources of Nevada: Reno, Nevada Newsletter Publishing Co., 295 p.
- Longwell, C. R., Pampeyan, E. H., Bowyer, Ben, and Roberts, R. J., 1965, Geology and mineral deposits of Clark County, Nevada: Nevada Bureau of Mines and Geology Bulletin 62, 218 p.
- Mann, R. L., 1916, Owl Head manganese deposit, San Bernardino County, California: Mining and Engineering World, v. 44, p. 743-744.
- Mardirosian, C. A., 1974a, Mining districts and mineral deposits of California: Mineral Research Co., Laredo, Texas, scale 1:1,000,000, 2 sheets.
- 1974b, Mining districts and mineral deposits of Nevada (exclusive of oil, gas, and water): Mineral Research Co., Laredo, Texas, scale 1:1,000,000.
- Martin, Conrad, 1956, Structure and dolomitization in crystalline magnesite deposits, Paradise Range, Nye County, Nevada (abs.): Geological Society of America Bulletin, v. 67, no. 12, pt. 2, p. 1774.
- McAllister, J. F., 1970, Geology of the Furnace Creek borate area, Death Valley, Inyo County, California: California Division of Mines and Geology Map Sheet 14, scale 1:24,000, 9 p.
- 1973, Geologic map and sections of the Amargosa Valley borate area--Southeast continuation of the Furnace Creek area--Inyo County, California: U.S. Geological Survey Miscellaneous Investigations Map I-782, scale 1:24,000.

- Norman, L. A., and Stewart, R. M., 1951, Mines and mineral resources of Inyo County: California Journal of Mines and Geology, v. 47, no. 1, p. 17-23.
- Norton, J. J., 1973, Lithium, cesium and rubidium--The rare alkali metals, in Brobst, D. A., and Pratt, W. P., eds., United States mineral resources: U.S. Geological Survey Professional Paper 820, p. 365-378.
- Olson, J. C., Shawe, D. R., Pray, L. C. and Sharp, W. N., 1954, [1955], Rare-earth mineral deposits of the Mountain Pass district, San Bernardino County, California: U.S. Geological Survey Professional Paper 261, 75 p.
- Oxnam, T. H., 1936, Weepah gold: Engineering and Mining Journal, v. 137, no. 6, p. 300-306.
- Papke, K. G., 1979, Fluorspar in Nevada: Nevada Bureau of Mines and Geology Bulletin 93, 77 p.
- Ransome, F. L., 1907, Preliminary account of Goldfield, Bullfrog, and other mining districts in southern Nevada, with notes on The Manhattan district, by G. H. Garrey, and W. H. Emmons: U.S. Geological Survey Bulletin 303, 98 p.
- 1909, The geology and ore deposits of Goldfield, Nevada: U.S. Geological Survey Professional Paper 66, 258 p.
- Ransome, F. L., Emmons, W. H., and Garrey, G. H., 1910, Geology and ore deposits of the Bullfrog district, Nevada: U.S. Geological Survey Bulletin 407, 130 p.
- Leaves, R. G., Shawe, F. R., and Kral, V. E., 1958, Iron ore deposits of west-central Nevada: Nevada Bureau of Mines Bulletin 53, pt. B, 46 p.

- Root, W. A., 1909, The Lida mining district of Nevada: Mining World, v. 31, no. 2, p. 123-125.
- Ross, D. C., 1961, Geology and mineral deposits of Mineral County, Nevada: Nevada Bureau of Mines Bulletin 58, 98 p.
- Sampson, R. J., and Tucker, W. B., 1940, Mineral resources of Mono County: California Journal of Mines and Geology, v. 36, no. 2, p. 117-156.
- Schilling, J. H., 1962, An inventory of molybdenum occurrences in Nevada: Nevada Bureau of Mines Report 2, 48 p.
- 1963, Tungsten mines in Nevada: Nevada Bureau of Mines Map 18, scale 1:1,000,000.
- 1968, The Gabbs magnesite-brucite deposit, Nye County, Nevada, in Ridge, J. D., ed., Ore deposits in the United States 1933/1967: Graton-Sales, v. 2, p. 1607-1622.
- Smith, G. I., Troxel, B. W., Gray, C. H., Jr., and Von Huene, Roland, 1968, Geologic reconnaissance of the Slate Range, San Bernardino and Inyo Counties, California: California Division of Mines and Geology Special Report 96, 33 p.
- Smith, W. C., 1964, Borates, in Mineral and water resources of Nevada: U.S. Geological Survey, 88th Congress, second session, Senate Committee on Interior and Insular Affairs, Senate Document 87, p. 180-184.
- Spurr, J. E., 1905, Geology of the Tonopah mining district, Nye County, Nevada: U.S. Geological Survey Professional Paper 42, 295 p.
- 1906, Ore deposits of the Silver Peak Quadrangle, Nevada: U.S. Geological Survey Professional Paper 55, 174 p.

- Stager, H. K., 1966, Silver, in Mineral and water resources of California, U.S. Geological Survey, 89th Congress, second session, Part 2, Senate Committee on Interior and Insular Affairs, Committee Print, p. 381-385.
- Stretch, R. H., 1904, The Montezuma district, Nevada: Engineering and Mining Journal, v. 78, no. 1, p. 5-6.
- Thurston, W. R., 1949, The Daisy fluorspar deposit near Beatty, Nye County, Nevada: U.S. Geological Survey Strategic Minerals Preliminary Investigation Report 3-209, 10 p.
- Toenges, A. L., Turnbull, L. A., and Schopf, J. M., 1946, Exploration, origin, and constitution of the coal deposit: U.S. Bureau of Mines Technical Paper 687, pt. 1, p. 1-35.
- Trask, P. D., 1950, Geologic description of the manganese deposits of California: California Division of Mines Bulletin 152, 378 p.
- Trengove, R. R., 1959, Reconnaissance of Nevada manganese deposits: U.S. Bureau of Mines Report of Investigations 5446, 40 p.
- Tschanz, C. M., and Pampeyan, E. H., 1970, Geology and mineral deposits of Lincoln County, Nevada: Nevada Bureau of Mines and Geology Bulletin 73, 187 p.
- U.S. Bureau of Land Management, 1983a, Nevada land classification status map: Washington, D.C., scale 1:500,000.
- 1983b, California, south half, land classification status map: Washington, D.C., scale 1:500,000.

- U.S. Geological Survey, 1964, Mineral and water resources of Nevada: U.S. Geological Survey, 88th Congress, second session, Senate Committee on Interior and Insular Affairs, Senate Document 87, 314 p.
- 1966, Mineral and water resources of California, Part I: U.S. Geological Survey, 89th Congress, second session, Senate Committee on Interior and Insular Affairs, Committee Print, 450 p.
- Vanderburg, W. O., 1936, Placer mining in Nevada: Nevada University Bulletin, v. 30, no. 4, 180 p.
- 1937a, Reconnaissance of mining districts in Clark County, Nevada: U.S. Bureau of Mines Information Circular 6964, 81 p.
- 1937b, Reconnaissance of mining districts in Mineral County, Nevada: U.S. Bureau of Mines Information Circular 6941, 79 p.
- Ver Planck, W. E., 1950, Boron, in Mineral commodities of California: California Division of Mines Bulletin 156, p. 215-219.
- 1958, Salt in California: California Division of Mines Bulletin 175, 168 p.
- Vitaliano, C. J., and Callaghan, Eugene, 1956, Geologic map of the Gabbs magnesite and brucite deposits, Nye County, Nevada: U.S. Geological Survey Mineral Investigations Field Study Map MF-35, scale 1:24,000.
- 1963, Geology of the Paradise Peak Quadrangle, Nevada: U.S. Geological Survey Geologic Quadrangle Map GQ-250, scale 1:62,500.

- Vitaliano, C. J., Callaghan, Eugene, and Silberling, N. L., 1957, Geology of Gabbs and vicinity, Nye County, Nevada: U.S. Geological Survey Mineral Investigations Field Study Map MF-52, scale 1:24,000.
- Walker, G. W., Lovering, T. G., and Stephens, H. G., 1956, Radioactive deposits in California: California Division of Mines and Geology Special Report 49, 38 p.
- Waring, C. A., and Huguenin, Emile, 1919, Inyo County, in 15th report of the State Mineralogist: California Journal of Mines and Geology, v. 15, p. 29-134.
- White, D. E., 1940, Antimony deposits of the Wildrose Canyon area, Inyo County: U.S. Geological Survey Bulletin 992-K, p. 307-325.
- Withington, C. F., 1966, Gypsum and anhydrite, in Mineral and water resources of California, Part I: U.S. Geological Survey, 89th Congress, second session, Part 2, Senate Committee on Interior and Insular Affairs, Committee Print, p. 188-198.
- Wong, George, 1983a, Preliminary map of the resource areas in the Basin and Range area of California: U.S. Geological Survey Open-File Report 83-719, map scale 1:500,000, 15 p.
- 1983b, Preliminary map of the resource areas in the Basin and Range area of Nevada: U.S. Geological Survey Open-File Report 83-721, map scale 1:500,000, 46 p.
- Wright, L. A., 1964, Talc and soapstone, in Mineral and water resources of California, Part I: U.S. Geological Survey, 89th Congress, second session, Senate Committee on Interior and Insular Affairs, Committee Print, p. 414-420.

----- 1968, Talc deposits of the southern Death Valley-Kingston Range region, California: California Division of Mines and Geology Special Report 95, 79 p.

Wright, L. A., Stewart, R. M., Gay, T. E., Jr., and Hazenbush, G. C., 1953, Mines and mineral deposits of San Bernardino County, California: California Journal of Mines and Geology, v. 49, nos. 1 and 2, p. 49-257.

Young, G. J., 1920, Divide silver-gold district of Nevada: Engineering and Mining Journal, v. 109, no. 2, p. 62-66.

Open-File Report 84-743

Plates 1 to 7

Also published as
Professional Paper 1370-F

POCKET CONTAINS
7 ITEMS

