Potassium-argon age determinations from the Port Moller
and Stepovak Bay quadrangles, Alaska Peninsula
by
Gregory D. DuBois, Frederic H. Wilson, and Nora B. Shew

Reported here are 17 new potassium-argon age determinations on 11 volcanic rocks and 6 intrusive rocks from the Port Moller, Steptovak Bay and Simeonof Island quadrangles. These were done in conjunction with an Alaska Mineral Resources Assessment Program (AMRAP) study for these quadrangles. The samples dated yielded ages of Eocene or younger, ranging from 33 to 37.13 Ma.

Table 1 shows the analytical dates. Each sample was extracted twice and the mean value is reported as the final age determination. Argon extraction and measurement were done by the University of Arizona, Tucson, where the samples were also analyzed for lead by the method of McDougall and Lanthier (1960). Sample preparation, argon extraction and gas reduction were done at the University of California, Los Angeles. Potassium ages were determined by the electron microprobe technique of Fleck et al. (1970) or by the laser fluorescence photometry using a lithium metaborate fusion technique (Engel and Ingalls, 1970). Potassium concentrations were determined by atomic absorption spectrophotometry. All potassium analyses were replicated 2-4 times and the mean is shown as well. Potassium analytical errors are given in parentheses. The error for the age calculation is estimated to be about ±1% or assigned to each age (Table 1), is an estimate of the standard deviation of analytical precision using the method of Cox and Dalrymple (1967) and the authors calculated estimates of the error for the age calculation from the error in the potassium concentration and the error in the gas reduction from hand sample and thin section for each sample. Hand sample colors were determined using Gifford (1948) except that we noted that the Mussett Soil Color Chart (1975) used.

References cited

5. Allan, and Dalrymple, G.B., 1967, Statistical analysis of geomagnetic reversal data and the precision of potassium-argon dating: *Journal of Geophysical Research*, v. 72, n. 10, p. 2603-2604.

rymple, G.B., and Lanphere, M.A., 1969, Potassium-argon dating: San Francisco, W.H. Freeman and Co., 258p.

els J.C., and Inganells, C.O., 1970, Effect of sample inhomogeneity in K-Ar dating: *Geochimica et Cosmochimica Acta*, v.34, p. 1007-1017.

sell Soil Color Chart, 1975: Munsell Color, Macbeth Division, Kollmorgen Corporation, Baltimore, 8 plates.

son, F.H., 1980, Late Mesozoic and Cenozoic tectonics and the age of porphyry copper prospects; Chignik and Sutwik Island quadrangles, Alaska Peninsula: U.S. Geological Survey Open-file Report 80-543, 94 p., 5 plates, scales 1:250,000 and 1:12,000.

Table 1. Potassium-argon age determinations.

Sample and Locality	Latitude and Longitude	Rock Type	Mineral	Fe ₂ O ₃	K ₂ O	⁴⁰ K _{rad} x10 ⁻¹²	²⁴⁰ Am _{rad}	Apparent Age and error (m.y.)
82AS49 Popof Strait	55 19.0°N 160 34.0°W	Tuff	BIO	7.88 7.92	374.4 7.915	374.4 34.31	27.55 31.16	32.56 ± .45 34.51 ± .18
82AS50 Popof Strait	55 19.4°N 160 34.2°W	Andesite	PLAG-HF	0.226 0.234 0.230 0.226	12.23 0.229	12.23 12.23	36.07 71.38	37.73 ± .08 37.53 ± .76
82AW10 Cape Alaiensis	55 30.0°N 160 47.8°W	Andesite	PLAG-HF	0.404 0.402 0.404 0.402	5.068 4.624	5.068 5.068	39.48 56.24	10.47 ± .09 10.73 ± .13
82AW15 Lake Priest	55 10.1°N 160 46.7°W	Andesite	QSER	3.51 3.61	167.45 166.76	167.45 166.76	92.60 92.60	10.40 ± .49 31.52 ± .39
82AW16 Crater Unga Is.	55 10.1°N 160 46.7°W	Andesite	QSER	3.59 3.59	167.45 166.76	167.45 166.76	92.60 92.60	31.78 ± .61
82AW32 Swedena Peninsula	55 33.5°N 160 32.5°W	Andesite	PLAG-HF	0.390 0.593 0.596 0.593	7.907 8.963	7.907 8.963	39.09 66.68	9.20 ± .11 10.43 ± .12
82AW36 Swedena Peninsula	55 33.1°N 160 33.7°W	Dacite	PLAG-HF	2.511 2.504	32.18 32.50	32.18 32.50	76.07 76.88	8.88 ± .04 8.98 ± .06
82AW41b American Bay	55 43.1°N 160 12.4°W	Dacite	HBL	0.539 0.539	6.33 6.33	6.33 6.33	20.91 26.24	8.40 ± .11 9.26 ± .07
82AW45 South of West Bay	55 50.2°N 160 01.78°W	Andesite	PLAG-HF	0.356 0.355	0.32 0.336	0.32 0.336	7.07 5.01	0.63 ± .03 0.44 ± .02
82AW46 Rasey Bay	55 49.0°N 158 53.4°W	Granite		9.02 9.03	125.09 120.63	125.09 120.63	68.31 64.38	9.60 ± .07 9.26 ± .07
82AW100 Rasey Bay	55 50.4°N 159 54.8°W	Andesite	HBL	0.325 0.326	2.45 2.45	2.45 2.45	13.60 11.42	5.15 ± .10 5.50 ± .13
82AW125 Ukaiyuk "Onion"	55 12.0°N 161 08.1°W	Basalt	PLAG-HF	0.082 0.079	1.71 1.88	1.71 1.88	28.44 27.22	14.95 ± .82 14.95 ± .82
82AW126 Mesa Cape plateau	55 37.1°N 161 39.1°W	Quartz- diorite	BTD	9.51 9.58	42.87 45.46	42.87 45.46	34.95 32.70	3.12 ± .04 3.31 ± .04
82AW129a Island	55 11.4°N 161 20.2°W	Diorite	PLAG-HF	0.313 0.317	8.00 7.94	8.00 7.94	71.36 64.92	17.55 ± .21 17.62 ± .22
82AW129b Island	55 11.2°N 161 20.3°W	Diorite	PLAG-HF	0.313 0.302	8.00 7.94	8.00 7.94	71.36 64.92	17.55 ± .21 17.62 ± .22
82AW130 Island	55 12.0°N 161 20.3°W	Basalt	WR-HF	0.251 0.251	2.95 3.16	2.95 3.16	16.10 10.10	8.14 ± .13 8.71 ± .14
82AW135 Point	55 47.8°N 159 51.3°W	Diorite	HBL	0.232 0.302	2.60 2.62	2.60 2.62	12.42 12.91	5.96 ± .12 6.17 ± .14
82AW212 Pinck Peak	55 44.5°N 160 45.2°W	PLAG-HF		0.517 0.501	0.494 0.486	0.494 0.486	10.62 10.62	0.86 ± .03 0.66 ± .04
82AW213 Peak	55 44.5°N 160 45.2°W	PLAG-HF		0.517 0.501	0.494 0.486	0.494 0.486	10.62 10.62	0.67 ± .05

Rate determinations calculated with constants given below:

$= 4.963 \times 10^{-10} \text{ year}^{-1}$	$= 5.72 \times 10^{-11} \text{ year}^{-1}$	$= 8.78 \times 10^{-13} \text{ year}^{-1}$
---	--	--

K/K = 1.167×10^{-4} atom percent.
O = biotite, HBL = hornblende, PLAG = plagioclase, QSER = whole-rock with quartz-sericite
terration, WR = whole-rock. HF = hydrofluoric acid treated, see Wilson (1980).

from U.S. Geological Survey topo sheets, 1:250,000.
Miller; Stepovak Bay; Simeonof Island, Alaska.
Prepared by Thematic Mapping in Menlo Park, CA, 1982

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards.

SCALE 1:250 000

0 5 10 15 20 25 MILES

0 5 10 15 20 25 KILOMETERS

CONTOUR INTERVALS 200, 250, 500, AND 1000 FEET

POTASSIUM-ARGON AGE DETERMINATIONS FROM THE PORT MOLLER AND STEPOVAK BAY QUADRANGLES, ALASKA PENINSULA

by
Gregory D. DuBois, Frederic H. Wilson, and Nora B. Shew
1987