

Base: Prepared by the U.S. Geological Survey
Data Sources: The World Data Bank II; Scripps Institution
and the U.S. Geological Survey
Lambert Azimuthal Equal-Area
Center point: 120° E, 15° S
Gridlines: 1 degree interval

BATHYMETRIC CONTOURS IN METERS
DATUM: MEAN SEA LEVEL
PALEOGENE ISOPACHS IN METERS

SOURCES OF DATA

87-495-A Thickness map of the Tertiary sequence of the circum-Borneo region, Southeast Asia.
87-495-B Palinspastic thickness map of the Paleogene sequence of the circum-Borneo region, Southeast Asia.
87-495-C Palinspastic thickness map of the Paleogene sequence of the circum-Borneo region, Southeast Asia.
87-495-D Palinspastic thickness map of the Paleogene sequence of the circum-Borneo region, Southeast Asia.
87-495-E Palinspastic thickness map of the Paleogene sequence of the circum-Borneo region, Southeast Asia.
87-495-F Location map of major Tertiary sedimentary basins and structural elements of the circum-Borneo region, Southeast Asia.

INTRODUCTION

This series of U.S. Geological Survey Open-File Reports, numbered 87-495-A through 87-495-F, constitutes a pilot study in the basic requirements of a basin analysis. The study was designed to provide an example of the rudimentary geologic data required as the first step in all hydrocarbon resource assessment programs.

The pilot study was prepared in cooperation with the U.S. Geological Survey World Energy Resources Program; the International Union of Geological Sciences; the Committee for Coordination of Joint Prospecting for Mineral Resources in Asian Offshore Areas; and the Circum-Pacific Map Project of the Circum-Pacific Council.

The data have been compiled on a 1:2,000,000 scale, Lambert azimuthal equal-area map base, prepared by the U.S. Geological Survey, from World Data Bank II, Scripps Institution of Oceanography, and U.S. Geological Survey sources.

The area chosen for the pilot study is the offshore area of the island of Borneo, situated in Southeast Asia. Borneo is divided politically into three countries: Indonesia, Malaysia, and Brunei. The portion of Indonesian territory that comprises the State of Sarawak and Sabah, and the Sultanate of Brunei is an entity by itself.

All thickness contours, faults, and sedimentary province boundaries are constructed, and are subject to change as modification. The discrepancies in terminology, such as shelf or platform, basin depression or trough, and the use of subunits, result from the compilation of many sources of data. They are reproduced from published reports exactly as the original author described them. These terms are familiar to those concerned with petroleum exploration in this region. The nomenclature and spelling of place names, structures, and other geographic and geologic features vary considerably from publication to publication; however, the phonetic pronunciation is invariably the same. Discrepancies in the terminology and description of geographic names between residents of local towns and villages affect nomenclature. The multilingual petroleum companies engaged in petroleum exploration in Borneo also have employed independent means of translation and spelling.

This text is included in its entirety on all maps for the reader's convenience and information.

GEOLOGIC SETTING

In Southeast Asia, many of the sedimentary basins of a size and volume conducive to hydrocarbon generation and accumulation are located totally or partially in offshore areas. This description fits the study area of thicknesses of Indonesia, Malaysia, and Brunei which constitute the island of Borneo.

Approximately 15 Tertiary depositional and four platform areas are peripheral to or straddle the coastal margin of Borneo. Only one major Tertiary basin complex, that of the Malay and Sulu Seas, is located entirely within the landward portion of Borneo. The Tertiary basin complex referred to as the Borneo Basin, many of the basins situated to the west of Borneo, and the Sulu Sea and Makassar Straits, are separated by lesser structural features. Despite certain similarities, each sedimentary basin complex is geologically distinct.

The geological development of Borneo itself has been the subject of several theories. In general it appears to have formed as a result of the collision of the Australian and Asian continental plates. The collision of the Australian continental plate, rifted from the southern margin of the Asian continent, possibly Indo-China, and India with the Australian plate, probably in the late Mesozoic time and culminated at the end of the Cretaceous. The blocks were probably transported along a complex series of transcurrent or transform faults, associated with or directly related to the collision of the plates. The development of major Tertiary sedimentary basins and stable shelf areas resulted from interaction between the plates. Collision and the subduction of microcontinents, the formation of a wedge and allochthonous wedges, and erosion and redeposition of sediments are envisioned for the development of most of Borneo.

At present, Tertiary clastic and carbonate rocks are the site of all significant hydrocarbon potential in the Asian Basin region. However, they do not preclude future discoveries in Mesozoic and older rocks. The thickness of Tertiary clastics in the study area varies from 1000 to 3000 meters. The maximum thickness of Tertiary clastics in the study area is 3000 meters (10 kilometers). The thickness of Mesozoic rocks varies from 1000 to 2000 meters. The thickness of Paleogene rocks varies from 1000 to 2000 meters.

That part of the Tertiary section referred to as a potentially petroliferous is limited to Tertiary sedimentary rocks, which are not only slightly deformed and moderately indurated. The sand bodies are commonly unconsolidated or extremely friable, whereas the limestone, silt, and clay strata are mostly well indurated. The shale sequences, although friable, are only weakly indurated.

In addition to the petroliferous sedimentary rocks, many thousands of feet of strongly deformed and highly indurated Tertiary rocks exist in Borneo. They are predominantly Palaeozoic.

TECTONIC AND DEPOSITIONAL SETTING

For major depositional basins in the circum-Borneo region, tectonic settings have variously been described as open-shelf cratonic, foreland, rifted pull-apart, spreading center, and continental margin basins. For a simplified classification of continental margin basins is accepted for all but the open-shelf basins along the south and possibly, the north coast of Borneo.

The structure of these continental margin basins is all fault controlled. The basins are formed by faulting along major zones of weakness in the pre-Tertiary basement, parallel to the coastline and generally associated with westward-trending transcurrent faulting. Mid-Miocene volcanics in an amount that ranges from 10 to 20 percent of the total thickness of the Tertiary section. The basins are generally filled with clastic sediments and are separated by lesser structural features. The continental margin collapsed in response to regional subsidence, and generally experienced stresses and to sedimentary loading. As a result of the collapse, a series of normal faults developed in the basin, and the basins were progressively filled by sediments derived from the high-relief margin of the Indonesian archipelago. The prograded thick fluvial, coastal plain, deltaic, and prodelta clastic sediments typically overlie older sequences. Intra-basin basins developed in this block-faulted terrain are ideally suited to the growth of carbonate platforms and pinnacles reefs.

This area substantiates the theory that continental margin basins exhibit highly variable structural styles, which bear strongly on depositional styles. Some of the thickened clastic wedges were deposited over the edge of the continental crust, where major delta systems pooled large volumes of sediments into the deep water of adjacent marginal seas. The continental margin basins are generally strongly asymmetrical, and were commonly deformed on the seaward, actively tectonic side.

The depositional centers of these clastic wedges lie near the boundary between continental and oceanic crust, such that flowage and downsinking of the continental crust occur on the continent side of the depositor. As stated in the literature, where rapid deposition into bathyal environments occurs, such as in the circum-Borneo region, sediment prograde over the continental margin, along down the continental slope, where strata are rapidly deposited and are associated with the continental margin. The continental margin collapsed in response to regional subsidence, and generally experienced stresses and to sedimentary loading. As a result of the collapse, a series of normal faults developed in the basin, and the basins were progressively filled by sediments derived from the high-relief margin of the Indonesian archipelago. The prograded thick fluvial, coastal plain, deltaic, and prodelta clastic sediments typically overlie older sequences. Intra-basin basins developed in this block-faulted terrain are ideally suited to the growth of carbonate platforms and pinnacles reefs.

Where the supply of sediment was abundant, as in the Sarawak, Kutei, and Tareban Basin deltas for example, the deltas have prograded rapidly since early Miocene time, accompanied by differential downwarping and complex faulting. Although the Makassar Delta of the Kutei Basin, Indonesia, contains more sediment by volume, it is probable that the Sarawak Delta of Brunei was the more tectonically active.

PALEOGENEOUS DEVELOPMENT

Paleogene through middle Eocene time was a period of major marine regression in the circum-Borneo region. In response to a period of intense orogenic uplift of a young mountain range along the present position of the Malay Peninsula-Indonesian border. Following this late Eocene to early Miocene time was a period of major marine transgression. Established at this time was an actively subsiding trough in the position of Sarawak, Brunei, and Sabah that was characterized by open marine Elych deposits; the northeast coarctation mountain range along the Malay-Indonesian border; and volcanic igneous conditions in the position of the present-day Sarawak and Tareban Basin. Flysch and rapid basin-fill deposits characterized the Sarawak Basin. "Indurated" basins in the southeast, whereas open marine and tectonic conditions occupied the Indonesian basins southeast of the coarctation range. Also, in late Eocene to early Miocene, a subsidence zone on the northeast coast of Borneo became active; and marine trenches and basins developed offshore northwest and east of Borneo. The basins were rapidly filled as marginal troughs in the northwest, although carbonate platforms with reef growth started to develop locally in response to fault-related horst-and-graben structures. In eastern Kalimantan open-marine conditions existed.


In contrast to Paleogene and early Miocene time, the Eocene time between middle Miocene and early Pliocene was a major period of regression, accompanied by cyclical periods of intense tectonics. This main part of the regression was marked by development of large delta systems, carbonate platforms and pinnacles reefs, shallow marine environments, and basin infill.

Finally, late Miocene to the present has been characterized by cyclical periods of regression and transgression. Renewed tectonic activity, resulting in continental subsidence, during Pliocene time led to the location of thick sedimentary deposits in basins like West Irian (located immediately west of the Balaigang province), Tareban Basin, etc.

EXPLANATION

The complexity of both depositional history and structural development, as illustrated in the circum-Borneo region, points to numerous exploration prospects in a variety of play types. The play types are characterized by a unique combination of stratigraphic setting and structural geometry. The controlling factors in play development are: first, the evolution of the depositional and the nature of sedimentary facies deposited in the play area; and second, the structural configuration or physical geometry of the features receiving the sediments. Structural styles associated with plays include fault-belt basins, structurally controlled deep-seated anticlines, carbonate platforms, pinnacles reef complexes, and fractured basement reservoir occurrences. Another controlling factor and important variables to basin analysis of prospective play types is the age of hydrocarbon traps. Any one basin may contain multiple play types of various ages.

Rapid burial of organic material, as in the Sarawak Delta, has a direct bearing on the ultimate yield of hydrocarbons, by limiting the degree of oxidation of the organic matter. Preferentially derived organic matter is considered to be the primary source for hydrocarbons in the circum-Borneo region.



PALINSPASTIC THICKNESS MAP OF THE PALEOGENE SEQUENCE OF THE CIRCUM-BORNEO REGION, SOUTHEAST ASIA

By
Keith Robinson
1987

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature.