



INTRODUCTION

This series of U.S. Geological Survey Open-File Reports, numbered 87-495-A through 87-495-F, constitutes a pilot study in the basic requirements of a basin analysis. The study was designed to provide an example of the rudimentary geologic data required as the first step in all hydrocarbon resource assessment programs.

The pilot study was prepared in cooperation with the U.S. Geological Survey World Energy Resources Program; the International Union of Geological Sciences; the Committee for Coordination of Joint Prospecting for Mineral Resources in Asian Offshore Areas; and the Circum-Pacific Map Project of the Circum-Pacific Council.

The data have been compiled on a 1:2,000,000 scale, Lambert azimuthal equal-area map base, prepared by the U.S. Geological Survey, from World Data Bank II, Scripps Institute of Oceanography, and U.S. Geological Survey sources.

The area chosen for the pilot study is the offshore area of the island of Borneo, situated in Southeast Asia. Borneo is divided politically into three countries: Indonesia, Malaysia, and Brunei. That portion of Indonesia comprises the State of Kalimantan; that portion of Malaysia comprises the States of Sarawak and Sabah; and the Sultanate of Brunei is an entity by itself.

All thickness contours, faults, and sedimentary province boundaries are conjectural, and are subject to change or modification. The discrepancies in names of various fields, basins, and provinces are given or proposed on the basis of substatin, result from the compilation of many sources of data. They are reproduced from published reports exactly as the original author described them. The names of fields and basins are given as they appear in petroleum exploration literature in this region. The nomenclature and spelling of place names, structures, and other geographic and geologic features vary considerably from publication to publication; however, the phonetic pronunciation is invariably the same. The names of fields and basins are designated in general by the name, section, and number of the well. The names of structures are designated by the name of the residents of local towns and villages affected nomenclature. The multinational petroleum companies engaged in petroleum exploration in Borneo also have employed independent means of translation and spelling.

This text is included in its entirety on all maps for the reader's

In Southeast Asia, many of the sedimentary basins of a size and volume conducive to hydrocarbon generation and accumulation are Tertiary in age and located totally or partially in offshore areas. This description fits the study area of these portions of the countries of Indonesia, Malaysia, and Brunei which constitute the island of Borneo.

Approximately 15 Tertiary depocenters and four platform areas lie peripheral to or straddle the coastal margins of Borneo. Only one major Tertiary basin, the complex, that of the Kuching and Ketungau basins, is located entirely within the interior of Borneo. The Ketungau Basin is frequently referred to as the Mandal Basin. Many of the basins are divided by major basement ridges and can be subdivided into geologic provinces or subbasins separated by lesser structural features. Despite certain similarities, each sedimentary region or subbasin is geologically unique.

The geological development of Borneo itself has been the subject of several theories. In general it appears to have formed as a result of the southward migration, tectonic collision, and subduction of the southern margin of the Asian continent, possibly Indo-China, and impeded in their southern migration by the Indian-Australian continent. The collision of the Asian and Australian continents and the subduction of the latter under the former took place in the Miocene and culminated at the end of the Miocene. The blocks were probably transported along a complex series of transcurrent or transform faults, which were active in the Miocene and Pliocene. The development and spreading, in both the South China Sea and Makassar Straits. The development of major tectonic sedimentary basins and stable shelf areas resulted from the collision of the Asian and Australian continents. The development of microcontinents, the formation of melange and allochthonous wedges, and the erosion and redeposition of sediments are envisioned for the development of most

At present, Tertiary clastic and carbonate rocks are the site of all known hydrocarbon accumulations in the circum-Borneo region. This is, however, does not preclude future discoveries in Mesozoic and older rocks. The thickness of potentially petroliferous Tertiary sedimentary rocks in the circum-Borneo region locally exceeds 30,000 feet (10 kilometers). The thickest sections are usually associated with areas of deltaic sedimentation and active basin subsidence; neighboring, horst-related carbonate platforms and pinnacle reef buildups, however, may also exceed 5,000 feet (1500 meters) in

That part of the Tertiary section referred to as potentially petroliferous is limited to Tertiary sedimentary rocks, which are at most only slightly indurated and moderately deformed. The sand bodies are commonly unconsolidated or extremely friable, whereas the limestone, marl, and chert stringers are mostly well indurated. The shale sequences, although fissile, are only weakly indurated.

In addition to the petroliferous sedimentary rocks, many thousands of feet of strongly deformed and highly indurated Tertiary rocks exist in Borneo. They are predominantly Paleocene in age.

TECTONIC AND DEPOSITIONAL SETTING

For major depocenters in the circum-Borneo region, tectonic settings have variously been described as open-shelf cratonic, foreland, rifted pull-apart, spreading center, and continental margin basins. For this study a simplified classification of continental margin basins is accepted for all but the open-shelf basins along the south and, possibly, the southeast coasts of Borneo.

The structure of these continental margin basins is all fault-controlled. The basins are formed by faulting along major zones of weakness in the crust. The primary faults are generally normal faults associated with west-trending transcurrent faulting. Miocene volcanism is an element that cannot be ignored. It was probably the result of extensional tectonics in the forearc region. The forearc margin collapsed, and the formation of the Northwest Borneo trench, the continental shelf, and the accretionary prism were the result of the forearc margin collapse and the subsequent loading. As a result of the collapse, a series of normal faults dropped the basement successively seaward. The basins and the accretionary prism were formed by the deposition of sediment derived from the high cordilleran ranges on the landward side. The prograded shelf, fluvial, coastal plain, deltaic and glacial facies developed in the forearc margin. The accretionary prism and the high cordilleran ranges on the landward side are ideally suited to the growth of carbonate platforms and reefs.

This area substantiates the theory that continental margin basins exhibit highly variable structural styles, which bear strongly on depositional styles. Some of the thickest clastic wedges were deposited over the edge of the continental crust, where major delta systems poured large volumes of sediment into the deep water of adjacent marginal seas. The continental margin basins are generally strongly asymmetrical, and were commonly deformed on the seaward, actively tectonic side.

The depositional centers of these classic wedges lie near the boundaries between continental and oceanic crust, such that flexuring and subsinking of the oceanic crust occur on the continent side of the depositor. As has been stated in the literature, where rapid deposition into bathyal environments occurs, such as in the circum-rifted basin, sediments prograde onto the oceanic crust, which is still subsiding. In the case of the Taranaki Basin, the transiently faulting oceanic, involving basement beneath the thicker parts of the depositional wedge, had disappeared. Although diapirism cannot be confirmed in the Taranaki Basin, the requisite structural elements for hydrocarbon accumulation are mainly present in the Taranaki Basin. Reef trends developed on horsts, basement highs, or on sedimentary platforms where active sediment supply had ceased, such as in the Central Taranaki Basin.

Where the supply of sediment was abundant, as in the Karam, Kutei, and Tarakan Basin deltas for example, the deltas have prograded rapidly since early Miocene time, accompanied by differential downwarp, slumping, and complex faulting. Although the Mahakam Delta of the Kutei Basin, Indonesia contains more sediment by volume, it is probable that the Karam Delta of Brunei was more tectonically active.

PALEOGEOGRAPHIC DEVELOPMENT

Palaeocene through middle Eocene time was a period of major marine regression in the circum-Borneo region. In response to periodic marine transgression, uplift of a young mountain range along the present position of the Sulu Trench was initiated. The regression was followed by a period of marine transgression. Middle Eocene time was a period of sea level rise. Following this, Late Eocene time was actively subsiding trough in the position of Sarawak, Brunei, and Kalimantan. The subsiding character was maintained through the middle Eocene. Middle Eocene time was a period of marine transgression in the circum-Borneo mountain range along the Malaysian-Indonesian border; an equivalent open marine conditions in the position of the present-day Kutei and Kutai basins. The transgression was followed by a period of marine regression in the Malaysian and Brunei "subduction" basins to the northwest, whereas open marine conditions were maintained in the Sarawak and Kalimantan basins to the northeast. Also, in late Eocene to early Miocene, a subduction zone on the northwest coast of Borneo became active; and marine transgression and basin development were initiated in the Sarawak and Kalimantan basins. The region was filled as marginal troughs in the northwest, although carbonate platforms with shallow water reefs were maintained in the northeast. The transgression was followed by a period of marine regression. In early Miocene, the transgression was followed by a period of marine regression and open-basin structures. In early Miocene, the transgression was followed by a period of marine regression and open-basin structures.

In contrast to Paleogene and early Miocene time, the Neogene time between middle Miocene and early Pliocene was a major period of regression accompanied by cyclical periods of eustatic eustation. This was in part of the Neogene was marked by development of large delta systems, carbonate platforms and pinnacle reefs, shallow marine environments, and basin infill.

Finally, late Pliocene to the present has been characterized by cyclical periods of regression and transgression. Renewed tectonic activity, resulting in considerable subsidence, during Pleistocene time led to the formation of thick sedimentary deposits in basins like West Uluksua (located immediately west of the Balingian province), Tarakan and Kutai.

EXPLORATION PLAYS

The complexity of both depositional history and structural development, as illustrated in the circum-borneo region, points to numerous exploration prospects in a variety of sedimentary types and tectonic settings. The unique combination of stratigraphic setting and structural geology. The controlling factors in play development are, first, the environment of deposition and the nature of sedimentary facies deposited in the play area, and second, the structural configuration or physical geometry of the features receiving the sediment. Structural styles associated with plays include fault-tectonic thrust, graben and half-graben structures, gently folded to highly folded and asymmetrical, and steeply folded to overturned and rollover features, structurally controlled deep-marine turbidites, carbonate platforms, pinnacle reef complexes, and fractured basement reservoirs. The structural complexity of the sedimentary basins and the variety of play types of prospective plays types is the age of hydrocarbon traps. Any one

lay types of var
c material, as

bearing on the volumetric yield of hydrocarbons, by limiting the degree of oxidation of the organic matter. Petrocarystically derived organic matter is considered to be the primary source for hydrocarbons in the circum-Boreale region.

U. S. GEOLOGICAL SURVEY OPEN-FILE REPORTS

87-095-A Thickness map of the petroliferous Tertiary sequence of the circum-Boreale region, Southeast Asia.

87-095-B Palaeopneumatic thickness map of the Palaeogene sequence of the circum-Boreale region, Southeast Asia.

87-095-C Palaeopneumatic paleogeographic map of the Palaeogene sequence of the circum-Boreale region, Southeast Asia.

87-095-D Palaeopneumatic thickness map of the Palaeogene sequence of the circum-Boreale region, Southeast Asia.

87-095-E Palaeopneumatic paleogeographic map of the Mesozoic sequence of the circum-Boreale region, Southeast Asia.

87-495-F Location map of major Tertiary sedimentary provinces and structural elements of the circus-Borneo region, Southeast Asia.

SOURCES OF DATA

Acknowledgements are due to the ASEAN Council on Petroleum (ASCOPE); the Committee for Co-ordination of Joint Prospecting for Mineral Resources in Asian Offshore Areas (CCOP); the national oil corporations of Indonesia (PERTAMINA), Malaysia (PETRONAS), and the Philippines (PNOC); to many published and unpublished papers by the Indonesian Petroleum Association (IPA) and the Geological Society of Malaysia; to the Indonesian Petroleum Association (IPA) and the Geological Society of Malaysia; the University of Malaya Geological Department and other organizations; and to the Geological Surveys of Indonesia and Malaysia, for their assistance and support in providing the necessary data as the basis for these interpretative maps, and to the many oil companies and consultants active in exploration within the area, who are primarily

**By
Keith Robinson
1987**