

This series of U.S. Geological Survey Open-File Reports, numbered 87-495-A through 87-495-F, constitutes a pilot study in the basic requirements of a basin analysis. The study was designed to provide an example of the rudimentary geologic data required as the first step in all hydrocarbon resource assessment programs.

The pilot study was prepared in cooperation with the U.S. Geological Survey World Energy Resources Program; the International Union of Geological Sciences; the Committee for Coordination of Joint Prospecting for Mineral Resources in Asian Offshore Areas; and the Circum-Pacific Map Project of the Circum-Pacific Council.

The data have been compiled on a 1:2,000,000 scale, Lambert azimuthal equal-area map base, prepared by the U.S. Geological Survey, from World Data Bank II, Scripps Institute of Oceanography, and U.S. Geological Survey sources.

The area chosen for the pilot study is the offshore area of the island of Borneo, situated in Southeast Asia. Borneo is divided politically into three countries: Indonesia, Malaysia, and Brunei. That portion of Indonesia comprises the State of Kalimantan; that portion of Malaysia comprises the States of Sarawak and Sabah; and the Sultanate of Brunei is an entity by itself.

All thickness contours, faults, and sedimentary province boundaries are conjectural, and are subject to change or modification. The discrepancies in terminology, such as shelf or platform, basin depression or trough, and the use of subbasin, result from the compilation of many sources of data. They are represented from publications as early as the original maps of the region. These areas are familiar to those concerned with the oil and gas exploration in this region. The nomenclature and spelling of place names, structures, and other geographic and geologic features vary considerably from publication to publication, however, because of the lack of uniformity in the field.

Discrepancies in the terminology and description of geographic names between residents of local towns and villages affect nomenclature. The multinational companies responsible for the oil and gas exploration in the basin in some cases have employed independent means of translation and spelling.

This text is included in its entirety on all maps for the reader's convenience and information.

GEOLOGIC SETTING

In Southeast Asia, many of the sedimentary basins of a size and volume conducive to hydrocarbon generation and accumulation are Tertiary in age and located totally or partially in offshore areas. This description fits the study area of those portions of the countries of Indonesia, Malaysia, and Brunei which constitute the island of Borneo.

Approximately 15 Tertiary depocenters and four platform areas lie peripheral to or straddle the coastal margins of Borneo. Only one major Tertiary basin complex, that of the Melawi and Ketungau Basins, is located entirely within the interior of Borneo. The Ketungau Basin is frequently referred to as the Mandal Basin. Many of the basins are divided by major basement ridges and are separated into geologic provinces or subbasins separated by lesser structural features. Despite certain similarities, each sedimentary region or subbasin is geologically unique.

The geological development of Borneo itself has been the subject of several theories; in general it appears to have formed as a result of the southward migration, rotation, collision, and accretion of series of microcontinents. The first of these was the Indochinese Peninsula, which was possibly Indo-China, and impeded in their southerly migration by the Indian-Australian plate. This migration process originated in the Cretaceous and was transported along a complex series of transformant or transform faults, which were later displaced by the Sulu Islands. The collision of these plates spreading, in both the South China Sea and Makassar Straits. The development of major Tertiary sedimentary basins and stable shelf areas resulted from the collision of the Indochinese and Australian plates. The collision of the microcontinents, the formation of melange and allochthonous wedges, and the erosion and redeposition of sediments are envisioned for the development of most

At present, Tertiary clastic and carbonate rocks are the site of all significant hydrocarbon discoveries in the circum-Borneo region. This, however, does not preclude future discoveries in Mesozoic and older rocks. The thickness of potentially petroliferous Tertiary sedimentary rocks in the circum-Borneo region locally exceeds 30,000 feet (10 kilometers). The thickest sections are usually associated with areas of deltaic sedimentation and active basin subsidence; neighboring, horst-related carbonate platforms and pinnacles build up, however, may also exceed 5,000 feet (1,500 meters) in thickness.

That part of the Tertiary section referred to as potentially petroliferous is limited to Tertiary sedimentary rocks, which are at most only slightly indurated and moderately deformed. The sand bodies are commonly unconsolidated or extremely friable, whereas the limestone, marl, and chert stringers are mostly well indurated. The shale sequences, although fissile, are only weakly indurated.

In addition to the petroliferous sedimentary rocks, many thousands of feet of strongly deformed and highly indurated Tertiary rocks exist in Borneo. They are predominantly Paleogene in age.

TECTONIC AND DEPOSITIONAL SETTING

For major depocenters in the circum-Borneo region, tectonic settings have variously been described as open-shelf cratonic, foreland, rifted pull-apart, spreading center, and continental margin basins. For this study a simplified classification of continental margin basins is accepted for all but the open-shelf basins along the south and, possibly, the southeast coasts of Borneo.

The structure of these continental margin basins is all fundamentally controlled by the tectonic evolution of the continent. The basins are formed by subsiding along a major zone of weakness in the pretertiary basement, parallel to the coastline and generally associated with northwest-trending transcurrent faulting. Mid-Miocene volcanism is an element that cannot be ignored. It has produced a complex pattern of extensional tectonics, Miocene Late tectonics, collisional tectonics, and the formation of the Northwest Borneo trench. The continental margin collapsed in response either to regional subduction or pull-apart processes and to a lesser extent to the effects of the Miocene Late normal faults dropped the basement successively seaward. The basins and grabens that formed on the continental edge are progressively filled by clastic sediments. The basins are filled by a sequence of clastic facies that progressed thick fluvial, coastal plain, deltaic, and prodelta clastics and sediments typically overlap older sequences. Horst blocks developed in this period. The basins are ideally suited to the growth of carbonate platform and petroleum reefs.

This area substantiates the theory that continental margin basins exhibit highly variable structural styles, which bear strongly on depositional styles. Some of the thickest clastic wedges were deposited over the edge of the continental crust, where major delta systems poured large volumes of sediment into the deep water of adjacent marginal seas. The continental margin basins are generally strongly asymmetrical, and were commonly deformed on the seaward, actively tectonic side.

The depositional centers of these clastic wedges lie near the boundary between continental and oceanic crust, much that fluearing and downdrafting of the continental crust occur on the continent side of the depocenter. As stated in the literature, these regions are the depocenters of the clastic wedges. In the Florida region, sediments prograde over the continental shelf edge, slumping down the shelf slopes. Where strike-slip faults are present, the units are truncated and the depocenter is shifted. In the depositional wedge, shale diapirism commonly results, as generally accepted for the Barataria Delta. Although diapirism cannot be confirmed in areas such as the Barataria Delta, it is a possibility in the Florida region. Diapirism is not certainly correct, as is also true in the Mahabank Delta of the Texas Basin. Reef trends developed on barata, basement highs, or on sedimentary basins. The active sediment supply had ceased such as in the Central and southern Louisiana province.

Where the supply of sediment was abundant, as in the Baram, Kutai, and Tarakan Basin deltas for example, the deltas have prograded rapidly since early Miocene time, accompanied by differential downwarp, slumping, and complex faulting. Although the Mahakam Delta of the Kutai Basin, Indonesia contains more sediment by volume, it is probable that the Baram Delta of Brunei was the more tectonically active.

[illegible]

Finally, late Pliocene to the present has been characterized by cyclical periods of regression and transgression. Renewed tectonic activity, resulting in considerable subsidence, during Pleistocene time led to the formation of thick sedimentary deposits in basins like West Mazonia (located immediately west of the Balingian province), Tarakan and Kutai.

EXPLORATION PLAYS

The complexity of both depositional history and structural development, as illustrated in the circumstantial points to numerous exploration prospects in a variety of play options. All play types are characterized by a unique combination of stratigraphic setting and structural geometry. The depositional and structural factors that define the play types are: (1) the deposition and the nature of sedimentary facies deposited in the play area, and second, the structural configuration or physical geometry of the features that define the play type. The play types are: (1) the low-salinity, high-pH, fault-bound horst, graben and half-gaben structures; gently folded to highly faulted and asymmetrical anticlines, unsymmetrical growth fault and rollover structures; (2) the steeply faulted, steeply dipping, high-salinity platforms, pinnacle reef complexes, and fractured basement reservoir structures. Another controlling factor and important variable in basin analysis is the prospective play type.

Rapid burial of organic material, as in the Saram Delta, has a direct bearing on the volumetric yield of hydrocarbons, by limiting the degree of oxidation of the organic matter. Terrestrially derived organic matter is considered to be the primary source for hydrocarbons in the circum-Borneo basin may contain multiple play types of various ages.

U. S. GEOLOGICAL SURVEY OPEN-FILE REPORTS

87-495-A Thickness map of the petroliferous Tertiary sequence of the circum-Borneo region, Southeast Asia.


87-495-B Palinspastic thickness map of the Paleogene sequence of the circum-Borneo region, Southeast Asia.

87-495-C Palinspastic paleogeographic map of the Paleogene sequence of the circum-Borneo region, Southeast Asia.

87-495-D Palinspastic thickness map of the Neogene sequence of the circum-Borneo region, Southeast Asia.

87-495-E Palinspastic paleogeographic map of the Neogene sequence of the circum-Borneo region, Southeast Asia.

87-495-F Location map of major Tertiary sedimentary provinces and structural elements of the circum-Borneo region, Southeast Asia.



50 0 50 100 150
50 0 50 100 150

SCALE 1:2 000 000

BATHYMETRIC CONTOURS IN METERS
DATUM: MEAN SEA LEVEL
NEOGENE ISOPACHS IN METERS

SOURCE: U.S. BUREAU OF ECONOMIC ANALYSIS.

Acknowledgements are due to the ASEAN Council on Petroleum (ASCOPE); the Committee for Coordination of Joint Prospecting for Mineral Resources in Indonesia-Malaysia (COMCRIM); the national oil corporations of Indonesia (PERTAMINA), Malaysia (PETRONAS), and Brunei Darussalam (PETROS); the published and unpublished papers by the Indonesian Petroleum Association (IPA), the Southeast Asia Petroleum Exploration Society (SEAPEX), the Geological Survey of Malaysia, the University of Wales Geological Department, and other organizations; and to the Geological Surveys of Indonesia and Malaysia, for their assistance and support in providing the necessary data as the basis for these interpretative maps, and to the many oil companies and other interested parties in the area, who are primarily responsible for compilation of the original data.

**By
Keith Robinson
1987**

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature.