


Base from U.S. Geological Survey 1:250,000
Nogales, 1956-69 and Tucson, 1956-77

SCALE: 1:250,000
5 0 5 10 MILES
5 0 5 10 KILOMETERS

TOPOGRAPHIC CONTOUR INTERVAL 200 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

Figure 4.—Approximate areal distribution of clay- and silt-size particles in the upper Tinaja beds in Avra Valley.




Base from U.S. Geological Survey 1:250,000
Nogales, 1956-69 and Tucson, 1956-77

SCALE: 1:250,000
5 0 5 10 MILES
5 0 5 10 KILOMETERS

TOPOGRAPHIC CONTOUR INTERVAL 200 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

Figure 5.—Approximate areal distribution of clay- and silt-size particles in the Fort Lowell Formation in Avra Valley.



M(200)
R29o

87-685

sheet 2
c. 1

POTENTIAL FOR AQUIFER COMPACTION, LAND SUBSIDENCE, AND EARTH
FISSURES IN AVRA VALLEY, PIMA AND PINAL COUNTIES, ARIZONA

BY

S. R. ANDERSON

1988

OFR 87-685

MX274/168
Av82a
sheet 2
c. 1

USGS LIBRARY - RESTON VA
3 1818 00653184 0