

UNITED STATES DEPARTMENT OF THE INTERIOR

GEOLOGICAL SURVEY

GEOCHEMISTRY OF LARAMIDE AND MIDDLE TERTIARY PLUTONS

IN THE AJO MINING DISTRICT, PIMA COUNTY, ARIZONA

By Dennis P. Cox

Open-File Report

88- 672

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or stratigraphic nomenclature. Any use of trade names is for descriptive purposes only and does not imply endorsement by the USGS.

Menlo Park, California

1988

## Geochemistry of Laramide and Middle Tertiary Plutons in the Ajo Mining District, Pima County, Arizona

The Ajo Mining District comprises the Ajo porphyry copper deposit and various small vein deposits in the Little Ajo Mountains of western Pima County, Arizona. The New Cornelia Mine, developed on the Ajo deposit by Phelps Dodge Corporation has produced 2.5 million tonnes of copper since it was opened in 1917. Ajo is the westernmost deposit of a large cluster of porphyry Cu-Mo deposits in southern Arizona and, although molybdenum is recovered from parts of the deposit, it most closely matches the porphyry Cu-Au model of Cox (1986).

The geology of the Ajo 15 minute quadrangle and the New Cornelia Mine was described by Gilluly (1946), and of the surrounding region by Gray and others (1988). Precambrian gneiss and granite and felsic volcanic rocks of possible Mesozoic age are intruded by granitic rocks of quartz diorite to monzogranite composition which he called the Cornelia quartz monzonite. Copper mineralization is related to a granodiorite porphyry phase of this pluton.

Analyses of 19 rock samples from the Ajo district are presented in table 1 and 2 and their locations are shown on figure 1. They illustrate the differences and similarities between rocks of the Cornelia pluton (Gilluly, 1946), located around the New Cornelia Mine and on Camelback Mountain, and the pluton of Cardigan Peak, lying to the west of Gibson Arroyo (see Fig. 2). The first has K-Ar ages clustered around 61 Ma and the second has K-Ar and incomplete U-Pb ages of approximately 25 Ma (McDowell, 1980; Hagstrum and others, 1987). The two rock units are similar in appearance and major-element chemistry and were considered by Gilluly (1946) to be parts of the same intrusion.

The rock samples are grouped by age on tables 1 and 2: four samples belong to the Cornelia pluton; one sample of uncertain age and affinity closely resembles quartz diorite of the Cornelia pluton but crops out on the west margin of the Cardigan Peak pluton; 12 samples are from the Cardigan Peak pluton; and two samples represent porphyritic dikes intruding the Cardigan Peak pluton and the Cornelia pluton on Camelback Mountain.

X-ray spectroscopic analyses of major element oxides and Nb were made by A. J. Bartel, Kathleen Stewart, and J. E. Taggart in Lakewood Colorado, and L. F. Espos in Menlo Park, California. Determination of FeO, H<sub>2</sub>O, CO<sub>2</sub>, Cl, and F were made by D. V. Vivit in Menlo Park. Be, Co, Cr, Cu, Ga, Ni, Pb, and V were determined by emission spectrograph by Judith Kent and R. L. Lerner in Menlo Park. Sn, and Nb were determined colorimetrically by J. S. Kane and M. W. Doughten in Reston, Virginia, respectively. Remaining elements in table 2 were determined by instrumental neutron activation analysis by J. Storey, J. R. Budahn, R. J. Knight, S. Danehey, and R. B. Vaughn in Lakewood.

Rocks of the Cardigan Peak pluton are indistinguishable from those of the Cornelia pluton with respect to major elements and most trace elements. The Peacock index (57, see Figure 3) is the same for both plutons; they are calc-alkalic. The Cardigan Peak rocks differ in their higher content of thorium, hafnium, and zirconium (Fig. 4a-c), and rare earth elements (Fig. 4d-f and Fig. 5a-d). In plots of these elements against SiO<sub>2</sub>, sample 516 of quartz diorite (KT?fd) from the undated western margin of the Cardigan Peak pluton consistently falls within the cluster of Cornelia pluton samples. This suggests that this quartz diorite unit is the same age as the quartz diorite (KTfd) on Camelback Mountain to which it bears a close resemblance (Cox and Ohta, 1984).

## References cited

- Cox, D.P. and Ohta, E., 1984, Maps showing rock types, hydrothermal alteration, and distribution of fluid inclusions in the Cornelia pluton, Ajo mining district, Pima County, Arizona: U. S. Geological Survey Open-File Report 84-388, 12p., 3 maps, scale 1:25,000.
- Cox, D.P., 1986, Descriptive model of porphyry Cu-Au in Cox, D. P. and Singer, D. A., Mineral Deposit Models: U. S. Geological Survey Bulletin 1693, p. 110.
- Gilluly, J., 1946, The Ajo Mining district, Arizona: U. S. Geological Survey Professional Paper 209, 112 p.
- Gray, Floyd, Miller, R.J., Grubinsky, Michael, Tosdal, R.M., Haxel, Gordon, Peterson, D.W., May, D.J., and Silver, L.T., 1988, Preliminary geologic map of the Ajo and Lukeville 1° by 2° quadrangles, S. W. Arizona: U. S. Geological Survey Open-File Map 87-147, scale, 1:250,000.
- Hagstrum, J.T., Cox, D.P., and Miller, R.J., 1987, Structural reinterpretation of the Ajo mining district, Pima County, Arizona, based on paleomagnetic and geochronologic studies: Economic Geology, Vol. 82, p. 1348-1361.
- McDowell, F.W., 1971, K-Ar ages of igneous rocks from the western United States: Isochron/West, no. 2, p. 1-2.

Table 1. Major element chemistry of rocks from the Ajo district, Arizona  
[Values in percent]

Sample	Cornelia Pluton (61 Ma)				Unknown age		Cardigan Peak Pluton (25 Ma)						Dike rocks						
	214	540	805	806	516	436	434	264	44	278	289	270	101	252	24	315	311	503	517
SiO2	59.7	64.4	64.5	65.7	61.1	59.6	63.2	66	67.4	70.1	67.1	70.4	71.5	76.7	77.5	73	77.5	64.3	63.1
Al2O3	16.8	16.3	15.6	15.4	16.1	16	16.4	15.9	15.4	14.8	15.5	14.8	14.4	12.6	12.1	14.1	12.2	16	16.2
Fe2O3	3	2.09	2.1	2.29	2.44	2.48	2.02	2.16	1.52	1.18	1.661	1.28	1.42	0.154	0.492	0.304	0.364	2.06	2.45
FeO	2.43	1.71	1.68	1.57	2.03	3.19	1.5	1.23	1.46	0.38	1.25	0.89	0.44	0.05	0.07	0.32	0.05	1.63	1.36
MgO	3.05	1.69	1.6	1.79	3.06	3.85	2.31	1.53	1.47	0.95	1.45	0.79	0.63	0.14	0.17	0.57	0.14	1.81	1.93
CaO	4.8	3.96	3.27	3.43	4.94	5.63	4.65	3.03	2.89	2.58	3	2.07	1.73	0.75	0.36	1.45	0.53	3.45	4.05
Na2O	4.31	4.31	4	3.98	3.88	3.47	3.91	3.78	3.62	3.57	3.63	3.37	3.35	3.28	3.11	2.95	3.38	4.06	3.88
K2O	2.42	2.98	3.42	3.1	2.92	3.18	3.68	3.97	4.14	4.46	4.17	4.65	4.85	4.8	5.23	5.54	4.74	3.52	4.03
TiO2	0.87	0.57	0.54	0.57	0.57	0.79	0.67	0.43	0.44	0.34	0.44	0.28	0.23	0.12	0.1	0.21	0.1	0.51	0.59
P2O5	0.28	0.23	0.2	0.21	0.22	0.38	0.29	0.19	0.18	0.14	0.18	0.1	0.09	<0.05	<0.05	0.07	<0.05	0.23	0.24
MnO	0.09	0.05	0.03	0.06	0.04	0.09	0.03	0.04	0.05	<0.02	0.03	0.03	0.04	<0.02	<0.02	<0.02	<0.02	0.03	0.02
H2O+	1.1	0.59	1.11	0.67	0.93	0.5	0.38	0.46	0.31	0.32	0.39	0.08	0.35	0.19	0.08	0.25	0.44	0.88	0.61
H2O-	0.12	0.12	0.16	0.16	0.27	0.09	0.14	0.2	0.17	0.17	0.16	0.16	0.18	0.08	0.14	0.22	0.03	0.22	0.18
CO2	0.1	0.07	0.64	0.03	0.18	0.03	0.08	0.07	0.04	0.05	0.15	0.11	0.03	0.03	0.03	0.07	0.11	0.08	0.1
Cl	0.02	0.02	0.012	0.02	0.033	0.07	0.05	0.03	0.03	0.03	0.02	0.02	0.03	0.04	0.02	0.02	<0.02	0.034	0.033
F	0.03	0.02	0.04	0.04	0.08	0.07	0.09	0.06	0.06	0.05	0.07	0.05	0.05	<0.02	<0.02	0.04	<0.02	0.05	0.23
Sum	99.12	99.11	98.90	99.02	98.79	99.42	99.40	99.08	99.18	99.12	99.20	99.08	99.32	98.93	99.40	99.11	99.58	98.86	99.00

Table 2. Trace element chemistry of rocks from the Ajo district, Arizona  
[Values in parts per million]

Sample No.	Cornelia Pluton (61 Ma)					Unknown age	Cardigan Peak Pluton (25 Ma)					Dike rocks								
	214	540	805	806	516		436	434	264	44	278	289	270	101	252	24	315	311	503	517
Be	1.9	2	1.8	1.7	2		2.4	3.1	3	3.3	3.6	2.8	2.4	2.1	3.9	2.8	2.5	7.1	2.2	2.4
Co	19	13	9	12	14		16	12	9.3	9.7	4.3	7	5.4	4.3	1.8	3.8	2.9	3.2	12	9.9
Cr	31	19	17	14	110		89	49	22	23	11	15	10	<10	<10	<10	<10	<10	30	39
Cu	32	17	60	7.8	7.1		50	71	12	1.6	1.6	18	6.3	2.1	2.3	7.9	130	1.5	15	6.2
Ga	26	25	20	18	21		21	24	21	22	21	19	20	19	17	19	18	21	19	20
Ni	18	10	9	9.2	46		44	30	17	17	10	13	8.5	6.3	2.5	4.8	5.7	3.2	18	19
Pb	18	18	20	<10	16		14	15	16	23		26	22	35	<10	13	<10	16	18	<10
V	140	100	83	81	84		120	110	66	71	48	48	45	31	19	33	29	29	64	68
Y	16	16	14	13	19		26	25	21	24	19	20	21	19	<10	10	23	12	19	19
Zr	150	99	120	93	240		210	200	160	190	160	140	180	150	53	86	130	65	220	280
Sn	1.5	1.2	1.6	2.8	1.2		1.3	1.9	1.8	2	1.2	1.6	2.3	2.6	1.1	2.1	1.4	2	1.9	1.3
Nb	1	12	12	13	10		13	16	14	15	15	15	12	12	17	15	15	25	15	16
U	1.19	3.72	3.49	3.28	2.12		2.71	3.81	3.66	4.6	4.57	4.89	3.07	6.12	6.08	6.94	6.17	8.19	2.21	2.29
Ba	836	927	973	846	1240		1450	1270	1600	1460	1380	1450	1440	1180	420	512	1080	90	1260	1610
Co	16.5	8.95	7.45	9.53	11		18.9	8.73	7.14	7.27	2.2	6.82	4.03	3.05	0.278	0.826	1.6	0.5	8.97	7.21
Cr	28.3	13.6	12.6	12.3	111		97.3	41	20.5	19.4	8.5	17	8	4.61	0.69	0.25	2.79	0.5	24.8	34.2
Cs	1.83	1.43	2.59	1.47	0.491		0.938	0.666	1.69	1.87	0.704	1.7	2.1	5.11	0.621	0.994	1.07	3.97	2.3	0.548
Hf	4.06	3.96	3.65	3.64	4.9		6.24	8.07	6.69	7.16	5.92	7.09	5.04	4.47	3.15	3.93	5	4.44	5.03	6.23
Rb	58.8	77.9	94.7	73.8	49.9		81	81.3	111	128	99.3	125	164	174	92.9	136	149	195	88.5	76.8
Sb	0.302	0.108	1.66	0.232	0.151		0.133	0.105	0.184	0.159	0.1	0.1	0.107	0.102	0.181	0.318	0.083	0.804	0.722	0.211
Sr	790	680	560	620	920		1040	903	723	669	603	660	385	344	186	105	270	<75	710	880
Ta	0.426	0.751	0.826	0.82	0.553		0.793	0.983	0.977	1.09	1.17	1.04	0.995	1.02	1.5	1.89	1.38	2.31	0.729	0.797
Th	3.68	7.2	7.99	8.43	6.42		11.3	16.3	17.1	19.8	27.8	19.7	19.3	25.4	29.9	40	24.8	45.4	10	10.7
Zn	59.1	27.2	28.5	26.8	17.7		54	56	<50	<50	<50	<50	<50	<50	<50	<50	<50	<50	<50	<50
Zr	139	136	124	124	172		241	321	252	259	202	265	180	170	89	110	151	83.2	198	267
Sc	10.7	6.67	6.29	6.49	9.79		13.9	8.54	5.99	5.56	3.5	5.55	3.36	2.32	0.547	0.423	2.89	0.738	6.67	7.29
La	28.6	30.9	27.7	29.5	43.4		58.2	52.6	60	59.5	60.9	64.2	50.8	45.8	9.85	21.3	32.2	26.4	44.9	53.4
Ce	57.4	61.6	56.2	59.3	73		120	120	110	111	110	116	89.4	79.5	17.5	36.2	58.2	40.7	81.6	95.5
Nd	24.1	23.7	22.7	24.7	26.3		49.9	47.3	34.9	37.2	32.8	39.5	29.4	25.5	8.98	11.5	20.5	10.5	32.1	32.9
Sm	4.52	4.62	4.14	4.24	4.58		8.01	7.07	5.25	5.53	4.78	5.58	4.19	3.83	1.21	1.7	3.94	1.27	4.66	5.39
Eu	1.36	1.2	1.05	1.12	1.14		1.85	1.67	1.17	1.2	1	1.02	0.873	0.759	0.344	0.31	0.726	0.141	1.24	1.23
Gd	3.51	3.24	2.98	2.98	3.67		6.6	5.69	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	3.75	3.59
Tb	0.492	0.445	0.401	0.452	0.482		0.824	0.765	0.547	0.575	0.466	0.561	0.483	0.415	0.109	0.132	0.512	0.138	0.508	0.528
Dy	<0.1	<0.1	<0.1	<0.1	<0.1		5.1	4.4	3.3	3.4	2.9	3.5	2.9	2.5	0.7	0.9	3.5	1	<0.1	<0.1
Tm	0.197	0.188	0.19	0.203	0.232		0.358	<0.5	0.26	0.29	0.265	0.27	0.255	0.241	<0.5	<0.5	0.37	<0.5	0.239	0.229
Yb	1.06	1.17	1.09	1.15	1.44		2.12	2.13	1.67	1.81	1.66	1.74	1.58	1.55	0.546	0.561	2.38	1.2	1.44	1.51
Lu	0.158	0.179	0.167	0.178	0.221		0.301	0.29	0.24	0.27	0.24	0.26	0.234	0.225	0.09	0.088	0.35	0.21	0.214	0.23




Figure 1, Locations of analyzed rock samples in the Ajo district, Arizona  
Base from U.S. Geological Survey, Ajo, 15 minute Quadrangle,  
scale 1:62,500, 1963




Figure 2, Geologic map of the Ajo district, Arizona modified from Cox and Ohta (1984)

## Explanation for Figure 2


<div>Qal</div>	Alluvium and mine waste, undifferentiated, Quaternary	
<div>Thp</div>	Hospital porphyry	Tertiary
<div>Ts</div>	Sneed Andesite	
<div>Ta</div>	Ajo Volcanics	
<div>Tlf</div>	Locomotive Fanglomerate	
<div>pd</div>	Feldspar porphyry and hornblende andesite dikes	
<div>ap</div>	Andesite porphyry dikes	
<div>Tmp</div> <div>Tfg</div> <div>Tmo</div> <div>Tm</div> <div>Tmd</div>	Pluton of Cardigan Peak: Monzogranite porphyry, Tmp; fine-grained granite, Tfg; monzogranite, Tm, and monzogranite with oikocrystic quartz and K-feldspar, Tmo; and quartz monzodiorite, Tmd	
<div>KT?fd</div>	Fine-grained diorite and quartz diorite, age uncertain	
<div>KTgp</div> <div>KTg</div> <div>KTmd</div> <div>KTfd</div>	Cornelia pluton: Granodiorite porphyry, KTgp; granodiorite, KTg; quartz monzodioite, KTmd; and fine grained diorite and quartz diorite, KTfd	Late Cretaceous and early Tertiary
<div>Kc</div>	Concentrator Volcanics, Cretaceous (?)	
<div>pCc</div>	Cardigan gneiss and Chico Shuni Quartz Monzonite, Proterozoic	

 60

Strike and dip of beds

 40

Strike and dip of monzogranite porphyry dikes



Geologic contact, dashed where uncertain



Fault, dashed where uncertain




Figure 3. Weight percent CaO and Na2O+K2O plotted against SiO2 for all pluton samples in the Ajo mining district, Arizona. Peacock indices for calc-alkalic rocks fall between vertical lines.






Figure 4a-c. Thorium, zirconium, and hafnium (in parts per million, ppm) plotted against weight percent SiO<sub>2</sub>. Closed symbols represent samples from Cardigan peak pluton; open symbols, Cornelia pluton






Figure 4d-f. Lanthanum, cerium, and yttrium (ppm) plotted against weight percent SiO<sub>2</sub>. Closed symbols represent samples from Cardigan peak pluton; open symbols, Cornelia pluton




Figure 5a. Chondrite-normalized rare-earth element abundances for 4 samples from the Cornelia pluton.

1 516

AJO, Q DIORITE, FAR WEST




Figure 5b. Chondrite-normalized rare-earth element abundances for sample 516 from the western margin of the Cardigan Peak pluton.




Figure 5c. Chondrite-normalized rare-earth element abundances for 12 samples from the Cardigan Peak pluton.




Figure 5d. Chondrite-normalized rare-earth element abundances for 2 dike rocks intruding the Cornelia pluton and Cardigan Peak pluton.