

(200)
R290
W. 91-353

U.S. DEPARTMENT OF THE INTERIOR

U.S. GEOLOGICAL SURVEY

GLOBAL GEOSCIENCE TRANSECT 8

QUEBEC-MAINE-GULF OF MAINE TRANSECT, SOUTHEASTERN CANADA,
NORTHEASTERN UNITED STATES OF AMERICA

by


D.B. STEWART¹, B.E. WRIGHT¹, J.D. UNGER¹, J.D. PHILLIPS¹, AND D.R. HUTCHINSON¹

PRINCIPAL COMPILERS

with contributions by:

J.H. LUETGERT¹, W.A. BOTHNER², K.D. KLITGORD¹, L.M. LIBERTY¹, AND C. SPENCER³

Open-File Report 91-353



Prepared in cooperation with Geological Survey of Canada and
Geological Survey of Maine

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards (or with the North American Stratigraphic Code). Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

¹U.S. Geological Survey; ²University of New Hampshire; ³Geological Survey of Canada

INTRODUCTION

Transects produced by the North American Continent-Ocean Transect Program (Speed and others, 1982; U.S. Geodynamics Committee, 1989) describe the complex orogen formed along the Late Proterozoic to Early Cambrian rifted margin of the Laurentian craton, including continental rocks of the Grenville Province, which themselves had been earlier formed by a Middle Proterozoic continent-continent collision. Continental accretion was followed by continental separation with the formation of a passive continental margin. This transect, about 900 km in length, crosses the entire Appalachian orogen and the passive continental margin from the craton to Atlantic oceanic crust. It is supported by abundant seismic reflection and refraction data that were gathered specifically for this transect, in part to resolve questions posed by the earlier transects across the Atlantic passive continental margin. Thus, it is a 'second generation' transect compared to those in the North American Continent-Ocean Transect Program. This transect has been accepted in the Global Geoscience Transects Project (CC-7) of the Inter-Union Commission on the Lithosphere, International Council of Scientific Unions (Monger, 1986) and when published will be called transect number 8.

The Quebec-Maine-Gulf of Maine transect provides excellent data which can be used for understanding the processes of continental accretion and separation. Accretion is shown by the uncertain number, possibly ten or more, of tectonostratigraphic terranes of predominantly continental affinity that make up the part of the orogen shown. In addition, at least two oceanic terranes also have been accreted. In general, these terranes were accreted episodically to the southeastern (current direction) part of the craton by thrust or strike-slip faulting, or both, at successively younger times during the Paleozoic, although some terranes had been joined together to form composite terranes before accretion (Boone and Boudette, 1989, describe an example). The terranes differ principally in the nature of their Late Proterozoic to Middle Paleozoic history and stratigraphy (Keppie, 1989) and paleontology (Neuman and others, 1989). Accretion in the part of the orogen shown on this transect took place during multiple collisional or transpressive episodes in the early and middle Paleozoic.

A very thick continental crust was produced during these episodes as shown by geologic evidence that includes Early Devonian garnet rhyolites extruded after fractional melting of felsic rocks at depths of greater than 50 km (Stewart, 1989), a hinge zone (Hatch and others, 1983) between a thin and a very thick (~10 km) sedimentary prism deposited in a successor basin formed after the mid-Ordovician (Taconian) collisional episode (the Central Maine synclinorium), and kyanite-bearing assemblages indicating the burial of even the youngest of these sediments to mid-crustal depths. Direct geophysical evidence today indicates that thick continental crust remaining from Paleozoic accretion is lacking. However, strong seismic reflectors at the base of allochthonous sheets can be traced to depths of ~25 km, and 10 km or more of allochthonous sheets may initially have been present but since have been eroded. It is a reasonable inference that the underlying continental crust must have been depressed by such thick tectonic loading. However, most stratigraphic evidence for a foreland basin in the Ordovician has been buried under later thrust sheets, and evidence for a foreland basin related to the Acadian orogeny is lacking in the region of this transect.

The present Appalachian crust was shaped principally by late Paleozoic to Early Jurassic extension as shown by abundant seismic reflection and refraction data. This extension led to rupture of continental crust and formation of the present Atlantic Ocean, which was followed by deposition of sedimentary rocks upon the passive continental margin. Extension began with the formation of Carboniferous transtensional rift basins, and continued with the development of a 400 km-wide province of rift basins of Triassic to Early Jurassic age that was accompanied across a wider region by the intrusion of numerous dike swarms and widespread normal faulting. Extension reversed the throw on many inclined Paleozoic thrust faults. The new seismic data suggest that during extension the lower crust and Moho concurrently acquired their principal present characteristics, such as shape and internal features, by ductile shear and recrystallization. Many of the faults bounding rift basins become listric into the top of the lower crust (Stewart and Luetgert, 1990), in some places paired with a rise of Moho by as much as 4 km over a distance of 15 km. Numerous but discontinuous subhorizontal reflectors in the lower crust indicate that during extension laminar packets of contrasting lithologies were formed, and simultaneously through-going older faults were obliterated. The strong seismic reflections from the allochthons in the upper crust die out at depths of 20 to 25 km into the uppermost newly reformed lower crust. The top of the lower crust is a gradational zone as much as 4 km thick where seismic velocity increases about 0.4 km/sec to 6.8 km/sec; such a large increase in velocity must correspond to a decrease in contents of quartz and potassium feldspar. The extended region has newly reformed lower crust and a well-defined reflection and refraction Moho, which contrasts with the indistinct Moho beneath Grenvillian crust to the northwest.

The Quebec-Maine-Gulf of Maine transect demonstrates that the record of Paleozoic orogenies in this part of the orogen is preserved almost exclusively in the presently brittle upper crust. Paleozoic terrane boundaries can only be identified as deep as the top of lower crust, which appears to pass smoothly beneath all terranes. Any genetic relationship of the lower crust to the overlying crust becomes increasingly uncertain as the amount of extension increases progressively southeastward toward the locus of oceanic rupture. The lower crust was attenuated and extended southeastward from its original position as it was ductilely deformed. The high proportion of the upper crust that now consists of exposed granitic rocks formed by partial melting of lower crust in the Paleozoic indicates that most of the lower crust was partially melted then, given any reasonable figure for melt extracted. The bulk composition of the lower crust must have been changed toward lower contents of quartz and alkali-feldspar by the gravitational ascent of magmas. For these reasons we have shown the lower crust as a discrete geologic entity on our interpretive cross section.

The transect graphic is a hard copy guide to the data in an underlying digital geographic information system where all map units shown and their boundaries are assigned attributes according to a scheme devised by Wright and Stewart (1990). More geologic data are preserved in the geographic information system than can be shown on maps at a scale as small as 1:1,000,000. Examples include the specific names and symbolic abbreviations of stratified formations and their members as well as their lithologies, greater detail about intrusive rock names, compositions, and mineralogies, and more complete details concerning age assignments and their ranges, if any. Another attribute assigned in the data base is the tectonic setting of

original deposition or igneous intrusion. We have used these data to display a different kind of map, tectonic setting, which affords a geologic map that conforms to the interpretive cross section. Digital optical discs (CD-ROM) will be used to archive the different geological and geophysical data sets used to prepare this transect. The geologic data archived will include all the digitized detail from the original 1:250,000 and 1:500,000 scale maps (Globensky, 1987; Klitgord and others, 1988, plate 2C; Osberg and others, 1985; St. Julien and Slivitsky, 1985; Weed and others, 1974). The complete data archived for the seismic reflection profiles will include reflection intensities, and for the refraction data will include data for lines and fans not shown because they were parallel to regional strike. Many other new kinds of combinations of geologic and geophysical data sets can be performed using digital data. Some examples given by Phillips (1990) include a color shaded-relief aeromagnetic map with superimposed geologic contacts and faults from the geologic map, and color composites of different wavelenghts of bandpass-filtered aeromagnetic or gravity maps (or combinations of both kinds of potential-field data) with superimposed geologic contacts and faults. A three-dimensional model of parts of this transect in western Maine has been prepared also (Unger and others, 1989).

ACKNOWLEDGEMENTS

This transect required the efforts of a large team of geologists, geophysicists, and cartographers in the United States, Canada, Maine, New Hampshire, and Quebec Geological Surveys and numerous universities starting in 1983. Only the principal authors of the transect graphic and pamphlet are listed as compilers. This group was substantially augmented for specific tasks by the scientists listed as contributors. Many of the contributors have or will publish their work in greater detail elsewhere; most published work is cited in this report. Most field data were collected in 1983-1985, though some geophysical data were collected as recently as 1988. In addition, we obtained important data, advice or other valuable assistance from W.A. Anderson, M.E. Loiselle, and R.G. Marvinney of the Maine Geological Survey; P. Charlesworth, J. Glynn, A. Green, P. Morel-a-l'Huissier, W.H. Poole, and M.D. Thomas of the Geological Survey of Canada; G.M. Boone of Syracuse University; R.A. Phinney, and C. Bottcher of Princeton University; W.E. Doll of Colby College; S.L. Klemperer, then at Cornell University; C.V. Guidotti and P.H. Osberg of the University of Maine-Orono; A.M. Hussey II of Bowdoin College; C.E. Mann of Stanford University; and R.A. Ayuso, Richard Goldsmith, and A.M. Trehu of the U.S. Geological Survey. Joshua Comenetz and Sara Minnich of the U.S. Geological Survey helped to prepare the transect graphic.

SOURCES OF DATA USED

Extensive new compilations of regional magnetic and gravity data on land and at sea were recently prepared for much of the transect region (Macnab and others, 1990; Hutchinson and others, unpublished data). Included were gravity and magnetic data collected by transect participants in support of this transect. Recently published geologic maps for Maine (Osberg and others, 1985) and southeastern Quebec (St. Julien and Slivitsky, 1985; Globensky, 1987) were used. Coverage of each of these primary sources is shown on the transect graphic on small-scale index maps. The regional compilations were prepared at scales larger than the transect, mostly at 1:500,000 or 1:250,000,

so only a minor part of each regional compilation was used. Digitizing these many regional data sets was one of the most time-consuming aspects of transect preparation. All data were converted to the map projection for the base used for this transect, the Lambert Conformal Conic having standard parallels at 33°N. and 45°N. and a central meridian at 69°W. Gravity and magnetic data were incorporated in gridded form.

Most of the geologic data were digitized by machine-scanning of stable-base separates of the maps. Attribute codes were then manually attached to these data after editing and conversion to line-graph data. Details and the codes are given in Wright and Stewart (1990). The minimum size of a geologic object shown on the transect graphic is approximately 1 mm by 1 mm, or 1 km² at 1:1,000,000. For clarity, some similar-aged formations and all members of formations had to be combined on the geologic map within a polygon with one color. A few places, volcanic units within formations were generalized into map units large enough to display overprinted patterns. Although some of the detail shown on the source maps was lost to make the generalized transect maps legible, all of the attributes of the individual geologic map units were preserved in the geographic information system. Geologic point data, such as strikes and dips, could not be shown at map scale and were not digitized. Other geologic point data, such as drill holes at sea and sites sampled by the submersible ALVIN (Hermes and others, 1978), were digitized manually.

The geologic data sets from the original sources were modified by the transect compilers in many places so as to make geologic units continuous across political boundaries, to accommodate new geophysical, geochronologic, and geologic mapping data, and to reflect new understanding of regional geologic processes deduced from the transect effort. Surficial and glacial deposits are not shown although all the land area shown was glaciated in the Quaternary. Coastal and south-central Maine were isostatically depressed by ice loading in excess of 100 m relative to present sea level, so surficial deposits of marine character are found as far inland as Skowhegan. Widespread surficial and glacial deposits include sand, gravel, marine clay and till (Thompson and Borns, 1985); together these reach 100 meters in thickness locally.

Seismic reflection profiles and refraction lines, fans, and related shot points shown on the gravity map are the loci of vertical planes of seismic data that traverse the entire crust. Only those vertical planes that lie close to the line of the interpretive cross section are shown on the transect graphic. Data for approximately 1000 km of deep (15 sec) seismic reflection lines and nearly twice again as many kilometers of deep seismic refraction data were available; nearly all were collected specifically to support this transect (Stewart and others, 1986; Klemperer and Luetgert, 1987; Unger, Stewart, and Phillips, 1987; Luetgert, Mann, and Klemperer, 1987; Hutchinson and others, 1988; Spencer and others, 1989; Luetgert and Mann, 1990; and papers in preparation).

DISPLAY OF GEOPHYSICAL DATA AND ITS PROCESSING

The transect graphic displays several kinds of geophysical data as maps or vertical profiles through the crust at a common scale and with no vertical exaggeration. The coverage for potential-field data is complete for the transect and overall is of high quality. Aeromagnetic and gravity profiles for the line of cross section are displayed to facilitate direct comparison with each other and with geologic structures on the interpretive cross

section. Migrated normal-incidence seismic reflection data are displayed as projected as much as 10 km onto the trace of the composite seismic reflection profile, except for line 3B, which is projected as much as 30 km along the regional strike. All of the available data on land appear on the composite profile. Only the marine data for USGS lines 1 and 1A (Hutchinson and others, 1988), USGS line Gyre 85-11-8 (Hutchinson, unpublished data, 1988), and USGS line 19 (Klitgord and others, 1982) were used. Many other deep seismic reflection profiles for Georges Bank show that line 19 is typical for this region (Schlee and Klitgord, 1988). Seismic refraction data are displayed from a 300-km-long profile on land nearly normal to regional strike (Spencer and others, 1989; Luetgert, unpublished data). This profile has been extended seaward because, according to J.H. Luetgert (written communication, 1990), travel times computed for the coastal part of the profile match well with travel times collected at Matinicus Rock during the 1961 Maine cooperative onshore-offshore experiment in crustal seismology. This experiment was the first fixed array-moving shot point study (Meyer and others, 1962; Steinhart and others, 1962a; Steinhart and others, 1962b; Nakamura and Howell, 1964; Suzuki, 1965). As only preliminary velocity models are available for the marine ocean-bottom seismometer profiles (Sattel and Trehu, 1989), and the results contain large uncertainty in the lower crustal thicknesses and velocities, we have displayed only the locations of these profiles. Velocity-depth profiles of along-strike refraction profiles are plotted on the seismic reflection profile at the point where the along-strike refraction profile crossed the section. Prominent wide-angle reflections from seismic refraction fans (shots fired broadside into refraction deployments) are also shown on the seismic reflection profile where the line connecting midpoints of the refraction fan profile intersected the reflection profile.

The limited available heat flow data are mostly in Maine (Decker, 1987) but have not been plotted on the graphic. They indicate that a normal reduced flux characterizes all of the region. This implies that the lithosphere beneath Maine has equilibrated since it was perturbed by Paleozoic and early Mesozoic orogenic epochs. Decker concluded from his data that there are not significant differences in heat flow between terranes having different basements, and that extension beneath Maine was brief or did not warm large vertical thicknesses of the lithosphere.

Gravity values were corrected for terrain effects onshore (density 2.67 g/cc), earth-tide and instrument drift, and are referenced to the international gravity standardization net of 1971 (Morelli, 1974). Anomaly values were calculated using the 1967 formula of the Geodetic Reference System (International Association of Geodesy, 1971). Complete Bouguer anomalies were used onshore, and free-air anomalies were used offshore. The two types of anomalies are equivalent at the coastline. The data set was gridded at 0.8 km intervals and contoured at 10 milligal intervals. Estimated error is ± 2 milligals. Nearly 2200 new gravity measurements to support this transect were made in 1983 and 1984 along the lines surveyed and vibrated during seismic reflection data collection programs (Snyder and Phillips, 1986; Phillips and others, 1988) and density and magnetic susceptibility measurements were performed on about 150 rock samples collected near measured gravity stations. These closely spaced data insure that the gravity profile shown for the cross section is of very high quality. An intensive collaborative revision of gravity data offshore was performed by colleagues from the Bedford Research Institute of the Geological Survey of Canada, the USGS Marine Geology Branch

at Woods Hole, MA, and the Woods Hole Oceanographic Institution (Hutchinson and others, unpublished data). Marine gravity line spacing was ~14 km in the Gulf of Maine and ~20 km for Georges Bank. The offshore data were initially gridded at 7.5 km. Aeromagnetic data for individual surveys were projected and gridded at an 0.8 km interval. If necessary, the data were continued to a surface 1000 feet above the terrain, and regional trends were removed. The grids for individual surveys were smoothly merged into a master grid, which was machine contoured at 100 nanotesla intervals. Estimated error is ± 25 nanoteslas.

The seismic refraction data on land were obtained in 1984 by the U.S. Geological Survey and the Geological Survey of Canada using arrays of 200 2-Hz seismometers at approximately 800 m spacing having an estimated maximum survey location error of 25 m. Shot points were spaced 20-45 km apart using 900-1800 kg of chemical explosive shot mostly in drilled 90 m holes (Murphy and Luetgert, 1986, 1987). Shallow (<10 km) seismic wave velocities were computed from refraction first-arrival times and amplitudes using two-dimensional ray tracing techniques and synthetic seismograms (Luetgert, 1988a, 1988b, 1988c; Klemperer and Luetgert, 1987). Wide-angle reflection data from mid- to deep-crustal reflectors present in the records were interpreted from normal move-out corrected data (Klemperer and Luetgert, 1987; Luetgert and others, 1987). Detailed explanations of data processing and interpretations of results for the cross-strike lines are given in Spencer and others (1989) and Luetgert and others (1987). Similar details and results for the along-strike lines are given by Klemperer and Luetgert (1987), Luetgert and Mann (1990), and Hennet and others (in press). Observed arrival times fit computed arrival times to within 0.1 sec, so that the uncertainty on any velocity determination is $\sim \pm 0.1$ km/sec. The depth uncertainty of features in the upper crust based upon agreement within 0.1 sec between predicted and observed travel times is probably approximately ± 0.8 km, increasing to ± 2 km at Moho depths. Seismic anisotropy is shown by the differences in velocities at the same site, commonly with slower across-strike and higher along-strike velocities. At some sites, however, the differences are small and could have arisen from different approaches or methods used for data processing. At other sites, the differences exceed the uncertainties of the two compared velocities. It might seem that the site near 325 km on the cross section where the two longest refraction profiles cross would be the best point for comparison. In fact, that point was a common shot point for two different seismometer deployments to the northwest and southeast, respectively, so there are not good data to define the cross-strike velocity at this point. Seismic anisotropy having faster velocities parallel to regional strike also has been observed by Kafka and Reiter (1987) for the group velocity dispersion of short-period Rayleigh waves generated by the same shot points. Kafka and Ebel (1988) also obtained similar results using short-period surface-waves and body-waves generated by the same explosions.

Seismic reflection data were gathered with vibrators on land and by air guns at sea. Recording was for 15 sec for onshore lines 1, 2, 3A, 3B, and 4 and offshore lines 1 and 1A; fold was normally 133 onshore and 30 offshore. Onshore line MERQ was recorded to 4 sec for the northwesternmost 58 km, and to 6 sec for the southeastern 107 km. Marine line Gyre-85-11-8 (24 fold) was recorded to 6 sec, and line 19 (36 fold) was recorded to 12 sec. The variety of problems solved during processing of seismic reflection data are described by Hutchinson and others (1988), Spencer and others (1989), and Hutchinson and

Lee (1989). Velocities for stacking seismic reflection data were obtained by trial velocity stacks at numerous intervals along the profiles and from the seismic refraction studies.

A plot of the compressive wave velocity versus depth for all of the velocity profiles derived from the refraction experiments in Maine by Luetgert (in Unger and others, 1987) showed that the crustal velocities for the region could be reasonably modeled by a linearly increasing function. Digitized line drawings of the principal reflectors on all profiles except line 19 were migrated using a ray-tracing migration program (Unger, 1988a) using the function $V = V_0 + kz$ (where V_0 is the initial surface velocity, taken as 5.98 km sec^{-1} ; k is the velocity gradient, taken as 0.027 sec^{-1} ; and z is the depth in km). Line 19 at the southeastern end of the transect, where thick Mesozoic and Tertiary strata overlie older rocks, could not be migrated using a single linear velocity function, so the digitized reflection section was divided into three sections. Within each section a different linear velocity model, which fit the observed velocity data, was used for migrating the reflectors. The Moho reflectors were migrated with the function $V = 5.75 \text{ km sec}^{-1} + 0.001 \text{ sec}^{-1}z$, and the reflectors within 0 to 5 sec two-way travel time out to the edge of the continental shelf were migrated with the function $V = 2.8 \text{ km sec}^{-1}z + 0.3 \text{ sec}^{-1}z$. The reflectors outboard of the edge of the continental shelf also required use of the function $V = 1.0 \text{ km sec}^{-1} + 0.5 \text{ sec}^{-1}z$ in order to fit the observed velocity profiles and to account for the increased water depth.

MAJOR TECTONOSTRATIGRAPHIC TERRANES CROSSED BY THE TRANSECT

This transect from craton to Atlantic oceanic crust crosses the six major tectonostratigraphic terranes of this part of the Appalachian orogen. The colored map shown on the lower right corner of the transect graphic displays and labels these major terranes and thirteen of their principal components. Different shades of each of the principal colors form belts on this map and show that most of the terranes are composite. This means that rocks with significantly different tectonic settings of origin or early history can be identified within each composite terrane. A careful effort has been made to show the bounding faults between the major tectonostratigraphic terranes on the transect maps and sections. Red ticks along both long edges of all the transect maps mark these faults, and the same red ticks are used along the top of the sections where the bounding faults reach the surface. A tectonostratigraphic evolution diagram gives our interpretation of the time-space history of the different terranes. A description of each of the major tectonostratigraphic terranes from craton to ocean is given below.

Laurentian craton

The basement of the Laurentian craton within the transect area was formed by the late Middle Proterozoic Grenville continent-continent collision. The many types of crystalline rocks exposed at the surface include high grade gneisses, migmatites, and deformed intrusive rocks of a wide range of compositions and ages (Moore and others, 1986; Rankin and others, 1989). The craton was extended in Late Proterozoic time. This rifting formed graben that were filled by Late Proterozoic sediments and bimodal volcanic rocks (Rankin and others, 1989). Following deposition of this sequence, platformal and slope-rise sediments were transgressively deposited upon it from Early Cambrian to Early Ordovician time. The tectonic setting and geologic maps and cross sections for this transect show the craton with unconformably overlying

platformal rocks of Cambrian and Ordovician age. Compression toward the craton during the late Early to Middle Ordovician Taconian orogeny successively thrust the platformal sediments onto the craton along faults such as the Aston and Logan's Line. Continued thrusting imbricated the slope-rise sediments, brought the external nappes by gravitation sliding, and then the internal nappes onto the craton, successively reaching rocks deposited further outboard and cutting deeper through the slope-rise prism to bring metamorphosed early rift-stage volcanics, foreland basin deposits and even slices of the Grenvillian basement itself back onto the craton (St.-Julien and Hubert, 1975; St.-Julien and others, 1983). Crustal processes involved in the Taconian orogeny are well summarized in Stanley and Ratcliffe (1985) and are in good agreement with the interpretation of the MERQ 2001 seismic reflection profile by St.-Julien and others (1983) and Ando and others (1983). We have used these syntheses practically unchanged for the interpretation of the crust at depths less than 10 km shown on the transect cross section except near the southeast end of their profile where new seismic data are more complete (Spencer and others, 1989).

Grenvillian basement continues at depth far to the southeast of its surface outcrop. Seismic reflection profiles (Spencer and others, 1989) show a strong reflector at the top of Grenvillian rocks for a distance of about 175 km southeast of autochthonous outcrops. This is interpreted as mylonite along a decollement possibly initiated along a Late Proterozoic rift fault reactivated in the Taconian orogeny and again reactivated in the Acadian orogeny. Southwest of the transect area, Trzcienski and Marchildon (1989) described Grenvillian xenoliths from dikes that straddle the Guadeloupe fault, more than 100 km southeast of exposed Grenvillian rocks. Our interpretive cross section indicates that, at a structurally equivalent site to that for Trzcienski and Marchildon's samples in this transect, Grenvillian rocks are involved in imbricate crystalline thrusts; autochthonous Grenvillian crust occurs at a depth of about 15 km. The shape of the strong reflector at the top of Grenvillian crust beneath the Chain Lakes block has been determined in three-dimensions from seismic reflection data (Unger, 1988b); it dips southeasterly at approximately 13° , and it can be traced to depths of ~25 km southeast of the Chain Lakes block. Beyond this point, the identity of the lower crust as Grenvillian becomes more speculative and its southeastern limit is conjectural due to deep crustal modification during and after the Paleozoic orogenies. This topic is discussed further below.

Cameron-Brompton (Boundary Mountains) composite terrane

The Cameron-Brompton (Boundary Mountains) composite tectonostratigraphic terrane (Zen, 1989; Boone and Boudette, 1989) occurs southeast of the Brompton part of the Baie Verte-Brompton line of Williams and St.-Julien (1982). These authors interpreted the Brompton line as the suture between the Laurentian craton and accreted terranes unrelated to Laurentia. Terranes included in the composite terrane consist of rocks deposited in oceanic and island-arc tectonic settings, and a highly metamorphosed diamictite or sedimentary melange of granodioritic or feldspathic graywacke bulk composition. The latter, known as the Chain Lakes massif or block, was formerly thought (Naylor and others, 1973) to be about 1500 Ma old, significantly older than Grenvillian crust. However, on the basis of Pb-U dating of single zircons from Chain Lakes rocks, Dunning and Cousineau (1990; and in press), Olszewski and others (1990), and Cousineau (1991) have demonstrated that the Chain Lakes

rocks are actually younger than Grenvillian rocks, and are possibly as young as Early Cambrian. The Chain Lakes block could have originated as a continental margin deposit or as a sedimentary melange from a continental terrane. The continental terrane could have been the Laurentian craton to the northwest (Dunning and Cousineau, in press), a rifted part of the Laurentian craton to the southeast, or even some other Proterozoic craton. On the interpretive cross section for the transect we speculate that the rocks of the Chain Lakes block were deposited upon Grenvillian crust. The increase of the seismic refraction velocity of about 0.25 km/sec at a depth of about 4 km may be caused by the change from Chain Lakes rocks to Grenvillian rocks. However, as there is no reflector on the seismic reflection profile at that depth, the increase in seismic velocity must occur over a broad depth interval.

The Cameron-Brompton composite terrane is interpreted to be entirely allochthonous upon the Laurentian craton. Neither of the faults that bound this terrane is a strong seismic reflector nor penetrates to mid-crustal depths. The Brompton line that forms the northwestern terrane boundary was mapped as a near-vertical thrust fault zone (St.-Julien and Slivitsky, 1985; Moench and St.-Julien, 1989, fig. 1, p.21). The fault zone is not a seismic reflector near the surface (St.-Julien and others, 1983), but appears at depth to be a weak reflector with southeast dip. The Squirtgun fault, a northwest-directed Acadian thrust fault, forms the boundary between rocks of the Cameron-Brompton and Central Maine terranes. This fault is not imaged near the surface on seismic reflection line 2 (Spencer and others, 1989), but may be one of the many southeast dipping reflectors imaged at depth shown on reflection profile 3A on the graphic.

Relationships among constituent terranes of the Cameron-Brompton composite terrane are complex and their history is only partly deciphered. The Quebec ophiolite and related Cambrian intrusive rocks are overlain by mélanges and olistostromes that are as much as 8 km thick (Hébert and Laurent, 1989). This tract of rocks formed in an oceanic tectonic setting, the Iapetus Ocean, which was of unknown width, although commonly inferred to have been many hundreds of kilometers wide. Geochemical and isotopic evidence (Laurent and Hébert, 1989; Harnois and others, 1990; Church, 1987; Crockett and Oshin, 1987; Shaw and Wasserburg, 1984) suggests that the igneous rocks associated with the Quebec ophiolite were formed by subduction-related magmatism in a fore-arc basin. This would seem to require an oceanic tract at least hundreds of kilometers wide to have been formed and then closed, which is problematic within apparent time constraints of formation in the Cambrian and closure in Middle Ordovician time. Possibly significant strike-slip motion along some faults occurred to bring oceanic crust closer to Laurentia. A small tract of lower Paleozoic calc-alkaline island-arc volcanic rocks has been thrust northwest an unknown distance to its present position over the ophiolite-melange terrane. The Chain Lakes continental terrane with its cover sequence could not have been far outboard from these terranes because kilometer-sized blocks of distinctive Chain Lakes rocks are found within the Quebec olistostromes, notably in the small triangular region on the northeastern side of the transect (Cousineau, 1991). The constituent Cameron-Brompton (Boundary Mountains) terranes were joined together and transported northwesterly by the late Early Ordovician to Middle Ordovician Taconian orogeny (St.-Julien and Hubert, 1975; Spencer and others, 1989). Their space-time relationships are shown on the tectonostratigraphic evolution diagram. The Hurricane Mountain ophiolite and olistostromes and other constituent terranes beneath the Late

Ordovician and younger sedimentary rocks of the Central Maine composite terrane may have been joined to the Cameron-Brompton terrane prior to the accretion of that composite terrane to Laurentia during the Taconian orogeny (Boone and Boudette, 1989).

Terranes within the Cameron-Brompton composite terrane deformed during the Taconian orogeny are unconformably overlain by a sequence of black pelite and basalt of probable Silurian age, taken here to have been deposited in an incipient rift, the Frontenac rift. Conformably overlying rocks (Marvinney, 1989) of the Connecticut Valley trough, a successor basin, include Early Devonian rocks (Hueber and others, 1990) in part coeval with rocks of another successor basin to the southeast in the Central Maine synclinorium. These rocks are thin above the Chain Lakes block, but thicken markedly southeastward across a tectonic hinge zone (Boone and others, 1970; Hatch and others, 1983; Moench and Pankiwskyj, 1988). The tectonic hinge may have resulted from bouyant rise of the overthickened crust to the northwest of the Central Maine synclinorium after the Taconic orogeny (Stewart, 1989). Younger rocks on the Chain Lakes block include sandstones, some with volcanic clasts, and thick rhyolites and garnet rhyolites of late Early Devonian (Siegenian) age (Rankin, 1968, 1980). Such garnet rhyolites can originate only by fractional melting of felsic rocks at lower-crustal or upper-mantle pressures, probably at depths of more than 50 km. This implies that a very thick crust was formed during Taconian and early Acadian compression and by crustal loading of thrust sheets. These distinctive rhyolites are shown on the tectonic setting map to result from continent-continent collision. The onset of the Acadian orogeny terminated sedimentation. The crust was significantly foreshortened by tight folding and extensive thrust faulting. Associated metamorphic grades increase to the southwest to facies as high as middle amphibolite. Numerous calc-alkaline plutons were emplaced immediately after the major deformational phase of the Acadian orogeny. The depth of pluton emplacement also increases to the southwest; southwest of the transect in Maine, both high-grade Carboniferous metamorphism and granitic plutonism result from crustal melting because of overthickening during the Acadian orogeny (DeYoreo and others, 1989). These authors conclude that regional uplift of overthickened crust across the transect in Maine was ~6 km per 100 km along regional strike to the southwest. Subsequently, steep normal faults, such as these along the northwest side of the Chain Lakes block with displacements as much as 5 km, formed in late Paleozoic and early Mesozoic times. Finally, a composite pluton of gabbro and granite was emplaced late in the Mesozoic (Eby, 1984).

Central Maine composite terrane

In the Central Maine composite terrane rocks older than Late Ordovician that were deposited in different tectonic settings are covered by the thick (up to ~13 km) sequence of conglomerates, sandstones, and calcareous siltstones of the Central Maine synclinorium, a post-Taconian successor basin. The rocks of the Central Maine synclinorium are very voluminous, and possibly once contained as much as 2200 cubic km per km of strike length along the synclinorium (Stewart, 1989). Most of these rocks were deposited in Silurian and Early Devonian time, and were severely deformed shortly thereafter during the Acadian orogeny. Thrust faults bound both margins of the Central Maine terrane. The suture line between the Cameron-Brompton (Boundary Mountains) composite terrane and the Central Maine composite terrane is a northwest-directed Penobscottian or Taconian thrust fault, the Squirtgun fault, that

forms the northwest boundary of the Hurricane Mountain belt of ophiolite, olistostromal deposits and flysch. The Hackmatack Pond thrust fault and related faults mark the southeastern boundary of the Central Maine composite terrane. Clearly all the terranes had been assembled by the end of Acadian orogeny. Most of the relationships between these terranes and their basements beneath the blanketing sediments of the Central Maine synclinorium are poorly known or conjectural despite the availability of modern high quality seismic reflection and refraction data. Some of the relations shown on the tectonostratigraphic diagram were projected along strike into the transect.

The Hurricane Mountain belt of ophiolite and olistostromes (Boone and Boudette, 1989; Boone and others, 1989) is as much as 4 km thick and differs from the Quebec ophiolite and olistostromes in the types of rocks found in the ophiolite sequence and the clast assemblages of the olistostromes. The ophiolite may be slightly younger in age, possibly Middle Cambrian. The geochemistry of the ophiolite also differs from that of the Quebec ophiolite (Coish and Rogers, 1987). The lower part of the ophiolite is chemically similar to island arc rocks, and was interpreted to have formed near a subduction zone. The upper part is chemically similar to mid-ocean ridge rocks, and may have formed in a proximal marginal basin. The ophiolite and olistostromes are covered by a volcanoclastic flysch carapace of pre-Middle Ordovician age that may be several km thick.

Several different suites of Cambrian or Ordovician rocks underlie the Central Maine composite terrane. Southeast of the Hurricane Mountain belt, volcanogenic rocks of the Bronson Hill anticlinorium crop out along the edge of the transect. These rocks are interpreted to be a Middle to Late Ordovician volcanic arc (Tucker and Robinson, 1990; Leo, 1985, and in press) and thus represent a different tectonic setting than the Hurricane Mountain belt. Compressional orogenic movements brought these two different volcanogenic assemblages into contact, but the timing and nature of this event is controversial (Boone and Boudette, 1989; Tucker and Robinson, 1990). Thick sections of Cambrian and Ordovician sedimentary rocks are exposed within the Central Maine composite terrane northeast of the transect (Osberg and others, 1985) and are shown only on the tectonostratigraphic section. Some of these rocks contain distinctive fauna of non-North American type (Neuman and others, 1989), as well as a pre-Taconian deformation (Neuman and Max, 1989). Boone and Boudette (1989) interpret these rocks as a separate terrane that was joined to terranes to the northwest in the Penobscottian orogeny. Lower Paleozoic rocks of the Nashoba-Casco-Miramichi composite terrane to the southeast of the Central Maine terrane are known from magnetic and seismic reflection data to underlie the Central Maine composite terrane for at least 25 km northwest of the Hackmatack Pond fault.

The Precambrian basement beneath the Central Maine composite terrane is poorly known, though it is sialic and appears to be broadly homogeneous on the basis of its seismic refraction velocity and its internal seismic reflectors. Where basement is exposed some distance from this transect in the core of one of the domes on the Bronson Hill anticlinorium (the Pelham dome in northern Massachusetts), high grade granulites not unlike Grenvillian gneisses are found, but with a Pb/U zircon age of 613 ± 3 Ma (Tucker and Robinson, 1990). The Grenvillian crust beneath the Cameron-Brompton composite terrane is known (Rankin and others, 1989) to have been extended prior to rupturing to form the Iapetus Ocean and a passive margin at about 650 to 570 Ma. As illustrated on our interpretative cross section for the present Atlantic passive margin, such

extension drastically thins the continental crust prior to rupture. The thinned crust was also heated during Late Proterozoic extension, and may have reached granulite facies at mid-crustal depths, which would have reset most isotopic systems. The core of the Pelham dome in Massachusetts may be Grenvillian granulite reset during Iapetan crustal rupture. This interpretation also is supported by Pb isotopic data from zircons in younger granitic plutons in New Hampshire (Harrison and others, 1987) and by characteristic Pb isotopic compositions of feldspars from plutons that formed by partial melting of deep crustal rocks (Ayuso and Bevier, 1991) in the northern Appalachians. The Ayuso and Bevier Pb data indicate that the isotopic composition of lead in feldspars changes gradually across the Central Maine composite terrane, but has a more uniform radiogenic composition near the coastline in the Atlantica¹ composite terrane. One interpretation of these data is that a crustal boundary or suture exists beneath the Central Maine composite terrane. The gradual nature of the change in Pb isotopic compositions of feldspars from plutonic rocks, however, does not permit a precise location or a determination of what type of crust might overlie another of different Pb isotopic composition. It has been suggested previously that a third type of basement that differs from both Grenvillian and Atlantican (the "Central" basement of Osberg, 1978, or "X" of Zen, 1983) lies beneath the Central Maine composite terrane. We therefore show this possible basement by a different color on our interpretive cross section. However, as N. Rast (oral communication, 1989) has remarked, a third type of basement between very long segments of the two known basements presents a serious topological problem in the context of the genesis of the entire Appalachian-Caledonide orogen. Given the available geologic, isotopic, and seismic evidence, and our new information on the nature and post-Paleozoic age of the reformation of the lower crust, we are skeptical that the third type of basement actually exists. We prefer to interpret most of the Late Proterozoic ages of basement rocks in the parts of the region that experienced large amounts of extension in the Late Proterozoic and extensive Paleozoic deformations to be reset Grenvillian ages. We reserve Atlantican basement to be those rocks upon which sediments were deposited at high paleolatitudes that contain a distinctive Late Proterozoic (Ediacaran) to Early Devonian fauna. We think that the lower crust gained its principal features such as thickness and internal structures during late Paleozoic and early Mesozoic extensional processes, and have shown it separately. This concept is discussed more thoroughly below.

Several large normal faults break the rocks of the Central Maine synclinorium and its basement to mid-crustal depths as shown by the seismic reflection profile. These faults are post-metamorphic and likely are Mesozoic in age. A buried mafic pluton in the upper crust near Waterville, Maine, was inferred from gravity data by Potts and Doll (1987) and is shown as Mesozoic in age on the interpretive cross section, though it could be of late Paleozoic age. The geologic map of Maine (Osberg and others, 1985) shows other such plutons in southwestern Maine. McHone (1978, 1984) reports numerous Mesozoic

¹Atlantica has been proposed by Zen and others (1986; see also Zen, 1989, 1991) as a more inclusive name for terranes commonly called Avalonian. This topic is discussed in greater detail below in connection with the description of numerous terranes within the Atlantica composite terrane.

dikes oriented normal to the principal direction of extension in the transect region, although these dikes are too small to be shown on the geologic map.

Nashoba-Casco-Miramichi composite terrane

The Nashoba-Casco-Miramichi composite terrane (Zen and others, 1986) in this transect is a complex narrow fault-bounded belt of lower Paleozoic metavolcanic and metasedimentary rocks, the Orrington-Liberty welt, that was deposited upon a sialic non-Grenvillian basement (Ayuso and Bevier, 1991), that is neither exposed in the transect nor for hundreds of km along strike to the northeast. Basement to this terrane may be like the Late Proterozoic metasedimentary and metavolcanic rocks in eastern Massachusetts and Connecticut (Zen, 1989). The Nashoba-Casco-Miramichi composite terrane in the transect contains Ordovician metasedimentary and felsic and mafic metavolcanic rocks that have been metamorphosed to kyanite grade (Hussey, 1988), as well as Silurian anatectic granites and migmatites that were modified by middle-to-high grade amphibolite Acadian metamorphism and granitic plutonism. The seismic reflection profile shows many strong reflectors in the crust that dip northwestward beneath the Central Maine composite terrane. These are interpreted to be stacked southeast-directed thrust sheets of a substantial pre-Acadian orogen whose character was not obliterated by Acadian events. Evidence for strong pre-Acadian orogeny along the southeastern side of the orogen along strike in the northern Appalachian is extensive (Hepburn and others, 1987; Van der Pluijm and Van Staal, 1988; Stewart, 1989; Dunning and others, 1990; Bevier and Whalen, 1990; Ayuso and Bevier, 1991).

The Norumbega fault zone is a near vertical zone within the Nashoba-Casco-Miramichi terrane that experienced 10 to 20 km of right lateral strike slip movement (Wones, 1978). It extends across the transect and for additional hundreds of kilometers both southwest and northeast. It has had a long history of movement as it cuts Carboniferous plutons and conglomerates of Carboniferous (or early Mesozoic?) age are found in the fault zone. Motion along this fault continued into the Mesozoic (West and others, 1988), and involved dip-slip as well as strike-slip movement. Distinctive lithotectonic sequences of rift volcanic and successor basin rocks are juxtaposed along the Norumbega fault. Metamorphic assemblages in pre-Silurian rocks northwest of the fault zone experienced as much as 4 kb higher pressures than rocks to the southeast. There is, however, no seismic reflection or refraction evidence along the transect cross section that the Norumbega fault cuts entirely through the present crust; instead it appears to be cut off by a Mesozoic listric fault at a depth of about 20 km. Doll and others (1989) describe seismic reflection profiles that cross part of the Norumbega fault zone about 150 km northeast of the transect profile. There the Norumbega fault zone cuts through the crust and appears to offset the Moho by ~1.5 km. The complex history of motions along the Norumbega fault zone remains to be deciphered, but it is not the fundamental suture with the Atlantican continental block that many workers have taken it to be.

The Hackmatack Pond fault forms the northwest boundary of the Nashoba-Casco-Miramichi composite terrane. A strong seismic reflector that originates at the mapped fault trace dips northwest at about 25° to depths of about 10 km, though it is offset at a depth of about 3 km by a normal fault with about 2 km of south-side down movement. Metasedimentary rocks of the Central Maine synclinorium lie above the reflector, and rocks of the Falmouth-Brunswick sequence of the Cushing Formation of Hussey (1988) occur below it. The

Falmouth-Brunswick rocks are sulfidic gneisses, granofels and amphibolites, possibly of Early Ordovician age (Olszewski and Gaudette, 1988). These rocks contain kyanite and staurolite and have been metamorphosed to significantly higher pressures and temperatures than rocks of the Central Maine synclinorium. Newberg (1986) and Hussey (1988) interpret the Hackmatack Pond fault as a southeast-directed thrust fault. Osberg (1988) considered it to be an unconformity; but in section G-G' in Hatcher and others (1990), Osberg shows it as a folded thrust fault. Our data support the interpretation as a southeast-directed thrust fault. The reflection from the fault trace at depth crosses the seismic reflection profile as a nearly linear feature and is sub-parallel to several strong deeper reflectors also interpreted to be thrust faults. One of the deeper reflectors reaches the surface along the Liberty-Orrington fault. The magnetic anomaly map on the transect and especially bandpass-filtered intermediate wavelength magnetic data (J.D. Phillips, unpublished data) show that the fault surface at the top of the magnetic Falmouth-Brunswick rocks has broad undulations, but dips northwest overall like the strong seismic reflector.

The southeastern border of the Nashoba-Casco-Miramichi composite terrane is marked by northwest-directed thrust faults, the principal of which is the Sennebec Pond fault (St. George fault of Berry and Osberg, 1989). Thrusting along this fault brought a suite of Early Paleozoic slope-rise conglomerates, impure carbonate rocks, sulfidic pelitic rocks and minor basalt onto the earlier formed buttress of the Orrington-Liberty welt. Seismic reflection profile 4 on land has a moderately strong reflector that projects to the trace of the Sennebec Pond fault. The fault dips to the southeast at about 20°, and is only imaged to ~6 km where the profile ends. A similar reflector imaged at 8 to 10 km depth on line 8 beneath west Penobscot Bay may be the base of the obducted rock sequence and, if so, indicates that the Turtle Head fault is not the fundamental suture between different continental blocks postulated by Stewart and Lux (1988).

Atlantica composite terrane

The Atlantica composite terrane southeast of the Sennebec Pond fault consists of several component terranes, each of which contains Cambrian and Ordovician strata deposited in different tectonic settings. All, however, are non-Laurentian. Atlantica contains an assemblage of terranes that share major paleogeographic and faunal similarities that are quite different from those of North American terranes and proximal areas during the Late Proterozoic and early Paleozoic. Zen and others' (1986) concept of Atlantica derives from the early recognition by Walcott (1889, 1891) that Cambrian fauna from eastern New England and the Maritime Provinces of Canada constituted a distinctive Atlantic Province quite different from faunas of typical North American aspect. As summarized by Neuman and others (1989), faunal distinctions have been found to occur from at least the Late Proterozoic (Ediacaran) through, locally, the earliest Devonian (Gedinnian). The composite terrane here called Atlantica has been called Avalonia or Avalon by others (for example, Nance, 1986; Rast and Skehan, 1983) and has been interpreted to be composite as faunal and stratigraphic evidence from the typical platformal rocks argues for their separation into two, or more, terranes (Secor and others, 1983). In the region called Atlantica on this transect, several terranes share paleogeographic and faunal similarities with parts of "Avalonia" but are from different tectonic settings and have different geologic histories, thus

demonstrating the composite nature of Atlantica. We have not followed the usage of Zen (1989) to include the Meguma terrane within the Atlantica composite terrane because its basement, though poorly known, may be different.

Onshore in this transect, the Atlantica composite terrane southeast of the Sennebec Pond thrust fault contains the St. Croix and the Ellsworth-Mascarene terranes. There is little specific evidence to identify the offshore terranes, but samples collected by the submersible Alvin are undeformed alkalic plutonic rocks typical of the Boston-Avalon zone of Hepburn and others (1987), which are equivalent to most of the rocks of the Esmond-Dedham terrane of Gromet (1989). It is not known if the high-grade basement gneisses typical of Gromet's Hope Valley terrane are present, but the highly magnetic rocks within and just southeast of the Gulf of Maine fault zone (Hutchinson and others, 1989) are interpreted here to be gneisses like those of the Hope Valley terrane.

The St. Croix terrane (Ludman, 1987) contains rocks of Cambrian through Late Ordovician age. Conglomerates, quartzite, impure carbonate rocks, and abundant black pelitic rocks and minor bimodal volcanic rocks suggest a continental slope-rise sequence that passes upward into a marginal basin formed upon rifted continental crust (Fyffe and others, 1988). We speculate that the continental basement involved was of part of Atlantica, and that the slope-rise sediments were deposited on its western side. However, no fossils having diagnostic Atlantica affinities have been found in these rocks. The St. Croix terrane had been accreted to the Ellsworth-Mascarene terrane to its southeast by mid- to late-Silurian time as Late Silurian conglomerate was deposited across the boundary faults (Gates, 1989).

The accretion of the already-joined St. Croix and Ellsworth-Mascarene terranes to the Nashoba-Casco-Miramichi composite terrane occurred early in the Acadian orogeny. Accretion may have resulted from a continent-continent collision that caused large thrusts such as the Sennebec Pond fault as well as faults with combined strike-slip and thrust movements such as the Turtle Head fault zone. The Turtle Head fault zone separates the St. Croix terrane from the Islesboro block, which is taken to be a fragment of the Ellsworth-Mascarene terrane. The Islesboro block has a Late Proterozoic basement and a thin platformal cover sequence of conglomerate, quartzite, and impure carbonate rocks and slate that grades up into volcanogenic sedimentary rocks interpreted to be distal to typical Ellsworth-Mascarene rocks (Stewart and Lux, 1988; Stewart, 1974). Blocks of St. Croix terrane rocks occur southeast of the Turtle Head fault zone in juxtaposition with blocks of the volcanoclastic lithologies of the Ellsworth-Mascarene terrane. Rocks in the southwestern part of the St. Croix terrane west of the Turtle Head fault zone reached upper amphibolite facies of Acadian metamorphism prior to completion of right-lateral strike slip and thrusting movement along the fault zone; lower Paleozoic rocks southeast of the fault zone have not exceeded greenschist facies of metamorphism (Guidotti, 1989). The high-grade part of the St. Croix terrane is the only part of Atlantica known to have been significantly affected by Acadian regional metamorphism. The youngest Devonian plutons intrude across the Sennebec Pond and Turtle Head faults and establish that this part of Atlantica was joined to North America by Early Carboniferous time.

The Ellsworth-Mascarene terrane in the transect has a continental Proterozoic basement that is intruded by Late Proterozoic granitic pegmatite (Stewart and Lux, 1988). These rocks are unconformably covered by two

different volcanic-rich sequences. The older sequence is unfossiliferous and consists predominantly of a bimodal continental rift suite of metabasalt and metarhyolite containing interbedded metasiltstone, quartzite, and marble. This sequence is at least 7 km thick on the bases of field mapping and seismic reflection data. The rocks are Cambrian or Ordovician in age and were thoroughly recrystallized to greenschist, probably in the Early Ordovician, prior to their erosion and deposition in unconformably overlying Middle Silurian conglomerate. The younger volcanic sequence is of Silurian and Devonian age, is relatively unmetamorphosed, and contains a characteristic Atlantica (Baltic) fauna. The sequence is mostly a bimodal basalt-rhyolite suite that contains interbedded shallow marine volcanoclastic sedimentary rocks. It is as much as 8 km thick where well exposed along strike northeast of the transect. Numerous mafic and granitic Silurian and Devonian plutons have intruded their consanguineous volcanic cover.

Seismic reflection and refraction data indicate that the St. Croix and Ellsworth-Mascarene terranes have been thrust northwestward to their present positions. Seismic reflection profiles 4 and 8 show reflectors that dip to the southeast and can be interpreted to mark the base of these terranes. Seismic refraction line 8 as interpreted by Luetgert and Mann (1990) suggests that the lower Paleozoic rocks of the Ellsworth-Mascarene terrane rest on basement at depths of 7 to 10 km, and that at about 20 km velocities increase to those characteristic of the lower crust. However, Luetgert and Mann found no unambiguous refraction evidence for a mid-crustal detachment fault as is shown in the interpretive cross section on the basis of reflection lines 4 and 8. Seismic reflection profile 3B across the strike of the Ellsworth-Mascarene terrane (Stewart and others, 1985) images numerous subhorizontal reflectors throughout the upper crust at depths equivalent to the 7 to 10 km thick Paleozoic cover sequences known from regional mapping and Luetgert and Mann's refraction studies. One of the deepest of these reflectors is moderately strong to weak and north of the end of seismic reflection profile 3B projects toward the trace of the Sennebec Pond thrust fault. It could be interpreted to be the detachment fault along which these rocks were transported northwestward. There are few reflectors in the interval below the reflector interpreted to be the Sennebec Pond thrust fault and depths of about 20 km. Below 20 km numerous strong reflectors occur throughout the lower crust to the Moho.

A major fault zone mapped in southern New Brunswick and just off the southeastern Maine coast is projected to cross the transect along a band of strong magnetic highs (Hutchinson and others, 1989). Strong seismic reflectors dipping 25° to 30° south at mid- to deep-crustal levels reach the surface in the same region. They have been interpreted (Hutchinson and others, 1988) to be thrust faults of the Gulf of Maine fault zone, a probable Alleghanian feature by analogy to the geology of eastern Massachusetts and Connecticut (Gromet, 1989). Still another band of magnetic highs and the projection of another seismic reflection to the surface mark the trace of the Fundy fault. This fault probably experienced Alleghanian thrust motion as well as the subsequent normal slip motion, later during the Carboniferous, that formed the large East Narragansett rift basin. This basin crosses the transect as a half graben. Normal slip along the Fundy fault may also have formed a Mesozoic rift basin to the northeast of the transect. The region between the Fundy fault and the northwestern side of the Gulf of Maine fault zone may well be made up of high grade basement gneisses and metamorphosed

cover rocks like those in the Hope Valley terrane of southeastern New England. West and northwest dips are common in the Hope Valley terrane and reflect Alleghanian compression as well as subsequent extension (Goldstein, 1989; Gromet, 1989; Getty and Gromet, 1991). The interpretive cross section shows imbrication of the northwest-dipping structures by younger northwestward-directed thrusts of the Esmond-Dedham terrane. In Massachusetts, geochronologic evidence (Hepburn and others, 1987) indicates that the Esmond-Dedham terrane was joined to cratonal North American about 370 m.y. (Late Devonian) to 340 m.y. (Mississippian). In Massachusetts, fragments of the Ellsworth-Mascarene terrane (the Newbury Volcanic Complex) are caught in faults between the Nashoba-Casco-Miramichi composite terrane and the Esmond-Dedham terrane. The keel of the Mesozoic Middleton basin formed along another fault is exposed nearby. It is clear that faults in the Esmond-Dedham terrane have had a long history of movement as a result of several episodes of compression, transtension, and extension.

Weak, discontinuous arcuate sub-horizontal reflections are imaged on the seismic reflection profile below the reflection Moho in the interval between 485 and 530 km. These resemble sub-Moho reflections observed near 100-150 km on deep seismic reflection profiles 86-1 and 86-2 in the Gulf of St. Lawrence (Marillier and others, 1989). These reflections occur near the deepest parts of the Carboniferous Magdalen basin where Marillier and Verhoef (1989) proposed, from consideration of gravity and seismic refraction velocity data together with the depth to reflection Moho, that a thick layer of mafic rocks having a high density (3.05 g/cm^3) and a high velocity (7.35 km/s) underplated the lower crust during the Carboniferous extension that formed the basin. A swath of broad high magnetic and gravity anomalies extends southwest from the deep part of the Magdalen basin across southern New Brunswick and southeasternmost Maine and passes southwesterly close to the coastline in the region of this transect, just northwest of the Gulf of Maine fault zone. Carboniferous plutons intruded the rocks of this swath in southwestern New Brunswick, and the Carboniferous Fredericton basin lies just north of it. The Gulf of Maine fault zone, interpreted by Hutchinson and others (1988) to be the late Paleozoic compressional Alleghanian (Variscan) front, is underlain by the arcuate sub-Moho reflections as is the extensional Carboniferous East Narragansett basin. It is speculated on the interpretive cross section that the sub-Moho reflections result from Carboniferous mafic underplating. At the very least, these reflectors imply that the upper mantle is laterally heterogeneous locally.

The Esmond-Dedham terrane is mostly underlain by Late Proterozoic calc-alkaline volcanic rocks and coeval plutonic rocks that intruded mildly metamorphosed earlier alkaline volcanic rocks and quartz-rich slope-rise sedimentary rocks deposited in rifts along the margin of an extending craton. Sedimentary cover rocks of Late Proterozoic age in rift basins consist of conglomerate, tillite and basaltic to intermediate volcanic rocks, overlain by platformal Cambrian sedimentary rocks containing characteristic Atlantic (Baltic) fauna representative of cold high-latitude (south polar?) marine environments. All these rocks were intruded by Late Ordovician through Devonian peralkaline plutons that have minor associated volcanic rocks. One of these plutons, the Cashes Ledge, forms a large magnetic high on the southwestern side of the transect. All rocks of the Esmond-Dedham terrane are little deformed, and there was no regional metamorphism above the lower greenschist facies.

Meguma terrane

The Cape Cod-Nauset fault is taken to be the boundary between the Esmond-Dedham block of the Atlantica composite terrane and the Meguma terrane, which underlies the bulk of Nova Scotia. Magnetic and gravity anomaly maps allow the identification of the Meguma terrane at least as far as 67°W longitude (Pe-Piper and Loncarevic, 1989), but farther from land and especially where younger sediments are present, the location of the terrane boundary and even the presence of the Meguma terrane beneath the sea are not well established. Our location is based on the presence of a large paired gravity and magnetic low that is indicative of a very extensive granitic batholith, the Franklin batholith (Hutchinson and others, 1988). The Franklin batholith is comparable to the South Mountain batholith that intrudes the Meguma terrane in Nova Scotia.

The Meguma terrane in Nova Scotia is made up of a very thick section (as much as 10 km) of turbiditic sandstone and shale of lower Paleozoic age, unconformably overlain by shallow water marine sedimentary and bimodal volcanic rocks (Keppie, 1989). Large peraluminous granitic batholiths as young as Late Devonian intrude all these rocks. Some of these plutons have been deformed by strike-slip faults associated with the accretion of Meguma to North America. Clasts from them occur in conglomerates in deep Mississippian basins that overlap the suture. Within this transect no drill hole has penetrated the crystalline basement interpreted here as a southwestern extension of the Meguma terrane, so its nature is virtually unknown. However, immediately to the west of the transect, at about 68°20'W, 40°50'N, drill hole G-1 bottomed in 540 to 450 Ma old graphitic slate, schist, and phyllite that could correspond to Meguma rocks (Klitgord and others, 1988; Schlee and Klitgord, 1988; Schenk, 1991).

The part of this transect south of approximately 42°N latitude overlaps Continent-Ocean Transect E-1 by Thompson and others (in press). Our interpretive cross section is based on seismic reflection profile 19 as is theirs, and from km 700 southeast to its end our cross section coincides with theirs. However, our transect stops north of theirs on what Klitgord and others (1988) call marginal oceanic crust. Our transect terminates about 15 km west of oceanic crust that shows the hyperbolic echoes in seismic reflection profiles taken to typify oceanic crust. The summary below is based on the discussion of the Gulf of Maine embayment and Georges Bank basin given by Klitgord and others (1988).

Mesozoic extension was so extensive that it led to continental breakup. This is the dominant feature of the Meguma terrane within the transect and quite different from the type Meguma terrane in Nova Scotia. Numerous rift basins were formed from the Triassic to late Early Jurassic. They are elongated northeast-southwest along the regional structural grain of the basement, and were probably controlled by the reactivation of older faults. The rift basins contain as much as 5 km of terrestrial sediments and some contain voluminous basaltic volcanic rocks as well as plutonic rocks locally. Some of the rift basins also contain evaporite beds near the top of the section. The rift basins in the northwestern part of Meguma on this transect have been deeply eroded so that only their lower parts remain. A tectonic hinge zone strikes southwest-northeast across the transect at about km 650 to 675. South of the hinge zone, the rift basins are much more completely preserved. The hinge zone developed shortly before crustal break-up produced oceanic crust (late Early Jurassic = 175 Ma). The Georges Bank basin is a

marginal basin that is a successor basin to the rift basins. It contains as much as 7.5 km of sediments in the transect region, and becomes even deeper to the southwest. Northwest of the hinge zone, onlapping, platformal, Coastal Plain sediments are up to 2 km thick.

The continent-ocean boundary is marked by the East Coast Magnetic Anomaly, which is near the present continental shelf edge. The Jurassic and Early Cretaceous paleoshelf edge was a carbonate-bank complex that built upward and seaward as the crust cooled after continental rupture. The carbonate-bank complex separates shelf sediment deposits from slope-rise deposits seaward. Salt diapirs intrude this sediment wedge just seaward of the paleoshelf edge.

Seismic data suggest that the continental crust of Meguma was thinned to about half by extension in the Mesozoic prior to rupture. The propagation of seismic energy is impeded by the thick blanket of post-rupture sediments, and no deep crustal refraction studies have yet been made in this region. By analogy to seismic refraction studies along the same margin off the southeastern United States it is probable that the deep crust was extensively intruded by mafic dikes and that it has higher average density (and seismic refraction velocity) than continental crust (Trehu and others, 1989; Austin and others, 1990). Underplating by upwelling mafic magma generated by pressure relief during rupture is quite probable, but not demonstrated here. If underplated, the lower crust could be continuous from broken former continental to oceanic crust. We surmise that the strong magnetic anomaly about km 830 on the section represents the rise of Mesozoic mafic magma to form a magnetic pluton in the rifted crust. The interpretive cross section is based on control from nearby deep wells and the limited seismic reflection data.

Atlantic oceanic crust

Oceanic crust at the southern end of the transect contains seaward-dipping seismic reflectors (mafic volcanics?) in the basement rocks beneath the sedimentary cover. This type of oceanic crust, called marginal oceanic crust by Klitgord and others (1988), forms a swath about 25 km wide just seaward of the salt diapirs. Individual reflectors can be traced only part of the way across this swath. All the reflectors merge upward into a conformable unit that covers the entire swath. The reflectors disappear at depth, and the zone of reflectors has no distinct base. This type of oceanic crust lacks the hyperbolic echoes in seismic reflection profiles that are typical of oceanic crust generated by sea-floor spreading processes. The East Coast Magnetic Anomaly that marks the continent-ocean crustal boundary is poorly defined by anomalies at the southeast of the transect. The label for this anomaly on the magnetic map follows the trend shown on maps for larger regions (Committee for the Magnetic Anomaly Map of North America, 1987). The strong magnetic anomaly at the southwestern corner of the transect is associated with Bear Seamount, one of the extensive chain of New England seamounts of late Mesozoic age.

EVOLUTION OF THE PRESENT CRUSTAL FRAMEWORK

The cumulative effect of the various Paleozoic orogenies undoubtedly was the formation of thick (~60 km) continental crust as terrane after terrane was accreted. Structural features that are still preserved in the upper crust attest to this: allochthonous sheets can be imaged to depths of more than 20 km, areas of younger metamorphic rocks containing mineral assemblages

indicative of pressures as high as 9 kb (~30 km) are now exposed on the surface, and garnet rhyolite formed by fractional melting of felsic rocks deeper than 50 km was extruded. Our composite seismic reflection profile demonstrates that the upper crust still retains coherent regional structural dips; these may be opposed to each other in adjacent terranes as in the case of the Nashoba-Casco-Miramichi composite terrane and the Atlantica composite terrane between 325 and 550 km on the interpretive cross section.

The crust today, however, lacks overthickened parts. The crust beneath the craton is about 43 km thick over broad regions of the northern Appalachians. In this transect the composite of seismic reflection profiles shows a stepwise crustal thinning toward the ocean basin having broad steps 75 to hundreds of km wide, and narrow risers 2 to 4 km high and 10 to 20 km wide where the Moho shallows. Other profiles elsewhere in the northern Appalachians (Marillier and others, 1989; Phinney and Roy-Chowdhury, 1989) also do not show overthickened crust, but show broad regions of crust having little variation in thickness and narrow regions where the Moho rises or falls from 2 km to as much as 6 km over a few tens of km. There is no systematic rise seaward as shown by this transect. In the northern Appalachians steps on the Moho are not simply related to any terrane boundary mapped at the surface. Evidently, since the late Paleozoic deep crustal processes have obliterated evidence of earlier Paleozoic orogenies, and the crust has been restored to normal continental thickness overall but thinned toward the ocean. Undoubtedly some of the continental crust has been removed by erosion; estimates based on fission track dates (Doherty and Lyons, 1980) suggest the erosion of about 6 ± 1 km of rock since the Jurassic for western Maine. This must be a maximum. Obviously the amount of erosion was not uniform along the transect, as the Coastal Plain (shown by overprints on the geologic map) was the site of deposition for much of the time since continental break-up. Zen (1991) has described denudation in the New England Appalachians in detail.

Tectonic restoration of normal crustal thicknesses had begun in mid-Paleozoic times, as evidenced by the hinge line in the rocks of the Central Maine synclinorium, a Silurian and Early Devonian successor basin. Thermal recovery of thick crust led to partial melting and the diapiric rise of magma into and upon the cover sequences of deep basement, depleting the deep crust in quartz and alkali feldspar and thus imposing the compositional transition in the lower crust toward the higher density and seismic velocity observed now. The lower crust lost its stiffness as a result of added internal and external heat, some possibly resulting from mafic underplating. The upper crust remained stiff and brittle, though some heating may have occurred by the mechanical transport of heat by ascending magmatic diapirs and fluids. The softening of the lower crust together with global plate tectonic forces led to the beginning of crustal transtension in the Carboniferous and the formation of deep half graben such as the East Narragansett basin near 530 km on the interpretive cross section.

This transect demonstrates that the tectonic processes that disrupted Pangea, the supercontinent that was assembled near the end of the Paleozoic, imprinted their effects upon a swath of continental crust that is now about 650 km wide in the part still attached to North America. These effects increase in intensity from the craton toward the oceanic basin, and are less apparent in the upper crust than in the lower crust. The effects in the upper crust are occult toward the craton and consist principally of normal faults and the intrusion of oriented dike swarms and small plutonic complexes. Only

limited extension is indicated, possibly only a few percent. This appears to be the case from about 175 to 325 km along the interpretive cross section. As the amount of extension in the upper crust increased, crustal blocks were rotated along favorably oriented older faults that were reactivated and probably become listric at mid-crustal depths; rift basins were formed where the extension was localized in the upper crust. The upper crust may have been thinned by as much as 20 percent as is shown from about 325 to 625 km along our interpretive cross section. More extensive rifting accompanied by basaltic volcanism occurred close to the locus of continental break-up. The upper crust may have been thinned as much as 40 percent when break-up occurred. This thinning is shown from 625 to 875 km along the interpretive cross section.

The effects of extension upon the lower crust and Moho were profound as is most evident in the seismic reflection and refraction data, as well as the broad southeasterly increase in the regional gravity field, most easily seen on the gravity profile. One of the most important features was the demarcation of the lower crust itself. The top of the lower crust is defined by a gradual increase in the seismic compressional wave velocity from about 6.4 km/s to about 6.8 km/s over a depth of as much as 4 km (Stewart and Luetgert, 1990). The top of the lower crust is not a reflector at near-vertical angles of incidence as can be seen by comparison of the seismic refraction and seismic reflection profiles. Seismic refraction data both parallel and normal to regional strike show that the top of the lower crust has little relief, rising smoothly from about 24 km depth under the Cameron-Brompton (Boundary Mountains) composite terrane to about 20 km beneath the northern part of the Atlantica composite terrane, a distance of over 300 km. The top of the lower crust is not offset by any terrane boundaries or faults. Using seismic refraction data collected after this transect was completed, J.H. Luetgert (written communication, 1990) has contoured the top of the lower crust for much of Maine, New Hampshire, and Vermont. He found that the top of the lower crust has a slight dip to the southwest beneath this transect in Maine, being as much as 2 km deeper near Farmington. Over a region in northern New England 600 km wide the relief on the top of the lower crust is only about 4 km, and no terrane boundaries are offset.

The velocity change at the top of the lower crust is sufficient to be observed only within the part of the orogen that experienced late Paleozoic or Mesozoic extension as shown by oriented dike swarms, rift basins, or steps in the regional gravity gradient parallel to the regional structural grain. As the seismic velocity in shallow Grenvillian crust is higher (~6.6 km/s), the top of the lower crust cannot be clearly identified therein (J.H. Luetgert, written communication, 1990). The interpretation that the lower crust took its present configuration during late Paleozoic and early Mesozoic extension is supported by several other lines of evidence.

Extension seems to have affected the properties of the Moho and the lower crust even where the effects in the upper crust may be occult. The Moho beneath the craton commonly is an indistinct gradational velocity change that does not yield a strong reflection at any angle of incidence or a sharp refraction, whereas a sharp crust-mantle boundary is reported (Taylor, 1989, fig. 5) in regions where the orogen has been extended. This results in a stronger Moho signal in extended regions. The transition can be seen from about 200 to 240 km of the composite seismic reflection profile, and more distinctly in the larger more detailed figures in Spencer and others (1989).

The Moho becomes stronger further to the southeast and remains strong to near the end of the transect where it is lost beneath thick carbonate-bank sediments that inhibit transmission of reflected sound energy.

As the amount of extension increases, reflectors in the lower crust become more numerous and stronger, although commonly they are sub-horizontal and discontinuous, and individual reflectors are only about 10 to 15 km long. This phenomenon has been observed widely, though seldom have corresponding refraction velocity profiles been available for the identification of the top of the lower crust, as is the case for this transect for over 300 km between ~200 and ~540 km. The numerous reflectors observed in the lower crust of this transect suggest that during deformation much of the lower crust yielded ductilely along subhorizontal anastomosing shear zones, causing the reflective velocity contrasts. Not all crustal blocks yielded in the same way. The bands of reflections in the Gulf of Maine fault zone and the Fundy fault pass continuously from the bottom of the upper crust through most of the lower crust until they merge into a layer of subhorizontal reflections 3 or 4 km thick that marks the bottom of the lower crust and reflection Moho. The crust has the same thickness in this region as in blocks to the north and south, suggesting that ductile extension of the lower crust was concentrated at the bottom of the lower crust. It is possible that the dip of the bands of reflectors was reduced by southeastward plastic flow of the deep crust.

Two observations gained by comparing coincident seismic reflection and refraction data support the interpretation that most of the lower crust was homogenized by plastic flow and recrystallization during extension so that it lacks detectable through-going faults.

The first observation is that the strong seismic reflector at the top of autochthonous Grenvillian crust beneath the Cameron-Brompton (Boundary Mountains) composite terrane dies out at the same depth as the top of the lower crust as determined by seismic refraction data. This would be expected if the boundary between the upper crust and the lower crust was a zone of rheological contrast that localized intense deformation and recrystallization, thus obliterating the strong fabric and compositional contrasts that cause the reflector. The disappearance of this reflector at the top of the lower crust makes interpretation of the southeastern extent of Grenvillian crust speculative; gravity and magnetic data are not definitive either. Geologic evidence from the northwest end of the transect indicates that the Grenvillian crust was extended, so that it thinned and ruptured to form the Iapetan passive margin. The later thrust imbrication of thinned Grenvillian blocks, interpreted to underlie the Cameron-Brompton composite terrane, must represent a considerable shortening of the Iapetan passive margin, and geologic and isotopic evidence suggests that the basement that constitutes the upper crust of the Central Maine terrane may not be Grenvillian. However, the nature of the lower crust is not constrained. Therefore the interpretive cross section shows an arbitrary boundary for the southeastern limit of Grenvillian lower crust. Another possible interpretation is that the top of Grenvillian lower crust continues to deepen at about the dip of the decollement surface until it pinches out near 320 km on the interpretive cross section. Such a tapered edge would be the result of extensive plastic flow, as the end of the ruptured Iapetan continental block probably would have been blunt like crust at the present Atlantic continental edge.

The second line of evidence is that young normal faults that break the upper crust, such as the one in southeastern part of the central Maine synclinorium, become listric and are lost at or near the top of the lower crust as indicated by the rapid increase of seismic refraction velocity at a depth of 20 km. A geographically coincident rise of the Moho of about 4 km

occurs below the footwall of this listric fault as indicated by well-constrained seismic reflection and refraction data. The intervening lower crust is strongly laminated with subhorizontal reflections. All of these observations support the interpretation of extensive plastic flow in the lower crust, in the direction of ultimate crustal rupture. Examples of the rise of Moho beneath the footwall of listric faults and the development of a well-laminated lower crust are numerous on seismic reflection profiles from the northern Appalachians (Phinney and Roy-Chowdhury, 1989; Marillier and others, 1989), although determinations of the refraction velocity changes are lacking. A conspicuous example occurs beneath the Franklin rift basin at about 650 km on the interpretive cross section. Phinney and Roy-Chowdhury (1989, fig. 12d) give another good example from the east end of line USGS-36 on the Long Island platform beneath the early Mesozoic Nantucket rift basin. If the large normal displacements on mapped faults are considered (Lyons and others, 1991), COCORP line 6 in southwestern New Hampshire beneath the Bronson Hill anticlinorium and Central Maine synclinorium (Phinney and Roy-Chowdhury, 1989, fig. 16) can be interpreted to show the same phenomenon. Seismic refraction data from northern New Hampshire (Hughes and Luetgert, in press) suggest that the top of the lower crust is at about 18 km depth. This corresponds closely to the depth on COCORP line 6 at which east-dipping reflectors, interpreted as decollements, die out, and where laminated subhorizontal reflectors appear in the lower crust in conjunction with a distinct Moho reflection.

The lower crust part of the continental crust that has been extended more than 10 or 15 percent cannot be assigned with confidence to a terrane that lies above it. As reflectors that can be traced downward from structures at the surface are obliterated, identity is lost as one block of lower crust has much the same velocity structure and bulk composition as another. The problem becomes more difficult the greater the distance from the craton, as the lower crust is thinned significantly more than the upper crust (Keen and others, 1987). Hence we have shown lower crust in a different pattern from the upper crust outboard from the arbitrary boundary of Grenvillian lower crust. The isotopic evidence of Ayuso and Bevier (1991) best describes the crust as it was in the middle Paleozoic when it was being sampled by partial melting and the ascent of granitic magmas. These data are not directly applicable to the lower crust after it was reformed in the late Paleozoic and early Mesozoic. Suites of deep crustal xenoliths from Mesozoic dikes and plutons would provide more reliable samples of lower crustal compositional variation now. Deep seismic refraction velocities below 7.1 km/s show that the lower crust is of sialic composition, probably plagioclase-rich pyroxene and amphibole granulites. No tract of transitional or oceanic crust has been identified to underlie any terrane northwest of Meguma.

REFERENCES CITED

- Ando, C.J., Cook, F.A., Oliver, J.E., Brown, L.D., and Kaufman, Sidney, 1983, Crustal geometry of the Appalachian orogen from seismic reflection studies: in Hatcher, R.D., Jr., Williams, Harold, and Zeitz, Isidore, eds., Contributions to the Tectonics and Geophysics of Mountain Chains, Geological Society of America Memoir 158, p. 83-101.
- Austin, J.A., Jr., Stoffa, P.L., Phillips, J.D., Oh, Jinyong; Sawyer, D.S., Purdy, G.M., Reiter, Ed, and Makris, Janis, 1990, Crustal structure of the southeast Georgia embayment-Carolina trough: Preliminary results of a composite seismic image of a continental suture(?) and a volcanic passive margin: *Geology*, v. 18, 1023-1027.
- Ayuso, R.A., and Bevier, M.L., 1991, Regional differences in Pb isotopic composition of feldspars in plutonic rocks of the northern Appalachian Mountains, USA and Canada: A geochemical method of terrane correlation: *Tectonics*, v. 10, p. 191-212.
- Berry, H.N., IV, and Osberg, P.H., 1989, A stratigraphic synthesis of eastern Maine and western New Brunswick, in Tucker, R.D., and Marvinney, R.G., eds., *Studies in Maine Geology*, v. 2: Structure and Stratigraphy, p. 1-32.
- Bevier, M.L., and Whalen, J.B., 1990, Tectonic significance of Silurian magmatism in the Canadian Appalachians: *Geology*, v. 18, p. 411-414.
- Boone, G.M., Boudette, E.L., and Moench, R.H., 1970, Bedrock geology of the Rangeley Lakes-Dead River Basin region, western Maine, in Boone, G.M., ed., *Guidebook for field trips in the Rangeley Lakes-Dead River Basin region, western Maine*, 62nd New England Intercollegiate Field Conference, Rangeley, Maine, p. 1-24.
- Boone, G.M., and Boudette, E.L., 1989, Accretion of the Boundary Mountains Terrane within the northern Appalachians orotectonic zone, in Horton, J.W., Jr., and Rast, N., eds., *Melanges and olistostromes of the U.S. Appalachians*: Geological Society of America Special Paper 228, p. 17-42.
- Boone, G.M., Doty, D.T., and Heizler, M.T., 1989, Hurricane Mountain Formation melange: Description and tectonic significance of a Penobscottian accretionary complex, in Tucker, R.D., and Marvinney, R.G., eds., *Studies in Maine Geology*, v. 2: Structure and Stratigraphy, Maine Geological Survey, Augusta, Maine, p. 33-83.
- Church, W.R., 1987, The geochemistry and petrogenesis of ophiolitic volcanic rocks from Lac de l'Est, Thetford Mines Complex, Quebec, Canada: Discussion: *Canadian Journal of Earth Sciences*, v. 24, p. 1270-1273.
- Coish, R.A., and Rogers, N.W., 1987, Geochemistry of the Boil Mountain ophiolitic complex, northwest Maine, and tectonic implications: *Contributions to Mineralogy and Petrology*, v. 97, p. 51-65.
- Cousineau, P.A., 1991, The Riviere Des Plante ophiolitic melange: Tectonic setting and melange formation in the Quebec Appalachians: *Journal of Geology*, v. 99, p. 81-96.
- Crocket, J.H., and Oshin, I.O., 1987, The geochemistry and petrogenesis of ophiolitic volcanic rocks from Lac de l'Est, Thetford Mines Complex, Quebec, Canada: Reply: *Canadian Journal of Earth Sciences*, v. 24, p. 1273-1275.
- Decker, E.R., 1987, Heat flow and basement radioactivity in Maine: First-order results and preliminary interpretation: *Geophysical Research Letters*, v. 14, p. 256-259.

- DeYoreo, J.J., Lux, D.R., Guidotti, C.V., Decker, E.R., and Osberg, P.H., 1989, The Acadian thermal history of western Maine: *Journal of Metamorphic Geology*, v. 7, p. 169-190.
- Doherty, J.T., and Lyons, J.B., 1980, Mesozoic erosion rates in New England: *Geological Society of America Bulletin*, v. 91, p. 16-20.
- Doll, W.E., Costain, J.K., Coruh, C., Domoracki, W.D., Potts, S.S., Ludman, A., and Hopeck, J., 1989, Results of a seismic reflection and gravity study of the Bottle Lake complex: *EOS*, v. 70, p. 401.
- Dunning, G.R., and Cousineau, P.A., 1990, U/Pb ages of single zircons from Chain Lakes massif and a correlative unit in ophiolite mélange in Quebec: *Geological Society of America Abstracts with Programs*, v. 22, no. 2, p. 13.
- Dunning, G.R., and Cousineau, P.A., in press, U/Pb ages of single zircons from the Chain Lakes massif, Maine, and equivalent rocks in ophiolitic mélange in Quebec. *Geological Society of America Bulletin*, v. ___, p. ___.
- Dunning, G.R., O'Brien, S.J., Colman-Sadd, S.P., Blackwood, R.F., Dickson, W.L., O'Neill, P.P., and Krogh, T.E., 1990, Silurian orogeny in the Newfoundland Appalachians: *Journal of Geology*, v. 98, p. 895-913.
- Eby, G.N., 1984, Geochronology of the Monteregian Hills alkaline igneous province, Quebec: *Geology*, v. 12, p. 468-470.
- Fyffe, L.R., Stewart, D.B., and Ludman, Allan, 1988, Tectonic significance of black pelites and basalts in the St. Croix terrane, coastal Maine and New Brunswick: *Maritime Sediments and Atlantic Geology*, v. 24, p. 281-288.
- Gates, Olcott, 1989, Silurian roundstone conglomerates of coastal Maine and adjacent New Brunswick, *in* Tucker, R.D., and Marvinney, R.G., eds., *Studies in Maine Geology*, v. 2: Structure and stratigraphy, p. 127-144.
- Getty, S.R., and Gromet, L.P., 1991, Late Paleozoic tectonism in Avalonian basement rocks: a view from beneath the Devonian (Acadian) metamorphic belt [abs.]: *Geological Society of America Abstracts with Programs*, v. 23, no. 1, p. A34.
- Globensky, Yvon, 1987, *Geologie des Basses-Terres du St.-Laurent*: Ministère de l'Énergie et des Ressources, Quebec, Canada, Report MM 85-02, map 1999, scale 1:250,000.
- Goldstein, A.G., 1989, Tectonic significance of multiple motions on terrane-bounding faults in the northern Appalachians: *Geological Society of America*, v. 101, p. 927-938.
- Gromet, L.P., 1989, Avalonian terranes and late Paleozoic tectonism in southeastern New England; Constraints and problems: *in* Dallmeyer, R.D., ed., *Terranes in the Circum-Atlantic Paleozoic Orogens*, *Geological Society of America Special Paper* 230, p. 193-211.
- Guidotti, C.V., 1989, Metamorphism in Maine: An overview, *in* Tucker, R.D., and Marvinney, R.G., eds., *Studies in Maine Geology*, v. 3: Igneous and metamorphic geology, p. 1-17.
- Harnois, Luc, Trottier, Jacques, and Morency, Maurice, 1990, Rare earth element geochemistry of Thetford Mines ophiolite complex, Northern Appalachians, Canada: *Contributions to Mineralogy and Petrology*, v. 105, p. 433-445.
- Harrison, T.M., Aleinikoff, J.N., and Compston, W., 1987, Observations and controls in the occurrence of inherited zircon in Concord-type granitoids, New Hampshire: *Geochimica et Cosmochimica Acta*, v. 51, p. 2549-2558.

- Hatch, N.L., Jr., Moench, R.H., and Lyons, J.B., 1983, Silurian-Lower Devonian stratigraphy of eastern and south-central New Hampshire: Extensions from western Maine: *American Journal of Science*, v. 283, p. 739-761.
- Hatcher, R.D., Jr., Osberg, P.H., Drake, A.A., Jr., Robinson, Peter, and Thomas, W.A., 1990, Tectonic map of the U.S. Appalachians, in Hatcher, R.D., Jr., Thomas, L.A., and Viele, G.W., eds., *The Appalachians-Ouachita Orogen in the United States: Geology of North America*, v. F-2, plate 1.
- Hébert, Rejean and Laurent, Roger, 1989, Mineral chemistry of ultramafic and mafic plutonic rocks of the Appalachian ophiolites, Quebec, Canada: *Chemical Geology*, v. 77, p. 265-285.
- Hennet, C.G., Luetgert, J.H., and Phinney, R.A., in press, The crustal structure in central Maine from coherency processed refraction data: *Journal of Geophysical Research*, v. 96, p. ____.
- Hepburn, J.C., Hill, Malcolm, and Hon, Rudolph, 1987, The Avalonian and Nashoba terranes, eastern Massachusetts, USA: An overview: *Maritime Sediments and Atlantic Geology*, v. 23, p. 1-12.
- Hermes, O.D., Ballard, R.D., and Banks, P.O., 1978, Upper Ordovician peralkalic granites from the Gulf of Maine: *Geological Society of America Bulletin*, v. 89, p. 1761-1774.
- Hueber, F.M., Bothner, W.A., Hatch, N.L., Jr., Finney, S.C., and Aleinikoff, J.N., 1990, Devonian plants from southern Quebec and northern New Hampshire and the age of the Connecticut Valley trough: *American Journal of Science*, v. 250, p. 360-395.
- Hughes, Stephen, and Luetgert, J.H., in press, Crustal structure of the western New England Appalachians and Adirondack Mountains: *Journal of Geophysical Research*, v. ____ p. ____.
- Hussey, A.M., III, 1988, Lithotectonic stratigraphy, deformation, plutonism, and metamorphism, greater Casco Bay region, southwestern Maine, in Tucker, R.D., and Marvinney, R.G., eds., *Studies in Maine Geology*, v. 1: *Structure and Stratigraphy*, p. 17-34.
- Hutchinson, D.R., Klitgord, K.D., Lee, M.W., and Trehu, A.M., 1988, U.S. Geological Survey deep seismic reflection profile across the Gulf of Maine: *Geological Society of America Bulletin*, v. 100, p. 172-184.
- Hutchinson, D.R., and Lee, M.W., 1989, Processing and attenuation of noise in deep seismic reflection data from the Gulf of Maine: *Marine Geophysical Research*, v. 11, p. 51-67.
- Hutchinson, D.R., Klitgord, K.D., Bothner, W.A., and Brooks, J.A., 1989, Gulf of Maine fault zone--a reactivated crustal suture in Avalonian rocks beneath the Gulf of Maine: *Geological Society of America Abstracts with Programs*, v. 21, no. 2, p. 24.
- International Association of Geodesy, 1971, Geodetic reference system 1967: *International Association of Geodesy Special Publication No. 3*, 116 p.
- Kafka, A.L., and Ebel, J.E., 1988, Seismic structure of the Earth's crust underlying the State of Maine, in Tucker, R.D., and Marvinney, R.G., eds., *Studies in Maine Geology*, v. 1, p. 137-156.
- Kafka, A.L., and Reiter, E.C., 1987, Dispersion of Rg waves in southeastern Maine: Evidence for lateral anisotropy in the shallow crust: *Bulletin of the Seismological Society of America*, v. 77, p. 925-941.
- Keen, C.E., Stockmal, G.S., Welsink, H., Quinlan, G., and Mudford, B., 1987, Deep crustal structure and evolution of the rifted margin northeast of Newfoundland: results from Lithoprobe East: *Canadian Journal of Earth Sciences*, v. 24, p. 1537-1549.

- Keppie, J.D., 1989, Northern Appalachian terranes and their accretionary history: in Dallmeyer, R.D., ed., Terranes in the Circum-Atlantic Paleozoic Orogens, Geological Society of America Special Paper 230, p. 159-191.
- Klemperer, S.L., and Luetgert, J.H., 1987, A comparison of reflection and refraction processing and interpretation methods applied to conventional refraction data from coastal Maine: Bulletin of the Seismological Society of America, v. 77, p. 614-630.
- Klitgord, K.D., Hutchinson, D.R., and Schouten, Hans, 1988, U.S. continental margin; Structural and tectonic framework, in Sheridan, R.E., and Grow, J.A., eds., The Atlantic Continental Margin: U.S. Geological Society of America, The Geology of North America, v. I-2, p. 19-55.
- Klitgord, K.D., Schlee, J.S., and Hinz, K., 1982, Basement structure, sedimentation, and tectonic history of the Georges Bank basin, in Scholle, P.A. and Wenkham, C.R., eds., United States Outer Continental Shelf; Geological studies of the COST nos. G-1 and G-2 wells: U.S. Geological Survey Circular 861, p. 160-186.
- Laurent, Roger, and Hébert, Réjean, 1989, The volcanic and intrusive rocks of the Quebec Appalachian ophiolites (Canada) and their island-arc setting: Chemical Geology, v. 77, p. 287-302.
- Leo, G.W., 1985, Trondhjemite and metamorphosed quartz keratophyre tuff of the Ammonoosuc Volcanics (Ordovician), western New Hampshire and adjacent Vermont and Massachusetts: Geological Society of America Bulletin, v. 96, p. 1493-1507.
- _____, in press, Oliverian domes, related plutonic rocks, and mantling Ammonoosuc volcanics of the Bronson Hill anticlinorium, New England Appalachians: U.S. Geological Survey Professional Paper 1516, _____ p.
- Ludman, Allan, 1987, Pre-Silurian stratigraphy and tectonic significance of the St. Croix Belt, southeastern Maine: Canadian Journal of Earth Science, v. 24, p. 2459-2469.
- Luetgert, J.H., 1988a, Users manual for RAY84/R83PLT, Interactive two-dimensional raytracing/synthetic seismogram package: U.S. Geological Survey Open File Report 88-238, 52 p.
- _____, 1988b, Users manual for R1D84, Interactive modeling of one-dimensional velocity-depth functions: U.S. Geological Survey Open File Report 88-247, 30 p.
- _____, 1988c, Users manual for RSEC88, Interactive computer program for plotting seismic refraction record sections: U.S. Geological Survey Open File Report 88-262, 89 p.
- Luetgert, J.H., Mann, C.E., and Klemperer, S.L., 1987, Wide-angle deep crustal reflections in the northern Appalachians: Geophysical Journal Royal Astronomical Society, v. 89, p. 183-188.
- Luetgert, J.H., and Mann, C.E., 1990, Avalon terrane in eastern coastal Maine: Seismic refraction--wide-angle reflection data: Geology, v. 18, p. 878-881.
- Lyons, J.B., Bothner, W.A., Moench, R.H., and Thompson, J.B., Jr., 1991, A new bedrock geologic map of New Hampshire: Geological Society of America, Abstracts with Programs, v. 23, no. 1, p. 60.
- Macnab, Ron, Shih, K.-G., Bothner, W.A., Brooks, John, Delorey, Cathy, and Klitgord, K.D., 1990, Magnetic data over Gulf of Maine and adjacent land areas: Preparation of a data base for construction of a 1:500,000 magnetic anomaly base: Geological Survey of Canada Open File 2295.

- Magnetic Anomaly Map of North America Committee, 1987, Magnetic Anomaly Map of North America, Geological Society of America, Boulder, CO, Centennial Scale Map--003, 4 sheets, scale 1:5,000,000.
- Marillier, Francois, Keen, C.E., Stockmal, G.S., Quinlan, Gary, Williams, Harold, Colman-Sadd, S.P., and O'Brien, S.J., 1989, Crustal structure and surface zonation of the Canadian Appalachians: implications of deep seismic reflection data: Canadian Journal of Earth Sciences, v. 26, p. 305-321.
- Marillier, Francois, and Verhoef, J., 1989, Crustal thickness under the Gulf of St. Lawrence, northern Appalachians, from gravity and seismic data: Canadian Journal of Earth Sciences, v. 26, p. 1517-1532.
- Marvinney, R.G., 1989, Bedrock geology of the southeastern margin of the Connecticut Valley-Gaspé Synclinorium in Somerset County, Maine, in Berry, A.W., Jr., ed., Guidebook for field trips in southern and west-central Maine, New England Intercollegiate Geological Conference, 81st Annual Meeting, Farmington, Maine, p. 212-237.
- McHone, J.G., 1978, Distributions, orientations, and ages of mafic dikes in central New England: Geological Society of America Bulletin, v. 89, p. 1645-1655.
- _____, 1984, Mesozoic igneous rocks of northern New England and adjacent Quebec: Geological Society of America Map and Chart Series MC-49, 5 p.
- Meyer, R.P., Steinhart, J.S., Howell, B.F., Bonini, W.E., Fahiquist, D.A., and Aldrich, L.T., 1962, Cooperative Maine experiment in crustal seismology: Methods and application of fixed linear recording arrays to crustal measurements [abs.]: J. Geophys. Res., v. 67, p. 3580-81.
- Moench, R.H., and Pankiwskyj, K.A., 1988, Geologic map of western interior Maine: U.S. Geological Survey Miscellaneous Investigation Series, Map I-1692 and text 21 p.
- Moench, R.H., and St.-Julien, Pierre, 1989, Northern Appalachian transect: southeastern Quebec, Canada through western Maine, U.S.A. Field Trip Guidebook T358, 28th International Geological Congress, Washington, D.C., American Geophysical Union, 52 p.
- Monger, J.W.H., 1986, The Global Geoscience Transects Project: International Union of Geological Sciences. Episodes, v. 9, p. 217-222.
- Moore, J.M., Davidson, A., and Baer, A.J., 1986, The Grenville Province: Geological Association of Canada Special Paper 31, 358 p.
- Morelli, C., ed., 1974, The international gravity standardization net 1971: International Association of Geodesy Special Publication No. 4, 49 p.
- Murphy, J.M., and Luetgert, J.H., 1986, Data report for the Maine-Quebec cross-strike seismic-refraction profile: U.S. Geological Survey Open File Report 86-47, 71 p.
- _____, 1987, Data report for the 1984 Maine along-strike seismic-refraction profiles: U.S. Geological Survey Open File Report 87-133, 134 p.
- Nance, R.D., 1986, Precambrian evolution of the Avalon terrane in the northern Appalachians: A review: Maritime Sediments and Atlantic Geology, v. 22, p. 214-238.
- Nakamura, Yosio, and Howell, B.F., Jr., 1964, Maine seismic experiment: Frequency spectra of refraction arrivals and the nature of the Mohorovičić discontinuity. Bulletin Seismological Society of America, v. 54, p. 9-18.
- Naylor, R.S., Boone, G.M., Boudette, E.L., Ashenden, D.D., and Robinson, Peter, 1973, Pre-Ordovician rocks in the Bronson Hill and Boundary Mountain anticlinoria, New England, U.S.A.: EOS, Transactions American Geophysical Union, v. 50, no. 4, p. 495.

- Neuman, R.B., Palmer, A.R., and Dutro, J.T., Jr., 1989, Paleontological contributions to Paleozoic paleogeographic reconstructions of the Appalachians, in Hatcher, R.D., Jr., Thomas, W.A., and Viele, G.W., eds., The Appalachian-Ouachita Orogen in the United States: Geological Society of America, The Geology of North America, Volume F-2, p. 375-384.
- Neuman, R.B., and Max, M.D., 1989, Penobscottian-Grampian-Finnmarkian orogenies as indicators of terrane linkages: in Dallmeyer, R.D., ed., Terranes in the Circum-Atlantic Paleozoic Orogens, Geological Society of America Special Paper 230, p. 31-45.
- Newberg, D.W., 1985, Bedrock geology of the Pelermo 7.5' quadrangle, Maine: Maine Geological Survey, Open-file Report 85-84, 14 p.
- Olszewski, W.J., Gaudette, H.E., and Cheatham, M.M., 1990, 207Pb/206Pb ages from the Chain Lakes massif, N.W. Maine using single grain direct thermal ionization: Geological Society of America Abstracts with Programs, v. 22, no. 2, p. 61.
- Olszewski, W.J. and Gaudette, H.E., 1988, Early Paleozoic thermotectonic history of eastern New England: Cambro-Ordovician metamorphism and plutonism -- a distinctive feature of the "Casco" terrane: Geological Society of America Abstracts with Programs, v. 20, p. 60.
- Osberg, P.H., 1978, Synthesis of the geology of the northeastern Appalachians, in Tosier, E.T., and Schenk, P.E., eds., Caledonian-Appalachian Orogen in the North Atlantic region, Geological Survey of Canada Paper 78-13, p. 137-147.
- Osberg, P.H., 1988, Geologic relations within the shale-wacke sequence in south-central Maine, in Tucker, R.D., and Marvinney, R.G., eds., Studies in Maine Geology, v. 1, p. 51-73.
- Osberg, P.H., Tull, J.F., Robinson, Peter, Hon, Rudolph, Butler, J.R., 1989, The Acadian orogen, in Hatcher, R.D., Jr., Thomas, W.A., and Viele, G.W., eds., The Appalachian-Ouachita Orogen in the United States: Geological Society of America, Geology of North America, v. F-2, p. 179-232.
- Osberg, P.H., Hussey, A.M., III, and Boone, G.M., 1985, Bedrock geologic map of Maine: Maine Geological Survey, Augusta, Maine, scale 1:500,000.
- Pe-Piper, Georgia, and Loncarevic, B.D., 1989, Offshore continuation of Meguma terrane, southwestern Nova Scotia. Canadian Journal of Earth Sciences, v. 26, p. 176-191.
- Phillips, J.D., 1990, Integration of potential-field and digital geologic data for two North American geoscience transects: Journal of Geological Education, v. 38, p. 330-338.
- Phillips, J.D., Thomas, M.D., and Jahrling, C.E., 1988, Principal facts for gravity stations along lines 1, 2, 3A, and 3B of the Quebec-Western Maine seismic reflection profile, and along MERQ seismic reflection line 2001, southeastern Quebec and west-central Maine: U.S. Geological Survey Open File Report 88-425, 46 p.
- Phinney, R.A., and Roy-Chowdhury, Kabir, 1989, Reflection seismic studies of crustal structure in the eastern United States, in Pakiser, L.C., and Mooney, W.D., eds., Geophysical Framework of the continental United States: Geological Society of America Memoir 172, p. 613-652.
- Potts, S.S., and Doll, W.E., 1987, Bouguer gravity map of the Waterville, Maine, 15' quadrangle: Maine Geological Survey Open-File Report 87-21, 16 p. and map.

- Rankin, D.W., 1968, Volcanism related to tectonism in the Piscataquis volcanic belt, an island arc of Early Devonian age in north-central Maine, in Zen, E-an, White, W.S., Hadley, J.B., and Thompson, J.B., Jr., eds., Studies in Appalachian geology: Northern and Maritime, Interscience Publishers, New York, p. 355-369.
- _____, 1980, The Traveler Rhyolite and its Devonian setting, Traveler Mountain area, Maine, in Roy, D.C., and Naylor, R.S., eds., Guidebook to the geology of northeastern Maine and neighboring New Brunswick, 72nd New England Intercollegiate Geological Conference, Presque Isle, Maine, p. 98-113.
- Rankin, D.W., Drake, A.A., Jr., Glover, L., III, Goldsmith, R., Hall, L.M., Murray, D.P., Ratcliffe, N.M., Read, J.F., Secor, D.T., Jr., and Stanley, R.S., 1989, Pre-orogenic terranes, in Hatcher, R.D., Jr., Thomas, W.A., and Viele, G.W., eds., The Appalachian-Ouachita Orogen in the United States: Geological Society of America, The Geology of North America, v. F-2, p. 7-100.
- Rast, Nicholas, and Skehan, J.W., S.J., 1983, The evolution of the Avalonian plate: Tectonophysics, v. 100, p. 257-286.
- Sattell, Daniel, and Trehu, A.M., 1989, P and S velocity structure beneath the Gulf of Maine [abs.]: EOS Transactions, American Geophysical Union, v. 70, p. 1204.
- Schenk, P.E., 1991, Events and sea-level changes on Gondwana's margin: The Meguma zone (Cambrian to Devonian) of Nova Scotia, Canada. Geological Society of America Bulletin, v. 103, p. 512-521.
- Schlee, J.S., and Klitgord, K.D., 1988, Georges Bank Basin: A regional synthesis, in Sheridan, R.E., and Grow, J.A., eds.: The Geology of North America, v. I-2, The Atlantic Continental Margin: U.S., p. 243-268.
- Secor, D.T., Jr., Samson, S.L., Snoke, A.W., and Palmer, A.R., 1983, Confirmation of the Carolina Slate Belt as an exotic terrane: Science, v. 221, p. 649-650.
- Shaw, H.F., and Wasserburg, G.J., 1984, Isotopic constraints on the origin of Appalachian mafic complexes: American Journal of Science, v. 284, p. 319-349.
- Snyder, S.L., and Phillips, J.D., 1986, Principal facts for gravity stations along line 4A and 4B of the Quebec-Western Maine seismic reflection profile, west-central Maine: U.S. Geological Survey Open File Report 86-460, 21 p.
- Speed, R.C., Guerrero, J., Rankin, D.W., and Yorath, J.C., 1982, North American Continent-Oceans Transect Program: Geological Society of America, Abstracts with Programs, v. 14, p. 622.
- Spencer, C., Green, A., Morel-a-l'Huissier, P., Luetgert, J.H., Stewart, D.B., Unger, J.D., and Phillips, J.D., 1989, The extension of Grenville basement beneath the northern Appalachians: Results from the Quebec-Maine seismic reflection and refraction surveys: Tectonics, v. 8, p. 677-696.
- St.-Julien, Pierre, and Hubert, Claude, 1975, Evolution of the Taconian orogen in the Quebec Appalachians: American Journal of Science, v. 275-A, p. 337-362.
- St.-Julien, Pierre, and Slivitsky, Anne, 1985, Compilation geologique de la region de l'Estrie-Beauce: Ministere de l'Energie et des Ressources, Quebec, Canada, Resport MM 85-04, map 2030, scale 1:250,000.

- St.-Julien, Pierre, Slivitsky, Anne, and Feininger, Thomas, 1983, A deep structural profile across the Appalachians of southern Quebec: in Hatcher, R.D., Jr., Williams, Harold, and Zeitz, Isidore, eds., Contributions to the Tectonic and Geophysics of Mountain Chains, Geological Society of America Memoir 158, p. 103-111.
- Stanley, R.S., and Ratcliffe, N.M., 1985, Tectonic synthesis of the Taconian orogeny in western New England: Geological Society of America Bulletin, v. 96, p. 1227-1250.
- Steinhart, J.S., Green, R., Asada, T., Rodriquez, B.A., Aldrich, L.T., and Tuve, M.A., 1962a, The earth's crust: The Maine seismic experiment: Carnegie Institution of Washington Yearbook 61, p. 221-234.
- Steinhart, J.S., Meyer, R.P., Howell, B.F., Fahlquist, D.A., Bonini, W.E., and Asada, T., 1962b, Maine seismic experiment: General crustal results [abs]: Journal of Geophysical Research, v. 67, p. 3601.
- Stewart, D.B., 1974, Precambrian Rocks of Seven Hundred Acre Island and development of cleavage in the Islesboro Formation, in Osberg, P.H., ed., Guidebook for trips in east-central and north-central Maine, 66th New England Intercollegiate Geological Conference, Orono, Maine, p. 86-96.
- _____, 1989, Crustal processes in Maine: American Mineralogist, v. 74, p. 698-714.
- Stewart, D.B., and Luetgert, J.H., 1990, The formation of the lower crust beneath the Quebec-Maine-Gulf of Maine Global Geoscience Transect: Geological Society of America Abstracts with Programs, v. 22, no. 7, p. A192.
- Stewart, D.B., and Lux, D.R., 1988, Lithologies and metamorphic age of the Precambrian rocks of Seven Hundred Acre Island and vicinity, Penobscot Bay, Maine, [abs.]: Geological Society of America Abstracts with Programs, v. 20, p. 73.
- Stewart, D.B., Unger, J.D., Phillips, J.D., Goldsmith, R., Poole, W.H., Spencer, C.P., Green, A.G., Loiselle, M.C., and St.-Julien, P., 1986, The Quebec-Western Maine seismic reflection profile: Setting and first year results, in Barazangi, M., and Brown, L., eds., Reflection seismology: The Continental Crust: American Geophysical Union, Geodynamics Series, v. 14, p. 189-199.
- Stewart, D.B., Unger, J.D., and Phillips, J.D., 1985, Quebec-western Maine seismic reflection profile 3B across the coastal volcanic belt, Maine [abs.]: Geological Society of America, Abstracts with Programs, v. 17, p. 64.
- Suzuki, Z., 1965, Maine seismic experiment: A study of shear waves: Bulletin Seismological Society of America, v. 55, p. 425-439.
- Taylor, S.R., 1989, Geophysical framework of the Appalachians and adjacent Grenville province, in Pakiser, L.C., and Mooney, W.D., eds., Geophysical Framework of the Continental United States: Geological Society of America Memoir 172, p. 317-348.
- Thompson, J.B., Jr., Bothner, W.A., Robinson, Peter, Isachsen, Y.W., and Klitgord, K.D., in press, A guide to Continent-Ocean Transect E-1, Adirondacks to Georges Bank: Geological Society of America Centennial Continent/Ocean Transect.
- Thompson, W.B., and Borns, H.W., Jr., 1985, Surficial geologic map of Maine: Maine Geological Survey, Augusta, Maine, scale 1:500,000.

- Trehu, A.M., Ballard, A., Dorman, L.M., Gettrust, J.F., Klitgord, K.D., and Schreiner, A., 1989, Structure of the lower crust beneath the Carolina trough, U.S. Atlantic continental margin. *Journal of Geophysical Research*, v. 94, p. 10585-10600.
- Trzcienski, W.E., Jr., and Marchildon, Nathalie, 1989, Kyanite-garnet-bearing Cambrian rocks and Grenville granulites from the Ayer's Cliff, Quebec, Canada, lamprophyre dike suite: Deep crustal fragments from the northern Appalachians: *Geology*, v. 17, p. 637-640.
- Tucker, R.D., and Robinson, Peter, 1990, Age and setting of the Bronson Hill magmatic arc: A re-evaluation based on U-Pb zircon ages in southern New England: *Geological Society of America Bulletin*, v. 102, p. 1404-1419.
- Unger, J.D., 1988a, A PC program for migration of seismic reflection profiles: *Geobyte*, August 1988, p. 42-54.
- _____, 1988b, Orientation of the decollement surface beneath the Chain Lakes massif: Results of a 3-D seismic survey: *Geological Society of America, Abstracts with Programs*, v. 20, p. 77.
- Unger, J.D., Stewart, D.B., and Phillips, J.D., 1987, Interpretation of migrated seismic reflection profiles across the northern Appalachians in Maine: *Geophysical Journal Royal Astronomical Society*, v. 89, p. 171-176.
- Unger, J.D., Liberty, L.M., Phillips, J.D., and Wright, B.E., 1989, Creating a 3-dimensional transect of the earth's crust from craton to ocean basin across the North Appalachian Orogen, *in* Raper, Johnathan, ed., *Three dimensional applications in Geographic Information Systems*: Taylor and Francis, London, Ch. 10, p. 137-148.
- U.S. Geodynamics Committee, 1989, North American Continent-Ocean Transects Program: National Academy Press, Washington, DC, 87 p.
- Van der Pluijm, Ben, and Van Staal, C.R., 1988, Characteristics and evolution of the central mobile belt, Canadian Appalachians: *Journal of Geology*, v. 96, p. 535-547.
- Walcott, C.D., 1889, Stratigraphic position of the Olenellus faunas in North America and Europe: *American Journal of Science*, v. 38, p. 29-42.
- _____, 1891, Correlation papers--Cambrian: *U.S. Geological Survey Bulletin* 81, 447 p.
- Weed, E.G.A., Minard, J.P., Perry, W.J., Jr., Rhodehamel, E.C., and Robbins, E.I., 1974, Generalized Pre-Pleistocene Geologic Map of the northern United States: Atlantic Continental Margin. U.S. Geological Survey, Miscellaneous Investigations Series Map I-861, scale 1:1,000,000.
- West, D.P., Jr., Lux, D.R., Guidotti, C.V., Hussey, A.M., II, and Newberg, D.W., 1988, $^{40}\text{Ar}/^{39}\text{Ar}$ mineral ages from the Casco Bay Group, southwestern Maine [abs.]: *Geological Society of America Abstracts with Programs*, v. 20, p. 78.
- Williams, Harold, and St.-Julien, Pierre, 1982, The Baie Verte-Brompton Line: Early Paleozoic continent-ocean interface in the Canadian Appalachians, *in* Major Structural Zones and Faults of the Northern Appalachians, St.-Julien, Pierre, and Beland, Jacques, eds., *Geological Association of Canada Special Paper* 24, p. 177-207.
- Wones, D.R., 1978, Norumbega Fault zone, Maine, *in* Seiders, Wanda, and Thomson, Jean, compilers, U.S. Geological Survey, Office of Earthquake Studies, *Summaries of Technical Reports*, v. 7, p. 108-111.
- Wright, B.E., and Stewart, D.B., 1990, Digitization of a geologic map for the Quebec-Maine-Gulf of Maine Global Geoscience Transect: *U.S. Geological Survey Circular* 1041, 16 p.

- Zen, E-an, 1983, Exotic terranes in the New England Appalachians--limits, candidates, and ages: A speculative essay: in Hatcher, R.D., Jr., Williams, Harold, and Zeitz Isidore, eds., Contributions to the Tectonics and Geophysics of Mountain Chains, Geological Society of America Memoir 158, p. 55-81.
- ____ 1989, Tectonostratigraphic terranes in the Northern Appalachians: Their distribution, origin and age, evidence for their existence: Field Trip Guidebook T359, 28th International Geological Congress: American Geophysical Union, Washington, D.C., 68 p.
- ____ 1991, Phanerozoic denudation history of the southern New England Appalachians deduced from pressure data: American Journal of Science, v. 291, p. 401-424.
- Zen, E-an, Stewart, D.B., and Fyffe, L.R., 1986, Paleozoic tectonostratigraphic terranes and their boundaries in the mainland northern Appalachians [abs.]: Geological Society of America Abstracts with Programs, v. 18, p. 800.