

- INSET C**
- 67 ●
 - 66 ●
 - 65 ●
 - 64 ●
 - 63 ●
 - 62 ●
 - 73 ●
 - 72 ●
 - 75 ●
 - 74 ●
 - 68 ●
 - 69 ●
 - 70 ●
 - 71 ●



INSET A

INSET B

EXPLANATION

SAMPLED WELL OR SPRING AND SITE NUMBER (TABLE 2) - Range of concentration of radon-222, in picocuries per liter, is indicated

WELL	SPRING
○ <500	○ <500
● 500-1,000	● 500-1,000
⊙ 1,010-10,000	⊙ 1,010-10,000
⦿ >10,000	⦿ >10,000



Base from U.S. Geological Survey, Reno, Nevada, 1957, Revised 1971, Transverse Mercator Projection.

Scale 1:250,000

0 5 10 15 20 Miles

0 5 10 15 20 Kilometers

CONTOUR INTERVAL 200 FEET
WITH SUPPLEMENTARY CONTOURS AT 100 FOOT INTERVALS
NATIONAL GEODETIC VERTICAL DATUM OF 1929

LOCATION OF GROUND-WATER SITES IN THE CARSON, TRUCKEE, AND WALKER RIVER BASINS, NORTHWESTERN NEVADA, WHERE WATER SAMPLES WERE COLLECTED AND ANALYZED FOR RADIOACTIVE CONSTITUENTS