



U.S. DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

GEOLOGIC RADON POTENTIAL OF EPA REGION 10

Alaska Idaho Oregon Washington



OPEN-FILE REPORT 93-292-J

**Prepared in Cooperation with the
U.S. Environmental Protection Agency**



1993

U.S. DEPARTMENT OF THE INTERIOR
U. S. GEOLOGICAL SURVEY
GEOLOGIC RADON POTENTIAL OF EPA REGION 10
Alaska, Idaho, Oregon, and Washington

R. Randall Schumann
EDITOR

OPEN-FILE REPORT 93-292-J

Prepared in cooperation with the U.S. Environmental Protection Agency

1993

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code.

CONTENTS

<u>SECTION</u>	<u>PAGE</u>
1. The USGS/EPA State Radon Potential Assessments: An Introduction <i>Linda C.S. Gundersen, R. Randall Schumann, and Sharon W. White</i>	1
Appendix A: Geologic Time Scale	19
Appendix B: Glossary of Terms	20
Appendix C: EPA Regional Offices, State Radon Contacts, and State Geological Surveys	26
2. EPA Region 10 Geologic Radon Potential Summary <i>James K. Otton, Kendell A. Dickinson, Douglass E. Owen, and Sandra L. Szarzi</i>	36
3. Preliminary Geologic Radon Potential Assessment of Alaska <i>Kendell A. Dickinson</i>	47
4. Preliminary Geologic Radon Potential Assessment of Idaho <i>Douglass E. Owen</i>	78
5. Preliminary Geologic Radon Potential Assessment of Oregon <i>James K. Otton</i>	103
6. Preliminary Geologic Radon Potential Assessment of Washington <i>James K. Otton</i>	125

ACKNOWLEDGMENTS

These reports are the culmination of a 3-year cooperative project involving, almost literally, a "cast of thousands." Sharon W. White and R. Thomas Peake of the U.S. Environmental Protection Agency coordinated the project for EPA and provided data, comments, assistance, and support. They did an exemplary job and we appreciate their work and our association with the EPA Radon program. The Association of American State Geologists (AASG), through each of the State geological surveys, provided reviews and information that greatly aided in the preparation of the reports and enhanced the final products. Special thanks are due to the AASG committee, consisting of Walter Schmidt (chairman), Robert Fakundiny, Mark Davis, Michael Mudrey, Jonathan Price, Charles Robertson, and John Rold. We also thank the State radon program representatives, whose comments and technical assistance are greatly appreciated.

Technical assistance in preparation of the maps and materials was ably provided by Michele Killgore, Debra Mickelson, Michele Murray, Mark Pyle, and Sandra Szarzi. We thank Helen Britton, Carole Buntenbah, Marge Cunningham, Marian Nance, and Gwen Pilcher for their assistance in manuscript processing. The project chiefs, Linda Gundersen and Randall Schumann, express thanks to those named here and to many others unnamed, but appreciated nonetheless.

THE USGS/EPA RADON POTENTIAL ASSESSMENTS: AN INTRODUCTION

by

Linda C.S. Gundersen and R. Randall Schumann
U.S. Geological Survey

and

Sharon W. White
U.S. Environmental Protection Agency

BACKGROUND

The Indoor Radon Abatement Act of 1988 (Public Law 100-551) directed the U.S. Environmental Protection Agency (EPA) to identify areas of the United States that have the potential to produce harmful levels of indoor radon. These characterizations were to be based on both geological data and on indoor radon levels in homes and other structures. The EPA also was directed to develop model standards and techniques for new building construction that would provide adequate prevention or mitigation of radon entry. As part of an Interagency Agreement between the EPA and the U.S. Geological Survey (USGS), the USGS has prepared radon potential estimates for the United States. This report is one of ten booklets that document this effort. The purpose and intended use of these reports is to help identify areas where states can target their radon program resources, to provide guidance in selecting the most appropriate building code options for areas, and to provide general information on radon and geology for each state for federal, state, and municipal officials dealing with radon issues. *These reports are not intended to be used as a substitute for indoor radon testing, and they cannot and should not be used to estimate or predict the indoor radon concentrations of individual homes, building sites, or housing tracts. Elevated levels of indoor radon have been found in every State, and EPA recommends that all homes be tested for indoor radon.*

USGS geologists are the authors of the booklets. The booklets are organized by EPA Federal boundaries (Regions). Each Regional booklet consists of several components, the first being this introduction to the project, including a general discussion of radon (occurrence, transport, etc.), and details concerning the types of data used. The second component is a summary chapter outlining the general geology and geologic radon potential of the EPA Region. The third component is an individual chapter for each state in the Region. Each state chapter discusses the state's specific geographic setting, soils, geologic setting, geologic radon potential, indoor radon data, and a summary outlining the radon potential rankings of geologic areas in the state. A variety of maps are presented in each chapter—geologic, geographic, population, soils, aerial radioactivity, and indoor radon data by county.

Because of constraints on the scales of maps presented in these reports and because the smallest units used to present the indoor radon data are counties, some generalizations have been made in order to estimate the radon potential of each area. Variations in geology, soil characteristics, climatic factors, homeowner lifestyles, and other factors that influence radon concentrations can be quite large within any particular geologic area, so these reports cannot be used to estimate or predict the indoor radon concentrations of individual homes or housing tracts. Within any area of a given geologic radon potential ranking, there are likely to be areas where the radon potential is lower or higher than that assigned to the area as a whole, especially in larger areas such as the large counties in some western states.

In each state chapter, references to additional reports related to radon are listed for the state, and the reader is urged to consult these reports for more detailed information. In most cases the

best sources of information on radon for specific areas are state and local departments of health, state departments responsible for nuclear safety or environmental protection, and U.S. EPA regional offices. More detailed information on state or local geology may be obtained from the state geological surveys. Addresses and telephone numbers of state radon contacts, geological surveys, and EPA regional offices are listed in Appendix C at the end of this chapter.

RADON GENERATION AND TRANSPORT IN SOILS

Radon (^{222}Rn) is produced from the radioactive decay of radium (^{226}Ra), which is, in turn, a product of the decay of uranium (^{238}U) (fig. 1). The half-life of ^{222}Rn is 3.825 days. Other isotopes of radon occur naturally, but, with the exception of thoron (^{220}Rn), which occurs in concentrations high enough to be of concern in a few localized areas, they are less important in terms of indoor radon risk because of their extremely short half-lives and less common occurrence. In general, the concentration and mobility of radon in soil are dependent on several factors, the most important of which are the soil's radium content and distribution, porosity, permeability to gas movement, and moisture content. These characteristics are, in turn, determined by the soil's parent-material composition, climate, and the soil's age or maturity. If parent-material composition, climate, vegetation, age of the soil, and topography are known, the physical and chemical properties of a soil in a given area can be predicted.

As soils form, they develop distinct layers, or horizons, that are cumulatively called the soil profile. The A horizon is a surface or near-surface horizon containing a relative abundance of organic matter but dominated by mineral matter. Some soils contain an E horizon, directly below the A horizon, that is generally characterized by loss of clays, iron, or aluminum, and has a characteristically lighter color than the A horizon. The B horizon underlies the A or E horizon. Important characteristics of B horizons include accumulation of clays, iron oxides, calcium carbonate or other soluble salts, and organic matter complexes. In drier environments, a horizon may exist within or below the B horizon that is dominated by calcium carbonate, often called caliche or calcrete. This carbonate-cemented horizon is designated the K horizon in modern soil classification schemes. The C horizon underlies the B (or K) and is a zone of weathered parent material that does not exhibit characteristics of A or B horizons; that is, it is generally not a zone of leaching or accumulation. In soils formed in place from the underlying bedrock, the C horizon is a zone of unconsolidated, weathered bedrock overlying the unweathered bedrock.

The shape and orientation of soil particles (soil structure) control permeability and affect water movement in the soil. Soils with blocky or granular structure have roughly equivalent permeabilities in the horizontal and vertical directions, and air and water can infiltrate the soil relatively easily. However, in soils with platy structure, horizontal permeability is much greater than vertical permeability, and air and moisture infiltration is generally slow. Soils with prismatic or columnar structure have dominantly vertical permeability. Platy and prismatic structures form in soils with high clay contents. In soils with shrink-swell clays, air and moisture infiltration rates and depth of wetting may be limited when the cracks in the surface soil layers swell shut. Clay-rich B horizons, particularly those with massive or platy structure, can form a capping layer that impedes the escape of soil gas to the surface (Schumann and others, 1992). However, the shrinkage of clays can act to open or widen cracks upon drying, thus increasing the soil's permeability to gas flow during drier periods.

Radon transport in soils occurs by two processes: (1) diffusion and (2) flow (Tanner, 1964). Diffusion is the process whereby radon atoms move from areas of higher concentration to




Figure 1. The uranium-238 decay series, showing the half-lives of elements and their modes of decay (after Wanty and Schoen, 1991). α denotes alpha decay, β denotes beta decay.

areas of lower concentration in response to a concentration gradient. Flow is the process by which soil air moves through soil pores in response to differences in pressure within the soil or between the soil and the atmosphere, carrying the radon atoms along with it. Diffusion is the dominant radon transport process in soils of low permeability, whereas flow tends to dominate in highly permeable soils (Sextro and others, 1987). In low-permeability soils, much of the radon may decay before it is able to enter a building because its transport rate is reduced. Conversely, highly permeable soils, even those that are relatively low in radium, such as those derived from some types of glacial deposits, have been associated with high indoor radon levels in Europe and in the northern United States (Akerblom and others, 1984; Kunz and others, 1989; Sextro and others, 1987). In areas of karst topography formed in carbonate rock (limestone or dolomite) environments, solution cavities and fissures can increase soil permeability at depth by providing additional pathways for gas flow.

Not all radium contained in soil grains and grain coatings will result in mobile radon when the radium decays. Depending on where the radium is distributed in the soil, many of the radon atoms may remain imbedded in the soil grain containing the parent radium atom, or become imbedded in adjacent soil grains. The portion of radium that releases radon into the pores and fractures of rocks and soils is called the emanating fraction. When a radium atom decays to radon, the energy generated is strong enough to send the radon atom a distance of about 40 nanometers ($1 \text{ nm} = 10^{-9} \text{ meters}$), or about 2×10^{-6} inches—this is known as alpha recoil (Tanner, 1980). Moisture in the soil lessens the chance of a recoiling radon atom becoming imbedded in an adjacent grain. Because water is more dense than air, a radon atom will travel a shorter distance in a water-filled pore than in an air-filled pore, thus increasing the likelihood that the radon atom will remain in the pore space. Intermediate moisture levels enhance radon emanation but do not significantly affect permeability. However, high moisture levels can significantly decrease the gas permeability of the soil and impede radon movement through the soil.

Concentrations of radon in soils are generally many times higher than those inside of buildings, ranging from tens of pCi/L to more than 100,000 pCi/L, but typically in the range of hundreds to low thousands of pCi/L. Soil-gas radon concentrations can vary in response to variations in climate and weather on hourly, daily, or seasonal time scales. Schumann and others (1992) and Rose and others (1988) recorded order-of-magnitude variations in soil-gas radon concentrations between seasons in Colorado and Pennsylvania. The most important factors appear to be (1) soil moisture conditions, which are controlled in large part by precipitation; (2) barometric pressure; and (3) temperature. Washington and Rose (1990) suggest that temperature-controlled partitioning of radon between water and gas in soil pores also has a significant influence on the amount of mobile radon in soil gas.

Homes in hilly limestone regions of the southern Appalachians were found to have higher indoor radon concentrations during the summer than in the winter. A suggested cause for this phenomenon involves temperature/pressure-driven flow of radon-laden air from subsurface solution cavities in the carbonate rock into houses. As warm air enters solution cavities that are higher on the hillslope than the homes, it cools and settles, pushing radon-laden air from lower in the cave or cavity system into structures on the hillslope (Gammage and others, 1993). In contrast, homes built over caves having openings situated below the level of the home had higher indoor radon levels in the winter, caused by cooler outside air entering the cave, driving radon-laden air into cracks and solution cavities in the rock and soil, and ultimately, into homes (Gammage and others, 1993).

RADON ENTRY INTO BUILDINGS

A driving force (reduced atmospheric pressure in the house relative to the soil, producing a pressure gradient) and entry points must exist for radon to enter a building from the soil. The negative pressure caused by furnace combustion, ventilation devices, and the stack effect (the rising and escape of warm air from the upper floors of the building, causing a temperature and pressure gradient within the structure) during cold winter months are common driving forces. Cracks and other penetrations through building foundations, sump holes, and slab-to-foundation wall joints are common entry points.

Radon levels in the basement are generally higher than those on the main floor or upper floors of most structures. Homes with basements generally provide more entry points for radon, commonly have a more pronounced stack effect, and typically have lower air pressure relative to the surrounding soil than nonbasement homes. The term "nonbasement" applies to slab-on-grade or crawl space construction.

METHODS AND SOURCES OF DATA

The assessments of radon potential in the booklets that follow this introduction were made using five main types of data: (1) geologic (lithologic); (2) aerial radiometric; (3) soil characteristics, including soil moisture, permeability, and drainage characteristics; (4) indoor radon data; and (5) building architecture (specifically, whether homes in each area are built slab-on-grade or have a basement or crawl space). These five factors were evaluated and integrated to produce estimates of radon potential. Field measurements of soil-gas radon or soil radioactivity were not used except where such data were available in existing, published reports of local field studies. Where applicable, such field studies are described in the individual state chapters.

GEOLOGIC DATA

The types and distribution of lithologic units and other geologic features in an assessment area are of primary importance in determining radon potential. Rock types that are most likely to cause indoor radon problems include carbonaceous black shales, glauconite-bearing sandstones, certain kinds of fluvial sandstones and fluvial sediments, phosphorites, chalk, karst-producing carbonate rocks, certain kinds of glacial deposits, bauxite, uranium-rich granitic rocks, metamorphic rocks of granitic composition, silica-rich volcanic rocks, many sheared or faulted rocks, some coals, and certain kinds of contact metamorphosed rocks. Rock types least likely to cause radon problems include marine quartz sands, non-carbonaceous shales and siltstones, certain kinds of clays, silica-poor metamorphic and igneous rocks, and basalts. Exceptions exist within these general lithologic groups because of the occurrence of localized uranium deposits, commonly of the hydrothermal type in crystalline rocks or the "roll-front" type in sedimentary rocks. Uranium and radium are commonly sited in heavy minerals, iron-oxide coatings on rock and soil grains, and organic materials in soils and sediments. Less common are uranium associated with phosphate and carbonate complexes in rocks and soils, and uranium minerals.

Although many cases of elevated indoor radon levels can be traced to high radium and (or) uranium concentrations in parent rocks, some structural features, most notably faults and shear zones, have been identified as sites of localized uranium concentrations (Deffeyes and MacGregor, 1980) and have been associated with some of the highest reported indoor radon levels (Gundersen,

1991). The two highest known indoor radon occurrences are associated with sheared fault zones in Boyertown, Pennsylvania (Gundersen and others, 1988a; Smith and others, 1987), and in Clinton, New Jersey (Henry and others, 1991; Muessig and Bell, 1988).

NURE AERIAL RADIOMETRIC DATA

Aerial radiometric data are used to quantify the radioactivity of rocks and soils. Equivalent uranium (eU) data provide an estimate of the surficial concentrations of radon parent materials (uranium, radium) in rocks and soils. Equivalent uranium is calculated from the counts received by a gamma-ray detector from the 1.76 MeV (mega-electron volts) emission energy corresponding to bismuth-214 (^{214}Bi), with the assumption that uranium and its decay products are in secular equilibrium. Equivalent uranium is expressed in units of parts per million (ppm). Gamma radioactivity also may be expressed in terms of a radium activity; 3 ppm eU corresponds to approximately 1 picocurie per gram (pCi/g) of radium-226. Although radon is highly mobile in soil and its concentration is affected by meteorological conditions (Kovach, 1945; Klusman and Jaacks, 1987; Schery and others, 1984; Schumann and others, 1992), statistical correlations between average soil-gas radon concentrations and average eU values for a wide variety of soils have been documented (Gundersen and others, 1988a, 1988b; Schumann and Owen, 1988). Aerial radiometric data can provide an estimate of radon source strength over a region, but the amount of radon that is able to enter a home from the soil is dependent on several local factors, including soil structure, grain size distribution, moisture content, and permeability, as well as type of house construction and its structural condition.

The aerial radiometric data used for these characterizations were collected as part of the Department of Energy National Uranium Resource Evaluation (NURE) program of the 1970s and early 1980s. The purpose of the NURE program was to identify and describe areas in the United States having potential uranium resources (U.S. Department of Energy, 1976). The NURE aerial radiometric data were collected by aircraft in which a gamma-ray spectrometer was mounted, flying approximately 122 m (400 ft) above the ground surface. The equivalent uranium maps presented in the state chapters were generated from reprocessed NURE data in which smoothing, filtering, recalibrating, and matching of adjacent quadrangle data sets were performed to compensate for background, altitude, calibration, and other types of errors and inconsistencies in the original data set (Duval and others, 1989). The data were then gridded and contoured to produce maps of eU with a pixel size corresponding to approximately 2.5 x 2.5 km (1.6 x 1.6 mi).

Figure 2 is an index map of NURE 1° x 2° quadrangles showing the flight-line spacing for each quadrangle. In general, the more closely spaced the flightlines are, the more area was covered by the aerial gamma survey, and thus, more detail is available in the data set. For an altitude of 400 ft above the ground surface and with primary flightline spacing typically between 3 and 6 miles, less than 10 percent of the ground surface of the United States was actually measured by the airborne gamma-ray detectors (Duval and others, 1989), although some areas had better coverage than others due to the differences in flight-line spacing between areas (fig. 2). This suggests that some localized uranium anomalies may not have been detected by the aerial surveys, but the good correlations of eU patterns with geologic outcrop patterns indicate that, at relatively small scales (approximately 1:1,000,000 or smaller) the National eU map (Duval and others, 1989) gives reasonably good estimates of average surface uranium concentrations and thus can assist in the prediction of radon potential of rocks and soils, especially when augmented with additional geologic and soil data.

FLIGHT LINE SPACING OF NURE AERIAL SURVEYS




Figure 2. Nominal flightline spacings for NURE aerial gamma-ray surveys covering the contiguous United States (from Duval and others, 1990). Rectangles represent 1°x2° quadrangles.

The shallow (20-30 cm) depth of investigation of gamma-ray spectrometers, either ground-based or airborne (Duval and others, 1971; Durrance, 1986), suggests that gamma-ray data may sometimes underestimate the radon-source strength in soils in which some of the radionuclides in the near-surface soil layers have been transported downward through the soil profile. In such cases the concentration of radioactive minerals in the A horizon would be lower than in the B horizon, where such minerals are typically concentrated. The concentration of radionuclides in the C horizon and below may be relatively unaffected by surface solution processes. Under these conditions the surface gamma-ray signal may indicate a lower radon source concentration than actually exists in the deeper soil layers, which are most likely to affect radon levels in structures with basements. The redistribution of radionuclides in soil profiles is dependent on a combination of climatic, geologic, and geochemical factors. There is reason to believe that correlations of eU with actual soil radium and uranium concentrations at a depth relevant to radon entry into structures may be regionally variable (Duval, 1989; Schumann and Gundersen, 1991). Given sufficient understanding of the factors cited above, these regional differences may be predictable.

SOIL SURVEY DATA

Soil surveys prepared by the U.S. Soil Conservation Service (SCS) provide data on soil characteristics, including soil-cover thickness, grain-size distribution, permeability, shrink-swell potential, vegetative cover, generalized groundwater characteristics, and land use. The reports are available in county formats and State summaries. The county reports typically contain both generalized and detailed maps of soils in the area.

Because of time and map-scale constraints, it was impractical to examine county soil reports for each county in the United States, so more generalized summaries at appropriate scales were used where available. For State or regional-scale radon characterizations, soil maps were compared to geologic maps of the area, and the soil descriptions, shrink-swell potential, drainage characteristics, depth to seasonal high water table, permeability, and other relevant characteristics of each soil group noted. Technical soil terms used in soil surveys are generally complex; however, a good summary of soil engineering terms and the national distribution of technical soil types is the "Soils" sheet of the National Atlas (U.S. Department of Agriculture, 1987).

Soil permeability is commonly expressed in SCS soil surveys in terms of the speed, in inches per hour (in/hr), at which water soaks into the soil, as measured in a soil percolation test. Although in/hr are not truly units of permeability, these units are in widespread use and are referred to as "permeability" in SCS soil surveys. The permeabilities listed in the SCS surveys are for water, but they generally correlate well with gas permeability. Because data on gas permeability of soils is extremely limited, data on permeability to water is used as a substitute except in cases in which excessive soil moisture is known to exist. Water in soil pores inhibits gas transport, so the amount of radon available to a home is effectively reduced by a high water table. Areas likely to have high water tables include river valleys, coastal areas, and some areas overlain by deposits of glacial origin (for example, loess).

Soil permeabilities greater than 6.0 in/hr may be considered high, and permeabilities less than 0.6 in/hr may be considered low in terms of soil-gas transport. Soils with low permeability may generally be considered to have a lower radon potential than more permeable soils with similar radium concentrations. Many well-developed soils contain a clay-rich B horizon that may impede vertical soil gas transport. Radon generated below this horizon cannot readily escape to the

surface, so it would instead tend to move laterally, especially under the influence of a negative pressure exerted by a building.

Shrink-swell potential is an indicator of the abundance of smectitic (swelling) clays in a soil. Soils with a high shrink-swell potential may cause building foundations to crack, creating pathways for radon entry into the structure. During dry periods, desiccation cracks in shrink-swell soils provide additional pathways for soil-gas transport and effectively increase the gas permeability of the soil. Soil permeability data and soil profile data thus provide important information for regional radon assessments.

INDOOR RADON DATA

Two major sources of indoor radon data were used. The first and largest source of data is from the State/EPA Residential Radon Survey (Ronca-Battista and others, 1988; Dziuban and others, 1990). Forty-two states completed EPA-sponsored indoor radon surveys between 1986 and 1992 (fig. 3). The State/EPA Residential Radon Surveys were designed to be comprehensive and statistically significant at the state level, and were subjected to high levels of quality assurance and control. The surveys collected screening indoor radon measurements, defined as 2-7 day measurements using charcoal canister radon detectors placed in the lowest livable area of the home. The target population for the surveys included owner-occupied single family, detached housing units (White and others, 1989), although attached structures such as duplexes, townhouses, or condominiums were included in some of the surveys if they met the other criteria and had contact with the ground surface. Participants were selected randomly from telephone-directory listings. In total, approximately 60,000 homes were tested in the State/EPA surveys.

The second source of indoor radon data comes from residential surveys that have been conducted in a specific state or region of the country (e.g. independent state surveys or utility company surveys). Several states, including Delaware, Florida, Illinois, New Hampshire, New Jersey, New York, Oregon, and Utah, have conducted their own surveys of indoor radon. The quality and design of a state or other independent survey are discussed and referenced where the data are used.

Data for only those counties with five or more measurements are shown in the indoor radon maps in the state chapters, although data for all counties with a nonzero number of measurements are listed in the indoor radon data tables in each state chapter. In total, indoor radon data from more than 100,000 homes nationwide were used in the compilation of these assessments. Radon data from State or regional indoor radon surveys, public health organizations, or other sources are discussed in addition to the primary data sources where they are available. Nearly all of the data used in these evaluations represent short-term (2-7 day) screening measurements from the lowest livable space of the homes. Specific details concerning the nature and use of indoor radon data sets other than the State/EPA Residential Radon Survey are discussed in the individual State chapters.

RADON INDEX AND CONFIDENCE INDEX

Many of the geologic methods used to evaluate an area for radon potential require subjective opinions based on the professional judgment and experience of the individual geologist. The evaluations are nevertheless based on established scientific principles that are universally applicable to any geographic area or geologic setting. This section describes the methods and conceptual framework used by the U.S. Geological Survey to evaluate areas for radon potential

STATE/EPA RESIDENTIAL RADON SURVEY SCREENING MEASUREMENTS




Figure 3. Percent of homes tested in the State/EPA Residential Radon Survey with screening indoor radon levels exceeding 4 pCi/L.

based on the five factors discussed in the previous sections. The scheme is divided into two basic parts, a Radon Index (RI), used to rank the general radon potential of the area, and the Confidence Index (CI), used to express the level of confidence in the prediction based on the quantity and quality of the data used to make the determination. This scheme works best if the areas to be evaluated are delineated by geologically-based boundaries (geologic provinces) rather than political ones (state/county boundaries) in which the geology may vary across the area.

Radon Index. Table 1 presents the Radon Index (RI) matrix. The five factors—indoor radon data, geology, aerial radioactivity, soil parameters, and house foundation type—were quantitatively ranked (using a point value of 1, 2, or 3) for their respective contribution to radon potential in a given area. At least some data for the 5 factors are consistently available for every geologic province. Because each of these main factors encompass a wide variety of complex and variable components, the geologists performing the evaluation relied heavily on their professional judgment and experience in assigning point values to each category and in determining the overall radon potential ranking. Background information on these factors is discussed in more detail in the preceding sections of this introduction.

Indoor radon was evaluated using unweighted arithmetic means of the indoor radon data for each geologic area to be assessed. Other expressions of indoor radon levels in an area also could have been used, such as weighted averages or annual averages, but these types of data were not consistently available for the entire United States at the time of this writing, or the schemes were not considered sufficient to provide a means of consistent comparison across all areas. For this report, charcoal-canister screening measurement data from the State/EPA Residential Radon Surveys and other carefully selected sources were used, as described in the preceding section. To maintain consistency, other indoor radon data sets (vendor, state, or other data) were not considered in scoring the indoor radon factor of the Radon Index if they were not randomly sampled or could not be statistically combined with the primary indoor radon data sets. However, these additional radon data sets can provide a means to further refine correlations between geologic factors and radon potential, so they are included as supplementary information and are discussed in the individual State chapters. If the average screening indoor radon level for an area was less than 2 pCi/L, the indoor radon factor was assigned 1 point, if it was between 2 and 4 pCi/L, it was scored 2 points, and if the average screening indoor radon level for an area was greater than 4 pCi/L, the indoor radon factor was assigned 3 RI points.

Aerial radioactivity data used in this report are from the equivalent uranium map of the conterminous United States compiled from NURE aerial gamma-ray surveys (Duval and others, 1989). These data indicate the gamma radioactivity from approximately the upper 30 cm of rock and soil, expressed in units of ppm equivalent uranium. An approximate average value of eU was determined visually for each area and point values assigned based on whether the overall eU for the area falls below 1.5 ppm (1 point), between 1.5 and 2.5 ppm (2 points), or greater than 2.5 ppm (3 points).

The geology factor is complex and actually incorporates many geologic characteristics. In the matrix, "positive" and "negative" refer to the presence or absence and distribution of rock types known to have high uranium contents and to generate elevated radon in soils or indoors. Examples of "positive" rock types include granites, black shales, phosphatic rocks, and other rock types described in the preceding "geologic data" section. Examples of "negative" rock types include marine quartz sands and some clays. The term "variable" indicates that the geology within the region is variable or that the rock types in the area are known or suspected to generate elevated radon in some areas but not in others due to compositional differences, climatic effects, localized

TABLE 1. RADON INDEX MATRIX. "ppm eU" indicates parts per million of equivalent uranium, as indicated by NURE aerial radiometric data. See text discussion for details.

FACTOR	<div> <div>INCREASING RADON POTENTIAL</div> <div>→</div> </div> POINT VALUE		
	1	2	3
INDOOR RADON (average)	< 2 pCi/L	2 - 4 pCi/L	> 4 pCi/L
AERIAL RADIOACTIVITY	< 1.5 ppm eU	1.5 - 2.5 ppm eU	> 2.5 ppm eU
GEOLOGY*	negative	variable	positive
SOIL PERMEABILITY	low	moderate	high
ARCHITECTURE TYPE	mostly slab	mixed	mostly basement

*GEOLOGIC FIELD EVIDENCE (GFE) POINTS: GFE points are assigned in addition to points for the "Geology" factor for specific, relevant geologic field studies. See text for details.

Geologic evidence supporting:	HIGH radon	+2 points
	MODERATE	+1 point
	LOW	-2 points
No relevant geologic field studies		0 points

SCORING:

Radon potential category	Point range	Probable average screening indoor radon for area
LOW	3-8 points	< 2 pCi/L
MODERATE/VARIABLE	9-11 points	2 - 4 pCi/L
HIGH	12-17 points	> 4 pCi/L

POSSIBLE RANGE OF POINTS = 3 to 17

TABLE 2. CONFIDENCE INDEX MATRIX

FACTOR	<div> <div>INCREASING CONFIDENCE</div> <div>→</div> </div> POINT VALUE		
	1	2	3
INDOOR RADON DATA	sparse/no data	fair coverage/quality	good coverage/quality
AERIAL RADIOACTIVITY	questionable/no data	glacial cover	no glacial cover
GEOLOGIC DATA	questionable	variable	proven geol. model
SOIL PERMEABILITY	questionable/no data	variable	reliable, abundant

SCORING:	LOW CONFIDENCE	4 - 6 points
	MODERATE CONFIDENCE	7 - 9 points
	HIGH CONFIDENCE	10 - 12 points

POSSIBLE RANGE OF POINTS = 4 to 12

distribution of uranium, or other factors. Geologic information indicates not only how much uranium is present in the rocks and soils but also gives clues for predicting general radon emanation and mobility characteristics through additional factors such as structure (notably the presence of faults or shears) and geochemical characteristics (for example, a phosphate-rich sandstone will likely contain more uranium than a sandstone containing little or no phosphate because the phosphate forms chemical complexes with uranium). "Negative", "variable", and "positive" geology were assigned 1, 2, and 3 points, respectively.

In cases where additional reinforcing or contradictory geologic evidence is available, Geologic Field Evidence (GFE) points were added to or subtracted from an area's score (Table 1). Relevant geologic field studies are important to enhancing our understanding of how geologic processes affect radon distribution. In some cases, geologic models and supporting field data reinforced an already strong (high or low) score; in others, they provided important contradictory data. GFE points were applied for geologically-sound evidence that supports the prediction (but which may contradict one or more factors) on the basis of known geologic field studies in the area or in areas with geologic and climatic settings similar enough that they could be applied with full confidence. For example, areas of the Dakotas, Minnesota, and Iowa that are covered with Wisconsin-age glacial deposits exhibit a low aerial radiometric signature and score only one RI point in that category. However, data from geologic field studies in North Dakota and Minnesota (Schumann and others, 1991) suggest that eU is a poor predictor of geologic radon potential in this area because radionuclides have been leached from the upper soil layers but are present and possibly even concentrated in deeper soil horizons, generating significant soil-gas radon. This positive supporting field evidence adds two GFE points to the score, which helps to counteract the invalid conclusion suggested by the radiometric data. No GFE points are awarded if there are no documented field studies for the area.

"Soil permeability" refers to several soil characteristics that influence radon concentration and mobility, including soil type, grain size, structure, soil moisture, drainage, slope, and permeability. In the matrix, "low" refers to permeabilities less than about 0.6 in/hr; "high" corresponds to greater than about 6.0 in/hr, in U.S. Soil Conservation Service (SCS) standard soil percolation tests. The SCS data are for water permeability, which generally correlates well with the gas permeability of the soil except when the soil moisture content is very high. Areas with consistently high water tables were thus considered to have low gas permeability. "Low", "moderate", and "high" permeability were assigned 1, 2, and 3 points, respectively.

Architecture type refers to whether homes in the area have mostly basements (3 points), mostly slab-on-grade construction (1 point), or a mixture of the two. Split-level and crawl space homes fall into the "mixed" category (2 points). Architecture information is necessary to properly interpret the indoor radon data and produce geologic radon potential categories that are consistent with screening indoor radon data.

The overall RI for an area is calculated by adding the individual RI scores for the 5 factors, plus or minus GFE points, if any. The total RI for an area falls in one of three categories—low, moderate or variable, or high. The point ranges for the three categories were determined by examining the possible combinations of points for the 5 factors and setting rules such that a majority (3 of 5 factors) would determine the final score for the low and high categories, with allowances for possible deviation from an ideal score by the other two factors. The moderate/variable category lies between these two ranges. A total deviation of 3 points from the "ideal" score was considered reasonable to allow for natural variability of factors—if two of the five factors are allowed to vary from the "ideal" for a category, they can differ by a minimum of 2

(1 point different each) and a maximum of 4 points (2 points different each). With "ideal" scores of 5, 10, and 15 points describing low, moderate, and high geologic radon potential, respectively, an ideal low score of 5 points plus 3 points for possible variability allows a maximum of 8 points in the low category. Similarly, an ideal high score of 15 points minus 3 points gives a minimum of 12 points for the high category. Note, however, that if both other factors differ by two points from the "ideal", indicating considerable variability in the system, the total point score would lie in the adjacent (i.e., moderate/variable) category.

Confidence Index. Except for architecture type, the same factors were used to establish a Confidence Index (CI) for the radon potential prediction for each area (Table 2). Architecture type was not included in the confidence index because house construction data are readily and reliably available through surveys taken by agencies and industry groups including the National Association of Home Builders, U.S. Department of Housing and Urban Development, and the Federal Housing Administration; thus it was not considered necessary to question the quality or validity of these data. The other factors were scored on the basis of the quality and quantity of the data used to complete the RI matrix.

Indoor radon data were evaluated based on the distribution and number of data points and on whether the data were collected by random sampling (State/EPA Residential Radon Survey or other state survey data) or volunteered vendor data (likely to be nonrandom and biased toward population centers and/or high indoor radon levels). The categories listed in the CI matrix for indoor radon data ("sparse or no data", "fair coverage or quality", and "good coverage/quality") indicate the sampling density and statistical robustness of an indoor radon data set. Data from the State/EPA Residential Radon Survey and statistically valid state surveys were typically assigned 3 Confidence Index points unless the data were poorly distributed or absent in the area evaluated.

Aerial radioactivity data are available for all but a few areas of the continental United States and for part of Alaska. An evaluation of the quality of the radioactivity data was based on whether there appeared to be a good correlation between the radioactivity and the actual amount of uranium or radium available to generate mobile radon in the rocks and soils of the area evaluated. In general, the greatest problems with correlations among eU, geology, and soil-gas or indoor radon levels were associated with glacial deposits (see the discussion in a previous section) and typically were assigned a 2-point Confidence Index score. Correlations among eU, geology, and radon were generally sound in unglaciated areas and were usually assigned 3 CI points. Again, however, radioactivity data in some unglaciated areas may have been assigned fewer than 3 points, and in glaciated areas may be assigned only one point, if the data were considered questionable or if coverage was poor.

To assign Confidence Index scores for the geologic data factor, rock types and geologic settings for which a physical-chemical, process-based understanding of radon generation and mobility exists were regarded as having "proven geologic models" (3 points); a high confidence could be held for predictions in such areas. Rocks for which the processes are less well known or for which data are contradictory were regarded as "variable" (2 points), and those about which little is known or for which no apparent correlations have been found were deemed "questionable" (1 point).

The soil permeability factor was also scored based on quality and amount of data. The three categories for soil permeability in the Confidence Index are similar in concept, and scored similarly, to those for the geologic data factor. Soil permeability can be roughly estimated from grain size and drainage class if data from standard, accepted soil percolation tests are unavailable; however, the reliability of the data would be lower than if percolation test figures or other

measured permeability data are available, because an estimate of this type does not encompass all the factors that affect soil permeability and thus may be inaccurate in some instances. Most published soil permeability data are for water; although this is generally closely related to the air permeability of the soil, there are some instances when it may provide an incorrect estimate. Examples of areas in which water permeability data may not accurately reflect air permeability include areas with consistently high levels of soil moisture, or clay-rich soils, which would have a low water permeability but may have a significantly higher air permeability when dry due to shrinkage cracks in the soil. These additional factors were applied to the soil permeability factor when assigning the RI score, but may have less certainty in some cases and thus would be assigned a lower CI score.

The Radon Index and Confidence Index give a general indication of the relative contributions of the interrelated geologic factors influencing radon generation and transport in rocks and soils, and thus, of the potential for elevated indoor radon levels to occur in a particular area. However, because these reports are somewhat generalized to cover relatively large areas of States, it is highly recommended that more detailed studies be performed in local areas of interest, using the methods and general information in these booklets as a guide.

EPA COUNTY RADON POTENTIAL MAPS

EPA has produced maps of radon potential, referred to as "radon zone maps", using counties as the primary geographic units. The maps were produced by adapting the results of the geologic radon potential evaluations of the approximately 360 geologic provinces defined for the United States, to fit county boundaries. Because the geologic province boundaries cross State and county boundaries, a strict translation of counties from the geologic province map was not possible. When a county fell within varying radon potential areas, the radon potential designation that covers the most area was chosen as the county designation. The geologic province assessments were adapted to a county map format because many planning, outreach, and information programs are based on political boundaries such as counties. The county-based EPA Radon Zone Maps are not included in the USGS geologic radon potential booklets. They are available from EPA headquarters and regional offices or through the state radon program offices.

REFERENCES CITED

- Akerblom, G., Anderson, P., and Clavensjo, B., 1984, Soil gas radon--A source for indoor radon daughters: *Radiation Protection Dosimetry*, v. 7, p. 49-54.
- Deffeyes, K.S., and MacGregor, I.D., 1980, World uranium resources: *Scientific American*, v. 242, p. 66-76.
- Durrance, E.M., 1986, *Radioactivity in geology: Principles and applications*: New York, N.Y., Wiley and Sons, 441 p.
- Duval, J.S., 1989, Radioactivity and some of its applications in geology: *Proceedings of the symposium on the application of geophysics to engineering and environmental problems (SAGEEP)*, Golden, Colorado, March 13-16, 1989: Society of Engineering and Mineral Exploration Geophysicists, p. 1-61.
- Duval, J.S., Cook, B.G., and Adams, J.A.S., 1971, Circle of investigation of an airborne gamma-ray spectrometer: *Journal of Geophysical Research*, v. 76, p. 8466-8470.
- Duval, J.S., Jones, W.J., Riggle, F.R., and Pitkin, J.A., 1989, Equivalent uranium map of conterminous United States: U.S. Geological Survey Open-File Report 89-478, 10 p.
- Duval, J.S., Reimer, G.M., Schumann, R.R., Owen, D.E., and Otton, J.K., 1990, Soil-gas radon compared to aerial and ground gamma-ray measurements at study sites near Greeley and Fort Collins, Colorado: U.S. Geological Survey Open-File Report 90-648, 42 p.
- Dziuban, J.A., Clifford, M.A., White, S.B., Bergstein, J.W., and Alexander, B.V., 1990, Residential radon survey of twenty-three States, *in* *Proceedings of the 1990 International Symposium on Radon and Radon Reduction Technology*, Vol. III: Preprints: U.S. Environmental Protection Agency report EPA/600/9-90/005c, Paper IV-2, 17 p.
- Gammage, R.B., Wilson, D.L., Saultz, R.J., and Bauer, B.C., 1993, Subterranean transport of radon and elevated indoor radon in hilly karst terranes: *Atmospheric Environment* (in press).
- Gundersen, L.C.S., Reimer, G.M., and Agard, S.S., 1988a, Correlation between geology, radon in soil gas, and indoor radon in the Reading Prong, *in* Marikos, M.A., and Hansman, R.H., eds., *Geologic causes of natural radionuclide anomalies*: Missouri Department of Natural Resources Special Publication 4, p. 91-102.
- Gundersen, L.C.S., Reimer, G.M., Wiggs, C.R., and Rice, C.A., 1988b, Map showing radon potential of rocks and soils in Montgomery County, Maryland: U.S. Geological Survey Miscellaneous Field Studies Map MF-2043, scale 1:62,500.
- Gundersen, Linda C.S., 1991, Radon in sheared metamorphic and igneous rocks, *in* Gundersen, Linda C.S., and Richard B. Wanty, eds., *Field studies of radon in rocks, soils, and water*: U.S. Geol. Survey Bulletin no. 1971, p. 39-50.

- Henry, Mitchell E., Kaeding, Margaret E., and Monteverde, Donald, 1991, Radon in soil gas and gamma-ray activity of rocks and soils at the Mulligan Quarry, Clinton, New Jersey, *in* Gundersen, Linda C.S., and Richard B. Wanty, eds., *Field studies of radon in rocks, soils, and water*: U.S. Geol. Survey Bulletin no. 1971, p. 65-75.
- Klusman, R. W., and Jaacks, J. A., 1987, Environmental influences upon mercury, radon, and helium concentrations in soil gases at a site near Denver, Colorado: *Journal of Geochemical Exploration*, v. 27, p. 259-280.
- Kovach, E.M., 1945, Meteorological influences upon the radon content of soil gas: *Transactions, American Geophysical Union*, v. 26, p. 241-248.
- Kunz, C., Laymon, C.A., and Parker, C., 1989, Gravelly soils and indoor radon, *in* Osborne, M.C., and Harrison, J., eds., *Proceedings of the 1988 EPA Symposium on Radon and Radon Reduction Technology, Volume 1: U.S. Environmental Protection Agency Report EPA/600/9-89/006A*, p. 5-75--5-86.
- Muessig, K., and Bell, C., 1988, Use of airborne radiometric data to direct testing for elevated indoor radon: *Northeastern Environmental Science*, v. 7, no. 1, p. 45-51.
- Ronca-Battista, M., Moon, M., Bergsten, J., White, S.B., Holt, N., and Alexander, B., 1988, Radon-222 concentrations in the United States--Results of sample surveys in five states: *Radiation Protection Dosimetry*, v. 24, p. 307-312.
- Rose, A.W., Washington, J.W., and Greeman, D.J., 1988, Variability of radon with depth and season in a central Pennsylvania soil developed on limestone: *Northeastern Environmental Science*, v. 7, p. 35-39.
- Schery, S.D., Gaeddert, D.H., and Wilkening, M.H., 1984, Factors affecting exhalation of radon from a gravelly sandy loam: *Journal of Geophysical Research*, v. 89, p. 7299-7309.
- Schumann, R.R., and Owen, D.E., 1988, Relationships between geology, equivalent uranium concentration, and radon in soil gas, Fairfax County, Virginia: U.S. Geological Survey Open-File Report 88-18, 28 p.
- Schumann, R.R., and Gundersen, L.C.S., 1991, Regional differences in radon emanation coefficients in soils: *Geological Society of America Abstracts With Programs*, v. 23, no. 1, p. 125.
- Schumann, R.R., Peake, R.T., Schmidt, K.M., and Owen, D.E., 1991, Correlations of soil-gas and indoor radon with geology in glacially derived soils of the northern Great Plains, *in* *Proceedings of the 1990 International Symposium on Radon and Radon Reduction Technology, Volume 2, Symposium Oral Papers*: U.S. Environmental Protection Agency report EPA/600/9-91/026b, p. 6-23--6-36.

- Schumann, R.R., Owen, D.E., and Asher-Bolinder, S., 1992, Effects of weather and soil characteristics on temporal variations in soil-gas radon concentrations, *in* Gates, A.E., and Gundersen, L.C.S., eds., *Geologic controls on radon: Geological Society of America Special Paper 271*, p. 65-72.
- Sextro, R.G., Moed, B.A., Nazaroff, W.W., Revzan, K.L., and Nero, A.V., 1987, Investigations of soil as a source of indoor radon, *in* Hopke, P.K., ed., *Radon and its decay products: American Chemical Society Symposium Series 331*, p. 10-29.
- Sterling, R., Meixel, G., Shen, L., Labs, K., and Bligh, T., 1985, Assessment of the energy savings potential of building foundations research: Oak Ridge, Tenn., U.S. Department of Energy Report ORNL/SUB/84-0024/1.
- Smith, R.C., II, Reilly, M.A., Rose, A.W., Barnes, J.H., and Berkheiser, S.W., Jr., 1987, Radon: a profound case: *Pennsylvania Geology*, v. 18, p. 1-7.
- Tanner, A.B., 1964, Radon migration in the ground: a review, *in* Adams, J.A.S., and Lowder, W.M., eds., *The natural radiation environment: Chicago, Ill., University of Chicago Press*, p. 161-190.
- Tanner, A.B., 1980, Radon migration in the ground: a supplementary review, *in* Gesell, T.F., and Lowder, W.M. (eds), *Natural radiation environment III, Symposium proceedings, Houston, Texas*, v. 1, p. 5-56.
- U.S. Department of Agriculture, 1987, Principal kinds of soils: Orders, suborders, and great groups: U.S. Geological Survey, National Atlas of the United States of America, sheet 38077-BE-NA-07M-00, scale 1:7,500,000.
- U.S. Department of Energy, 1976, National Uranium Resource Evaluation preliminary report, prepared by the U.S. Energy Research and Development Administration, Grand Junction, Colo.: GJO-11(76).
- Wanty, Richard B., and Schoen, Robert, 1991, A review of the chemical processes affecting the mobility of radionuclides in natural waters, with applications, *in* Gundersen, Linda C.S., and Richard B. Wanty, eds., *Field studies of radon in rocks, soils, and water: U.S. Geological Survey Bulletin no. 1971*, p. 183-194.
- Washington, J.W., and Rose, A.W., 1990, Regional and temporal relations of radon in soil gas to soil temperature and moisture: *Geophysical Research Letters*, v. 17, p. 829-832.
- White, S.B., Bergsten, J.W., Alexander, B.V., and Ronca-Battista, M., 1989, Multi-State surveys of indoor ^{222}Rn : *Health Physics*, v. 57, p. 891-896.

APPENDIX A GEOLOGIC TIME SCALE

Subdivisions (and their symbols)						Age estimates of boundaries in mega-annum (Ma) ¹		
Eon or Eonothem	Era or Erathem	Period, System, Subperiod, Subsystem		Epoch or Series				
Phanerozoic ²	Cenozoic ² (Cz)	Quaternary ² (Q)		Holocene		0.010		
				Pleistocene		1.6 (1.6–1.9)		
		Tertiary (T)	Neogene ² Subperiod or Subsystem (N)	Pliocene		5 (4.9–5.3)		
				Miocene		24 (23–26)		
			Paleogene ² Subperiod or Subsystem (Pt)	Oligocene		38 (34–38)		
				Eocene		55 (54–56)		
				Paleocene		66 (63–66)		
						96 (95–97)		
						138 (135–141)		
				Mesozoic ² (Mz)	Cretaceous (K)		Late	Upper
					Early	Lower		
	Jurassic (J)				Late	Upper		
					Middle	Middle		
					Early	Lower		
	Triassic (Tr)				Late	Upper	205 (200–215)	
					Middle	Middle		
					Early	Lower		
							~240	
							290 (290–305)	
	Paleozoic ² (Pz)	Permian (P)		Late	Upper			
				Early	Lower			
		Carboniferous Systems (C)		Pennsylvanian (P)	Late	Upper		
			Middle		Middle			
			Mississippian (M)	Early	Lower			
				Late	Upper	~330		
				Early	Lower	360 (360–365)		
		Devonian (D)		Late	Upper			
				Middle	Middle			
				Early	Lower			
		Silurian (S)		Late	Upper	410 (405–415)		
				Middle	Middle			
				Early	Lower			
		Ordovician (O)		Late	Upper	435 (435–440)		
				Middle	Middle			
				Early	Lower			
		Cambrian (C)		Late	Upper	500 (495–510)		
		Middle	Middle					
		Early	Lower					
Proterozoic (P)		Late Proterozoic (Z)	None defined				~570 ³	
	Middle Proterozoic (Y)	None defined				900		
	Early Proterozoic (X)	None defined				1600		
	Early Proterozoic (X)	None defined				2500		
Archean (A)	Late Archean (W)	None defined				3000		
	Middle Archean (V)	None defined				3400		
	Early Archean (U)	None defined				3800		
	Early Archean (U)	None defined				3800 ?		
pre-Archean (pA) ⁴								

¹ Ranges reflect uncertainties of isotopic and biostratigraphic age assignments. Age boundaries not closely bracketed by existing data shown by ~. Decay constants and isotopic ratios employed are cited in Steiger and Jäger (1977). Designation m.y. used for an interval of time.

² Modifiers (lower, middle, upper or early, middle, late) when used with these items are informal divisions of the larger unit; the first letter of the modifier is lowercase.

³ Rocks older than 570 Ma also called Precambrian (p-C), a time term without specific rank.

⁴ Informal time term without specific rank.

APPENDIX B GLOSSARY OF TERMS

Units of measure

pCi/L (picocuries per liter)- a unit of measure of radioactivity used to describe radon concentrations in a volume of air. One picocurie (10^{-12} curies) is equal to about 2.2 disintegrations of radon atoms per minute. A liter is about 1.06 quarts. The average concentration of radon in U.S. homes measured to date is between 1 and 2 pCi/L.

Bq/m³ (Becquerels per cubic meter)- a metric unit of radioactivity used to describe radon concentrations in a volume of air. One becquerel is equal to one radioactive disintegration per second. One pCi/L is equal to 37 Bq/m³.

ppm (parts per million)- a unit of measure of concentration by weight of an element in a substance, in this case, soil or rock. One ppm of uranium contained in a ton of rock corresponds to about 0.03 ounces of uranium. The average concentration of uranium in soils in the United States is between 1 and 2 ppm.

in/hr (inches per hour)- a unit of measure used by soil scientists and engineers to describe the permeability of a soil to water flowing through it. It is measured by digging a hole 1 foot (12 inches) square and one foot deep, filling it with water, and measuring the time it takes for the water to drain from the hole. The drop in height of the water level in the hole, measured in inches, is then divided by the time (in hours) to determine the permeability. Soils range in permeability from less than 0.06 in/hr to greater than 20 in/hr, but most soils in the United States have permeabilities between these two extremes.

Geologic terms and terms related to the study of radon

aerial radiometric, aeroradiometric survey A survey of radioactivity, usually gamma rays, taken by an aircraft carrying a gamma-ray spectrometer pointed at the ground surface.

alluvial fan A low, widespread mass of loose rock and soil material, shaped like an open fan and deposited by a stream at the point where it flows from a narrow mountain valley out onto a plain or broader valley. May also form at the junction with larger streams or when the gradient of the stream abruptly decreases.

alluvium, alluvial General terms referring to unconsolidated detrital material deposited by a stream or other body of running water.

alpha-track detector A passive radon measurement device consisting of a plastic film that is sensitive to alpha particles. The film is etched with acid in a laboratory after it is exposed. The etching reveals scratches, or "tracks", left by the alpha particles resulting from radon decay, which can then be counted to calculate the radon concentration. Useful for long-term (1-12 months) radon tests.

amphibolite A mafic metamorphic rock consisting mainly of pyroxenes and(or) amphibole and plagioclase.

argillite, argillaceous Terms referring to a rock derived from clay or shale, or any sedimentary rock containing an appreciable amount of clay-size material, i.e., argillaceous sandstone.

arid Term describing a climate characterized by dryness, or an evaporation rate that exceeds the amount of precipitation.

basalt A general term for a dark-colored mafic igneous rocks that may be of extrusive origin, such as volcanic basalt flows, or intrusive origin, such as basalt dikes.

batholith A mass of plutonic igneous rock that has more than 40 square miles of surface exposure and no known bottom.

carbonate A sedimentary rock consisting of the carbonate (CO_3) compounds of calcium, magnesium, or iron, e.g. limestone and dolomite.

carbonaceous Said of a rock or sediment that is rich in carbon, is coaly, or contains organic matter.

charcoal canister A passive radon measurement device consisting of a small container of granulated activated charcoal that is designed to adsorb radon. Useful for short duration (2-7 days) measurements only. May be referred to as a "screening" test.

chert A hard, extremely dense sedimentary rock consisting dominantly of interlocking crystals of quartz. Crystals are not visible to the naked eye, giving the rock a milky, dull luster. It may be white or gray but is commonly colored red, black, yellow, blue, pink, brown, or green.

clastic pertaining to a rock or sediment composed of fragments that are derived from preexisting rocks or minerals. The most common clastic sedimentary rocks are sandstone and shale.

clay A rock containing clay mineral fragments or material of any composition having a diameter less than 1/256 mm.

clay mineral One of a complex and loosely defined group of finely crystalline minerals made up of water, silicate and aluminum (and a wide variety of other elements). They are formed chiefly by alteration or weathering of primary silicate minerals. Certain clay minerals are noted for their small size and ability to absorb substantial amounts of water, causing them to swell. The change in size that occurs as these clays change between dry and wet is referred to as their "**shrink-swell**" potential.

concretion A hard, compact mass of mineral matter, normally subspherical but commonly irregular in shape; formed by precipitation from a water solution about a nucleus or center, such as a leaf, shell, bone, or fossil, within a sedimentary or fractured rock.

conglomerate A coarse-grained, clastic sedimentary rock composed of rock and mineral fragments larger than 2 mm, set in a finer-grained matrix of clastic material.

cuesta A hill or ridge with a gentle slope on one side and a steep slope on the other. The formation of a cuesta is controlled by the different weathering properties and the structural dip of the rocks forming the hill or ridge.

daughter product A nuclide formed by the disintegration of a radioactive precursor or "parent" atom.

delta, deltaic Referring to a low, flat, alluvial tract of land having a triangular or fan shape, located at or near the mouth of a river. It results from the accumulation of sediment deposited by a river at the point at which the river loses its ability to transport the sediment, commonly where a river meets a larger body of water such as a lake or ocean.

dike A tabular igneous intrusion of rock, younger than the surrounding rock, that commonly cuts across the bedding or foliation of the rock it intrudes.

diorite A plutonic igneous rock that is medium in color and contains visible dark minerals that make up less than 50% of the rock. It also contains abundant sodium plagioclase and minor quartz.

dolomite A carbonate sedimentary rock of which more than 50% consists of the mineral dolomite ($\text{CaMg}(\text{CO}_3)_2$), and is commonly white, gray, brown, yellow, or pinkish in color.

drainage The manner in which the waters of an area pass, flow off of, or flow into the soil. Also refers to the water features of an area, such as lakes and rivers, that drain it.

eolian Pertaining to sediments deposited by the wind.

esker A long, narrow, steep-sided ridge composed of irregular beds of sand and gravel deposited by streams beneath a glacier and left behind when the ice melted.

evapotranspiration Loss of water from a land area by evaporation from the soil and transpiration from plants.

extrusive Said of igneous rocks that have been erupted onto the surface of the Earth.

fault A fracture or zone of fractures in rock or sediment along which there has been movement.

fluvial, fluvial deposit Pertaining to sediment that has been deposited by a river or stream.

foliation A linear feature in a rock defined by both mineralogic and structural characteristics. It may be formed during deformation or metamorphism.

formation A mappable body of rock having similar characteristics.

glacial deposit Any sediment transported and deposited by a glacier or processes associated with glaciers, such as glaciofluvial sediments deposited by streams flowing from melting glaciers.

gneiss A rock formed by metamorphism in which bands and lenses of minerals of similar composition alternate with bands and lenses of different composition, giving the rock a striped or "foliated" appearance.

granite Broadly applied, any coarsely crystalline, quartz- and feldspar-bearing igneous plutonic rock. Technically, granites have between 10 and 50% quartz, and alkali feldspar comprises at least 65% of the total feldspar.

gravel An unconsolidated, natural accumulation of rock fragments consisting predominantly of particles greater than 2 mm in size.

heavy minerals Mineral grains in sediment or sedimentary rock having higher than average specific gravity. May form layers and lenses because of wind or water sorting by weight and size

and may be referred to as a "placer deposit." Some heavy minerals are magnetite, garnet, zircon, monazite, and xenotime.

igneous Said of a rock or mineral that solidified from molten or partly molten rock material. It is one of the three main classes into which rocks are divided, the others being sedimentary and metamorphic.

intermontane A term that refers to an area between two mountains or mountain ranges.

intrusion, intrusive The processes of emplacement or injection of molten rock into pre-existing rock. Also refers to the rock formed by intrusive processes, such as an "intrusive igneous rock".

kame A low mound, knob, hummock, or short irregular ridge formed by a glacial stream at the margin of a melting glacier; composed of bedded sand and gravel.

karst terrain A type of topography that is formed on limestone, gypsum and other rocks by dissolution of the rock by water, forming sinkholes and caves.

lignite A brownish-black coal that is intermediate in coalification between peat and subbituminous coal.

limestone A carbonate sedimentary rock consisting of more than 50% calcium carbonate, primarily in the form of the mineral calcite (CaCO_3).

lithology The description of rocks in hand specimen and in outcrop on the basis of color, composition, and grain size.

loam A permeable soil composed of a mixture of relatively equal parts clay, silt, and sand, and usually containing some organic matter.

loess A fine-grained eolian deposit composed of silt-sized particles generally thought to have been deposited from windblown dust of Pleistocene age.

mafic Term describing an igneous rock containing more than 50% dark-colored minerals.

marine Term describing sediments deposited in the ocean, or precipitated from ocean waters.

metamorphic Any rock derived from pre-existing rocks by mineralogical, chemical, or structural changes in response to changes in temperature, pressure, stress, and the chemical environment. Phyllite, schist, amphibolite, and gneiss are metamorphic rocks.

moraine A mound, ridge, or other distinct accumulation of unsorted, unbedded glacial material, predominantly till, deposited by the action of glacial ice.

outcrop That part of a geologic formation or structure that appears at the surface of the Earth, as in "rock outcrop".

percolation test A term used in engineering for a test to determine the water permeability of a soil. A hole is dug and filled with water and the rate of water level decline is measured.

permeability The capacity of a rock, sediment, or soil to transmit liquid or gas.

phosphate, phosphatic, phosphorite Any rock or sediment containing a significant amount of phosphate minerals, i.e., minerals containing PO_4 .

physiographic province A region in which all parts are similar in geologic structure and climate, which has had a uniform geomorphic history, and whose topography or landforms differ significantly from adjacent regions.

placer deposit See heavy minerals

residual Formed by weathering of a material in place.

residuum Deposit of residual material.

rhyolite An extrusive igneous rock of volcanic origin, compositionally equivalent to granite.

sandstone A clastic sedimentary rock composed of sand-sized rock and mineral material that is more or less firmly cemented. Sand particles range from 1/16 to 2 mm in size.

schist A strongly foliated crystalline rock, formed by metamorphism, that can be readily split into thin flakes or slabs. Contains mica; minerals are typically aligned.

screening level Result of an indoor radon test taken with a charcoal canister or similar device, for a short period of time, usually less than seven days. May indicate the potential for an indoor radon problem but does not indicate annual exposure to radon.

sediment Deposits of rock and mineral particles or fragments originating from material that is transported by air, water or ice, or that accumulate by natural chemical precipitation or secretion of organisms.

semiarid Refers to a climate that has slightly more precipitation than an arid climate.

shale A fine-grained sedimentary rock formed from solidification (lithification) of clay or mud.

shear zone Refers to a roughly linear zone of rock that has been faulted by ductile or non-ductile processes in which the rock is sheared and both sides are displaced relative to one another.

shrink-swell clay See clay mineral.

siltstone A fine-grained clastic sedimentary rock composed of silt-sized rock and mineral material and more or less firmly cemented. Silt particles range from 1/16 to 1/256 mm in size.

sinkhole A roughly circular depression in a karst area measuring meters to tens of meters in diameter. It is funnel shaped and is formed by collapse of the surface material into an underlying void created by the dissolution of carbonate rock.

slope An inclined part of the earth's surface.

solution cavity A hole, channel or cave-like cavity formed by dissolution of rock.

stratigraphy The study of rock strata; also refers to the succession of rocks of a particular area.

surficial materials Unconsolidated glacial, wind-, or waterborne deposits occurring on the earth's surface.

tablelands General term for a broad, elevated region with a nearly level surface of considerable extent.

terrace gravel Gravel-sized material that caps ridges and terraces, left behind by a stream as it cuts down to a lower level.

terrain A tract or region of the Earth's surface considered as a physical feature or an ecological environment.

till Unsorted, generally unconsolidated and unbedded rock and mineral material deposited directly adjacent to and underneath a glacier, without reworking by meltwater. Size of grains varies greatly from clay to boulders.

uraniferous Containing uranium, usually more than 2 ppm.

vendor data Used in this report to refer to indoor radon data collected and measured by commercial vendors of radon measurement devices and/or services.

volcanic Pertaining to the activities, structures, and extrusive rock types of a volcano.

water table The surface forming the boundary between the zone of saturation and the zone of aeration; the top surface of a body of unconfined groundwater in rock or soil.

weathering The destructive process by which earth and rock materials, on exposure to atmospheric elements, are changed in color, texture, composition, firmness, or form with little or no transport of the material.

APPENDIX C

EPA REGIONAL OFFICES

EPA Regional Offices	State	EPA Region
EPA Region 1 JFK Federal Building Boston, MA 02203 (617) 565-4502	Alabama.....	4
	Alaska.....	10
	Arizona.....	9
	Arkansas.....	6
	California.....	9
	Colorado.....	8
	Connecticut.....	1
	Delaware.....	3
	District of Columbia.....	3
	Florida.....	4
	Georgia.....	4
	Hawaii.....	9
	Idaho.....	10
	Illinois.....	5
	Indiana.....	5
	Iowa.....	7
	Kansas.....	7
	Kentucky.....	4
	Louisiana.....	6
	Maine.....	1
	Maryland.....	3
	Massachusetts.....	1
	Michigan.....	5
	Minnesota.....	5
	Mississippi.....	4
	Missouri.....	7
	Montana.....	8
	Nebraska.....	7
	Nevada.....	9
	New Hampshire.....	1
	New Jersey.....	2
	New Mexico.....	6
	New York.....	2
	North Carolina.....	4
	North Dakota.....	8
	Ohio.....	5
	Oklahoma.....	6
	Oregon.....	10
	Pennsylvania.....	3
	Rhode Island.....	1
	South Carolina.....	4
	South Dakota.....	8
	Tennessee.....	4
	Texas.....	6
	Utah.....	8
	Vermont.....	1
	Virginia.....	3
	Washington.....	10
	West Virginia.....	3
	Wisconsin.....	5
	Wyoming.....	8
EPA Region 2 (2AIR:RAD) 26 Federal Plaza New York, NY 10278 (212) 264-4110		
Region 3 (3AH14) 841 Chestnut Street Philadelphia, PA 19107 (215) 597-8326		
EPA Region 4 345 Courtland Street, N.E. Atlanta, GA 30365 (404) 347-3907		
EPA Region 5 (5AR26) 77 West Jackson Blvd. Chicago, IL 60604-3507 (312) 886-6175		
EPA Region 6 (6T-AS) 1445 Ross Avenue Dallas, TX 75202-2733 (214) 655-7224		
EPA Region 7 726 Minnesota Avenue Kansas City, KS 66101 (913) 551-7604		
EPA Region 8 (8HWM-RP) 999 18th Street One Denver Place, Suite 1300 Denver, CO 80202-2413 (303) 293-1713		
EPA Region 9 (A-3) 75 Hawthorne Street San Francisco, CA 94105 (415) 744-1048		
EPA Region 10 1200 Sixth Avenue Seattle, WA 98101 (202) 442-7660		

STATE RADON CONTACTS

May, 1993

<u>Alabama</u>	James McNees Division of Radiation Control Alabama Department of Public Health State Office Building Montgomery, AL 36130 (205) 242-5315 1-800-582-1866 in state	<u>Connecticut</u>	Alan J. Siniscalchi Radon Program Connecticut Department of Health Services 150 Washington Street Hartford, CT 06106-4474 (203) 566-3122
<u>Alaska</u>	Charles Tedford Department of Health and Social Services P.O. Box 110613 Juneau, AK 99811-0613 (907) 465-3019 1-800-478-4845 in state	<u>Delaware</u>	Marai G. Rejai Office of Radiation Control Division of Public Health P.O. Box 637 Dover, DE 19903 (302) 736-3028 1-800-554-4636 In State
<u>Arizona</u>	John Stewart Arizona Radiation Regulatory Agency 4814 South 40th St. Phoenix, AZ 85040 (602) 255-4845	<u>District of Columbia</u>	Robert Davis DC Department of Consumer and Regulatory Affairs 614 H Street NW Room 1014 Washington, DC 20001 (202) 727-71068
<u>Arkansas</u>	Lee Gershner Division of Radiation Control Department of Health 4815 Markham Street, Slot 30 Little Rock, AR 72205-3867 (501) 661-2301	<u>Florida</u>	N. Michael Gilley Office of Radiation Control Department of Health and Rehabilitative Services 1317 Winewood Boulevard Tallahassee, FL 32399-0700 (904) 488-1525 1-800-543-8279 in state
<u>California</u>	J. David Quinton Department of Health Services 714 P Street, Room 600 Sacramento, CA 94234-7320 (916) 324-2208 1-800-745-7236 in state	<u>Georgia</u>	Richard Schreiber Georgia Department of Human Resources 878 Peachtree St., Room 100 Atlanta, GA 30309 (404) 894-6644 1-800-745-0037 in state
<u>Colorado</u>	Linda Martin Department of Health 4210 East 11th Avenue Denver, CO 80220 (303) 692-3057 1-800-846-3986 in state	<u>Hawaii</u>	Russell Takata Environmental Health Services Division 591 Ala Moana Boulevard Honolulu, HI 96813-2498 (808) 586-4700

Idaho Pat McGavarn
Office of Environmental Health
450 West State Street
Boise, ID 83720
(208) 334-6584
1-800-445-8647 in state

Louisiana Matt Schlenker
Louisiana Department of
Environmental Quality
P.O. Box 82135
Baton Rouge, LA 70884-2135
(504) 925-7042
1-800-256-2494 in state

Illinois Richard Allen
Illinois Department of Nuclear Safety
1301 Outer Park Drive
Springfield, IL 62704
(217) 524-5614
1-800-325-1245 in state

Maine Bob Stilwell
Division of Health Engineering
Department of Human Services
State House, Station 10
Augusta, ME 04333
(207) 289-5676
1-800-232-0842 in state

Indiana Lorand Magyar
Radiological Health Section
Indiana State Department of Health
1330 West Michigan Street
P.O. Box 1964
Indianapolis, IN 46206
(317) 633-8563
1-800-272-9723 In State

Maryland Leon J. Rachuba
Radiological Health Program
Maryland Department of the
Environment
2500 Broening Highway
Baltimore, MD 21224
(410) 631-3301
1-800-872-3666 In State

Iowa Donald A. Flater
Bureau of Radiological Health
Iowa Department of Public Health
Lucas State Office Building
Des Moines, IA 50319-0075
(515) 281-3478
1-800-383-5992 In State

Massachusetts William J. Bell
Radiation Control Program
Department of Public Health
23 Service Center
Northampton, MA 01060
(413) 586-7525
1-800-445-1255 in state

Kansas Harold Spiker
Radiation Control Program
Kansas Department of Health and
Environment
109 SW 9th Street
6th Floor Mills Building
Topeka, KS 66612
(913) 296-1561

Michigan Sue Hendershott
Division of Radiological Health
Bureau of Environmental and
Occupational Health
3423 North Logan Street
P.O. Box 30195
Lansing, MI 48909
(517) 335-8194

Kentucky Jeana Phelps
Radiation Control Branch
Department of Health Services
Cabinet for Human Resources
275 East Main Street
Frankfort, KY 40601
(502) 564-3700

Minnesota Laura Oatmann
Indoor Air Quality Unit
925 Delaware Street, SE
P.O. Box 59040
Minneapolis, MN 55459-0040
(612) 627-5480
1-800-798-9050 in state

Mississippi Silas Anderson
Division of Radiological Health
Department of Health
3150 Lawson Street
P.O. Box 1700
Jackson, MS 39215-1700
(601) 354-6657
1-800-626-7739 in state

Missouri Kenneth V. Miller
Bureau of Radiological Health
Missouri Department of Health
1730 East Elm
P.O. Box 570
Jefferson City, MO 65102
(314) 751-6083
1-800-669-7236 In State

Montana Adrian C. Howe
Occupational Health Bureau
Montana Department of Health and
Environmental Sciences
Cogswell Building A113
Helena, MT 59620
(406) 444-3671

Nebraska Joseph Milone
Division of Radiological Health
Nebraska Department of Health
301 Centennial Mall, South
P.O. Box 95007
Lincoln, NE 68509
(402) 471-2168
1-800-334-9491 In State

Nevada Stan Marshall
Department of Human Resources
505 East King Street
Room 203
Carson City, NV 89710
(702) 687-5394

New Hampshire David Chase
Bureau of Radiological Health
Division of Public Health Services
Health and Welfare Building
Six Hazen Drive
Concord, NH 03301
(603) 271-4674
1-800-852-3345 x4674

New Jersey Tonalee Carlson Key
Division of Environmental Quality
Department of Environmental
Protection
CN 415
Trenton, NJ 08625-0145
(609) 987-6369
1-800-648-0394 in state

New Mexico William M. Floyd
Radiation Licensing and Registration
Section
New Mexico Environmental
Improvement Division
1190 St. Francis Drive
Santa Fe, NM 87503
(505) 827-4300

New York William J. Condon
Bureau of Environmental Radiation
Protection
New York State Health Department
Two University Place
Albany, NY 12202
(518) 458-6495
1-800-458-1158 in state

North Carolina Dr. Felix Fong
Radiation Protection Division
Department of Environmental Health
and Natural Resources
701 Barbour Drive
Raleigh, NC 27603-2008
(919) 571-4141
1-800-662-7301 (recorded info x4196)

North Dakota Arlen Jacobson
North Dakota Department of Health
1200 Missouri Avenue, Room 304
P.O. Box 5520
Bismarck, ND 58502-5520
(701) 221-5188

Ohio Marcie Matthews
Radiological Health Program
Department of Health
1224 Kinnear Road - Suite 120
Columbus, OH 43212
(614) 644-2727
1-800-523-4439 in state

Oklahoma Gene Smith
Radiation Protection Division
Oklahoma State Department of
Health
P.O. Box 53551
Oklahoma City, OK 73152
(405) 271-5221

South Dakota Mike Pochop
Division of Environment Regulation
Department of Water and Natural
Resources
Joe Foss Building, Room 217
523 E. Capitol
Pierre, SD 57501-3181
(605) 773-3351

Oregon George Toombs
Department of Human Resources
Health Division
1400 SW 5th Avenue
Portland, OR 97201
(503) 731-4014

Tennessee Susie Shimek
Division of Air Pollution Control
Bureau of the Environment
Department of Environment and
Conservation
Customs House, 701 Broadway
Nashville, TN 37219-5403
(615) 532-0733
1-800-232-1139 in state

Pennsylvania Michael Pyles
Pennsylvania Department of
Environmental Resources
Bureau of Radiation Protection
P.O. Box 2063
Harrisburg, PA 17120
(717) 783-3594
1-800-23-RADON In State

Texas Gary Smith
Bureau of Radiation Control
Texas Department of Health
1100 West 49th Street
Austin, TX 78756-3189
(512) 834-6688

Puerto Rico David Saldana
Radiological Health Division
G.P.O. Call Box 70184
Rio Piedras, Puerto Rico 00936
(809) 767-3563

Utah John Hultquist
Bureau of Radiation Control
Utah State Department of Health
288 North, 1460 West
P.O. Box 16690
Salt Lake City, UT 84116-0690
(801) 536-4250

Rhode Island Edmund Arcand
Division of Occupational Health and
Radiation
Department of Health
205 Cannon Building
Davis Street
Providence, RI 02908
(401) 277-2438

Vermont Paul Clemons
Occupational and Radiological Health
Division
Vermont Department of Health
10 Baldwin Street
Montpelier, VT 05602
(802) 828-2886
1-800-640-0601 in state

South Carolina Bureau of Radiological Health
Department of Health and
Environmental Control
2600 Bull Street
Columbia, SC 29201
(803) 734-4631
1-800-768-0362

Virgin Islands Contact the U.S. Environmental
Protection Agency, Region II
in New York
(212) 264-4110

Virginia Shelly Ottenbrite
Bureau of Radiological Health
Department of Health
109 Governor Street
Richmond, VA 23219
(804) 786-5932
1-800-468-0138 in state

Washington Kate Coleman
Department of Health
Office of Radiation Protection
Airdustrial Building 5, LE-13
Olympia, WA 98504
(206) 753-4518
1-800-323-9727 In State

West Virginia Beattie L. DeBord
Industrial Hygiene Division
West Virginia Department of Health
151 11th Avenue
South Charleston, WV 25303
(304) 558-3526
1-800-922-1255 In State

Wisconsin Conrad Weiffenbach
Radiation Protection Section
Division of Health
Department of Health and Social
Services
P.O. Box 309
Madison, WI 53701-0309
(608) 267-4796
1-800-798-9050 in state

Wyoming Janet Hough
Wyoming Department of Health and
Social Services
Hathway Building, 4th Floor
Cheyenne, WY 82002-0710
(307) 777-6015
1-800-458-5847 in state

STATE GEOLOGICAL SURVEYS

May, 1993

<u>Alabama</u>	Ernest A. Mancini Geological Survey of Alabama P.O. Box 0 420 Hackberry Lane Tuscaloosa, AL 35486-9780 (205) 349-2852	<u>Florida</u>	Walter Schmidt Florida Geological Survey 903 W. Tennessee St. Tallahassee, FL 32304-7700 (904) 488-4191
<u>Alaska</u>	Thomas E. Smith Alaska Division of Geological & Geophysical Surveys 794 University Ave., Suite 200 Fairbanks, AK 99709-3645 (907) 479-7147	<u>Georgia</u>	William H. McLemore Georgia Geologic Survey Rm. 400 19 Martin Luther King Jr. Dr. SW Atlanta, GA 30334 (404) 656-3214
<u>Arizona</u>	Larry D. Fellows Arizona Geological Survey 845 North Park Ave., Suite 100 Tucson, AZ 85719 (602) 882-4795	<u>Hawaii</u>	Manabu Tagomori Dept. of Land and Natural Resources Division of Water & Land Mgt P.O. Box 373 Honolulu, HI 96809 (808) 548-7539
<u>Arkansas</u>	Norman F. Williams Arkansas Geological Commission Vardelle Parham Geology Center 3815 West Roosevelt Rd. Little Rock, AR 72204 (501) 324-9165	<u>Idaho</u>	Earl H. Bennett Idaho Geological Survey University of Idaho Morrill Hall, Rm. 332 Moscow, ID 83843 (208) 885-7991
<u>California</u>	James F. Davis California Division of Mines & Geology 801 K Street, MS 12-30 Sacramento, CA 95814-3531 (916) 445-1923	<u>Illinois</u>	Morris W. Leighton Illinois State Geological Survey Natural Resources Building 615 East Peabody Dr. Champaign, IL 61820 (217) 333-4747
<u>Colorado</u>	Pat Rogers (Acting) Colorado Geological Survey 1313 Sherman St., Rm 715 Denver, CO 80203 (303) 866-2611	<u>Indiana</u>	Norman C. Hester Indiana Geological Survey 611 North Walnut Grove Bloomington, IN 47405 (812) 855-9350
<u>Connecticut</u>	Richard C. Hyde Connecticut Geological & Natural History Survey 165 Capitol Ave., Rm. 553 Hartford, CT 06106 (203) 566-3540	<u>Iowa</u>	Donald L. Koch Iowa Department of Natural Resources Geological Survey Bureau 109 Trowbridge Hall Iowa City, IA 52242-1319 (319) 335-1575
<u>Delaware</u>	Robert R. Jordan Delaware Geological Survey University of Delaware 101 Penny Hall Newark, DE 19716-7501 (302) 831-2833	<u>Kansas</u>	Lee C. Gerhard Kansas Geological Survey 1930 Constant Ave., West Campus University of Kansas Lawrence, KS 66047 (913) 864-3965

Kentucky Donald C. Haney
Kentucky Geological Survey
University of Kentucky
228 Mining & Mineral Resources
Building
Lexington, KY 40506-0107
(606) 257-5500

Louisiana William E. Marsalis
Louisiana Geological Survey
P.O. Box 2827
University Station
Baton Rouge, LA 70821-2827
(504) 388-5320

Maine Walter A. Anderson
Maine Geological Survey
Department of Conservation
State House, Station 22
Augusta, ME 04333
(207) 289-2801

Maryland Emery T. Cleaves
Maryland Geological Survey
2300 St. Paul Street
Baltimore, MD 21218-5210
(410) 554-5500

Massachusetts Joseph A. Sinnott
Massachusetts Office of
Environmental Affairs
100 Cambridge St., Room 2000
Boston, MA 02202
(617) 727-9800

Michigan R. Thomas Segall
Michigan Geological Survey Division
Box 30256
Lansing, MI 48909
(517) 334-6923

Minnesota Priscilla C. Grew
Minnesota Geological Survey
2642 University Ave.
St. Paul, MN 55114-1057
(612) 627-4780

Mississippi S. Cragin Knox
Mississippi Office of Geology
P.O. Box 20307
Jackson, MS 39289-1307
(601) 961-5500

Missouri James H. Williams
Missouri Division of Geology &
Land Survey
111 Fairgrounds Road
P.O. Box 250
Rolla, MO 65401
(314) 368-2100

Montana Edward T. Ruppel
Montana Bureau of Mines & Geology
Montana College of Mineral Science
and Technology, Main Hall
Butte, MT 59701
(406) 496-4180

Nebraska Perry B. Wigley
Nebraska Conservation & Survey
Division
113 Nebraska Hall
University of Nebraska
Lincoln, NE 68588-0517
(402) 472-2410

Nevada Jonathan G. Price
Nevada Bureau of Mines & Geology
Stop 178
University of Nevada-Reno
Reno, NV 89557-0088
(702) 784-6691

New Hampshire Eugene L. Boudette
Dept. of Environmental Services
117 James Hall
University of New Hampshire
Durham, NH 03824-3589
(603) 862-3160

New Jersey Haig F. Kasabach
New Jersey Geological Survey
P.O. Box 427
Trenton, NJ 08625
(609) 292-1185

New Mexico Charles E. Chapin
New Mexico Bureau of Mines &
Mineral Resources
Campus Station
Socorro, NM 87801
(505) 835-5420

New York Robert H. Fakundiny
New York State Geological Survey
3136 Cultural Education Center
Empire State Plaza
Albany, NY 12230
(518) 474-5816

<u>North Carolina</u>	Charles H. Gardner North Carolina Geological Survey P.O. Box 27687 Raleigh, NC 27611-7687 (919) 733-3833	<u>South Carolina</u>	Alan-Jon W. Zupan (Acting) South Carolina Geological Survey 5 Geology Road Columbia, SC 29210-9998 (803) 737-9440
<u>North Dakota</u>	John P. Bluemle North Dakota Geological Survey 600 East Blvd. Bismarck, ND 58505-0840 (701) 224-4109	<u>South Dakota</u>	C.M. Christensen (Acting) South Dakota Geological Survey Science Center University of South Dakota Vermillion, SD 57069-2390 (605) 677-5227
<u>Ohio</u>	Thomas M. Berg Ohio Dept. of Natural Resources Division of Geological Survey 4383 Fountain Square Drive Columbus, OH 43224-1362 (614) 265-6576	<u>Tennessee</u>	Edward T. Luther Tennessee Division of Geology 13th Floor, L & C Tower 401 Church Street Nashville, TN 37243-0445 (615) 532-1500
<u>Oklahoma</u>	Charles J. Mankin Oklahoma Geological Survey Room N-131, Energy Center 100 E. Boyd Norman, OK 73019-0628 (405) 325-3031	<u>Texas</u>	William L. Fisher Texas Bureau of Economic Geology University of Texas University Station, Box X Austin, TX 78713-7508 (512) 471-7721
<u>Oregon</u>	Donald A. Hull Dept. of Geology & Mineral Indust. Suite 965 800 NE Oregon St. #28 Portland, OR 97232-2162 (503) 731-4600	<u>Utah</u>	M. Lee Allison Utah Geological & Mineral Survey 2363 S. Foothill Dr. Salt Lake City, UT 84109-1491 (801) 467-7970
<u>Pennsylvania</u>	Donald M. Hoskins Dept. of Environmental Resources Bureau of Topographic & Geologic Survey P.O. Box 2357 Harrisburg, PA 17105-2357 (717) 787-2169	<u>Vermont</u>	Diane L. Conrad Vermont Division of Geology and Mineral Resources 103 South Main St. Waterbury, VT 05671 (802) 244-5164
<u>Puerto Rico</u>	Ramón M. Alonso Puerto Rico Geological Survey Division Box 5887 Puerta de Tierra Station San Juan, P.R. 00906 (809) 722-2526	<u>Virginia</u>	Stanley S. Johnson Virginia Division of Mineral Resources P.O. Box 3667 Charlottesville, VA 22903 (804) 293-5121
<u>Rhode Island</u>	J. Allan Cain Department of Geology University of Rhode Island 315 Green Hall Kingston, RI 02881 (401) 792-2265	<u>Washington</u>	Raymond Lasmanis Washington Division of Geology & Earth Resources Department of Natural Resources P.O. Box 47007 Olympia, Washington 98504-7007 (206) 902-1450

West Virginia Larry D. Woodfork
West Virginia Geological and
Economic Survey
Mont Chateau Research Center
P.O. Box 879
Morgantown, WV 26507-0879
(304) 594-2331

Wisconsin James Robertson
Wisconsin Geological & Natural
History Survey
3817 Mineral Point Road
Madison, WI 53705-5100
(608) 263-7384

Wyoming Gary B. Glass
Geological Survey of Wyoming
University of Wyoming
Box 3008, University Station
Laramie, WY 82071-3008
(307) 766-2286

EPA REGION 10 GEOLOGIC RADON POTENTIAL SUMMARY

by

James K. Otton, Kendall A. Dickinson, Douglass E. Owen, and Sandra L. Szarzi
U.S. Geological Survey

EPA Region 10 includes the states of Alaska, Idaho, Oregon, and Washington. For each state, geologic radon potential areas were delineated and ranked on the basis of geologic, soils, housing construction, and other factors. Areas in which the *average screening indoor radon level of all homes within the area* is estimated to be greater than 4 pCi/L were ranked high. Areas in which the average screening indoor radon level of all homes within the area is estimated to be between 2 and 4 pCi/L were ranked moderate/variable, and areas in which the average screening indoor radon level of all homes within the area is estimated to be less than 2 pCi/L were ranked low. Information on the data used and on the radon potential ranking scheme is given in the introduction to this volume. More detailed information on the geology and radon potential of each state in Region 10 is given in the individual state chapters. The individual chapters describing the geology and radon potential of the states in EPA Region 10, though much more detailed than this summary, still are generalized assessments and there is no substitute for having a home tested. Within any radon potential area homes with indoor radon levels both above and below the predicted average likely will be found.

The geology and radon potential of the Pacific Northwest (fig. 1) and Alaska (fig. 2) is diverse; thus the two areas will be considered separately. The Pacific Northwest includes eight distinct major radon geologic provinces: the Coastal Range-Klamath Mountains, the Puget Lowland-Willamette River Valley, the Cascade Range, the Columbia Plateau-High Lava Plains-Blue Mountains, the northern Rocky Mountains, the Snake River plain, the middle Rocky Mountains, and the northern Basin and Range-Owyhee Plateau (fig. 1). Maps showing indoor radon averages for counties in the Pacific Northwest and boroughs in Alaska are shown in figures 3a and 3b. Averages range from less than 1.0 pCi/L to 14.9 pCi/L. Details of the indoor radon studies are described in the individual state chapters.

PACIFIC NORTHWEST

Coastal Range-Klamath Mountains

The Coastal Range Province (1, fig. 1) extends from the Olympic Peninsula of Washington south to the coastal parts of the Klamath Mountains in southwestern Oregon. In Washington, the Coast Ranges are underlain principally by Cretaceous and Tertiary continental and marine sedimentary rocks and pre-Miocene volcanic rocks. In Oregon, the northern part of the Coastal Ranges is underlain principally by marine sedimentary rocks and mafic volcanic rocks of Tertiary age. The southern part of the Coast Range is underlain by Tertiary estuarine and marine sedimentary rocks, much of them feldspathic and micaceous. The Klamath Mountains (2, fig. 1) are dominated by Triassic to Jurassic metamorphic, volcanic, and sedimentary rocks, with some Cretaceous intrusive rocks. These metamorphic and volcanic rocks are largely of mafic composition. Large masses of ultramafic rocks occur throughout the Klamath area. Sand dunes and marine terraces are common along the coastal areas of this province.

The radon potential of the Coastal Range Province is low overall. Most of the area has high rainfall and, as a consequence, high soil moisture. Uranium in the soils is typically low, although soils of the Oregon part of the Coast Ranges tend to be higher in uranium than do soils of




Figure 1- Radon geologic provinces of the Pacific Northwest (generalized from state chapters included in this report). 1- Coast Ranges; 2- Klamath Mountains; 3- Puget Lowland; 4- Willamette River Valley; 5- Cascade Range; 6- Northern Rocky Mountains; 7- Columbia Plateau; 8- Blue Mountains; 9- High Lava Plains; 10- Basin and Range; 11- Owyhee Plateau; 12- Snake River Plain; and 13- Middle Rocky Mountains.

the Washington part. A few communities along the river valleys near the coast of Oregon may have locally elevated indoor radon where highly permeable, excessively well-drained soils occur on river alluvium with a modestly elevated uranium content. The northeastern corner of the Olympic Peninsula has lower rainfall and lower soil moisture than does the rest of the Coastal Range Province. Here, highly permeable, excessively well-drained soils may cause locally elevated indoor radon levels.

Puget Lowland-Willamette River Valley

The Puget Lowland (3, fig. 1) is underlain almost entirely by glacial deposits and Holocene alluvium. Most of the glacial and alluvial material of the Puget Lowland is derived from the Cascades to the east, and from the mountains of the Olympic peninsula to the west. River alluvium and river terraces underlie most of the Willamette River valley (4, fig. 1). However, many of the hills that rise above the plains of the Lowland are underlain by Tertiary basalts and marine sediments.

The Puget Lowland overall has very low radon potential because of low uranium content of soils and because high rainfall produces high soil moisture, which slows radon movement. Houses in most townships in the Bonneville Power Administration study from Tacoma northward average less than 1 pCi/L radon. Structures built on locally very steep or well-drained soils, especially on the east side of the lowland area, may be among the few likely to have elevated indoor radon levels. The geologic radon potential is moderate only in the southern part of the Puget Lowland, south of Tacoma, where excessively drained soils and somewhat elevated uranium in soils occur.

The Willamette River Valley has moderate radon potential overall. Much of the area has somewhat elevated uranium in soils, and many areas have excessively drained soils and soils with high emanating power. Studies by the Oregon Department of Health and the Bonneville Power Administration indicate that houses in many counties and townships in the valley average between 2 and 4 pCi/L radon.

Cascade Range

The Cascade Range (5, fig. 1) can be divided into two geologic terranes: a northern terrane composed principally of Mesozoic metamorphic rocks intruded by Mesozoic and Tertiary granitic rocks, and a southern terrane composed of Tertiary and Holocene volcanic rocks. The Holocene volcanic centers are responsible for locally thick volcanic-ash deposits east of the Cascade Mountains. Within the southern terrane, the western Cascades are dominated by Tertiary andesite flows, basalt flows, and pyroclastic rocks, whereas the eastern Cascades have many recently active volcanoes and are underlain by late Tertiary to Quaternary basaltic and andesitic volcanic rocks.

Overall, the sparsely populated Cascade Range Province has low radon potential because of the low uranium and high moisture contents of the soils. Areas that are exceptions to this include the Columbia River Gorge, where highly permeable, excessively well drained soils underlie many of the communities, and thus the radon potential is moderate. Much of the alluvium in the Gorge is also derived from the upper Columbia River valley, where the uranium content of the geologic materials is higher than the rocks within the Cascade Mountain Province itself. Studies by the Oregon Department of Health and the Bonneville Power Administration show that indoor radon levels in homes in population centers along the Columbia River average 2 to 4 pCi/L.

Columbia Plateau, High Lava Plains, and Blue Mountains

The Columbia Plateau (7, fig. 1) is underlain principally by Miocene basaltic and andesitic volcanic rocks, tuffaceous sedimentary rocks and tuff. An extensive veneer of Pleistocene glaciofluvial outwash, eolian, and lacustrine deposits covers these volcanic rocks. The High Lava Plains (9, fig. 1) are underlain by Miocene basaltic and volcanic rocks like those of the Columbia Plateau without the veneer of younger sedimentary rocks. The Blue Mountains (8, fig. 1) have similar basaltic and andesitic rocks and also include significant outcrop areas of Triassic and Jurassic sedimentary and volcanic rocks, weakly metamorphosed in many areas, and younger intrusive rocks.

The Columbia Plateau, with its areas of extensive Pleistocene glacio-fluvial outwash, eolian, and lacustrine deposits, contains locally highly permeable soils, soils with high emanating coefficients, and elevated soil uranium levels. This area has generally moderate radon potential. Although the Blue Mountains have relatively low uranium in soils, average indoor radon levels are in the 2-4 pCi/L range, probably because most population centers occur in alluviated valleys with highly permeable soils. This area has moderate radon potential. In contrast, the High Lava Plains, with much lower uranium in soils and only local areas of highly permeable soils, have low overall radon potential.

Northern Rocky Mountains

The Northern Rocky Mountains (6, fig. 1) comprise the mountainous terrane of the northeast and north-central parts of Washington and northern and central Idaho. This area is underlain by Precambrian and Paleozoic sedimentary rocks, and by Mesozoic metamorphic rocks; all are intruded by Mesozoic and Tertiary granitic rocks. The largest intrusive mass, the Idaho Batholith, is a complex of granitic rock units that range from diorite to granite. Highly uraniumiferous, Late Cretaceous to early Tertiary granites crop out throughout the Northern Rocky Mountains. An extensive, though dissected, veneer of Tertiary volcanic rocks crops out over much of the central Idaho portion of the Northern Rocky Mountains.

The Northern Rocky Mountains Province has high radon potential. Excessively well drained glaciofluvial outwash or coarse gravels in alluvial fans underlie many of the valleys throughout the area. The granitic material in much of the outwash contains moderate to locally high concentrations of uranium. Areas where uranium occurrences are found, such as in the granitic and metamorphic terranes in the mountains north of Spokane, may have structures with extreme levels of indoor radon. Buildings in most of the alluvial valleys in Washington and Idaho north, northwest, and east of Spokane may be expected to have average indoor radon screening measurements above 4 pCi/L.

Snake River Plain

The Snake River Plain (12, fig. 1) forms an arcuate depression in southern Idaho that is underlain principally by basaltic volcanic rocks of generally low eU (1 ppm or less). However, alluvium from neighboring mountains and silicic tuffaceous sedimentary rocks covers much of the upper Snake River Valley near Wyoming and the western end of the Snake River Plain near Boise and south of Mountain Home. These materials have eU values that range from 1.5-5.0 ppm. Those areas underlain by basalt have low to locally moderate radon potential. However, those areas where basalt is overlain by silicic tuffaceous sedimentary rocks and alluvium along the Snake River Valley have high overall radon potential. Most populous areas are in the latter category.

Middle Rocky Mountains

The Middle Rocky Mountains Province (13, fig. 1) forms a strip along the border between Wyoming and Idaho and comprises two areas. The northern area is the Yellowstone Plateau, a high-standing plateau area underlain mostly by rhyolites containing moderate amounts of uranium. To the south are complexly faulted and folded mountain ranges of Paleozoic and Mesozoic sedimentary rocks, including uranium-bearing phosphatic rocks.

The high average uranium content of the volcanic rocks of the Yellowstone area and the coarse alluvium in the valleys of the southern mountain areas suggest that this province has high geologic radon potential.

Basin and Range Province, Owyhee Plateau

The very sparsely populated northern part of the Basin and Range Province (10, fig. 1) lies along the southern and southeastern edge of Region 10. It is composed of tectonically extended areas where linear mountain ranges alternate with valleys and less extended plateau areas. It is underlain mainly by basaltic to andesitic volcanic rocks, silicic ash-flow tuffs, including some welded tuffs, and sediments derived from these units. Several playa basins occupy the centers of the valleys. The Owyhee Plateau of southwestern Idaho (11, fig. 1) consists of Tertiary and Quaternary basalt, andesite, and rhyolite, and sediments derived from these units. A few caldera complexes, some of them with associated uranium mineralization, occur within the Owyhee Plateau. Some mountain ranges in the eastern part of this province are underlain mainly by Paleozoic and Mesozoic sedimentary rocks. Based on the high aeroradiometric signature of most of the exposed rock units and the presence of many highly permeable soil units, the radon potential of this area is generally high.

ALASKA

Alaska can be divided from north to south into eight geologic radon provinces: the Arctic Coastal Plain, the Arctic Foothills, the Arctic Mountains, Central Alaska, the Northern Plateaus (a subprovince of Central Alaska), the Alaska-Aleutian Ranges, the Coastal Trough, and the Border Ranges Provinces (fig. 2).

Arctic Coastal Plain

The Arctic Coastal Plain Province (North Slope, 1, fig. 2) consists primarily of Quaternary sediment, most of which is composed of alluvium, glacial debris, and eolian sand and silt. A belt of Tertiary sedimentary rocks along the eastern third of the area separates the coastal plains from the foothills to the south.

This area has low radon potential. No significant uranium occurrences are known in this area, and the number of gamma-ray anomalies is low when compared with other parts of Alaska. The coastal plain is unglaciated and contains tundra soils and permafrost. These soils probably have low gas transmissivity because of water or ice saturation.

Arctic Foothills

The Arctic Foothills Province (2, fig. 2) is largely composed of marine and nonmarine Cretaceous sandstone and shale. The Cretaceous beds are folded into west-trending anticlines and synclines. Part of the area was covered by glaciers.




Figure 2- Physiographic provinces of Alaska (from the state chapter included in this report). 1- Arctic Coastal Plain; 2- Arctic Foothills; 3- Arctic Mountains; 4- Central Province, 4a- Seward Peninsula, 4b- Bering Shelf, 4c- Ahklun Mountains, 4d- Western Alaska, 4e- Northern Plateaus; 5- Alaska-Aleutian Province; 6- Coastal Trough; 7- Pacific Border Ranges; and 8- Coast Mountains.

This area has low radon potential overall. The Cretaceous sandstone and shale that makes up the foothills could produce relatively large amounts of radon but no evidence that they do is on hand. The area contains no known uranium occurrences or deposits, and the part of the area where airborne gamma-ray measurements were made shows a low number of anomalies. The tundra soils have permafrost and apparent low gas transmissivity.

Arctic Mountains

The Arctic Mountains Province (3, fig. 2) is composed largely of upper Precambrian and Paleozoic marine sedimentary rocks. They are cut by west-trending thrust faults with upthrown sides to the south.

This area has moderate radon potential. The Precambrian and Paleozoic marine sedimentary rocks that make up the Arctic Mountains probably are not producers of high levels of radon as there is little or no phosphate rock or black shale in these sequences. There are no known significant uranium occurrences in this area. However, stream sediments in this province contain moderately high levels of uraniferous resistate minerals. The area has been glaciated, but much of the terrane is bare rock without surficial glacial material. The soils are classified as rock land, which includes glacial ice.

Central Province (exclusive of the Northern Plateaus subprovince)

The Central Province, an area of plains, plateaus, and rounded mountains, is geologically complex. The Central Province is divided into five subprovinces: Western Alaska, Seward Peninsula, Aklun Mountains, the Bering Shelf (4a-d, fig. 2) and the Northern Plateaus (5, fig. 2). The Northern Plateaus are considered separately below.

Western Alaska is underlain mostly by Cretaceous marine sedimentary rocks and lower Paleozoic sedimentary and metamorphic rocks. A large area of Cretaceous and Tertiary volcanic rock is present in the western part of this subprovince. The Seward Peninsula consists mostly of Precambrian and Paleozoic metamorphic rocks, with lesser amounts of Precambrian and Paleozoic sedimentary rocks, Quaternary sediments, and Tertiary and Quaternary mafic volcanic rocks. The Aklun Mountains are composed mostly of marine sedimentary rocks and small intrusive masses of Jurassic and Tertiary age. The Bering Shelf is covered almost entirely by Quaternary surficial sediments, with minor areas of Tertiary volcanic rocks.

Overall the Central Province has moderate radon potential as many radon-producing rocks occur there. There are, for instance, several areas of uraniferous granites together with felsic intrusive and volcanic rocks. In addition, the area contains a few uranium deposits of potentially commercial size at Death Valley on the Seward Peninsula and in the Healy Creek coal basin. The area also contains a significant number of gamma-ray anomalies. Nearly all of the area falls within a belt of uraniferous stream sediments. The schist that produces high indoor radon near Fairbanks is in this area. Little of the province has been glaciated. The soils are mostly of the Tundra type with variable permafrost. Significant areas of rockland and subarctic brown forest soils occur. The latter soils may have high gas transmissivity.

Northern Plateaus

The Northern Plateaus subprovince (5, fig. 2) is covered by flat-lying Tertiary basin-fill (nonmarine clastic rocks), Quaternary surficial deposits, Precambrian through Cretaceous mostly marine sedimentary rocks, Paleozoic and Precambrian metamorphic rocks, and Mesozoic intrusive and volcanic rocks. The metamorphic rocks include metamorphosed granites and amphibolite.

The mesozoic intrusives are mostly gabbro and diabase. The Tintina and Denali fault zones cross this subprovince.

The Northern Plateaus subprovince has a moderate radon potential overall. A moderate number of aeroradiometric anomalies occurs in the subprovince. Although indoor radon data are sparse, indoor radon in parts of the Fairbanks and Fairbanks Northstar Boroughs is high. Felsic intrusives are scattered in two belts, one intruding Paleozoic and Precambrian metamorphic rocks in the southeast one-third of the subprovince and one intruding Lower Paleozoic and (or) Precambrian sedimentary rocks along the northwest margin of the subprovince. The area contains one known significant uranium and thorium deposit at Mount Prindle. Uranium is high in stream sediments in the south-central part and along the northwest border of the subprovince.

Alaska-Aleutian Ranges and Coastal Mountains

The Alaska-Aleutian Ranges and Coastal Mountains Province (6, fig. 2) includes the Aleutian Peninsula, a northeast-trending mountain belt in south-central Alaska that includes Mt. McKinley, a southeast-trending mountain belt that extends from the Mt. McKinley area southeastward to Canada, and the Coast Mountains in the southeast. On the Aleutian Peninsula from Unimak Pass westward, the bedrock consists mostly of Quaternary and Tertiary volcanic rocks and Tertiary sedimentary rocks. Tertiary and Quaternary volcanic rocks are also common northeast of the Pass, but other rocks, including Jurassic and Cretaceous sedimentary rocks and Jurassic intrusive rocks of intermediate and felsic composition, are also common in this area. In addition, large masses of Tertiary mafic volcanic rocks and Jurassic or Cretaceous intermediate intrusives are found in the area west of Cook Inlet and southwest of Mount McKinley. A varied assortment of Phanerozoic rocks are present in the Talkeetna Mountains and southeastward to the Canadian border. These include Paleozoic mafic volcanic rocks together with their sedimentary and metamorphic derivatives; Mesozoic mafic volcanic flows and tuffs, together with various units of shale, conglomerate, graywacke, and slate; and Tertiary and Quaternary intermediate volcanic rocks, Tertiary felsic intrusives, and Quaternary glacial deposits including eolian sand and silt. The Coastal Mountains are composed mostly of ultramafic, intermediate, and silicic volcanic intrusive rocks of varying ages, and Paleozoic through Mesozoic sedimentary rocks. These rocks are highly deformed and variably metamorphosed.

This area has moderate radon potential overall, although the uncertainty is high. The Aleutian-Alaska Range contains felsic intrusives and other rocks that are likely to be uranium-rich, although no significant uranium occurrences are known in this area. However, the area has a moderate to substantial number of anomalously uranium-rich stream sediment samples. Most of the area is or was covered by glaciers and glacial outwash may be highly permeable in many areas. Soils are mostly classified as rockland or tundra.

Coastal Trough

The Coastal Trough Province (7, fig. 2) includes a series of Cenozoic depositional basins containing thick sequences of Tertiary continental clastic and volcanic rocks that generally overlie Cretaceous or older sedimentary rocks penetrated by Tertiary intrusive rocks. Mesozoic sedimentary rocks and Pleistocene, mostly glacial, deposits, occur in some areas.

The radon potential of this area is moderate overall, but locally high indoor radon levels could occur near uranium occurrences. The Coastal Trough Province contains Tertiary continental clastic rocks similar to units that produce uranium in the western conterminous United States. The overall uranium content of these rocks is not high, but small uranium occurrences are found in the

Susitna Lowlands and in the Admiralty trough in southeastern Alaska. Soils are mostly brown and gray-brown podzolic forest soils, which could have high gas transmissivity. Heavy rainfall and saturated soils in southeast Alaska likely retards soil gas migration.

Pacific Border Ranges

The Border Ranges Province (8, fig. 2) is generally south and west of the Coastal Trough Province. Jurassic and Cretaceous sedimentary and metamorphic rocks with interbedded mafic volcanic rocks and some gabbro make up most of the Border Ranges rocks. A fairly large area of early Tertiary sedimentary, volcanogenic sedimentary rocks, and volcanic rocks is found in the Prince William Sound area.

The Border Ranges Province generally has low radon potential, although some uranium-bearing rocks and uranium occurrences are likely to be present. The uranium deposit at Bokan Mountain is associated with a uranium-rich peralkaline granite. The uranium content of stream sediments in the Border Ranges is intermediate for Alaska, although data are absent from many areas. Podzolic brown and gray-brown forest soils are common in the Border Ranges, and they could have high gas permeability. However, in this part of Alaska annual rainfall is about 14 feet, and water saturation likely retards gas flow in soils on all but the steepest slopes.




Figure 3A. Screening indoor radon data from the State/EPA Residential Radon Survey and the Oregon Radon Project, for counties with 5 or more measurements in the conterminous part of EPA Region 10. Histograms in map legends show the number of counties in each category. The number of samples in each county may not be sufficient to statistically characterize the radon levels of the counties, but they do suggest general trends. Unequal category intervals were chosen to provide reference to decision and action levels.




Figure 3B. Screening indoor radon data from the State/EPA Residential Radon Survey of Alaska, for counties with 5 or more measurements. Data are from 2-7 day charcoal canister tests. Histograms in map legends show the number of counties in each category. The number of samples in each county may not be sufficient to statistically characterize the radon levels of the counties, but they do suggest general trends. Unequal category intervals were chosen to provide reference to decision and action levels.

PRELIMINARY GEOLOGIC RADON POTENTIAL ASSESSMENT OF ALASKA

by
Kendell A. Dickinson
U.S. Geological Survey

INTRODUCTION

The state of Alaska is characterized by complex geology and soils developed on rugged terrain in cool, moist climates. It is a large, sparsely populated area. Indoor radon data are only available for the populated areas. For many areas conclusions are very general because of the lack of field studies and the general lack of data. It is, however, hoped that the study will suffice as a general guide to future studies and planning in terms of radon potential.

This is a generalized assessment of geologic radon potential of rocks, soils, and surficial deposits of Alaska. The scale of this assessment is such that it is inappropriate for use in identifying the radon potential of small areas such as neighborhoods, individual building sites, or housing tracts. Any localized assessment of radon potential must be supplemented with additional data and information from the locality. Within any area of a given radon potential ranking, there are likely to be areas with higher or lower radon levels than characterized for the area as a whole. Indoor radon levels, both high and low, can be quite localized, and there is no substitute for testing individual homes. Elevated levels of indoor radon have been found in every State, and EPA recommends that all homes be tested. For more information on radon, the reader is urged to consult the local or State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

GEOGRAPHIC SETTING

Alaska occupies the great northwestern peninsula of North America; it is an extension of the North American Cordillera. Alaska can be divided from north to south into the following physiographic provinces (Wahrhaftig, 1965): The Arctic Coastal Plain, Arctic Foothills, Arctic Mountains, Central province (including Northern Plateaus, Western Alaska, Seward Peninsula, Bering Shelf, and Aklun Mountains subprovinces), Alaska-Aleutian Ranges (including the Coast Mountains of southeastern Alaska), Coastal Trough, and Pacific Border Ranges (fig. 1).

The southeastern corner of southeastern Alaska lies about 500 miles from the State of Washington, the nearest part of the conterminous 48 states. The southeast corner of southeast Alaska lies about 1800 miles from the northwest end of the north slope and about 2300 miles from Attu Island, Alaska's and the USA's most eastern point (west of 180 degrees E. longitude). Alaska is the largest state in the USA and is the least populated of all the states. It occupies 591,000 square miles.

According to the 1990 census there were 550,043 people in Alaska, about one third of whom live in Anchorage and about one half of whom live in Anchorage and Fairbanks combined. Alaska averages about 1.1 people per square mile (fig. 2).




Figure 1. Physiographic provinces of Alaska (Wahrhaftig, 1965).




Figure 2. Population of boroughs in Alaska (1990 U.S. Census data).

GEOLOGIC SETTING

The geology of Alaska varies widely (fig. 3). As many as 60 tectonostratigraphic terranes have been recognized in Alaska (Jones and others, 1987; Monger and Berg, 1987). Terranes were defined by Howell (1985) as "fault-bounded geologic entities of regional extent each characterized by a geologic history distinct from neighboring terranes". These terranes were accreted to the western North American craton during the last 200 million years and now form Alaska. Diverse assemblages of igneous, metamorphic, and sedimentary rocks ranging from Precambrian to Holocene make up these terranes. Numerous Tertiary basins that contain largely continental deposits of coal-bearing clastic rock formed after the terranes were in place. Stone and Wallace (1987) grouped the terranes into framework provinces each containing several terranes of similar tectonic or geologic style. The tectonostratigraphic terrane boundaries are coincident with fault zones in many areas.

The geology of Alaska is summarized below on the basis of the physiographic provinces and subprovinces of Wahrhaftig (1955). Terrane summaries included for each physiographic division were abstracted from Jones and others (1987) and Monger and Berg (1987). Rocks characteristic of each terrane form the bedrock in much of Alaska, but they are not part of the surface geology in some areas and do not play a direct role in radon generation in those areas.

Surficial deposits in the Arctic Coastal Plain consist mostly of alluvium, eolian sand, and silt of Quaternary age. A belt of marine and nonmarine conglomerate, sandstone, shale, and mudstone of Tertiary age that separates the coastal plain from the Arctic Foothills to the south is found along the eastern 1/3 of the area. The Arctic Coastal Plain province is underlain by the North Slope subterrane (Precambrian to Lower Paleozoic basement rocks overlain by Mississippian through Triassic and younger Mesozoic sedimentary rocks).

Surface rocks in the Arctic Foothills province are largely composed of marine and nonmarine Cretaceous sandstone and shale, but some Paleozoic and Mesozoic rocks are also present. The Cretaceous beds are folded into west-trending anticlines and synclines. The North Slope (see above), Endicott Mountains (Sequences of Devonian clastic rocks, Mississippian shale and carbonate rocks, and Upper Paleozoic and Mesozoic chert, argillite and calcareous rocks), and DeLong Mountains (thick Devonian and Mississippian carbonates and younger sequences of chert and argillite) subterrane of the Arctic terrane underlie the Arctic Foothills province.

The Arctic Mountains province (Brooks Range) is composed largely of upper Precambrian and Paleozoic marine sedimentary rocks. These rocks are cut by west-trending thrust faults with their upthrown sides to the south. The Endicott Mountains (see above) and Hammond subterrane (polymetamorphosed assemblage of Middle Paleozoic and older carbonate rocks, calc-schist, quartz-mica schist, quartzite, and metarhyolite intruded by Devonian gneissic granitic rocks) underlie most of the province. Several other terranes underlie small parts of the province. Among these are the Angayucham terrane, (complex assemblage of oceanic rocks, including: gabbro, diabase, pillow basalt, tuff, chert, graywacke, argillite, and minor limestone; Mississippian to Jurassic sedimentary rocks; Late Carboniferous and Late Jurassic volcanic basalts; and thrust sheets of ultramafic rocks which are found throughout the section).

The Central province, a large area of plains, plateaus and rounded mountains, is geologically complex. The Central province was divided into five physiographic subprovinces by Wahrhaftig (1965). These are the Northern Plateaus, Western Alaska, Seward Peninsula, Aklun Mountains, and Bering Shelf subprovinces. More than 20 different terranes underlie the Central province (Jones and others, 1987).




Figure 3. Generalized geologic map of Alaska (Kinney, 1965).

The Seward Peninsula, on the western end of the Central province, consists mostly of Precambrian and Paleozoic metamorphic rocks. There is, however, an area of Precambrian and Paleozoic sedimentary rocks in the northwest part of the peninsula, a belt of Quaternary sediments along its northern edge, and an area of Tertiary and Quaternary mafic volcanic rocks generally in the center of the northeast part of the Peninsula. The Seward Peninsula is primarily underlain by the Seward terrane (Mica schist, micaceous calc-schist, metavolcanics, marble, and high grade gneissic rocks; rocks of probable Precambrian and known Devonian age are present but some ages are uncertain), but parts are underlain by the Crazy Mountains (Cambrian quartzitic sandstone and other younger Paleozoic sedimentary rocks) and Koyukuk terranes (andesitic volcanic rocks together with conglomerate, graywacke, and mudstone). Small parts of the Peninsula are underlain by other terranes (Jones and others, 1987).

The Bering Shelf, on the western end of the Central province and south of the Seward Peninsula, is covered almost entirely by Quaternary surficial sediments and scattered areas of Tertiary volcanic rocks. The Bering Shelf is underlain almost entirely by the Crazy Mountain terrane (see above).

The Aklun Mountains subprovince lies south of the Bering Shelf. It is covered mostly by Precambrian through Cretaceous marine sedimentary rocks, consisting of sandstone, shale, and limestone. The Aklun Mountain subprovince also contains small felsic intrusive bodies of Jurassic and Tertiary age. Southwest-trending faults and folds and at least one major southeast-trending fault are present in the Aklun Mountains. The subprovince is mostly underlain by the Togiak (Jurassic and Lower Cretaceous volcanic and volcanoclastic rocks including pillowed flows, tuffs, breccias, conglomerates, graywackes, chert and a minor amount of Cretaceous limestone) and GoodNews (chert, pillow basalt, tuff, limestone, and blocks of ultramafic rock; dated limestones vary from Ordovician to Permian; lawsonite-bearing metamorphic rocks in some areas) terranes.

The eastern third of the Central province is covered by the Northern Plateaus subprovince. About the western half of this subprovince underlies flat-lying Tertiary basin fill (nonmarine clastic rocks) and Quaternary surficial deposits. The northeastern part of the northern plateaus subprovince, along the Canadian border, is a complex area of Precambrian through Cretaceous mostly marine sedimentary rocks and Paleozoic metamorphic rocks that contain fairly large areas of Mesozoic intrusive and volcanic rocks. Much of the southern and southeastern parts of the subprovince is composed of a large variety of Paleozoic and Precambrian metamorphic rocks that were penetrated by felsic intrusions. The Tintina fault, which is generally followed by the Yukon River, trends southeast through the central part of the area. The Denali fault, which forms the southeast boundary of the subprovince, more or less parallels the Tintina. A complex grouping of terranes underlies most of the northern plateaus. The southeastern part is underlain by Yukon-Tanana (metamorphic rock including Paleozoic granitoid protoliths and Devonian marble) terrane. The northern part is underlain by Crazy Mountain (see above); Tozitna (Paleozoic (?) and Mesozoic gabbro, basalt, diabase, argillite, tuff, chert, and conglomerate; also includes Permian Limestone and Mississippian radiolarian cherts); and Porcupine terrane (phyllite, slate, quartzite and carbonate rocks overlain by sedimentary complex ranging from Cambrian (?) to Devonian in age) terranes. The western part of the Northern Plateaus province includes the Ruby (phyllite, schist, marble, quartzite, marble, amphibolite, granite, and metachert) and other terranes. The easternmost part of this province, near the Canadian Border, is underlain by phyllite, slate, and siltstone that is part of the North American craton (not an accreted terrane).

The largest subprovince of the Central province is Western Alaska (fig. 1). Large areas of Cretaceous and lower Paleozoic sedimentary and metamorphic rocks are present in the western half

of this subprovince. The Cretaceous rocks are mostly marine conglomerate, graywacke, sandstone, and siltstone. The lower Paleozoic rocks are both sedimentary and metamorphic and consist of limestone, dolomite, siltstone, argillite, chert, schist, quartzite, and greenstone. A large area of Cretaceous and Tertiary volcanic rock is present in the western part of this subprovince just east of the Seward Peninsula. Prominent faults cross the western Alaska subprovince from east to west. The Mulchatna, Kaltag, Farewell, and Iditarod-Nixon faults trend southwest through the western half of this subprovince. The Western Alaska subprovince is underlain by a large number of terranes. The largest of these are Koyukuk (andesitic flow tuffs, breccias, agglomerates, conglomerates, graywacke, and mudstone; locally interbedded limestone contains Early Cretaceous fossils); Kahiltna (Late Jurassic to Early Cretaceous deep marine graywacke, tuffs, pelitic rocks and minor amounts of chert, limestone, and conglomerate); Nixon Fork (reefal and platform carbonate rocks of Ordovician to Upper Devonian age overlain by Permian, Triassic, and Cretaceous sedimentary rocks); Dilinger (Paleozoic graptolitic shale, basinal carbonate rocks, and turbiditic sandstone and shale together with overlying Jurassic sandstone and siltstone); Innoko (Late Paleozoic and early Mesozoic chert, argillite, graywacke, and volcanic sandstone, conglomerate, and tuff); and Minchumina (Ordovician chert, argillite, and quartzite together with chert as young as Devonian).

The Alaska-Aleutian province includes the Aleutian Peninsula, a northeast-trending belt in south-central Alaska that includes Mt. McKinley, a southeast-trending belt that extends from the Mt. McKinley area south eastward to Canada, and the Coast Mountains in southeastern Alaska. On the Aleutian Peninsula from Unimak Pass westward, the bedrock consists mostly of Quaternary and Tertiary volcanic rocks and Tertiary sedimentary rocks. Quaternary and Tertiary volcanic rocks are also important east of the Unimak Pass but other rocks including Jurassic and Cretaceous sedimentary rocks and Jurassic intrusive rocks of intermediate and felsic composition are also important in this area. In addition, in the area west of Cook Inlet and southwest of Mount McKinley, large bodies of Tertiary mafic volcanics and Cretaceous or Jurassic intermediate intrusives are found. A varied assortment of Phanerozoic rocks is present in the area from Mt. McKinley southeastward to the Canadian border. This assortment includes Paleozoic mafic volcanic rocks and their sedimentary and metamorphic derivatives; Mesozoic mafic volcanic flows and tuffs, together with various units of shale, conglomerate, graywacke, and slate; and Tertiary and Quaternary intermediate volcanic rocks, Tertiary felsic intrusives, and Quaternary glacial deposits including eolian sand and silt. The Alaska-Aleutian province is mostly underlain in the west by the Peninsular (Paleozoic and Mesozoic igneous and sedimentary rocks including limestone, argillite, basalt, tuff, andesitic flows, breccias, volcanic sandstone and siltstone, and batholithic granitic rocks; also included is the andesitic arc assemblage), Kahiltna (see above), and Crazy Mountain terranes (see above), and in the north by the Yukon-Tanana terrane (see above). The Coastal Mountains in southeastern Alaska consist mostly of ultramafic, intermediate, and felsic intrusives, Mesozoic mafic volcanic rock, and Mesozoic through Paleozoic sedimentary rocks. The Tracy Arm (metamorphosed pelitic and quartzofeldspathic schist and paragneiss, marble, amphibolite, and minor serpentinite) and Taku (variably metamorphosed upper Paleozoic and Triassic basalt and local felsic volcanic rock, carbonate rock, and pelite; includes some undated metamorphosed clastic and volcanic rocks) terranes underlie most of the Coast Mountains.

The Coastal Trough province includes a row of basins that were active centers of deposition during the Cenozoic. The province includes the Cook Inlet and Copper River Basins in southern Alaska and the Admiralty Trough in southeastern Alaska. These depositional basins contain thick sequences of Tertiary continental clastic rocks that generally overlie Cretaceous or

older sedimentary rocks and have been penetrated and covered by Tertiary volcanic rocks. Mesozoic sedimentary rocks are abundant in the area north of Cook Inlet basin and large areas of Pleistocene, mostly glacial deposits, are found at the north end of Cook Inlet basin and in the Copper River basin. Large areas of Tertiary volcanic rocks are present in the vicinity of Shelikof Strait and the Wrangell Mountains in southern Alaska and on Admiralty, Kupreanof, and Wrangell Islands in southeastern Alaska. The coastal trough province is bounded by regional faults such as the Bruin Bay and Border Ranges faults that trend southwest in the Cook Inlet area and the Clarence Strait and Chatham Strait faults that trend southeast in southeastern Alaska. In southern Alaska the Coastal Trough province is underlain by Crazy Mountains (see above), Wrangellia (in ascending order, Upper Paleozoic arc-related volcanic breccias, flows, and clastic rocks; Permian limestone, pelitic rocks, and chert; Triassic black cherty argillite; a thick sequence of pillow basalt; basinal spiculitic argillaceous and calcareous rocks; and predominantly clastic rocks of Jurassic and Cretaceous age); Peninsular (see above); and Kahiltna (see above) terranes. In southeast Alaska much of the Coastal Trough province is underlain by the Stikinia terrane (Mississippian, Permian, Triassic and Jurassic marine and nonmarine volcanic and sedimentary rocks together with coeval granodioritic batholithic rocks).

The Border Ranges province is generally south and west of the coastal trough province. This province includes the Chilcat and Baranof Mountains and Prince of Wales Island in southeastern Alaska. Cretaceous and Jurassic sedimentary and metamorphic rocks with interbedded mafic volcanic rocks and some gabbro comprise most of the Border Ranges rocks. A fairly large area of lower Tertiary sedimentary, volcanogenic, and volcanic rocks is found in the Prince William Sound area. In southeastern Alaska, the Border ranges consist mostly of Paleozoic metamorphic and Paleozoic to Mesozoic sedimentary rock together with some intermediate intrusives of Cretaceous and Tertiary age. The metamorphic rocks are mainly Devonian Schist, phyllite, marble, and amphibolite. The sedimentary rocks are mainly Paleozoic shale, siltstone, graywacke, conglomerate, and limestone and a Cretaceous melange containing blocks of flysch, greenstone, limestone, chert, granodiorite, schist, layered gabbro, and serpentinite in a pelitic matrix. In southern Alaska the Border Ranges province is underlain primarily by the Chugach (mostly weakly metamorphosed Cretaceous graywacke and slate interbedded locally with radiolarian chert, gabbro, pillow basalt, and ultramafic rocks); Yakutat (Upper Mesozoic graywacke and shale intercalated with lenses of chert, argillite, and volcanic rocks); Ghost Rocks (strongly deformed assemblage of pillow lava, pillow breccia, and tuff with andesitic to basaltic composition interbedded with Late Cretaceous to Oligocene sandstone and mudstone and intruded by sparse plutons) terranes. In southeastern Alaska most of the Border Ranges province is underlain by the Alexander terrane which includes the Admiralty (Devonian and Mississippian basalt, carbonate rocks, and chert in contact with Ordovician flysch); Annette (variably metamorphosed Ordovician to Triassic intrusive, extrusive, clastic and carbonate rocks); and Craig (Pre-Ordovician metamorphic complex and Ordovician to Triassic mafic and felsic volcanic rocks and terrigenous clastic and carbonate rocks) subterranean.

Uranium deposits and radioactive anomalies: Table 1 lists the major uranium deposits and other significant uranium occurrences in Alaska. Both igneous and sedimentary types are present, but none are found in populated areas (fig. 4; table 1). Only the Bokan Mountain deposit has actually produced uranium (MacKevett, 1963). As presently known, only the Death Valley and Bokan Mountain deposits are potential U producers. In addition to the deposits listed in table 1, many small radioactive mineral occurrences (several as U and Th bearing placer deposits) and radioactive anomalies were reported by Eakins (1969, 1975). National Uranium Resource




Figure 4. Uranium deposits of Alaska.

Evaluation (NURE) surveys enumerated 1520 radioactive anomalies in an area that covers about 2/3 of the State (Dickinson and others, 1983). It should be noted, however, that uranium enrichment in the rock is not necessary to produce elevated radon to be in homes, and in the case of very young uranium deposits, even high concentrations of uranium may not cause elevated radon in homes because the uranium would be in gross disequilibrium with its daughter products.

TABLE 1. Uranium deposits in Alaska.

Area	Type of deposit	Reserves	Reference
Death Valley	Epigenetic, sandstone	1 M lbs. U ₃ O ₈	Dickinson (1988)
Bokan Mountain	Peralkaline granite	800 K lbs. U ₃ O ₈	MacKevett (1963)
Nenana Coal Field	Epigenetic and other	unknown	Dickson (1981)
Mount Prindle	Igneous	unknown	Armbrustmacher (1989)
Keku Strait	Epigenetic sandstone and phosphate	unknown	Dickinson and Vuletich (1990)

SOILS AND SURFICIAL DEPOSITS

The soils of Alaska are controlled largely by the cool moist climate and the rugged terrane. The soil terminology used below generally follows the nomenclature of the National Cooperative Soil Survey Classification of 1967. The distribution of soils is summarized from the National Atlas of the United States (U.S. Department of Agriculture, 1987; fig. 5).

About 2/3 of Alaska is covered by tundra. The tundra soils are largely aquepts and cryaquepts of the Inceptisol order. These soils are characterized by undecomposed plant material, generally high moisture content, and the presence of permafrost in the area generally north of the arctic circle (fig 6). Radon transmissivity in these soils is presumed to be poor because of the water and ice content. No large population centers are found in these areas.

Soils classified as rockland, which includes glacial ice, cover about 1/4 of Alaska (fig. 5). These surfaces are generally found in the rugged mountain ranges such as the Brooks, Alaska, Aleutian, Chugach, Kenai, Wrangell, St. Elias, and Coast Ranges. No significant populated areas in Alaska are found on rockland.

Another group of Inceptisols, cryochrepts (subarctic brown forest soils) and cryandepts (brown podzolic or gray-brown podzolic soils) cover about 5 percent of the Alaskan area. The cryochrepts are mostly distributed along the valleys of the Yukon and Tanana Rivers in central Alaska and the cryandepts are on the Alaskan Peninsula. These soils rank high in radon transmissivity, but are not common in populated areas.

Spodosols make up about 3-1/2 percent of the surface area of Alaska. These soils are characterized by a low base content and a high content of amorphous material, aluminum, and probably iron. They are formed under acid conditions, generally on porous parent material, and they probably have relatively high gas transmissivity. Anchorage and Juneau are located in areas with spodosols.




Figure 5. Generalized soils map of Alaska (after U.S. Department of Agriculture, 1987)

FIGURE 5 (continued). GENERALIZED SOILS MAP OF ALASKA
EXPLANATION

1. Cryaquepts (formerly Tundra). These are the Aquepts of cold climates. Aquepts are seasonally wet Inceptisols that have an organic surface horizon. Inceptisols are soils that have weakly differentiated horizons; materials in the soil have been altered or removed but have not accumulated. These soils are usually moist, but during the warm season of the year some are dry part of the time (66.5 percent).
2. Rock Land is a miscellaneous land type that includes Cryandepts (see below), Cryumbrepts (Umbrepts of cold regions), Cryaquepts (see above), Cryorthods (see below), ice fields, and glaciers. These are generally shallow soils formed on moderately sloping or steep slopes. Umbrepts are Inceptisols with crystalline clay minerals, thick dark-colored surface horizons, and altered subsurface horizons that have lost mineral materials and that are low in bases (24 percent).
3. Cryandepts (formerly brown podzolic or gray brown podzolic soils) are Andepts of cold regions. Andepts are Inceptisols (see above) that have formed in ashy materials, have low bulk density and large amounts of amorphous materials or both.
4. Cryorthods (formerly podzols)-- Orthods of cold regions. Orthods are Spodosols that have a horizon in which organic matter plus compounds of iron and aluminum have accumulated. Spodosols with low base supply that have in subsurface horizons an accumulation of amorphous materials consisting of organic matter plus compounds of aluminum and usually iron; formed in acidic, mainly coarse-textured materials in humid and mostly cool or temperate climates (3.5 percent).
5. Cryopsamments and Quartzipsamments (formerly regosols)-- Cryopsamments are Psamments of cold regions, Quartzipsamments are Psamments that consist almost entirely of minerals that are highly resistant to weathering, mainly quartz. Psamments are Entisols that have textures of loamy fine or coarser sand. Entisols are soils with no pedogenic horizons (1 percent)
6. Cryochrepts (former subarctic brown forest soils) are the Ochrepts of cold regions. Ochrepts are Inceptisols that have formed in materials with crystalline clay minerals, have light-colored surface horizons, and have altered subsurface horizons that have lost mineral materials (5 percent).

Entisols, soils with no pedogenic horizons, make up <1 percent of the Alaskan soils. They are the cryopsamments and quartzipsamments that are found only along the south central and southeast coastal plains. They probably have high gas transmissivity, but are not found in populated areas except for the village of Cordova (fig. 5).

Glacial deposits: Nearly half (48 percent) of Alaska was previously covered by glaciers and as a result nearly half of the State is covered by glacial deposits. About 7 percent of the land area is presently covered by glacial ice (fig. 6). Glacial outwash areas have more permeable sediment than do areas of till, but for most glaciated areas the distribution of various sediments has not been mapped.

Stream and River deposits: Stream and river sediments cover much of Alaska especially in delta and basinal areas. Examples are the Yukon flats area and the Yukon delta. There also large areas of Quaternary sediments along the drainages of most of the major rivers. These deposits are complex mixtures of fine- and coarse-grained sediment. In general the fine-grained sediment is impervious (has low gas transmissivity) and the coarse-grained sediments are pervious (have high gas transmissivity). No data is available to allow generalization on the distribution of fine versus coarse-grained sediment on an area by area basis.

INDOOR RADON DATA

Indoor radon data from Alaska are principally from the State/EPA Residential Radon Survey. This survey was conducted during the fall and winter of 1988 and 1989 by the Division of Geology and Geophysics (DGGS) of Alaska and the Environmental Protection Agency (EPA). The EPA analyzed the detectors and provided survey design and consultation through its contractor, Research Triangle Institute (RTI). DGGS provided information on demography, geology, and geography, selected participating households, and distributed the canisters (Nye and Kline, 1990).

The data for the Alaska indoor radon survey are organized by borough or by other area designations (fig. 7). Figure 8 shows the indoor radon concentration data for those areas where sufficient data was obtained and table 2 lists the indoor radon data for all areas in which data were collected in the State/EPA Residential Radon Survey. The southeast Fairbanks census area had the highest percent of radon values over 4 pCi/L. Forty-eight percent of the measurements in the southeastern Fairbanks area exceeded 4 pCi/L (fig. 8; table 2). Boroughs or other areas with average screening indoor radon levels exceeding 4 pCi/L in the State/EPA survey include Southeast Fairbanks, Yukon-Koyukuk, and Skagway-Yakutat-Angoon, although the latter is based on only three measurements (fig. 8; table 2) and should not be considered representative of all indoor radon levels in the area.




Figure 6. Permafrost and glaciers map of Alaska (Wahrhaftig, 1965).

TABLE 2. Screening indoor radon data from the EPA/State Residential Radon Survey of Alaska conducted during 1988-89. Data represent 2-7 day charcoal canister measurements from the lowest level of each home tested.

BOROUGH	NO. OF MEAS.	MEAN	GEOM. MEAN	MEDIAN	STD. DEV.	MAX	%>4 pCi/L	%>20 pCi/L
ANCHORAGE	282	1.0	0.6	0.6	1.6	16.4	3	0
FAIRBANKS NORTHSTAR	281	3.5	1.7	1.6	12.3	191.9	13	2
HAINES	12	1.0	0.5	0.6	1.2	4.1	8	0
JUNEAU	137	0.4	0.3	0.3	0.6	4.1	1	0
KENAI PENINSULA	135	2.6	1.1	1.3	3.9	26.3	18	1
KETCHIKAN GATEWAY	56	0.2	0.2	0.2	0.3	2.0	0	0
KODIAK ISLAND	27	0.4	0.2	0.2	0.6	2.4	0	0
MATANUSKA-SUSITNA	60	2.8	1.6	1.5	3.2	15.3	22	0
SITKA	24	0.3	0.2	0.1	0.4	1.8	0	0
SKAGWAY-YAKUTAT-ANGOON	3	4.2	2.6	5.3	3.3	6.9	67	0
SOUTHEAST FAIRBANKS	31	6.4	4.1	4.0	6.4	28.5	48	6
VALDEZ-CORDOVA	31	1.6	0.6	0.7	2.1	9.0	6	0
WRANGELL-PETERSBURG	35	0.3	0.2	0.2	0.5	2.6	0	0
YUKON-KOYUKUK	13	5.5	2.5	2.0	7.8	27.9	31	8

GEOLOGIC RADON POTENTIAL

The occurrence of radon in Alaska is related to bedrock type, surface sediment type, soil type, and fault locations. Black shale and phosphatic rocks commonly have relatively high uranium contents. Phosphate deposits were found in Paleozoic and Mesozoic rocks in the Arctic Foothills province and in southeastern Alaska (Patton and Matzko, 1959; Dickinson, 1979a, 1979b). In addition, black shale beds were found in the Arctic Foothills province. Black shales are generally high in U, but no data is available for these rocks. Felsic igneous rocks, scattered around Alaska (Forbes, 1975; Jones and Forbes, 1977), generally contain 5-7 ppm U, which is higher than the average of 3.5 ppm for crustal igneous rocks (Wedepohl, 1971). One example is the Darby pluton on the Seward Peninsula which contains about 7 ppm U and is believed to be the source of U for the Death Valley deposit (Dickinson and others, 1987). Similar rocks are found in other parts of Alaska (Jones and Forbes, 1977). The Darby pluton is, like the rock in the Arctic Foothills province, located in an area with very low population density.

There are examples of high radon potential along fault zones in the conterminous 48 states (Gundersen, 1991). Special attention should be paid to faults as radon hazards in Alaska because of their abundance. Many if not most of the nearly 60 tectonostratigraphic terranes are separated from one another by fault zones. Most of the faults are not in populated areas, however. One exception is the Knik Fault which runs along the east side of suburban Anchorage. The Knik Fault passes near a radiometric anomaly along Turnagain Arm near Anchorage. The anomaly is over Quaternary sediment, but the exact location of the fault in this area is not known. A relation between the fault and the anomaly could exist.



Figure 7. Boroughs and other subdivisions for plotting indoor radon data of Alaska.




Figure 8. Screening indoor radon data from the EPA/State Residential Radon Survey of Alaska, 1988-89, for boroughs with 5 or more measurements. Data are from 2-7 day charcoal canister tests. Histograms in map legends show the number of boroughs or other areas in each category. The number of samples in each borough (See Table 2) may not be sufficient to statistically characterize the radon levels of the areas, but they do suggest general trends. Unequal category intervals were chosen to provide reference to decision and action levels.

An eU radiometric map is not available for Alaska. However, the number of radioactive anomalies for each of the 1:250,000 quadrangle map areas in about-two thirds of the State are available (Dickinson and others, 1983; fig. 9). These results are based on the National Uranium Resource Evaluation (NURE) studies (LKB Resources, 1978a, 1978b, 1979). Most of the anomalies represent contrasts between rock types and in some cases between bedrock and ice or water. None of these anomalies has been identified as an important uranium deposit; however, their distribution is significant for radon evaluation. In each of 13 quadrangles from a total of 91, more than 40 anomalies were discovered (fig. 9). These anomalies can be grouped into two belts and two isolated areas. The largest anomaly belt begins in the Kateel River quadrangle and extends eastward through Melozitna, Tanana, Bettles, Beaver, Fort Yukon, Black River and Coleen. This area generally follows the drainage basin of the Yukon River and its tributaries, the Tanana, Porcupine, and Koyukuk. The other anomaly belt follows the Anchorage, Valdez and McCarthy quadrangles that are occupied by the Chugach and Wrangell Mountains and a southern extension of the Copper River Basin. The Medfra quadrangle, west of the Mt. McKinley quadrangle, by itself exhibited 58 anomalies, and the Tanacross quadrangle southeast of Fairbanks exhibited 42. The North Slope, Northern Foothills, Brooks Range, and the Alaska Peninsula were not covered by this study.

Figure 10 shows the uranium content of stream sediments based on analyses from the Geochemical Atlas (Weaver, 1983). This data is summarized as follows: A 200-mile-wide area of high values extends westward across the State from Norton Sound and the Seward Peninsula eastward through central Alaska, including most of the Brooks Range, to the Yukon Flats area and the Canadian Border. A smaller belt covers the Alaska Range with Mount McKinley approximately at its center. Small areas of intermediate values are found in the Chugach, Kenai, and Wrangell Mountains. Scattered areas of relatively high values are present in southeastern Alaska. Three belts of generally low values can be seen. These are approximately the North Slope, the Kuskokwin River drainage basin in the central part of the State, and the Cook Inlet-Matanuska Valley-Copper River basin area.

The following section is reproduced directly from Nye and Kline (1990):

"Interior Alaska has the highest proportion of homes with elevated radon concentrations as well as the individual homes with highest concentrations. In the interior, 3 percent of homes with the sample population had screening levels higher than 20 pCi/L and 17.6 percent of homes had radon screening levels that were higher than 4 pCi/L. Figure 3 [fig. 11, this report] summarizes the responses to a request for home site geographic information which was included with the report of test results that was sent to part participants in the survey. This figure shows that 30 to 35 percent of homes built in the hills around Fairbanks have elevated radon concentrations.

In the Fairbanks area, homes built in the hills adjacent to the valley floor with concrete slab or basement structures which are in contact with bedrock yielded the highest measured radon levels. These areas also included the highest proportion of homes with high radon. We do not yet know the proportion of homes with basements in contact with bedrock that do not have elevated radon concentrations. The data shown in Figure 3 [fig. 12, this report] homes located on hillside sites, includes homes which are built on thick accumulations of windblown glacial silt (loess). Thick accumulations of loess appear to be an effective barrier to radon migration. Homes built on alluvium from the Tanana and Chena Rivers are also at lower risk. High radon concentrations in homes which are sited in bedrock are likely to result from high fracture




Figure 9. eU anomalies map of Alaska. From LKB Resources Inc. (1978a, 1978b, 1979).




Figure 10. Uranium in stream sediments map of Alaska. Modified from Weaver (1983).




Figure 11. Histograms summarizing results of radon screening measurements (Nye and Kline, 1990).




Figure 12. Histograms summarizing radon screening measurements and local geography of Fairbanks area homes (modified from Nye and Kline, 1990).

permeability of the bedrock as well as relatively high uranium concentration in the schist which comprises local bedrock. Low radon concentration in homes built on loess and alluvium may reflect low soil gas permeability, low uranium concentrations of soils, or both.

Throughout interior Alaska homes constructed on bedrock seem to be at higher risk for elevated radon concentrations. However, homes built on coarse glacial outwash deposits which are well drained can also have high radon concentrations. Outwash gravels may have sufficient porosity and permeability to allow significant amounts of soil gas to migrate into overlying structures. Outwash gravel may also contain abundant pebbles and boulders of rock types with higher uranium concentrations, including granitic rocks.

Based on survey data, south-central Alaska includes communities with a significant proportion of homes with elevated radon concentrations. The maximum radon concentrations detected in south-central Alaskan homes, however, are not as high as those measured in the Fairbanks area. The geologic setting of south-central Alaskan homes with elevated radon concentrations appears to be diverse. In some cases high radon screening levels can probably be attributed to the presence of coarse, well-sorted, granite-rich outwash. In other cases such concentrations appear to reflect proximity to bedrock.

Survey data indicate that Anchorage, southeastern Alaska, and northern and western Alaska have very few homes with elevated radon concentrations. In Anchorage this may reflect the abundance of fine-grained glacio-fluvial and glacio-marine sediments which underlie most of the borough. Fine-grained sediments such as these are characterized by low permeability, slow soil gas transport, and much reduced reservoir volume of soil gas; thus reducing the amount of radon which can be drawn into the home. Some of the homes included in the survey were located in the hills surrounding Anchorage. The fact that these homes also have low radon concentrations suggests that the local bedrock has low concentrations of uranium. The lack of homes with high radon in Anchorage probably reflects low gas permeability and pore volume of local soils, and low rates of radon production.

The low concentrations of radon in homes throughout southeastern Alaska are not well understood, geologically. Many of the underlying bedrock types should have sufficient uranium to produce a significant amount of radon, and many homes are built on extremely porous soils and fractured bedrock. The sample base includes homes in Juneau built on coarse talus fans and homes in the Mendenhall Valley built on coarse outwash with a high proportion of granitic cobbles. Similar materials in other regions have relatively high source concentrations of radon in soil gas. A high proportion of southeastern homes are built on pilings. Since these homes were structurally ineligible for the survey they do not influence the data presented here.

The small number of homes with elevated radon concentrations in northern and western Alaska may be due to several factors, including local soil conditions such as shallow permafrost and high water tables. Also, most of the communities included in this survey are sited on fine grained alluvial deposits such as overbank silt, rather than in upland areas where bedrock is exposed at or near the surface. Additionally, air exchange is probably more efficient in small homes during routine entry and exit" (Nye and Kline, 1990).

The data on which this study is based "do not reflect the fact that most homes in rural Alaska were structurally ineligible for this survey because they are built on pilings. Also there are many Alaskan communities which do not include homes which were eligible for this study under EPA guidelines" (Nye and Kline, 1990).

A compelling geological reason for the high indoor radon values obtained from the Fairbanks area is difficult to determine. The bedrock in the hills around Fairbanks is Precambrian to Paleozoic metamorphic rock (formerly termed the Birch Creek Schist). No data is at hand to suggest that the Birch Creek Schist is an abnormally high radon producer in the vicinity of Fairbanks. The Fairbanks 1:250,000 quadrangle produced only two gamma-ray anomalies in studies carried out during the NURE program. The Anchorage quadrangle produced 43 anomalies and the Juneau quadrangle produced 12. It should be noted that these airborne gamma-ray anomalies are based on measurement of Bismuth-214 decay and are more closely related to radon than to uranium. If abnormal amounts of radon are being generated by a particular rock unit such as the Birch Creek Schist, these values should show up in airborne Bi-214 measurements.

One of the two airborne radioactivity anomalies in the Fairbanks quadrangle was obtained over Quaternary sediments about 35 miles northwest of Fairbanks, the other was obtained over the Mississippian Totatlinika Schist about 55 miles south of Fairbanks. In addition, Eakins (1969) reported four uranium occurrences in stream sediment concentrates in the Fairbanks quadrangle. One is in the Easter Dome area where minor granitic bodies intruded the Birch Creek Schist about 10 miles northwest of Fairbanks. Stream concentrates contained as much as 70 ppm U in this area. Another is in the Pedro Dome-Gilmore Dome area about 15 miles northeast of Fairbanks, where high Au, Bi, and W contents are found in lodes and high Au contents are found in placers that are associated with the Birch Creek Schist. Panned concentrates from this area contained up to 660 ppm U and outcrop samples contained as much as 10 ppm U (Eakins, 1969). Placer deposits are difficult to assess for radon potential because the amount of radon that escapes uraniferous resistate minerals through the uranium decay series is not well known. In addition, houses are not often built on placer deposits and houses built on the alluvium in the Fairbanks area were low in radon. Two other U occurrences are the Liberty Bell Mine, 64 miles southwest of Fairbanks, and at Grubstake Creek 57 miles southwest of Fairbanks. They are too remote to suggest high uranium values in the Fairbanks area. None of these occurrences suggest abnormally high radon potential from the Birch Creek Schist in the hills around Fairbanks.

In the Anchorage quadrangle, most of the 43 airborne radiometric anomalies identified during the NURE studies are in the Chugach Mountains, away from populated areas. One strong anomaly, however, was located along Turnagain Arm near residential areas of Anchorage. This anomaly, as noted above, is near the Knik fault. It is also located on alluvium that is related to Cretaceous and/or Upper Jurassic deep water clastic rocks consisting of siltstone, graywacke, arkose, conglomeratic sandstone, pillow basalt, and other rocks. This sequence and a melange of other Cretaceous and Jurassic (?) rocks including flysch, greenstone, limestone, chert, granodiorite, schist, layered gabbro, and serpentinite form the hills above Anchorage to the west. Houses in these hills generally have low indoor radon readings.

The Juneau quadrangle contains two NURE airborne radiometric anomalies. Only one of these was near Juneau. It is located over the waters of Taku Inlet about 10 miles east of Juneau. The anomaly is next to Mesozoic and Paleozoic onshore rocks and may have resulted from the migration of subaerial radon gas from these rocks. The rocks consist of greenstone, graphitic schist, slate, chert, graywacke, quartzite, phyllite interlayered with marble, and a small amount of amphibolite.

SUMMARY

The major physiographic provinces of Alaska are the land units used for radon potential estimates, except for the Central province, which is divided into the Northern Plateaus subprovince and an area made up of the remaining subprovinces (fig. 1). Each of these areas has a variety of geology and probably contains rock units that produce significantly high levels of radon. A brief summary of the estimated potential for radon in each province or group of subprovinces together with a radon potential index is given below (table 3).

The scores, particularly those for the confidence index, are low, in part because of dearth of data considering the large size of the areas. The indoor radon data is limited to populated areas and does not represent the entire land area of the province evaluated. Nevertheless, in the absence of adequate data these data are given some credibility. The aerial radioactivity is presented only in number of significant anomalies compiled by the NURE program for each 1:250,000 quadrangle area (LKB resources, Inc., 1978a, 1978b, 1979). The geology for these anomalies has been incompletely compiled, but is known in some cases. The aerial radiometric data is applicable within the constraints of the method of compilation. The geology for each province or subprovince is well known, but in some of the areas it is complex and indoor and/or soil-gas radon information is lacking. Broad assumptions on radon potential based on geology have been used for the score compilation. Broad assumptions on water and ice saturation of the soils and its effect on soil-gas transmissivity were also used in the evaluations. No points were assigned for geologic field evidence because no documented field studies of radon in the geologic environment in Alaska were available.

The relatively flat-lying Cretaceous and Tertiary sedimentary rocks that make up the Arctic Coastal Plain could contain high radon producers, although there is no data to suggest they do. A radon potential score (RI) of 6 and confidence index (CI) of 6 is given for this province. These low values stem mostly from lack of data and the assumption that soil saturation with water and ice reduces radon transmissivity (figs. 5, 6). No significant uranium occurrences are known in this area and the number of gamma-ray anomalies is low for this area when compared with other parts of Alaska (figs. 4, 9). The area also is a low in radioactive resistate minerals in stream sediments (fig. 10). The coastal plain is unglaciated and contains tundra soils and permafrost.

The folded Cretaceous sandstone and shale that makes up most of the Arctic Foothills province could produce relatively large amounts of radon but, again, no evidence that they do is at hand. The province was assigned an RI of 7 and a CI of 6. The area contains no known U deposits (fig. 4), and the part of the area where airborne gamma-ray measurements were made shows a relatively low number of anomalies (fig. 9). The stream sediments show varied amounts of uraniferous placer sediments with some high values in the eastern part of the area (fig. 10). Part of the area was covered by glaciers (fig. 6). The soils are tundra soils with permafrost and apparent low gas transmissivity (fig. 5).

The Precambrian and Paleozoic marine sedimentary rocks that make up most of the Arctic Mountains province probably are not producers of large amounts of radon inasmuch as there is little phosphate rock or black shale in these sequences. An RI of 9 and a CI of 6 was estimated for this area. There are no known significant uranium occurrences in this area. The stream sediments are, however, moderately high in uraniferous resistate minerals (fig. 10). The area has been glaciated (fig. 6), but much of the terrane is bare rock without surficial glacial material. The soils are classified as rockland, which includes glacial ice.

The Central province excluding the Northern Plateaus subprovince is a vast area of varied geology and undoubtedly contains many rocks that are significant radon producers and many that are not. The area was assigned an RI of 11 and a CI of 6. Indoor radon was given a rank of 3, but with a very low confidence level because of the small number of measurements for such a vast area. The number of gamma-ray anomalies ranged from one in the Lake Clark quadrangle to 71 in the Kateel River quadrangle (fig. 9), and the area was given 2 RI points in this category. For geology, the area was given 3 points. It contains uranium deposits of potentially commercial size at Death Valley on the Seward Peninsula and in the Healy Creek Coal basin (fig. 4). There are several areas of uraniferous granites, felsic intrusives, and volcanic rocks in parts of this province together with various uranium occurrences in the Darby Mountains (Foley and Barker, 1986). Nearly all of the area lies within the belt of uraniferous stream sediment minerals (fig. 10). Little of the Central Province has been glaciated. The soils are mostly of the tundra type with variable permafrost. There are significant areas of rockland and Cryochrepts (subarctic brown forest soils), which may have high gas transmissivity. Water and ice saturation probably limits soil-gas transmissivity in some parts of the area. Soil permeability was ranked 2.

The Northern Plateaus subprovince, a large, geologically complex area, was assigned an RI of 11 and a CI of 7. Three points were given for indoor radon because of the high levels in parts of the Fairbanks and Fairbanks-Northstar boroughs. The Northern Plateaus subprovince was separated from the rest of the Central province in order to emphasize the high indoor radon levels in the Fairbanks area, even though that area represents only a small part of the subprovince, and to point out that high indoor radon levels are likely to occur locally in other parts of the Northern Plateaus as well as the remainder of the Central Province, though both areas are ranked moderate in geologic radon potential overall. Radioactivity, as measured by the number of anomalies, was rated 2. The number of anomalies for 1:250,000 quadrangles covering the Northern Foothills ranged from lows of two each in Fairbanks and Big Delta to highs of 59 in Tanana and 63 in Beaver. The schist that apparently produces high indoor radon near Fairbanks is in the Northern Plateaus subprovince, but no airborne gamma-ray anomalies were found over the schist in the Fairbanks area. There are, however, 10 anomalies over the schist in the Tanacross quadrangle 130 to 200 miles to the southeast of Fairbanks. Felsic intrusives are scattered in two belts, one intruding Paleozoic and Precambrian metamorphic rocks in the southeast one third of the subprovince and one intruding Lower Paleozoic and (or) Precambrian sedimentary rocks along the northwest margin of the subprovince. For geology the subprovince was rated 3. The area contains one known significant uranium and thorium deposit at Mount Prindle (fig. 4; Armbrustmacher, 1989). Uranium is high in stream sediments in the south-central part and along the northwest border of the subprovince (fig. 10). Soil permeability was ranked 2. In order of decreasing abundance the soils consist of Cryaquepts (tundra), Cryochrepts (former subarctic brown forest soils), and rockland. Soil permeability is variable, depending on water and ice saturation. The Cryochrepts and rockland may have fairly high gas transmissivity.

The Alaska-Aleutian Range was assigned an RI of 9 and a CI of 6 overall (table 3), though the area contains some felsic intrusives and other rocks that could be locally significant radon producers. Indoor radon was given a RI score of one and confidence index of one. No significant uranium occurrences are known in this province. The number of radiometric anomalies is relatively low. The McKinley quadrangle, for instance, only has six (fig. 9). The province, however, is an intermediate to high producer of uraniferous stream sediments (fig. 10). Most of the area is or was covered by glaciers. Soils are mostly classified as rockland or tundra (figs. 5, 6).

The Coastal Trough contains thick sequences of carbonized wood-bearing Tertiary continental clastic rocks similar to units that produce uranium in the western conterminous United States. Uranium content of these rocks is not high, however, especially in the Cook Inlet and Copper River basins. There is a slight northward increase in uranium content of the Tertiary sedimentary rocks in the Cook Inlet basin and small uranium occurrences are found in the Susitna Lowlands (Dickinson and Campbell, 1978). There are also small uranium deposits in the Admiralty trough in southeastern Alaska (Dickinson and Campbell, 1984; Dickinson and Vuletich, 1990). Stream sediments are low in radioactive minerals in the Cook Inlet and Copper River basin areas, and data from Admiralty Trough, where there are very few surface exposures, is sparse. Soils are mostly brown and gray-brown podzolic forest soils which could have high gas transmissivity. Heavy rainfall in southeast Alaska may retard soil gas migration. The Coastal Trough area was given an RI of 9 and a CI of 6 (table 3).

The rocks of the Border Ranges, which consist mostly of Cretaceous and Jurassic sedimentary and metamorphic rocks, together with some mafic volcanic and intrusive rocks, generally have low radon potential. The RI for this area is 7 and the CI is 6 (table 3). The uranium deposit at Bokan Mountain is associated with a peralkaline granite and it illustrates that because of the great geologic diversity some high radon producers are undoubtedly present. Uraniferous mineral content of stream sediments is intermediate for the Border Ranges, although data are absent for many parts of the area (fig. 10). The radioactive anomaly density is also low for the Border Ranges, although the Port Alexander quadrangle contains 17 and over half of the area is ocean water. Podzolic brown and gray-brown forest soils are common in the Border Ranges and they could have high gas transmissivity. In southern Southeastern Alaska, where annual precipitation is about 14 feet, water saturation probably retards gas flow in soils. Nearly all of the Border Ranges province was glaciated or is presently occupied by glaciers.

This is a generalized assessment of the State's geologic radon potential and there is no substitute for having a home tested. The conclusions about radon potential presented in this report cannot be applied to individual homes or building sites. Indoor radon levels, both high and low, can be quite localized, and within any radon potential area there will likely be areas with higher or lower radon potential than assigned to the area as a whole. Any local decisions about radon should not be made without consulting all available local data. For additional information on radon and how to test, contact your State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

TABLE 3. RI and CI scores for geologic radon potential areas of Alaska.

PROVINCE	Arctic Coastal Plain		Arctic Foothills		Arctic Mountains		Central*	
FACTOR	RI	CI	RI	CI	RI	CI	RI	CI
INDOOR RADON	1	1	1	1	2	1	3	2
RADIOACTIVITY	1	1	2	1	2	1	2	1
GEOLOGY	2	2	2	2	2	2	3	2
SOIL PERM.	1	2	1	2	2	2	2	1
ARCHITECTURE	1	-	1	-	1	-	1	-
GFE POINTS	0	-	0	-	0	-	0	-
TOTAL	6	6	7	6	9	6	11	6
RANKING	LOW	LOW	LOW	LOW	MOD LOW		MOD	LOW

PROVINCE	Northern Plateaus		Alaska-Aleutian Ranges**		Coastal Trough		Border Ranges	
FACTOR	RI	CI	RI	CI	RI	CI	RI	CI
INDOOR RADON	3	2	2	2	2	2	1	2
RADIOACTIVITY	2	1	2	1	2	1	2	1
GEOLOGY	3	2	2	2	2	2	2	2
SOIL PERM.	2	2	2	1	2	1	1	1
ARCHITECTURE	1	-	1	-	1	-	1	-
GFE POINTS	0	-	0	-	0	-	0	-
TOTAL	11	7	9	6	9	6	7	6
RANKING	MOD	MOD	MOD	LOW	MOD	LOW	LOW	LOW

*Excluding Northern Plateaus

**including the Coast Mountains

RADON INDEX SCORING:

Radon potential category	Point range	Probable screening indoor radon average for area
LOW	3-8 points	< 2 pCi/L
MODERATE/VARIABLE	9-11 points	2 - 4 pCi/L
HIGH	> 11 points	> 4 pCi/L

Possible range of points = 3 to 17

CONFIDENCE INDEX SCORING:

LOW CONFIDENCE	4 - 6 points
MODERATE CONFIDENCE	7 - 9 points
HIGH CONFIDENCE	10 - 12 points

Possible range of points = 4 to 12

REFERENCES CITED IN THIS REPORT
AND GENERAL REFERENCES PERTAINING TO RADON IN ALASKA

- Armbrustmacher, T.J., 1989, Minor element content, including radioactive elements and rare-earth elements, in rocks from the syenite complex at Roy Creek, Mount Prindle area, Alaska: U.S. Geol. Survey, Open-File Report 89-0146, 11 p.
- Beikman, H.M., (Compiler), 1980: Geologic Map of Alaska: U.S. Geological Survey, scale 1:2,500,000.
- Bennison, A.P.(Compiler), 1974, Geologic highway map of the State of Alaska and the State of Hawaii: American Association of Petroleum Geologists, scale 1:3,500,000
- Conner, Cathy, and O'Haire, Daniel, 1988, Roadside geology of Alaska: Mountain Press Publishing Co., Missoula, Montana, 250 p.
- Dickinson, K.A., 1978, Uraninite in siderite nodules from Tertiary continental sedimentary rocks in the Healy Creek basin area, central Alaska: U.S. Geological Survey Circular 804B, p. B98-B99.
- _____, 1979a, Uraniferous phosphate occurrence on Kupreanof Island, southeast Alaska: U.S. Geological Survey, Open-File Report, 2 p.
- _____, 1979b, A uraniferous occurrence in the Tertiary Kootznahoo Formation on Kuiu Island, southeast Alaska: U.S. Geological Survey, Open-File Report 79-1427, 5 p.
- _____, 1984, Uranium geology of the Tertiary Kootznahoo Formation of the Admiralty Trough, southeastern Alaska: Journal of the Alaskan Geological Society, v. 4, p. 1-11.
- _____, and Campbell, J.A., 1978, Epigenetic mineralization and areas favorable for uranium exploration in Tertiary continental sedimentary rock in south-central Alaska-A preliminary report: U.S. Geological Survey Open-File Report 78-757, 13 p.
- _____, 1984, Uranium geology of the Tertiary Kootznahoo Formation of the southern part of the Admiralty Trough, Southeastern Alaska: Alaska Geological Society Journal, v. 4, p. 1-11.
- _____, and Vuletich, April, 1990, Diagenesis and uranium mineralization of the Lower Tertiary Kootznahoo Formation in the northern part of Admiralty Trough, Southeastern Alaska: U.S. Geological Survey Bulletin 1888, 12 p.
- _____, Cunningham, K.D., and Ager, T.A., 1987, Geology and origin of the Death Valley uranium deposit, Seward Peninsula, Alaska: Economic Geology, v. 82, pp. 1558-1574.
- _____, Morone, J.F., and Roberts, M.E., 1983, Summary of radiometric anomalies in Alaska: U.S. Geological Survey, Open-File Report 83-169.
- Dickson, R.K. 1981, Uranium mineralization in the Nenana Coal Field, Alaska *in* short notes on Alaskan Geology: Division of Geological and Geophysical Surveys, Geologic Report 73, p. 37-42.

- Eakins, G.R., 1969, Uranium in Alaska: Division of Mines and Geology, Department of Natural Resources, State of Alaska, Report number 38, 49 p.
- _____, 1975, Investigations of Alaska's uranium potential: Alaska State Department of Natural resources, Division of Geological and Geophysical Surveys, GJO-1627, Energy research and development Administration, 37 p.
- _____, 1977, Investigation of Alaska's uranium potential, Part 1, Reconnaissance program, west-Central Alaska and Copper River Basin: Alaska State Department of Natural Resources, Division of Geological and Geophysical Surveys, U.S. Energy Research and Development Administration, Grand Junction Colorado, 58 p.
- Fleischer R.L., and Mogro-Campero, Antonio, 1985, Association of subsurface radon changes in Alaska and the northeastern United States with earthquakes: *Geochimica et Cosmochimica Acta*, v. 49, p.1061-1071.
- Foley J.Y., and Barker, J.C., 1986, Uranium Occurrences in the Northern Darby Mountains, Seward Peninsula, AK: U.S. Bureau of Mines Information Circular, 26 p.
- Forbes, R.B., 1975, Investigation of Alaska's uranium potential, map of the granitic rocks of Alaska, regional distribution and tectonic setting of Alaskan alkaline intrusive igneous rocks: GJO-1627, Part 2, 44p.
- Gundersen, L.C.S., 1991, Radon in sheared metamorphic and igneous rocks, *in* Gundersen, Linda C.S., and Richard B. Wanty, eds., *Field studies of radon in rocks, soils, and water*: U.S. Geol. Survey Bulletin 1971, p. 39-50.
- Howell, D.G. (editor), 1985, Tectonostratigraphic terranes of the Circum-Pacific region: Council for Energy and Mineral Resources, Earth Science Series no. 1, p. 3-30.
- Jones, B.K., and Forbes, R.B., 1977, Investigation of Alaska's uranium potential, Part 2, Uranium and thorium in Granitic and alkaline rocks in western Alaska: Alaska State Department of Natural Resources, Division of Geological and Geophysical Surveys, 66 p.
- Jones, D.L., Silberling, N.J., Berg, H.C., and Plafker, George, 1981, Map showing tectonostratigraphic terranes of Alaska, columnar sections, and summary description of terranes: U.S. Geological Survey Open-File Report 81-792, 20 p.
- Jones, D. L., Silberling, N. J., Coney, P. J., and Plafker, George, 1987, Lithotectonic terrane map of Alaska (west of the 141st Meridian): U.S. Geological Survey Miscellaneous Field Studies Map MF-1874-A.
- Kinney, D.M., 1966, Geology: U.S. Geological Survey, National Atlas of the United States of America, Sheet no. 74.
- Larson, R.E., 1974, Radon profiles over Kilauea, the Hawaiian Islands and Yukon Valley Snow Cover: *Pure Applied Geophysics*, v. 112, p. 204-208.

- LKB Resource Inc., 1978a, NURE aerial gamma ray and magnetic reconnaissance survey, Cook Inlet, Alaska area: U.S. Department of Energy, Open-File Report GJBX-108 (78), v. 1-2.
- _____, 1978b, NURE aerial gamma ray and magnetic reconnaissance survey, Eagle-Dillingham area, U.S. Dept. of Energy, Open-File Report, GJBX-113 (78), v. 1-2.
- _____, 1979, NURE aerial gamma ray and magnetic reconnaissance survey, southeastern area, Alaska: U.S. Dept. of Energy, Open-File Report, GJBX-48 (79), v. 1-2.
- Monger, J.W.H., and Berg, H.C., 1987, Lithotectonic terrane map of western Canada and southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1874-B.
- Nye, C.J., and Kline, J.T., 1990, Preliminary evaluation of data derived from the recently completed Alaska home radon survey: Alaska Division of Geological and Geophysical Surveys, Public Data File 90-6, 14 p.
- MacKevett, E.M., Jr., 1963, Geology and ore deposits of the Bokan Mountain Uranium-Thorium Area, Southeastern Alaska: U.S. Geological Survey, Bulletin 1154, 125 p.
- Patton, W.P., and Matzko, J.J., 1959, Phosphate deposits in northern Alaska: U.S. Geological Survey, Professional Paper 302-A, 17 p.
- Staatz, M.H., Hall, R.B., Macke, D.L., and Brownfield, I.K., 1980, Thorium resources of selected regions in the United States: U.S. Geological Survey Circular 824, 32 p.
- Stone, David B., and Wallace Wesley K., 1987, A geological framework of Alaska: Episodes, v. 10, no. 4, p. 284-289.
- U.S. Department of Agriculture, Soil Conservation Service, 1987, Soils: U.S. Geological Survey National Atlas sheet 38077-BE-NA-07M-00, scale 1:7,500,000.
- Wahrhaftig, Clyde, 1965, Physiographic divisions of Alaska: U.S. Geological Survey, Professional Paper 482, 52 p.
- Weaver, T.A. (project leader), 1983, Geochemical atlas of Alaska: Los Alamos National Laboratory, Los Alamos, New Mexico, LA-9897-MS, UC-51, 49 plates.
- Wedepohl, K.H., 1971, Geochemistry: New York, Holt, Rhinehart, Winston, 231 p.
- Wiess, H.V., and Naidu, A.S., 1986, 210 Pb Flux in an Arctic coastal region: Arctic, v. 39, p. 59-64.
- Wilkening, M.H., Clements, W.E., and Stanley, D., 1972, 38 Radon-222 flux measurements in widely separated regions, *in* the Natural radiation environment II, Conference 720805-P2: United States Energy Research and Development Administration, v. 2, p 717-729.

PRELIMINARY GEOLOGIC RADON POTENTIAL ASSESSMENT OF IDAHO

by
Douglass E. Owen
U.S. Geological Survey

INTRODUCTION

Idaho is located in the Pacific Northwest (fig. 1A) and politically is divided into a number of counties (fig. 1B). Idaho is a rural state and farming, grazing, forestry, and mining are major economic activities. Figure 1C shows the population distribution within the state.

This is a generalized assessment of geologic radon potential of rocks, soils, and surficial deposits of Idaho. The scale of this assessment is such that it is inappropriate for use in identifying the radon potential of small areas such as neighborhoods, individual building sites, or housing tracts. Any localized assessment of radon potential must be supplemented with additional data and information from the locality. Within any area of a given radon potential ranking, there are likely to be areas with higher or lower radon levels than characterized for the area as a whole. Indoor radon levels, both high and low, can be quite localized, and there is no substitute for testing individual homes. Elevated levels of indoor radon have been found in every State, and EPA recommends that all homes be tested. For more information on radon, the reader is urged to consult the local or State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

GEOGRAPHIC SETTING

Idaho is divided into 4 physiographic provinces (fig. 2). The largest province, the Northern Rocky Mountain Province, covers about half of the state and includes most of northern and central Idaho. This province is characterized by high mountains and deep intermontane valleys. The next largest province is the Columbia Intermontane Province, an area of plains and plateaus. This province contains six sub-provinces, or sections, which are shown and described on figure 2. Southeastern Idaho contains a triangle-shaped desert area 100-130 miles on a side, which is part of the Basin and Range Province. This province is characterized by block-faulted mountains separated by gravel-floored basins. The Middle Rocky Mountain Province forms a 10-35 mile wide strip along the Wyoming border and is characterized by 2 landform types. The first type is the Yellowstone Plateau, which is of volcanic origin, and the second is the complexly faulted and folded mountain ranges found in the southeastern corner.

Soil moisture conditions affect both radon emanation and transport (see introduction to this volume). Because of this, a brief summary of Idaho's precipitation patterns is presented below. The Columbia Intermontane Province is largely arid, receiving less than 10 inches of rain per year (fig. 3). The Northern Rocky Mountain Province receives from 20 to more than 60 inches of precipitation annually (fig. 3). The Middle Rocky Mountain Province and the Basin and Range Province generally receive between 10 and 30 inches of precipitation per year (fig. 3). Areas with greater than 30 inches of precipitation per year probably experience some capping effects and also blockage of gas transport through soil pores during periods when soil moisture levels are high.

IDAHO'S

LOCATION



(modified from Caldwell, 1970)

Fig. 1A Location Map



Fig. 1B Counties





Figure 1C. Population of counties in Idaho (1990 U.S. Census data).

Physiographic Provinces






Fig. 3. Precipitation Map

GEOLOGIC SETTING

The geology of Idaho is dominated by intrusive and extrusive igneous rocks (fig. 4). The largest intrusive body, the Idaho batholith (fig. 5), forms a major part of the Northern Rocky Mountain Province. The Idaho batholith is a complex of granitic rock units ranging from diorite to granite (Mackin and Schmidt, 1957). North of the batholith in the panhandle of the state Precambrian metamorphic rocks dominate (fig. 4). The Columbia Intermontane Province is dominated by extrusive rocks (igneous rocks that have cooled at or near the surface); it is floored by a series of nearly horizontal basalt flows. The Yellowstone Plateau (Middle Rocky Mountain Province) also contains a large number of volcanic rocks. A number of the valleys in the Northern Rocky Mountain Province contain glacial lake deposits (fine grained) that accumulated behind ice dams and (or) glacial flood deposits (coarse grained) that were laid down when ice dams were breached.

SOILS

Idaho has 10 major soil types (fig. 6), but more than 600 soils have been mapped in Idaho (Caldwell, 1970) making generalizations about soil permeabilities difficult. Some soils found in Major Soil Type #4 (fig. 6) are clayey (Barker and others, 1983) and somewhat poorly drained (possible low permeability), but develop deep cracks on drying (secondary porosity--increased gas permeability). Almost all (44 of 54) of the soils and soil series described in the Idaho Soils Atlas are listed as either well-drained or excessively-drained (Barker and others, 1983, p. 132). The slope of the land influences drainage (i.e., steep slopes increase drainage) and enters into the determination of a soil's drainage class along with the soil's permeability. Taking slope into consideration, it is not unreasonable to assume that these well-drained and excessively drained soils have at least moderate permeabilities and allow diffusive transport of radon. Highly permeable soils allow convective transport of radon (Tanner, 1964; Sextro and others, 1987). Soil permeabilities greater than 6 inches per hour (listed in soil surveys) are considered highly permeable. Duval and others (1989), in a study for the Bonneville Power Administration, examined the available soil surveys for Idaho. Figure 7 shows the counties for which detailed soil surveys were available and examined for permeability data, and figure 8 shows the areas larger than a few square miles that were determined to have highly permeable soils. Tables in Duval and others (1989) list the soil associations and the specific intervals that were described in the soil reports as having permeabilities greater than 6 inches per hour. The following counties contained soil associations with intervals in the highly permeable range: Ada, Bannock, Benewah, Bingham, Blaine, Bonneville, Bonner, Boundary, Canyon, Camas, Cassia, Caribou, Gem, Idaho, Kootenai, Latah, Lincoln, Madison, Minidoka, Owyhee, Power, and Teton. Many of these highly permeable soils have formed on windblown sand, alluvial fans, river terraces, or on floodplains.

Geology




Fig. 4. Geologic Map




Fig. 5. Map showing Idaho batholith and Phosphoria Formation.

MAJOR SOILS




Fig. 6. Soils Map:

INDOOR RADON DATA

Figure 9 graphically summarizes the indoor radon data acquired during 1989-90 in the State/EPA Residential Radon Survey. Most of the counties (24) in Idaho had an insufficient number (less than five) of radon measurements to show on the maps (fig. 9). In six counties (Butte, Caribou, Custer, Lemhi, Madison, and Teton), more than 40 percent of the homes tested had indoor radon concentrations greater than 4 pCi/L. Eight counties (Bear Lake, Caribou, Cassia, Custer, Kootenai, Lemhi, Madison, and Teton) had average (mean) indoor radon concentrations greater than 4 pCi/L.

Table 1 presents a summary of the State/EPA indoor radon data. Table 1 lists the following: the number of measurements in each county; the mean, geometric mean, median, and standard deviation for each county; the maximum indoor radon concentration sampled in each county; the percentage of homes with indoor radon concentrations greater than 4 pCi/L for each county; and the percentage of homes with indoor radon concentrations greater than 20 pCi/L. In only 6 counties (Blaine, Cassia, Kootenai, Madison, Shoshone, and Teton) with 5 or more measurements, more than 1 percent of the homes tested had indoor radon concentrations greater than 20 pCi/L (Table 1).

The Idaho Department of Health and Welfare conducted a study of indoor radon in the State (Idaho Department of Health and Welfare, 1988). Volunteers were solicited for this study and therefore the degree of randomness of the sampling is not known. Also, many kinds of homes were tested including mobile homes, condominiums, apartments, and earth-covered homes. Considering the above information, the statistical representativeness of these data is uncertain, nevertheless their results show means greater than 4 pCi/L for Adams, Bear Lake, Blaine, Butte, Caribou, Cassia, Custer, Elmore, Franklin, Kootenai, Lemhi, Madison, Shoshone, and Teton counties. This data set was not used in the radon matrix.

GEOLOGIC RADON POTENTIAL

Areas in the vicinity of known uranium occurrences have a high radon potential for several reasons other than the unlikely occurrence that homes would be built over an ore body itself: (1) Noncommercial concentrations of uranium are often also present in an area that contains ore grade deposits; (2) Even minor mineralization (primary or secondary) of uranium along faults and fractures is commonly enough to produce a radon hazard in homes built above them; (3) Sediments shed and transported from rocks with elevated uranium are also likely to have elevated uranium levels. Figure 10 shows known uranium deposits in Idaho. The Phosphoria Formation, which contains beds of uranium-bearing phosphate rock, is exposed in southeastern Idaho (fig. 5) and contains up to 600 parts per million (ppm) uranium (Cook, 1957). Up to 530 ppm uranium has been found in uranium-bearing coal, carbonaceous shale, and limestone in the Caribou Mountains east of Idaho Falls in Bonneville County. Uranium-bearing lignites also occur in Payette, Twin Falls, and Cassia Counties (Cook, 1957). Placer deposits (Black Sand Districts--Fig. 10) lie within and around the borders of the Idaho batholith (fig. 5). Armstrong (1957) believes that the uranium-bearing black minerals are weathering from pegmatites within the batholith, while Kiilsgaard and Hall (1986) believe that they are coming from granitic rocks in the batholith. Up to 1,000 ppm uranium has been found in concentrates associated with gold extraction from these placers (Armstrong and Weis, 1957).




Fig. 7

Map showing counties in the Pacific Northwest for which soil surveys were available for pre-1900 data. Counties studied are lined. Many soil surveys did not cover all of the county but focused on the populated and privately owned parts of the county. Indian reservations, national forest land, and other government lands within many of the counties are not included.

(from Duval and others, 1989)

COUNTY KEY

Washington: 1-Whitman; 2-Okanogan; 3-Ferry; 4-Stevens; 5-Snell; 6-Clallam; 7-Snohomish; 8-Chelan; 9-Douglas; 10-Lincoln; 11-Spokane; 12-Jefferson; 13-Fitzgerald; 14-King; 15-Grays Harbor; 16-Thurston; 17-Pierce; 18-Kittitas; 19-Grant; 20-Adams; 21-Whitman; 22-Pacific; 23-Lewis; 24-Takima; 25-Pendleton; 26-Clark; 27-Whitman; 28-Coville; 29-Bhannia; 30-Clark

Idaho: 1-Boundary; 2-Bonner; 3-Rooten; 4-Beneish; 5-Latah; 6-Idaho; 7-Cam; 8-Canyon; 9-Bolae; 10-Cass; 11-Blaine; 12-Bingham; 13-Bonneville; 14-Madison; 15-Teton; 16-Lincoln; 17-Minidoka; 18-Power; 19-Bannock; 20-Caribou; 21-Cassia; 22-Blaine; 23-Phillips; 24-Missoula; 25-LeWelle; 26-Clark; 27-Whitman; 28-Pacific; 29-Lewis; 30-Coville; 31-Bannock; 32-Clark; 33-Columbia; 34-Tillamook; 35-Washington; 36-Multnomah; 37-Hood River; 38-Wasco; 39-Yacholt; 40-Clackamas; 41-Clallam; 42-Morrow; 43-Umatilla; 44-Polk; 45-Marion; 46-Grant; 47-Jefferson; 48-Lane; 49-Crook; 50-DeWitt; 51-Curry; 52-Imperial; 53-Klamath




Fig. 8. Map showing areas of highly permeable soil associations for the soil surveys in the counties shown in Figure 1A. Very small areas (less than 100 square miles) of highly permeable soils are indicated by the shaded areas. These areas are shaded and designed to show the general area where highly permeable soils dominate in the study area. These highly permeable soils include both rapidly permeable (6 to 20 in/hr) and very rapidly permeable (>20 in/hr) soils. The reader is referred to Appendix 1 for a survey by survey discussion of the highly permeable soils and to the original soil surveys for more exact location information.





Figure 9. Screening indoor radon data from the State/EPA Residential Radon Survey of Idaho, 1989-90, for counties with 5 or more measurements. Data are from 2-7 day charcoal canister tests. Histograms in map legends show the number of counties in each category. The number of samples in each county (See Table 1) may not be sufficient to statistically characterize the radon levels of the counties, but they do suggest general trends. Unequal category intervals were chosen to provide reference to decision and action levels.

TABLE 1. Screening indoor radon data from the EPA/State Residential Radon Survey of Idaho conducted during 1989-90. Data represent 2-7 day charcoal canister measurements from the lowest level of each home tested.

COUNTY	NO. OF MEAS.	MEAN	GEOM. MEAN	MEDIAN	STD. DEV.	MAXIMUM	%>4 pCi/L	%>20 pCi/L
ADA	769	2.8	1.9	2.1	2.8	29.5	18	0
ADAMS	4	1.2	0.9	1.1	0.9	2.3	0	0
BANNOCK	16	3.0	1.8	2.1	2.8	10.4	19	0
BEAR LAKE	10	4.6	3.6	3.4	3.0	8.8	40	0
BENEWAH	3	0.3	0.3	0.4	0.2	0.5	0	0
BINGHAM	10	2.9	1.9	2.0	2.7	8.3	20	0
BLAINE	6	8.4	5.5	7.6	7.4	21.4	67	17
BOISE	1	4.5	4.5	4.5	0.0	4.5	100	0
BONNER	17	3.1	1.0	0.8	5.3	16.7	18	0
BONNEVILLE	21	2.6	1.3	1.6	3.6	13.3	10	0
BOUNDARY	6	2.2	1.7	2.2	1.2	3.9	0	0
BUTTE	7	3.9	1.9	3.6	3.6	10.9	43	0
CAMAS	1	3.9	3.9	3.9	0.0	3.9	0	0
CANYON	37	1.7	1.0	1.2	1.8	9.8	5	0
CARIBOU	19	5.8	4.4	5.7	4.0	13.8	58	0
CASSIA	15	10.1	2.9	5.9	16.1	59.3	60	13
CLARK	2	5.9	5.9	5.9	0.6	6.3	100	0
CLEARWATER	8	3.4	1.1	2.4	4.9	14.6	25	0
CUSTER	5	5.9	4.6	5.1	4.0	12.0	80	0
ELMORE	4	0.9	0.6	0.9	0.7	1.7	0	0
FRANKLIN	16	4.1	2.4	2.4	4.2	15.0	25	0
FREMONT	1	2.6	2.6	2.6	0.0	2.6	0	0
GEM	5	2.0	1.4	1.5	1.6	4.0	0	0
GOODING	5	4.7	1.2	1.3	8.0	18.7	20	0
IDAHO	18	1.6	0.5	0.6	2.2	6.8	17	0
JEFFERSON	2	3.0	2.1	3.0	2.9	5.0	50	0
JEROME	3	0.9	0.5	0.4	1.0	2.0	0	0
KOOTENAI	29	7.0	2.5	2.6	10.0	49.0	41	3
LATAH	55	2.1	0.8	0.9	3.5	18.1	13	0
LEMHI	17	4.1	3.4	3.0	2.6	10.5	41	0
LEWIS	2	13.5	11.2	13.5	10.7	21.1	100	50
LINCOLN	1	3.0	3.0	3.0	0.0	3.0	0	0
MADISON	32	6.4	3.8	4.8	6.1	23.5	59	6
MINIDOKA	9	1.6	1.2	1.5	1.0	3.9	0	0
NEZ PERCE	50	1.9	1.3	1.6	1.5	5.9	16	0
OWYHEE	2	5.8	1.8	5.8	7.8	11.3	50	0
PAYETTE	5	0.8	0.6	0.8	0.4	1.2	0	0
POWER	2	0.6	0.2	0.6	0.8	1.1	0	0
SHOSHONE	11	6.7	2.8	2.8	9.3	30.1	45	9
TETON	6	14.9	4.6	8.1	20.0	54.3	83	17
TWIN FALLS	30	1.7	1.1	1.5	1.3	5.0	7	0
WASHINGTON	4	5.4	3.5	4.5	5.0	11.4	50	0



(modified from Breckenridge and others, 1980)

Fig. 10. Uranium Deposits

Figure 11 is a radiometric map of Idaho showing equivalent uranium (eU) in surficial deposits. The average uranium content for the upper continental crust is 2.5 ppm (Carmichael, 1989). In comparison, the Idaho batholith and the rocks to the west of the batholith have low radiometric signatures (less than 2 ppm eU). The panhandle has a mixture of low and higher uranium concentrations. The eastern part of the Northern Rocky Mountain Province, the Columbia Intermontane Province, the Basin and Range Province, and the Middle Rocky Mountain Province all have large proportions of their areas indicating uranium concentrations higher than the upper continental crust average.

Ogden and others (1987) concluded that areas of suspected higher radon levels include homes built on granite, gneiss, or valleys filled with erosional products of these rock types. Also suspect are homes built on phosphate-rich rocks and processed ore of the Phosphoria Formation in southeastern Idaho, as well as the geothermal areas of central Idaho. Their data also suggested that radon levels may be higher in homes located on or near faults.

Figure 12 shows the soil-gas radon potential estimated by Duval and others (1989); they believe that most of the state has a moderate radon potential in soil gas (based strictly on radiometric data). Where the Columbia River Basalt Group extends into Idaho (fig. 4) they believe there is a low radon potential. Their high potential areas seem to compare well with the areas that have a 3.5 ppm or greater eU signature on the radiometric map (fig. 11).

Comparing figure 9 and Table 1 to figures 2, 4, and 5, there appears to be a good correlation between the geology and the indoor radon values. Counties within the Northern Rocky Mountain Province (fig. 2) and that surround the Idaho Batholith (fig. 5) have a fair percentage of homes above 4 pCi/L. The rocks are dominantly Precambrian metamorphic rocks and Mesozoic intrusives (igneous rocks) in the northern part of the province. On the east side of the batholith there is a mixture of Precambrian metamorphic rocks, Cenozoic volcanic rocks, and some Paleozoic, Mesozoic, and Cenozoic sedimentary rocks and Cenozoic sediments. These rocks contain many known uranium occurrences (fig. 10) and sediments derived from these rocks are likely to have elevated uranium.

Apparently, some of the lacustrine and fluvial sediments in section 4B of province 4 (fig. 2) are producing indoor radon levels greater than 4 pCi/L (fig. 9) in 18 percent of the homes in Ada county and 5 percent of the homes in Canyon County. Cenozoic volcanic rocks (fig. 4), which are part of the Yellowstone Plateau in province 2 (fig. 2), appear to be causing a significant portion of the homes to have radon concentrations exceeding 4 pCi/L (fig. 9). Finally, the complexly faulted and folded section of the Middle Rocky Mountain Province (fig. 2) in southern Idaho and the Cenozoic volcanics and sediments in south-central Idaho (fig. 4) are also producing a number of homes with radon concentrations exceeding 4 pCi/L (fig. 9).

SUMMARY

Geologic radon potential areas (fig. 13) were delineated for ranking the radon potential in Idaho. These areas reflect the geology of the State (fig. 4) and generally combine some of the physiographic sections shown in figure 2. The 8 radon potential areas (fig. 13) were evaluated using the Radon Index (RI) and the Confidence Index (CI) discussed in the introduction to this volume. Table 2 shows the ranking of the 8 areas using these two matrices (Note: indoor Rn index number is based only on the State/EPA data).




Figure 11. Aerial radiometric map of Idaho (after Duval and others, 1989). Contour lines at 1.5 and 2.5 ppm equivalent uranium (eU). Pixels shaded from 0 to 6.0 ppm eU at 0.5 ppm eU increments; darker pixels have lower eU values; white indicates no data.




Fig. 12. Radon soil gas potential determined in BPA study (generalized from Duval and others 1989).




Fig. 13. Radon Potential Areas (See Table 2).

Table 2 shows that 5 of the 8 areas rank as having a moderate radon potential, 2 rank high, and 1 ranks low. Area 5, the Yellowstone Plateau, and area 3, the Northern Rockies, both rank as having a high radon potential at a high confidence level. Area 3 contains numerous uranium occurrences (fig. 10) and generally has elevated eU (fig. 11). Area 5 contains volcanic rocks with consistently high eU (fig. 11).

Areas 1, 4, 6, 7, and 8 (fig. 13 and Table 2) have moderate radon potentials at a high confidence levels. These areas tend to have mixed geology and elevated eU. Area 2 contains the Columbia River Basalt Group which is very low in uranium, and it has a ranking of low radon potential at a high confidence level.

This is a generalized assessment of Idaho's geologic radon potential and there is no substitute for having a home tested. The conclusions about radon potential presented in this report cannot be applied to individual homes or building sites. Indoor radon levels, both high and low, can be quite localized, and within any radon potential area there will likely be areas with higher or lower radon potential than assigned to the area as a whole. Any local decisions about radon should not be made without consulting all available local data. For additional information on radon and how to test, contact your State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

TABLE 2. Radon Index (RI) and Confidence Index (CI) scores for Radon Potential Areas, figure 13, based on the geology found within the physiographic provinces and physiographic sections shown in figure 2.

FACTOR	AREA 1		AREA 2		AREA 3		AREA 4	
	RI	CI	RI	CI	RI	CI	RI	CI
INDOOR RADON	2	3	2	3	2	3	2	3
RADIOACTIVITY	2	3	1	3	3	3	2	3
GEOLOGY	2	2	1	2	3	3	2	2
SOIL PERM.	2	2	2	2	2	2	2	2
ARCHITECTURE	2	-	2	-	2	-	2	-
GFE POINTS	0	-	0	-	0	-	0	-
TOTAL	10	10	8	10	12	11	10	10
RANKING	MOD	HIGH	LOW	HIGH	HIGH	HIGH	MOD	HIGH

FACTOR	AREA 5		AREA 6		AREA 7		AREA 8	
	RI	CI	RI	CI	RI	CI	RI	CI
INDOOR RADON	2	3	2	3	2	3	2	3
RADIOACTIVITY	3	3	3	3	3	3	3	3
GEOLOGY	3	2	2	2	2	2	2	2
SOIL PERM.	2	2	2	2	2	2	2	2
ARCHITECTURE	2	-	2	-	2	-	2	-
GFE POINTS	0	-	0	-	0	-	0	-
TOTAL	12	10	11	10	11	10	11	10
RANKING	HIGH	HIGH	MOD	HIGH	MOD	HIGH	MOD	HIGH

RADON INDEX SCORING:

<u>Radon potential category</u>	<u>Point range</u>	<u>Probable screening indoor radon average for area</u>
LOW	3-8 points	< 2 pCi/L
MODERATE/VARIABLE	9-11 points	2 - 4 pCi/L
HIGH	> 11 points	> 4 pCi/L

Possible range of points = 3 to 17

CONFIDENCE INDEX SCORING:

LOW CONFIDENCE	4 - 6 points
MODERATE CONFIDENCE	7 - 9 points
HIGH CONFIDENCE	10 - 12 points

Possible range of points = 4 to 12

REFERENCES CITED IN THIS REPORT
AND GENERAL REFERENCES RELEVANT TO RADON IN IDAHO

- AAPG, 1972, Geologic Highway Map-NORTHERN ROCKY MOUNTAIN REGION-Idaho, Montana, Wyoming: AAPG, Tulsa, Oklahoma, scale 1:1,000,000.
- Anderson, A.L., 1958, Uranium, thorium, columbium, and rare earth deposits in the Salmon Region, Lemhi County, Idaho: Idaho Bureau of Mines and Geology Pamphlet No. 115, 81 p.
- Armstrong, F.C., 1957, Dismal Swamp placer deposit, Elmore County, Idaho: U.S. Geological Survey Bulletin 1042-K, 11 p.
- Armstrong, F.C. and Weis, P.L., 1957, Uranium-bearing minerals in placer deposits of the Red River Valley, Idaho County, Idaho: U.S. Geological Survey Bulletin 1046-C, 36 p.
- Barker, R.J., McDole, R.E., and Logan, G.H., 1983, Idaho soils atlas: University Press of Idaho, 148 p.
- Bates, R.G., 1964, Natural gamma aeroradioactivity of the national reactor testing station area, Idaho: U.S. Geological Survey Map GP-446.
- Bond, J.G. and Wood, C.H., 1978, Geologic map of Idaho: Idaho Department of Lands, Bureau of Mines and Geology.
- Breckenridge, R.M., Bennett, E.H., and Harbour, J.L., 1980, Map of energy resources of Idaho: Idaho Department of Lands, Bureau of Mines and Geology Map 3.
- Caldwell, H.H. ed., 1970, Idaho economic atlas: Idaho Bureau of Mines and Geology, Moscow, Idaho, 82 p.
- Carmichael, R.S., 1989, Practical handbook of physical properties of rocks and minerals: CRC Press, Inc., 741 p.
- Cook, E.F., 1957, Radioactive minerals in Idaho: Idaho Bureau of Mines and Geology Mineral Resources Report No. 8, 5 p.
- Durrance, E.M., 1986, Radioactivity in geology, principles and applications: John Wiley & Sons, 441 p.
- Duval, J.S., 1989, Radioactivity and some of its applications in geology, in Proceedings of the Symposium on the Application of Geophysics to Engineering and Environmental Problems: Society of Engineering and Mineral Exploration Geophysicists, p. 1-61.
- Duval, J.S., Otton, J.K., and Jones, W.J., 1989, Estimation of radon potential in the Pacific Northwest using geological data: U.S. Department of Energy, Bonneville Power Administration, 146 p.
- Eisenbud, M., 1987, Environmental radioactivity from natural, industrial, and military sources: Academic Press, Inc., 475 p.
- Hunt, C.B., 1979, Surficial geology: U.S. Geological Survey National Atlas of the United States, Sheet NAC-P-0204-75M-O.

- Idaho Department of Health and Welfare, 1988, Screening survey of radon in Idaho homes: Idaho Department of Health and Welfare Division of Environmental Quality and Division of Health C4491, 10-88, 300, 42-40-100, 15 p.
- Kern, B.F., 1959, Geology of the uranium deposits near Stanley, Custer County, Idaho: Idaho Bureau of Mines and Geology Pamphlet No. 117, 40 p.
- Kiilsgaard, T.H. and Hall, W.E., 1986, Radioactive black sand placer deposits of the Challis 1°x2° Quadrangle, Idaho: U.S. Geological Survey open-File Report 86-0633, 13 p.
- Mackin, J.H. and Schmidt, D.L., 1957, Uranium and thorium-bearing minerals in placer deposits in Idaho: Idaho Bureau of Mines and Geology Mineral Resources Report No. 7, 9 p.
- Ogden, A.E., Welling, W.B., Funderburg, R.D., and Boschult, L.C., 1987, A preliminary assessment of factors affecting radon levels in Idaho, *in* Graves, B. (ed.) Radon, Radium, and Other Radioactivity in Ground Water--Hydrologic Impact and Application to Indoor Airborne Contamination: Lewis Publishers, p. 83-96.
- Ross, C.P. and Forrester, J.D., 1958, Outline of the geology of Idaho: Idaho Bureau of Mines and Geology Bulletin No. 15, 74 p.
- Savage, C.N., 1968, Lexicon of Idaho geologic names: Idaho Bureau of Mines and Geology Information Circular No. 20, 78 p.
- Sextro, R.G., Moed, B.A., Nazaroff, W.W., Revzan, K.L., and Nero, A.V., 1987, Investigations of soil as a source of indoor radon, *in* Hopke, P.K., ed., Radon and its decay products: American Chemical Society Symposium Series 331, p. 10-29.
- Sharp, W.N. and Cavender, W.S., 1962, Geology and thorium-bearing deposits of the Lemhi Pass area, Lemhi County, Idaho, and Beaverhead County, Montana: U.S. Geological Survey Bulletin 1126, 76 p.
- Tanner, A.B., 1964, Radon migration in the ground: a review, *in* Adams, J.A.S., and Lowder, W.M., eds., The natural radiation environment: Chicago, Ill., University of Chicago Press, p. 161-190.
- Trites, A.F. and Tooker, E.W., 1953, Uranium and thorium deposits in East-Central Idaho, Southwestern Montana: U.S. Geological Survey Bulletin 988-H, 52 p.
- University of Idaho, 1983, Compact atlas of Idaho: Cart-O-Graphics Laboratory, Department of Geography, College of Mines and Earth Resources, University of Idaho, Moscow, Idaho, 108 p.
- Weis, P.L., Armstrong, F.C., and Rosenblum, S., 1958, Reconnaissance for radioactive minerals in Washington, Idaho, and Western Montana 1952-1955: U.S. Geological Survey Bulletin 1074-B, 88 p.
- Witt D.A., 1984, Background gamma radiation measurements on various geologic terrains of the Rocky Mountain Region: Geological Society of America Abstracts with Programs, v. 16, No. 4, p. 260.
- Wodzicki, A. and Krasen, J., 1981, National uranium resource evaluation Dillon Quadrangle Montana and Idaho: DOE GJQ-007(81), 81 p.

Young, H.W. and Mitchell, J.C., 1973, Geothermal investigations in Idaho Part I-- Geochemistry and Geologic Setting of Selected Thermal Waters: Idaho Department of Water Administration Water Information Bulletin No. 30, 43 p.

PRELIMINARY GEOLOGIC RADON POTENTIAL ASSESSMENT OF OREGON

by
James K. Otton
U.S. Geological Survey

INTRODUCTION

Oregon is a state of varied geologic and climatic settings and varied radon potential. This assessment of the radon potential of the state relied heavily on data provided by studies of indoor radon by the Oregon Division of Health and the Bonneville Power Administration (BPA), previous work by Duval, Otton, and Jones (1989) and Otton and Duval (1990), and geologic information derived from publications of the U.S. Geological Survey and the Oregon Department of Geology and Mineral Industries. Much information in the geographic setting section is derived from the National Atlas of the United States.

This is a generalized assessment of geologic radon potential of rocks, soils, and surficial deposits of Oregon. The scale of this assessment is such that it is inappropriate for use in identifying the radon potential of small areas such as neighborhoods, individual building sites, or housing tracts. Any localized assessment of radon potential must be supplemented with additional data and information from the locality. Within any area of a given radon potential ranking, there are likely to be areas with higher or lower radon levels than characterized for the area as a whole. Indoor radon levels, both high and low, can be quite localized, and there is no substitute for testing individual homes. Elevated levels of indoor radon have been found in every State, and EPA recommends that all homes be tested. For more information on radon, the reader is urged to consult the local or State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

GEOGRAPHIC SETTING

Oregon is divisible into several physiographic provinces (fig. 1). The Coast Ranges lie along the Pacific coast and are dominated by low mountains with a relief of 1000-3000 feet. Southward the Coast Ranges give way to the Klamath Mountains where relief is generally greater than 3000 feet. Throughout these two areas less than 20 percent of the surface is gently sloping. The Willamette River Valley lies to the east of and parallels the Coast Ranges, but the valley dies out a little south of the central part of the state. The river valley is an area of plains and hills where 50-80 percent of the land's surface is gently sloping and the rest is hilly. The local relief ranges 300-500 feet. The Cascade Range lies east of the Willamette River Valley and the Klamath Mountains. The western Cascades are characterized by high mountains where greater than 80 percent of the land is steeply sloping and relief exceeds 3000 feet. The east side of the Cascades (the high Cascades) is marked by open high mountains where 20-50 percent of the land is gently sloping and the relief exceeds 3000 feet. The Deschutes-Umatilla Plateau (fig. 1) is characterized by tablelands of high relief (1000-3000 feet) where 50-80 percent of the land is gently sloping and greater than 75 percent of the gently sloped areas are in the uplands. The Blue Mountains and adjacent Joseph Upland are marked by open low (1000-3000 feet of relief) to open high (more than 3000 feet of relief) mountains where 20-50 percent of the area is gently sloping (fig. 1) and the rest is steeply sloping. The eastern boundary of the latter two areas is formed by the Snake River




Figure 1- Physiographic divisions of Oregon. (Modified from U.S. Geological Survey, 1969).

Canyon along the Oregon-Idaho border. South of the Blue Mountains are the High Lava Plains, which are characterized by plains with high hills (500-1000 feet of relief) to low mountains (1000-3000 feet of relief) (fig. 1). The Basin and Range area of south-central Oregon is characterized by plains with high hills (500-1000 feet of relief), low mountains (1000-3000 feet of relief), and two high mountain ridges (greater than 3000 feet of relief) near the southern border of the state. The Owyhee Upland is underlain by tablelands of considerable relief (500-1000 feet). Fifty to eighty percent of these latter three areas are gently sloping and the rest is steeply sloping.

Precipitation ranges from 48 to 96 inches per year throughout most of the Coast Ranges and the Cascade Mountains. The Willamette River Valley receives 32 to 48 inches of rain per year. Most of the rest of Oregon is relatively dry with the interior areas east of the Cascades receiving only 8 to 16 inches of precipitation per year except for the Blue Mountains where 16 to 48 inches of precipitation fall.

Most people in Oregon live in cities and towns extending along the Willamette River valley from Portland to Eugene (fig. 2). The remainder of the population lives in coastal towns, in a series of communities in valleys south of Eugene, and in scattered communities in north-central and northeastern Oregon. Desert areas of southeastern Oregon are sparsely populated.

Most of the Coast Ranges and the Cascade Mountains are forest and woodland areas, but some cropland and pasture areas occur adjacent to the coast. The Willamette River valley is used mostly for cropland and, to a lesser extent, for pasture, woodland, and forest. The Deschutes-Umatilla Plateau includes cropland, and grasslands used for grazing. The Blue Mountains contain areas of grassland and semiarid shrublands used for grazing at lower altitudes and forest and woodlands used for grazing at higher altitudes. The Basin and Range and Owyhee Plateau of southeastern Oregon is dominated by desert shrubland used for grazing.

GEOLOGIC SETTING

The northern Coast Ranges are underlain principally by marine sedimentary rocks and mafic volcanic rocks of Tertiary age (fig. 3). The southern part of the Coast Range is underlain by Tertiary estuarine and marine sedimentary rocks that are commonly feldspathic and micaceous. The Klamath Mountains are dominated by Triassic to Jurassic metamorphic, volcanic, and sedimentary rocks but also contain some Cretaceous intrusive rocks. The metamorphic and volcanic rocks are largely of mafic composition. Large masses of ultramafic rocks occur throughout the Klamath area.

River alluvium and river terraces occur along most of the Willamette River valley, but many of the hills that rise above the plains are underlain by Tertiary basalts and marine sedimentary rocks. The western Cascades are dominated by Tertiary andesite and basalt flows and pyroclastic rocks. The eastern Cascades are underlain by late Tertiary to Quaternary basaltic and andesitic volcanic rocks and several recently active volcanoes. Extensive areas underlain by pumice and ash occur near Crater Lake.

The Deschutes-Umatilla Plateau is underlain principally by Tertiary basalt, basaltic breccia, tuffaceous sedimentary rocks and tuff. Thin surficial deposits composed of lacustrine sediments, glaciofluvial outwash, loess, and pediment gravels form much of the northeastern part of this plateau. The Blue Mountains and Joseph Upland are comprised mainly of Tertiary basalt and andesite, tuffaceous sedimentary rocks, and tuff; however, Triassic and Jurassic sedimentary and volcanic rocks and later intrusive rocks core uplifts in this area. The sedimentary and volcanic rocks are weakly metamorphosed in many areas.




Figure 2. Population of counties in Oregon (1990 U.S. Census data).




Figure 3- Generalized geologic map for the State of Oregon. (From U.S. Geological Survey, 1969).

The High Lava Plains are underlain by basalt flows of late Tertiary and Quaternary age and Tertiary silicic ash-flow tuff. The Basin and Range Province of Oregon is an area of basaltic to andesitic volcanic rocks, silicic ash-flow tuffs, some of them welded tuffs, and sediments derived from these units. Playa basins dot the landscape. The Owyhee Plateau is underlain by Tertiary and Quaternary basalt, andesite, and rhyolite and sediments derived from these units. A few caldera complexes, some of them with associated uranium mineralization, occur within the Owyhee Plateau.

Aeroradiometric data for the state of Oregon (fig. 4) show that surface materials from the Cascade Range westward contain generally less than 1.5 ppm equivalent uranium (eU), with the exception of the Willamette River Valley, where eU values as much as 2.5 ppm are recorded locally. The northern and southern Coast Ranges show significantly different signatures, with the northern parts, underlain mostly by mafic volcanic rocks, being less than 0.5 ppm eU whereas the southern part, underlain by micaceous and feldspathic sedimentary rocks, ranging from 0.5-1.5 ppm eU. The Klamath Mountains are mixed in their eU signature with some metamorphosed sedimentary rocks and granites showing as much as 2.5 ppm, whereas the ultramafic rocks and other metamorphosed mafic volcanic rocks and sedimentary rocks show less than 1 ppm eU. East of the Cascades, much of the western part of the Basin and Range, and the Blue Mountains are very low in eU, ranging from 0.0 to 1.5 ppm. A broad area of elevated eU values (as much as 3.5 ppm in Umatilla County) occurs northwest of the Blue Mountains in the Deschutes-Umatilla Plateau in Sherman, Gilliam, Morrow and Umatilla Counties (fig. 5). Underlying bedrock in this area consists of basalts of the Columbia basin. Basalts are generally low in uranium content (<1 ppm), however, it appears that the elevated eU signatures in this area are due to surficial deposits of loess and eolian sand derived from the nearby Columbia River Valley. Sediment in the Columbia River Valley is derived in large part from more uraniferous rocks in the northeast corner of Washington and northern Idaho, transported to this part of the area by major late Pleistocene floods.

In the eastern Basin and Range and the Owyhee Plateau, a patchwork of elevated (1.5-4.5 ppm) and low eU (0.5-1.5) values occurs. Elevated eU values appear to be associated with Tertiary silicic volcanic centers and sediments derived from those rocks whereas lower eU values are associated with basaltic and andesitic rocks. Two areas of uranium deposits near the Lakeview area (southern Harney County) and the northern McDermitt caldera area (southern Malheur County) are marked by fairly intense eU signatures (as much as 5.0 ppm).

Most uranium occurrences in Oregon are found in the Basin and Range Province where they are associated with rhyolitic volcanic centers; however, they are also scattered elsewhere in the Blue Mountains, the Klamath Mountains, and the southern part of the Coast Ranges, where they are associated with various sedimentary, volcanic, and intrusive rocks.

SOILS

As part of an earlier study for the Bonneville Power Administration, Duval and others (1989) evaluated permeability data in the county soil surveys for 24 of the 36 counties in the state (fig. 6). They compiled a map of the highly permeable soil associations (those soil associations dominated by soils with percolation tests greater than 6 in/hr) in those 24 counties (fig. 7). Highly permeable soils occur mainly in the upper Willamette River valley, along parts of the Columbia River valley, in the Milton-Freewater area, and in some coastal areas.




Figure 4- Aerial radiometric map of Oregon (after Duval and others, 1989). Contour lines at 1.5 and 2.5 ppm equivalent uranium (eU). Pixels shaded from 0 to 6.0 ppm eU at 0.5 ppm eU increments; darker pixels have lower eU values; white indicates no data.




Figure 5- Map showing counties in the state of Oregon



Oregon: 1-Clatsop; 2-Columbia; 3-Tillamook; 4-Washington; 5-Multnomah; 6-Hood River; 7-Wasco; 8-Yamhill; 9-Clackamas; 10-Gilliam; 11-Morrow; 12-Umatilla; 13-Polk; 14-Marion; 15-Benton; 16-Linn; 17-Jefferson; 18-Grant; 19-Lane; 20-Crook; 21-Malheur; 22-Curry; 23-Josephine; 24-Klamath

Figure 6- Map showing counties studied by Duval and others (1989) to assess areas underlain by soil associations dominated by highly permeable soils (>6 in/hr in percolation tests).




Figure 7- Map showing areas underlain by soil associations dominated by highly permeable soils (>6 in/hr in a percolation test) within the State of Oregon as assessed by Duval and others (1989).

Soils of the west flank of the Coast Range and all of the Cascades are typically very moist in the wintertime (pore space saturation ranges from 56-96 percent for a sandy loam and 74-99 percent for a silty clay loam) and moderately moist in the summertime (pore space saturation ranges 44-56 percent for a sandy loam and 58-74 percent for a silty clay loam) (Rose and others, 1990). Within the Willamette River Valley, on the east flank of the Coast Ranges, and on the Deschutes-Umatilla Plateau, soils are typically very moist in the winter (pore space saturation ranges 56-96 percent for a sandy loam and 74-99 percent for a silty clay loam) and slightly moist in the summer (pore space saturation ranges 24-44 percent for a sandy loam and 39-58 for a silty clay loam). In the Blue Mountains and the westernmost Basin and Range the soils are very moist in the winter (pore space saturation ranges from 56-96 percent for a sandy loam and 74-99 percent for a silty clay loam) and moderately moist in the summer (pore space saturation ranges 44-56 percent for a sandy loam and 58-74 percent for a silty clay loam). In the southeastern quadrant of the state soils are typically slightly moist in the winter (pore space saturation ranges 24-44 percent for a sandy loam and 39-58 for a silty clay loam) and slightly dry in the summer (pore space saturation ranges 4-24 percent in a sandy loam and 6-39 percent for a silty clay loam).

INDOOR RADON DATA

Indoor radon data for Oregon were gathered during the Oregon Radon Project conducted by the Oregon Division of Health during 1988-1990 (Table 1, fig. 8). Data are composed of 1,954 randomly-sampled 12-month alpha-track detector measurements. These data show that counties whose major population centers lie along the Columbia River Gorge and the Willamette River Valley generally have average indoor radon levels in the 2-4 pCi/L range. Polk County averages 7.2 pCi/L. Highest county maximum readings for the State also occur in this area. Three Pacific coastal counties also range 2-4 pCi/L. Averages for three northeastern counties range 2-4 pCi/L and Union County in the northeast has a single value of 3.3 pCi/L. Lake County in the south-central part of the State averages 2.8 pCi/L.

Another source for indoor radon data in the State of Oregon is an extended study conducted by the Bonneville Power Administration (BPA). Indoor radon tests were offered as part of BPA's weatherization program, extended to homeowners across its service area (parts of the states of Washington, Oregon, Idaho, Montana, and Wyoming). Three month, wintertime alpha-track measurements were made in participants' homes starting in the fall of 1985. The resultant data were gathered on a township location system and are portrayed in figure 9. In this figure all those townships with less than 5 measurements are shown but no data are given. For those townships with at least 5 measured houses, the percentage of those homes greater than 5 pCi/L is represented in 4 categories: 0 percent, 0-5 percent, 5-10 percent, and greater than 10 percent (5 pCi/L was used as an index by BPA under the assumption that a 5 pCi/L wintertime measurement would approximately equal a 4 pCi/L year-long measurement). Most of the indoor radon data for the State of Oregon in the BPA data set come from the populous areas of western Oregon.

Examination of data tables for the townships in Oregon from this study (Bonneville Power Administration, 1990) show the following:

1. Of 151 townships in Oregon with at least 5 measurements, 17 townships averaged greater than 2 pCi/L and 2 averaged greater than 4 pCi/L.
2. Overall, 513 of 12,079 measurements (4.2 percent) for the State equalled or exceeded 4 pCi/L.




Figure 8. Indoor radon data from the Oregon Radon Project conducted by the Oregon Division of Health, 1988-90, for counties with 5 or more measurements. Data are from 12-month alpha-track tests. Histograms in map legends show the number of counties in each category. The number of samples in each county (See Table 1) may not be sufficient to statistically characterize the radon levels of the counties, but they do suggest general trends. Unequal category intervals were chosen to provide reference to decision and action levels.

TABLE 1. Indoor radon data from the Oregon Radon Project conducted by the Oregon Division of Health. Data represent randomly-sampled 12-month Alpha-track detector measurements collected during 1988-1990.

COUNTY	NO. OF MEAS.	MEAN	GEOM. MEAN	MEDIAN	STD. DEV.	MAXIMUM	%>4 pCi/L	%>20 pCi/L
BAKER	19	3.2	1.9	1.5	3.2	10.0	32	0
BENTON	108	1.2	0.9	0.9	1.1	6.0	4	0
CLACKAMAS	156	2.2	1.6	1.6	2.7	25.5	15	1
CLATSOP	29	1.4	0.8	0.5	2.4	8.7	14	0
COLUMBIA	9	2.6	2.3	2.2	1.6	6.0	11	0
COOS	38	1.9	1.2	1.1	2.6	15.6	11	0
CROOK	1	0.1	0.1	0.1	***	0.1	0	0
CURRY	1	0.8	0.8	0.8	***	0.8	0	0
DESCHUTES	42	0.8	0.5	0.5	1.2	7.2	2	0
DOUGLAS	21	0.7	0.6	0.4	0.9	3.4	0	0
GRANT	1	1.0	1.0	1.0	***	1.0	0	0
HARNEY	8	1.4	1.0	1.0	1.2	3.6	0	0
HOOD RIVER	21	1.5	1.1	0.8	1.1	3.3	0	0
JACKSON	135	1.0	0.7	0.7	1.2	6.0	4	0
JEFFERSON	2	0.7	0.6	0.7	0.4	0.9	0	0
JOSEPHINE	63	0.7	0.5	0.4	0.9	5.3	2	0
KLAMATH	31	0.6	0.5	0.4	0.7	3.1	0	0
LAKE	13	2.8	1.7	1.8	3.1	11.5	15	0
LANE	79	1.6	1.0	0.9	2.3	15.4	8	0
LINCOLN	11	1.6	1.2	0.9	2.2	7.8	9	0
LINN	57	1.3	0.9	0.9	1.4	5.9	7	0
MALHEUR	10	1.3	1.4	1.4	0.7	2.7	0	0
MARION	136	2.0	1.3	1.2	2.8	25.2	13	1
MORROW	2	1.4	1.3	1.4	0.5	1.7	0	0
MULTNOMAH	611	3.3	2.1	2.1	3.9	35.5	25	1
POLK	29	7.2	3.1	2.4	12.2	48.8	31	14
TILLAMOOK	9	2.3	1.7	2.4	1.7	5.7	11	0
UMATILLA	40	2.6	1.6	1.7	3.0	14.0	20	0
UNION	1	3.3	3.3	3.3	***	3.3	0	0
WALLOWA	8	2.2	0.8	0.3	3.4	9.9	13	0
WASCO	8	0.9	0.6	0.5	1.1	3.4	0	0
WASHINGTON	225	1.7	1.2	1.2	1.8	13.9	6	0
WHEELER	2	0.6	1.1	0.6	0.8	1.1	0	0
YAMHILL	28	3.4	2.1	1.9	4.5	20.9	18	4

BPA'S RESIDENTIAL WEATHERIZATION PROGRAM PERCENT DISTRIBUTION OF INDOOR RADON ABOVE 5 pCi/l




Figure 9- BPA's residential weatherization program, percent distribution of indoor radon above 5 pCi/L for the State of Oregon.
Source- Bonneville Power Administration, 1990.

3. Townships averaging greater than 2 pCi/L are located along the Columbia River Valley; in scattered areas in the upper Willamette River Valley; in some river valleys in the central and southern Coast Range; and in the Milton-Freewater area (northeast Umatilla County).

GEOLOGIC RADON POTENTIAL

The available data suggest that selected parts of several counties across the State are likely to have average indoor radon levels above 2 pCi/L and a few areas are likely to have average levels above 4 pCi/L. In some counties these local areas may be sufficiently widespread to cause the countywide average to exceed 2 or even 4 pCi/L.

In the central and southern Coast Range, eU values are sufficiently elevated that, in spite of high soil moisture due to high precipitation, elevated indoor radon levels are likely where soils are highly permeable or steeply sloping. This can occur in many soils developed on coarse alluvial deposits along the several river valleys that cut through the Coast Range. Soils formed on sand dunes in coastal areas, although locally highly permeable, are less likely to produce high indoor radon levels because of probable low soil-gas radon concentrations and high soil moisture.

Similarly, precipitation is also relatively high in the Willamette River Valley and the eU signature of many soils is elevated compared to other areas west of the Cascades in Oregon and Washington. Scattered areas in the upper Willamette River Valley where soils are well-drained, steep, or soil textures permit high emanation of radon are likely to produce elevated indoor radon levels. Such soils are most common in river terrace areas near the Willamette River but they also occur in some broad valley areas without significant river deposits.

Areas along the Columbia River Gorge in various counties in northwest Oregon seem susceptible to elevated indoor radon where the soils formed on river terrace deposits. Such soils are often highly permeable, well-drained, and locally steep.

Houses sited on the highly permeable Yakima gravelly loam in the town of Milton-Freewater in northeastern Umatilla County are likely to have elevated indoor radon levels. The Yakima gravelly loam and the Pilot Rock silt loam (thin silt loam over gravel) underlie much of the city of Pendleton in north-central Umatilla County and elevated indoor radon levels are likely in this city.

Areas of elevated eU associated with silicic volcanic rocks in Harney and Malheur Counties in southeast Oregon probably cause average indoor radon levels above 4 pCi/L. This area is very sparsely populated. Most people in this area live on ranches and in small communities in river and stream valleys. Few homes are located directly on soils developed on the uraniferous volcanic rocks themselves. Where alluvium is derived from silicic volcanic rocks, elevated indoor radon levels are likely.

SUMMARY

There are nine distinct geologic provinces in Oregon for which radon potential may be evaluated: the Coast Range, the Klamath Mountains, the Willamette River Valley, the Cascade Range, the Deschutes Umatilla Plateau, the Blue Mountains and Joseph Upland, the High Lava Plains, the Basin and Range Province, and the Owyhee Plateau. A relative index of radon potential (RI) and an index of the level of confidence in the available data (CI) have been established (see discussion in chapter 1 of this volume). The nine geologic provinces in Oregon are evaluated in

Table 2. The Columbia River Gorge, although not a distinctive geologic province, is considered separately (see discussion below).

The radon potential of the Coast Range is low overall due to high rainfall and high soil moisture, and low to moderate levels of uranium in the soils and rocks of the area. Locally, however, structures on steep or excessively well-drained soils especially along many of the river valleys may have elevated indoor radon levels (>4 pCi/L). For example, houses in Clatsop, Tillamook, and Coos Counties show elevated indoor radon levels. The radon potential of the Klamath Mountains is similar to that of the Coast Ranges.

The overall radon potential of the Willamette River Valley is moderate. The uranium content of the soils is moderate, the soils locally include some with high emanating power, and some of the soils are excessively well-drained (fig. 7).

Radon potential in the Cascade Range is low due to low uranium and high soil moisture. However, houses sited on steep, excessively well-drained slopes are likely to have higher indoor radon levels including many areas along the Columbia River gorge.

The Deschutes Umatilla Plateau has moderate radon potential. This area has elevated uranium in soils, includes many dry alluvial soils with high permeability, and includes many soils that are likely to have high emanating power.

The Blue Mountains and Joseph Upland have moderate radon potential. The uranium content of soils and rocks is generally low (fig. 4), but most populated areas are located on valley-floor alluvium where highly permeable soils are producing locally elevated indoor radon levels.

The High Lava Plains have low radon potential overall, but the confidence in this assessment is low due to sparse data. The western part of the High Lava Plains has generally low uranium in soils. The eastern part of the Lava Plains includes some uraniferous silicic volcanic rocks and they or alluvium associated with them may be responsible for elevated indoor radon levels locally.

The Basin and Range has moderate radon potential, but the data are sparse and the confidence level is low. Although the soil uranium content is generally low in the western part of the province, soils are fairly dry, river alluvium may be locally highly permeable, and some are likely to have high radon emanating power. In the eastern part of the province many areas are uraniferous and some volcanic rocks host uranium deposits. These volcanic rocks and sediment derived from them are likely to produce elevated indoor radon levels. Structures inadvertently sited on uranium occurrences found locally in this area may have very high indoor radon levels.

The Owyhee Plateau has high radon potential, although the confidence level is low. Much of the area is underlain by uraniferous soils and rocks and alluvium derived from them which are likely to produce elevated indoor radon levels. Uranium prospects occur in many areas and structures sited on them are likely to have very high indoor radon levels.

This is a generalized assessment of the State's geologic radon potential and there is no substitute for having a home tested. The conclusions about radon potential presented in this report cannot be applied to individual homes or building sites. Indoor radon levels, both high and low, can be quite localized, and within any radon potential area there will likely be areas with higher or lower radon potential that assigned to the area as a whole. Any local decisions about radon should not be made without consulting all available local data. For additional information on radon and how to test, contact your State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

TABLE 2. Radon Index (RI) and Confidence Index (CI) for geologic radon potential areas of Oregon. See figure 1 for locations of areas. See the introductory chapter for discussion of RI and CI.

FACTOR	Coast Range		Klamath Mountains		Willamette Valley	
	RI	CI	RI	CI	RI	CI
INDOOR RADON	1	2	1	2	2	2
RADIOACTIVITY	1	3	1	3	2	3
GEOLOGY	2	2	2	2	2	2
SOIL PERMEABILITY	2	2	2	2	2	2
ARCHITECTURE	1	-	1	-	1	-
GFE POINTS	0	-	0	-	0	-
TOTAL	7	9	7	9	9	9
RANKING	LOW MOD		LOW MOD		MOD MOD	

FACTOR	Cascade Range		Deschutes Umatilla		Blue Mountains	
	RI	CI	RI	CI	RI	CI
INDOOR RADON	1	2	2	2	2	1
RADIOACTIVITY	1	3	2	3	1	3
GEOLOGY	2	2	2	2	2	1
SOIL PERMEABILITY	2	2	2	3	3?	1
ARCHITECTURE	1	-	1	-	1	-
GFE POINTS	0	-	0	-	0	-
TOTAL	7	9	9	10	9	6
RANKING	LOW MOD		MOD HIGH		MOD MOD	

- Not used in CI.

RADON INDEX SCORING:

Radon potential category	Point range	Probable screening indoor radon average for area
LOW	3-8 points	< 2 pCi/L
MODERATE/VARIABLE	9-11 points	2 - 4 pCi/L
HIGH	> 11 points	> 4 pCi/L

Possible range of points = 3 to 17

CONFIDENCE INDEX SCORING:

LOW CONFIDENCE	4 - 6 points
MODERATE CONFIDENCE	7 - 9 points
HIGH CONFIDENCE	10 - 12 points

Possible range of points = 4 to 12

TABLE 2 (continued)--Radon Index (RI) and Confidence Index (CI) for geologic radon potential areas of Oregon. See figure 1 for locations of areas. See the introductory chapter for discussion of RI and CI.

FACTOR	High Lava Plains		Basin & Range		Owyhee Upland		Columbia River Gorge	
	RI	CI	RI	CI	RI	CI	RI	CI
INDOOR RADON	1?	1	2?	1	1?	1	3	2
RADIOACTIVITY	1	3	1	3	3	3	1	3
GEOLOGY	2	1	3	1	2	1	2	2
SOIL PERMEABILITY	2	1	2	1	2	1	3	1
ARCHITECTURE	1	-	1	-	1	-	1	-
GFE POINTS	0	-	0	-	0	-	0	-
TOTAL	7	6	9	6	9	6	10	8
RANKING	LOW LOW		MOD LOW		MOD LOW		MOD MOD	

- Not used in CI.

RADON INDEX SCORING:

Radon potential category	Point range	Probable screening indoor radon average for area
LOW	3-8 points	< 2 pCi/L
MODERATE/VARIABLE	9-11 points	2 - 4 pCi/L
HIGH	> 11 points	> 4 pCi/L

Possible range of points = 3 to 17

CONFIDENCE INDEX SCORING:

LOW CONFIDENCE	4 - 6 points
MODERATE CONFIDENCE	7 - 9 points
HIGH CONFIDENCE	10 - 12 points

Possible range of points = 4 to 12

REFERENCES CITED IN THIS REPORT
AND GENERAL REFERENCES RELAVENT TO RADON IN OREGON

- Berry, M. R., 1981, Geology of the Lakeview uranium district, Oregon *in* Goodell, P. C. and Waters, A. C., eds., Uranium in volcanic and volcanoclastic rocks: AAPG Studies in Geology 13, p. 55-62.
- Bonneville Power Administration, 1990, Radon monitoring results from BPA's residential conservation programs: Report No. 12: Portland, Oregon, Bonneville Power Administration, unpaginated.
- Bradley, R., 1982, Mining districts and mineral deposits of the Basin and Range Province of Oregon: United States Geological Survey Open-File Report 82-0058, 15 p.
- Cathrall, J. B. and Tuckek, E. T., Charles Sheldon Antelope Range and Sheldon National Antelope Refuge, Nevada and Oregon: *in* Marsh, S. P., Kropscot, S. J. and Dickinson, R. G., eds., Wilderness mineral potential; assessment of mineral-resource potential in U.S. Forest Service lands studied 1964-1984: U.S. Geological Survey Professional Paper 1300, p. 765-767.
- Dayvault, R. D., Castor, S. B. and Berry, M. R., 1985, Uranium associated with volcanic rocks of the McDermitt Caldera, Nevada and Oregon, *in* Anonymous, Uranium deposits in volcanic rocks; proceedings of a technical committee meeting, El Paso, TX, Apr. 2-5, 1984: Panel Proceedings Series-International Atomic Energy Agency STI/PUB/690, p. 379-409.
- Duval, J. S., Otton, J. K., and Jones, W. J., 1989, Estimation of radon potential in the Pacific Northwest using geological data: Portland, Oregon, Bonneville Power Administration, 146 p.
- Duval, J. S., Jones, W. J., Riggle, F. R., and Pitkin, J. A., 1989, Equivalent uranium map of the conterminous United States: U.S. Geological Survey Open-File Report 89-478, 10 p.
- Erdman, J. A. and Harrach, G. H., 1981, Uranium in big sagebrush from Western U.S. and evidence of possible mineralisation in the Owyhee Mountains of Idaho: Journal of Geochemical Exploration, v. 14, n. 1, p. 83-94.
- Erikson, E. H., 1977, Preliminary study of the uranium favorability of Malheur County, Oregon: U. S. Department of Energy Report No.: GJBX-91(77), 16 p.
- Moed, B.A., Nazaroff, A.V., Nero, M.B., Schwehr, M.B., and Van Heuvelen, A., 1984, Identifying areas with potential for high indoor radon levels: analysis of the National Airborne Radiometric Reconnaissance data for California and the Pacific Northwest: Berkeley, California, Lawrence Berkeley Laboratory Report No. LBL-16955, 70 p.

- Nero, A.V., Schwehr, M.B., Nazaroff, W.W., and Revzan, K.L., 1986, Distribution of airborne radon-222 concentrations in U.S. homes: *Science*, v. 234, p.992-997.
- Otton, J. K., 1987, Indoor radon; geologic controls in Pacific Northwest: *Geological Society of America Abstracts with Programs*, v. 19, n. 2, p. 122.
- Otton, J. K. and Duval, J. S., 1990, Geologic controls on indoor radon in the Pacific Northwest, *in* U.S. Environmental Protection Agency, The 1990 international symposium on radon and radon reduction technology: Volume III. Preprints, unpaginated.
- Peck, D. L., 1961, Geologic map of Oregon west of the 121st meridian: U.S. Geological Survey Miscellaneous Investigations Map I-325. Scale 1:500,000.
- Peterman, Z. E., Coleman, R. G. and Bunker, C. M., 1981, Provenance of Eocene graywackes of the Flournoy Formation near Agness, Oregon; a geochemical approach: *Geology*, v. 9, n. 2, p. 81-86.
- Reimer, G. M., 1981, Helium soil-gas survey of a portion of the McDermitt Caldera complex, Malheur County, Oregon: United States Geological Survey Open-File Report 81-0565, 10 p.
- Ronkos, C. J., 1981, Geology, alteration, and mineralization in the pyroclastic and sedimentary deposits of the Bretz-Aurora Basin, McDermitt Caldera, Nevada-Oregon: Master' thesis, Univ. of Nevada, Reno, unknown p.
- Roper, M. W. and Wallace, A. B., 1981, Geology of the Aurora uranium prospect, Malheur County, Oregon *in* Goodell, P. C. and Waters, A. C., eds., Uranium in volcanic and volcanoclastic rocks: AAPG Studies in Geology 13, p.81-88
- Rose, A. W., Ciolkosz, E. J., and Washington, J. W., 1990, Effects of regional and seasonal variations in soil moisture and temperature on soil gas radon, *in* U.S. Environmental Protection Agency, The 1990 international symposium on radon and radon reduction technology: Volume III. Preprints, unpaginated.
- Rytuba, J. J., 1977, Uranium content of tuffaceous sediments and opalite mercury deposits within the McDermitt Caldera, Oregon-Nevada: *Geological Society of America Abstracts with Programs*, v. 9, n. 4, p. 492.
- Rytuba, J. J., 1979, Alteration aureole in McDermitt caldera in Nevada and Oregon: U. S. Geological Survey Professional Paper 1150, 79 p.
- Rytuba, J. J. and Conrad, W. K., 1981, Petrochemical characteristics of volcanic rocks associated with uranium deposits in the McDermitt caldera complex *in* Goodell, P. C. and Waters, A. C., eds., Uranium in volcanic and volcanoclastic rocks: AAPG Studies in Geology, v. 13, p. 63-72

- Rytuba, J. J. and Glanzman, R. K., 1979, Relation of mercury, uranium, and lithium deposits to the McDermitt Caldera complex, Nevada-Oregon *in* Ridge, J. D., ed., Papers on mineral deposits of western North America; proceedings of the Fifth quadrennial symposium of the International Association on the Genesis of Ore Deposits; Volume 2: Nevada Bureau of Mines Report 33, p. 109-117.
- Rytuba, J. J., Glanzman, R. K. and Conrad, W. K., 1979, Uranium, thorium, and mercury distribution through the evolution of the McDermitt Caldera complex *in* Newman, G. W. and Goode, H. D., eds., Basin and Range symposium and Great Basin field conference, Oct. 7-11, 1979: Rocky Mountain Assoc. Geol., p. 405-412.
- Rytuba, J. J., Minor, S. A. and McKee, E. H., 1981, Geology of the Whitehorse Caldera and caldera-fill deposits, Malheur County, Oregon: United States Geological Survey Open-File Report 81-1092, 22 p.
- U.S. Geological Survey, 1969, Mineral and Water Resources of Oregon: Report of the Committee on Interior and Insular Affairs, Washington, U.S. Government Printing Office, 462 p.
- U. S. Geological Survey and U. S. Bureau of Mines, 1984, Mineral resources of the Charles Sheldon Wilderness Study Area, Humboldt and Washoe counties, Nevada, and Lake and Harney counties, Oregon: U.S. Geological Survey Bulletin 1538, 139 p.
- Walker, G. W., 1977, Geologic map of Oregon east of the 121st meridian: U.S. Geological Survey Miscellaneous Investigations Map I-902, Scale 1:500,000.
- Walker, G. W., 1980, Preliminary report on the Lakeview uranium area, Lake County, Oregon: U. S. Geological Survey Open-File Report 80-532, 59 p.
- Walker, G. W., 1980, Uranium in the Lakeview area, Oregon: U.S. Geological Survey Professional Paper 1175, 43 p.
- Walker, G. W., 1985, Geology of the Lakeview uranium area, Lake County, Oregon, *in* Anonymous, Uranium deposits in volcanic rocks; proceedings of a technical committee meeting, El Paso, TX, Apr. 2-5, 1984: Panel Proceedings Series-International Atomic Energy Agency STI/PUB/690, p.411-447 p.
- Wallace, A. B. and Roper, M. W., 1981, Geology and uranium deposits along the northeastern margin, McDermitt caldera complex, Oregon *in* Goodell, P. C. and Waters, A. C., eds., Uranium in volcanic and volcanoclastic rocks: AAPG Studies in Geology 13, p. 73-79.
- Weissenberger, K. W., 1984, Lakeview uranium area, Lake County, Oregon; constraints on genetic modelling from a district-scale perspective: Ph.D. thesis, Stanford Univ., Stanford, CA, 367 p.

- Wilkening, R. M. and Cummings, M. L., 1987, Mercury and uranium mineralization in the Clarno and John Day formations, Bear Creek Butte area, Crook County, Oregon: Oregon Geology, v. 49, n. 9, p. 103-110.
- Wilson, R. D., Purdom, William, and Welton, Richard, 1989, Radon study of the Rogue Valley region of southern Oregon: Ashland, Oregon, Southern Oregon State College, unknown p.

PRELIMINARY GEOLOGIC RADON POTENTIAL ASSESSMENT OF WASHINGTON

by
James K. Otton
U.S. Geological Survey

INTRODUCTION

Washington is a State of varied geologic and climatic settings and varied radon potential. This assessment of the radon potential of the State is based largely on data provided by a study of indoor radon by the State/EPA Indoor Radon Survey of Washington and the Bonneville Power Administration (BPA), previous work by Duval and others (1989) and Otton and Duval (1990), and geologic information derived from publications of the Washington Department of Natural Resources, Division of Geology and Earth Resources. Much of the information in the geographic setting section is derived from the National Atlas of the United States of America.

This is a generalized assessment of geologic radon potential of rocks, soils, and surficial deposits of Washington. The scale of this assessment is such that it is inappropriate for use in identifying the radon potential of small areas such as neighborhoods, individual building sites, or housing tracts. Any localized assessment of radon potential must be supplemented with additional data and information from the locality. Within any area of a given radon potential ranking, there are likely to be areas with higher or lower radon levels than characterized for the area as a whole. Indoor radon levels, both high and low, can be quite localized, and there is no substitute for testing individual homes. Elevated levels of indoor radon have been found in every State, and EPA recommends that all homes be tested. For more information on radon, the reader is urged to consult the local or State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

GEOGRAPHIC SETTING

Washington is divided into several distinctive physiographic provinces (fig. 1). The Okanogan Highlands area in northeastern and north-central Washington is part of the northern Rocky Mountains. This highland area is dissected by a few north-trending river valleys including the Columbia River valley. This highland is characterized mostly by low mountains where less than 20 percent of the area is gently sloping and the local relief is 1000-3000 feet. High mountains with greater relief occur in the northeastern corner of this highland area and low open mountains occur in the highland area close to Spokane.

The Columbia Basin is in the southeastern part of the State. This terrain is bounded to the north by the Columbia and Spokane Rivers and to the west by the foothills of the Cascade Range. It is characterized by irregular plains of low relief (100-300 feet), tablelands of moderate to high relief (300-3000 feet), and open hills and open low mountains of moderate to high relief (300-3000 feet). The percentage of area underlain by gentle slopes ranges from 50 to 80 percent in the plains and 20-50 percent in the open hills and open low mountains.

The Blue Mountains area lies in the southeasternmost corner of the State. This area is characterized by open low mountains with 1000-3000 feet of relief, and 20-50 percent of the area is gently sloping.




Figure 1- Physiographic divisions of Washington. (Modified from Lasmanis, 1991).

The Cascade Mountains separate these eastern areas from the western part of the State. The Cascades are high mountains and open high mountains where relief exceeds 3000 feet. The Columbia River has cut a gorge through these mountains along the border between Washington and Oregon.

Immediately west of the Cascades are the Puget Lowlands, which extend from the Canadian border southward nearly to the Columbia River. This area is marked by tablelands of moderate relief (300-500 feet) in the area around Seattle; by plains with low mountains (local relief of 1000-3000 feet) in the areas near the Canadian border; and by open hills (local relief 300-500 feet) from southern Tacoma southward to the Columbia River. Fifty to eighty percent of the land is gently sloping in the first two areas whereas 20-50 percent of the latter area is gently sloping.

The Portland Basin, a small physiographic area near Vancouver, is a northward extension of the Willamette River Valley. This area resembles the southern part of the Puget Lowlands physiographically.

Westernmost Washington is dominated by the Coastal Mountains and Coastal Hills. The Coastal Mountains are high mountains (relief greater than 3000 feet) in the eastern half and lower mountains (relief 100-3000 feet) in the western, coastal part. An area of open hills separates the Coastal Mountains from the Coastal Hills, which are low mountains of 1000-3000 feet local relief.

The western Coastal Mountains, the Coastal Hills, and the Cascades are areas of heavy rainfall, with amounts ranging from 40 to 200 inches per year. Only the northeasternmost part of the Coastal Mountains are relatively dry. There, precipitation ranges from 20 to 30 inches per year. Rainfall in the Puget Lowlands and the Portland Basin generally ranges from 30 to 50 inches per year. In Seattle, the wettest months are the winter months and summer is somewhat drier, but even in summer 2 to 3 inches of rain may fall per month.

East of the Cascade Mountains, rainfall is typically in the range of 9 to 15 inches per year, with the exception of the Okanogan Highlands, which receive as much as 30 to 40 inches per year. In Spokane, rainfall is fairly evenly distributed through the year, except for July and August, which are the driest months.

Most of the population of the State is located in the Puget Lowlands and the Portland area at the west end of the Columbia River Gorge (fig. 2). A lengthy urban corridor in the Puget Lowlands extends from Bellingham near the Canadian border to the Tacoma/Olympia area at the south end of Puget Sound. Other scattered population centers occur east of the Cascades, including Spokane and adjacent areas, the lower Yakima River Valley, and Walla Walla. Most of the populated areas east of the Cascades lie along the river valleys.

Mountainous areas of the State are used principally for timber or timber and grazing. The Puget Lowlands outside of the urban areas is used for cropland, with lesser grazing and timber. The Columbia Basin is dominated by dryland farming and grazing. Some irrigated cropland is found along the Columbia River valley in the western part of the Columbia Basin.

GEOLOGIC SETTING

The coastal mountains and hills of Washington are underlain principally by Cretaceous and Tertiary continental and marine sedimentary rocks and pre-Miocene volcanic rocks (fig. 3). The Puget Lowlands is underlain almost entirely by glacial deposits and Holocene alluvium. Most of the glacial and alluvial material of the Puget Lowlands is derived from the Cascades to the east and the mountains of the Olympic Peninsula to the west. The Portland Basin is underlain primarily by alluvial material. The Cascade Mountains can be divided into two geologic terranes: a northerly





Figure 2. Population of counties in Washington (1990 U.S. Census data).




Figure 3- Generalized geologic map for the State of Washington. (From Schuster, 1990).

GEOLOGIC MAP OF WASHINGTON

compiled by
J. ERIC SCHUSTER
1990

GEOLOGIC UNITS

Unconsolidated Deposits

- Qs** Quaternary sediments, dominantly nonglacial:
includes alluvium and volcanoclastic, glacial
outburst flood, eolian, landslide, and coastal
deposits
- Qg** Quaternary sediments, dominantly glacial
drift; includes alluvium

Neogene Sedimentary Rocks

- uTs** Upper Tertiary (Pliocene-Miocene)
lTs Lower Tertiary (Oligocene-Paleocene)
Mzs Mesozoic
MzPzs Mesozoic-Paleozoic
Pzs Paleozoic
pCs Precambrian

Volcanic Rocks

- Qv** Quaternary
QPy Quaternary-Pliocene
uTv Upper Tertiary (Pliocene-Miocene)
uTv2 Columbia River Basalt Group
lTv Lower Tertiary (Oligocene-Paleocene)
Mzv Mesozoic

Intrusive Igneous Rocks

- Ti** Tertiary
Mzi Mesozoic
Pzi Paleozoic

Metamorphic Rocks

- pTm** Pre-Tertiary
pKm Pre-Cretaceous
Pzm Paleozoic
pCm Precambrian
u Ultramafic rocks

Note: Some pre-Tertiary sedimentary and volcanic rock units
include low-grade metamorphic rocks.
Ages assigned to metamorphic rocks are protolith ages.

EXPLANATION

- Contact**
- Fault**
- Thrust fault, sawteeth on upper plate**
- Detachment fault, blocks on upper plate**
- Strike-slip fault, arrows show relative movement**
- Anticline, dotted where concealed**

terrane composed principally of Mesozoic metamorphic rocks intruded by Mesozoic and Tertiary granitic rocks, and a southerly terrane composed of Tertiary and Holocene volcanic rocks. The Holocene volcanic centers are responsible for locally thick volcanic ash deposits east of the Cascade Mountains. The Columbia Basin is underlain principally by Miocene basaltic volcanic rocks overlain by extensive veneer of glaciofluvial outwash and eolian and lacustrine deposits. The mountainous terrain of the northeast and north-central part of the State is underlain by Precambrian sedimentary rocks, Paleozoic sedimentary rocks, and Mesozoic metamorphic rocks, all intruded by Mesozoic and Tertiary granites.

During the late Pleistocene, most of the Cascades, the Olympic Peninsula, and the Rocky Mountains north and northwest of Spokane were glaciated. Extensive glacial deposits now cover most of the Puget Lowlands. As ice receded from northeastern Washington, large lakes formed behind ice dams along the ancestral Columbia River valley. When these dams broke, massive floods poured down many of the river valleys across the Columbia Basin. These floods carried substantial quantities of alluvium which was deposited along the entire river drainages. In some places, valleys were scoured clean of surficial sediments down to the underlying basaltic bedrock. Subsequent westerly winds blew much of the finer-grained river alluvium from river bottoms and deposited this sand and silt across many of the adjacent valley floors.

Aerial radiometric data for the State of Washington (fig. 4) show that surficial materials within the Puget Lowlands and the coastal areas to the west contain entirely less than 1.5 ppm equivalent uranium (eU). Within the Cascades, the rocks have mostly less than 1.5 ppm eU; however, some of the Holocene volcanic centers in the southern Cascades include volcanic rocks containing as much as 2.5 ppm eU. In addition, some of the Tertiary granitic plutons in the central and northern Cascades are significantly enriched in uranium, as much as 5.5 ppm eU. In the westernmost part of the Okanogan Highlands, Mesozoic and early Tertiary granites have unusually low eU signatures (as low as 0.5 ppm eU), whereas to the east, many of the granites and some of the metamorphic rocks have significantly elevated eU signatures (as much as 5 ppm eU).

Although much of the Columbia Basin is underlain by basaltic rocks that are generally low in uranium content, most of this area has eU signatures of 2.0 to 3.0 ppm because of the extensive surficial cover of Quaternary and Holocene glaciofluvial, lacustrine, and eolian deposits. This surficial cover has been derived in large part from more uraniferous rocks in northeastern Washington and northern Idaho. The larger areas of 0.5 to 1.5 ppm eU within the Columbia Basin represent areas on the east flank of the southern Cascades and the Blue Mountains where this surficial cover was not deposited or areas within the Columbia basin where cover was stripped by late Pleistocene floods.

Uranium occurrences and deposits in Washington, a potential source of elevated radon levels, are largely associated with granitic and metamorphic rocks in the Okanogan Highland area north and northwest of Spokane and with Tertiary sedimentary rocks derived from these crystalline rocks, although some uranium occurrences are located in the central Cascade Mountains. Granitic rocks in the hills immediately north of Spokane host numerous uranium occurrences.

Organic-rich Holocene and late Pleistocene alluvium along stream valleys in many of the granitic terranes in the Okanogan Highland area is host to surficial uranium deposits (Johnson and others, 1987). Because of their relative youth, these Holocene uranium deposits are typically low in radioactivity (and radioactive decay products such as radon) in spite of average grades that range from 100 to 800 ppm uranium (Zielinski and others, 1986). Fortunately, most of these deposits are on the floodplains of streams or are in wetlands, and thus are not suitable for construction.




Figure 4- Aerial radiometric map of Washington (after Duval and others, 1989). Contour lines at 1.5 and 2.5 ppm equivalent uranium (eU). Pixels shaded from 0 to 6.0 ppm eU at 0.5 ppm eU increments; darker pixels have lower eU values; white indicates no data.

SOILS

As part of an earlier study for the BPA, Duval and others (1989) evaluated permeability data in the county soil surveys for 30 of the 37 counties in the State (fig. 5). They compiled a map of the highly permeable soil associations (those soil associations dominated by soils with permeabilities greater than 6 in/hr in percolation tests) in those 30 counties (fig. 6).

The low coastal areas and coastal mountains and hills of western Washington and the Cascades of west-central Washington are characterized by very moist soils in the winter (56-96 percent pore saturation in sandy loams, and 74-99 percent saturation in a silty clay loam) and moderately moist soils in the summer (44-56 percent saturation in sandy loams, and 58-74 percent in a silty clay loam) (Rose and others, 1990). The Puget Lowlands and the northern and western edges of the Columbia Basin and the Blue Mountains are characterized by very moist soils in the winter (56-96 percent pore saturation in sandy loams, and 74-99 percent pore saturation in a silty clay loam) and slightly moist soils in the summer (24-44 percent pore saturation in sandy loams, and 39-58 percent pore saturation in silty clay loams). Soils in the rest of the Columbia Basin are slightly moist in the winter and slightly dry in the summer (4-24 percent pore saturation in sandy loams and 6-39 percent pore saturation in silty clay loams). Soils in the Okanogan Highlands are characterized by very moist soils in the winter and moderately moist soils in the summer.

INDOOR RADON DATA

The EPA and the State of Washington completed a population-weighted survey of indoor radon levels in Washington during the winter of 1990-1991 (fig. 7; table 1). Sampled houses were randomly selected from existing housing stock, which means that homes sampled tend to cluster in the more populated areas. Interpretations of population-weighted data must be made with caution, because the measured houses are typically only from a relatively few population centers within a given county and do not provide geographic coverage of the county's surface area.

Highest indoor radon readings occur in the northeastern part of the State, with three counties in that area averaging greater than 4 pCi/L. Much of the east and north-central part of the State has indoor radon averages between 2 and 4 pCi/L. All counties west of the Cascade Range have very low indoor radon averages, generally less than 1 pCi/L. A few readings exceed 4 pCi/L in three counties in the southern part of the Puget Lowland.

Another source for indoor radon data in the State of Washington is a long-term study conducted by the BPA. Indoor radon tests were offered as part of BPA's weatherization program, extended to homeowners across its service area (parts of the states of Washington, Oregon, Idaho, Montana, and Wyoming). Three month, wintertime alpha-track measurements were made in participants' homes starting in the fall of 1985. The resulting data were plotted on a township location system and are presented in figure 8. In this figure all those townships with less than 5 measurements are shown but no data are given. For those townships with at least 5 measured houses, the percentage of those homes greater than 5 pCi/L is represented (this number was used as an index by BPA under the assumption that a 5 pCi/L wintertime measurement would approximately equal a 4 pCi/L year-long measurement). The coverage of the State is less complete than for the State/EPA data set, however because a smaller geographic area was sampled, more detailed correlations with geology and soils is possible.




Figure 5- Map showing counties studied by Duval and others (1989) to assess areas underlain by soil associations dominated by highly permeable soils (>6 in/hr in percolation tests).



Figure 6- Map showing areas underlain by soil associations dominated by highly permeable soils (>6 in/hr in a percolation test) within the State of Washington as assessed by Duval and others (1989).





Figure 7. Screening indoor radon data from the EPA/State Residential Radon Survey of Washington, 1991-92, for counties with 5 or more measurements. Data are from 2-7 day charcoal canister tests. Histograms in map legends show the number of counties in each category. The number of samples in each county (See Table 1) may not be sufficient to statistically characterize the radon levels of the counties, but they do suggest general trends. Unequal category intervals were chosen to provide reference to decision and action levels.

TABLE 1. Screening indoor radon data from the EPA/State Residential Radon Survey of Washington conducted during 1991-92. Data represent 2-7 day charcoal canister measurements from the lowest level of each home tested.

COUNTY	NO. OF MEAS.	MEAN	GEOM. MEAN	MEDIAN	STD. DEV.	MAXIMUM	%>4 pCi/L	%>20 pCi/L
ADAMS	11	3.5	2.3	2.0	4.4	15.9	18	0
ASOTIN	18	2.9	1.5	1.3	6.3	27.8	6	6
BENTON	106	1.8	1.0	1.1	3.5	28.1	7	2
CHELAN	9	2.2	1.4	1.8	2.2	7.4	11	0
CLALLAM	22	0.3	0.1	0.2	0.4	1.6	0	0
CLARK	69	2.4	1.1	1.3	3.5	23.0	12	1
COLUMBIA	5	3.4	3.0	4.0	1.7	5.4	40	0
COWLITZ	35	0.5	0.2	0.2	1.4	8.2	3	0
DOUGLAS	17	1.6	1.0	1.2	1.3	4.0	0	0
FERRY	28	3.2	1.5	1.3	3.9	10.8	25	0
FRANKLIN	26	1.6	1.1	1.2	1.4	5.1	12	0
GARFIELD	5	0.8	0.5	0.6	0.7	1.9	0	0
GRANT	54	1.9	0.9	1.0	2.9	17.7	9	0
GRAYS HARBOR	29	0.7	0.2	0.2	1.4	6.8	3	0
ISLAND	10	0.1	0.1	0.2	0.2	0.4	0	0
JEFFERSON	11	0.3	0.2	0.2	0.4	1.1	0	0
KING	215	0.4	0.2	0.3	0.8	6.2	2	0
KITSAP	34	0.4	0.3	0.3	0.5	1.8	0	0
KITTITAS	4	1.2	0.9	1.4	0.7	1.9	0	0
KLICKITAT	22	1.7	1.1	0.8	2.3	10.1	14	0
LEWIS	24	0.7	0.4	0.5	0.7	2.8	0	0
LINCOLN	15	2.2	1.7	1.8	1.6	5.7	7	0
MASON	18	0.6	0.3	0.3	0.7	2.5	0	0
OKANOGAN	43	2.8	1.5	1.3	3.7	18.7	21	0
PACIFIC	11	0.2	0.1	0.1	0.2	0.6	0	0
PEND OREILLE	55	7.2	2.2	2.0	19.1	136.9	31	5
PIERCE	132	0.8	0.3	0.4	1.7	13.6	5	0
SKAGIT	9	0.4	0.2	0.0	0.6	1.3	0	0
SKAMANIA	35	8.3	1.9	1.4	17.4	79.5	34	11
SNOHOMISH	63	0.2	0.1	0.1	0.3	1.4	0	0
SPOKANE	449	9.9	5.1	4.9	14.2	152.2	59	13
STEVENS	47	5.1	2.1	1.7	11.3	73.7	32	4
THURSTON	45	0.8	0.3	0.3	1.8	9.1	4	0
WAHKIAKUM	21	1.0	0.4	0.4	1.8	6.7	10	0
WALLA WALLA	56	2.6	1.7	1.9	2.3	9.4	23	0
WHATCOM	17	0.4	0.3	0.3	0.3	1.3	0	0
WHITMAN	31	5.9	2.2	2.3	14.0	79.2	39	3
YAKIMA	134	1.8	1.1	1.1	2.5	17.8	6	0

BPA'S RESIDENTIAL WEATHERIZATION PROGRAM PERCENT DISTRIBUTION OF INDOOR RADON ABOVE 5 pCi/l

WASHINGTON



LEGEND

- DATA AVAILABLE - LESS THAN 5 SITES PER TOWNSHIP
- 0.00%
- 0.01 - 4.99%
- 5.00 - 9.99%
- 10.00% Or More

Note: Radon concentrations depicted are for townships monitored throughout the Pacific Northwest. Townships without monitoring data are not shown.

Total number of sites = 3513 - January, 1990
For Washington, Oregon, Idaho, Montana, and Wyoming.

Source data: Bonneville Power Administration.

BPA GEOGRAPHIC INFORMATION SYSTEM
BONNEVILLE POWER ADMINISTRATION

Figure 8- BPA's residential weatherization program, percent distribution of indoor radon above 5 pCi/L for the Pacific Northwest. (Source- Bonneville Power Administration).

Within the Puget Lowlands, townships extending from the northwestern corner of Snohomish County southward to northernmost Pierce County show that either 0 percent or 1-5 percent of measured homes are over 5 pCi/L (see fig. 9 for a map of the counties in the State). This highly populated area has the lowest average indoor radon levels in the entire study. In one township in Seattle, the highest reading in more than 1000 measurements was about 2.8 pCi/L.

Townships in which 10 percent or more of homes exceed 5 pCi/L (fig. 8) include the southern suburbs of Tacoma, where they appear to be associated with the highly permeable Spanaway soils; the Sequim area in western Clallam County, where the soils are relatively dry and locally highly permeable; the Columbia River gorge in southern Skamania County, where they are associated with highly permeable, steeply sloped soils; west-central Benton County where highly permeable river alluvium and dune sands occur; the Wenatchee area in southern Chelan County, where highly permeable river terrace deposits occur; the city of Spokane and its eastern suburbs, which are built on highly permeable glacial outwash; and the Curlew area in northern Ferry County, where soils have formed on highly permeable glacial outwash, river terraces, and fans. The highly permeable soils that cause elevated indoor radon levels in the eastern Spokane suburbs are widespread throughout the Spokane metropolitan area. In some townships near Spokane as many as 70 percent of the homes are over 4 pCi/L and indoor radon levels are locally as high as 200 pCi/L.

Townships in which 5 to 10 percent of the measured homes are over 5 pCi/L occur in west-central Lewis County where they seem to be associated with highly permeable soils on glacial outwash plains and high terraces, and in the Ellensburg area in Kittitas County where they likely are associated with river alluvium and glacial outwash.

This analysis suggests that the northeastern corner of the State has the highest overall radon potential, and specific areas elsewhere in the State, generally small parts of counties where unusual soils are found, may also have significant geologic radon potential.

GEOLOGIC RADON POTENTIAL

The State of Washington can be subdivided into three areas of varying overall geologic radon potential. Areas within and west of the Cascade Mountains are generally characterized by very low uranium in soils (less than 1.5 ppm eU) and high soil moisture due to the high annual rainfall. Under these conditions only modest levels of radon are generated by the soil and radon migrates slowly through the soil. Only in local areas where the soils are extremely permeable ($>> 20$ in/hr) or where the topography is steeply sloped are indoor radon levels likely to be significant. Such areas might include the easternmost and southernmost parts of the Puget Lowlands, where the combination of steep slopes and extremely permeable glacial outwash occur locally, or areas in the Portland Basin near Vancouver or along the Columbia River gorge where highly permeable river or lacustrine delta deposits, volcanic debris flows, and steep slopes locally occur.

In the Columbia Basin area east of the Cascade Mountains, most soils are drier (rainfall is typically in the range of 9 to 15 inches per year), the uranium content of soils is moderately elevated (most areas range from 1.5 to 2.5 ppm eU), and many river valleys contain soils that are rapidly (6-20 in/hr) or very rapidly (>20 in/hr) permeable because they have developed on sand, river alluvium, or glacial outwash (fig. 6). Lacustrine deposits are present in many areas and soils



Figure 9. County boundaries and names in Washington.

developed on these units often have high porosity and emanating power, and show development of secondary permeability by soil cracking. Many of these areas have significant potential for producing elevated indoor radon levels.

In the Okanogan Highland, soils in the valley areas have formed on highly permeable, coarse glacial outwash, fluvial terraces, and similar materials. Although these soils are generally wetter than soils in the Columbia Basin to the south, the very high permeability (>20 in/hr) of the majority of the soils permits rapid drainage of soil water and radon transport by diffusion and convective flow is significant. These areas have significant geologic potential for elevated indoor radon levels.

Of special interest are buildings that may be inadvertently sited on uranium concentrations in the hills north of Spokane. Very high indoor radon levels are likely in these buildings. Such uranium occurrences are likely in the granitic rocks that underlie much of the area. Many of these uranium occurrences are described in the National Uranium Resource Evaluation literature for the area.

SUMMARY

There are six distinct geologic provinces in Washington for which radon potential may be evaluated (Table 2): the Coastal mountains and hills, the Puget Lowlands, the Cascade Mountains, the Okanogan Highlands, the Columbia Basin, and the Blue Mountains. A separate area of moderate radon potential, the Columbia River Gorge, is also evaluated in Table 2. A relative index of radon potential (RI) and an index of the level of confidence in the available data (CI) have been established (see discussion in chapter 1 of this volume).

Low uranium in soils and very high rainfall throughout most of the area contribute to low radon potential overall in the coastal mountains and hills of western Washington. The northeastern corner of the Olympic Peninsula has lower rainfall and lower soil moisture than the rest of the Coastal Range Province. Highly permeable, excessively well drained soils may cause locally elevated indoor radon levels here.

The Puget lowland overall has very low radon potential because of high rainfall and consequently high soil moisture, and low uranium content of soils. Most townships from Tacoma northward average less than 1 pCi/L radon in the BPA study. Structures built on local very steep or excessively well-drained soils, especially on the east side of the lowland area, may be among the few likely to have elevated indoor radon levels. Only in the southern part of the Puget lowland south of Tacoma where excessively drained soils and somewhat elevated uranium in soils occurs is the potential moderate.

Overall, the sparsely populated Cascade Mountain Province has low radon potential because of the low uranium content and high soil moisture of the soils. A major exception to this is the Columbia River Gorge area where highly permeable, excessively well drained soils underlie many of the communities and the radon potential is moderate. Moreover, much of the alluvium in the Gorge is derived from the upper Columbia River valley where the uranium content of the geologic materials is higher than within the Cascade Mountain Province itself. In the State/EPA survey 85 percent of the basements in homes in Skamania County exceeded 4 pCi/L. Indoor radon levels in townships along the Columbia River average 2 to 5 pCi/L in the BPA study.

In spite of the fact that bedrock across the Columbia Basin is often composed of low-uranium basaltic rocks, surficial deposits include extensive Pleistocene glaciofluvial outwash, eolian, and lacustrine deposits, which locally have highly permeable soils, soils with high emanating coefficients, and higher soil uranium levels. This area has moderate to locally high radon potential.

The Okanogan Highlands has high radon potential. This assessment is supported by several studies (Otton and Duval, 1990; Moed and others, 1984; R. Sextro, written commun., 1985; Otton and Reimer, unpub. data, 1986). Excessively well drained glaciofluvial outwash underlies most of the valleys throughout the area. The granitic material in much of the outwash is highly uraniferous. Areas where uranium occurrences are found, such as the granitic and metamorphic terranes in the mountains north and northwest of Spokane, may have buildings with extreme levels of indoor radon. Homes in most of the alluvial valleys in Washington north and northwest of Spokane may be expected to have average indoor radon above 4 pCi/L.

The Blue Mountains probably have moderate radon potential in spite of the low soil uranium content; however, little indoor radon data or soil information is available. Indoor radon data for the counties in this province mostly come from the populated parts of those counties in the adjacent Columbia Basin. However, populated areas of this geologic province in nearby counties in Oregon have indoor radon values averaging 2-4 pCi/L. In those Oregon counties, most housing is on river valley alluvium and it is likely that soil permeabilities in these areas are high, which contributes to modestly elevated indoor radon levels in spite of low soil uranium values.

This is a generalized assessment of the State's geologic radon potential and there is no substitute for having a home tested. The conclusions about radon potential presented in this report cannot be applied to individual homes or building sites. Indoor radon levels, both high and low, can be quite localized, and within any radon potential area there will likely be areas with higher or lower radon potential than assigned to the area as a whole. Any local decisions about radon should not be made without consulting all available local data. For additional information on radon and how to test, contact your State radon program or EPA regional office. More detailed information on state or local geology may be obtained from the State geological survey. Addresses and phone numbers for these agencies are listed in chapter 1 of this booklet.

TABLE 2. Radon Index (RI) and Confidence Index (CI) for geologic radon potential areas of Washington. See figure 1 for locations of areas. See chapter 1 for discussion of RI and CI.

FACTOR	Coastal mountains and hills		Puget Lowlands		Cascade Range		Columbia River Gorge	
	RI	CI	RI	CI	RI	CI	RI	CI
Indoor radon	1	3	1	3	1	3	2	3
Radioactivity	1	3	1	3	1	2	1	3
Geology	1	2	1	2	1	2	2	2
Soil permeability	2	3	2	3	2	2	3	2
Architecture	1	-	1	-	1	-	1	-
GFE points	0	-	0	-	0	-	0	-
Total	6	10	6	10	6	8	9	10
Ranking	Low High		Low High		Low Mod		Mod High	

FACTOR	Okanogan Highland		Columbia Basin		Blue Mountains	
	RI	CI	RI	CI	RI	CI
Indoor radon	3	3	2	3	2?	1
Radioactivity	2	2	2	3	1	3
Geology	3	3	2	2	2	1
Soil permeability	3	3	2	3	2	1
Architecture	3	-	2	-	2	-
GFE points	+2	-	0	-	0	-
Total	16	11	10	11	9?	6
Ranking	High High		Mod High		Mod Low	

- Not used in CI.

RADON INDEX SCORING:

Radon potential category	Point range	Probable screening indoor radon average for area
LOW	3-8 points	< 2 pCi/L
MODERATE/VARIABLE	9-11 points	2 - 4 pCi/L
HIGH	> 11 points	> 4 pCi/L

Possible range of points = 3 to 17

CONFIDENCE INDEX SCORING:

LOW CONFIDENCE	4 - 6 points
MODERATE CONFIDENCE	7 - 9 points
HIGH CONFIDENCE	10 - 12 points

Possible range of points = 4 to 12

REFERENCES CITED IN THIS REPORT
AND OTHER REFERENCES RELEVANT TO RADON IN WASHINGTON

- Asmerom, Yemane, 1981, Geochemistry of biotite from a part of the Loon Lake Batholith and its relationship to uranium mineralization at the Midnite uranium mine, Stevens County, Washington: Master's thesis, Eastern Washington University, Cheney, WA, 110 p.
- Beyer, W. C., 1981, Petrology and genesis of a uranium-bearing system of pegmatite dikes, Nancy Creek area, northeastern Washington: Master's thesis, University of Montana, Missoula, MT, 82 p., 1 plate.
- Cady, J. W. and Fox, K. F., Jr., 1984, Geophysical interpretation of the gneiss terrane of northern Washington and southern British Columbia, and its implications for uranium exploration: U.S. Geological Survey Professional Paper 1260, 29 p., 1 plate.
- Castor, S. B., Berry, M. A. and Robins, J. W., 1977, Uranium and thorium content of intrusive rocks in northeastern Washington and northern Idaho, *in* Bendix Field Engineering Corporation, 1977 NURE uranium geology symposium; abstracts and visual presentations, Grand Junction, CO, Dec. 7-8, 1977: U.S. Department of Energy Report No. GJBX-12(78), p. 167-172.
- Chou, Gin, 1983, Geochemistry of the Togo metapelites from the Midnite uranium mine, northeastern Washington: Master's thesis, Eastern Washington University, Cheney, WA, 94 p.
- Duval, J. S., Jones, W. J., Riggle, F. R., and Pitkin, J. A., 1989, Equivalent uranium map of the conterminous United States: U.S. Geological Survey Open-File Report 89-478, 10 p.
- Duval, J. S., Otton, J. K. and Jones, W. J., 1989, Radium distribution map and radon potential in the Bonneville Power Administration service area: U. S. Geological Survey Open-File Report 89-0340, 125 p.
- Flanigan, V. J., 1976, Geophysical survey of uranium mineralization in alaskitic rocks, eastern Washington: U. S. Geological Survey Open-File Report 76-0679, 25 p.
- Gay, S. P., Jr., 1981, Spokane Mountain Deposit, NW U.S.A.; a uranium discovery resulting from aeromagnetic lineaments analysis in a Precambrian metamorphic terrane: *in* Gabrielson, R. H., Ramberg, I. B., Roberts, D. and Steinlein, O. A., eds., Proceedings of the International Conference on Basement Tectonics, n. 4, p. 145-156.
- Johnson, S. Y., Otton, J. K. and Macke, D. L., 1987, Geology of the Holocene surficial uranium deposit of the north fork of Flodelle Creek, northeastern Washington: Geological Society of America Bulletin, v. 98, n. 1, p. 77-85.

- Kinart, K. P. and Ikramuddin, Mohammed, 1980, Geochemistry of part of the Loon Lake batholith and its relationship to uranium mineralization at the Midnite mine, northeastern Washington: Geological Society of America Abstracts with Programs, v. 12, n. 7, p. 463.
- Lasmanis, Raymond, 1991, Geology of Washington: Rocks and Minerals Magazine, v. 66, n. 4, p.262-277.
- Lott, Thomas L., Jr., 1982, Petrography, major element chemistry, and geology of uraniferous igneous rocks in the Turtle Lake Quadrangle, Washington: Master's thesis, University of Georgia, Athens, GA, 107 p.
- Milne, P. C., 1980, Uranium in Washington: Washington Geologic Newsletter, v. 8, n. 2, p. 1-5
- Mitchell, T. E., 1981, Uranium mineralization of the metamorphic aureole of the Spirit pluton, Stevens County, Washington: Master's thesis, Oregon State University, Corvallis, OR, 109 p., 2 plates.
- Moed, B.A., Nazaroff, A.V., Nero, M.B., Schwehr, M.B., and Van Heuvelen, A., 1984, Identifying areas with potential for high indoor radon levels: analysis of the National Airborne Radiometric Reconnaissance data for California and the Pacific Northwest: Berkeley, California, Lawrence Berkeley Laboratory Report No. LBL-16955, 70 p.
- Nash, J. T. and Lehrman, N. J., 1975, Geology of the Midnite uranium mine, Stevens County, Washington; a preliminary report: U. S. Geological Survey Open-File Report 75-0402, 36 p.
- Nash, J. T. and Lehrman, N., 1975, The Midnite uranium mine, Stevens County, Washington: U. S. Geological Survey Professional Paper 975, 29 p.
- Nash, J. T. and Lehrman, N. J., 1976, Secondary enrichment of uranium, Midnite Mine: U. S. Geological Survey Professional Paper 1000, 38 p.
- Nero, A.V., Schwehr, M.B., Nazaroff, W.W., and Revzan, K.L., 1986, Distribution of airborne radon-222 concentrations in U.S. homes: Science, v. 234, p.992-997.
- Otton, James K., 1987, Indoor radon; geologic controls in Pacific Northwest: Geological Society of America Abstracts with Programs, v.19, n.2, p. 122.
- Otton, J.K. and Duval, J.S., 1990, Geologic controls on indoor radon in the Pacific Northwest, in U.S. Environmental Protection Agency, The 1990 international symposium on radon and radon reduction technology: Volume III. Preprints, unpaginated.
- Pitkin, J. A., 1975, Truckborne gamma-ray spectrometry in northeastern Washington: in Craig, L. C., Brooks, R. A. and Patton, P. C., eds., Abstracts of the 1975 Uranium and Thorium Research and Resources Conference: U.S. Geological Survey Open-File Report 75-595, p. 35.

Rose, A.W., Ciolkosz, E.J., and Washington, J.W., 1990, Effects of regional and seasonal variations in soil moisture and temperature on soil gas radon, *in* U.S. Environmental Protection Agency, The 1990 international symposium on radon and radon reduction technology: Volume III. Preprints, unpaginated.

Schuster, E.J., 1990, Geologic map of Washington: Washington Division of Geology and Earth Resources, Olympia, 1 plate.

U.S. Geological Survey, 1978, Geology of Washington: Washington Division of Geology and Earth Resources Reprint 12, p. 13-51, 1 plate.

Wicklund, M. A., 1984, Geology and genesis of the rocks of the Graeber uranium area, Stevens County, Washington: Master's thesis, Eastern Washington University, Cheney, WA, 160 p., 2 plates.

Zielinski, R. A., Bush, C. A. and Rosholt, J. N., 1986, Uranium series disequilibrium in a young surficial uranium deposit, northeastern Washington, U.S.A.: Applied Geochemistry, v. 1, n. 4, p. 503-511.