



PRELIMINARY GEOLOGIC MAP OF THE ROGERS LAKE NORTH QUADRANGLE, KERN COUNTY, CALIFORNIA

by A.W. Ward and G.L. Dixon

Open-file Report 93-694

2002

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

**U. S. DEPARTMENT OF THE INTERIOR
U. S. GEOLOGICAL SURVEY**

INTRODUCTION

The Rogers Lake North Quadrangle is located in the Antelope Valley region of the western Mojave Desert of Southern California. The Antelope Valley is a small, closed basin, rimmed by the San Gabriel Mountains, the Sawmill-Liebre Mountains, and the Tehachapi Mountains to the south, southwest, and northwest, respectively, and by various buttes and pediments to the north and east. The most notable boundaries, however, are the Garlock Fault to the north, which separates the Mojave province from the Basin and Range province, and the San Andreas Fault to the south, which separates the Transverse Range province (Fig. 1).

The western Mojave province, in which the Antelope Valley Basin lies, is characterized by many deep Cenozoic sedimentary basins that are separated by low hills of Mesozoic granitic and metasedimentary rocks and Tertiary volcanic rocks. These basins, which surface and geophysical observations show to trend northeast (parallel to the Garlock Fault), were depocenters for upper Tertiary and lower Quaternary sediments; many contain more than 3 km of alluvium. During Tertiary time, the area drained southwest into the Pacific Ocean. Tectonic activity along the San Andreas and Garlock faults and the uplifting of the Transverse, Tehachapi, and southern Sierra Nevada ranges reversed this drainage pattern and formed new closed basins in the Mojave Desert. In Quaternary time these basins were occupied by pluvial lakes; one in particular, Lake Thompson, covered about 500 km² and was perhaps as deep as 20 m. Today Lake Thompson is represented by the playas Buckhorn, Rosamond, and Rogers dry lakes (Thompson, 1929).

STRATIGRAPHIC UNITS

Local rock units in the area include tuffs, lava flows, and sediments of the Tertiary Tropico Formation, Tertiary fanglomerates, Tertiary intrusive rhyolites and dacites, and deeply weathered Mesozoic granitic rocks. Dibblee (1967) gives excellent background on this subject. Exposed local Quaternary geologic units include sandy alluvium, playa clay, and beach deposits, most of which are late Quaternary in age and related to pluvial Lake Thompson.

Late Quaternary and Holocene drainages into the playas are from the west and south, along the Garlock and San Andreas faults. Much of the surface west of the playas is exposed granitic rocks and fine pediment gravel, cut by well-defined shallow arroyos. The eastern side of Rogers Lake has a gentler slope characterized by several long, low alluvial fans and fan segments, some of which were periodically inundated with fluctuations in Lake Thompson. South of the playas, long channels have supplied sediment to both the former and modern lakes, channels, such as Big Rock Creek and Little Rock Creek, associated with large alluvial fans along the front of the San Gabriel Mountains. Ponti and Burke (1980) and Ponti and others (1981) have characterized the exposed alluvium as late Pleistocene (90,000 ybp) to Holocene (4,000 ybp) medium grained, moderately to well sorted, moderately to well stratified sediments with textural and soil development characteristics that are dominated by the addition of calcium carbonate during groundwater fluctuations of pluvial lake stands.

Other Quaternary and Tertiary units in the area have been described by well data (Dutcher and Worts, 1963; Neal and Motts, 1967; Neal, 1968). These units are principally alluvial fan and channel deposits from the south as the Transverse Range (San Gabriel Mts.) were uplifted, and the north as the Tehachapis shed debris.




Figure 1. Sketch map showing major features in the western Mojave Desert.

STRUCTURAL SETTING

Maybey (1960) first recognized the East Antelope Basin from a 40-mgal gravity low and suggested that the steep gravity gradients may be fault controlled. The gravity signature suggests that this is a northeast-trending basin, which is deepest just southwest of Rosamond Lake and shallows to the northeast. Confirmation of this being a fault-bounded basin (at least on the northwest side) has come from gravity data (Morin and others, in press) and also from a series of resistivity anomalies (Zohdy, 1990), as well as subtle surface escarpments, alignment of historical spring discharge mounds trending N30-45E, and steep hydrologic gradients (Ward and others, 1991). The southeast boundary of the basin is less apparent, however, but is definable from field observations and geophysics.

Known and inferred faults in the immediate area trend northwest. Major faults can be recognized from juxtaposed bedrock units and associated with long, linear valleys (with Quaternary and Holocene alluvium) that lead to the playa. Dibblee (1967) considers that several such features may have controlled the development and shape of Rogers Lake; the El Mirage fault and several unnamed faults along the north and south edges of the dry lake define and give the playa its hourglass shape. We consider further that the El Mirage Fault probably extends beneath the central portion of the playa and connects with the Bissell Hills fault.

We consider that a geographic "corridor" in the southern part of Edwards AFB is a geologic fault zone, which we herein informally called the Antelope Valley Fault Zone (AVFZ; Ward and Dixon, 1994; Dixon and Ward, 1994a,b). This corridor trends approximately N30E to N40E, and extends from south of Rosamond Playa to the southwestern part of Rogers Lake, a distance of at least 10 km.

Ward and Dixon (1994) and Dixon and Ward (1994a,b) recognize no disturbances of the surface in the AVFZ. This corridor is characterized, however, by several depressions that are former (historically active) spring discharge areas in the adjacent Redman Quadrangle. These springs (approximately 30 m in diameter) are circular depressed features, containing loamy soil and peat and commonly surrounded by dead mesquite. In the Rogers Lake North Quadrangle we recognize complex deposits on Rogers Lake at South Base that overlie the projected AVFZ.

GEOLOGIC UNITS IN THE ROGERS LAKE NORTH QUADRANGLE

- D Disturbed Ground (Holocene)-- Surfaces disrupted by construction and development
- Qah Alluvium (Holocene)--Sand and silt (and minor gravel) in active channels and dry washes. Contains unconsolidated and unsorted granitic sediments; fines often have been reworked by wind into ripples
- Qes Eolian Sand (Holocene and Late Pleistocene)--Fine to medium and coarse sand composed of feldspar and quartz, principally on the lee of hills marginal to Rogers Lake

- Qaf Fan Alluvium (Late Pleistocene)**--Thin deposits of alluvium in fans on slopes immediately adjacent to pluvial Lake Thompson. Moderately sorted sediment derived principally from local granitic rocks, with minor amounts of volcanic and metamorphic rock fragments. On northwest side of Rogers Lake, deposits are undifferentiated (Qaf). On east side, however, several deposits can be distinguished tentatively by topographic expression and stratigraphic position (older deposits tend to be higher and incised by channels leading to lower, younger fans. Mapped on east side as undifferentiated (Qafu) and older to younger (Qaf1-Qaf4) deposits. Undifferentiated units Qaf and Qafu are considered equivalent to Quca of Ponti and Burke (1980) and Ponti and others (1981) -- late Pleistocene (90,000 ybp) to Holocene (4,000 ybp), undifferentiated alluvial sediments around pluvial lake shorelines; medium grained, moderately to well sorted, moderately to well stratified, but with textural and soil development characteristics that are dominated by the addition of calcium carbonate during groundwater fluctuations of pluvial lake stands. Differentiated units Qaf1-4 correspond to Q4-Q5 sediments of Ponti and Burke (1980) and Ponti and others (1981) in age ranges of 90,000-17,000 ybp (Q4-Q5). Units may also be truncated by wave action during high lake stands.
- Qpc Playa Deposits (Late Pleistocene)**--Clay and silt of Lake Thompson, thinly stratified on flat floor of Rogers Dry Lake. New exposure occurs in 800-m fissure in southeast part of dry lake (Ward and others, 1991)
- Qbb Beach Bar (Late Pleistocene)**--Ridges and hills of sand and gravel marking shoreline of Lake Thompson, stratified, with 1-2-cm-thick gravel layers alternating with 1-m-thick sand beds. May have eolian sand cap of <1m
- Qcs Clayey Sand (Late Pleistocene)**--Moderately indurated, cohesive beach deposits, composed of coarse to fine sands, silt, and clay, at margins of Rogers Lake. Deposits are indurated by the desiccation of clays but also contain CaCO₃. Stratified layers and bedforms suggest these were parabolic lunette dunes formed by southwesterly winds off Lake Thompson. At northeast side of Rogers Lake, these deposits are carved by wind into yardangs (Ward and Greeley, 1984)
- Qsc Sandy Clay (Late Pleistocene)**--Stratified, silty clay deposits interbedded with clayey sand deposits (Qcs), but may be transitional between playa clay deposits (Qpc) and clayey sands (Qcs). Also carved into yardangs at northeast side of Rogers Lake
- Kiq1 Quartz Latite Dike (Cretaceous?)**--Dike intruding quartz monzonite (Klqm). Quartz latite is light reddish gray to white, contains less than 10% phenocrysts. Slightly altered. Dike is faintly flow banded. Approximately 1m wide
- Kihd Hornblende Diorite Dike (Cretaceous?)**--Dikes intruding leucogranite (Klg) and quartz monzonite (Klqm). Dark green to gray, coarse grained. Hornblende up to 5 mm
- Klg Leucogranite (Cretaceous?)**-- Leucocratic, biotitic granite and granodiorite, weakly foliated; fine to medium grained; color index 1-5. Composed primarily of quartz, potassic feldspar, and plagioclase. Minor minerals (chiefly biotite) less than 1%. Intrudes underlying quartz monzonite (Klqm). Intruded by dikes of granite and hornblende diorite (Tihd)


Klqm **Quartz Monzonite/Monzodiorite (Cretaceous?)**--Light grayish white, medium grained granitic rocks. Color index 5-10; dark where altered. Composed of quartz, potassic feldspar, and plagioclase. Biotite is dark where fresh, minor hornblende, and sphene (3-4 mm). Surface is weathered to depths of three to ten m (altered gneiss) at most exposures

REFERENCES


- Dibblee, T.W., Jr., 1967, Areal geology of the western Mojave Desert, California: U.S. Geological Survey Professional Paper 522, 153 p.
- Dixon, G.L. and Ward, A.W., 1994a, Preliminary geologic map of the Edwards Quadrangle, Kern County, California, U.S. Geological Survey open-file Map; map scale 1:24,000, in press.
- Dixon, G.L. and Ward, A.W., 1994b, Preliminary geologic map of the Rogers Lake South Quadrangle, Los Angeles and Kern Counties, California, U.S. Geological Survey open-file map; map scale 1:24,000, in press.
- Dutcher, L.C. and Worts, G.F., Jr., 1963, Geology, Hydrology, and Water Supply of Edwards Air Force Base, Kern County, California, U.S. Geological Survey Open-File Report 63-146, 240 p.
- Maybey, D.R., 1960, Gravity survey of the western Mojave Desert, California: U.S. Geological Survey Professional Paper 316-D, p. 51-73.
- Morin, Robert L., Mariano, John, and Jachens, Robert C., in press, Isostatic residual gravity map of Edwards Air Force Base and vicinity, Kern, Los Angeles, and San Bernardino counties, California, U.S. Geological Survey open-file map; scale 1:62,500.
- Neal, J.T. (ed.) 1968, Playa Surface Morphology: Miscellaneous Investigations, Air Force Cambridge Research Laboratories Environmental Research Papers, 235, 150 p.
- Neal, J.T. and Motts, W.S., 1967, Recent geomorphic changes in playas of western United States, Jour. Geology, 75, 511-525.
- Ponti, D.J. and Burke, D.B., 1980, Map showing Quaternary geology of the eastern Antelope Valley and vicinity, California: U.S. Geol. Surv. Open-file map 80-1064, scale 1:62,500.
- Ponti, D.J., Burke, D.B., and Hedel, C.W., 1981, Map showing Quaternary geology of the central Antelope Valley and vicinity, California: U.S. Geol. Surv. Open-file map 81-737, scale 1:62,500.
- Thompson, D.G., 1929, The Mojave Desert Region, California, U.S. Geological Survey Water-Supply Paper 578, 759 p.
- Ward, A.W. and Dixon, G.L., 1993, Preliminary geologic map of the Redman Quadrangle, Los Angeles and Kern Counties, California, U.S. Geological Survey open-file map; map scale 1:24,000.
- Ward, A.W., Dixon, G.L., Galloway, D.L., and Mariano, John, 1991, Land subsidence at Edwards AFB, Mojave Desert, California: Groundwater withdrawal or neotectonics?: Geol. Soc. Amer., Abstr. w/prog., 23, 5, A477.
- Ward, A.W. and Greeley, Ronald, 1984, Evolution of the Yardangs at Rogers Lake, California: Bull. Geol. Soc. Amer., 95, 829-837.
- Zohdy, A.A.R., 1990, Groundwater exploration using deep Schlumberger soundings at Edwards AFB, California, Part I: Graham Ranch and Rogers Lake: U.S. Geological Survey Open-File Report 90-536, 95 p.


_____ . . . Contact, dashed where inferred, dotted where concealed.

_____ • • Fault, dashed where inferred, dotted where concealed.

 Igneous dike.

 Strike of vertical foliation.

 Strike and dip of joint.

 Strike and dip of fault plane.

CORRELATION OF UNITS

Qah	D							Holocene	QUATERNARY
Qes							Holocene/Pleistocene		
Qaf4	Qafu	Qaf	Qsc	Qcs	Qpc	Qbb	Late Pleistocene		
Qaf3									
Qaf2									
Qaf1									

unconformity

Kihd	Kiql
Klg	Klqm

CRETACEOUS?