

U.S. DEPARTMENT OF THE INTERIOR  
U.S. GEOLOGICAL SURVEY

PRELIMINARY GEOLOGIC MAP OF THE SITKA QUADRANGLE AND PARTS OF THE SITKA  
SW, SUGARLOAF, AND TROUT CREEK QUADRANGLES, CLARK COUNTY, KANSAS

By

MORRIS W. GREEN<sup>1</sup> and GEORGE M. FAIRER<sup>1</sup>

Open-File Report 94-658

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government

<sup>1</sup> Denver, Colorado



1994

U.S. Geological Survey Open-File Report 94-658



Preliminary geologic map of the Sitka  
 a quadrangle and parts of the Sitka SW, Sugarloaf, and Trout Creek  
 quadrangles, Clark County, Kansas

By Morris W. Green and George M. Fairer



CORRELATION OF MAP UNITS



Unconformity



Unconformity



## DESCRIPTION OF MAP UNITS

Qa1

Alluvium (Holocene)--Unconsolidated gravel, sand, silt, and clay within and adjacent to major stream channels and flood plains. Locally includes dune sand, slopewash, colluvium, and terrace deposits; locally intertongues with loess (Q1) and eolian sand (Qds) deposits

Q1

Loess deposits (Holocene and Pleistocene?)--Eolian deposits of nonstratified silt and minor amounts of sand (5-15 percent) and clay; deposits are porous and calcareous; commonly contain nodules, carbonate-filled root casts, fossil-plants, and fossil-animals. Cementation and the cohesiveness of sediment particles cause loess deposits to form characteristic near-vertical scarps in dissected deposits; locally includes slopewash and colluvium; mantles bedrock and other associated surficial deposits; locally forms dunes. Thickness 0-40 ft

Qds

Eolian dune sand (Holocene and Pleistocene?)--Eolian deposits of stratified and nonstratified fine to very fine sand and minor amounts of silt and clay in irregularly shaped eolian dunes which may be locally altered and reworked in ephemeral ponds and on flood plains by fluvial processes; deposits are porous and locally calcareous; commonly contain nodules, sand concretions, and carbonate-filled root casts. Locally intertongues with slopewash, alluvium, and colluvium; mantles bedrock and associated loess deposits (Q1) and occurs dominantly in proximity to major streams and tributary drainages; locally includes deposits of sandy loess. Thickness 0-60 ft

Kk

Kiowa Formation (Lower Cretaceous)--Dark-gray to black gypsiferous shale; locally contains thin beds of light- to medium-gray coquinoidal limestone and light-yellowish-brown calcareous sandstone. Commonly includes thin beds and lenses of selenite and limestone with cone-in-cone concretionary structure. Shale contains finely disseminated flecks of coaly carbonaceous plant matter in the silt and clay fractions of the formation sediment. Deposition of the formation apparently occurred a considerable distance offshore in a shallow epeiric sea or marine embayment. Formation is unconformable with formations above and below. Pre-Ogallala erosion has truncated the formation in the eastern and southeastern parts of the map area. Formation ranges from 0 to about 20 ft in thickness

--Pdc--

**Day Creek Dolomite (Upper Permian)**--Light-gray, pink, and white, dense, very fine grained crystalline dolomite; locally contains chert nodules and disseminated chert grains. Deposition apparently occurred by chemical precipitation from sea water in a restricted marine embayment under arid conditions. Formation present only locally in northwestern part of the map area where the unit is shown on the map as a dashed contact between the Whitehorse below and the Kiowa Formation above. The unit also forms a caprock on Sugarloaf hill in the NE¼NE¼ sec. 15, T. 33 S., R. 21 W. Conformable with underlying Whitehorse Formation. Formation forms an erosion resistant bench or caprock and is as much as 2 ft thick

Pw

**Whitehorse Formation (Upper Permian)**--Reddish-orange, nonstratified, argillaceous siltstone; locally includes some thin beds of very fine grained, locally crossbedded, silty sandstone and thin beds of dolomite; contains beds of gypsum and anhydrite in subsurface. Unit is locally bleached white along bedding planes, is moderately friable, and contains sandy calcareous concretions that resemble musket balls. Silt and sand are well sorted, well rounded, and consist mostly of iron oxide-stained quartz but also some dark accessory minerals. Upper part of unit is locally impregnated with secondary calcium carbonate. Sediment deposition apparently occurred subaqueously from the settling of suspended sediment in relatively shallow marine waters of a restricted embayment. The character of the detrital fraction of Whitehorse rocks suggests that detrital sediment was transported into the embayment primarily as loess winnowed by eolian activity from dune fields adjacent to the embayment; apparently subaqueous currents acted locally to form low-angle crossbedded units. Beds are characteristically wavy because of deformation caused by dissolution of gypsum and anhydrite. Thickness about 250 ft

Pb

**Blaine Formation (Lower Permian)**--Light-gray to white gypsum locally interbedded with thin beds of reddish-orange siltstone. Deposition apparently occurred by chemical precipitation during the evaporation of sea water in a restricted marine embayment. Base of formation not exposed in map area. Exposed thickness about 50 ft

—————

CONTACT

- - - 1 - - -

NORMAL FAULT--Dashed where  
approximately located.  
Bar and ball on downthrown  
side


5  
———  
1

STRIKE AND DIP OF BEDS

Inclined

⊕

Horizontal



Index map of Clark County showing the location of 7.5' quadrangles (1-20), this geologic map report (F), and other published geologic maps (A-E) covering the county. \*, map report listed by letter in References Cited

7.5' quadrangles:

- |                    |                   |
|--------------------|-------------------|
| (1) Minneola       | (11) Profitt Lake |
| (2) Bloom          | (12) Ashland NW   |
| (3) Simmons Creek  | (13) Ashland      |
| (4) Mount Jesus NW | (14) Sitka        |
| (5) Turkey Creek   | (15) Sugarloaf    |
| (6) Fowler SE      | (16) Mount Helen  |
| (7) Bloom SW       | (17) Englewood    |
| (8) Bloom SE       | (18) Ashland SE   |
| (9) Mount Jesus    | (19) Sitka SW     |
| (10) Lexington     | (20) Trout Creek  |

## References Cited

- (A) Fairer, George M. and Green, Morris W., 1992, Preliminary geologic map of the Bloom SW quadrangle and parts of the Bloom, Minneola, and Fowler SE quadrangles, Clark County, Kansas: U.S. Geological Survey Open-file Report 92-698, 1 pl., 10 p., scale 1:24,000.
- (D) Fairer, George M. and Green, Morris W., 1994, Preliminary geologic map of the Ashland NW quadrangle and parts of the Englewood, Profitt Lake, and Mount Helen quadrangles, Clark County, Kansas: U.S. Geological Survey Open-file Report 94-655, 1 pl., 7 p., scale 1:24,000.
- (E) Fairer, George M. and Green, Morris W., 1994, Preliminary geologic map of the Ashland quadrangle and part of the Ashland SE quadrangle, Clark County, Kansas: U.S. Geological Survey Open-file Report 94-657, 1 pl., 6 p., scale 1:24,000.
- (B) Green, Morris W. and Fairer, George M., 1992, Preliminary geologic map of the Bloom SE quadrangle and part of the Simmons Creek quadrangle, Clark County, Kansas: U.S. Geological Survey Open-file Report 92-697, 1 pl., 7 p., scale 1:24,000.
- (C) Green, Morris W. and Fairer, George M., 1994, Preliminary geologic map of the Mount Jesus quadrangle and parts of the Mount Jesus NW, Lexington, and Turkey Creek quadrangles, Clark County, Kansas: U.S. Geological Survey Open-file Report 94-656, 1 pl., 8 p., scale 1:24,000.