

U. S. DEPARTMENT OF THE INTERIOR
U. S. GEOLOGICAL SURVEY

**PRELIMINARY GEOLOGIC MAP OF THE CALABASAS
7.5' QUADRANGLE, SOUTHERN CALIFORNIA:
A DIGITAL DATABASE**

Compiled by R. F. Yerkes¹ and R. H. Campbell²

Open - File Report 95-51

This report is preliminary and has not been reviewed for conformity with U. S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U. S. Government.

This database, identified as "Preliminary Geologic Map of the Calabasas 7.5' Quadrangle, southern California: A Digital Database," has been approved for release and publication by the Director of the USGS. Although this database has been reviewed and is substantially complete, the USGS reserves the right to revise the data pursuant to further analysis and review. This database is released on condition that neither the USGS nor the U. S. Government may be held liable for any damages resulting from its use.

1997

¹ Menlo Park, CA 94025

² Reston, VA 20192

INTRODUCTION

This Open-File report is a digital geologic map database. This pamphlet serves to introduce and describe the digital data. There is no paper map included in the Open-File report.

This digital map database is compiled from previously published sources combined with some new mapping and modifications in nomenclature. The geologic map database delineates map units that are identified by general age and lithology following the stratigraphic nomenclature of the U. S. Geological Survey. For detailed descriptions of the units, their stratigraphic relations and sources of geologic mapping consult Yerkes and Campbell (1993). More specific information about the units may be available in the original sources.

The content and character of the database and methods of obtaining it are described herein. The geologic map database itself, consisting of four ARC coverages and one base layer, can be obtained over the Internet or by magnetic tape copy as described below. The processes of extracting the geologic map database from the tar file, and importing the ARC export coverages (procedure described herein), will result in the creation of an ARC workspace (directory) called '**calab.**'

The database was compiled using ARC/INFO version 7.0.3, a commercial Geographic Information System (Environmental Systems Research Institute, Redlands, California), with version 3.0 of the menu interface ALACARTE (Fitzgibbon and Wentworth, 1991, Fitzgibbon, 1991, Wentworth and Fitzgibbon, 1991). It is stored in uncompressed ARC export format (ARC/INFO version 7.x) in a compressed UNIX tar (tape archive) file. The tar file was compressed with gzip, and may be uncompressed with gzip, which is available free of charge via the Internet from the gzip Home Page (<http://w3.teaser.fr/~jlgailly/gzip>). A tar utility is required to extract the database from the tar file. This utility is included in most UNIX systems, and can be obtained free of charge via the Internet from Internet Literacy's Common Internet File Formats Webpage (<http://www.matisse.net/files/formats.html>). ARC/INFO export files (files with the .e00 extension) can be converted into ARC/INFO coverages in ARC/INFO (see below) and can be read by some other Geographic Information Systems, such as MapInfo via ArcLink and ESRI's ArcView (version 1.0 for Windows 3.1 to 3.11 is available for free from ESRI's web site: <http://www.esri.com>).

This document replaces the original digital database description published in 1995. Changes in this release include the following:

1. Different base layer - The original digital database included separates clipped out of the Los Angeles 1:100,000 sheet. This release includes a vectorized scan of a scale-stable negative of the Calabasas 7.5 minute quadrangle.
2. Map projection - The files in the original release were in polyconic projection. The projection used in this release is state plane, which allows for the tiling of adjacent quadrangles.
3. File compression - The files in the original release were compressed with UNIX compression. The files in this release are compressed with gzip.

OBTAINING THE DIGITAL DATA

The digital database package can be obtained in any of three ways:

1. Sending a tape with request

A 18 MB compressed tar file of the geologic map database and related files can be obtained by sending a tape with request and return address to:

Calabasas, CA 7.5' Geologic Database
c/o Database Coordinator
U.S. Geological Survey
345 Middlefield Road, M/S 975
Menlo Park, CA 94025

The compressed tar file will be returned on the tape. The acceptable tape types are:

1/4 inch, 150 MB cartridge tape
2.3 or 5.0 GB, 8mm Exabyte tape.

2. Anonymous ftp over the Internet

To obtain the tar file by anonymous ftp do the following:

cd local_directory	• go to a directory to receive the tar file
ftp wrgis.wr.usgs.gov	• make ftp connection with the USGS computer WRGIS
Name: anonymous	• use "anonymous" as your user name
Password: your email address	• use your email address as a password
cd pub/geologic	• go down to the pub/geologic directory
cd ca/of95-51	• go down to the open file directory
type binary	• change transfer type to binary
get calab.tar.gz	• copy the compressed tar file across Internet to the receiving directory
quit	• close the ftp connection

3. From the Western Region Geologic Information Web Page.

To obtain the tar file via the World Wide Web:

The U.S. Geological Survey now supports a set of graphical pages on the World Wide Web. Digital geologic publications (including this one) can be accessed via these pages. The web page for digital geologic publications from the Western Region (including this one) is "<http://wrgis.wr.usgs.gov>". Once at the main page, click on '**Geologic Map Databases**' under the heading '**Data Online**'; next click on '**California**.' Scroll down to get to the listing for this database. Set your web browser to save to a local disk and click on the link labeled 'calab.tar.gz' to download the compressed tar file that contains the Calabasas geologic map database.

EXTRACTING THE GEOLOGIC MAP DATABASE FROM THE TAR FILE

If you obtained the database package on tape:

put the tape in your tape drive	
cd local_directory	-go to a directory to receive the tar file
tar xvfb /dev/rstn 20	-/dev/rstn is the tape device with n an integer, this puts the tar file in local_directory; 20 is the block size of the tar file
gzip -d calab.tar.gz	-makes a 73 MB uncompressed tar file calab.tar
cd local_directory2	-go to the directory that will hold the directory calab (if different from local_directory)
tar xvfb {path to tar file}/calab.tar 20	-extract the calab directory from the tar file; 20 is the block size of the tar file.

If you obtained the database package by anonymous ftp or from the web page:

gzip -d calab.tar.gz	-makes a 73 MB uncompressed tar file calab.tar
cd local_directory2	-go to the directory that will hold the directory calab (if different from local_directory)
tar xvfb {path to tar file}/calab.tar 20	-extract the calab directory from the tar file.

Each of the processes described above will create a directory "calab" that contains the ARC export files and supporting files as described below. The directory structure at this point will look like this:

```
calab/
  cb-geol.e00
  cb-strc.e00
  cb-wells.e00
  cb-foss.e00
  cb-topo.e00
  calab.ps
  calab.txt
  import.aml
```

GEOLOGIC MAP DATABASE CONTENTS

The geologic map database consists of four separate layers and one base layer. Each of these layers (coverages) has been converted to uncompressed ARC/INFO export files. The ARC export files and the associated ARC/INFO coverages, as well as the additional digital material included in the database package, are described below:

ARC/INFO export file	Resultant Coverage	Description of Coverage
cb-geol.e00	cb-geol	Depositional contacts, faults, and unit labels
cb-strt.e00	cb-strt	Strike and dip information, fold axes
cb-wells.e00	cb-wells	Well localities, supporting data given in Yerkes and Campbell (1993), and Yerkes and Showalter (1990)
cb-foss.e00	cb-foss	Fossil localities, supporting data given in Yerkes and Campbell (1993)
cb-topo.e00	cb-topo	Topographic base map taken from a scan of a composite negative of the Calabasas 7.5 minute quadrangle (contour interval = 25 ft.)

ASCII text files and PostScript plot files:

calab.ps	This file.
calab.txt	A text-only file containing an unformatted version of calab.ps
import.aml	ASCII text file in ARC Macro Language to convert these ARC export files to ARC coverages in ARC/INFO.

The following directory is produced in the process of converting the export files into ARC coverages:

info/	INFO directory containing the database files that accompany each ARC/INFO layer (coverage).
-------	---

Once the ARC export coverages have been imported (see discussion below), the Calabasas (calab) directory, or ARC workspace, will look like this:

```
calab/
  info/
  cb-geol/
  cb-strt/
  cb-foss/
  cb-wells/
  cb-topo/
  calab.ps
  calab.txt
  import.aml
```

CONVERTING ARC EXPORT FILES

ARC export files are converted to ARC coverages using the ARC command IMPORT with the option COVER. In order to ease conversion and to maintain naming conventions, we have included an ASCII text file in ARC Macro Language that will convert all of the export files in the database into coverages and create the associated INFO directory. Change directories to the calab/ directory. From the ARC command line type:

Arc: &run import.aml

ARC export files can also be read by some other Geographic Information Systems. Please consult your GIS documentation to see if you can use ARC export files and the procedure to import them.

DIGITAL COMPILATION

The geologic map information was digitized from stable originals of the geologic maps at 1:24,000 scale. The author manuscripts (pencil on mylar) were scanned using a monochrome scanner with a resolution of 800 dots per inch. The scanned images were vectorized and transformed from scanner coordinates to projection coordinates with digital tics placed by hand at quadrangle corners. The scanned lines were edited interactively by hand using graphical user interface ALACARTE (Fitzgibbon, 1991, Fitzgibbon and Wentworth, 1991, Wentworth and Fitzgibbon, 1991). Scanning artifacts significant for display at a scale of 1:24,000 were corrected.

BASE MAPS

The base map layer (cb-topo) was prepared by scanning a scale-stable composite negative image of the Calabasas 7.5 minute topographic map (1967 edition). The negative was scanned using a monochrome scanner with a resolution of 400 dots per inch. The raster scan was converted to a GRID in ARC/INFO. The GRID was then vectorized and reprojected into State Plane Projection using the four corner tics as reference points. The arcs in the base layer have not been attributed. The base map is provided for reference only.

SPATIAL RESOLUTION

Uses of this digital geologic map should not violate the spatial resolution of the data. Although the digital form of the data removes the constraint imposed by the scale of a paper map, the detail and accuracy inherent in map scale are also present in the digital data. The fact that this database was edited at a scale of 1:24,000 means that higher resolution information is not present in the dataset. Plotting at scales larger than 1:24,000 will not yield greater real detail, although it may reveal fine-scale irregularities below the intended resolution of the database. Similarly, where this database is used in combination with other data of higher resolution, the resolution of the combined output will be limited by the lower resolution of these data.

FAULTS AND LANDSLIDES

This database is not sufficiently detailed or comprehensive to identify or characterize site-specific hazards represented by faults or landslides shown; the faults shown do not take the place of fault-rupture hazard zones designated by the California State Geologist (see Hart, 1988).

DATABASE SPECIFICS

The map databases consist of ARC coverages which are stored in State Plane (California coordinate system) projection (Table 1). Digital tics define a 2.5 minute grid of latitude and longitude in the geologic coverages corresponding with quadrangle corners and internal tics. In the base map layer, the tics define a 7.5 minute grid, corresponding with quadrangle corners.

Table 1 - Map Projection

The map is stored in State Plane projection

PROJECTION	STATE PLANE
UNITS	METERS
ZONE	3426
DATUM	NAD27
PARAMETERS	

-on the ground
-Arc/Info State Plane zone
corresponding to California
coordinate system zone 7

-none

The content of the geologic database can be described in terms of the lines and the areas that compose the map. Descriptions of the database fields use the terms explained in Table 2.

Table 2 - Field Definition Terms

ITEM NAME	name of the database field (item)
WIDTH	maximum number of digits or characters stored
OUTPUT	output width
TYPE	B-binary integer, F-binary floating point number, I-ASCII integer, C-ASCII character string
N. DEC.	number of decimal places maintained for floating point numbers

LINES

The lines (arcs) are recorded as strings of vectors and are described in the arc attribute table (Table 3). They define the boundaries of the map units, the boundaries of open bodies of water, and the map boundaries. These distinctions, including the geologic identities of the unit boundaries, are recorded in the LTYPE field according to the line types listed in Tables 4 and 5.

Table 3 - Content of the Arc Attribute Tables (CB-GEOL.AAT, CB-STRC.AAT)

ITEM NAME	WIDTH	OUTPUT	TYPE	N. DEC	
FNODE#	4	5	B		starting node of arc (from node)
TNODE#	4	5	B		ending node of arc (to node)
LPOLY#	4	5	B		polygon to the left of the arc
RPOLY#	4	5	B		polygon to the right of the arc
LENGTH	4	12	F	3	length of arc in meters
<coverage>#	4	5	B		unique internal control number
<coverage>-ID	4	5	B		unique identification number
LTYPE	35	35	C		line type (see Tables 4 and 5)
SEL	1	1	I		user defined field used to save a selected set
SYMB	3	3	I		user defined field used to save symbol assignments (such as color)

Table 4 - Line Types Recorded in the LTYPE Field (CB-GEOL)

contact, approx. located
 fault, approx. located
 fault, concealed
 fault, inferred
 map boundary, certain
 thrust fault, approx. located

Table 5 - Line Types Recorded in the LTYPE Field (CB-STRC)

f.a., anticline, approx. located
 f.a., anticline, concealed
 f.a., syncline, approx. located
 f.a., syncline, concealed

AREAS

Map units (polygons) are described in the polygon attribute table (Table 6). The identities of the map units from compilation sources are recorded in the PTYPE field by map label (Table 7). For a full description of the map units see Yerkes and Campbell (1993).

Table 6 - Content of the Polygon Attribute Tables (CB-GEOL.PAT)

ITEM NAME	WIDTH	OUTPUT	TYPE	N. DEC	
AREA	4	12	F	3	area of polygon in square meters
PERIMETER	4	12	F	3	length of perimeter in meters
<coverage>#	4	5	B		unique internal control number
<coverage>-ID	4	5	B		unique identification number
PTYPE	35	35	C		unit label (see Table 7)
SEL	1	1	I		user defined field used to save a selected set
SYMB	3	3	I		user defined field used to save symbol assignments (such as color)

Table 7 - Map Units (CB-GEOL)

Kc	Tc	Tmb	Ttcc
Qal	Tcab	Tmc	Tw
Qay1	Tcb	Tmd	af
Qay2	TI	Tms	pond
Qls	Tlv	Tsc	res
Qt	Tm	Tss	

POINTS

Point information (strikes and dips, well localities, fossil localities) is recorded as coordinate and related information and is described in the Point Attribute Table (Tables 8, 10 and 11). The identities of point types recorded in the PTYPE field of the CB-STRC.PAT table are shown in Table 9.

Table 8 - Content of the Point Attribute Tables (CB-STRC.PAT)

ITEM NAME	WIDTH	OUTPUT	TYPE	N. DEC	
AREA	4	12	F	3	not used
PERIMETER	4	12	F	3	not used
<coverage>#	4	5	B		unique internal control number
<coverage>-ID	4	5	B		unique identification number
PTYPE	35	35	C		point type (see Table 9)
DIP	3	3	I		dip angle in degrees
STRIKE	3	3	I		strike angle in degrees
SEL	1	1	I		user defined field used to save a selected set
SYMB	3	3	I		user defined field used to save symbol assignments (such as color)

Table 9 - Point Types (CB-STRC)

bedding
flat bedding
ot bedding
vert bedding

Table 10 - Content of the Point Attribute Tables (CB-FOSS.PAT)

ITEM NAME	WIDTH	OUTPUT	TYPE	N. DEC	
AREA	4	12	F	3	not used
PERIMETER	4	12	F	3	not used
<coverage>#	4	5	B		unique internal control number
<coverage>-ID	4	5	B		unique identification number
QUAD.	6	6	C		abbrev. quad. name
SEC	2	2	I		section number
SOURCE	35	35	C		source of data
MAP_NO	4	4	C		point ID used on map
COLL-NO	5	5	C		collector's number
T	2	2	C		township north
RW	2	2	C		range west
AGE	6	6	C		indicated age of collection
UNIT	3	3	C		host map unit

Table 11 - Content of the Point Attribute Tables (CB-WELLS.PAT)

ITEM NAME	WIDTH	OUTPUT	TYPE	N. DEC	
AREA	4	12	F	3	not used
PERIMETER	4	12	F	3	not used
<coverage>#	4	5	B		unique internal control number
<coverage>-ID	4	5	B		unique identification number
QUADRANGLE	6	6	C		abbrev. quad. name
MAP_NUMBER	4	4	I		point ID used on map
TOWNSHIP	2	2	C		township north
RANGE_WEST	2	2	C		range west
SECTION	2	2	I		section number
OPERATOR	35	35	C		operator name
NAME/NUMBER	35	35	C		name/number of well
ELEV.	4	4	I		elevation (ft.)
YEAR	4	4	I		year drilled
TD	5	5	I		total depth (ft.)
BOTTOM	1	1	C		bottom-hole geology

ACKNOWLEDGMENTS

R. H. Campbell, Reston, VA assisted in field interpretations and compilation; P. K. Showalter and C. M. Wentworth supplied critical custom programming; and S. E. Graham updated supporting files for completeness and consistency and prepared this report.

REFERENCES

- Fitzgibbon, T.T., 1991, ALACARTE installation and system manual (version 1.0): U.S. Geological Survey, Open-File Report 91-587B.
- Fitzgibbon, T.T., and Wentworth, C.M., 1991, ALACARTE user interface - AML code and demonstration maps (version 1.0): U.S. Geological Survey, Open-File Report 91-587A.
- Hart, E.W., 1988, Fault-rupture hazard zones in California; Alquist-Priolo Special Studies Zones Act of 1972 with index to special studies zones maps: California Division of Mines and Geology Special Publication 42.
- Wentworth, C. M., and Fitzgibbon, T. T., 1991, ALACARTE user manual (version 1.0): U. S. Geological Survey Open-File Report 91-587C.
- Yerkes, R. F., and Campbell, R. H., 1993, Preliminary geologic map of the Calabasas 7.5 minute quadrangle, southern California: U. S. Geological Survey Open-File Report 93-205.
- Yerkes, R. F., and Showalter, 1990, Exploratory wells drilled in the Los Angeles, California 1:100,000 quadrangle, southern California: U. S. Geological Survey Open-File Report 90-627.