

# SURFACE-WATER-QUALITY ASSESSMENT OF THE YAKIMA RIVER BASIN IN WASHINGTON: SPATIAL AND TEMPORAL DISTRIBUTION OF TRACE ELEMENTS IN WATER, SEDIMENT, AND AQUATIC BIOTA, 1987-91



U.S. GEOLOGICAL SURVEY  
Open-File Report 95-440



SURFACE-WATER-QUALITY ASSESSMENT OF THE YAKIMA RIVER  
BASIN IN WASHINGTON: SPATIAL AND TEMPORAL DISTRIBUTION  
OF TRACE ELEMENTS IN WATER, SEDIMENT, AND AQUATIC  
BIOTA, 1987-91

By Gregory J. Fuhrer, Daniel J. Cain, Stuart W. McKenzie, Joseph F. Rinella,  
J. Kent Crawford, Kenneth A. Skach, and Michelle I. Hornberger

*With a section on Geology*

By Marshall W. Gannett



U.S. GEOLOGICAL SURVEY

Open-File Report 95-440

Portland, Oregon

1996

U.S. DEPARTMENT OF THE INTERIOR  
BRUCE BABBITT, Secretary

U.S. GEOLOGICAL SURVEY  
Gordon P. Eaton, Director

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government

---

For additional information  
write to:

District Chief  
U.S. Geological Survey, WRD  
10615 S.E. Cherry Blossom Drive  
Portland, Oregon 97216

Copies of this report can  
be purchased from:

U.S. Geological Survey  
Branch of Information Services  
Box 25286, Denver Federal Center  
Denver, Colorado 80225



# FOREWORD

The mission of the U.S. Geological Survey (USGS) is to assess the quantity and quality of the earth resources of the Nation and to provide information that will assist resource managers and policymakers at Federal, State, and local levels in making sound decisions. Assessment of water-quality conditions and trends is an important part of this overall mission.

One of the greatest challenges faced by water-resources scientists is acquiring reliable information that will guide the use and protection of the Nation's water resources. That challenge is being addressed by Federal, State, interstate, and local water-resource agencies and by many academic institutions. These organizations are collecting water-quality data for a host of purposes that include: compliance with permits and water-supply standards; development of remediation plans for specific contamination problems; operational decisions on industrial, wastewater, or water-supply facilities; and research on factors that affect water quality. An additional need for water-quality information is to provide a basis on which regional- and national-level policy decisions can be based. Wise decisions must be based on sound information. As a society we need to know whether certain types of water-quality problems are isolated or ubiquitous, whether there are significant differences in conditions among regions, whether the conditions are changing over time, and why these conditions change from place to place and over time. The information can be used to help determine the efficacy of existing water-quality policies and to help analysts determine the need for and likely consequences of new policies.

To address these needs, the U.S. Congress appropriated funds in 1986 for the USGS to begin a pilot program in seven project areas to develop and refine the National Water-Quality Assessment (NAWQA) Program. In 1991, the USGS began full implementation of the program. The NAWQA Program builds upon an existing base of water-quality studies of the USGS, as well as those of other Federal, State, and local agencies. The objectives of the NAWQA Program are to:

- Describe current water-quality conditions for a large part of the Nation's freshwater streams, rivers, and aquifers.

- Describe how water quality is changing over time.

- Improve understanding of the primary natural and human factors that affect water-quality conditions.

This information will help support the development and evaluation of management, regulatory, and monitoring decisions by other Federal, State, and local agencies to protect, use, and enhance water resources.

The goals of the NAWQA Program are being achieved through ongoing and proposed investigations of 60 of the Nation's most important river basins and aquifer systems, which are referred to as study units. These study units are distributed throughout the Nation and cover a diversity of hydrogeologic settings. More than two-thirds of the Nation's freshwater use occurs within the 60 study units and more than two-thirds of the people served by public water-supply systems live within their boundaries.

National synthesis of data analysis, based on aggregation of comparable information obtained from the study units, is a major component of the program. This effort focuses on selected water-quality topics using nationally consistent information. Comparative studies will explain differences and similarities in observed water-quality conditions among study areas and will identify changes and trends and their causes. The first topics addressed by the national synthesis are pesticides, nutrients, volatile organic compounds, and aquatic biology. Discussions on these and other water-quality topics will be published in periodic summaries of the quality of the Nation's ground and surface water as the information becomes available.

This report is an element of the comprehensive body of information developed as part of the NAWQA Program. The program depends heavily on the advice, cooperation, and information from many Federal, State, interstate, Tribal, and local agencies and the public. The assistance and suggestions of all are greatly appreciated.

*Robert M. Hirsch*

Robert M. Hirsch  
Chief Hydrologist

# CONTENTS

Foreword .....	iii
Abstract .....	1
Introduction .....	3
Background .....	4
Purpose and Scope .....	4
Acknowledgments.....	5
The Yakima River Basin.....	5
Stream Reaches.....	7
Geologic Overview <i>By</i> Marshall W. Gannett .....	10
Land Use .....	11
Previous Studies .....	11
Approach and Methods .....	15
Water and Suspended Sediment.....	27
Streambed Sediment .....	28
Aquatic Biota .....	28
Comparison of Trace-Element Concentrations in Streambed Sediment, Water, and Fish Muscle to Water-Quality Guidelines .....	30
Streambed Sediment .....	30
Water.....	37
Ambient Stream Water .....	37
Aquatic Life.....	37
Human Health.....	51
Drinking Water .....	53
Regulations.....	53
Health Advisories .....	54
Fish Muscle.....	55
Spatial and Temporal Distribution of Trace Elements in the Aquatic Environment .....	55
Antimony .....	58
Arsenic and Lead .....	71
Barium.....	88
Cadmium.....	91
Chromium and Nickel.....	100
Cobalt.....	112
Copper.....	113
Mercury .....	130
Selenium .....	136
Silver.....	140
Zinc .....	144
Temporal Variations in Element Concentrations for Aquatic Biota Sampled at Sites in Common in 1989 and 1990 .....	150
Summary and Conclusions.....	151
Selected References .....	158
Supplemental Data Tables .....	163
Appendix .....	185
Glossary of Selected Terms .....	189

## FIGURES

1. Map showing the Yakima River Basin, Washington .....	6
2. Schematic diagram showing relative positions of selected tributaries, diversion canals, return flows, and stream-gaging stations in the Yakima River Basin, Washington .....	8
3. Elevation profile and distinctive hydrologic reaches of the Yakima River, Washington .....	9
4. Map showing land use and land cover by subbasin in the Yakima River Basin, Washington, 1981 .....	12
5. Map showing sampling-site locations for major and trace elements in sediment, water, and aquatic biota, Yakima River Basin, Washington, 1987–91 .....	16
6.–19. Graphs showing:	
6. Distribution of antimony concentrations in suspended sediment at fixed sites in the Yakima River Basin, Washington, 1987–90 .....	70
7. Streamflow and antimony concentrations in suspended sediment for selected time periods for the Yakima River at Cle Elum, Yakima River Basin, Washington, 1988–90 .....	71
8. Arsenic concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987–90 .....	72
9. Comparison of arsenic concentrations between agricultural and nonagricultural land uses or streambed sediment of lower-order streams in the Quaternary deposits and loess geologic unit, and in the nonmarine sedimentary rocks geologic unit, Yakima River Basin, Washington, 1987 .....	73
10. Distribution of arsenic concentrations in suspended sediment at fixed sites in the Yakima River Basin, Washington, 1987–90 .....	74
11. Concentrations of suspended sediment and arsenic in suspended sediment at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90 .....	75
12. Distribution of suspended sediment finer than 62 micrometers in diameter and suspended organic carbon concentrations at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90 .....	75
13. Percentage of suspended sediment finer than 62 micrometers in diameter and the concentration of arsenic in suspended sediment and the concentration of suspended organic carbon at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90 .....	76
14. Percentages of the annual load of arsenic in suspended sediment for selected time periods at Sulphur Creek Wasteway near Sunnyside, the Yakima River at Umtanum, and the Yakima River at Kiona, Yakima River Basin, Washington, 1989 .....	78
15. Distribution of arsenic concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90 .....	79
16. Arsenic concentrations in filtered-water samples at Sulphur Creek Wasteway near Sunnyside, the Yakima River at Euclid Bridge at river mile 55 near Grandview, and the Yakima River at Kiona, Yakima River Basin, Washington, 1989–90 .....	80
17. Arsenic concentrations in filtered-water samples and streamflow for selected time periods at the Yakima River at Euclid Bridge at river mile 55 near Grandview and the Yakima River at Kiona, Yakima River Basin, Washington, 1989–90 .....	81
18. Arsenic loads in filtered-water samples and in suspended-sediment samples for selected time periods at fixed sites, Yakima River Basin, Washington, 1989 .....	83
19. Distribution of lead concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	86
20. Maps showing the distribution of lead concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	88
21.–25. Graphs showing:	
21. Barium concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987 .....	91
22. Distribution of cadmium concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	92

23. Cadmium concentrations in suspended sediment and suspended-sediment concentrations, percentage of suspended sediment finer than 62 micrometers in diameter, and streamflow in the Naches River near North Yakima, Yakima River Basin, Washington, 1987–90 .....	93
24. Percentages of the annual load of cadmium in suspended sediment for the Yakima River at Umtanum, the Yakima River above Ahtanum Creek at Union Gap, the Yakima River at Euclid Bridge at river mile 55 near Grandview, and the Yakima River at Kiona, Yakima River Basin, Washington, 1989 .....	96
25. Cadmium loads in filtered-water samples and in suspended-sediment samples for selected time periods at the Yakima River at Umtanum, Yakima River Basin, Washington, 1989 .....	98
26. Maps showing the distribution of cadmium concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	100
27.–33. Graphs showing:	
27. Chromium concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987 .....	101
28. Distribution of chromium concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	102
29. Chromium concentrations in suspended sediment at the Yakima River at Umtanum and the Yakima River at Cle Elum, Yakima River Basin, Washington, 1987–90 .....	103
30. Chromium concentrations in suspended sediment at the Yakima River above Ahtanum Creek at Union Gap, Yakima River Basin, Washington, 1987–90 .....	104
31. Distribution of nickel concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	105
32. Nickel concentrations in suspended sediment at the Yakima River above Ahtanum Creek at Union Gap and at the Yakima River at Umtanum, Yakima River Basin, Washington, 1987–90 .....	106
33. Chromium and nickel concentrations in suspended sediment at the Yakima River at Umtanum, Yakima River Basin, Washington, 1987–90 .....	106
34. Maps showing the distribution of nickel concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	112
35. Maps showing the distribution of cobalt concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	113
36.–43. Graphs showing:	
36. Copper concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987 .....	114
37. Copper concentrations in suspended sediment at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90 .....	115
38. Copper concentrations in suspended sediment and the percentage of suspended sediment finer than 62 micrometers in diameter at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90 .....	115
39. Distribution of copper concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90 .....	118
40. Relation between copper concentrations in filtered-water samples and streamflow for selected time periods at the Yakima River at Cle Elum, Yakima River Basin, Washington, 1987–90 .....	119
41. Copper concentrations in filtered-water samples for the Yakima River at Cle Elum and the Yakima River at Umtanum, Yakima River Basin, Washington, 1989–90 .....	120
42. Percentages of the annual load of copper in filtered-water samples for selected time periods at selected fixed sites, Yakima River Basin, Washington, 1989 .....	122
43. Copper loads in filtered-water samples and in suspended-sediment samples for selected time periods at fixed sites, Yakima River Basin, Washington, 1989 .....	123
44. Maps showing the distribution of copper concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, caddisflies, and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	125

45. Maps showing the distribution of mercury concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, whole sculpins, and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	133
46. Graphs showing the relation between fish length and mercury concentrations of mountain-whitefish livers and largescale sucker livers, Yakima River Basin, Washington, 1990.....	134
47. Graphs showing mercury in mountain whitefish livers and rainbow trout livers compared to mercury in streambed sediment normalized to total organic carbon, Yakima River Basin, Washington, 1989–90 .....	135
48. Maps showing the distribution of selenium concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, whole sculpins, and streambed sediment, Yakima River Basin, Washington, 1987–90 .....	137
49.–56. Graphs showing:	
49. Selenium concentrations in whole sculpins at sites in Satus Creek and Ahtanum Creek, Yakima River Basin, Washington, 1990 .....	138
50. Relation between fish length and selenium concentrations of whole sculpins from the American River at Hells Crossing near Nile and Rattlesnake Creek above North Fork Rattlesnake Creek near Nile, Yakima River Basin, Washington, 1990.....	139
51. Distribution of silver concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	143
52. Silver concentrations in suspended sediment at the Yakima River above Ahtanum Creek at Union Gap and at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1989 .....	143
53. Suspended-sediment concentrations and the percentage of suspended sediment finer than 62 micrometers in diameter at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1989 .....	144
54. Zinc concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987 .....	145
55. Distribution of zinc concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	146
56. Zinc concentrations in suspended sediment and suspended-sediment concentrations at the Yakima River at Umtanum, Yakima River Basin, Washington, 1989 .....	150

## TABLES

1. Sampling-site locations and media sampled for major and trace elements, Yakima River Basin, Washington, 1987–91 .....	17
2. Major and trace elements analyzed in aquatic biota, water, and sediment, Yakima River Basin, 1987–91 .....	20
3. Types of samples analyzed for major and trace elements in the Yakima River Basin, Washington, 1987–91 .....	21
4. Sampling frequency for major and trace elements in aquatic biota, Yakima River Basin, Washington, 1989–91 .....	26
5. Elements analyzed in whole fish, fish liver, fish muscle, clams, and aquatic insects, Yakima River Basin, Washington, 1989–91 .....	29
6. Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90 .....	31
7. Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90.....	39
8. Summary of trace-element concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90 .....	45
9. Summary of estimated total-recoverable iron and manganese concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90.....	50

10. Concentrations of mercury in muscle of rainbow trout, largescale sucker, and mountain whitefish, Yakima River Basin, Washington, October 29–31, 1991 .....	56
11. Concentrations of mercury in fish muscle relative to Environmental Protection Agency screening values, Yakima River Basin, Washington, October 29–31, 1991 .....	57
12. Summary of major- and trace-element concentrations in streambed sediment, Yakima River Basin, Washington, 1987–91 .....	59
13. Summary of major- and trace-element concentrations in suspended sediment, Yakima River Basin, Washington, 1987–90 .....	61
14. Summary of major- and trace-element concentrations in filtered-water samples, Yakima River Basin, Washington, 1987–90 .....	62
15. Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91 .....	63
16. Estimated arsenic loads in suspended sediment at selected fixed sites, Yakima River Basin, Washington, 1987–90 .....	77
17. Estimated arsenic loads in filtered-water samples at selected fixed sites, Yakima River Basin, Washington, 1989–90 .....	82
18. Comparison of low and high arsenic concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	84
19. Comparison of low and high lead concentrations in sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	89
20. Estimated cadmium loads in suspended sediment at selected fixed sites, Yakima River Basin, Washington, 1987–90 .....	95
21. Frequency of occurrence of cadmium concentrations equalling or exceeding 0.2 micrograms per liter in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–89 .....	97
22. Estimated cadmium loads in filtered-water samples at selected fixed sites, Yakima River Basin, Washington, 1987–90 .....	98
23. Comparison of low, medium, and high chromium concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	107
24. Comparison of low and high nickel concentrations in sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	110
25. Estimated copper loads in suspended sediment at selected-fixed sites, Yakima River Basin, Washington, 1987–90 .....	117
26. Estimated copper loads in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90 .....	121
27. Comparison of low and high copper concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	127
28. Comparison of low and high mercury concentrations in streambed sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	131
29. Comparison of low and high selenium concentrations in streambed sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	141
30. Comparison of low and high zinc concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91 .....	147
31. Comparison of selected element concentrations in caddisflies sampled at sites in common in 1989 and 1990, Yakima River Basin, Washington .....	151
32. Comparison of selected element concentrations in Asiatic clams sampled at sites in common in 1989 and 1990, Yakima River Basin, Washington .....	151
33. Comparison of selected element concentrations in fish livers sampled at sites in common in 1989 and 1990, Yakima River Basin, Washington .....	152

#### SUPPLEMENTAL DATA TABLES

34. Comparison of selected element concentrations in streambed-sediment samples of the Yakima River Basin to streambed-sediment samples in the Lower Kansas River Basin, Upper Illinois River Basin, and the Kentucky River Basin .....	165
---	-----

35. Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 .....	166
36. Comparison of selected element concentrations in filtered-water samples from surface waters of the Yakima River Basin to surface waters in the United States .....	173
37. Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90 .....	174
38. Comparison of arsenic, mercury, and selenium concentrations in freshwater fish collected in 1984 for the National Contaminant Biomonitoring Program to concentrations in whole-body sculpin collected from the Yakima River Basin, Washington, 1990.....	181
39. Comparison of selected trace-element concentrations in <i>Corbicula</i> species collected from uncontaminated or minimally contaminated aquatic environments in other basins to Asiatic clams collected from the Yakima River Basin, Washington, 1990 .....	182
40. Comparison of selected trace-element concentrations in benthic insects collected from uncontaminated or minimally contaminated aquatic environments in other basins to benthic insects collected from the Yakima River Basin, Washington, 1990 .....	183

## CONVERSION FACTORS

Multiply	By	To obtain
acre	4,047	square meter (m <sup>2</sup> )
acre-foot (acre-ft)	1,233	cubic meter (m <sup>3</sup> )
cubic foot per second (ft <sup>3</sup> /s) (m <sup>3</sup> /s)	0.02832	cubic meter per second
inch	25.4	millimeter (mm)
foot (ft)	0.3048	meter (m)
mile (mi)	1.609	kilometer (km)
square mile (mi <sup>2</sup> )	2.590	square kilometer (km <sup>2</sup> )
cubic yard (yd <sup>3</sup> )	0.7646	cubic meter (m <sup>3</sup> )
cubic foot per second (ft <sup>3</sup> /s) (m <sup>3</sup> /s)	0.02832	cubic meter per second

Temperature in degrees Celsius (°C) as follows:

$$^{\circ}\text{C} = (^{\circ}\text{F} - 32) / 1.8.$$

**Sea level:** In this report “sea level” refers to the National Geodetic Vertical Datum of 1929 (NGVD of 1929)—a geodetic datum derived from a general adjustment of the first-order level nets of both the United States and Canada, called Mean Sea Level of 1929.





# Surface-Water-Quality Assessment of the Yakima River Basin in Washington: Spatial and Temporal Distribution of Trace Elements in Water, Sediment, and Aquatic Biota, 1987–91

By Gregory J. Fuhrer, Daniel J. Cain, Stuart W. McKenzie, Joseph F. Rinella, J. Kent Crawford, Kenneth A. Skach, and Michelle I. Hornberger

*With a section on Geology*  
By Marshall W. Gannett

## Abstract

In 1986, the U.S. Geological Survey implemented a pilot program to test and refine concepts for a National Water Quality Assessment (NAWQA) Program. Washington State's Yakima River Basin was selected as one of the Nation's four surface-water pilot studies.

One of the objectives of the NAWQA program, and the subject of this report, is to determine the occurrence and distribution of major and trace elements in sediment, water, and aquatic biota. Between 1987 and 1990, fine-grained (less than 62 micrometers in diameter) streambed sediment was collected from 448 sites in the Yakima River Basin. During 1-week periods in July and November 1987, trace elements were measured in filtered- (0.45 micrometer-pore size) water samples from 44 sites during steady-state flow conditions. Of the 44 sites, 7 were termed fixed sites, and were sampled monthly and during some storms for the period 1987–90. These sites provide the coverage necessary to describe temporal variations in element concentrations and loads. Fixed sites were sampled for trace elements in streambed sediment, water (filtered-water samples and unfiltered-water samples), suspended sediment, and aquatic biota. Aquatic biota from 34 sites were sampled for trace elements during

the period 1989–91. Aquatic biota include plant tissue (algae, curlyleaf pondweed, waterweed, and coontail), fish (rainbow trout, mountain whitefish, sculpin, brook trout, bridgelip sucker, largescale sucker, chiselmouth, carp, and cut-throat trout), clams (Asiatic), and aquatic insects (caddisflies, stoneflies, and mayflies).

The results of this study will provide multiple lines of evidence to:

- Define spatial and temporal variability for major and trace elements in several media;
- Identify sources and describe the transport and fate of trace elements;
- Evaluate the effects of trace elements on fish, benthic invertebrates, and algae.

The Yakima NAWQA data will be comparable with data collected at 59 other NAWQA basins throughout the United States so that the status of trace elements in the Nation's water can be addressed.

Most element enrichment in the Yakima River Basin results from natural geologic sources in the forested landscapes of the Kittitas and mid-Yakima Valley—primarily in the Cle Elum, Upper

Naches, Teanaway and Tieton Subbasins. These areas are geologic sources of antimony, arsenic, chromium, copper, mercury, nickel, selenium, and zinc. Concentrations of arsenic, chromium, and nickel in streambed sediment (as high as 45, 212, and 260  $\mu\text{g/g}$  [micrograms per gram], respectively) are nearly 4 to 13 times higher than their respective median concentrations in streambed sediment of agricultural land-use areas in the lower Yakima Valley. As a result of geologic sources, several of these elements, including arsenic, chromium, and nickel, leave chemical signatures that are measurable in streambed sediment and suspended sediment of higher-order streams, including the main stem. For example, the median concentration of chromium in streambed sediment in the Kittitas Valley (near the geologic source of chromium) is nearly twice that in the Yakima River at Kiona, located in the lower Yakima Valley. Some of the geologically derived elements, including chromium, nickel, and selenium, are measurable in aquatic biota of higher-order streams. In the Teanaway Subbasin alone, concentrations of chromium and nickel in benthic insects are, respectively, 4 to 52 times and 43 to 102 times higher than the minimum concentration in the Yakima River Basin.

Distributions of element concentrations that increase in areas affected by human activities include arsenic, cadmium, copper, lead, mercury, selenium, and zinc. These element concentrations frequently are high in the streambed sediment of Wide Hollow Creek, which drains urbanized and lightly industrialized lowlands as well as agricultural land. In Wide Hollow Creek, concentrations of lead in sediment (63  $\mu\text{g/g}$ ) and benthic insects (5.6  $\mu\text{g/g}$ ), respectively, are more than twice that in streambed sediment from local geologic sources and 15 times higher than that in caddisflies from a reference site not affected directly by human activity. In addition to suspected urban sources, lead may result from past applications of lead arsenate. Concentrations of lead in soils of apple orchards in the Yakima River Basin are as much as 60 times higher than the median concen-

tration of lead in streambed sediment from local geologic sources.

Suspended-arsenic concentrations in Sulphur Creek Wasteway, an agricultural drain, range from 4.9 to 20  $\mu\text{g/g}$  and are the highest in the Yakima River Basin. Arsenic may result from past applications of lead arsenate—soils of apple orchards in the Yakima River Basin have as much as 36 times more arsenic than the median concentration in streambed sediment from local geologic sources. During the irrigation season, the June contributions of suspended arsenic from Sulphur Creek Wasteway (2 lbs [pounds] per day) typically account for most of the suspended-arsenic load (measured 6 miles downstream) in the main stem. Between the Kittitas Valley and the mid-Yakima Valley, the annual suspended-arsenic loads during 1987–90 increased by as much as threefold. During the irrigation season, in particular, about 2.2 lbs of suspended arsenic per day enter the mid-Yakima Valley over a 9.4 river-mile reach that receives irrigation return flow from Moxee Subbasin and Wide Hollow Subbasin. This arsenic load represents about one-half the irrigation-season load in the Yakima River above Ahtanum Creek at Union Gap.

Concentrations of arsenic in filtered-water samples from Sulphur Creek Wasteway and in the main stem of the lower Yakima Valley are high (exceed the 90th percentile (3  $\mu\text{g/L}$ ) for the Yakima River Basin) in comparison to fixed sites (<1  $\mu\text{g/L}$ ) in the Kittitas Valley. These high concentrations reside in waters affected primarily by agricultural return flow. In addition to higher concentrations of arsenic in filtered-water samples from agriculturally affected parts of the Yakima River Basin, the load of arsenic in agricultural drains probably represents a large proportion of the arsenic load passing the Yakima River at Kiona, the terminus of the basin. For example, Sulphur Creek Wasteway has an annual streamflow representing only about 8 percent of the annual streamflow at the Kiona fixed site, yet accounts for nearly 20 percent of the filtered-arsenic load at Kiona. Comparisons, between loads determined from filtered-water

samples (an operational approximation of dissolved load) and loads determined from arsenic in suspended sediment, show that the annual dissolved-arsenic loads at fixed sites in the lower Yakima Valley are from four to nine times higher than their respective suspended loads.

Fish taxa provide the most comprehensive spatial coverage for arsenic, mercury, and selenium; however, no single fish taxon is widely distributed across the Yakima River Basin. The aquatic-insect taxon *Hydropsyche* spp. provides the most comprehensive spatial coverage of any single insect taxon. Concentrations of several elements, including cadmium, mercury, and selenium, in various taxa (except sculpin) were higher in the main stem of the lower Yakima Valley than in the Kittitas and mid-Yakima Valley. In mountain whitefish livers from the lower Yakima Valley, the concentrations of mercury (1.3 µg/g) and selenium (15 µg/g) are nearly twice that measured in mountain whitefish in the Kittitas Valley—similar patterns also were observed for largescale suckers. Compared with other studies of mercury in liver tissue of pike and mud fish, mercury concentrations in some mountain whitefish in the lower Yakima Valley are indicative of moderate enrichment. Sculpin were not sampled in the main stem; however, concentrations in some mid-Yakima Valley tributaries that drain the Northern Cascades are about 10 times higher (because of geologic sources of selenium) than in Satus Creek, a lower Yakima Valley tributary. Concentrations of selenium in sculpin in the Yakima River Basin (mean and 85th percentile values) also exceed those for sculpin in the National Contaminant Biomonitoring Program.

## INTRODUCTION

One of the most difficult issues facing water managers today is protecting the Nation's water resources while maintaining viable industrial and agricultural activities. Over the last several decades, concern about the water quality of our Nation's waterways has intensified. Federal, State, and local governments, as well as the public in general,

recognize the detrimental effects of major and especially trace elements that enter surface waters from point and nonpoint sources. When present in excessive concentrations, these elements may cause acute toxicity in the water column as well as accumulate in the tissues of aquatic organisms to toxic amounts, thereby altering the aquatic-community structure. Trace elements enter the aquatic environment from sources that include the weathering of rocks and human activities.

**Streambed sediment** is derived primarily from the physical and chemical weathering of rocks at the Earth's surface. Initially, rock is transformed into an aggregate of loose material by physical weathering processes, such as frost. The freshly disaggregated rock surfaces are then subject to chemical-weathering processes when water is present. Chemical-weathering processes (hydration and hydrolysis, oxidation and reduction, and the action of carbon dioxide) enable some of the physically disaggregated rocks to dissolve in water. Additionally, other rocks may be altered chemically by reacting with ions present in water or may remain totally unaltered by the action of water. The physically, and sometimes chemically, disaggregated rocks ultimately are transported to surface water by the action of rain, ice, wind, and animals (including man). Once in surface water, streambed sediment is known as **suspended sediment**. Trace-element concentrations in suspended sediment often are inversely related to grain size because they tend to sorb on sediment-particle surfaces (Horowitz, 1991; Forstner and Wittmann, 1979). Larger streamflows (often capable of suspending larger grain-sized sediment) normally carry lower concentrations of suspended-trace elements; conversely, higher streamflows (normally capable of suspending only fine-grained-sized sediment) normally carry higher concentrations of suspended-trace elements.

Once in the aquatic environment, streambed sediment becomes an important sink, or accumulator, of potentially toxic trace elements like arsenic, cadmium, copper, mercury, and zinc that enter streams from point and nonpoint sources. Many of these elements can be remobilized from streambed sediment to overlying **water**. Some of these trace elements are essential to **aquatic biota**. At the cellular level, elements like copper and zinc chemically bond to protein molecules and catalyze enzymatic reactions; additionally, without these trace elements, organisms

would fail to grow or complete their life cycles (Forstner and Wittmann, 1979). The same trace elements, however, can be toxic at the cellular level in aquatic biota when concentrations exceed those required for cellular metabolism and, additionally, can accumulate in aquatic biota and pass up the food chain. Thus, contaminants can disrupt the structure of the aquatic-biological community and through trophic transfers and biomagnification can pose a risk to consumers near the top of the food chain, including humans.

## Background

In 1986, Congress appropriated funds for the U.S. Geological Survey (USGS) to implement a pilot program to test and refine concepts for a National Water-Quality Assessment (NAWQA) program (Hirsch and others, 1988). The Yakima River Basin was one of four surface-water-pilot studies selected to refine NAWQA concepts (McKenzie and Rinella, 1987). The Yakima River Basin study included a planning phase in 1986, a data-collection phase from 1987 to 1990, and a report-writing phase that began in 1991. This report, one of several topical reports for the Yakima NAWQA study, presents the spatial and temporal distribution of major and trace elements in water, sediment, and aquatic biota.

The full-scale NAWQA program, which entails operation of 60 combined surface-water and ground-water study unit and covers about 60 to 70 percent of the Nations water use, began operation in 1991 (Leahy and others, 1990). The NAWQA program will provide results that are useful in understanding and managing water resources, and will address national water-quality issues. Specifically, the goals of the NAWQA program are to:

1. Provide a nationally consistent description of current water-quality conditions for a large part of the Nation's water resources;
2. Define long-term trends (or lack of trends) in water quality; and
3. Identify, describe, and explain, as possible, the major factors affecting observed water-quality conditions and trends.

The program is perennial and will involve a cyclic pattern of 3 years of active sampling followed by a period of low-level sampling. This cyclic pattern of sampling is sufficient to define long-term trends in water quality. The water-quality issues addressed in the program are broad, covering topics, such as eutrophication, pesticides, major and trace elements<sup>1</sup>, fecal indicator bacteria, suspended sediment, temperature, and aquatic biota.

## Purpose and Scope

Most trace elements in the aquatic environment result from natural sources; however, some trace elements result from anthropogenic sources or a mixture of natural and anthropogenic sources. With traditional environmental studies, it is often difficult to separate these sources and to assess the effects on the aquatic environment. Similarly, with traditional studies, it is often difficult to relate elements in water and sediment media to land use or geology and even more difficult to relate elements in water and sediment media to element concentrations in aquatic biota. Determining activities and conditions that affect the distribution and bioaccumulation of trace elements, therefore, entails a basinwide sampling of lower- and higher-order<sup>2</sup> streams. The purpose of this report is to describe, to the extent possible:

- (1) The occurrence and distribution of selected elements in water, sediment, and aquatic biota of the Yakima River Basin;

---

<sup>1</sup>Although definitions of the terms "major" and "trace" in reference to element concentration are not precise, substances typically occurring in concentrations of less than 1,000 parts per million (< 0.1 percent) are considered trace elements (Forstner and Wittmann, 1979, p. 5). Elements typically occurring in concentrations of greater than 1,000 parts per million are considered major elements. In this report major elements are reported in concentration units of percent and minor elements are reported in concentration units of micrograms per gram.

<sup>2</sup>In this report, lower-order streams are defined as first- or second-order tributaries and higher-order streams are defined as third order or larger tributaries—the largest being the main stem of the Yakima River. The smallest unbranched mapped (1:24,000 map scale) tributaries are first-order tributaries, streams receiving only first-order tributaries are second-order tributaries, larger streams receiving only first- and second-order tributaries are third-order and so on (Horton, 1945).

- (2) The temporal variation for element concentrations in filtered water and in suspended sediment at selected sites;
- (3) The suitability of surface water for preservation of aquatic life and protection of human health;
- (4) The major natural and anthropogenic sources in the Yakima River Basin that affect observed water-quality conditions; and
- (5) Implications of the assessment study with regard to future monitoring activities, assessment studies, and water management.

## Acknowledgments

The authors wish to acknowledge the aid and advice provided by members of the Yakima NAWQA Liaison Committee. During 1992, this committee included:

David W. Zimmer	Bureau of Reclamation
Don Schram	Bureau of Reclamation
Richard Albright	U.S. Environmental Protection Agency
Bill Garrigues	U.S. Forest Service
Terry W. Berkompas	Bureau of Indian Affairs
Kate Benkert	U.S. Fish and Wildlife Service
Wendell Hanigan	Yakima Indian Nation
Jannine Jennings	Yakima Indian Nation
Bob Barwin	Washington State Department of Ecology
Perry Harvester	Washington State Department of Fisheries
Brent Renfrow	Washington State Department of Wildlife
Glen Patrick	Washington State Department of Health
Dr. L. Clint Duncan	Washington Water Research Center, Central Washington State University
Skip Steinmetz	Yakima County Health Department
Ronald L. Van Gundy	Yakima River Basin Association of Irrigation Districts
Ray L. Wondercheck	U.S. Soil Conservation Service,

Elaine Taylor

Don Chaplin

Mike Tobin

Richard C. Bain Jr.

Yakima Valley  
Conference of  
Governments

County Extension  
Service


North Yakima  
Conservation District

Kittitas County  
Conservation District

Special thanks to (1) the Yakima Indian Nation for their cooperation in providing staff time for electrofishing, reviewing reports, and giving the USGS permission to collect water-quality samples from waterways in the Yakima Indian Nation; (2) U.S. Fish and Wildlife Service, Washington State Department of Fisheries, Washington State Department of Game for assistance in electrofishing; (3) Jean-Pierre Wilson (Heritage College, Toppenish, Washington) for assistance and cooperation in electrofishing; (4) Washington State Department of Wildlife for providing refrigeration and freezer space at the Naches Fish Hatchery; and (5) Shen Xian Chen (Institute of Water Conservancy and Hydroelectric Power Research, Chinese Academy of Sciences, Ministry of Water Resources and Electric Power, Beijing, China) for assistance in sampling.

## THE YAKIMA RIVER BASIN

The Yakima River flows 214.5 miles from the outlet of Keechelus Lake, in the central Washington Cascades, southeasterly to the Columbia River, draining an area of 6,155 mi<sup>2</sup> (square miles) (fig. 1); (Columbia Basin Inter-Agency Committee, 1964). The Yakima River Basin is one of the most intensively irrigated areas in the United States. The main stem and its largest tributary, the Naches River, have perennial streamflow with peak runoff occurring during peak snowmelt, usually in April and May. The Bureau of Reclamation's Yakima Project has seven irrigation divisions and provides water to irrigate almost one-half million acres. The project facilities include 6 storage dams, 416 miles of canals, 1,701 miles of laterals, 30 pumping plans, 145 miles of drains, 2 small power plants, and 74 miles of transmission lines (Bonneville Power Administration, 1985). Many of these waterways, most of which are natural streams, convey agricultural runoff and drainage, livestock wastes, and sewage-treatment-plant effluent to the



**Figure 1.** The Yakima River Basin, Washington

main stem. Surface-water diversions account for about 60 percent of the mean annual streamflow from the basin. Return flows, downstream of the city of Yakima, contribute as much as 80 to 90 percent of the flow in the lower main stem during irrigation season. A schematic diagram of selected inflows and outflows is shown in figure 2. Many of these inflows carry agricultural return flow.

The Yakima River Basin contains a variety of landforms, including the high peaks and deep valleys of the Cascade Range, the broad valleys and basalt ridges of the Columbia River Plateau, and the lowlands. Altitude in the basin ranges from about 340 ft (feet) at the mouth of the Yakima River, to 8,184 ft in the headwaters located in the Cascade Range. Glaciation has carved deep valleys in the high mountains, and currently, erosion by streams and small glaciers continues on the steep gradients. Mean annual precipitation in the basin ranges from 140 inches per year in the mountains to less than 10 inches per year in Kennewick, near the mouth of the basin. The eastern part of the basin consists of basalt flows, which form ridges trending northwest-southeast with valleys between them, and is more arid than the western part that is forested and mountainous.

It is estimated that before 1880, anadromous fish runs were more than 500,000 annually in the Yakima River Basin. By 1905, however, construction of irrigation projects, including reservoirs, had affected fish habitat and fish migrations. By 1920, anadromous fish runs had declined to 12,000 per year and in the 1980s declined to less than 4,000 adults of all species (Bonneville Power Administration, 1988). Currently, the major factors considered to affect fisheries in the basin are loss of fish habitat, mortality of smolts as they migrate down the Yakima and Columbia Rivers to the ocean, fishing pressures on the Columbia River and in the ocean, and poor water-quality conditions.


## Stream Reaches

The Yakima River descends from a water-surface altitude of 2,449 ft at the foot of Keechelus Dam to 340 ft at its mouth, downstream of Horn Rapids Dam near Richland (fig. 3). The headwater of Keechelus Lake and other tributaries flowing to the lake range in altitude from about 2,500 ft to more than 6,000 ft on the eastern slopes of the Cascade Mountains. Because of physical characteristics, the

Yakima River Basin can be divided into three distinctive river reaches (fig. 3). The upper reach drains the Kittitas Valley, is high gradient, and has an average streambed slope of 14 ft/mi (feet per mile) that extends 67.5 miles from the foot of Keechelus Dam to just upstream of Wilson Creek (river mile [RM] 147). In this reach, the river is shallow and the streambed is composed mostly of cobble and large gravel with some boulders, sand, and silt. Rocks are covered lightly with periphyton and slightly embedded in sediment.



The middle reach drains the mid-Yakima Valley and extends a distance of 39.5 miles from Wilson Creek to Union Gap (RM 107), also is high gradient, and has an average streambed slope of 11 ft/mi (fig. 3). Located on the Yakima River in the mid-Yakima Valley are the Roza (RM 127.9), Wapato (RM 106.6), and Sunnyside (RM 103.8) Dams that raise the hydraulic head of the river to divert water into irrigation canals. Upstream of the irrigation diversion dams, small stream segments of 1.0 RM or less are in backwater and accumulate fine-fraction sediment. Several waterways, including Wilson Creek and Moxee Drain, carry sediment-laden, irrigation return flow to the middle reach during the irrigation season (March 15 through October 15). Typical suspended-sediment concentrations during the irrigation season were about 100 mg/L (milligrams per liter) and 650 mg/L for Wilson Creek and Moxee Drain, respectively. Some of this sediment-laden water is, in turn, diverted into the Roza, Wapato, and Sunnyside irrigation canals. Sediment also is deposited in low-velocity backwaters of the middle reach. The fine-fraction sediment is transported farther down the main stem. Similar to the upper reach, the middle reach is shallow and the streambed is composed mostly of cobble and large gravel with some boulders, sand, and silt. In general, the rocks are covered lightly with periphyton and are free of rooted aquatic plants. Conversely, physical characteristics in reaches affected by backwaters of diversion dams are notably different. For example, the substrate in backwater of the Roza Dam is predominantly silt/clay with some organic matter and supports rooted aquatic plants.

The Naches River, a major tributary with 1,106 mi<sup>2</sup> (square miles) of drainage area, flows into the mid-Yakima Valley at RM 116.3. The Naches River is a high-gradient stream with an average streambed slope of 36 ft/mi. The Naches River ranges in altitude from 2,560 ft at the confluence of the Little Naches



**Figure 2.** Schematic diagram showing relative positions of selected tributaries, diversion canals, return flows, and stream-gaging stations in the Yakima River Basin, Washington.





**Figure 3.** Elevation profile and distinctive hydrologic reaches of the Yakima River, Washington.

and Bumping Rivers to 1,070 ft at its mouth (headwaters of the Naches River have water-surface altitudes as high as 6,000 ft). The river is shallow and the streambed is composed mostly of cobble and large gravel with some boulders, sand, and silt. Rocks are covered lightly to moderately with periphyton and are embedded slightly in sediment. The vegetative cover and thin soil mantle of the upper Naches Subbasin limit the amount of suspended sediment in the stream. Steep river gradients tend to keep most sediment suspended until the Naches River flows into the middle reach where velocities decrease in the backwaters of the Wapato and Sunnyside Diversion Dams.

The lower reach of the Yakima River drains the lower Yakima Valley and has an average streambed slope of 7 ft/mi and extends for 107.5 miles from the city of Union Gap to the mouth of the Yakima River (fig. 3). The streambed slope, streamflow, and average velocity are highly variable through this reach. During the irrigation season, streamflow in the main stem below the Wapato and Sunnyside Diversion Dams commonly is less than a few hundred cubic feet per second; streamflows remain low until irrigation water returns to the main stem by waterways between Parker (RM 104.6) and Mabton (RM 59.8). During the 1974–81 irrigation seasons, as much as 80 percent of the mean monthly irrigation return at Kiona was from tributaries carrying irrigation return flow between Parker and Kiona (Rinella and others, 1992). The upstream part of the lower reach of the Yakima River has a steep channel slope (12.8 ft/mi) which decreases midway through the reach (0.9 ft/mi) and results in a slow moving, meandering pool. The pool, located upstream of the Euclid Street Bridge (RM 55) near Grandview, is hydraulically characterized as a stilling basin behind a bedrock control. The pool is a depositional reach and contains predominantly silt/clay and some small gravel and organic matter. The substrate in the higher gradient sections, preceding and following the pool, is similar to that of the upper and middle reaches. Numerous rooted aquatic plants exist along the lower reach, especially in the vicinity of Horn Rapids Dam (RM 18.0).

## Geologic Overview

By Marshall W. Gannett, U.S. Geological Survey

The Yakima River Basin comprises parts of the Columbia Plateau and the Cascade geologic provinces. The largest segment of the basin, including

the entire southern and eastern parts, is in the Columbia Plateau, a province that consists primarily of basaltic lava flows with minor interbedded and overlying sediment. The western and northern margins of the basin are in the Cascade Range (fig. 1). The Cascade Mountains in the basin consist of a complex assemblage of volcanic, sedimentary, metamorphic, and intrusive rocks.

Approximately two-thirds of the Yakima River Basin is in the Columbia Plateau province. In the Yakima River Basin, this province is dominated by lavas of the Columbia River Basalt Group, which include the Grande Ronde, Wanapum, and Saddle Mountain Basalts (Walsh and others, 1987). The basalt occurs as multiple flows, each ranging in thickness from 10 to over 100 ft. Compressional forces in the Earth's crust during and after the emplacement of Columbia River Basalt Group lavas have warped and faulted the basalt into a series of east-northeast to east-southeast trending valleys and ridges. The ridges include the Horse Heaven Hills, the Rattlesnake Hills, and Toppenish, Ahtanum, Umtanum, Manastash, Naneum, and Yakima Ridges (fig. 1). Some of the lowlands between these basalt highlands have accumulated significant amounts of sediment. Major sediment accumulations, such as the Ellensburg Formation, are in structural lows of the Kittitas, Selah, Yakima, and Toppenish sedimentary basins according to Smith and others (1989).

Basalt flows of the Columbia River Basalt Group are overlain by, and locally interbedded with, sedimentary deposits. The major sedimentary unit in the Columbia Plateau province, in the Yakima River Basin, is the Ellensburg Formation, which consists chiefly of volcanoclastic material derived from the Cascade Range. Smith and others (1989) report that more than 1,000 ft of coarse-grained volcanoclastic sediment has accumulated over many parts of the Yakima River Basin.

A variety of unconsolidated surficial deposits of Quaternary age is present on the Columbia Plateau in the Yakima River Basin. These deposits include alluvial deposits along rivers and streams, alluvial terrace deposits, loess, and deposits resulting from catastrophic glacial outburst floods that inundated the lower part of the basin during the Pleistocene Epoch (Waite, 1985). These catastrophic flood deposits are present up to an altitude of about 1,000 ft in parts of the basin (Waite, 1985).

Approximately one-third of the Yakima River Basin is located in the Cascade Range geologic province. The Cascade Range province includes parts of the western and northern margins of the basin. The southern part of the Cascade Range in the basin, south of the Naches River, is dominated by Tertiary volcanic rocks, which include basalt and andesite flows, flow breccias, and related pyroclastic and volcanoclastic rocks (Walsh and others, 1987). Tertiary volcanic units are predominant in the middle part of the Tieton drainage, the upper part of the Rattlesnake Creek, most of the American River, Bumping River, and Crow Creek drainages. Older Jurassic to early Cretaceous marine sedimentary rocks are present in the Cascades Range south of the Naches River, most notably in the upper Tieton River drainage. These nonvolcanic rocks consist of sandstone and mudstone with lesser conglomerate (Walsh and others, 1987).

North of the Naches River, the Cascade Range province in the Yakima River Basin is dominated by Tertiary nonmarine sedimentary rocks, and pre-Tertiary metamorphic and intrusive rocks with small amounts of Tertiary volcanic rocks. Major sedimentary units in this area include the Eocene Roslyn and Swauk Formations.

The Roslyn Formation, which underlies a large part of the Teanaway River drainage, consists primarily of nonmarine sandstone with a smaller amount of conglomerate and thin coal seams (Tabor and others, 1982). The Swauk Formation, which is older than the Roslyn Formation, is located in the upper parts of the Teanaway River, Cle Elum River, and Swauk Creek drainages and consists primarily of nonmarine sandstone and a small amount of siltstone, shale, and conglomerate. The Swauk and Roslyn Formations are separated by the Teanaway Basalt, which consists primarily of basaltic flows, tuff, and breccia.

The upper parts of the south fork of Manastash Creek and the north and south forks of Taneum Creek drain areas where pre-Tertiary metamorphic rocks are found, including gneiss, schist, phyllite and amphibolite. These metamorphic rocks are surrounded and locally overlain by Tertiary volcanic rocks and nonmarine sedimentary rocks similar to the Swauk and Roslyn Formations.

In the far northern part of the Yakima River Basin, the uppermost sections of the Cle Elum River and the north fork of the Teanaway River drain an area underlain by ultramafic rocks adjacent to the Mount

Stuart batholith. These ultramafic rocks include serpentinite, serpentized peridotite, metaserpentinite, metaperidotite, diabase, and gabbro (Tabor and others, 1982).

Unconsolidated surficial deposits in the Cascade Province in the Yakima River Basin include alluvium along rivers and streams, alluvial fans, landslides, and glacial drift and outwash.


## Land Use

Major land-use activities in the Yakima River Basin include growing and harvesting timber, grazing on nonirrigated land, intensively irrigated agriculture, and urbanization (fig. 4). Intense water use for agriculture and cities makes these land-use categories of primary importance to water-quality issues. Population in the Yakima River Basin was about 250,000 in 1990 (Elaine Taylor, Yakima Valley Conference of Governments, Yakima, Washington, written commun., October 1992).

The forested northern and western areas in the Yakima River Basin lie in the Wenatchee and Snoqualmie National Forests, in the eastern slope of the Cascade Range. These forest lands are used for recreation, wildlife habitat, grazing, and timber harvesting. About one-fourth of this area is wilderness land, which has been designated for nonmotorized recreation. Rangelands are used for cattle grazing, wildlife habitat, and military training (at the Yakima Firing Center, northeast of the city of Yakima).

## Previous Studies

Water-quality investigations in the Yakima River Basin historically have focused on specific issues and objectives rather than a river-basin assessment of water quality. However, data from previous studies did provide useful historical perspectives when compared with present-study data. Most streambed-sediment chemistry studies in the Yakima River Basin were mineral-resource investigations and mostly cover the mountainous regions in the northern and southwestern Yakima River Basin. The following studies are noteworthy: the Goat Rocks Wilderness area (Church and others, 1983), the Alpine Lakes Wilderness area (U.S. Geological Survey, 1989), and the mineral-resource study of the headwaters of the Yakima and Naches Rivers (Moen, 1969).



**Figure 4.** Land use and land cover by subbasin in the Yakima River Basin, Washington, 1981 (U.S. Geological Survey, 1986).

The Alpine Lakes study (U.S. Geological Survey, 1989), the largest of the three previous studies, included collection of 4,702 streambed-sediment and rock samples between 1971–72. The Alpine Lakes report contains geology, mineral resources, geochemical data, and a detailed chronology of historical-mining activities. Streambed-sediment samples were screened by semiquantitative-spectrographic analysis for 30 elements; anomalous data were further analyzed by atomic absorption spectrophotometry. Streambed sediment collected in the northeastern part of the Yakima River Basin is enriched in concentrations of copper and zinc—concentrations varied in proportion to the distance from the contact between the Mount Stuart batholith and local-ultramafic rocks. Many of the anomalous data collected from the northwestern part of the basin—in the vicinity of Gold Creek—indicate the presence of hydrothermally altered and mineralized rock containing proportionally more lead and zinc and less copper. The second largest study of the headwaters of the Yakima and Naches Rivers (Moen, 1969) included collection and analysis of 182 streambed-sediment samples for copper, molybdenum, lead, and zinc, between 1965–68. Fries and Ryder (*in* Rinella and others, 1992) interpreted the data from these headwater studies and concluded that concentrations of copper (206 µg/g [micrograms per gram] in Easton Subbasin), and zinc (330 µg/g in the Upper Naches Subbasin and 120 µg/g in the Tieton Subbasin) exceed the 95-percent range shown for soils in the Western United States (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). The Goat Rocks study (Church and others, 1983) included collection and analysis of 43 streambed-sediment samples for 31 elements (using a direct current arc emission spectrographic method) in 1981. Fries and Ryder (*in* Rinella and others, 1992) interpreted data for the Goats Rocks study and concluded that the 90-percentile concentrations of copper (50 µg/g), vanadium (300 µg/g), and zinc (200 µg/g) in the Tieton Subbasin were anomalous.

Few studies of trace elements in filtered-water samples or in suspended sediment and water mixtures were made in the Yakima River Basin. For the most part, historical data were collected in the Yakima River at Kiona (RM 29) as part of the USGS and U.S. Environmental Protection Agency (EPA) monitoring programs from 1966–72. In 1973, samples were

collected from the Yakima River above Ahtanum Creek at Union Gap (RM 107) and in 1974 from the Yakima River at Kiona (RM 29.9). Both sites were operated as part of the USGS's National Stream Quality Accounting Network (NASQAN).

The Washington State Department of Ecology did a study in 1987 of low-flow conditions in Wide Hollow Creek which flows into the Yakima River (RM 107.4). This creek receives urban runoff from the city of Yakima, as well as agricultural return flow. Water and streambed sediment were analyzed for trace metals and trace-organic compounds. Water-quality measurements also were made, and fish and macroinvertebrate communities were surveyed, in order to relate the abundance and diversity of taxa to water-quality conditions. Fish abundance generally increased upstream, while macroinvertebrate distribution varied more with canopy shading than with water-quality conditions. Six trace elements, analyzed in unfiltered-water samples, were either below or only slightly above the lower limit of determination<sup>3</sup> (LLD). Streambed sediment was sampled at two sites and analyzed for 13 elements. The upstream site (RM 7.2) was subject to orchard and pasture effects, and the downstream site (RM 0.9) was subject to urban and livestock-pasturing activities. Arsenic, lead, mercury and zinc concentrations at the upstream site were respectively, 2.5 µg/g, 32 µg/g, 0.028 µg/g, and 92 µg/g. Concentrations of arsenic, lead, mercury, and zinc at the downstream sites were respectively, 0.9 µg/g, 11 µg/g, 0.036 µg/g, and 132.5 µg/g (Kendra, 1988). The concentrations of arsenic and lead were higher in close proximity to orchard and pasture effects. Mercury and zinc concentrations were higher in close proximity to urban and livestock-pasturing activities.

Filtered-water samples<sup>4</sup> analyzed for iron, manganese, lead, and mercury equaled or exceeded EPA National Primary and Secondary Drinking-Water Regulations at one or more sites in the Yakima River Basin, from 1953–85 water years; filtered

---

<sup>3</sup>The limit of determination is three times the standard deviation of a blank water sample added to the average of the blank.

<sup>4</sup>The term "filtered water" is operational defined as that portion of a water-suspended sediment sample that passes through a nominal 0.45 µm (micrometer) filter. Conversely, the term "unfiltered water" refers to a water sample that has not been filtered or centrifuged, nor in any way altered from the original matrix.

concentrations of cadmium, chromium, copper, lead, mercury, silver, and zinc equaled or exceeded EPA acute and (or) chronic standards established for the protection of freshwater-aquatic life (at four or more sites Rinella and others, 1992). Dissolved elements with the largest percentage of drinking-water exceedances (relative to the total number of determinations) were iron (8 percent), manganese (less than 1 percent), lead (3 percent), and mercury (less than 1 percent). Similarly, elements with the largest percentage of aquatic-life exceedances were lead (59 percent), mercury (41 percent), copper (24 percent), cadmium (11 percent), zinc (3 percent), chromium (1 percent), and silver (less than 1 percent). Whole-water and whole-water recoverable concentrations with the largest percentage of drinking-water exceedances (relative to the total number of determinations) are manganese (34 percent), iron (6 percent), and lead (2 percent). Similarly, elements with the largest percentages of acute and (or) chronic aquatic-life exceedances were lead (77 percent), mercury (50 percent), copper (46 percent), and nickel (less than 1 percent).

Few monitoring programs for trace elements in aquatic biota exist in the Yakima River Basin. Most programs emphasize the collection of fish tissue and cover only a small number of sites (Hopkins and others, 1985; Johnson and others, 1986; Lowe and others, 1985; Walsh and others, 1977; Schmitt and Brumbaugh, 1990). Washington State Department of Ecology did a study in the Yakima River Basin in 1985, targeting mercury in addition to trace-organic compounds. Fish, water, and streambed sediments were analyzed. The highest mercury concentrations in fish occurred in Northern squawfish. The maximum concentrations occurred at Wymer (RM 134–136), and the concentrations were 0.33 µg/g for whole fish and 0.45 µg/g for fish muscle, in wet weight. Mercury concentrations at downstream sites tended to increase in other species of fish. Whole fish, excluding northern squawfish, had mercury concentrations ranging from 0.02 µg/g for mountain whitefish at Cle Elum (RM 179–181) to 0.05 µg/g for mixed suckers at Buena (RM 93–95). Fish muscle, excluding northern squawfish, had mercury concentrations, in wet weight, ranging from 0.03 µg/g for mountain whitefish at Wymer to 0.13 µg/g for largescale sucker at Kiona (Johnson and others, 1986).

Since 1978, the Washington State Department of Ecology has analyzed fish tissue at selected sites in

Washington through the Basic Water Monitoring Program. In 1984, streambed-sediment sampling was initiated in addition to the fish sampling. Ten sites were sampled in 1984, two of which were on the Yakima River (below Moxee Drain at RM 107.6 and at Kiona at RM 29.9). Fish muscle was analyzed for pesticides and seven trace metals. Northern squawfish from the Yakima River below Moxee Drain had the highest concentration of mercury (0.78 µg/g, wet weight) found during the 1984 sampling. Mountain whitefish at the same site had 0.64 µg/g, wet weight, of mercury (Hopkins and others, 1985). These concentrations exceeded the National Academy of Sciences/National Academy of Engineering (NAS/NAE) recommendation for the protection of selected species of fish and predatory aquatic organisms. The recommendation states that the concentration of total mercury should not exceed a total body burden of 0.5 µg/g, wet weight, in any aquatic organism (National Academy of Sciences/National Academy of Engineering, 1972, p. 173).

The U.S. Fish and Wildlife Service has participated in the National Contaminant Biomonitoring Program (before 1984, called the National Pesticide Monitoring Program) since 1967. In this program, samples of bottom-feeding and predator fish were taken at over 100 sites throughout the United States and analyzed for potentially toxic elements and selected organochlorine compounds. One of these sites is in the Yakima River Basin (Yakima River at Granger). From 1971–73, concentrations of mercury in northern squawfish were below the U.S. Food and Drug Administration (FDA) guideline (1.0 µg/g, dry weight) for human consumption (Walsh and others, 1977). In 1978, a concentration of lead (1.4 µg/g, wet weight), which exceeded the 85th percentile value determined for various fish species in the United States, was found in a sample of white crappie at Granger (Lowe and others, 1985). In 1980 and 1984, however, none of the samples collected at Granger contained concentrations of lead that exceeded 85th percentile values, or concentrations of mercury that exceeded FDA guidelines (Lowe and others, 1985; and Schmitt and Brumbaugh, 1990). Different species were collected during different years, which might account for some of the variation reported among years (Schmitt and Brumbaugh, 1990). Additionally, some of the variation may have resulted from a change in the



location of the sampling site (Don Kane, U.S. Fish and Wildlife Service, oral commun., 1991). Samples were collected in 1984–88 from a pond located near the main stem of the Yakima River at Granger. In 1978 and 1980, however, it is unknown whether or not the main stem or the pond was sampled. The pond was believed to have been connected hydrologically to the main stem by a culvert; however, based on a site inspection in 1992, no evidence of a culvert remained.

## APPROACH AND METHODS

Trace-element concentrations were determined in media from the Yakima River Basin (fig. 5). These media included *streambed sediment*, *suspended sediment*, *water* (filtered and unfiltered), and *aquatic biota*. One or more of these media were measured at 57 sites in the Yakima River Basin, 1987–91, (table 1) for selected elements (table 2). Data pertaining to these media have been published for streambed sediment by Ryder and others (1992) and for sediment, water, and biota by Fuhrer, Fluter, and others (1994). To characterize temporal variations for trace-element concentrations and loads, trace-element data for suspended-sediment and water media were categorized according to season. The seasons defined for use in this report are *snowmelt*, *irrigation*, and *non-irrigation*. The irrigation season (March 15 to October 15) and snowmelt season, however, are not mutually exclusive from one another. Snowmelt generally is a major contributor to streamflow during April and May; consequently, the snowmelt portion of the irrigation season is considered separately. As a result, the snowmelt season is defined as April and May, the irrigation season as June through September, and the non-irrigation season as October through March.


Spatially, the most extensively sampled medium was streambed sediment. In 1987, as part of the Yakima NAWQA occurrence and distribution survey, streambed sediment was collected from 448 locations in the Yakima River Basin. These locations represented a variety of sampling sites that covered lower- and higher-order streams and included limited samplings of urban-storm drains and agricultural soils. Results from this streambed-sediment study were published earlier (Fuhrer, McKenzie, and others, 1994). Many of the sites measured for trace elements measured in aquatic biota, water, and suspended

sediment corresponded to streambed-sediment sites sampled as part of the occurrence and distribution survey in 1987. Additionally, some new streambed-sediment sites were sampled in 1989–90 to complement sites where trace elements were measured in aquatic biota. During 1987–90, one streambed-sediment sample was collected at each of the 32 sites at which an aquatic biota sample also was collected. Results from the 1987 occurrence and distribution survey are used in this report to describe the spatial distribution of trace elements in water, aquatic biota, and sediment from sites where aquatic biota were sampled.

Sampling frequency for filtered water varied for some sites. Forty-four sites were sampled at least once for filtered trace elements; a majority of these sites were sampled during synoptic surveys in July and (or) November 1987. Synoptic surveys were made over a short period of time (during steady-state streamflow conditions) and provided a broad spatial coverage for occurrence and distribution of trace-element concentrations in filtered water. Seven of the 44 sites also were sampled monthly and during hydrologic events (including snowmelt and winter rainstorms) for the period March 1987 to April 1990. These seven sites are referred to in this report as *fixed sites* 6, 19, 26, 32, 50, 52, and 56 (table 3 and fig. 5). The monthly and event-sampling frequency provides the temporal coverage necessary to describe seasonal variations for trace-element concentrations in filtered water.

Of the sites sampled, fixed sites were sampled with the greatest frequency. Generally, trace elements in all media were measured at these sites (fig. 5), with the exception of the Yakima River above Ahtanum Creek at Union Gap. (This site is not wadable; consequently, aquatic biota and streambed sediment samples were collected 2.6 miles downstream at the Yakima River at Parker.) Five of the seven fixed sites were on the main stem of the Yakima River; one site was located at the mouth of the Naches River—a major tributary; and the other site was located at the mouth of Sulphur Creek Wasteway—a major drain for carrying irrigation return flow and urban runoff (fig. 5). Fixed sites were sampled in a systematic-downstream order to simulate the movement of surface water passing through the Yakima River Basin. Operation of fixed sites in the basin is described by Fuhrer, Fluter, and others (1994).

Sampling frequency for suspended sediment and filtered water was identical for the period March 1987



**Figure 5.** Sampling-site locations for major and trace elements in sediment, water, and aquatic biota, Yakima River Basin, Washington, 1987–91 (site reference numbers are shown on map and are listed in Table 1; shaded areas of individual pie charts correspond to media that were sampled at the site).



**Table 1. Sampling-site locations and media sampled for major and trace elements, Yakima River Basin, Washington, 1987–91**

[S, streambed sediment; SS, suspended sediment; T, aquatic-biota tissue; F, filtered water; U, unfiltered water; RM, river mile; STP, sewage treatment plant; and DID, drainage irrigation district; --, not analyzed]

Site reference number	Site name	Station number <sup>1</sup>	Latitude	Longitude	Media				
					Sediment			Water	
					S	SS	T	F	U
1	Waptus River at mouth near Roslyn	12478100	472513	1210515	x	--	x	--	--
2	Cle Elum River above Cle Elum Lake near Roslyn	12478300	472119	1210622	--	--	--	x	--
3	Jungle Creek near mouth near Cle Elum	12479720	472030	1205159	x	--	x	x	--
4	North Fork Teanaway River below bridge at Dickey Creek Campground	12479750	471721	1205130	--	--	x	--	--
5	Teanaway River below Forks near Cle Elum	12480000	471448	1205136	x	x	x	--	--
6	Yakima River at Cle Elum	12479500	471135	1205655	x	x	x	x	x
7	Naneum Creek below High Creek near Ellensburg	12483750	471055	1202644	x	--	x	--	--
8	Taneum Creek at Taneum Meadow near Thorp	12481900	470647	1205201	x	--	x	--	--
9	Yakima River at Thorp Highway bridge at Ellensburg	12482800	470020	1203543	--	--	--	x	--
10	Little Naches River at mouth near Cliffdell	12487200	465920	1210555	x	--	x	x	--
11	American River near Nile	12488500	465839	1211005	--	--	--	x	--
12	South Fork Manastash Creek near Ellensburg	<sup>2</sup> 12483190	465818	1204832	x	--	x	x	--
13	American River at Hells Crossing near Nile	12488250	465804	1211545	x	--	x	--	--
14	Cherry Creek above Wipple Wasteway at Thrall	12484440	465544	1202948	x	--	x	--	--
15	Wilson Creek above Cherry Creek at Thrall	12484100	465535	1203001	--	--	--	x	--
16	Cherry Creek at Thrall	12484480	465534	1202951	--	--	x	x	--
17	Bumping River at Soda Springs Walkway near Nile	12488050	465527	1211250	--	--	--	x	--
18	Naches River at Cottonwood Campground near Cliffdell	12489050	465424	1210133	--	--	--	x	--
19	Yakima River at Umtanum	12484500	465146	1202844	x	x	x	x	x
20	Umtanum Creek near mouth at Umtanum	12484550	465127	1202946	x	--	x	x	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	<sup>3</sup> 12489150	464850	1205658	x	--	x	x	--

**Table 1.** Sampling-site locations and media sampled for major and trace elements, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Station number <sup>1</sup>	Latitude	Longitude	Media				
					Sediment			Water	
					S	SS	T	F	U
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	12489100	464834	1210408	x	--	x	--	--
23	Yakima River above canal diversion at RM 128 at Roza Dam	12484950	464503	1202752	--	--	--	x	--
24	Tieton River at mouth near Naches	12493100	464439	1204706	--	--	--	x	--
25	Yakima River above Selah Creek at Pomona	12485550	464232	1202825	--	--	--	x	--
26	Naches River near North Yakima	12499000	463742	1203110	x	x	x	x	x
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	12500437	463456	1203634	x	--	x	x	--
28	Tributary to Moxee Drain at Bell Road near Union Gap	12500415	463326	1202632	--	--	--	x	--
29	Wide Hollow Creek at old STP at Union Gap	<sup>4</sup> 12500442	463235	1202826	x	--	x	x	--
30	Moxee Drain at Thorp Road near Union Gap	12500430	463218	1202719	x	--	x	x	--
31	Ahtanum Creek at Union Gap	12502500	463210	1202820	x	--	x	--	--
32	Yakima River above Ahtanum Creek at Union Gap	12500450	463204	1202758	--	x	--	x	x
33	Yakima River at Parker	<sup>5</sup> 12503950	463022	1202707	x	--	x	x	--
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	12500900	462932	1205723	x	--	x	--	--
35	Unnamed drain at Progressive Road near Harrah	12507050	462509	1203545	--	--	--	x	--
36	Yakima River at RM 91 at Zillah	12505320	462407	1201654	--	--	--	x	--
37	East Toppenish Drain at Wilson Road near Toppenish	12505350	462204	1201500	--	--	--	x	--
38	Yakima River at Bridge Avenue at Granger	12505440	462016	1201148	--	--	--	x	--
39	Sub 35 Drain at Parton Road near Granger	12505410	462011	1201348	--	--	--	x	--
40	Granger Drain at mouth near Granger	12505460	462010	1201138	x	--	x	x	--
41	Marion Drain at Indian Church Road at Granger	12505510	461952	1201154	--	--	--	x	--
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	12507525	461858	1200913	x	--	x	--	--

**Table 1.** Sampling-site locations and media sampled for major and trace elements, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Station number <sup>1</sup>	Latitude	Longitude	Media				
					Sediment			Water	
					S	SS	T	F	U
43	Toppenish Creek at Indian Church Road near Granger	12507508	461852	1201153	x	--	x	x	--
44	Toppenish Creek near Fort Simcoe	12506000	461840	1204713	--	--	--	x	--
45	Yakima River at Van Geison Bridge near Richland	12511800	461750	1191956	--	--	--	x	--
46	DID 3 Drain below STP at Midvale Road at Sunnyside	12508838	461728	1200148	--	--	--	x	--
47	Satus Creek at Gage at Satus	12508620	461626	1200832	x	--	x	x	--
48	Yakima River at RM 72 above Satus Creek near Sunnyside	12507585	461611	1200530	x	--	x	--	--
49	Yakima River above Chandler Pump at RM 35.9 near Whitstran	12509900	461558	1193518	--	--	--	x	--
50	Yakima River at Kiona	12510500	461513	1192837	x	x	x	x	x
51	Yakima River below Satus Creek at RM 68 near Satus	12508625	461506	1200545	--	--	--	x	--
52	Sulphur Creek Wasteway near Sunnyside	12508850	461503	1200107	x	x	x	x	x
53	Satus Creek below Dry Creek near Toppenish	12508500	461500	1202240	x	--	x	x	--
54	Spring Creek at mouth at Whitstran	12509710	461400	1194038	x	--	x	--	--
55	Yakima River at Mabton	12508990	461353	1195954	--	--	--	x	--
56	Yakima River at Euclid Bridge at RM 55 near Grandview	12509050	461301	1195500	x	x	x	x	x
57	Satus Creek above Wilson-Charley Canyon near Toppenish	12507594	460100	1204054	x	--	x	--	--

<sup>1</sup>This number can be used for computer retrieval of suspended-sediment, filtered-water, and unfiltered-water chemical data from either the U.S. Geological Survey's WATER data STORage and RETrieval system (WATSTORE) or U.S. Environmental Protection Agency's STORage and RETrieval system (STORET).

<sup>2</sup>For filtered-water data retrieval, use station number 12483200.

<sup>3</sup>For filtered-water data retrieval, use station number 12489300.

<sup>4</sup>For filtered-water data retrieval, use station number 12500445.

<sup>5</sup>For filtered-water data retrieval, use station number 12505000.

**Table 2.** Major and trace elements analyzed in aquatic biota, water, and sediment, Yakima River Basin, Washington, 1987–91  
[--, not analyzed]

Element analyzed	Aquatic biota				Water		Sediment	
	Insects	Fish	Clams	Plants	Filtered	Unfiltered	Streambed	Suspended
<b>Major Elements</b>								
Aluminum	X	X	X	X	X	X	X	X
Calcium	--	X	X	--	--	--	X	X
Carbon, inorganic	--	--	--	--	--	--	X	--
Carbon, organic	--	--	--	--	X	--	--	X
Carbon, total	--	--	--	--	--	--	X	--
Iron	X	X	X	X	X	X	X	X
Magnesium	X	X	X	X	--	--	X	X
Phosphorous	--	--	--	--	--	--	X	X
Potassium	--	--	X	--	--	--	X	X
Sodium	--	--	X	--	--	--	X	X
Sulfur	--	--	--	--	--	--	X	--
Titanium	X	X	X	--	--	--	X	X
<b>Trace Elements</b>								
Antimony	--	--	--	--	X	--	X	X
Arsenic	X	X	X	X	X	--	X	X
Barium	X	X	X	X	X	X	X	--
Beryllium	X	X	X	X	X	X	X	X
Bismuth	--	--	--	--	--	--	X	--
Boron	X	X	X	X	X	X	X	--
Bromide	--	--	--	--	X	--	--	--
Cadmium	X	X	X	X	X	X	X	X
Cerium	--	--	--	--	--	--	X	--
Chromium	X	X	X	X	X	X	X	X
Cobalt	X	X	X	--	X	--	X	X
Copper	X	X	X	X	X	X	X	X
Cyanide	--	--	--	--	X	--	--	--
Europium	--	--	--	--	--	--	X	--
Gallium	--	--	--	--	--	--	X	--
Gold	--	--	--	--	--	--	X	--
Lanthanum	--	--	--	--	--	--	X	--
Lead	X	X	X	X	X	X	X	X
Lithium	--	--	--	--	X	--	X	--
Manganese	X	X	X	X	X	X	X	X
Mercury	X	X	X	X	X	X	X	--
Molybdenum	X	X	X	X	X	X	X	X
Neodymium	--	--	--	--	--	--	X	--
Nickel	X	X	X	X	X	X	X	X
Niobium	--	--	--	--	--	--	X	--
Scandium	--	--	--	--	--	--	X	--
Selenium	X	X	X	X	X	--	X	--
Silver	X	X	X	X	X	X	X	X
Strontium	X	X	X	X	X	--	X	--
Thallium	X	X	X	X	--	--	--	X
Thorium	--	--	--	--	--	--	X	--
Tin	--	--	--	--	--	--	X	--
Uranium	--	--	--	--	--	--	X	--
Vanadium	X	X	X	X	X	--	X	X
Ytterbium	--	--	--	--	--	--	X	--
Yttrium	--	--	--	--	--	--	X	--
Zinc	X	X	X	X	X	X	X	X

**Table 3. Types of samples analyzed for major and trace elements in the Yakima River Basin, Washington, 1987–91**

[OT, sample was collected at the site at least one time; M, sample was collected at a fixed site once a month from March 1987 to April 1990; RT, rainbow trout (*Oncorhynchus mykiss*); MW, mountain whitefish (*Prosopium williamsi*); BT, brook trout (*Salvelinus fontinalis*); BS, bridgeline sucker (*Catostomus columbianus*); LS, largescale sucker (*Catostomus macrocheilus*); CM, chiselmouth (*Acrocheilus alutaceus*); CT, cutthroat trout (*Oncorhynchus clarki*); CP, carp (*Cyprinus carpio*); SN, sculpin (*Cottus* spp.); CF, caddisflies (Trichoptera: Hydropsychidae); SF, stoneflies (Plecoptera: Perlidae; Perlodidae; Pteronarcidae); MF, mayflies (Ephemeroptera); WP, western pearlshell clam (Unionoida: Unionidae *Margaritifera falcata*); AC, Asiatic clam (Veneroida: Corbiculidae *Corbicula fluminea*); AG, algae (unidentified); CL, curlyleaf pondweed (*Potamogeton crispus*); WW, waterweed (*Elodea* sp.); CO, coontail (*Ceratophyllum demersum*); Q, sample was collected at the site approximately four times in a year and (or) during storm events; “–”, not analyzed; N.F., North Fork; R, River; RM, river mile; MS, Middle School; STP, sewage treatment plant; DID, drainage irrigation district]

Site reference number	Site name	Sediment		Aquatic-biota tissue <sup>1</sup>				Water		
		Streambed	Suspended	Fish			Insects	Clams	Plants	Unfiltered
				Liver	Whole fish	Muscle				
1	Wapatus River at mouth near Roslyn	OT	--	RT	--	--	CF	--	--	--
2	Cle Elum River above Cle Elum Lake near Roslyn	--	--	--	--	--	--	--	OT	--
3	Jungle Creek near mouth near Cle Elum	OT	--	RT	--	--	MF	--	AG	OT
4	N.F. Teanaway R below bridge at Dickey Creek Campground	--	--	--	--	--	CF,SF	--	--	--
5	Teanaway River below Forks near Cle Elum	OT	--	RT	--	--	--	--	--	OT
6	Yakima River at Cle Elum	OT	M	MW	SN	--	CF,SF	--	--	M
7	Naneum Creek below High Creek near Ellensburg	OT	--	BT	SN	--	CF,SF	--	--	--
8	Taneum Creek at Taneum Meadow near Thorp	OT	--	RT	SN	RT	CF,SF	--	--	--
9	Yakima River at Thorp Highway bridge at Ellensburg	--	--	--	--	--	--	--	OT	--
10	Little Naches River at mouth near Cliffdell	OT	--	--	--	--	CF,SF	--	--	OT
11	American River near Nile	--	--	--	--	--	--	--	--	OT
12	South Fork Manastash Creek near Ellensburg	OT	--	RT	SN	--	CF,SF	--	--	OT
13	American River at Hells Crossing near Nile	OT	--	--	RT,SN	--	CF,SF	--	--	--
14	Cherry Creek above Wipple Wasteway at Thrall	OT	--	MW	--	--	CF	--	--	--

**Table 3.** Types of samples analyzed for major and trace elements in the Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Sediment		Aquatic-biota tissue <sup>1</sup>						Water	
				Fish			Insects	Clams	Plants		
		Streambed	Sus- pended	Liver	Whole fish	Muscle				Whole organism	Soft tissue
15	Wilson Creek above Cherry Creek at Thrall	--	--	--	--	--	--	--	--	OT	--
16	Cherry Creek at Thrall	--	--	BS	--	--	CF	--	CL	OT	--
17	Bumping River at Soda Springs Walkway near Nile	--	--	--	--	--	--	--	--	OT	--
18	Naches River at Cottonwood Campground near Cliffdell	--	--	--	--	--	--	--	--	OT	--
19	Yakima River at Umtanum	OT	M	MW,RT	--	RT	CF,SF	WP	CL	M	Q
20	Umtanum Creek near mouth at Umtanum	OT	--	RT	SN	--	CF,SF	--	--	OT	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	OT	--	--	--	--	CF,SF	--	--	OT	--
22	Rattlesnake Creek above N.F. Rattlesnake Creek near Nile	OT	--	RT	SN	RT	CF,SF	--	--	--	--
23	Yakima River above canal diver- sion at RM 128 at Roza Dam	--	--	--	--	--	--	--	--	OT	--
24	Tieton River at mouth near Naches	--	--	--	--	--	--	--	--	OT	--
25	Yakima River above Selah Creek at Pomona	--	--	--	--	--	--	--	--	OT	--
26	Naches River near North Yakima	OT	M	LS,MW	--	--	CF,SF	--	--	M	Q
27	Wide Hollow Creek at West Val- ley MS near Ahtanum	OT	--	BS	--	--	CF	--	--	OT	--
28	Tributary to Moxee Drain at Bell Road near Union Gap	--	--	--	--	--	--	--	--	OT	--
29	Wide Hollow Creek at old STP at Union Gap	OT	--	CM,RT	--	--	CF	--	CL,WW	OT	--

**Table 3.** Types of samples analyzed for major and trace elements in the Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Sediment		Aquatic-biota tissue <sup>1</sup>						Water	
				Fish			Insects	Clams	Plants		
		Streambed	Suspended	Liver	Whole fish	Muscle	Whole organism	Soft tissue	Stems and leaves	Filtered <sup>2</sup>	Unfiltered
30	Moxee Drain at Thorp Road near Union Gap	OT	--	CM	--	--	--	--	--	OT	--
31	Ahtanum Creek at Union Gap	OT	--	--	SN	--	CF	--	--	--	--
32	Yakima River above Ahtanum Creek at Union Gap	--	M	--	--	--	--	--	--	M	Q
33	Yakima River at Parker	OT	--	LS,CP, MW	--	--	CF	--	CL	OT	--
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	OT	--	CT	SN	--	CF,SF	--	--	--	--
35	Unnamed drain at Progressive Road near Harrah	--	--	--	--	--	--	--	--	OT	--
36	Yakima River at RM 91 at Zillah	--	--	--	--	--	--	--	--	OT	--
37	East Toppenish Drain at Wilson Road near Toppenish	--	--	--	--	--	--	--	--	OT	--
38	Yakima River at Bridge Avenue at Granger	--	--	--	--	--	--	--	--	OT	--
39	Sub 35 Drain at Parton Road near Granger	--	--	--	--	--	--	--	--	OT	--
40	Granger Drain at mouth near Granger	OT	--	BS	--	--	CF	--	--	OT	--
41	Marion Drain at Indian Church Road at Granger	--	--	--	--	--	--	--	--	OT	--
42	Yakima R below Toppenish Creek at RM 79.6 near Granger	OT	--	LS,MW	--	--	CF	AC	--	--	--
43	Toppenish Creek at Indian Church Road near Granger	OT	--	LS	--	--	CF	--	WW	OT	--

**Table 3.** Types of samples analyzed for major and trace elements in the Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Sediment		Aquatic-biota tissue <sup>1</sup>						Water	
				Fish			Insects	Clams	Plants		
		Streambed	Sus- pended	Liver	Whole fish	Muscle	Whole organism	Soft tissue	Stems and leaves	Filtered <sup>2</sup>	Unfiltered
44	Toppenish Creek near Fort Simcoe	--	--	--	--	--	--	--	--	OT	--
45	Yakima River at Van Geison Bridge near Richland	--	--	--	--	--	--	--	--	OT	--
46	DID 3 Drain below STP at Midvale Road at Sunnyside	--	--	--	--	--	--	--	--	OT	--
47	Satus Creek at Gage at Satus	OT	--	LS	SN	--	CF	--	WW,CL	OT	--
48	Yakima River at RM 72 above Satus Creek near Sunnyside	OT	--	LS,CP	--	--	--	AC	CL	--	--
49	Yakima R above Chandler Pump at RM 35.9 near Whitstran	--	--	--	--	--	--	--	--	OT	--
50	Yakima River at Kiona	OT	M	MW,LS,C P	--	LS,MW	CF	AC	CL,CO	M	Q
51	Yakima River below Satus Creek at RM 68 near Satus	--	--	--	--	--	--	--	--	OT	--
52	Sulphur Creek Wasteway near Sunnyside	OT	M	LS	--	--	CF	--	--	M	Q
53	Satus Creek below Dry Creek near Toppenish	OT	--	--	SN	--	CF,SF	--	--	OT	--
54	Spring Creek at mouth at Whit- stran	OT	--	LS	--	--	CF	AC	WW	--	--
55	Yakima River at Mabton	--	--	--	--	--	--	--	--	OT	--
56	Yakima River at Euclid Bridge at RM 55 near Grandview	OT	M	LS	--	--	CF	AC	CL	M	Q
57	Satus Creek above Wilson- Charley Canyon near Toppenish	OT	--	RT	SN	--	CF,SF	--	--	--	--

<sup>1</sup> Tissue samples were collected during one or more of the following time periods: May and October–November of 1989, October–November of 1990, and October of 1991.

<sup>2</sup> Elements analyzed varied with frequency of sampling. For example, only cadmium, copper, mercury, and lead were analyzed monthly. All other elements were analyzed once or twice in a given year.



to April 1990. The seven fixed sites were sampled monthly and during hydrologic events, including snowmelt and winter rainstorms, and analyzed for major and trace elements. At the fixed sites, suspended sediment and water (filtered and unfiltered) were sampled simultaneously. The sampling frequency for unfiltered water, however, varied from that of the filtered portion and likewise, from that of the suspended sediment. Generally, unfiltered samples were collected quarterly during 1987 and not at all during 1988–90; however, some sites were sampled less frequently in 1987.

Of the seven fixed sites, the Yakima River above Ahtanum Creek at Union Gap (site 32) and the Yakima River at Kiona (site 50) were sampled most frequently. Additionally, unfiltered samples had been collected previously at these sites as part of the USGS's NASQAN program. Trace-element concentrations in unfiltered water were used to determine the suitability of surface water for supporting aquatic life, based on EPA's guidelines for freshwater aquatic life (U.S. Environmental Protection Agency, 1986, 1992c). These guidelines are based on a *total recoverable* method (trace elements removed by a mild-acid extraction of an unfiltered-water sample) in which complete digestion of all particulate matter is not achieved and the determination actually represents something less than 95 percent of the trace-element concentration being sought. This extraction method is similar to that used on unfiltered-water samples analyzed by the USGS's National Water Quality Laboratory (Fuhrer, Fluter, and others, 1994). The analytical method for determining trace-element concentrations in suspended sediment is termed a *total* method (elements removed by a harsh-acid extraction of a suspended sediment sample) in which complete digestion is nearly achieved and the determination actually represents something greater than 95 percent of the trace-element concentration being sought (Fuhrer, Fluter, and others, 1994). Trace-element concentrations determined using the total-recoverable method for unfiltered-water samples will be compared to trace-element concentrations from the total method for suspended-sediment and filtered-water samples. If the sum of the trace-element concentrations from the suspended-sediment sample and the filtered-water sample were similar to the trace-element concentrations in unfiltered water, then the sum of the trace-element concentrations in filtered water and suspended sediment, for the period March 1987 to April 1990, were

evaluated against EPA's guidelines for freshwater aquatic life.

Sampling frequency for the aquatic biota medium varied among sampling sites (table 4). A preliminary sampling for trace elements in the tissue of aquatic biota was made in May 1989 (sites 3, 19, 20, and 27) to test and refine collection and processing methods for aquatic biota. Aquatic biota were sampled at 34 sites from 1989–90—seven sites were located on the main stem. The aquatic biota medium included analyses of plant tissues, such as algae (unidentified specimens), curlyleaf pondweed (*Potamogeton crispus*), waterweed (*Elodea sp.*), and coontail (*Ceratophyllum demersum*); fish tissues, such as rainbow trout (*Oncorhynchus mykiss*), mountain whitefish (*Prosopium williamsoni*), sculpin (*Cottus spp.*), brook trout (*Salvelinus fontinalis*), bridgelip sucker (*Catostomus columbianus*), largescale sucker (*Catostomus macrocheilus*), chiselmouth (*Acrocheilus alutaceus*), carp (*Cyprinus carpio*), and cutthroat trout (*Oncorhynchus clarki*), Asiatic clam (Veneroida: Corbiculidae *Corbicula fluminea*), western pearlshell (*margaritifera falcata*); and aquatic insects, such as caddisflies (*Arctopsyche spp.*) (*Cheumatopsyche spp.*) (*Parapsyche spp.*) (*Hydropsyche ambilis*) (*Hydropsyche californica*) (*Hydropsyche cockerelli*) (*Hydropsyche occidentalis*), stoneflies (*Calineuria spp.*) (*Claassennia spp.*) (*Doroneuria spp.*) (*Hesperoperla spp.*) (*Isoperla spp.*) (*Megarcys spp.*) (*Perlinoles spp.*) (*Skwala spp.*) (*Pteronarcys spp.*), and mayflies (unidentified) [see table 3].

Streamflow was usually measured during each site visit for sites not equipped with continuous stage recorders. All fixed sites were equipped with stage recorders—stage was recorded every 30 minutes. Streamflow rating tables were developed and updated for fixed sites according to methods described by Buchanan and Somers (1969). At sites not equipped with stage recorders, either suspension- or wading-streamflow measurements were made according to methods described by Buchanan and Somers (1969).

Loads were determined for selected trace elements using the ESTIMATOR load computation model version 92.07 (Cohn and others, 1992). The ESTIMATOR model uses the minimum variance unbiased estimator for estimating constituent transport. Daily mean streamflow and trace-element concentrations for the period 1987–89 were used to calculate monthly and annual loads at the fixed sites in the Yakima River Basin. Equations used by

**Table 4.** Sampling frequency for major and trace elements in aquatic biota, Yakima River Basin, Washington, 1989–91  
[Nov., November; Oct., October; X, aquatic biota sampled in the month and year indicated]

Site reference number	Site name	May 1989	Nov. 1989	Nov. 1990	Oct. 1991
1	Waptus River at mouth near Roslyn		X		
3	Jungle Creek near mouth near Cle Elum	X			
4	North Fork Teanaway River below bridge at Dickey Creek Campground			X	
5	Teanaway River below Forks near Cle Elum		X		
6	Yakima River at Cle Elum		X	X	
7	Naneum Creek below High Creek near Ellensburg			X	
8	Taneum Creek at Taneum Meadow near Thorp			X	X
10	Little Naches River at mouth near Cliffdell			X	
12	South Fork Manastash Creek near Ellensburg			X	
13	American River at Hells Crossing near Nile		X	X	
14	Cherry Creek above Wipple Wasteway at Thrall			X	
16	Cherry Creek at Thrall		X	X	
19	Yakima River at Umtanum	X	X	X	X
20	Umtanum Creek near mouth at Umtanum	X	X	X	
21	Rattlesnake Creek above Little Rattlesnake near Nile			X	
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile		X	X	X
26	Naches River near North Yakima			X	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	X		X	
29	Wide Hollow Creek at old sewage treatment plant at Union Gap		X	X	
30	Moxee Drain at Thorp Road near Union Gap			X	
31	Ahtanum Creek at Union Gap			X	
33	Yakima River at Parker		X	X	
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico			X	
40	Granger Drain at mouth near Granger		X	X	
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger			X	
43	Toppenish Creek at Indian Church Road near Granger		X		
47	Satus Creek at Gage at Satus		X	X	
48	Yakima River at river mile 72 above Satus Creek near Sunnyside		X		
50	Yakima River at Kiona		X	X	X
52	Sulphur Creek Wasteway near Sunnyside		X	X	
53	Satus Creek below Dry Creek near Toppenish			X	
54	Spring Creek at mouth at Whitstran		X	X	
56	Yakima River at Euclid Bridge at river mile 55 near Grandview		X	X	
57	Satus Creek above Wilson-Charley Canyon near Toppenish		X	X	

ESTIMATOR for calculating specific constituent loads are found in the appendix.

## Water and Suspended Sediment

Water and suspended sediment were sampled using an equal-width-increment method which requires a sample volume proportional to the amount of flow at each of several equally spaced verticals in the stream cross section (Edwards and Glysson, 1988). A minimum of 10 verticals was sampled in the stream cross section. Additional verticals were sampled at sites where water in the cross section may not have been equally mixed because of nearby tributary inflow. Samples were collected using USGS depth-integrating samplers fitted with polyethylene, 3-liter bottles. The D-77 (depth-integrating sampler) is an epoxy-coated brass sampler that is operated by a cable and reel assembly; this sampler was used most of the time for sampling. The DH-81 (depth-integrating, hand-held sampler) was an optional sampler used when stream-flow conditions permitted crossing the stream by wading.

Equipment used for trace-element sampling was washed in liquinox and rinsed in distilled deionized water. Sample containers and equipment, with the exception of depth-integrating samplers, were rinsed again in 10 percent (by volume) hydrochloric acid. The depth-integrating sampler was not rinsed because the water sample did not come into contact with the sampler body. The acid rinse was followed by several rinses with distilled deionized water. All equipment was thoroughly rinsed in native stream water prior to sample collection.

Analytical methods have become more sensitive to contamination because of decreases in the analytical limit of determination. Field-processing methods and sampling equipment were evaluated by the USGS's Office of Water Quality to determine if they were a source of contamination to water samples analyzed at concentrations of parts per billion in micrograms per liter. Personnel from the Yakima NAWQA study participated in two field-processing and equipment blanks for cadmium, copper, lead, and zinc in filtered-water samples (Fuhrer, Fluter, and others, 1994). Cadmium, copper, lead, and zinc were among a listing of trace elements derived by the Office of Water Quality that pose a risk of contaminating water samples that are processed using certain water-quality samplers and

processing techniques (D.A. Rickert, U.S. Geological Survey, written commun., 1991). Blanks (assayed distilled-deionized water) run on the type of sampler and processing equipment used to collect water samples at fixed sites contained concentrations of cadmium, copper, lead, and nickel that were lower than the analytical limit of determination for the Yakima study, and were similar to background concentrations found in the blank water prior to the test (Fuhrer, Fluter, and others, 1994). A second study was made by the Office of Water Quality (D.A. Rickert, U.S. Geological Survey, written commun., 1992) to determine the potential for trace-element contamination from a variety of surface-water-quality samplers including the D-77 (depth-integrating) sampler. The Office of Water Quality concluded that the D-77 sampler with a polyethylene sample bottle is suitable for trace-element sampling at the parts per billion level. Furthermore, specific cleaning techniques were recommended for sampling and processing equipment which include a detergent wash, water rinse, acid soak/rinse, and several distilled water rinses (D.A. Rickert, U.S. Geological Survey, written commun., 1992). Cleaning techniques similar to these were used on sampling equipment used at fixed-location sampling sites (Fuhrer, Fluter, and others, 1994).

Filtered and unfiltered-water samples were preserved according to methods described in Fishman and Friedman (1989) and shipped on ice from the field to the USGS National Water Quality Laboratory in Arvada, Colorado. The samples were analyzed for major and trace elements according to methods described by Fishman and Friedman (1989). For fixed sites, a small suite of trace elements (cadmium, copper, and lead) was determined with an atomic absorption spectrometer in conjunction with a graphite furnace containing a graphite platform (AAGF). This method of analysis can detect low concentrations of trace elements that may adversely affect aquatic organisms. A larger suite of dissolved trace elements, which included aluminum, antimony, barium, beryllium, nickel, selenium, silver, and zinc, was determined simultaneously on a single sample by a direct-reading emission spectrometric method by using an induction-coupled argon plasma as an excitation source (ICP). This ICP method of analysis was used periodically for fixed sites during the first year of operation. Also, ICP was used during synoptic surveys in July and November 1987; however, this method was not ideally suited to evaluate element concentra-

tions in filtered water against guidelines for the protection of freshwater-aquatic life. Ambient concentrations for some elements (for example, cadmium, copper, lead, and zinc) with potential to adversely affect aquatic biota, were below the analytical limit of determination for the ICP. Method reporting levels, analytical method codes, and listings of trace-element suites are published in Fuhrer, Fluter, and others, 1994.

## Streambed Sediment

Streambed sediment was collected from five to seven points in each cross section of the stream channel; sampling was confined to surficial sediment that is usually in the upper one-half inch of the streambed. All streambed-sediment samples were wet sieved through a 0.062-mm (millimeter) mesh polyethylene sieve, using a minimum amount of stream-site water.

Samples were analyzed for 48 constituents—see Ryder and others (1992) for analytic methods, limits of determination, and decomposition methods. The majority of elements were determined by inductively coupled plasma-atomic emission spectrometry (ICP-AES). Decompositions of the sediment samples for elements measured by atomic absorption spectroscopy (AAS) and ICP-AES are total digestions—greater than 95 percent of the element is recovered during analysis. Total carbon was determined by combustion techniques. The decomposition used for boron and inorganic carbon are partial techniques. Quality assurance was addressed by including standard reference materials, random sample splits, and analytical sample splits (Sanzolone and Ryder, 1989; Fuhrer, McKenzie, and others, 1994, appendix A). Protocols used for sample handling procedures, sample preparation, analytical methods, instrument use, and laboratory procedures and quality control are described in Arbogast (1990).

## Aquatic Biota

Aquatic biota were collected over the period 1989–91. The 1990 data, however, are more comprehensive in the number of sites sampled and in using analytical methods with low or sensitive limits of determination. As a result, the 1990 data are used primarily to describe the spatial distribution of trace

elements; data from other years also are used to complement the 1990 data.

For sites where duplicate and triplicate measurements of a single species of aquatic biota were made, the mean trace-element concentration value was used; these values also were used for constructing a statistical-summary table. In contrast, multiple species of *Hydropsyche* were combined and reported under the single name of caddisfly (*Hydropsyche* spp.). The caddisfly (*Hydropsyche* spp.) and Asiatic clam (*corbicula fluminea*) tissue concentrations were normalized by calculating the logarithm of the trace-element concentrations, taking the mean of these values, and then calculating the antilogarithm of the aggregated data. Thus, in the statistical-summary table, each site is represented by a single trace-element concentration for each species analyzed. Because of the small number of stations summarized, only minimum, median, and maximum values were reported in the statistical-summary table for aquatic biota. The modified 1990 data were used to create maps of trace-element distribution using a geographical information system (GIS).

Several species of aquatic biota were collected in order to spatially depict the distribution of trace elements in the Yakima River Basin. In 1990, for example, the aquatic insect caddisfly (*Hydropsyche* spp.) provided the greatest coverage of a single taxon for spatially depicting trace-element concentrations. Arsenic, mercury, and selenium, however, were not analyzed in *hydropsyche*. For these trace elements, analyses of fish liver provided the greatest spatial coverage where several different fish species were sampled because no single species was present at all sites. Conversely, Asiatic clams (*corbicula fluminea*) were sampled only at five sites in the lower Yakima Valley.

Chemical data for aquatic biota in the Yakima River Basin were compared to data from other basins for *Corbicula* sp. and caddisfly (*Hydropsyche* spp.). Several caveats exist when making comparisons between basins. Only general comparisons were possible because differences may exist in species collected, size and (or) age of individuals in the sample, the tissues analyzed (muscle, gills, whole body, and so forth), analytical methods, and the way data were reported. Data for fish liver and whole fish were considered separately, because liver tissue is a principal site for accumulation of many trace elements, and concentrations of trace elements in liver

tissue are typically higher than in whole fish (Kaiser and others, 1979; Finley 1985; Ogle and others, 1988). To make comparisons with other studies, data originally reported as wet-weight concentrations were converted to dry-weight concentrations; unless otherwise stated, wet-weight data were converted using an assumed percent-moisture value of 75 percent.

In 1989 and 1990–91, a U.S. Fish and Wildlife Service (USFWS) contract laboratory and USGS laboratories prepared and analyzed samples from the Yakima River Basin (Fuhrer, Fluter, and others, 1994). The trace elements analyzed among years are shown in table 5. In 1989 and 1990, benthic insect samples

**Table 5.** Elements analyzed in whole fish, fish liver, fish muscle, clams, and aquatic insects, Yakima River Basin, Washington, 1989–91

Sample	Number of sites	Year collected	Elements analyzed
Whole fish (sculpin [ <i>Cottus</i> spp] only)	13	1990	Arsenic, mercury, and selenium
Fish liver (multiple species)	29	1989–90	See footnote <sup>1</sup> and <sup>2</sup>
Fish muscle (multiple species)	4	1991	Mercury
Clams ( <i>Corbicula fluminea</i> and <i>Margaritifera falcata</i> )	6	1989–90	See footnote <sup>1</sup> and <sup>2</sup>
Aquatic insects (multiple species)	31	1989–90	See footnote <sup>1</sup> and <sup>2</sup> Arsenic, mercury, and selenium not determined in 1990

<sup>1</sup>1989: silver, aluminium, arsenic, boron, barium, beryllium, cadmium, chromium, copper, iron, mercury, magnesium, manganese, molybdenum, nickel, lead, selenium, strontium, titanium, vanadium, and zinc.

<sup>2</sup>1990: silver, arsenic, calcium, cadmium, cobalt, chromium, copper, iron, mercury, magnesium, manganese, molybdenum, nickel, lead, selenium, titanium, vanadium, and zinc.

were not sorted, prior to chemical analysis, to the same taxonomic level. In other studies, differences in the taxonomic composition of samples have been shown to affect comparisons of element concentrations between years (Cain and others, 1992) and may be a factor in the Yakima River Basin. Also, differences in analytical methods of laboratories resulted in different

analytical limits of determination. Differences in sample preparation or analysis between years are explained in greater detail in subsequent sections, as relevant to results of this study. Approach and methods, analytical methods, limits of determination, and quality-assurance data have been described by Fuhrer, Fluter, and others (1994) as well as trace-element data for the different media covered in this report.

Elemental analyses of tissue from aquatic biota are a widely accepted approach to assess temporal (year to year) and spatial variations in environmental contaminants (Phillips, 1980); this approach has been incorporated into the NAWQA program (Crawford and Luoma, 1993). Additionally, multiple species were sampled to provide the spatial coverage necessary for the Yakima River Basin. In this manner, complementary data from other species may aid in describing spatial variations in environmental contaminants (Moore and others, 1991; Cain and others, 1992). Data from streambed sediment, suspended sediment, and water media also were examined to help identify natural and anthropogenic sources contributing to the spatial distribution of trace elements in aquatic biota. Ancillary data were collected for some species as correlative measures of bioaccumulation. Age, size, sex, and the presence of abnormalities, for example, were recorded as ancillary data for fish. These data, however, were of limited use. Because a limited number of species of fish existed at many sites, acquiring 10 individual livers for each composite sample sometimes required collection of a wide range of ages and sizes of fish.

Polar substances, such as cadmium, copper, lead, and zinc, concentrate in the livers of fish, but are regulated to small concentrations in muscle (Crawford and Luoma, 1993). For this reason, fish liver was selected as the target organ for the Yakima River Basin assessment of trace elements. Obtaining liver samples from a single species that was pervasive over the basin was problematic. Only sculpin were found in numbers suitable for a basinwide survey; however, sculpin were not collected from the main stem. Sculpin were analyzed for arsenic, mercury, and selenium. Whole bodies were analyzed instead of the liver because sculpin are small. Therefore, comparing absolute concentrations of trace elements between fish liver (from other species) and whole sculpin should be made with caution. In 1991, fish muscle was analyzed for mercury (sites 8, 19, 22, and 50 [table 5]) to

determine the risk to human health from consumption of local game fish.

## COMPARISON OF TRACE-ELEMENT CONCENTRATIONS IN STREAMBED SEDIMENT, WATER, AND FISH MUSCLE WITH WATER-QUALITY GUIDELINES

Bioavailability and toxicity vary with the form of a trace element (Jenne and Luoma, 1977). Aquatic organisms that feed on detritus are exposed to trace elements in solution and from the ingestion of sediment (Luoma, 1989). Trace elements associated with sediment generally are believed to be less bioavailable than trace elements dissolved in water (U.S. Environmental Protection Agency, 1992a). The toxicity to aquatic organisms from trace elements associated with sediment, however, is not necessarily zero. The concentration of total metals in sediment often are orders of magnitude higher than in water; small geochemical changes in the chemistry of sediment can affect solution chemistry greatly, and thus enhance bioavailability (Luoma, 1989). For example, trace elements associated with suspended sediment may dissolve in the chemical environment of the gill or the gut of an aquatic organism (U.S. Environmental Protection Agency, 1992a; Luoma, 1983).

### Streambed Sediment

As of 1993, the Washington State Department of Ecology and EPA Region X did not have sediment-quality guidelines. However, the Water Resources Branch of the Ontario Ministry of the Environment, Canada, developed Provincial Sediment Quality Guidelines (Persaud and others, 1993). These guidelines are compared with trace-element concentrations in streambed sediment of the Yakima River Basin.

The Provincial Sediment Quality Guidelines for trace elements are based on two levels of toxic effects. The first level, termed the lowest effect level, represents trace-element concentrations that can be tolerated by a majority of benthic organisms and that are comparable to the low effect levels determined through a review of sediment-toxicity bioassays by the National Oceanic and Atmospheric Administration (Persaud and others, 1993). The second level, termed

the severe effect level, represents trace-element concentrations at which “pronounced disturbances of the sediment-dwelling community can be expected” and has been deemed a concentration that would be detrimental to the majority of benthic species (Persaud and others, 1993).

The lowest effect level and the severe effect level were derived from a survey of trace-element concentrations in bulk sediment in addition to a survey of *in-situ* benthic abundance. Because concentrations of trace elements in the Yakima River Basin were determined from sediment finer than 62  $\mu\text{m}$  (micrometers) in diameter (rather than bulk sediment) and because the 62- $\mu\text{m}$  size fraction tends to have higher element concentrations than bulk sediment, element concentrations in the Yakima River Basin may in some instances exceed the Provincial Sediment Quality Guidelines, solely due to differences in the quantity of fine-grained-size sediment. Consequently, instances where trace-element concentrations in the Yakima River Basin that exceed Provincial Sediment Quality Guidelines should be used only as an indication of potential sediment quality problems. The Provincial Sediment Quality Guidelines are derived from a large number of data sets—each data set represents a minimum of 10 sites where at least 10 species of interest reside. In addition, the sum total of the data sets is assumed to represent a complete range of trace-element concentrations for benthic species of interest. The 90th-percentile concentration is determined for the trace-element concentrations in each data set. These concentrations were then pooled and plotted as a frequency distribution. The concentration corresponding to the 5th percentile of the frequency distribution was termed the “lowest effect level” and, similarly, the concentration corresponding to the 95th percentile of the frequency distribution was termed the “severe effect level.” Application of the Provincial Sediment Quality guidelines was limited, however, because the guidelines were not based on a toxicological response and do not directly infer a cause and effect relation between the trace-element content of streambed sediment and the uptake of trace elements by aquatic organisms.

Streambed sediment was compared with the Provincial Sediment Quality Guidelines at 32 locations in the Yakima River Basin (table 6). Trace-element concentrations at a number of sites exceeded the severe effect level. Trace elements that exceeded the

**Table 6. Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90**

[The Provincial Sediment-Quality Guidelines were developed by the Water Resources Branch of the Ontario Ministry of the Environment, Canada, for use in evaluating sediments throughout the province of Ontario (Persaud and others, 1991); the guidelines define two levels of toxic effects which are based on protecting benthic organisms from the chronic, long-term effects of selected elements; these levels are: (1) lowest effect level, and (2) severe effect level; a lowest effect level is a level of contamination that can be tolerated by a majority of benthic organisms, and a severe effect level is a level of contamination that would be detrimental to a majority of benthic species; note, some sites have been sampled in duplicate or triplicate as part of the quality-assurance program that has been described along with results by Fuhrer and others (1993); for replicate sites, only the mean element concentration is reported here; percentages were calculated using all measurements (censored and detected) for all sites sampled; concentrations are provided in units of micrograms per gram, ( $\mu\text{g/g}$ ), except for iron, which is reported in units of percent; RM, river mile]

Site reference number	Site name	Streambed-sediment concentration	Samples exceeding guidelines	
			Lowest effect level	Severe effect level
Arsenic, guideline: lowest effect level: 6 µg/g; severe effect level: 33 µg/g				
1	Waptus River at mouth near Roslyn	45	X	X
3	Jungle Creek near mouth near Cle Elum	29	X	
6	Yakima River at Cle Elum	7.9	X	
13	American River at Hells Crossing near Nile	13	X	
21	Rattlesnake Creek above Little Rattlesnake near Nile	7.8	X	
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	9.5	X	
56	Yakima River at Euclid Bridge at RM 55 near Grandview	6.6	X	
Percentage of samples that exceed guidelines			22	3
Cadmium, guideline: lowest effect level: 0.6 mg/g; severe effect level: 10 mg/g				
13	American River at Hells Crossing near Nile	.8	X	
Percentage of samples that exceed guidelines			3	0
Chromium, guideline: lowest effect level: 26 µg/g; severe effect level: 110 µg/g				
1	Waptus River at mouth near Roslyn	61	X	
3	Jungle Creek near mouth near Cle Elum	93	X	
5	Teanaway River below Forks near Cle Elum	210	X	X
6	Yakima River at Cle Elum	210	X	X
7	Naneum Creek below High Creek near Ellensburg	58	X	
8	Taneum Creek at Taneum Meadow near Thorp	170	X	X
10	Little Naches River at mouth near Cliffdell	50	X	
12	South Fork Manastash Creek near Ellensburg	130	X	X
14	Cherry Creek above Wipple Wasteway at Thrall	52	X	
19	Yakima River at Umtanum	64	X	
20	Umtanum Creek near mouth at Umtanum	50	X	
21	Rattlesnake Creek above Little Rattlesnake near Nile	49	X	
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	62	X	
26	Naches River near North Yakima	90	X	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	42	X	
29	Wide Hollow Creek at old STP at Union Gap	67	X	



**Table 6.** Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Streambed- sediment concentration	Samples exceeding guidelines	
			Lowest effect level	Severe effect level
Chromium, guideline: lowest effect level: 26 µg/g; severe effect level: 110 µg/g—Continued				
30	Moxee Drain at Thorp Road near Union Gap	64	X	
31	Ahtanum Creek at Union Gap	56	X	
33	Yakima River at Parker	64	X	
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	33	X	
40	Granger Drain at mouth near Granger	60	X	
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	73	X	
43	Toppenish Creek at Indian Church Road near Granger	75	X	
47	Satus Creek at gage at Satus	54	X	
48	Yakima River at RM 72 above Satus Creek near Sunnyside	64	X	
50	Yakima River at Kiona	61	X	
52	Sulphur Creek Wasteway near Sunnyside	80	X	
53	Satus Creek below Dry Creek near Toppenish	48	X	
54	Spring Creek at mouth at Whitstran	62	X	
56	Yakima River at Euclid Bridge at RM 55 near Grandview	56	X	
57	Satus Creek above Wilson-Charley Canyon near Toppenish	53	X	
Percentage of samples that exceed guidelines			97	12
Copper, guideline: lowest effect level: 16 µg/g; severe effect level: 110 µg/g				
1	Waptus River at mouth near Roslyn	20	X	
3	Jungle Creek near mouth near Cle Elum	36	X	
5	Teanaway River below Forks near Cle Elum	32	X	
6	Yakima River at Cle Elum	43	X	
7	Naneum Creek below High Creek near Ellensburg	24	X	
8	Taneum Creek at Taneum Meadow near Thorp	47	X	
10	Little Naches River at mouth near Cliffdell	48	X	
12	South Fork Manastash Creek near Ellensburg	44	X	
13	American River at Hells Crossing near Nile	79	X	
14	Cherry Creek above Wipple Wasteway at Thrall	22	X	
19	Yakima River at Umtanum	28	X	
20	Umtanum Creek near mouth at Umtanum	30	X	
21	Rattlesnake Creek above Little Rattlesnake near Nile	39	X	
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	43	X	
26	Naches River near North Yakima	61	X	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	93	X	
29	Wide Hollow Creek at old STP at Union Gap	42	X	
30	Moxee Drain at Thorp Road near Union Gap	25	X	
31	Ahtanum Creek at Union Gap	25	X	



**Table 6.** Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Streambed- sediment concentration	Samples exceeding guidelines	
			Lowest effect level	Severe effect level
Copper, guideline: lowest effect level: 16 µg/g; severe effect level: 110 µg/g—Continued				
33	Yakima River at Parker	29	X	
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	29	X	
40	Granger Drain at mouth near Granger	21	X	
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	31	X	
43	Toppenish Creek at Indian Church Road near Granger	32	X	
47	Satus Creek at gage at Satus	29	X	
48	Yakima River at RM 72 above Satus Creek near Sunnyside	31	X	
50	Yakima River at Kiona	29	X	
52	Sulphur Creek Wasteway near Sunnyside	26	X	
53	Satus Creek below Dry Creek near Toppenish	32	X	
54	Spring Creek at mouth at Whitstran	17	X	
56	Yakima River at Euclid Bridge at RM 55 near Grandview	27	X	
57	Satus Creek above Wilson-Charley Canyon near Toppenish	25	X	
Percentage of samples that exceed guidelines			100	0
Iron, guideline: lowest effect level: 2 percent; severe effect level: 4 percent				
1	Waptus River at mouth near Roslyn	3.9	X	
3	Jungle Creek near mouth near Cle Elum	6.0	X	X
5	Teanaway River below Forks near Cle Elum	5.5	X	X
6	Yakima River at Cle Elum	5.1	X	X
7	Naneum Creek below High Creek near Ellensburg	4.9	X	X
8	Taneum Creek at Taneum Meadow near Thorp	5.0	X	X
10	Little Naches River at mouth near Cliffdell	6.0	X	X
12	South Fork Manastash Creek near Ellensburg	6.3	X	X
13	American River at Hells Crossing near Nile	4.9	X	X
14	Cherry Creek above Wipple Wasteway at Thrall	4.2	X	X
19	Yakima River at Umtanum	4.3	X	X
20	Umtanum Creek near mouth at Umtanum	6.7	X	X
21	Rattlesnake Creek above Little Rattlesnake near Nile	5.6	X	X
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	5.2	X	X
26	Naches River near North Yakima	4.9	X	X
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	4.5	X	X
29	Wide Hollow Creek at old STP at Union Gap	5.2	X	X
30	Moxee Drain at Thorp Road near Union Gap	4.3	X	X
31	Ahtanum Creek at Union Gap	5.7	X	X
33	Yakima River at Parker	4.6	X	X

**Table 6.** Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Streambed- sediment concentration	Samples exceeding guidelines	
			Lowest effect level	Severe effect level
Iron, guideline: lowest effect level: 2 percent; severe effect level: 4 percent— Continued				
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	6.9	X	X
40	Granger Drain at mouth near Granger	5.5	X	X
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	5.1	X	X
43	Toppenish Creek at Indian Church Road near Granger	5.1	X	X
47	Satus Creek at gage at Satus	5.9	X	X
48	Yakima River at RM 72 above Satus Creek near Sunnyside	4.9	X	X
50	Yakima River at Kiona	5.1	X	X
52	Sulphur Creek Wasteway near Sunnyside	5.1	X	X
53	Satus Creek below Dry Creek near Toppenish	5.2	X	X
54	Spring Creek at mouth at Whitstran	5.5	X	X
56	Yakima River at Euclid Bridge at RM 55 near Grandview	5.0	X	X
57	Satus Creek above Wilson-Charley Canyon near Toppenish	7.3	X	X
Percentage of samples that exceed guidelines			100	97
Lead, guideline: lowest effect level: 31 µg/g; severe effect level: 250 µg/g				
26	Naches River near North Yakima	36	X	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	63	X	
29	Wide Hollow Creek at old STP at Union Gap	48	X	
Percentage of samples that exceed guidelines			9	0
Manganese, guideline: lowest effect level: 460 µg/g; severe effect level: 1,100 µg/g				
1	Waptus River at mouth near Roslyn	1,400	X	X
3	Jungle Creek near mouth near Cle Elum near Cle Elum	910	X	
5	Teanaway River below Forks near Cle Elum near Cle Elum	1,100	X	X
6	Yakima River at Cle Elum	1,100	X	X
7	Naneum Creek below High Creek near Ellensburg	1,000	X	
8	Taneum Creek at Taneum Meadow near Thorp	820	X	
10	Little Naches River at mouth near Cliffdell	1,400	X	X
12	South Fork Manastash Creek near Ellensburg	1,100	X	X
13	American River at Hells Crossing near Nile	1,100	X	X
14	Cherry Creek above Wipple Wasteway at Thrall	830	X	
19	Yakima River at Umtanum	1,700	X	X
20	Umtanum Creek near mouth at Umtanum	1,100	X	X
21	Rattlesnake Creek above Little Rattlesnake near Nile	1,000	X	
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	1,200	X	X
26	Naches River near North Yakima	1,100	X	X
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	550	X	
29	Wide Hollow Creek at old STP at Union Gap	770	X	

**Table 6.** Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Streambed- sediment concentration	Samples exceeding guidelines	
			Lowest effect level	Severe effect level
Manganese, guideline: lowest effect level: 460 µg/g; severe effect level: 1,100 µg/g—Continued				
30	Moxee Drain at Thorp Road near Union Gap	740	X	
31	Ahtanum Creek at Union Gap	960	X	
33	Yakima River at Parker	780	X	
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	1,500	X	X
40	Granger Drain at mouth near Granger	1,000	X	
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	750	X	
43	Toppenish Creek at Indian Church Road near Granger	960	X	
47	Satus Creek at gage at Satus	1,100	X	X
48	Yakima River at RM 72 above Satus Creek near Sunnyside	1,000	X	
50	Yakima River at Kiona	1,500	X	X
52	Sulphur Creek Wasteway near Sunnyside	990	X	
53	Satus Creek below Dry Creek near Toppenish	820	X	
54	Spring Creek at mouth at Whitstran	930	X	
56	Yakima River at Euclid Bridge at RM 55 near Grandview	1,000	X	
57	Satus Creek above Wilson-Charley Canyon near Toppenish	1,400	X	X
Percentage of samples that exceed guidelines			100	44
Mercury, guideline: lowest effect level: 0.2 µg/g; severe effect level: 2 µg/g				
3	Jungle Creek near mouth near Cle Elum	.56	X	
5	Teanaway River below Forks near Cle Elum	.20	X	
8	Taneum Creek at Taneum Meadow near Thorp	.40	X	
21	Rattlesnake Creek above Little Rattlesnake near Nile	.24	X	
26	Naches River near North Yakima	.27	X	
29	Wide Hollow Creek at old STP at Union Gap	.26	X	
33	Yakima River at Parker	.26	X	
Percentage of samples that exceed guidelines			22	0
Nickel, guideline: lowest effect level: 16 µg/g; severe effect level: 75 µg/g				
1	Waptus River at mouth near Roslyn	29	X	
3	Jungle Creek near mouth near Cle Elum	43	X	
5	Teanaway River below Forks near Cle Elum	260	X	X
6	Yakima River at Cle Elum	150	X	X
7	Naneum Creek below High Creek near Ellensburg	22	X	
8	Taneum Creek at Taneum Meadow near Thorp	78	X	X
10	Little Naches River at mouth near Cliffdell	20	X	
12	South Fork Manastash Creek near Ellensburg	61	X	
14	Cherry Creek above Wipple Wasteway at Thrall	21	X	
19	Yakima River at Umtanum	48	X	

**Table 6.** Summary of major- and trace-element concentrations in streambed sediment that exceeded Provincial Sediment-Quality Guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Streambed- sediment concentration	Samples exceeding guidelines	
			Lowest effect level	Severe effect level
Nickel, guideline: lowest effect level: 16 µg/g; severe effect level: 75 µg/g—Continued				
20	Umtanum Creek near mouth at Umtanum at Umtanum	19	X	
21	Rattlesnake Creek above Little Rattlesnake near Nile	16	X	
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	27	X	
26	Naches River near North Yakima	43	X	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	19	X	
29	Wide Hollow Creek at old STP at Union Gap	28	X	
30	Moxee Drain at Thorp Road near Union Gap	22	X	
31	Ahtanum Creek at Union Gap	19	X	
33	Yakima River at Parker	36	X	
40	Granger Drain at mouth near Granger	20	X	
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	35	X	
43	Toppenish Creek at Indian Church Road near Granger	31	X	
47	Satus Creek at gage at Satus	25	X	
48	Yakima River at RM 72 above Satus Creek near Sunnyside	36	X	
50	Yakima River at Kiona	30	X	
52	Sulphur Creek Wasteway near Sunnyside	37	X	
53	Satus Creek below Dry Creek near Toppenish	24	X	
54	Spring Creek at mouth at Whitstran	22	X	
56	Yakima River at Euclid Bridge at RM 55 near Grandview	27	X	
57	Satus Creek above Wilson-Charley Canyon near Toppenish	23	X	
Percentage of samples that exceed guidelines			94	9
Zinc, guideline: lowest effect level: 120 µg/g; severe effect level: 820 µg/g				
3	Jungle Creek near mouth near Cle Elum	150	X	
13	American River at Hells Crossing near Nile	210	X	
26	Naches River near North Yakima	120	X	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	170	X	
29	Wide Hollow Creek at old STP at Union Gap	210	X	
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	120	X	
57	Satus Creek above Wilson-Charley Canyon near Toppenish	160	X	
Percentage of samples that exceed guidelines			22	0



severe-effect level with the greatest frequency were: iron > manganese > chromium > nickel > arsenic. Concentrations of cadmium, copper, lead, mercury, and zinc did not exceed the severe effect level at any of the sites. Numerous sites had trace-element concentrations that exceeded the lowest effect level but did not exceed the severe effect level.

## Water

Trace-element concentrations in filtered- and unfiltered-water samples collected from the Yakima River Basin, 1987–90, are screened against

- (1) EPA ambient water-quality criteria for the protection of aquatic life and human health (U.S. Environmental Protection Agency, 1986 and 1992c),
- (2) EPA primary and secondary drinking-water regulations (U.S. Environmental Protection Agency, 1992b), and
- (3) EPA drinking-water human-health advisories (U.S. Environmental Protection Agency, 1992b).

All EPA ambient water-quality criteria are nonenforceable guidelines that may provide the basis for State standards (Nowell and Resek, 1994) and are designed to protect human health and aquatic organisms from deleterious-element concentrations. Ambient water-quality data in the Yakima River Basin was screened in order to identify trace-element concentrations that may require study by State and local health agencies. State and local health agencies are responsible for issuing advice or formal advisories to protect the health of their constituents.

The primary drinking-water regulations have been established for contaminants that are known to be present in public-water systems and which may affect human health adversely; secondary drinking-water regulations are nonenforceable guidelines, based on aesthetics, for contaminants that can affect the odor or appearance of drinking water adversely (Nowell and Resek, 1994). Health advisories provide nonregulatory levels of contaminants in drinking water which would result in no known or anticipated health effects.

## Ambient Stream Water

### Aquatic life

According to EPA's interim guidance on aquatic-life criteria for metals (U.S. Environmental

Protection Agency, 1992a, p. 4), the toxicity tests that form the basis for EPA ambient water-quality criteria for the protection of aquatic life were generally done in water "lower in metal-binding particulate matter and dissolved organic carbon than most ambient waters[. Therefore], these toxicity tests may overstate the ambient toxicity of nonbiomagnified metals that interact with particulate matter or dissolved organic matter." Trace elements used for the toxicity tests by EPA were analyzed from an unfiltered-water sample using a total-recoverable method of analysis. This type of analysis for ambient water, with metal-binding phases, may extract trace elements from the particulate or carbon phases and, consequently, overstate ambient toxicity. Presently, EPA provides guidance on the use of two new methods that may be used to implement aquatic-life criteria without overstating ambient toxicity (U.S. Environmental Protection Agency, 1992c). The new methods can be used to adjust the numeric value of the criteria in proportion to the quantity of the metal-binding phases in the ambient water. These methods may increase the numeric value of the aquatic-life criteria (making it less stringent). The simplest and most conservative method, however, is to use the total-recoverable method for filtered- and unfiltered-water samples and to compare such measurements to EPA criteria.

Although EPA's ambient water-quality criteria are based on analyses of unfiltered-water samples, the trace-element concentrations in filtered-water samples analyzed for this report were often high enough to equal or exceed the criteria. Consequently, EPA's ambient water-quality criteria are used as screening values for the protection of aquatic organisms. For many trace elements, including cadmium, chromium, lead, nickel, silver, and zinc, aquatic toxicity is related to the hardness of the water—toxicity increases (the screening value decreases) as hardness decreases. For example, as water hardness ranges from 50 to 200 mg/L (milligrams per liter) as calcium carbonate, lead toxicity to aquatic organisms ranges from 1.3 to 7.7 µg/L (micrograms per liter) [U.S. Environmental Protection Agency, 1986]. For trace elements with toxicity that varies with water hardness, the ambient-water hardness at the time of sample collection was used to derive the appropriate screening value. Both acute (1-hour average concentrations) and chronic (4-day average concentrations) trace-element criteria for aquatic life are used to evaluate waters in the Yakima River Basin; however, trace-element con-

centrations in the basin are from discrete measurements, rather than multiple measurements to derive 1-hour or 4-day average concentrations.

Concentrations of cadmium, copper, lead, mercury, silver, and zinc in filtered-water samples exceeded the screening value (based on EPA's ambient water-quality criteria for the protection of aquatic organisms) at two or more sites. Additionally, concentrations of cadmium, mercury, and zinc at several sites exceeded acute or chronic screening values for the protection of aquatic life (table 7). Sites with zinc exceedances include those receiving irrigation return flow, such as East Toppenish Drain at Wilson Road near Toppenish (site 37), as well as those in mountainous areas, such as Bumping River at Soda Springs Walkway near Nile (site 17). Acute and chronic exceedances of copper exist in the Yakima River at Cle Elum (site 6) and the Yakima River at Umtanum (site 19). These exceedances, although infrequent, happen during periods of winter-storm runoff. Similar to copper, the number of chronic, aquatic-life exceedances of cadmium and mercury are few in number. Unlike copper, however, exceedances of cadmium and mercury do not appear to be associated with any particular season or streamflow condition.

Some of the above aquatic-life exceedances are related to hydrologic conditions or events. For example, concentrations of cadmium generally exceed the chronic aquatic-life criterion during periods of winter warming, in spring during snowmelt, and during storms. Of the seven fixed sites, exceedances are found principally in the Kittitas Valley at the Yakima River at Cle Elum and at the Yakima River at Umtanum. The exceedances, in part, are related to hardness, which generally is low in waters of the Kittitas Valley. Hardness in the Kittitas Valley is less than 50 mg/L as calcium carbonate and is noteworthy because aquatic-life screening values for cadmium decrease as hardness decreases. Consequently, concentrations of cadmium in the Kittitas Valley are more likely to exceed screening values for the protection of aquatic life than are cadmium concentrations in the mid- and lower Yakima Valley. Cadmium also was detected in the Kittitas Valley during synoptic samplings in July 1987–88—chronic aquatic-life exceedances at these times were measured in the main stem near Ellensburg (site 9) and in the main stem above Roza Dam (site 23).

Concentrations of cadmium, chromium, copper, iron, lead, mercury, and silver in unfiltered-water samples exceeded the screening value (based on EPA's ambient water-quality criteria for the protection of aquatic organisms) at two or more fixed sites (table 8). In addition, concentrations of chromium and copper exceeded both acute and chronic screening values at several sites. The aquatic-life screening values for chromium vary depending on valency; for example, assuming a water hardness of 50 mg/L as calcium carbonate, the criteria for hexavalent (+ 6) and trivalent (+ 3) chromium range from 11 µg/L to 120 µg/L, respectively. Because chromium was analyzed as the sum of the trivalent and hexavalent species, the chromium data—as a conservative measure—were evaluated against the more stringent hexavalent chromium criterion. The screening values for the protection of aquatic life from copper were exceeded during spring and summer at the following fixed stations: Yakima River at Cle Elum (site 6), Yakima River at Umtanum (site 19), Yakima River above Ahtanum Creek at Union Gap (site 32), and Sulphur Creek Wasteway near Sunnyside (site 52). The seasonality of these exceedances coincides with those measured from 1953–85 (Rinella and others, 1992). The 1953–85 exceedances of copper, in part, were attributed to the use, past and present, of copper sulphate—a herbicide used by some irrigation districts to control nuisance-aquatic growths in irrigation canals (Lee Henderson, Kittitas Reclamation District, oral commun., 1989).

In this study, relatively few determinations of unfiltered-water samples were made using the *total-recoverable* method of analysis—the method of analysis used by the EPA to derive aquatic-life criteria. Instead, element concentrations often are measured directly on suspended sediment using a *total* method of analysis, and on dissolved trace elements in filtered-water samples. To estimate a total-recoverable measurement in unfiltered-water samples (for the purpose of making comparisons with EPA guidelines), trace-element concentrations on suspended sediment are added to concentrations in filtered-water samples<sup>5</sup>. The resultant values, termed *calculated-total concentrations*, are regressed, in turn, against the relatively small number of trace-element

<sup>5</sup>Element concentrations in suspended sediment (in micrograms per gram) were converted to volumetric concentrations (in micrograms per liter) and added to the dissolved element concentration to obtain an estimate of the trace-element concentration in an unfiltered-water sample.

**Table 7. Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987-90**

[The term "filtered water" is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; element concentrations of antimony, barium, beryllium, boron, chromium, cyanide, iron, nickel, and selenium met: (1) U.S. Environmental Protection Agency (1986, 1992c) ambient water-quality criteria or State Standards (Washington State Administrative Code (1992)) for the protection of aquatic life and human health, and (2) U.S. Environmental Protection Agency (1992b) drinking-water guidelines; only detectable concentrations were evaluated against water-quality guidelines; percentages were calculated using all measurements (censored and detected) for all sites sampled; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate during a single site visit, only one element concentration per visit was evaluated; for reference purposes, the aquatic-life guidelines listed below are based on a hardness of 50 milligrams per liter (mg/L) as calcium carbonate; for evaluation of detected concentrations for aquatic life, the ambient hardness was used to calculate screening values; the listed acute and chronic criteria for arsenic and mercury and the chronic criteria for silver do not vary with water hardness; µg/L, micrograms per liter; \*, waterway not required to meet drinking-water guidelines; --, no current guideline exists; DID, drainage irrigation district; STP, sewage treatment plant]

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria			Drinking-water guidelines		
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human-health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Arsenic:</b>								
Ambient water-quality criteria:								
Aquatic life:								
Acute: 1-hour average 360 µg/L once in 3 years								
Chronic: 4-day average 190 µg/L once in 3 years								
Human health:								
Consumption of aquatic organisms and water: 0.18 µg/L								
Consumption of aquatic organisms only: 1.4 µg/L								
Note: Human-health guidelines are based on a slope factor ( $q_1$ ) of 1.75 (milligrams per kilogram per day) <sup>1</sup> , a bioconcentration factor of 44 liters per kilogram, a lifetime risk of cancer equivalent to 1 in 100,000, a human consumption rate of fish equal to 6.5 grams per day (about one 6-ounce fillet per month—the national average), a human consumption rate of water equal to 2 liters per day, a human-body weight of 70 kilograms (154 pounds), and a life expectancy of 70 years <sup>9</sup> .								
Drinking water:								
Regulation: 50.0 µg/L (Maximum Contaminant Level)								
Human-health advisory: 0.2 µg/L Risk-specific dose (RSD)								
Note: Human-health advisory is based on a slope factor ( $q_1$ ) of 1.75 (milligrams per kilogram per day) <sup>1</sup> , a lifetime risk of cancer equivalent to 1 in 100,000, a human consumption rate of water equal to 2 liters per day, a human-body weight of 70 kilograms (154 pounds), and a life expectancy of 70 years <sup>10</sup> .								
Note: Because the limit of determination for arsenic (1 µg/L) exceeds the human-health criterion for consumption of aquatic organisms and water (0.18 µg/L) and the human-health advisory for drinking water (0.2 µg/L), censored data (concentrations reported as <1 µg/L) were not counted as exceeding these screening values.								
3	Jungle Creek near mouth near Cle Elum	1	0	0	1	1	0	1
26	Naches River near North Yakima	15	0	0	1	0	0	1
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	1	0	0	1	0	0	1
32	Yakima River above Ahtanum Creek at Union Gap	23	0	0	4	0	0	4
50	Yakima River at Kiona	25	0	0	20	15	0	20
52	Sulphur Creek Wasteway near Sunnyside	15	0	0	15	15	*	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	14	0	0	11	6	0	11
Percentage of samples that exceed screening values			0	0	43	30	0	31



**Table 7.** Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987-90—Continued

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria				Drinking-water guidelines	
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Cadmium:</b> Ambient water-quality criteria: Aquatic life: Acute: 1-hour average 1.8 µg/L once in 3 years Chronic: 4-day average 0.66 µg/L once in 3 years Drinking water: Regulation: 5 µg/L (Maximum Contaminant Level)								
6	Yakima River at Cle Elum	42	0	6	--	--	0	--
9	Yakima River at Thorp Highway Bridge at Ellensburg	3	0	1	--	--	0	--
19	Yakima River at Umtanum	40	0	4	--	--	0	--
23	Yakima River above canal diversion at river mile 128 at Roza Dam	1	0	1	--	--	0	--
26	Naches River near North Yakima	38	1	1	--	--	0	--
43	Toppenish Creek at Indian Church Road near Granger	2	0	1	--	--	0	--
	Percentage of samples that exceed screening values		0.3	4	--	--	0	--
<b>Copper:</b> Ambient water-quality criteria: Aquatic life: Acute: 1-hour average 9.2 µg/L once in 3 years Chronic: 4-day average 6.5 µg/L once in 3 years Drinking water: Regulation: 1,300 µg/L (Maximum Contaminant Level Goal)								
9	Yakima River at Cle Elum	42	4	4	--	--	0	--
16	Yakima River at Umtanum	39	1	2	--	--	0	--
	Percentage of samples that exceed screening values		2	2	--	--	0	--



**Table 7.** Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria				Drinking-water guidelines	
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human-health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Lead:</b> Ambient water-quality criteria: Aquatic life: Acute: 1-hour average 34 µg/L once in 3 years Chronic: 4-day average 1.3 µg/L once in 3 years Drinking water: Regulation: 0 µg/L (Maximum Contaminant Level Goal) Note: Because the limit of determination for lead (0.5 µg/L) exceeds the drinking-water regulation (0 µg/L), censored data (concentrations reported as <0.5 µg/L) were not counted as drinking-water exceedances.								
19	Yakima River at Umtanum	39	0	1	--	--	0	--
26	Naches River near North Yakima	37	0	0	--	--	4	--
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	1	0	0	--	--	1	--
32	Yakima River above Ahtanum Creek at Union Gap	41	0	0	--	--	2	--
36	Yakima River at river mile 91 at Zillah	1	0	1	--	--	1	--
50	Yakima River at Kiona	39	0	1	--	--	6	--
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	38	0	1	--	--	2	--
	Percentage of samples that exceed screening values		0	1	--	--	6	--
<b>Manganese:</b> Drinking water: Regulation: 200 µg/L (Maximum Contaminant Level Goal)								
35	Unnamed drain at Progressive Road near Harrah	1	--	--	--	--	1	--
	Percentage of samples that exceed screening values		--	--	--	--	1	--

**Table 7. Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90—Continued**

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria			Drinking-water guidelines		
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human-health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Mercury:</b> Ambient water-quality criteria: Aquatic life: Acute: 1-hour average 2.4 µg/L Chronic: 4-day average 0.012 µg/L Human health: Consumption of aquatic organisms and water: 0.14 µg/L Consumption of aquatic organisms only: 0.15 µg/L Note: Human-health guidelines are based on a reference dose (RfD) of 6 x 10 <sup>-5</sup> milligrams per kilogram per day, a bioconcentration factor of 5,500 liters per kilogram, a human consumption rate of fish equal to 6.5 grams per day (about one 6-ounce filet per month—the national average), a human consumption rate of water equal to 2 liters per day, a human-body weight of 70 kilograms (154 pounds), and a life expectancy of 70 years <sup>11</sup> . Drinking water: Regulation: 2 µg/L (Maximum Contaminant Level) Human-health advisory: 0.4 µg/L Lifetime health advisory (relative-source contribution from drinking water is assumed to be 20 percent) Note: Human-health advisory is based on a reference dose (RfD) of 6 x 10 <sup>-5</sup> milligrams per kilogram per day, a human consumption rate of water equal to 2 liters per day, a human-body weight of 70 kilograms (154 pounds), and a life expectancy of 70 years <sup>12</sup> . Note: Because the limit of determination for mercury (0.1 µg/L) exceeds the chronic criterion (0.012 µg/L), censored data (concentrations reported as <0.1 µg/L) were not counted as exceeding the chronic-mercury screening value.								
6	Yakima River at Cle Elum	42	0	2	2	2	0	1
19	Yakima River at Umtanum	40	0	3	3	3	0	1
26	Naches River near North Yakima	38	0	1	1	1	0	0
32	Yakima River above Ahtanum Creek at Union Gap	42	0	2	2	2	0	0
50	Yakima River at Kiona	43	0	3	2	2	0	0
52	Sulphur Creek Wasteway near Sunnyside	41	0	2	1	1	*	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	37	0	1	0	0	0	0
Percentage of samples that exceed screening values			0	5	4	4	0	1

**Table 7. Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987-90—Continued**

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria				Drinking-water guidelines	
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Silver:</b> Ambient water-quality criteria: Aquatic life: Acute: 1.2 µg/L at any time Chronic: 0.12 µg/L at any time Drinking water: Regulation: 100 µg/L (Secondary Maximum Contaminant Level) Note: Because the limit of determination for silver (1 µg/L) exceeds the chronic criterion (0.12 µg/L), censored data (concentrations reported as <1 µg/L) were not counted as exceeding the chronic-silver screening value.								
9	Yakima River at Thorp Highway Bridge at Ellensburg	2	1	1	--	--	0	--
12	South Fork Manastash Creek near Ellensburg	1	1	1	--	--	0	--
19	Yakima River at Umtanum	6	1	2	--	--	0	--
26	Naches River near North Yakima	3	2	2	--	--	0	--
32	Yakima River above Ahtanum Creek at Union Gap	17	1	2	--	--	0	--
36	Yakima River at river mile 91 at Zillah	2	1	1	--	--	0	--
43	Toppenish Creek at Indian Church Road near Granger	2	0	1	--	--	0	--
46	DID 3 Drain below STP at Midvale Road at Sunnyside	1	0	1	--	--	0	--
50	Yakima River at Kiona	16	0	1	--	--	0	--
52	Sulphur Creek Wasteway near Sunnyside	7	0	2	--	--	*	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	4	0	1	--	--	0	--
Percentage of samples that exceed screening values			7	15	--	--	0	--

**Table 7. Summary of trace-element concentrations in filtered-water samples that exceed screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987-90—Continued**

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria				Drinking-water guidelines	
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Zinc:</b> Ambient water-quality criteria: Aquatic life: Acute: 1-hour average 65 µg/L once in 3 years Chronic: 4-day average 59 µg/L once in 3 years Drinking water: Regulation: 5,000 µg/L (Secondary Maximum Contaminant Level)								
13	American River near Nile	2	1	1	--	--	0	--
17	Bumping River at Soda Springs Walkway near Nile	1	1	1	--	--	0	--
23	Yakima River above canal diversion at river mile 128 at Roza Dam	1	1	1	--	--	0	--
30	Moxee Drain at Thorp Road near Union Gap	2	1	1	--	--	0	--
37	East Toppenish Drain at Wilson Road near Toppenish	2	1	1	--	--	0	--
55	Yakima River at Mabton	1	1	1	--	--	0	--
Percentage of samples that exceed screening values			6	6	--	--	0	--

<sup>1</sup>U.S. Environmental Protection Agency (1986, 1992c)

<sup>2</sup>U.S. Environmental Protection Agency (1992c)

<sup>3</sup>U.S. Environmental Protection Agency (1992b)

<sup>4</sup>U.S. Environmental Protection Agency (1992b), Nowell and Resek (1994)

<sup>5</sup>Element concentrations were obtained by instantaneous measurements rather than 1-hour averages.

<sup>6</sup>Element concentrations were obtained by instantaneous measurements rather than 3-day averages.

<sup>7</sup>Assumes that 100 percent of exposure to the element is from consumption of contaminated water and contaminated aquatic organisms.

<sup>8</sup>Assumes that 100 percent of exposure to the element is from consumption of contaminated water.

<sup>9</sup>U.S. Environmental Protection Agency (1992c, 1992d)

<sup>10</sup>U.S. Environmental Protection Agency (1992c, 1992d)

<sup>11</sup>U.S. Environmental Protection Agency (1992c, 1992d)

<sup>12</sup>U.S. Environmental Protection Agency (1992c, 1992d)



**Table 8. Summary of trace-element concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987-90**

[The term "unfiltered water" refers to the chemical analysis of a water sample that has not been filtered or centrifuged, nor in any way, altered from the original matrix; element concentrations of barium, beryllium, boron, nickel, and zinc met: (1) U.S. Environmental Protection Agency (1986, 1992c) ambient water-quality criteria or State Standards (Washington State Administrative Code, 1992) for the protection of aquatic life and human health, and (2) U.S. Environmental Protection Agency (1992b) drinking-water guidelines; only detectable concentrations were evaluated against water-quality guidelines; percentages were calculated using all measurements (censored and detected) for all sites sampled; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate during a single site visit, only one element concentration per visit was evaluated; for reference purposes, the aquatic-life guidelines listed below are based on a hardness of 50 milligrams per liter (mg/L) as calcium carbonate; for evaluation of detected concentrations for aquatic life, the ambient hardness was used to calculate screening values; the listed acute and chronic criteria for chromium and mercury and the chronic criteria for silver do not vary with water hardness; µg/L, micrograms per liter; \*, waterway not required to meet drinking-water guidelines; --, no current guideline exists]

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria				Drinking-water guidelines	
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Cadmium:</b>								
Ambient water-quality criteria:								
Aquatic life:								
Acute: 1-hour average 1.8 µg/L once in 3 years								
Chronic: 4-day average 0.66 µg/L once in 3 years								
Drinking water:								
Regulation: 5 µg/L (Maximum Contaminant Level)								
Note: Because the limit of determination for cadmium (1 µg/L) exceeds the chronic criterion (0.66 µg/L), censored data (concentrations reported as <1 µg/L) were not counted as exceeding the chronic-cadmium screening value.								
6	Yakima River at Cle Elum	3	0	1	--	0	--	--
32	Yakima River above Ahtanum Creek at Union Gap	4	0	1	--	0	--	--
Percentage of samples that exceed screening values								
			0	11	--	0	--	--
<b>Chromium:</b>								
Ambient water-quality criteria:								
Aquatic life:								
Acute: 1-hour average 16 µg/L once in 3 years								
Chronic: 4-day average 11 µg/L once in 3 years								
Drinking water:								
Regulation: 100 µg/L (Maximum Contaminant Level)								
6	Yakima River at Cle Elum	3	0	1	--	0	--	0
19	Yakima River at Umtanum	3	1	1	--	0	--	0
50	Yakima River at Kiona	3	1	1	--	0	--	0
52	Sulphur Creek Wasteway near Sunnyside	2	1	2	--	*	--	*
Percentage of samples that exceed screening values								
			17	28	--	0	--	0

**Table 8.** Summary of trace-element concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values						
			Ambient water-quality criteria				Drinking-water guidelines		
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>	
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>			
<b>Copper:</b>									
Ambient water-quality criteria:									
Aquatic life:									
Acute: 1-hour average 9.2 µg/L once in 3 years									
Chronic: 4-day average 6.5 µg/L once in 3 years									
Drinking water:									
Regulation: 1,300 µg/L (Maximum Contaminant Level Goal)									
Note: Because the limit of determination for copper (10 µg/L) exceeds the acute (9.2 µg/L) and chronic (6.5 µg/L) criteria, censored data (concentrations reported as <10 µg/L) were not counted as exceeding acute- or chronic-copper screening values.									
6	Yakima River at Cle Elum	3	1	1	--	--	0	--	--
19	Yakima River at Umtanum	3	1	1	--	--	0	--	--
32	Yakima River above Ahtanum Creek at Union Gap	4	2	2	--	--	0	--	--
50	Yakima River at Kiona	3	1	1	--	--	0	--	--
52	Sulphur Creek Wasteway near Sunnyside	2	1	1	--	--	*	--	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	2	1	1	--	--	0	--	--
Percentage of samples that exceed screening values			39	39	--	--	0	--	--
<b>Iron:</b>									
Ambient water-quality criteria:									
Aquatic life:									
Chronic: 1,000 µg/L									
Drinking water:									
Regulation: 300 µg/L (Secondary Maximum Contaminant Level)									
6	Yakima River at Cle Elum	3	--	1	--	--	2	--	--
19	Yakima River at Umtanum	3	--	2	--	--	3	--	--
26	Naches River near North Yakima	1	--	0	--	--	1	--	--
32	Yakima River above Ahtanum Creek at Union Gap	4	--	2	--	--	4	--	--
50	Yakima River at Kiona	3	--	3	--	--	3	--	--
52	Sulphur Creek Wasteway near Sunnyside	1	--	1	--	--	*	--	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	2	--	2	--	--	2	--	--
Percentage of samples that exceed screening values			--	65	--	--	94	--	--

**Table 8.** Summary of trace-element concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values							
			Ambient water-quality criteria			Drinking-water guidelines				
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human-health advisories <sup>4</sup>		
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>				
<b>Lead:</b>										
Ambient water-quality criteria:										
Aquatic life:										
Acute: 1-hour average 34 µg/L once in 3 years										
Chronic: 4-day average 1.3 µg/L once in 3 years										
Drinking water:										
Regulation: 0 µg/L (Maximum Contaminant Level Goal)										
Note: Because the limit of determination for lead (5 µg/L) exceeds the chronic criterion (1.3 µg/L) and the drinking-water regulation (0 µg/L), censored data (concentrations reported as <5 µg/L) were not counted as exceeding these guidelines.										
6	Yakima River at Cle Elum	3	0	1	--	--	1	--		
19	Yakima River at Umtanum	3	0	2	--	--	2	--		
32	Yakima River above Ahtanum Creek at Union Gap	4	0	1	--	--	1	--		
50	Yakima River at Kiona	3	0	1	--	--	1	--		
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	2	0	1	--	--	1	--		
Percentage of samples that exceed screening values			0	33	--	--	33	--		
<b>Manganese:</b>										
Drinking water:										
Regulation: 200 µg/L (Maximum Contaminant Level Goal)										
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	2	--	--	--	--	1	--		
Percentage of samples that exceed screening values			--	--	--	--	6	--		



**Table 8.** Summary of trace-element concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987-90—Continued

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values						
			Ambient water-quality criteria				Drinking-water guidelines		
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>	
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>			
<b>Mercury:</b>									
Ambient water-quality criteria:									
Aquatic life:									
Acute: 1-hour average 2.4 µg/L									
Chronic: 4-day average 0.012 µg/L									
Human health:									
Consumption of aquatic organisms and water: 0.14 µg/L									
Consumption of aquatic organisms only: 0.15 µg/L									
Note: Human-health guidelines are based on a reference dose (RfD) of 6 x 10 <sup>-5</sup> milligrams per kilogram per day, a bioconcentration factor of 5,500 liters per kilogram, a human consumption rate of fish equal to 6.5 grams per day (about one 6-ounce filet per month—the national average), a human consumption rate of water equal to 2 liters per day, a human-body weight of 70 kilograms (154 pounds), and a life expectancy of 70 years <sup>9</sup>									
Drinking water:									
Regulation: 2 µg/L (Maximum Contaminant Level)									
Human-health advisory: 0.4 µg/L Lifetime health advisory (relative-source contribution from drinking water is assumed to be 20 percent)									
Note: Human-health advisory is based on a reference dose (RfD) of 6 x 10 <sup>-5</sup> milligrams per kilogram per day, a human consumption rate of water equal to 2 liters per day, a human-body weight of 70 kilograms (154 pounds), and a life expectancy of 70 years <sup>10</sup>									
Note: Because the limit of determination for mercury (0.1 µg/L) exceeds the chronic criterion (0.012 µg/L), censored data (concentrations reported as <0.1 µg/L) were not counted as exceeding the chronic-mercury screening value.									
6	Yakima River at Cle Elum	3	0	1	0	0	0	0	0
19	Yakima River at Umtanum	3	0	2	2	2	0	0	0
26	Naches River near North Yakima	1	0	1	1	1	0	0	1
32	Yakima River above Ahtanum Creek at Union Gap	4	0	1	0	0	0	0	0
50	Yakima River at Kiona	3	0	1	1	1	0	0	0
52	Sulphur Creek Wasteway near Sunnyside	2	0	1	0	0	*	*	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	2	0	1	1	1	0	0	0
Percentage of samples that exceed screening values			0	44	28	28	0	0	6



**Table 8.** Summary of trace-element concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Total number of samples	Number of samples that exceed screening values					
			Ambient water-quality criteria			Drinking-water guidelines		
			Aquatic life <sup>1</sup>		Human health <sup>2</sup>		Regulations <sup>3</sup>	Human- health advisories <sup>4</sup>
			Acute <sup>5</sup>	Chronic <sup>6</sup>	Aquatic organisms and water <sup>7</sup>	Aquatic organisms only <sup>8</sup>		
<b>Silver:</b> Ambient water-quality criteria: Aquatic life: Acute: 1.2 µg/L at any time Chronic: 0.12 µg/L at any time Drinking water: Regulation: 100 µg/L (Secondary Maximum Contaminant Level) Note: Because the limit of determination for silver (1 µg/L) exceeds the chronic criterion (0.12 µg/L), censored data (concentrations reported as <1 µg/L) were not counted as exceeding the chronic-silver screening value								
19	Yakima River at Umtanum	3	0	1	--	--	0	--
32	Yakima River above Ahtanum Creek at Union Gap	4	1	1	--	--	0	--
52	Sulphur Creek Wasteway near Sunnyside	2	1	1	--	--	*	*
Percentage of samples that exceed screening values			7	15	--	--	0	--

<sup>1</sup>U.S. Environmental Protection Agency (1986, 1992c)

<sup>2</sup>U.S. Environmental Protection Agency (1992c)

<sup>3</sup>U.S. Environmental Protection Agency (1992b)

<sup>4</sup>U.S. Environmental Protection Agency (1992b); Nowell and Resek (1994)

<sup>5</sup>Element concentrations were obtained by instantaneous measurements rather than 1-hour averages.

<sup>6</sup>Element concentrations were obtained by instantaneous measurements rather than 3-day averages.

<sup>7</sup>Assumes that 100 percent of exposure to the element is from consumption of contaminated water and contaminated aquatic organisms.

<sup>8</sup>Assumes that 100 percent of exposure to the element is from consumption of contaminated water.

<sup>9</sup>U.S. Environmental Protection Agency (1992c, 1992d)

<sup>10</sup>U.S. Environmental Protection Agency (1992c, 1992d)

**Table 9.** Summary of estimated total-recoverable iron and manganese concentrations in unfiltered-water samples that exceeded screening values derived from water-quality guidelines, Yakima River Basin, Washington, 1987–90

[The term “unfiltered water” refers to the chemical analysis of a water sample that has not been filtered or centrifuged, nor in any way, altered from the original matrix; estimated total-recoverable concentrations of iron and manganese were derived using prediction equations generated from a least-squares regression made between the measured total-recoverable concentration in unfiltered water and the calculated-total concentrations; estimated concentrations were evaluated against (1) U.S. Environmental Protection Agency (EPA) (1986, 1992c) ambient water-quality criteria for the protection of aquatic life, and (2) U.S. EPA (1992b) drinking-water guidelines; µg/L, micrograms per liter; \*, waterway not required to meet drinking-water guidelines; --, no current guideline exists]

Site reference number	Site name	Number of estimated total-recoverable concentrations	Number of samples that exceed screening values	
			Aquatic life: Chronic	Drinking-water regulations
<b>Iron:</b> Ambient water-quality criteria: Aquatic life: Chronic: 1,000 µg/L Drinking water: Regulation: 300 µg/L (Secondary Maximum Contaminant Level)				
6	Yakima River at Cle Elum	1	1	1
19	Yakima River at Umtanum	3	2	2
32	Yakima River above Ahtanum Creek at Union Gap	12	3	10
50	Yakima River at Kiona	15	5	14
52	Sulphur Creek Wasteway near Sunnyside	6	5	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	4	1	4
Percentage of estimated concentrations that exceed screening values			41	90
<b>Manganese:</b> Drinking water: Regulation: 200 µg/L (Maximum Contaminant Level Goal)				
6	Yakima River at Cle Elum	1	--	0
19	Yakima River at Umtanum	4	--	0
32	Yakima River above Ahtanum Creek at Union Gap	12	--	1
50	Yakima River at Kiona	15	--	0
52	Sulphur Creek Wasteway near Sunnyside	6	--	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	4	--	0
Percentage of estimated concentrations that exceed screening values			--	2

concentrations derived using the total-recoverable method on unfiltered-water samples. There were nine unfiltered-water samples analyzed for iron and manganese. Only iron and manganese had an adequate number of uncensored data (detectable-trace-element concentrations) for making the regressions. The resulting prediction equation<sup>6</sup> is used to convert the calculated-total concentrations to **estimated total-recoverable concentrations**, which are then compared against screening values as an aid in understanding water-quality concerns. The estimated total-recoverable concentrations of iron exceeded screening values for ambient water quality (1,000 µg/L) at most fixed sites (table 9). Nearly one-third of the estimated iron concentrations for the Yakima River above Ahtanum Creek at Union Gap and for the Yakima River at Kiona exceeded the screening value for aquatic organisms.

### Human Health

The ambient stream-water criteria for the protection of human health (table 7) consist of ambient concentrations which, for noncarcinogens, prevent adverse health effects in humans, and represent various levels of incremental cancer risk for suspected or proven carcinogens. The human-health criteria are designed to indicate human exposure to a contaminant from (1) ingestion of water and aquatic organisms, and (2) ingestion of aquatic organisms (U.S. Environmental Protection Agency, 1992c). In the former, 100 percent of the exposure to humans is assumed to be from consumption of water containing a specified-contaminant concentration and aquatic organisms that have biologically concentrated a contaminant from ambient stream water according to an assumed biological concentration factor, (Nowell and Resek, 1994; U.S. Environmental Protection Agency, 1992c). In the latter, 100 percent of the exposure to humans is assumed to be from consumption of aquatic organisms that have biologically concentrated a contaminant from water on the basis of a bioconcentration factor. Equations for deriving ambient-contaminant concentrations for the protection of human health are given in Nowell and Resek (1994). The following discussion is focused

principally on arsenic (a carcinogen) and mercury (a noncarcinogen); concentrations of each exceeded screening values for human health.

For carcinogens, the human-health criteria are derived from a two-part evaluation in which the trace element is assigned a weight of evidence classification and a slope factor. The weight of evidence classification is the likelihood that a trace element is a human carcinogen. Arsenic, which is measured in filtered-water samples in the Yakima River Basin, has a "Group A" weight-of-evidence classification—a human carcinogen (U.S. Environmental Protection Agency, 1989). The slope factor is generally a plausible upper-bound estimate (95-percent confidence limit) of a human developing cancer as a result of a lifetime (70 years) of exposure to a particular level of a potential carcinogen. Slope factors are derived from mathematical models that are used on available data sets. These models extrapolate from carcinogenic responses observed at high doses in experimental animals to responses expected in humans from lower exposure levels in the environment. If the extrapolation model selected is EPA's linearized-multistage model (as in the case of arsenic) then the resultant slope factor is known as  $q_1^*$  (U.S. Environmental Protection Agency, 1989).

For example, the slope factor or  $q_1^*$  for arsenic is a 1.75 risk per milligram contaminant per kilograms body weight per day (mg/kg/day)<sup>-1</sup> (U.S. Environmental Protection Agency, 1992c). For carcinogens, the derivation of human-health criteria for ambient stream water is contingent on several additional assumptions which include:

**Risk Level (RL)** = an assigned level of maximum-acceptable individual-lifetime risk. Screening values for human health are based on a  $RL = 10^{-5}$ —a level of risk not to exceed one excess case of cancer per 100,000 individuals exposed over a 70-year lifetime (U.S. Environmental Protection Agency, 19993).

**Consumption Rate (CR)** = Mean daily consumption rate, in kilograms per day (kg/d), of the species of interest by the general population or subpopulation of concern averaged over a 70-year lifetime. Screening values for human health were derived using a CR of

<sup>6</sup> The regression equation for iron yielded an R-squared value of 0.87 and a prediction equation,  $y = 517 + 0.58x$  ( $y$  = estimated total-recoverable iron concentration;  $x$  = calculated total-iron concentration).

0.0065 kg/d—an estimate of the average fish and shellfish consumption by the general United States population (U.S. Environmental Protection Agency, 1993). The value is approximately one 6-oz (ounce) fillet of fish per month. In addition, screening concentrations which include a measure of chemical uptake from the consumption of water (for example, EPA's health advisories for drinking water), or from the consumption of water and fish, use a CR of 2 liters of water per day—an estimate of the average water consumption by the general United States population.

Body Weight (BW) = Mean body weight, in kilograms, of a standard adult within the general population or subpopulation of concern. Screening values for human health were derived using a BW of 70 kg (about 154 pounds, the average weight of the general United States population.

Biological Concentration Factor = the ratio of the contaminant concentration in an aquatic organism, in milligrams per kilograms to the contaminant concentration in the surrounding water, in milligrams per liter, and is reported in units of liters per kilogram. A weighted-average biological concentration factor (BCF), adjusted to the average percent lipids in fish and shellfish (3 percent) is used by EPA in deriving human-health guidelines (U.S. Environmental Protection Agency, 1992c).

Concentrations of arsenic in ambient stream water (filtered-water samples) in the Yakima River Basin exceeded the screening value (based on EPA's ambient stream-water-quality criteria) for consumption of aquatic organisms and water (0.18 µg/L) at seven sites, and overall, exceeded the screening value in 43 percent of the samples (table 7). Additionally, arsenic concentrations in ambient stream water did not meet the screening value for consumption of aquatic organisms (1.4 µg/L) at four sites, and overall, exceeded the screening values in 30 percent of the samples (table 7). Exceedances of arsenic were found predominantly in the lower

Yakima Valley. The screening value for arsenic is based only on the inorganic form (Federal Register, 1992, p. 60887). The arsenic determination in the Yakima River Basin study, however, is based on the organic and inorganic forms of arsenic. As a conservative assumption for screening, arsenic is assumed to reside in ambient water in the pentavalent and (or) trivalent forms—the former being most likely in surface water (Eisler, 1988, p. 8). Methylated forms of arsenic also reside in surface water; their exact proportions, however, are not known (Hem, 1989, p. 144). Methylated forms are significantly less toxic than inorganic forms of arsenic (U.S. Environmental Protection Agency, 1993, p. III-56; Federal Register, 1992, p. 60887).

For noncarcinogens, the screening values are based on a Reference Dose (RfD) which represents a daily exposure (with uncertainty spanning perhaps an order of magnitude or more) to the human population (including sensitive subpopulations) that is probably without appreciable risk of causing deleterious effects during a 70-year lifetime (U.S. Environmental Protection Agency, 1992d). Additionally, the assumptions listed for BW, CR, and BCF are used in determining screening values for human health. Screening values are derived from EPA ambient water-quality criteria for human health (U.S. Environmental Protection Agency, 1992c).

For the purpose of calculating screening values for mercury (a noncarcinogen), EPA recommends that the RfD for methylmercury be lowered from 0.0003 mg/kg/day to  $6.0 \times 10^{-5}$  mg/kg/day (U.S. Environmental Protection Agency, 1992d). The lowering of the RfD is based on evidence that the fetus, and possibly pregnant women, are at increased risk of adverse neurological effects from exposure to methylmercury (U.S. Environmental Protection Agency, 1992d).

Under the usual conditions of temperature and pressure, mercury in surface water exists in inorganic forms which include the liquid ( $\text{Hg}^0$ ) and the ionic ( $\text{Hg}_2^+$  and  $\text{Hg}^{2+}$ ) states. In addition, inorganic forms in sediment and water can be methylated to highly soluble and toxic methylmercury (Moore, 1991). The analytical technique used in the Yakima River Basin study for measuring mercury in ambient stream water is defined as a total- (inorganic plus organic) mercury analysis. For screening purposes, total-mercury concentrations are compared to the screening values for human health. Using total-mercury concentrations

for comparison to screening values is conservative, because 100 percent of the mercury that accumulates in fish tissue (based on a BCF) is assumed to be in the toxic methylmercury form. It is this methylated form of mercury that is highly toxic to humans (U.S. Environmental Protection Agency, 1992d).

Concentrations of filtered mercury in ambient stream water in the Yakima River Basin exceeded the human-health screening value for consumption of aquatic organisms and water (0.14 µg/L) at six of the seven fixed sites, and overall, exceeded the screening value in 4 percent of the samples. Because the criterion for consumption of aquatic organisms is similar (0.15 µg/L) to the human-health screening value, the frequency and occurrence of exceedances for consumption of aquatic organisms is identical to that for consumption of aquatic organisms and water (table 7).

Concentrations of mercury in unfiltered-ambient water in the Yakima River Basin exceed the human-health screening value for consumption of aquatic organisms and water at four of the seven fixed sites—exceedances are identical for consumption of aquatic organisms (table 8). Using mercury concentrations from unfiltered-water samples for comparison to screening values is not conservative and exceedances should be interpreted accordingly.

## Drinking Water

Trace-element concentrations determined from filtered- and unfiltered-water samples were screened by making comparisons with EPA drinking-water regulations (U.S. Environmental Protection Agency, 1992c) and EPA advisories for human health (U.S. Environmental Protection Agency, 1992b).

The water samples from the Yakima River Basin that were compared to drinking-water guidelines represent untreated water (ambient stream water, rather than finished or treated water available for distribution to community-water supplies). The principal sites for diverting stream water to water-treatment plants for the city of Cle Elum and Yakima, respectively, are located on the Yakima River (RM 183) and on the Naches River (RM 18.4). The Yakima River at Cle Elum (RM 183) was the only sampling site adjacent to a water-treatment-plant intake. Although nearly all sites sampled in this study were not sources for domestic-water supplies, water-quality exceedances were important because

many of these sites reside in streams classified by the State of Washington as AA- or A-type waters. Classifications AA and A require that water “shall markedly and uniformly exceed requirements for all uses,” which include domestic-water supplies (Washington State Administrative Code, 1992). It is important to note, however, that “although a surface water in Washington State may be designated as a potential domestic-water source”, in Chapter 173–201 in the Washington State Administrative Code (1992), “approval for such use must first be obtained from the Washington State Department of Health following an evaluation of the water quality” (Harriet Ammann, Denise Laflamme, and Glen Patrick, Washington State Department of Health, written commun., 1993). Thus, trace-element concentrations, which exceed screening values (drinking-water regulations), in filtered and unfiltered stream-water samples are not an indication that human health is directly at risk.

## Regulations

The types of primary and secondary drinking-water regulations set forth by EPA include Maximum Contaminant Levels (MCLs), Maximum Contaminant Level Goals (MCLGs), and Secondary Maximum Contaminant Levels (SMCLs) [U.S. Environmental Protection Agency, 1991]. The MCLs represent achievable levels of drinking-water quality that take into consideration health effects, treatment feasibility, and aesthetic considerations. The MCLGs are non-enforceable health goals that are not expected to cause any adverse human-health effects over a lifetime of exposure and include a margin of safety. The SMCLs are unenforceable guidelines regarding the taste, odor, color, and certain other nonaesthetic effects of drinking water.

Concentrations of lead in filtered-water samples did not meet the screening value (based on the MCLG for drinking water) at six sites, and overall, exceeded the screening value in 6 percent of the samples (table 7). In the Yakima River at Kiona, a fixed site, lead was detected in 6 of 40 determinations for 1987–90—the MCLG for lead is 0 µg/L. Kiona had the largest number of detectable-lead concentrations; the next largest concentrations of lead were found in the Naches River near North Yakima, also a fixed site, which had detectable lead in 4 of 37 determinations. Manganese did not meet the MCLG at site 35, a drain that receives agricultural runoff at Progressive Road

near Harrah. Analysis of historical water-quality data (1953–85) in the Yakima River Basin show that drinking-water exceedances for lead and manganese represent about 2 percent of the total number of historical determinations (Rinella and others, 1992). During the July 1987 synoptic sampling, 30 µg/L of lead was detected by ICP in the Yakima River at RM 91 at Zillah; however, during a follow-up sampling in November 1987, lead was below the analytical limit of determination. The high lead concentration in July 1987 corresponds to anomalous concentrations of chromium and zinc and probably indicates the inadvertent use of a brass sampler.

Concentrations of iron in unfiltered-water samples did not meet the screening value (based on the SMCL for drinking water) at six of the fixed sites and, overall, exceeded the SMCL in 94 percent of the samples (table 8). However, because the corresponding iron concentrations in filtered-water samples met the SMCL, the exceedances in unfiltered water probably resulted from iron associated with sediment. Additionally, iron associated with sediment probably would be removed in the water-treatment process. The percentage of estimated total-recoverable concentrations that did not meet the SMCL for iron (table 9) was similar to that for iron in unfiltered-water samples. Again, however, the exceedances probably resulted from iron associated with sediment. The concentration of suspended sediment associated with iron exceedances ranged from 19 to 212 mg/L; however, concentrations associated with non-exceedances were less than or equal to 16 mg/L. The concentration of manganese in an unfiltered-water sample collected in the Yakima River at Euclid Bridge at RM 55 near Grandview (table 8) did not meet the MCLG (200 µg/L). Of 42 estimated total-recoverable manganese concentrations<sup>7</sup>, only one value exceeded the MCLG (table 9). This value was determined from a sample collected in November 1987 at the Yakima River above Ahtanum Creek at Union Gap, which contained high concentrations of suspended sediment (137 mg/L) and suspended manganese (2,930 µg/g).

---

<sup>7</sup>The regression for manganese yielded an R-squared value of 0.78 and a prediction equation,  $y = 20.4 + 0.74x$  ( $y$  = estimated total-recoverable manganese concentration;  $x$  = calculated total-manganese concentration).

## Health Advisories

Concentrations of trace elements in filtered- and unfiltered-water samples are screened for human-health effects by making comparisons to human-health advisories for drinking water (U.S. Environmental Protection Agency, 1992b). For the carcinogen arsenic, the human-health advisory listed in table 7 is a risk specific dose (RSD) associated with a specified RL and is calculated from the  $q_1^*$  for arsenic (Nowell and Resek, 1994). For the non-carcinogen mercury, the human-health advisory listed in table 7 is a lifetime-health advisory which is equal to 20 percent of the drinking water equivalent level (Nowell and Resek, 1994). Unlike the ambient water-quality criteria, however, health advisories are based only on the consumption of domestic water. In the Yakima River Basin study, ambient stream water is used to screen for health effects. Additionally, the aforementioned assumptions for BW, CR (2 liters of water per day), and RL ( $10^{-5}$ ) are applicable to screening values for health advisories.

Concentrations of arsenic in ambient stream water (filtered-water samples) in the Yakima River Basin exceeded the screening value (EPA's RSD of 0.2 µg/L) for consumption of domestic-drinking water at 6 sites and overall exceeded the screening value in 31 percent of the samples (table 7). The largest number of exceedances were in the lower Yakima Valley. For example, 20 of 25 ambient stream-water samples collected from the Yakima River at Kiona from 1987–90 exceeded the screening value. Although all 15 arsenic determinations at Sulphur Creek Wasteway near Sunnyside would have exceeded the screening value, these determinations were omitted from table 7 because Sulphur Creek is designated Class B in the Washington State Administrative Code (1992). Class B waterways are not required to meet water-quality guidelines for domestic-water supplies. The arsenic concentrations in Sulphur Creek (especially during base-flow conditions), however, may be indicative of arsenic concentrations in shallow domestic-ground-water supplies in the Sulphur Creek drainage.

Concentrations of mercury in ambient stream water (filtered-water samples) rarely exceeded the screening value (EPA's lifetime-health advisory of 0.4 µg/L). The two exceedances measured were in the Kittitas Valley—one from the Yakima River at Cle Elum and the other from the Yakima River at Umtanum (table 7). These exceedances, however, represent only 1 percent of all the samples.



## Fish Muscle

Fish muscle was analyzed for mercury for various fish taxa collected from four sites in 1991 (table 10). Sites sampled in 1991 generally coincided with mercury anomalies in fish livers from samplings in 1989–90 (Fuhrer, Fluter, and others, 1994). The median mercury concentration in fish muscle and for each fish species from each site (table 10) were screened against mercury concentrations in fish that are of potential public-health concern. According to the EPA, “exceedance of screening values should be taken as an indication that more intensive site-specific monitoring and (or) evaluation of human-health risk should be done” (U.S. Environmental Protection Agency, 1992d). State and local health agencies are responsible for issuing advice and (or) formal advisories to protect the health of their constituents.

The screening values for mercury in fish muscle are calculated from a dose-response variable which, for noncarcinogens such as mercury, is the *RfD*. Additionally, the assumptions listed above for *BW* and *CR* also are applicable. Screening values are determined for noncarcinogens by the following equation:

$$SV_n = \frac{RfD \times BW}{CR},$$

where,  $SV_n$  is the screening value for a noncarcinogen in units of milligrams per kilogram and *BW* and *CR* are defined earlier. Recognizing that screening levels change as *CR* and (or) *BW* vary, screening levels are calculated for standard adults, children, recreational fishermen, and subsistence fishermen (table 11). The EPA applies an uncertainty factor of 1,000 to the *RfD* as a safeguard to account for uncertainty in projecting human-health effects over a 70-year lifetime from animal studies of less than a lifetime duration (Moore, 1991, p. 205). In addition to applying an uncertainty factor, the EPA also has lowered the *RfD* to  $6.0 \times 10^{-5}$  for screening purposes as a conservative measure to prevent adverse neurological effects from exposure to methylmercury (U.S. Environmental Protection Agency, 1992c).

Muscle samples collected from rainbow trout and mountain whitefish at four sites in the Yakima River Basin contained mercury concentrations that were below the screening-value standards for adults

(table 11). The standard-adult screening value is based on the consumption of about one 6-ounce fillet per month for a 70-kg adult over a 70-year lifetime. Mercury concentrations in fish muscle of largescale sucker collected in the Yakima River at Kiona, however, exceeded the EPA screening value ( $0.65 \mu\text{g/g}$ ) for standard adults. Because some individuals consume different quantities of fish and differ in body weight, EPA screening values are calculated for subpopulations that include children, recreational fishermen, and subsistence fishermen in addition to standard adults (table 11). The concentration of mercury in fish muscle consumed by children eating an average of one 6-ounce fillet per month exceeded the screening value ( $0.11 \mu\text{g/g}$ ) for all species sampled and at all sites sampled. Similarly, screening values of  $0.14$  and  $0.03 \mu\text{g/g}$  were exceeded, respectively, by recreational fishermen (consumers of an average of about five 6-ounce fillets per month) and subsistence fishermen (consumers of an average of about 25 6-ounce filets per month) for all fish species sampled and at all sites sampled.

The screening values derived for mercury are based on the assumption that mercury resides in or has been converted to methylmercury in fish muscle that is consumed by humans. Total mercury, rather than methylmercury, was measured in the Yakima River Basin study in accordance with the EPA recommendation that total mercury be determined and the conservation assumption be made that all mercury is present as methylmercury. This approach has been deemed the most protective of human health because methylmercury is the most toxic mercury form for humans, and also because methylmercury is the most common form measured in fish muscle (U.S. Environmental Protection Agency, 1992d).

## SPATIAL AND TEMPORAL DISTRIBUTION OF TRACE ELEMENTS IN THE AQUATIC ENVIRONMENT

The spatial distribution of trace elements in sediment, water, and aquatic organisms in the Yakima River Basin is governed by factors which include: natural weathering and erosion of rocks and soils of the Cascade Range; decomposition of plant and animal matter; atmospheric deposition affected by natural events (ash fallout from the volcanic eruption of Mount St. Helens); human activities (combustion of

**Table 10.** Concentrations of mercury in muscle of rainbow trout, largescale sucker, and mountain whitefish, Yakima River Basin, Washington, October 29–31, 1991

[Concentrations are reported in micrograms per gram ( $\mu\text{g/g}$ ), dry weight; sample species: rainbow trout (*Oncorhynchus mykiss*), largescale sucker (*Catostomus macrocheilus*), and mountain whitefish (*Prosopium williamsoni*)]

Site reference number	Site name	Species sampled	Concentrations
8	Taneum Creek at Taneum Meadow near Thorp	Rainbow trout	0.32
			.20
			.20
			.25
			.25
19	Yakima River at Umtanum	Rainbow trout	.30
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	Rainbow trout	.18
			.17
			.21
			.20
			.17
50	Yakima River at Kiona	Largescale sucker	.41
			.84
			1.05
			.89
			1.56
			.98
			1.00
		Mountain whitefish	.23
			.38
			.58
			.29
			.31
			.25
			.28



**Table 11. Concentrations of mercury in fish muscle relative to U.S. Environmental Protection Agency screening values, Yakima River Basin, Washington, October 29–31, 1991**

[Shaded areas represent instances where median mercury concentrations in fish muscle exceed the screening values determined by the U.S. Environmental Protection Agency (EPA) [1992d]; screening values are based on a reference dose (RfD) for mercury of 0.00006 milligrams per kilogram per day (mg/kg/day); the reference dose is defined as the preferred toxicity value set by EPA for evaluating noncarcinogenic effects resulting from exposure of the human population to the specified element (U.S. Environmental Protection Agency, 1989); human consumption rates of fish are equal to (1) 6.5 grams per day (g/day), about one 6-ounce filet per month—the national average, (2) 30 g/day, about five 6-ounce filets per month—the 50th percentile for recreational fishermen, and (3) 140 g/day, about 25 6-ounce filets per month—the 90th percentile for recreational fishermen (for example, subsistence fishermen); a body weight of 12 kilograms (26 pounds) is assumed for children of 3 years and younger; a body weight of 70 kilograms (154 pounds) is assumed for adults; sample species: rainbow trout (*Oncorhynchus mykiss*), largescale sucker (*Catostomus macrocheilus*), mountain whitefish (*Prosopium williamsoni*); kg, kilogram; µg/g, micrograms per gram]

Site reference number	Site name	Species sampled	Subpopulation	Consumption rate (g/day)	Body weight (kg)	Mercury	
						Screening value (µg/g)	Median concentration <sup>1</sup> (µg/g)
8	Taneum Creek at Taneum Meadow near Thorp	Rainbow trout	Children	6.5	12	.11	.25
			Standard adults	6.5	70	.65	.25
			Recreational fishermen	30	70	.14	.25
			Subsistence fishermen	140	70	.03	.25
19	Yakima River at Umtanum	Rainbow trout	Children	6.5	12	.11	.30
			Standard adults	6.5	70	.65	.30
			Recreational fishermen	30	70	.14	.30
			Subsistence fishermen	140	70	.03	.30
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	Rainbow trout	Children	6.5	12	.11	.18
			Standard adults	6.5	70	.65	.18
			Recreational fishermen	30	70	.14	.18
			Subsistence fishermen	140	70	.03	.18
50	Yakima River at Kiona	Largescale sucker	Children	6.5	12	.11	.98
			Standard adults	6.5	70	.65	.98
			Recreational fishermen	30	70	.14	.98
			Subsistence fishermen	140	70	.03	.98
		Mountain whitefish	Children	6.5	12	.11	.29
			Standard adults	6.5	70	.65	.29
			Recreational fishermen	30	70	.14	.29
			Subsistence fishermen	140	70	.03	.29

<sup>1</sup>Median mercury concentrations for fish muscle from the Yakima River Basin; see table 10 for individual values.

fossil fuels and air emissions from industrial processes), transportation; municipal and industrial wastewater; urban stormwater runoff; paints; fertilizers; and pesticides. Data for trace-element concentrations (including some major elements) and organic carbon in streambed sediment, suspended sediment, filtered water, and aquatic organisms were collated and statistically summarized as percentiles (tables 12, 13, 14, and 15).

## Antimony

Concentrations of antimony in streambed sediment at the 32 biological-sampling sites ranged from 0.2 to 1.4  $\mu\text{g/g}$  (table 12) and are within the range of concentration (0.1 to 1.5  $\mu\text{g/g}$ ) reported by Parker (1967) for igneous and sedimentary rock on the earth's surface. The median concentration (0.4  $\mu\text{g/g}$ ) of antimony at the biological sites is slightly smaller than that determined from analysis of fine-fraction streambed sediment in other river basins of the United States (table 34, at back of report). The concentration maximum (1.4  $\mu\text{g/g}$ ) for the biological sampling sites, found in streambed sediment at the Naches River near north Yakima (site 26), probably is related to a geologic source. The headwaters of the Naches River contained large concentrations of antimony. For example, 3.0  $\mu\text{g/g}$  of antimony were found in streambed sediment, formed from Miocene and older volcanic rock, in the American River (Fuhrer, McKenzie, and others, 1994). Although antimony has a variety of industrial sources (Sittig, 1981) and would be expected at enriched levels at the mouths of urban and industrialized tributaries to the main stem, few cases of antimony enrichment were found in the Yakima River Basin during the occurrence and distribution survey (Fuhrer, McKenzie, and others, 1994); additionally, antimony enrichment was attributed to geologic sources.

Concentrations of antimony in suspended sediment at the seven fixed sites ranged from 0.5 to 3.1  $\mu\text{g/g}$  (table 13). The highest concentrations were in the Yakima River at Cle Elum (site 6). Here, the median (0.8  $\mu\text{g/g}$ ) and maximum (3.1  $\mu\text{g/g}$ ) concentration exceeded concentrations measured at the other fixed sites in the Yakima River Basin (fig. 6 and table 35 at back of report). Large concentrations of antimony are not unprecedented in streambed sediment in the Cle Elum River drainage. Of the 407

sites sampled for antimony in streambed sediment during the occurrence and distribution survey (Fuhrer, McKenzie, and others, 1994), the maximum concentration (4.8  $\mu\text{g/g}$ ) is in the Cle Elum River drainage—an area in which the U.S. Geological Survey (1989) reported the presence of pyrite (iron sulfide), galena (lead sulphate), and sphalerite (zinc-iron sulfide), all known sources of antimony (Levinson, 1980).

Concentrations of suspended antimony as high as 2.6, 3.1, and 2.3  $\mu\text{g/g}$  were measured in May 1989, January 1990, and February 1990, respectively, in the Yakima River at Cle Elum. The highest concentration (3.1  $\mu\text{g/g}$ ) of antimony corresponds to a major storm (January 9–12, 1990) in which U.S. Geological Survey personnel, during sampling at the Yakima River at Cle Elum, reported rain falling on 5 to 8 inches of snow. Two samplings were made during this major storm. The first sampling was at the onset of the storm (January 9, 1990) when streamflow (1,450 cubic feet per second [ $\text{ft}^3/\text{s}$ ]) was rapidly rising, and the suspended-sediment concentration (12  $\text{mg/L}$ ) was twice the median concentration measured during the 1987–90 period of study at Cle Elum. The second sampling was at the peak of the storm (January 10, 1990) when streamflow was 3,130  $\text{ft}^3/\text{s}$ , and the suspended-sediment concentration (130  $\text{mg/L}$ ) was the maximum concentration at Cle Elum for the period of study 1987–90. The concentration of antimony decreases from 3.1  $\mu\text{g/g}$  at the onset of the storm to 0.6  $\mu\text{g/g}$  at the peak of the storm (fig. 7). Similarly, the fraction of fine-grained-sized suspended sediment (the percentage of suspended sediment finer than 0.062 mm [millimeters] in diameter) decreased from about 90 percent at the onset of the storm to less than 50 percent at the peak of the storm. Even when the storm concentrations of antimony are normalized to the percentage of fine-grained-sized sediment, grain size alone does not account for the increase in antimony concentration at the onset of the storm event. The variability of antimony concentrations during the January storm probably indicates a change in the source of suspended sediment and the distribution of fine-grained-sized suspended sediment.

Because streambed sediment in the Cle Elum River drainage is enriched in antimony (Fuhrer, McKenzie, and others, 1994), storms may be a source of measurable suspended antimony in the Yakima River at Cle Elum. It is doubtful that all suspended sediment, transported from upstream of Cle Elum

**Table 12.** Summary of major- and trace-element concentrations in streambed sediment, Yakima River Basin, Washington, 1987–91

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per site was statistically summarized; concentrations of major elements are given as percentages; concentrations of trace elements are in units of micrograms per gram (µg/g), dry weight; data statistically summarized in this table are from Fuhrer and others (1992); element names and percentile concentrations shown in bold print are U.S. Environmental Protection Agency (1992c) Priority Pollutants; <, less than]

Element	Number of sites	Minimum value	Value at indicated percentile						Maximum value
			10	25	50	75	90	95	
Major Elements									
Aluminum	32	5.8	6.3	6.8	7.1	7.6	8.0	8.2	8.2
Calcium	32	.8	1.8	2.1	2.5	2.9	3.0	3.1	3.2
Carbon, inorganic	32	<.01	<.01	<.01	<.01	<.01	.1	.1	.1
Carbon, total	32	.4	.7	1.5	2.3	4.0	5.7	7.5	9.2
Iron	32	3.9	4.3	4.9	5.1	5.7	6.6	7.1	7.3
Magnesium	32	.7	.9	1.2	1.3	1.4	1.8	2.3	2.6
Phosphorus	32	.1	.1	.1	.1	.1	.1	.2	.2
Potassium	32	.7	.8	.9	1.0	1.3	1.4	1.4	1.5
Sodium	32	1.1	1.2	1.4	1.6	1.7	1.9	2.0	2.0
Sulfur	32	<.05	<.05	<.05	<.05	.1	1.2	3.3	5.3
Titanium	32	.3	.6	.6	.7	.8	1.0	1.0	1.1
Trace Elements									
Antimony <sup>1</sup>	17	.2	.3	.3	.4	.6	.9	1.4	1.4
Arsenic	32	1.1	1.5	2.4	3.7	5.7	12	35	45
Barium	32	380	393	447	480	530	563	584	590
Beryllium	32	1	1	1	1	2	2	2	3
Bismuth	32	<10	<10	<10	<10	<10	<10	<10	<10
Boron	16	.4	.4	.5	.8	1.4	3.6	4.0	4.0
Cadmium <sup>2</sup>	27	<2	<2	<2	<2	<2	<2	<2	<2
Cadmium <sup>3</sup>	5	.2	.2	.2	.2	.2	.5	.6	.8
Cerium	32	35	37	39	45	56	74	91	94
Chromium	32	21	44	53	62	79	170	210	210



**Table 12.** Summary of major- and trace-element concentrations in streambed sediment, Yakima River Basin, Washington, 1987–91—Continued

Element	Number of sites	Minimum value	Value at indicated percentile						Maximum value
			10	25	50	75	90	95	
Trace Elements									
Cobalt	32	14	16	19	20	23	30	32	33
Copper	32	17	21	25	30	42	70	94	96
Europium	32	<2	<2	<2	<2	<2	<2	<2	2
Gallium	32	15	15	17	18	19	20	20	21
Gold	32	<8	<8	<8	<8	<8	<8	<8	<8
Lanthanum	32	18	20	22	24	29	38	48	51
Lead	32	9	11	12	14	17	32	49	63
Lithium	32	17	19	21	24	28	35	45	45
Manganese	32	550	760	860	1,000	1,200	1,500	1,600	1,700
Mercury	32	<.02	<.02	<.02	.1	.2	.3	.5	.6
Molybdenum	32	<2	<2	<2	<2	<2	<2	<2	2
Neodymium	32	19	21	23	26	30	35	45	45
Nickel	32	9	17	20	27	37	82	190	260
Niobium	32	<4	<4	5	8	9	10	13	15
Scandium	32	13	15	18	20	21	25	27	29
Selenium	23	<.4	<.4	<.4	.4	.7	.9	1.0	1.0
Silver	32	<2	<2	<2	<2	<2	<2	<2	3
Strontium	32	140	210	230	260	290	310	340	350
Thorium	32	<4	4	5	6	7	10	12	15
Tin	32	<10	<10	<10	<10	<10	<10	<10	<10
Uranium <sup>4</sup>	24	<.05	.1	.7	1.1	1.4	1.9	3.0	3.3
Vanadium	32	84	113	130	140	180	220	230	24
Ytterbium	32	2	2	2	3	3	3	4	4
Yttrium	32	19	19	21	23	26	29	35	380
Zinc	32	77	84	95	100	110	170	200	210

<sup>1</sup>For antimony, two limits of determination (LD) exist (0.1 µg/g and 0.7 µg/g). Less-than values for the higher LD of 0.7 µg/g were not statistically summarized.

<sup>2</sup>Analyzed by inductively coupled plasma-atomic emission spectroscopy (ICP-AES); limit of determination = 2.0 µg/g.

<sup>3</sup>Analyzed by ICP-AES with organometallic-halide extraction; limit of determination = 0.05 µg/g.

<sup>4</sup>For uranium, two different methods of determination were used with different LDs (0.05 µg/g and 100 µg/g). Less-than values for the higher LD of 100 µg/g were not statistically summarized.

**Table 13.** Summary of major- and trace-element concentrations in suspended sediment, Yakima River Basin, Washington, 1987–90

[Data statistically summarized in this table are from sites 6, 19, 26, 32, 50, 52, and 56 (see table 1); to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; concentrations of major elements are given as percentages, except suspended organic carbon, which is in milligrams per liter (mg/L); concentrations of trace elements are in units of micrograms per gram (µg/g); element names and percentile concentrations shown in bold print are U.S. Environmental Protection Agency (1992c) Priority Pollutants; <,less than]

Element	Number of samples	Minimum value	Value at indicated percentile						Maximum value
			10	25	50	75	90	95	
Major Elements									
Aluminum	211	4.6	6.3	6.6	6.9	7.3	7.7	8.3	9.7
Calcium	211	1.2	1.8	2.0	2.2	2.5	2.9	3.0	3.5
Carbon, suspended organic	203	<.1	.2	.4	.6	1.3	2.5	2.9	4.9
Iron	211	3.8	4.7	4.9	5.2	5.5	5.7	5.9	8.1
Magnesium	210	.8	1.1	1.2	1.4	1.4	1.6	1.7	2.4
Phosphorus	211	.10	.13	.14	.16	.18	.21	.24	1.2
Potassium	211	.70	.90	1.0	1.2	1.4	1.6	1.6	2.4
Sodium	211	.8	1.0	1.2	1.3	1.4	1.6	1.7	2.4
Titanium	211	.4	.5	.5	.6	.6	.7	.7	.8
Trace Elements									
Antimony	211	.3	.5	.5	.6	.7	.8	.9	3.1
Arsenic	211	2.8	4.7	5.4	6.6	8.2	11	14	20
Beryllium	211	<2	<2	<2	<2	<2	2	2	3
Cadmium	211	<.1	.2	.3	.5	.7	1.4	1.7	32.6
Chromium	184	28	46	55	60	83	110	120	160
Cobalt	211	13	18	19	21	22	24	25	31
Copper	211	21	33	39	44	55	74	96	680
Lead	211	6	12	15	19	24	27	30	410
Manganese	211	910	1,200	1,400	1,900	2,900	3,500	4,000	6,300
Molybdenum	211	<.1	<.1	.6	.6	.8	1.1	1.4	3.0
Nickel	184	12	22	29	37	55	82	105	170
Silver	211	<.1	.2	.2	.4	.5	.9	1.3	7.7
Thallium	211	.1	.2	.3	.4	0.4	.5	.5	.6
Vanadium	211	101	121	131	142	149	160	166	193
Zinc	184	88.0	112	123	142	172	202	231	521

**Table 14. Summary of major- and trace-element concentrations in filtered-water samples, Yakima River Basin, Washington, 1987–90**

[The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate, only one element concentration per visit was statistically summarized; data statistically summarized in this table are from sites 6, 19, 26, 32, 50, 52, and 56 (see table 1); all concentrations listed below are in micrograms per liter (µg/L), except organic carbon, hardness, and bromide, which are given in milligrams per liter (mg/L); element names and percentile concentrations shown in bold print are U.S. Environmental Protection Agency (1992c) Priority Pollutants; for cadmium, chromium, copper, and lead, only samples analyzed by atomic absorption spectroscopy with graphite furnace (AAGF) were statistically summarized; <, less than]

Element	Number of samples	Minimum value	Value at indicated percentile						Maximum value
			10	25	50	75	90	95	
Major Elements									
Aluminum	27	<10	<10	<10	10	20	50	150	210
Carbon, organic	275	.4	1.2	1.5	2.0	2.6	3.5	4.4	8.0
Hardness	292	18	25	34	54	95	120	250	270
Iron	36	8	13	18	28	39	57	101	250
Trace Elements									
Antimony	18	<1	<1	<1	<1	<1	<1	1	1
Arsenic	106	<1	<1	<1	<1	2	3	7	9
Barium	36	<2	3	6	10	26	40	74	79
Beryllium	36	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
Boron	23	<10	<10	10	20	20	40	40	40
Bromide	19	<.01	<.01	<.01	<.01	<.01	.02	.08	.08
Cadmium	279	<.2	<.2	<.2	<.2	<.2	.3	.5	2.2
Chromium	26	<.5	<.5	<.5	<.5	.6	1.0	1.1	1.1
Cobalt	36	<3	<3	<3	<3	<3	<3	<3	<3
Copper	280	<.5	<.5	.6	.9	1.3	1.9	3.0	20
Lead	279	<.5	<.5	<.5	<.5	<.5	<.5	.6	1.9
Lithium	36	<4	<4	<4	<4	<4	5	8	16
Manganese	36	<1	2	3	7	16	40	87	110
Mercury	283	<.1	<.1	<.1	<.1	<.1	<.1	.1	.6
Molybdenum	36	<10	<10	<10	<10	<10	<10	<10	<10
Nickel	36	<10	<10	<10	<10	<10	<10	<10	<10
Selenium	22	<1	<1	<1	<1	<1	<1	2	2
Silver	36	<1	<1	<1	<1	<1	1	2	2
Strontium	36	21	30	46	68	130	160	330	330
Vanadium	36	<6	<6	<6	<6	8	10	21	22
Zinc	36	<3	<3	<3	5	12	18	29	30

**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, the mean element concentration of each site was statistically summarized; concentrations are reported in units of micrograms per gram (µg/g), dry weight; livers from bridgelip sucker, carp, largescale sucker, mountain whitefish, and rainbow trout were analyzed from composites of usually 10 samples; whole bodies of sculpin, caddisflies, and stoneflies were analyzed from composites of usually 10 fish and 50 or more insects; soft parts of the Asiatic clam were analyzed from composites of usually 20 samples; curlyleaf-pondweed and waterweed samples consisted of a minimum of 5 grams of mass; only 1990 data are summarized for largescale suckers, *Arctopsyche* spp., *Hydropsyche* spp., and lead in Asiatic clams; Asiatic clams were of the order Veneroida and family Corbiculidae; organism taxa are listed as follows: common name

(*Genus species*)

Sample	Number of sites	Minimum	Median	Maximum
<b>Arsenic</b>				
Bridgelip sucker ( <i>Catostomus columbianus</i> )	3	0.20	0.85	1.2
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	.10	.30	.50
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	<.30	<.30	.40
Sculpin ( <i>Cottus</i> spp.)	12	<.10	.15	.37
Asiatic clam ( <i>Corbicula fluminea</i> )	5	3.6	4.1	5.2
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	.48	.79	1.5
Waterweed ( <i>Elodea</i> sp.)	4	1.0	1.6	2.6
<b>Cadmium</b>				
Carp ( <i>Cyprinus carpio</i> )	3	.46	.79	2.5
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	.03	.35	.43
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	<.20	<.20	1.2
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	<.40	<.40	1.0
Asiatic clam ( <i>Corbicula fluminea</i> )	5	<.20	.24	.38
Caddisfly ( <i>Arctopsyche</i> sp.)	12	<.41	<.41	.43
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	<.19	<.19	.19
Caddisfly ( <i>Hydropsyche</i> spp.)	24	<.18	<.18	.25
Stonefly ( <i>Calineuria</i> spp.)	6	<.14	<.14	.45
Stonefly ( <i>Claassenia</i> sp.)	7	<.15	.16	.40
Stonefly ( <i>Doroneuria</i> spp.)	5	<.15	<.15	.50
Stonefly ( <i>Hesperoperla</i> sp.)	10	<.12	<.12	.33

**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91—Continued

Sample	Number of sites	Minimum	Median	Maximum
<b>Cadmium—Continued</b>				
Stonefly ( <i>Megarcys</i> spp.)	3	.18	.24	.36
Stonefly ( <i>Perlinodes</i> spp.)	3	.12	.17	.25
Stonefly ( <i>Pteronarcys</i> spp.)	8	<.41	<.41	<.41
Stonefly ( <i>Skwala</i> spp.)	9	<.23	<.23	.23
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	<.80	<.80	.91
<b>Chromium</b>				
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	.50	.59	.78
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	<1.0	<1.0	1.0
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	<4.0	<4.0	<4.0
Asiatic clam ( <i>Corbicula fluminea</i> )	5	1.0	1.4	2.0
Caddisfly ( <i>Arctopsyche</i> sp.)	12	.71	1.6	10
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	2.3	4.3	5.9
Caddisfly ( <i>Hydropsyche</i> spp.)	24	.66	2.4	3.8
Stonefly ( <i>Calineuria</i> spp.)	6	<.11	1.1	3.9
Stonefly ( <i>Claassenia</i> sp.)	7	.49	1.4	2.2
Stonefly ( <i>Doroneuria</i> spp.)	5	.95	1.3	2.6
Stonefly ( <i>Hesperoperla</i> sp.)	10	.61	1.3	2.4
Stonefly ( <i>Megarcys</i> spp.)	3	.44	2.2	5.2
Stonefly ( <i>Perlinodes</i> spp.)	3	2.2	3.0	34
Stonefly ( <i>Pteronarcys</i> spp.)	8	.58	1.2	3.3
Stonefly ( <i>Skwala</i> spp.)	9	.31	1.8	16
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	2.0	3.0	4.0
Waterweed ( <i>Elodea</i> sp.)	4	3.8	6.4	8.3



**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91—Continued

Sample	Number of sites	Minimum	Median	Maximum
<b>Cobalt</b>				
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	0.12	0.31	0.50
Mountain whitefish ( <i>Prosopium williamsoni</i> )	6	.38	.46	.84
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	5	.18	.32	.46
Asiatic clam ( <i>Corbicula fluminea</i> )	4	.50	.58	1.1
Caddisfly ( <i>Arctopsyche</i> sp.)	12	.71	1.1	5.0
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	1.8	2.5	5.7
Caddisfly ( <i>Hydropsyche</i> spp.)	24	.75	2.7	9.1
Stonefly ( <i>Calineuria</i> spp.)	6	.37	.68	.86
Stonefly ( <i>Claassenia</i> sp.)	7	.39	.57	1.0
Stonefly ( <i>Doroneuria</i> spp.)	5	.20	.49	1.7
Stonefly ( <i>Hesperoperla</i> sp.)	10	.25	.43	1.8
Stonefly ( <i>Megarcys</i> spp.)	3	.58	.88	.97
Stonefly ( <i>Perlinodes</i> spp.)	3	1.8	2.2	5.7
Stonefly ( <i>Pteronarcys</i> spp.)	8	.74	.82	4.3
Stonefly ( <i>Skwala</i> spp.)	9	.89	1.6	3.0
<b>Copper</b>				
Bridgelip sucker ( <i>Catostomus columbianus</i> )	3	7.7	14	19
Carp ( <i>Cyprinus carpio</i> )	3	28	55	100
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	23	26	32
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	5.6	6.4	11
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	18	91	480
Asiatic clam ( <i>Corbicula fluminea</i> )	5	25	28	34

**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91—Continued

Sample	Number of sites	Minimum	Median	Maximum
<b>Copper—Continued</b>				
Caddisfly ( <i>Arctopsyche</i> sp.)	12	5.9	9.8	15
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	6.5	12	19
Caddisfly ( <i>Hydropsyche</i> spp.)	24	9.2	13	21
Stonefly ( <i>Calineuria</i> spp.)	6	18	22	24
Stonefly ( <i>Claassenia</i> sp.)	7	27	32	38
Stonefly ( <i>Doroneuria</i> spp.)	5	25	30	38
Stonefly ( <i>Hesperoperla</i> sp.)	10	18	24	28
Stonefly ( <i>Megarcys</i> spp.)	3	11	26	36
Stonefly ( <i>Perlinodes</i> spp.)	3	14	15	22
Stonefly ( <i>Pteronarcys</i> spp.)	8	8.0	21	32
Stonefly ( <i>Skwala</i> spp.)	9	14	19	26
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	9.2	11	22
Waterweed ( <i>Elodea</i> sp.)	4	13	19	65
<b>Lead</b>				
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	<.12	.18	.29
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	<4.0	<4.0	<4.0
Asiatic clam ( <i>Corbicula fluminea</i> )	4	.18	.31	.40
Caddisfly ( <i>Arctopsyche</i> sp.)	12	<2.1	<2.1	24
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	1.3	2.1	3.2
Caddisfly ( <i>Hydropsyche</i> spp.)	24	<.96	1.1	5.6
Stonefly ( <i>Claassenia</i> sp.)	7	<.68	<.68	1.8

**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91—Continued


Sample	Number of sites	Minimum	Median	Maximum
<b>Lead—Continued</b>				
Stonefly ( <i>Doroneuria</i> spp.)	5	<0.59	<0.59	0.65
Stonefly ( <i>Hesperoperla</i> sp.)	10	<.65	<.65	.92
Stonefly ( <i>Perlinodes</i> spp.)	3	1.8	2.8	3.3
Stonefly ( <i>Pteronarcys</i> spp.)	8	<2.1	<2.1	<2.1
Stonefly ( <i>Skwala</i> spp.)	9	<.80	<.80	2.8
<b>Mercury</b>				
Bridgelip sucker ( <i>Catostomus columbianus</i> )	3	.05	.06	.06
Carp ( <i>Cyprinus carpio</i> )	3	.30	.38	.42
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	.05	.32	.47
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	.40	.81	1.3
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	.12	.26	.35
Sculpin ( <i>Cottus</i> spp.)	12	.09	.19	.31
Asiatic clam ( <i>Corbicula fluminea</i> )	5	.10	.16	.17
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	.03	.05	.06
Waterweed ( <i>Elodea</i> sp.)	4	.06	.07	.09
<b>Nickel</b>				
Asiatic clam ( <i>Corbicula fluminea</i> )	5	<2.0	<2.0	<2.0
Caddisfly ( <i>Arctopsyche</i> sp.)	12	<1.0	<1.0	42
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	1.5	5.1	7.3
Caddisfly ( <i>Hydropsyche</i> spp.)	24	.58	2.4	5.5
Stonefly ( <i>Calineuria</i> spp.)	6	.28	.48	6.9
Stonefly ( <i>Claassenia</i> sp.)	7	<.38	<.38	6.4
Stonefly ( <i>Doroneuria</i> spp.)	5	<.15	1.4	1.6
Stonefly ( <i>Hesperoperla</i> sp.)	10	<.26	.43	7.1

**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91—Continued

Sample	Number of sites	Minimum	Median	Maximum
<b>Nickel—Continued</b>				
Stonefly ( <i>Megarcys</i> spp.)	3	0.26	2.0	2.4
Stonefly ( <i>Perlinodes</i> spp.)	3	2.2	2.3	76
Stonefly ( <i>Pteronarcys</i> spp.)	8	.74	1.2	7.5
Stonefly ( <i>Skwala</i> spp.)	9	.38	.86	34
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	3.0	5.8	20
Waterweed ( <i>Elodea</i> sp.)	4	8.5	14	23
<b>Selenium</b>				
Bridgelip sucker ( <i>Catostomus columbianus</i> )	3	1.9	2.0	5.2
Carp ( <i>Cyprinus carpio</i> )	3	2.2	3.8	4.2
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	1.9	3.5	4.8
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	4.2	5.0	13
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	2.2	7.0	31
Sculpin ( <i>Cottus</i> spp.)	12	.20	1.6	5.4
Asiatic clam ( <i>Corbicula fluminea</i> )	5	2.1	2.4	3.0
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	.20	.36	.70
Waterweed ( <i>Elodea</i> sp.)	4	.30	.59	1.2
<b>Silver</b>				
Carp ( <i>Cyprinus carpio</i> )	3	.24	.45	3.0
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	.06	.10	.14
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	<2.0	<2.0	<2.0
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	<7.0	<7.0	20
Stonefly ( <i>Calineuria</i> spp.)	6	<.09	.13	.21
Stonefly ( <i>Claassenia</i> sp.)	7	.13	.15	.30
Stonefly ( <i>Doroneuria</i> spp.)	5	<.07	.12	.30
Stonefly ( <i>Hesperoperla</i> sp.)	10	<.08	.08	.22

**Table 15.** Summary of selected trace-element concentrations in aquatic biota, Yakima River Basin, Washington, 1989–91—Continued

Sample	Number of sites	Minimum	Median	Maximum
Zinc				
Bridgelip sucker ( <i>Catostomus columbianus</i> )	3	55	56	94
Carp ( <i>Cyprinus carpio</i> )	3	160	634	890
Largescale sucker ( <i>Catostomus macrocheilus</i> )	6	60	81	102
Mountain whitefish ( <i>Prosopium williamsoni</i> )	7	57	72	79
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	10	75	99	226
Asiatic clam ( <i>Corbicula fluminea</i> )	5	96	108	452
Caddisfly ( <i>Arctopsyche</i> sp.)	12	96	141	192
Caddisfly ( <i>Cheumatopsyche</i> spp.)	4	81	96	102
Caddisfly ( <i>Hydropsyche</i> spp.)	24	67	105	152
Stonefly ( <i>Calineuria</i> spp.)	6	142	196	251
Stonefly ( <i>Claassenia</i> sp.)	7	174	218	352
Stonefly ( <i>Doroneuria</i> spp.)	5	216	231	254
Stonefly ( <i>Hesperoperla</i> sp.)	10	276	372	450
Stonefly ( <i>Megarcys</i> spp.)	3	127	256	271
Stonefly ( <i>Perlinodes</i> spp.)	3	84	106	141
Stonefly ( <i>Pteronarcys</i> spp.)	8	114	128	150
Stonefly ( <i>Skwala</i> spp.)	9	102	138	314
Curlyleaf pondweed ( <i>Potamogeton crispus</i> )	8	50	76	187
Waterweed ( <i>Elodea</i> sp.)	4	44	130	239



#### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.

- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value
- × 1.5 to 3 times the interquartile range from the 25-percentile value


**Figure 6.** Distribution of antimony concentrations in suspended sediment at fixed sites in the Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).

Lake, enters the Yakima River at Cle Elum. At times, Cle Elum Lake may act as a sieve, by passing small-grained-sized suspended sediment and trapping larger sediment particles. Contributions of suspended sediment, from areas less enriched than the Cle Elum River drainage, would dilute the concentration of suspended antimony in the Yakima River at Cle Elum. One possible source of less-enriched antimony is Crystal Creek (formerly Roslyn Creek) which flows

into the main stem 0.15 miles upstream from the Yakima River at Cle Elum sample site. Crystal Creek receives urban runoff from the towns of Cle Elum and Roslyn, in addition to runoff from numerous open-pit coal mines that are located east of Roslyn. During the January storm, Crystal Creek was creating a turbidity plume which, during the rising limb of the storm, obscured the shallow river bottom near the left bank of the main stem at Cle Elum. The Crystal Creek drainage, which is located in the nonmarine rock geologic unit, contains concentrations of antimony (0.5 µg/g and 0.8 µg/g) in streambed sediment that were small in comparison to the large concentration (4.8 µg/g) in streambed sediment in the Cle Elum River drainage (Ryder and others, 1992). During large storms, therefore, Crystal Creek may be diluting the concentration of suspended antimony at Cle Elum because of Crystal Creek's close proximity to the Yakima River at Cle Elum sampling site.

The peak of a rain-on-snow storm at the Yakima River at Cle Elum was sampled on April 6, 1989. Similar to the January 1990 storm, streamflow near the peak of the storm was large (2,150 ft<sup>3</sup>/s) and the concentration of suspended sediment also was large (85 mg/L). Similar to the sample collected near the peak of the storm in January 1990, the concentration of antimony near the peak of the April 1989 storm was only 0.6 µg/g (fig. 7). These two storm samplings, in addition to a third storm-event sampling (May 9, 1989), indicated that concentrations of suspended antimony are diluted during large storms, presumably by less enriched concentrations of antimony from drainages such as Crystal Creek. What remains unclear, however, is the transport potential of antimony to the main stem from sources in the Cle Elum River Drainage.

On the basis of a limited number of filtered-water determinations at fixed sites, antimony was measured in the Yakima River at Umtanum and in the Yakima River above Ahtanum Creek at Union Gap at the limit of determination (1 µg/L) in 2 of 22 samples. Although antimony was detected at the limit of determination (1 µg/L), antimony concentrations were high in comparison to background concentrations in natural water. Antimony measured at the Umtanum and Union Gap sites were 10 times greater than background concentrations in other basins that are minimally affected by human activities (0.1 µg/L; Forstner and Wittmann, 1979) [table 36 at back of report].




**Figure 7.** Streamflow and antimony concentrations in suspended sediment for selected time periods for the Yakima River at Cle Elum, Yakima River Basin, Washington, 1988–90 (snowmelt season occurs April through May; non-irrigation season occurs October through March).

## Arsenic and Lead

Concentrations of arsenic in streambed sediment at the 32 biological sampling sites ranged from 1.1 to 45 µg/g (table 12). Concentrations of arsenic as high as 45 µg/g and 29 µg/g were measured in streambed sediment of the Wapatus River at mouth near Roslyn (site 1) and Jungle Creek near mouth (site 3), respectively, which are probably from geologic sources. Streambed sediment in the Wapatus River drainage in the Wenatchee Mountains is formed from pre-Tertiary metamorphic and intrusive rocks of the Cascade Range geologic province. Streambed sediment concentrations of arsenic as large as 31 µg/g in this pre-Tertiary rock geologic unit were reported by Fuhrer and others (1994a) and were attributed to the presence of arsenopyrite. Streambed sediment in the Jungle Creek drainage is formed from nonmarine sedimentary rock of the Cascade Range geologic province. Concentrations of arsenic as high as 21 µg/g were reported in streambed sediment of the nonmarine rock geologic unit (Fuhrer, McKenzie, and others, 1994) which contains hydrothermally altered volcanic and granitic rock with concentrations of arsenic as large as 7,000 µg/g (Simmons and others, 1983; Gualtieri and Simmons, 1989).

Concentrations of arsenic in main-stem tributaries are as much as 16 times larger than concentrations in the main stem. The largest difference in arsenic concentration was between the Wapatus River at mouth near Roslyn (site 1; 45 µg/g) and the Yakima River at Umtanum (site 19; 2.8 µg/g) located 6.6 miles downstream of irrigation return flows entering the main stem from the Kittitas Valley (fig. 8). Arsenic-depleted sediment within the Kittitas Valley was probably responsible for the measurably lower arsenic concentrations in the Yakima River at Umtanum (Fuhrer, McKenzie, and others, 1994). Concentrations of arsenic as high as 13.0 µg/g and 9.8 µg/g were measured in the American River at Hell's Crossing and in Rattlesnake Creek above North Fork Rattlesnake Creek, respectively. Both streams are tributaries to the Naches River, and the presence of arsenic in these streams probably was the result of geologic sources. Although, arsenic concentrations at the mouths of many streams carrying irrigation return flow to the main stem are lower than in streams affected by geologic sources, some streams that receive irrigation return flow show evidence of arsenic enrichment. For example, arsenic concentrations increased more than 4 times in a downstream manner in Satus Creek. Arsenic concentrations increased from






**Figure 8.** Arsenic concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987–90 (“Waptus River” represents Waptus River at mouth near Roslyn; “Jungle Creek” represents Jungle Creek near mouth near Cle Elum; “Rattlesnake Creek above North Fork” represents Rattlesnake Creek above North Fork Rattlesnake Creek near Nile; “Rattlesnake Creek” represents Rattlesnake Creek above Little Rattlesnake Creek near Nile; “American River” represents American River at Hells Crossing near Nile; “Satus Creek at gage” represents Satus Creek at gage at Satus; “Satus Creek below Dry Creek” represents Satus Creek below Dry Creek near Toppenish; and “Satus Creek” represents Satus Creek above Wilson-Charley Canyon near Toppenish).

1.3 to 2.8 to 5.8 µg/g at Satus Creek sites above Wilson-Charley Canyon (site 57), below Dry Creek (site 53), and at Gage at Satus (site 47), respectively.

Potential sources of arsenic in agricultural land-use areas include the historical application of pesticides containing lead arsenate. In eastern Washington, beginning in 1908, lead-arsenate formulations were applied to control codling moth in apples. This practice continued until the introduction of dichlorodiphenyltrichloroethane (DDT) in 1947. Significant amounts of this pesticide have been applied and detected in the basin’s agricultural soils; arsenic concentrations as large as 140 µg/g were found in a former Yakima River Basin apple orchard (Fuhrer, McKenzie, and others, 1994). Using available land-use information, which does not separate agricultural land into orchard and non-orchard categories, and a spatially large sampling of Yakima River Basin streambed sediment, Fuhrer, McKenzie, and others, 1994, have shown that lands classified as agricultural had significantly larger ( $p \leq 0.05$ ) arsenic

concentrations than lands classified as nonagricultural (fig. 9). The enrichment of arsenic occurred in agricultural land-use areas residing in the Quaternary deposits and loess geologic unit and the nonmarine sedimentary rock geologic unit—geologic hosts to many of the basin’s apple orchards. The enrichment of arsenic in agricultural areas may result from a legacy of pesticide use.

Concentrations of lead in streambed sediment of the Yakima River Basin ranged from 9.0 to 63.0 µg/g (table 12) and slightly exceed the 5.2 to 55 µg/g range of concentration that encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). The median concentration of lead at the biological sampling sites (14 µg/g) is nearly one-half that in fine-fraction streambed sediment in the highly urbanized Upper Illinois River Basin (table 34, at back of report). Lead concentrations as large as 63 µg/g, 56 µg/g, and 36 µg/g, however, were measured in Wide Hollow



**Figure 9.** Comparison of arsenic concentrations between agricultural and nonagricultural land uses for streambed sediment of lower-order streams in the Quaternary deposits and loess geologic unit and in the nonmarine sedimentary rocks geologic unit, Yakima River Basin, Washington, 1987.


Creek at West Valley Middle School near Ahtanum (site 27), Wide Hollow Creek at old sewage treatment plant at Union Gap (site 29), and Naches River near north Yakima (site 26), respectively.

Lead in Wide Hollow Creek Subbasin probably results from urbanization, agricultural practices, and light industrial, rather than from geologic sources. Geologic sources of lead in Wide Hollow Creek Subbasin are few and consist of the Quaternary deposits and loess geologic unit, which contains a median lead concentration of only 14 µg/g (Fuhrer, McKenzie, and others, 1994). The 56 µg/g concentration of lead at Wide Hollow Creek at the old

Sewage Treatment Plant—located near the mouth of Wide Hollow Creek—is not unexpected considering that Wide Hollow Creek drains the urban area of Union Gap and that the median concentration of lead in sediment from urban drains of the Yakima River Basin is 71 µg/g (Fuhrer, McKenzie, and others, 1994). What is unexpected, however, is the 63 µg/g lead at the upstream site, Wide Hollow Creek at West Valley Middle School near Ahtanum, which is farther removed from the urban effects of Union Gap. In addition to urban sources, sources of lead enrichment in Wide Hollow Creek probably result from the legacy of lead-arsenate usage for control of codling moth in apple orchards. The application of lead arsenate, from 1908 to 1947, in eastern Washington increased from 50 lb (pounds) of lead and 18 lb of arsenic to 192 lb of lead and 71 lb of arsenic per acre (Peryea, 1989). A considerable reservoir of lead and arsenic may exist in the Yakima River Basin considering that 3,000 acres of apple orchards existed (primarily in the middle and lower basin) prior to 1955 (U.S. Department of Agriculture, 1986). Concentrations of lead in soils of former apple orchards, located in an adjacent subbasin, were as large as 890 µg/g—a concentration approximately 60 times that associated with geologic sources in the Ahtanum Subbasin (Fuhrer, McKenzie, and others, 1994). Lead also was reported in streambed sediment of Wide Hollow Creek in an earlier study by Washington State Department of Ecology (see Previous Studies section of this report).

Concentrations of suspended arsenic at the seven fixed sites ranged from 2.8 to 20 µg/g (table 13). Arsenic was enriched slightly (median 7.0 µg/g) in the Yakima River at Cle Elum and notably (median 10 µg/g) in Sulphur Creek Wasteway near Sunnyside (fig. 10; table 35, at back of report). The highest concentrations of arsenic were found in the lower Yakima Valley where median-arsenic concentrations ranged from 7.0 to 10 µg/g.

Concentrations of suspended arsenic in the Yakima River at Cle Elum ranged from 6.3 µg/g to 8.8 µg/g. Although these concentrations of arsenic were not high in comparison to Sulphur Creek Wasteway near Sunnyside, these concentrations probably are indicative of an upstream source of enrichment. Similar to antimony, arsenic enrichment from geologic sources is not uncommon in the Cle Elum Subbasin. As discussed previously, high arsenic concentrations reside in streambed sediment and rock in the pre-Tertiary and metamorphic rocks geologic unit which



### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.

- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value

**Figure 10.** Distribution of arsenic concentrations in suspended sediment at fixed sites in the Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).

contains arsenopyrite (Gualtieri and Simmons, 1989; Fuhrer, McKenzie, and others, 1994).


During storms, higher arsenic concentrations were found preceding the storm's peak than at the peak. During the peak of the January 9, 1990 storm at Cle Elum (rain-on-snow), the suspended arsenic concentration was 6.3 µg/g; preceding the storm's peak however, the suspended arsenic concentration was 7.6 µg/g. This variation, although low in comparison to antimony, represents approximately

one-half the range of concentrations measured over the 1987–90 monthly sampling period for Cle Elum.

The highest concentrations of suspended arsenic were found in Sulphur Creek Wasteway near Sunnyside (site 52). The median concentration was 10 µg/g (exceeding those of other fixed sites) and overall, arsenic concentrations range from 4.9 µg/g to 20 µg/g. Concentrations of suspended arsenic in Sulphur Creek Wasteway also varied seasonally to a greater extent than at the other fixed sites; the interquartile range for suspended arsenic in Sulphur Creek Wasteway was 7.4 µg/g in comparison to just 1.9 µg/g in the Yakima River at Kiona (site 50). Differences in the interquartile ranges among sampling sites probably were related to the size of the waterway and to the principal water use in the drainage. Sulphur Creek Wasteway, located in the Sunnyside Subbasin, is different from other fixed sites because it is a major conveyance of agricultural return flow, as well as a spillway for excess canal water during the irrigation season. Large variations in constituent concentrations, similar to those measured for arsenic, may not be unusual in Sulphur Creek Wasteway or in other waterways that principally receive irrigation return flow. Conversely, sites located along the main stem or on larger tributaries probably receive adequate dilution to dampen large fluxes in constituent concentrations from smaller waterways carrying agricultural return flow to the main stem.

During the irrigation season, concentrations of suspended arsenic in Sulphur Creek Wasteway near Sunnyside are significantly lower ( $p \leq 0.001$ ) than during the non-irrigation season (fig. 11). These concentrations also are in close agreement with arsenic concentrations measured in streambed sediment (finer than 62 µm, in diameter) of Sunnyside Subbasin (Fuhrer, McKenzie, and others, 1994). The median concentration of suspended arsenic during the irrigation season was 7.0 µg/g, and the median arsenic concentration of streambed sediment was 8.2 µg/g. The similarity between suspended- and streambed-sediment concentrations of arsenic, in addition to high concentrations of suspended sediment during the irrigation season, indicate the significance of agricultural soil erosion as a dampening mechanism which, during the irrigation season, limits concentrations of suspended arsenic in Sulphur Creek (fig. 11).


Conversely, during the non-irrigation season, the median concentration of suspended arsenic is 13.9 µg/g at Sulphur Creek Wasteway near Sunnyside.



**Figure 11.** Concentrations of suspended sediment and arsenic in suspended sediment at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987-90 (shaded area represents the irrigation season).

The higher concentrations of suspended arsenic, during the non-irrigation season, coincide with increases in the percentage of fine-grained suspended sediment and increases in the concentration of suspended-organic carbon (fig. 12). As a result, the higher arsenic concentrations probably are not from new or different sources, but rather from a change in the distribution of fine-grained-sized sediment and organic carbon content.


Differences in the concentrations of suspended arsenic between the irrigation and the non-irrigation seasons probably result from variations in the quantity of fine-grained suspended sediment and suspended organic carbon. Smaller grain-sized sediment contains larger element concentrations because of the increase in sediment surface area; larger grain-sized sediment,



**Figure 12.** Distribution of suspended sediment finer than 62 micrometers in diameter and suspended organic carbon concentrations at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987-90 (shaded area represents the irrigation season).

however, have comparatively small surface areas and low element concentrations (Horowitz, 1991, p. 32). The variations in suspended arsenic concentrations in Sulphur Creek Wasteway have a distinct pattern which is statistically ( $p \leq 0.001$ ) correlated to the percentage of suspended sediment finer than 62  $\mu\text{m}$ , in diameter (fig. 13). A similar relation and significant correlation exists between suspended-arsenic concentrations and suspended-organic carbon (fig. 13). Thus, the concentrations of suspended arsenic in Sulphur Creek Wasteway increased in proportion to the quantity of fine-grained suspended sediment and organic carbon.

High concentrations of suspended arsenic in Sulphur Creek Wasteway near Sunnyside may be a legacy



**Figure 13.** Percentage of suspended sediment finer than 62 micrometers in diameter and the concentration of arsenic in suspended sediment and the concentration of suspended organic carbon at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90.

of lead-arsenate formulations applied to apple orchards in the Sunnyside Subbasin. In addition to the Sunnyside Subbasin, other agricultural subbasins (although not measured in this study), also may be sources of suspended arsenic. Present day applications of phosphate fertilizer may leach arsenic from soils to shallow ground water (Peryea, 1989). Arsenic, therefore, may not occur only in streambed sediment from the erosion of soil treated with lead arsenate, but arsenic also may be discharged to waterways (in dissolved form) from shallow ground water. Once dissolved, arsenic subsequently can attach to streambed

sediment by adsorption or coprecipitation reactions with hydrous iron oxides (Hem, 1989, p. 144).

The significant contribution of Sulphur Creek Wasteway to the suspended-arsenic load in the main stem is perhaps more important than the relation that exists in Sulphur Creek Wasteway between suspended-arsenic concentrations and the physical and chemical measurements described earlier. During the irrigation season, Sulphur Creek had a suspended-arsenic load two-thirds that of the Yakima River at Kiona (table 16) but accounted for less than 20 percent of the streamflow at Kiona. The suspended-arsenic loads in Sulphur Creek also vary seasonally (fig. 14). In 1989, for example, the irrigation-season load of arsenic (1.5 pounds per day) was nearly 4 times higher than the non-irrigation season load (0.4 pounds per day). This increase was not caused directly by high concentrations of suspended arsenic, but rather by agricultural soil erosion that results in high suspended-sediment concentrations in Sulphur Creek Wasteway. As noted earlier, the concentration of arsenic in suspended sediment during the irrigation season was not large relative to that measured in streambed sediment of the Sunnyside Subbasin. Concentrations of suspended sediment were high; however, during the 1989 irrigation season, the average suspended-sediment concentration (162 mg/L) in Sulphur Creek was about 4 times higher than that measured during the non-irrigation season. Thus, large concentrations of suspended sediment, and not of suspended arsenic, result in high irrigation-season loads of suspended arsenic in Sulphur Creek. Farm practices that tend to reduce erosion also will reduce the suspended-arsenic load of Sulphur Creek Wasteway to the main stem.


The irrigation-season load of arsenic for the Yakima River at Umtanum, like Sulphur Creek Wasteway near Sunnyside, also is larger than the non-irrigation season load. In 1989, the irrigation-season load of arsenic (2.2 pounds per day) was more than three times the non-irrigation season load. The large irrigation-season load probably results from agricultural soil erosion in the Kittitas Valley, rather than from an increase in the concentration of suspended arsenic at Umtanum. In July 1988, Wilson Creek (RM 147) was discharging 83 mg/L of suspended sediment to the main stem above Umtanum (Embrey, 1992). Several other irrigation drains also discharge to the main stem, upstream of the Umtanum site (Bureau of Reclamation and U.S. Soil

**Table 16. Estimated arsenic loads in suspended sediment at selected fixed sites, Yakima River Basin, Washington, 1987–90**

[Loads reported as pounds per day; load estimates are based on calibration data collected from March 1987 to March 1990; --, indicates insufficient data; bold lines represent the irrigation season, and lightly shaded cells represent the snowmelt portion of the irrigation season; non-irrigation season, October through March]

Year	January	February	March	April	May	June	July	August	September	October	November	December	Daily mean
<b>Yakima River at Umtanum</b>													
1987	0.2	0.4	2	2	1	2	2	2	0.4	0.1	0.1	0.2	1.0
1988	.1	.5	.6	2	.9	1	3	3	1	.2	.4	.6	1.1
1989	.8	.6	.8	4	3	2	3	3	.9	.4	.6	.7	1.7
1990	1	2	2	5	3	5	3	3	.8	--	--	--	--
<b>Naches River near North Yakima</b>													
1987	--	--	1	2	9	.5	.1	<.05	.5	.3	<.05	.2	--
1988	.1	.2	.3	2	2	1	.2	.1	1	.6	.2	.3	.7
1989	.3	.1	.4	4	3	2	.1	.1	1	.3	.1	.6	1.0
1990	2	.9	.9	--	--	--	--	--	--	--	--	--	--
<b>Yakima River above Ahtanum Creek at Union Gap</b>													
1987	1	2	5	5	8	4	4	4	2	1	.4	1	3.2
1988	.7	2	2	6	4	4	4	5	3	2	2	2	3.0
1989	2	2	3	10	8	6	5	4	3	2	2	2	4.1
1990	3	3	4	10	7	11	5	5	5	4	--	--	--
<b>Sulphur Creek Wasteway near Sunnyside</b>													
1987	--	--	1	3	3	2	2	.9	.6	.3	.1	.2	--
1988	.2	.3	2	2	2	2	1	.9	.9	.4	.1	.1	1.0
1989	.2	.2	1	4	4	2	2	1	1	.6	.2	.2	1.4
1990	.2	.2	2	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Euclid Bridge at river mile 55 near Grandview</b>													
1987	--	--	7	6	10	2	2	2	1	1	.9	2	--
1988	1	2	2	6	4	3	2	2	2	2	2	2	2.5
1989	2	2	5	16	11	4	2	2	2	2	3	3	4.5
1990	4	4	5	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Kiona</b>													
1987	.5	1	12	4	18	1	1	1	.7	1	.5	1	3.5
1988	.5	1	.9	8	4	3	1	1	2	1	2	1	2.2
1989	.9	.7	3	30	17	3	2	2	2	2	3	2	5.6
1990	4	2	3	18	9	50	2	8	3	--	--	--	--





**Figure 14.** Percentages of the annual load of arsenic in suspended sediment for selected time periods at Sulphur Creek Wasteway near Sunnyside, the Yakima River at Umtanum, and the Yakima River at Kiona, Yakima River Basin, Washington, 1989 (non-irrigation season occurs October through March; snowmelt season occurs April through May; irrigation season occurs June through September).

Conservation Service, 1974); although not sampled in this study, these irrigation drains also may contribute to the suspended-arsenic load at Umtanum.

The dominant season for transport of suspended arsenic at most sites, however, is snowmelt. At Kiona, for example, the quantity of arsenic transported during the snowmelt season of 1989 (about 1,400 pounds) was 5 times larger than that transported during the irrigation season, and the load of arsenic during the snowmelt season (24 pounds per day) is more than 10 times larger than the irrigation-season load. Additionally, more than one-half the 1989 annual load of suspended arsenic at Kiona was transported during the snowmelt season (fig.14). Similar transport patterns were measured for the Yakima River at Euclid Bridge at RM 55 near Grandview and at the Naches River near North Yakima. For the Yakima River at Umtanum, however, the 1989 irrigation-season load was only 1.3 times larger than the snowmelt-season load. The smaller load of snowmelt at Umtanum indicates the capture of snowmelt runoff by irrigation-storage reservoirs in the Kittitas Valley. During the irrigation season, waters released from reservoirs in the Kittitas Valley are routed down the main stem and, at various points, diverted for irrigation. The combination of steep channel slope (14 ft/mi) and high streamflow in the main-stem portion of the Kittitas Valley, provide the high velocity streamflows for transporting suspended arsenic.

Annual loads of suspended arsenic increase as much as 3 times between the Yakima River at Umtanum and the Yakima River above Ahtanum Creek at Union Gap, but loads were similar between the Union Gap site and the Yakima River at Euclid Bridge at RM 55 near Grandview and between the Grandview site and the Yakima River at Kiona (table 16). The similarity in load between the Union Gap site and the Grandview site should not be interpreted as meaning that Sulphur Creek's load contribution to the main stem is negligible. Instead, during the irrigation season, a majority of the suspended-arsenic load, as well as streamflow, is diverted into irrigation canals downstream of the Union Gap site. As a result of these diversions, about 80 percent of the streamflow in the main stem of the lower Yakima Valley is from tributaries that carry agricultural return flow (Rinella and others, 1992), and suspended-arsenic loads measured at the Grandview site principally represent suspended arsenic returned to the main stem from agricultural drains.


The annual loads of suspended arsenic for most sites were larger in 1989 than in 1987–88. These differences were attributed to increased streamflows during 1989, rather than an increase in the concentration of suspended arsenic. For example, the annual average streamflow in 1989 for the Kiona site was 20 percent more in 1989 than in 1988.

In the Yakima River, between the Umtanum and Union Gap sites, the sources of arsenic that exist in Wide Hollow Subbasin, and possibly Moxee Subbasin may account for the increase in suspended-arsenic load between the Umtanum and Union Gap sites. Lead-arsenate formulations of pesticide were applied in these areas to control codling moths in apples from 1908 to 1947. To determine contributions in the reach of the main stem affected by Wide Hollow Subbasin and Moxee Subbasin during 1989, the sum of the loads for the Naches River near North Yakima and the Yakima River at Umtanum was subtracted from the load for the Yakima River above Ahtanum Creek at Union Gap. A small correction was applied to the Umtanum loads to account for the load diverted around Union Gap, via Roza Canal, during the irrigation season. During the 1989-irrigation season, approximately 2.2 lbs of suspended arsenic per day enter the main stem over the 9.4-mile reach that includes discharges from the Wide Hollow and Moxee Subbasins; this suspended arsenic represents about one-half the irrigation-season load of arsenic at the Union Gap site.

The suspended arsenic load was estimated for Moxee Drain, the major irrigation return flow in this reach, to determine if the irrigation season load of 2.2 lbs of suspended arsenic per day is a reasonable estimate for the reach of main stem affected by Wide Hollow and Moxee Subbasins. This estimate was made by using the arsenic concentration of streambed sediment instead of the suspended-arsenic concentration, because suspended-arsenic concentrations were not measured. For example, by using the Moxee Drain concentration of suspended sediment (597 mg/L) and discharge (76 ft<sup>3</sup>/s) measurements from July 28, 1988 (Embrey, 1992) and a streambed-sediment concentration of arsenic (3.6 µg/g) [Moxee Drain at Thorp Road; Fuhrer, McKenzie, and others, 1994], the estimated-instantaneous load of suspended arsenic was nearly 1 lb per day. Irrigation season contributions of suspended arsenic from waterways such as Moxee Drain easily could account for the observed increase in load over this reach. Additionally, because the concen-

tration of suspended sediment is high and the concentration of suspended arsenic was low, the load estimated for Moxee Drain was affected more by suspended sediment than by the suspended-arsenic concentration.

Arsenic concentrations in filtered-water samples ranged from <1 to 9 µg/L at the seven fixed sites (table 14). The largest concentrations were found in the lower Yakima Valley (fig. 15; and table 37 at back of report). Arsenic concentrations exceeding 3 µg/L, the 90th percentile for the Yakima River Basin, were found in Sulphur Creek Wasteway near Sunnyside, the




**Figure 15.** Distribution of arsenic concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).


Yakima River at Euclid Bridge at RM 55 near Grandview, and the Yakima River at Kiona—the three most downstream sites in the lower Yakima Valley. Arsenic concentrations in the Yakima River at Cle Elum, the Yakima River at Umtanum, and the Yakima River above Ahtanum Creek at Union Gap generally are below the limit of determination for arsenic ( $1.0 \mu\text{g/L}$ ).

The concentrations of arsenic during the irrigation season, at the Sulphur Creek site, are lower than during the non-irrigation season (fig. 16). In contrast, concentrations of arsenic during the irrigation season at the Grandview and Kiona sites generally were higher than during the non-irrigation season. The lower arsenic concentrations at the Sulphur Creek site are probably the result of dilution from excess canal water spilled into Sulphur Creek from Roza Canal and by irrigation return flow in the Sunnyside Subbasin. Although arsenic concentrations in Roza Canal were not measured directly, concentrations should be similar to those measured during the irrigation season in the Yakima River at Umtanum (RM 140.4). The Umtanum site is located 12.5 miles upstream from the Roza Canal diversion. Monthly arsenic concentrations at Umtanum were low ( $<1 \mu\text{g/L}$ ) for the period April 1989 to March 1990) and probably accounted for the dilution of arsenic in Sulphur Creek Wasteway. Sunnyside Subbasin also receives water from Sunnyside Canal. Again, arsenic concentrations were not measured directly in the canal, but were similar to concentrations in the main stem at the Union Gap site. The Union Gap site is located 3.4 miles upstream of the diversion to Sunnyside Canal and has monthly arsenic concentrations generally below the method reporting level.

Although, arsenic concentrations at the Sulphur Creek site were low during the irrigation season, concentrations generally were similar to higher arsenic concentrations measured at the Grandview and Kiona sites (fig. 16). These increases in arsenic concentrations during the irrigation season at the Grandview and Kiona sites coincide with decreased streamflows (fig. 17) in the main stem and probably result from dilution processes—higher concentrations of arsenic generally were associated with decreased streamflows and conversely, lower concentrations of arsenic generally were associated with increased streamflows. Arsenic concentrations in Sulphur Creek Wasteway, and possibly in other



**Figure 16.** Arsenic concentrations in filtered-water samples at Sulphur Creek Wasteway near Sunnyside, the Yakima River at Euclid Bridge at river mile 55 near Grandview, and the Yakima River at Kiona, Yakima River Basin, Washington, 1989–90 (shaded area represents the irrigation season; less-than values are graphically represented as one-half their value).



**Figure 17.** Arsenic concentrations in filtered-water samples and streamflow for selected time periods at the Yakima River at Euclid Bridge at river mile 55 near Grandview and the Yakima River at Kiona, Yakima River Basin, Washington, 1989–90 (concentrations lower than the limit of determination [1.0 micrograms per liter] are shown as 0.5 micrograms per liter; irrigation season occurs June through September; non-irrigation season occurs October through March).

tributaries that carry irrigation return flow (although not measured in this study), are important sources of arsenic in the main stem. The effect of these sources of arsenic is especially important during the irrigation season, because a majority of the streamflow in the main stem of the lower Yakima Valley is irrigation return flow (Rinella and others, 1992).

Intersite comparisons of arsenic loads were hampered in the Kittitas Valley because a majority of the arsenic determinations were below the method reporting level. In the lower Yakima Valley, however, detectable arsenic was measured and thus quantifiable loads were calculated and are reported in table 17. In Sulphur Creek Wasteway near Sunnyside, for example, the monthly arsenic loads ranged from 3 to 4 pounds per day in 1989 and were similar between the irrigation and non-irrigation seasons. The large variation in arsenic concentrations between the irrigation and the non-irrigation seasons, as well as the small-intrasite variability for arsenic loads, indicates that Sulphur Creek Wasteway is a constant year-round source of arsenic to the main stem. Agricultural lands historically treated with lead-arsenate formulations and present-day applications of phosphate fertilizers may be a source of arsenic to shallow ground water and to surface water.

Arsenic in Sulphur Creek Wasteway also represents a large proportion of the load in the main stem at the Grandview site. In 1989, the Sulphur Creek site (located about 5 miles upstream from the Grandview site) accounted for approximately 20 percent of the annual arsenic load at Grandview but accounted for only about 8 percent of the streamflow—ratios were similar at the Kiona site. In addition, during the non-irrigation season, the Sulphur Creek site accounts for approximately 18 percent of the annual load of arsenic at the Grandview site but accounts for less than 3 percent of the streamflow. The consistency among monthly arsenic loads and the intrasite variability for streamflow and arsenic concentrations confirmed that arsenic is diluted in Sulphur Creek Wasteway during the irrigation season. The consistency among monthly arsenic loads at the Grandview and Kiona sites also indicates that arsenic is discharged to the main stem at a relatively constant rate.

Arsenic in dissolved form (water sample filtered through a nominal 0.45-µm-pore-size filter) comprises the major transport phase in the lower Yakima Valley (fig. 18). For the Sulphur Creek, Grandview, and Kiona sites, the irrigation- and non-irrigation-

**Table 17.** Estimated arsenic loads in filtered-water samples at selected fixed sites, Yakima River Basin, Washington, 1989–90

[The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; loads reported as pounds per day; load values are based on calibration data collected from April 1989 to March 1990; --, indicates insufficient data; bold lines represent the irrigation season, and lightly shaded cells represent the snowmelt portion of the irrigation season; non-irrigation season, October through March]

Year	Jan- uary	Feb- ruary	March	April	May	June	July	Aug- ust	Sep- tember	Oct- ober	Nov- ember	Dec- ember	Daily mean
<b>Sulphur Creek Wasteway near Sunnyside</b>													
1989	2.9	3.0	3.2	3.8	3.8	3.6	3.6	3.7	3.7	3.4	3.0	3.0	3.4
1990	3.0	2.9	3.2	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Euclid Bridge at river mile 55 near Grandview</b>													
1989	17.2	17.0	17.5	18.1	17.7	17.0	16.5	16.5	16.6	16.7	17.3	17.3	17.1
1990	17.5	17.6	17.7	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Kiona</b>													
1989	19.6	18.8	21.2	25.4	21.9	17.8	15.8	16.1	16.7	18.0	19.8	20.4	19.3
1990	21.4	21.2	21.5	24.4	21.1	24.2	16.1	17.8	17.8	--	--	--	--


season loads of arsenic in dissolved forms are from 4 to 9 times higher than the respective suspended loads. Conversely, during the snowmelt season, arsenic is transported in near-equal proportions in both forms. Increases in suspended-sediment concentration, rather than increases in suspended-arsenic concentrations (mass/mass), however, are responsible for the increase in suspended-arsenic load. For example, the suspended-sediment concentration was nearly 8 times the median suspended-sediment concentration during the 1989 snowmelt season at the Kiona site; although, concentrations of suspended arsenic remained unchanged.

Arsenic concentrations in aquatic biota generally were low throughout the Yakima River Basin; however, a few sites in the lower Yakima Valley had relatively high concentrations (table 18). In curlyleaf pondweed, arsenic concentrations ranged from 0.48 to 1.5  $\mu\text{g/g}$  (table 15) and concentrations in the main stem of the lower Yakima Valley were 3 times higher than in the main stem of the Kittitas Valley. Enrichment of arsenic in aquatic vegetation in the lower Yakima Valley also coincides with the high concentrations of arsenic in filtered water described previously. Concentrations of arsenic from 0.1 to 0.4  $\mu\text{g/g}$  were typical in whole sculpins and in fish liver (table 15). Slightly higher concentrations were observed in livers

of bridgeline sucker at Granger Drain at mouth near Granger (0.8  $\mu\text{g/g}$ ) and in carp at the Yakima River at Kiona (0.7  $\mu\text{g/g}$ ). In 1990, arsenic concentrations in Asiatic clams varied little (3.6 to 4.6  $\mu\text{g/g}$ ) among sites. In 1989, arsenic concentrations in caddisflies from Granger Drain site (5.4  $\mu\text{g/g}$ ) and from Sulphur Creek Wasteway near Sunnyside (4.1  $\mu\text{g/g}$ ) exceeded the 85th percentile concentration for the basin.

Arsenic concentrations in fish in the Yakima River Basin were low in relation to concentrations measured in other basins. Concentrations in whole sculpin generally were less than the mean arsenic concentration in freshwater fish collected for the National Contaminant Biomonitoring Program between 1976 and 1984 (table 38, at back of report).

Arsenic concentrations in sculpin from the Yakima River Basin also were similar to background concentrations for bluegills and common carp in the San Joaquin River Basin in California (approximately 0.18 to 0.42  $\mu\text{g/g}$  for bluegills and 0.26 to 1.68  $\mu\text{g/g}$  for common carp, as calculated from original wet-weight data) (Saiki and May, 1988). Sampling sites for sculpin, however, are not representative of the entire Yakima River Basin. Sculpin were not collected from the main stem, and samplings in tributaries containing agricultural runoff were limited to Satus Creek at gage at Satus and Ahtanum Creek at Union Gap.



**Figure 18.** Arsenic loads in filtered-water samples and in suspended-sediment samples for selected time periods at fixed sites, Yakima River Basin, Washington, 1989 (the term "filtered water" represents the portion of a water sample passing through a nominal 0.45 micrometer pore-size filter; non-irrigation season occurs October through March; snowmelt season occurs April through May; irrigation season occurs June through September; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).

Arsenic concentrations in Asiatic clams from the four main-stem sites in the lower Yakima Valley and from Spring Creek at mouth near Whitstran were compared to concentrations reported for uncontaminated or minimally contaminated aquatic environments (table 39, at back of report). Arsenic concentrations in Asiatic clams of the Yakima River Basin were an order of magnitude higher than concentrations reported in the Apalachicola River in Florida (Elder and Matraw, 1984); arsenic concentrations were at least 3 times higher than in the Sacramento River Basin in California (McCleneghan and others, 1981), but were similar to arsenic concentrations in Asiatic clams in the San Joaquin River in California which are considered to be affected by minor anthropogenic sources (Johns and Luoma, 1990; Leland and Scudder, 1990). However, Asiatic clams of the Yakima River Basin had lower arsenic concentrations than the 8 to 13  $\mu\text{g/g}$  measured in Asiatic clams of the upper San Joaquin River in California that are affected by arsenic in agricultural drainage water (Leland and Scudder, 1990).

Median concentrations of lead in suspended sediment ranged from 15  $\mu\text{g/g}$  to 24  $\mu\text{g/g}$  at the seven fixed sites (table 13); the lowest and highest median values of lead, respectively, were in the Yakima River at Umtanum and in Sulphur Creek Wasteway near Sunnyside (fig. 19; and table 35, at back of report). The low concentrations of suspended lead at Cle Elum, Umtanum, and Naches indicate the lack of a significant geologic source of lead and contrast sharply with enrichment measured at these same sites for suspended elements such as arsenic, chromium, and nickel—all of which have geologic sources. Furthermore, the lack of a significant geologic source is consistent with the distribution of lead in streambed sediment of the Yakima River Basin (Fuhrer, McKenzie, and others, 1994). For example, the 90th percentile concentration of lead in streambed sediment from 272 first- or second-order streams in the basin was only 23  $\mu\text{g/g}$ . A few small and isolated geologic sources of lead were found in streambed sediment of first- and second-order streams in the Naches Subbasin; however, no evidence of enrichment, was found in corresponding higher-order streams (Fuhrer, McKenzie, and others, 1994), indicating that measurable quantities of lead were not being transported as suspended sediment and then later redeposited in streambed sediment of higher-order streams.




**Table 18. Comparison of low and high arsenic concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91**

[For filtered water and suspended sediment, the low- and high-concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For streambed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an “H” in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an “L” in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an “\*” in the table. The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale sucker livers, sample species: largescale sucker (*Catostomus macrocheilus*) mountain whitefish (*Prosopium williamsoni*), sculpin (*Cottus spp.*), Asiatic clam (*Veneroida: Corbiculidae Corbicula fluminea*), and curlyleaf pondweed (*Potamogeton crispus*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Yakima River above Ahtanum Creek at Union Gap, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987–90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]

Site reference number	Site name	Filtered water	Sediment		Aquatic biota				
			Streambed	Suspended	Largescale sucker	Mountain whitefish	Sculpin	Asiatic clam	Curlyleaf pondweed
1	Wapatus River at mouth near Roslyn	--	H	--	*	--	--	--	--
3	Jungle Creek near mouth near Cle Elum	--	H	--	--	--	--	--	--
5	Teanaway River below Forks near Cle Elum	--	*	--	--	--	--	--	--
6	Yakima River at Cle Elum	L	H	*	--	L	*	--	--
7	Naneum Creek below High Creek near Ellensburg	--	L	--	--	--	L	--	--
8	Taneum Creek at Taneum Meadow near Thorp	--	L	--	--	--	L	--	--
10	Little Naches River at mouth near Cliffdell	--	*	--	--	--	--	--	--
12	South Fork Manastash Creek near Ellensburg	--	L	--	--	--	L	--	--
13	American River at Hells Crossing near Nile	--	H	--	--	--	H	--	--
14	Cherry Creek above Wipple Wasteway at Thrall	--	*	--	--	*	--	--	L
19	Yakima River at Umtanum	L	*	L	--	L	--	--	L
20	Umtanum Creek near mouth at Umtanum	--	L	--	--	--	L	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	--	H	--	--	--	--	--	--
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	--	H	--	--	--	H	--	--
26	Naches River near North Yakima	L	*	*	--	*	--	--	--

**Table 18.** Comparison of low and high arsenic concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Filtered water	Sediment		Aquatic biota				
			Streambed	Suspended	Fish			Asiatic clam	Curlyleaf pondweed
					Largescale sucker	Mountain whitefish	Sculpin		
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	--	*	--	--	--	--	--	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	--	*	--	--	--	--	--	H
30	Moxee Drain at Thorp Road near Union Gap	--	*	--	--	--	--	--	--
31	Ahtanum Creek at Union Gap	--	L	--	--	--	H	--	--
32	Yakima River above Ahtanum Creek at Union Gap	L	--	L	--	--	--	--	--
33	Yakima River at Parker	--	*	--	L	H	--	--	*
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	--	L	--	--	--	L	--	--
40	Granger Drain at mouth near Granger	--	*	--	--	--	--	--	--
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger	--	*	--	H	H	--	*	--
43	Toppenish Creek at Indian Church Road near Granger	--	*	--	--	--	--	--	--
47	Satus Creek at gage at Satus	--	H	--	*	--	H	--	*
48	Yakima River at river mile 72 above Satus Creek near Sunnyside	--	*	--	--	--	--	*	*
50	Yakima River at Kiona	H	*	*	*	*	--	*	*
52	Sulphur Creek Wasteway near Sunnyside	H	*	H	--	--	--	--	--
53	Satus Creek below Dry Creek near Toppenish	--	*	--	--	--	H	--	--
54	Spring Creek at mouth at Whitstran	--	*	--	--	--	--	L	--
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	H	H	H	L	--	--	H	H
57	Satus Creek above Wilson-Charley Canyon near Toppenish	--	L	--	--	--	H	--	--



#### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.

- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value
- × 1.5 to 3 times the interquartile range from the 25-percentile value

**Figure 19.** Distribution of lead concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).

A few anomalous concentrations of suspended lead were measured in the Yakima River at Cle Elum and in the Yakima River at Umtanum (fig. 19). At the Cle Elum site, concentrations of suspended lead as high as 120 µg/g and 60 µg/g were measured during the rising limb of a May 9, 1989 snowmelt and a January 9–12, 1990 rainstorm or snowstorm, respectively. During the January storm, 60 µg/g of lead was measured on the rising limb, but decreased to just 6 µg/g at

the peak of the storm. The Cle Elum site, located 0.15 mi downstream from Crystal Creek, receives urban runoff from the towns of Roslyn and Cle Elum, and runoff from numerous open-pit coal mines east of Roslyn. During the storm, Crystal Creek created a turbidity plume which, during the rising limb of the storm, obscured the shallow river bottom in the main stem at the Cle Elum site. The Crystal Creek drainage, located in the nonmarine rocks geologic unit, contains concentrations of lead in streambed sediment (13 µg/g, 14 µg/g, and 18 µg/g; Fuhrer, McKenzie, and others, 1994) that were low when compared to the 60 µg/g of suspended lead measured during the rising limb of the storm. Considering the close proximity of the Cle Elum site to outfall from Crystal Creek and the lack of measurable enrichment for lead in streambed sediment, the high concentration of suspended lead on the rising limb of the storm may have resulted from local urban runoff.

The probable effect of urban runoff on suspended-lead concentrations is confirmed by computing ratios of lead:arsenic. Arsenic was selected because upstream of the Cle Elum site in the Kittitas Valley, it resides predominantly in geologic formations. Variations in lead relative to arsenic may be indicative of urban runoff. For the May snowmelt and January storms, the ratios of lead:arsenic were 15 and 8, respectively. Generally, the lead:arsenic ratios ranged from 2.2 to 2.6 for streambed sediment formed from geologic units in the Kittitas Valley (based on median lead and arsenic concentrations shown in table 10 in Fuhrer, McKenzie, and others, 1994). These ratios are similar to lead:arsenic ratios measured in suspended sediment samples. Lead concentrations during the May snowmelt and January storms, however, were approximately 3 to 6 times higher than expected and suggests that a small anthropogenic source of lead exists in close proximity to the Cle Elum site.

In addition to urban runoff, agricultural areas that historically were treated with pesticides containing lead arsenate possibly may be sources of suspended lead. The median concentration of suspended lead in Sulphur Creek Wasteway near Sunnyside, for example, was higher than that measured at the other fixed sites (fig. 19). Lead concentrations at the Sulphur Creek site, however, did not vary seasonally to the extent measured for arsenic and were generally in agreement with lead concentrations found in streambed sediment (17 to 21 µg/g) in the Sulphur Creek

drainage (Ryder and others, 1992). The lack of seasonal variability for lead may result from geochemical associations that retard the movement of lead to shallow ground water and subsequently to Sulphur Creek, during the non-irrigation season. The application of phosphate fertilizer tends to form insoluble lead-phosphate compounds (Peryea, 1989). In soils of agricultural areas historically treated with pesticides containing lead arsenate, lead was found to accumulate below the immediate soil surface (30 to 60 cm [centimeters]), but not to the deeper soil depths (50 to 120 cm) noted for arsenic (Peryea, 1989). Lead may be trapped effectively between subsurface ground water and the immediate soil surface. The immobile nature of lead may explain the absence of lead anomalies in suspended sediment and in filtered-water samples during the non-irrigation season, when subsurface ground-water contributions dominate Sulphur Creek streamflow.


Lead in filtered-water samples, determined by atomic-absorption spectrometry with graphite furnace, ranged from  $<0.5$  to  $1.9 \mu\text{g/L}$  at the seven fixed sites (table 14). Most lead determinations (approximately 94 percent of the 279 measurements made by AAGF), however, were below the limit of determination ( $0.5 \mu\text{g/L}$ ). Detectable lead concentrations were distributed evenly among most sampling sites. The Cle Elum site, however, had no detectable concentrations of lead. With one exception, spatial and temporal patterns generally were absent among sites. A few days after the close of the irrigation season on October 15, 1989 detectable concentrations of lead were measured at the Naches site ( $0.7 \mu\text{g/L}$ ), the Union Gap site ( $0.8 \mu\text{g/L}$ ), and the Sulphur Creek site ( $0.9 \mu\text{g/L}$ ). All sites had experienced sharp decreases in streamflow over the preceding days due to the close of the irrigation season. None of these lead concentrations, however, exceeded water-quality criteria, regulations, or standards. Additionally, only a small number of the 279 lead determinations for 1987–90 exceeded water-quality guidelines (table 7).

Because the quantity of rainfall in the lower Yakima Valley is small (less than 10 inches annually), storms would be expected to contain detectable lead if urban runoff was a significant source. On the basis of the absence of lead enrichment, in general, and storm related enrichment, in particular, urban sources of lead are probably limited over much of the Yakima River Basin.

The maximum lead concentrations in fish liver, Asiatic clams (*Corbicula fluminea*), and caddisflies (*Hydropsyche* spp.) were 0.29, 0.40, and  $5.6 \mu\text{g/g}$ , respectively (table 15). For *Hydropsyche* spp., the highest lead concentrations were measured in Wide Hollow Creek site (fig. 20). Lead in rainbow trout liver from the Wide Hollow Creek was below the limit of determination ( $<0.19 \mu\text{g/g}$ ). The apparent disparity in these results may be attributed to the tendency of lead to decrease in concentration in higher trophic-level consumers, such as fish (Luoma, 1986). Lead also was moderately elevated in *Hydropsyche* spp. in Ahtanum Creek ( $3.0 \mu\text{g/g}$ ), Sulphur Creek ( $2.6 \mu\text{g/g}$ ), and Spring Creek ( $2.6 \mu\text{g/g}$ ). The Ahtanum site, which is located near the mouth of Ahtanum Creek, receives irrigation return flow and some urban runoff. The Ahtanum site has lead concentrations in *Hydropsyche* spp. that are more than 5 times higher than at the upstream site (South Fork Ahtanum Creek above Conrad Ranch near Tampico) which does not receive irrigation return flow and is removed from urban effects. A substantial lead concentration ( $24 \mu\text{g/g}$ ) was measured in *Arctopsyche* sp. in Naneum Creek; however, other insect taxa collected from this site were not similarly enriched (Fuhrer, Fluter, and others, 1994).

Most samples collected in 1989 had lead concentrations below the limit of determination. Lead was detected in 1989 only caddisflies from Wide Hollow Creek at old Sewage Treatment Plant at Union Gap ( $8.0$  and  $9.0 \mu\text{g/g}$ ), and in Granger Drain at mouth near Granger ( $5.0 \mu\text{g/g}$ ) and in Sulphur Creek Wasteway near Sunnyside ( $5.0 \mu\text{g/g}$ ).

Compared with concentrations reported for benthic insects from other systems (Cain and others, 1992; Lynch and others, 1988), lead appears to be slightly enriched at Wide Hollow Creek, Ahtanum Creek, and possibly Naneum Creek. Most tissue samples, however, had little or no lead enrichment which is consistent with the geology as well as lead concentrations in streambed sediment (table 19; and table 40, at back of report). Concentrations of lead in aquatic plants were below limits of determination, except for Wide Hollow Creek at old Sewage Treatment Plant at Union Gap where a lead concentration of  $7.0 \mu\text{g/g}$  was measured in waterweed (Fuhrer, Fluter, and others, 1994). As described earlier, Wide Hollow Creek also had the largest concentrations of lead among the biological sampling sites in the Yakima River Basin.



**Figure 20.** Distribution of lead concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are graphically represented for caddisflies; sample species: caddisflies [*Trichoptera*: *Hydropsychidae* *Hydropsyche* spp.]).

Lead concentrations were comparatively low in fish livers from the Yakima River Basin (0.09 to 0.31  $\mu\text{g/g}$ ). For comparison, five fish species from two rivers (that receive agricultural and some urban runoff), had lead concentrations ranging from 0.12 to 0.44  $\mu\text{g/g}$  (calculated from original wet-weight data; Barak and Mason 1990). Mean lead concentrations in the liver of bream (*Abramis brama*) and in pike perch (*Stizostedion lucioperca*) from Lake Balaton, a heavily recreated, mildly contaminated lake in Hungary, were 12.9  $\mu\text{g/g}$  and 3.10  $\mu\text{g/g}$  (Salanki and others, 1982). The 85th percentile lead concentration for fish livers sampled in California's Toxic Substances Monitoring Program was approximately 0.4  $\mu\text{g/g}$  (calculated from original wet-weight data; Rasmussen, 1992).

Primarily on the basis of samplings from main-stem sites, lead does not appear to be enriched in Asiatic clams. Concentrations in the Yakima River Basin were at least 4 times lower than from an area of the San Joaquin River in California that is not subject to urban and industrial inputs (Luoma and others,

1990), but concentrations were somewhat higher than concentrations in the Apalachicola River in Florida (Elder and Matraw, 1984; table 39, at back of report).

## Barium

Concentrations of barium in streambed sediment of the Yakima River Basin ranged from 380 to 590  $\mu\text{g/g}$  (table 12) and are well within the 200 to 1,700  $\mu\text{g/g}$  range of concentration which encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). Concentrations of barium increase down the main stem from a minimum of 470  $\mu\text{g/g}$  (site 19) at the Yakima River at Umtanum to a maximum of 540  $\mu\text{g/g}$  (site 54) at the Yakima River at Grandview (fig. 21), and barium concentrations are notably higher at the mouths of several tributaries that carry agricultural return flow to the main stem. For example, concentrations of 510  $\mu\text{g/g}$ , 560  $\mu\text{g/g}$ , and 590  $\mu\text{g/g}$  (the concen-

**Table 19. Comparison of low and high lead concentrations in sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91**


[For filtered water and suspended sediment, the low and high concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; for streambed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an "H" in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an "L" in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an "\*" in the table. The term "filtered water" is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale sucker livers; sample species: largescale sucker (*Catostomus macrocheilus*), caddisfly (*Hydropsyche* spp.), and Asiatic clam (*Veneroida: Corbiculidae Corbicula fluminea*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Yakima River above Ahtanum Creek at Union Gap, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987–90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]

Site reference number	Site name	Sediment		Aquatic biota			
		Streambed	Suspended	Largescale-sucker liver	Caddisfly	Asiatic clam	
1	Waptus River at mouth near Roslyn	*	--	--	--	--	--
3	Jungle Creek near mouth near Cle Elum	H	--	--	--	--	--
5	Teaaway River below Forks near Cle Elum	L	--	--	--	--	--
6	Yakima River at Cle Elum	*	L	--	*	--	--
7	Naneum Creek below High Creek near Ellensburg	L	--	--	*	--	--
8	Taneum Creek at Taneum Meadow near Thorp	*	--	--	--	--	--
10	Little Naches River at mouth near Cliffdel	L	--	--	*	--	--
12	South Fork Manastash Creek near Ellensburg	*	--	--	--	--	--
13	American River at Hells Crossing near Nile	*	--	--	--	--	--
14	Cherry Creek above Wipple Wasteway at Thrall	L	--	--	*	--	--
16	Cherry Creek at Thrall	--	--	--	*	--	--
19	Yakima River at Umtanum	*	L	--	*	--	--
20	Umtanum Creek near mouth at Umtanum	L	--	--	L	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	L	--	--	*	--	--
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	*	--	--	L	--	--
26	Naches River near North Yakima	H	*	H	*	--	--

**Table 19.** Comparison of low and high lead concentrations in sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Sediment		Aquatic biota		
		Streambed	Suspended	Largescale- sucker liver	Caddisfly	Asiatic clam
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	H	--	--	H	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	H	--	--	H	--
30	Moxee Drain at Thorp Road near Union Gap	*	--	--	--	--
31	Ahtanum Creek at Union Gap	*	--	--	H	--
32	Yakima River above Ahtanum Creek at Union Gap	--	*	--	--	--
33	Yakima River at Parker	*	--	L	*	--
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	L	--	--	*	--
40	Granger Drain at mouth near Granger	*	--	--	H	--
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger	*	--	*	L	H
43	Toppenish Creek at Indian Church Road near Granger	*	--	--	--	--
47	Satus Creek at gage at Satus	L	--	*	*	--
48	Yakima River at river mile 72 above Satus Creek near Sunnyside	*	--	--	--	--
50	Yakima River at Kiona	H	H	*	*	L
52	Sulphur Creek Wasteway near Sunnyside	*	H	--	H	--
53	Satus Creek below Dry Creek near Toppenish	*	--	--	L	--
54	Spring Creek at mouth at Whitstran	*	--	--	H	*
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	H	*	H	*	*
57	Satus Creek above Wilson-Charley Canyon near Toppenish	L	--	--	*	--





**Figure 21.** Barium concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987 ("Waptus River" represents Waptus River at mouth near Roslyn; "Cherry Creek" represents Cherry Creek above Wipple Wasteway at Thrall; "Umtanum" represents Yakima River at Umtanum; "Moxee Drain" represents Moxee Drain at Thorp Road near Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; and "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview).

tration maximum) were measured at Cherry Creek above Wipple at Thrall (site 14), Moxee Drain at Thorp Road near Union Gap (site 30), and Sulphur Creek Wasteway near Sunnyside (site 52), respectively—all sites carry agricultural return flow to the main stem. The increased barium concentration in streambed sediment of agricultural drains and streambed sediment of the lower main stem probably results from the precipitation of barium as barite (barium sulphate). According to Hem (1989), barium solubility in water is limited by the solubility of barite. When the concentrations of dissolved barium and sulphate become large enough to exceed the solubility product for barite, the concentrations are precipitated to streambed sediment as barite. A common agricultural source of sulphate in the Yakima River Basin is the application of zinc sulphate to promote and retain the blossoms on fruit trees (Robert Wample, University of Washington, Prosser Experimental Station, oral commun., 1991). The solubility of barite is probably exceeded in many of the basin's agricultural waters as a result of various soil amendments that contain sul-

phate. The large concentration of barium in Waptus River streambed sediment is attributed to nonmarine sedimentary rock of the Swauk Formation which contains barium and potassium aluminosilicate minerals (Tabor and others, 1982).

Barium concentrations in filtered-water samples ranged from <2 to 79 µg/L at the seven fixed sites (table 14). The highest concentrations were in the lower Yakima Valley (table 37, at back of report). Barium concentrations exceeding 39 µg/L, the 90th percentile for the Yakima River Basin, were found in Sulphur Creek Wasteway near Sunnyside and in the Yakima River at Kiona. The highest concentrations, however, were found in Sulphur Creek Wasteway and coincide with high barium concentrations in streambed sediment.


## Cadmium

Most of the cadmium determinations were made using inductively coupled plasma-atomic emission spectrometry (ICP-AES) which had an analytical limit

of determination of 2  $\mu\text{g/g}$ . Cadmium concentrations were below the limit of determination for the 27 samples analyzed by using ICP-AES. Five of 32 determinations, however, were made by using an organometallic halide extraction prior to ICP-AES which had a method reporting level of 0.05  $\mu\text{g/g}$  (Arbogast, 1990). Concentrations of cadmium, determined by using the latter analytical method, ranged from 0.2 to 0.8  $\mu\text{g/g}$ . The largest concentration of cadmium (0.8  $\mu\text{g/g}$ ) was found in streambed sediment in the American River at Hells Crossing near Nile (site 13) and was well above the average crustal abundance for basalt of 0.19  $\mu\text{g/g}$  (Parker, 1967) that comprises a majority of the Yakima River Basin. Additionally, the concentration of cadmium in the American River site also exceeded that found in the anthropogenically affected main stem of the Willamette River at Portland, Oregon (Fuhrer, 1989).

Concentrations of suspended cadmium at the seven fixed sites ranged from  $<0.1$   $\mu\text{g/g}$  to 33  $\mu\text{g/g}$  (table 13). Cadmium enrichment was found at the Yakima River at Cle Elum in the Kittitas Valley, the Naches River near North Yakima in the mid-Yakima Valley at Naches, and at the Yakima River at Euclid Bridge at RM 55 near Grandview and the Yakima River at Kiona in the lower Yakima Valley (fig. 22; and table 35, at back of report). The highest median concentration (0.8  $\mu\text{g/g}$ ), however, was found in the Kittitas Valley at the Cle Elum site.

Concentrations of suspended cadmium in the Yakima River at Cle Elum ranged from 0.3 to 9.3  $\mu\text{g/g}$ . Based on limited data, suspended-cadmium concentrations were higher during storms (rising limb of the storm hydrograph) and during late snowmelt. In the late snowmelt season, during mid-April and May, cadmium concentrations increased by a factor of four but suspended-sediment concentrations decreased from 85 to 5  $\text{mg/L}$ . A similar pattern resulted during a January 9–10, 1990 storm. Relative to the preceding month's (December) sampling, suspended-cadmium concentrations during the rising limb of the January 9, 1990 storm increased to 1.0  $\mu\text{g/g}$ —2 times that of the December sampling and the suspended-sediment concentration increased to 12  $\text{mg/L}$ —4 times that of the December sampling. The following day (January 10, 1990), during the storm's peak, the suspended cadmium concentration decreased to 0.3  $\mu\text{g/g}$ —a decrease of 3 times in relation to the rising limb of the storm, but the suspended-sediment concentration increased from 12  $\text{mg/L}$  to 130  $\text{mg/L}$ . During storms



**Figure 22.** Distribution of cadmium concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).


and snowmelt, suspended-cadmium concentrations at Cle Elum decreased as suspended-sediment concentration increased.

Crystal Creek created a turbidity plume of suspended sediment that obscured the shallow-river bottom in the Yakima River at Cle Elum during the January 10, 1990 storm. The lower suspended-cadmium concentration during the storm's peak, noted earlier at the Cle Elum site, coincided with the presence of the turbidity plume of suspended sediment

from Crystal Creek and indicates that suspended-cadmium concentrations were diluted from mixing with incoming sediment from Crystal Creek. This hypothesis is possible especially if cadmium concentrations in the Crystal Creek drainage are low. The lower limit of determination of cadmium for the sites sampled in the Crystal Creek drainage, as part of the occurrence and distribution survey (Fuhrer, Flutter, and others, 1994), was too high to conclude that Crystal Creek sediment potentially could dilute suspended-cadmium concentrations at the Cle Elum site.

Suspended-cadmium concentrations in the Naches River near North Yakima may reflect the presence of cadmium in streambed sediment of the Upper Naches Subbasin. A cadmium concentration of 0.8  $\mu\text{g/g}$  was measured in streambed sediment of the American River at Hells Crossing near Nile (site 13) located in the Upper Naches Subbasin. This cadmium concentration was 4 times higher than the average crustal abundance of cadmium in basalt and was similar to the median suspended-cadmium concentration (0.7  $\mu\text{g/g}$ ) at site 13 in the Naches Subbasin. Suspended-cadmium concentrations at the Naches site probably originate from geologic sources in the Upper Naches and Tieton Subbasins. Most of the cadmium determinations for streambed sediment, however, had an analytical limit of determination (2.0  $\mu\text{g/g}$ ) that exceeded concentrations typical in basalt rock. As a result, cadmium concentrations at many sites were below the limit of determination and could not be used for making comparisons with suspended-cadmium concentrations or for locating geologic sources of cadmium.

The highest cadmium concentrations in the Naches River near North Yakima (site 26) generally were found during the non-irrigation season (November through March), when suspended-sediment concentrations were small (<10 mg/L) and the proportion of fine-grained-size sediment was large (>85 percent) (fig. 23). During the irrigation and snowmelt seasons at Naches River (site 26), the highest suspended-cadmium concentrations coincide with periods of increased streamflow (fig. 23). During the irrigation season at Naches, streamflow was augmented by withdrawing water from Rimrock Reservoir and Bumping Reservoir. Irrigation-season releases from these reservoirs probably assist in the transport of cadmium which originates from sources in the Upper Naches Subbasin.



**Figure 23.** Cadmium concentrations in suspended sediment and suspended-sediment concentrations, percentage of suspended sediment finer than 62 micrometers in diameter, and streamflow in the Naches River near North Yakima, Yakima River Basin, Washington, 1987-90 (irrigation season occurs June through September; non-irrigation season occurs October through March; and snowmelt season occurs April through May).

As noted earlier, the Kittitas Valley had concentrations of suspended cadmium which generally exceeded concentrations measured at other fixed sites. The annual load of cadmium in the Kittitas Valley, however, is similar to that measured in the mid-Yakima Valley at the Yakima River above Ahtanum Creek at Union Gap, the lower Yakima Valley at the Yakima River at Euclid Bridge at RM 55 near Grandview, and the Yakima River at Kiona (table 20). Although annual loads of cadmium generally are similar for these main-stem sites, the proportions of the annual load transported during the snowmelt, irrigation, and non-irrigation seasons vary; moreover, snowmelt loads generally exceed loads during the irrigation season and the non-irrigation season (fig. 24). At the Kiona site, nearly 68 percent of the 1989-annual-cadmium load was transported during the snowmelt season—this load is nearly 6 times the quantity of cadmium transported during the irrigation season. Similarly, at the Naches River near North Yakima more than 70 percent of the 1989-annual-cadmium load was transported during the snowmelt season—this load is 4 times the quantity of cadmium transported during the irrigation season. During the snowmelt season, the higher cadmium loads at the Naches site are the result of suspended-sediment concentrations that are 3 to 4 times larger than during the irrigation season.

In filtered-water samples, cadmium concentrations, determined by atomic-absorption spectrometry with graphite furnace, ranged from  $<0.2$  to  $2.2 \mu\text{g/L}$  at the seven fixed sites (table 14). Among fixed sites, the highest number of detections of cadmium were in the Yakima River at Cle Elum and in the Yakima River at Umtanum—cadmium was detected at least 25 percent of the time at both sites (table 37, at back of report). Conversely, in the mid- and lower Yakima Valley at least 75 percent of the determinations for cadmium were below the limit of determination ( $0.2 \mu\text{g/L}$ ). The higher concentrations of cadmium in filtered-water samples in the Kittitas Valley also coincide with cadmium enrichment in suspended sediment. Cadmium concentrations in suspended sediment are higher at Cle Elum during snowmelt and storms. Comparisons between cadmium concentration in filtered-water samples in the Yakima River Basin and cadmium concentrations in other studies (table 36, at back of report) are hampered by a large amount of censored data.

Cadmium concentrations, which exceed the limit of determination by at least a factor of two, were

selected as conservative measures of detectable cadmium in order to illustrate temporal and spatial patterns (table 21). These patterns indicate that sources of cadmium generally are not anthropogenic, but these patterns are related to periods of snowmelt and to storms in the Kittitas Valley. Cadmium was detected during periods of snowmelt at the Cle Elum, Umtanum, Naches, and Union Gap sites in concentrations that ranged from  $0.4$  to  $0.5 \mu\text{g/L}$  in April 1987 and at the same sites in 1988 at lower concentrations. In November 1988, fixed sites were visited during an early November winter-warming period (early snowmelt); and cadmium concentrations ranging from  $0.2$  to  $2.2 \mu\text{g/L}$  were measured at the Cle Elum, Umtanum, Naches, Union Gap, and Kiona sites. Similar patterns also were found during storms and during the irrigation season in June. The cadmium concentration ( $2.2 \mu\text{g/L}$ ) at the Naches site exceeded that of other sites by an order of magnitude.

During November 1988, the increase in cadmium concentration at the Naches site, in part, may have resulted from construction of a walking path/green way along the left bank, upstream and downstream of the Naches site. Fill material was placed into and alongside the right bank of the Naches River as a foundation for an asphalt pathway. Although, in this instance, the cadmium concentration probably was affected to a larger degree by the construction activity than by snowmelt; the snowmelt conditions probably enhanced the cadmium concentration. The effect of snowmelt conditions on cadmium concentrations was noted during an earlier sampling, prior to construction activity. In December 1987—again, during a period of winter warming and subsequent snowmelt runoff—dissolved cadmium ( $0.3 \mu\text{g/L}$ ) was detected at the Naches site on the rising limb of the streamflow hydrograph. This time, however, the concentration was similar to streamflow enhanced by snowmelt at other fixed sites.


Load computations for cadmium were hampered at mid- and lower-Yakima Valley sites by large amounts of censored data (element concentrations lower than the limit of determination). Consequently, load estimates were made only for the Cle Elum and Umtanum sites—sites with the most uncensored data. In 1988 and 1989, the annual load of cadmium between the Cle Elum and Umtanum sites increased 2 times (table 22), and the annual streamflow increased by a factor of about 1.5. The increase in load was primarily because of increases in streamflow and, to a

**Table 20.** Estimated cadmium loads in suspended sediment at selected-fixed sites, Yakima River Basin, Washington, 1987–90

[Loads reported as pounds per day; load estimates are based on calibration data collected from March 1987 to March 1990; --, indicates insufficient data; bold lines represent the irrigation season, and lightly shaded cells represent the snowmelt portion of the irrigation season; non-irrigation season, October through March; Jan, January; Feb, February; Mar, March; Apr, April; Aug, August; Sept, September; Oct, October; Nov, November; Dec, December]

Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Daily mean
<b>Yakima River at Umtanum</b>													
1987	<0.05	0.1	0.5	0.4	0.3	0.4	0.3	0.2	<0.05	<0.05	<0.05	<0.05	0.2
1988	<.05	.1	.2	.5	.2	.2	.3	.2	.1	<.05	<.05	.1	.2
1989	.1	.1	.2	1	.6	.4	.3	.2	.1	<.05	<.05	.1	.3
1990	.2	.3	.4	1	.6	.7	.3	.2	.1	--	--	--	--
<b>Naches River near North Yakima</b>													
1987	--	--	.2	.4	1	.1	<.05	<.05	<.05	<.05	<.05	<.05	--
1988	<.05	<.05	.1	.4	.3	.1	<.05	<.05	.1	<.05	<.05	<.05	.1
1989	<.05	<.05	.1	.7	.5	.2	<.05	<.05	.1	<.05	<.05	.1	.1
1990	.3	.2	.2	--	--	--	--	--	--	--	--	--	--
<b>Yakima River above Ahtanum Creek at Union Gap</b>													
1987	.2	.4	.8	.7	.9	.3	.2	.2	.1	.1	.1	.1	.3
1988	.2	.4	.3	.7	.5	.3	.2	.2	.1	.1	.2	.2	.3
1989	.3	.4	.5	1	.7	.4	.2	.2	.1	.1	.1	.2	.4
1990	.4	.5	.7	1	.7	.6	.3	.2	.2	--	--	--	--
<b>Sulphur Creek Wasteway near Sunnyside</b>													
1987	--	--	<.05	.1	.1	.1	.1	<.05	<.05	<.05	<.05	<.05	--
1988	<.05	<.05	.1	.1	.1	.1	<.05	<.05	<.05	<.05	<.05	<.05	<.05
1989	<.05	<.05	<.05	.1	.2	.1	.1	<.05	<.05	<.05	<.05	<.05	<.05
1990	<.05	<.05	.1	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Euclid Bridge at river mile 55 near Grandview</b>													
1987	--	--	.6	.6	.6	.2	.2	.1	.1	.1	.1	.2	--
1988	.3	.4	.4	.6	.4	.3	.1	.1	.1	.1	.2	.2	.3
1989	.4	.4	.6	1	.7	.3	.2	.1	.1	.1	.2	.3	.4
1990	.5	.6	.7	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Kiona</b>													
1987	.1	.1	1	.4	1	.1	.1	<.05	<.05	<.05	<.05	.1	.2
1988	.1	.1	.1	.7	.3	.2	.1	.1	.1	.1	.1	.1	.2
1989	.1	.1	.4	2	1	.2	.1	.1	.1	.1	.1	.1	.4
1990	.3	.3	.3	1	.7	2	.1	.2	.1	--	--	--	--





**Figure 24.** Percentages of the annual load of cadmium in suspended sediment for the Yakima River at Umtanum, the Yakima River above Ahtanum Creek at Union Gap, the Yakima River at Euclid Bridge at river mile 55 near Grandview, and the Yakima River at Kiona, Yakima River Basin, Washington, 1989 (non-irrigation season occurs October through March; snowmelt season occurs April through May; irrigation season occurs June through September).

lesser extent, increases in cadmium concentrations between sites.

A large proportion of the annual cadmium load is transported during the irrigation season. At the Cle Elum and Umtanum sites in 1989, for example, the irrigation season accounted for 63 and 53 percent, respectively, of the annual load. The importance of the irrigation season, in relation to the annual load of cadmium, is directly related to streamflow, because storage reservoirs are used to augment streamflow in the Kittitas Valley during the irrigation season. At the Cle Elum and Umtanum sites in 1989, the irrigation season accounted for 61 and 49 percent, respectively, of the annual streamflow. At these sites, smaller

proportions of the annual-cadmium load (17 and 28 percent, respectively) were transported during the snowmelt season in 1989. The smaller proportion of the load transported during the snowmelt season also is directly related to streamflow. During the 1989 snowmelt season, the Cle Elum and Umtanum sites accounted for only 19 and 31 percent, respectively, of the annual streamflow.

Based on comparisons between dissolved (water sample filtered through a nominal 0.45  $\mu\text{m}$ -pore-size filter) and suspended forms of cadmium, the dissolved form is the major transport phase in the Yakima River at Umtanum (fig. 25). Dissolved loads exceeded suspended loads by more than an order of magnitude

**Table 21.** Frequency of occurrence of cadmium concentrations equalling or exceeding 0.2 micrograms per liter in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–89

[The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; ref. no., reference number; Jan, January; Feb, February; Mar, March; Apr, April; Aug, August, Sept, September; --, no data; blank cells in table, cadmium concentrations less than 0.2 micrograms per liter; shaded cells represent months in which the total number of occurrences of cadmium concentrations that equal or exceed 0.2 µg/L is greater than or equal to seven; •, one cadmium concentration equalled or exceeded 0.2 micrograms per liter (µg/L); ••, two cadmium concentration equalled or exceeded 0.2 micrograms per liter; Cle Elum, Yakima River at Cle Elum; Umtanum, Yakima River at Umtanum; Naches, Naches River near North Yakima; Union Gap, Yakima River above Ahtanum Creek at Union Gap; Sulphur Creek, Sulphur Creek Wasteway near Sunnyside; Grandview, Yakima River at Euclid Bridge at river mile 55 near Grandview; Kiona, Yakima River at Kiona]


Year	Site ref. no.	Site name	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec
1987	6	Cle Elum	--	--	--	•		•	•	•				
1987	19	Umtanum	--	--	--	•	•	•			•			
1987	26	Naches	--	--	--	•		•						•
1987	32	Union Gap	--	--	--	•	•	•						
1987	50	Kiona	--	--	--			•					•	
1987	52	Sulphur Creek	--	--	--	•		•						
1987	56	Grandview	--	--	--			•			•			•
Subtotal			--	--	--	5	2	7	1	1	2	0	1	2
1988	6	Cle Elum	•			••	•	•		•	•		•	•
1988	19	Umtanum			••	•		•			•	•	•	•
1988	26	Naches				•				•			•	
1988	32	Union Gap				•							•	
1988	50	Kiona	•		•								•	•
1988	52	Sulphur Creek	••					•						
1988	56	Grandview	•											
Subtotal			5	0	3	5	1	3	1	2	2	1	5	3
1989	6	Cle Elum	•		•		•						•	
1989	19	Umtanum	•		•					•		•	•	
1989	26	Naches												
1989	32	Union Gap												
1989	50	Kiona												
1989	52	Sulphur Creek			•									
1989	56	Grandview												
Subtotal			2	0	3	0	1	0	0	1	0	1	2	0
Total			7	0	6	10	4	10	2	4	4	2	8	5



**Table 22.** Estimated cadmium loads in filtered-water samples at selected fixed sites, Yakima River Basin, Washington, 1987–90

[The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; loads reported as pounds per day; load values are based on calibration data collected from March 1987 to March 1990; bold lines represent the irrigation season, and lightly shaded cells represent the snowmelt portion of the irrigation season; non-irrigation season, October through March; Jan, January; Feb, February; Mar, March; Apr, April; Aug, August; Sept, September; Oct, October; Nov, November; Dec, December; --, indicates insufficient data;]

Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Daily mean
<b>Yakima River at Cle Elum</b>													
1987	--	--	0.8	0.9	1	2	3	3	0.7	0.3	0.2	0.3	--
1988	0.3	0.5	.6	.9	.7	1	3	3	1	.4	.6	.6	1.0
1989	.7	.5	.4	1	2	2	4	3	1	.4	.7	.7	1.4
1990	.8	.9	.8	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Umtanum</b>													
1987	.5	.8	2	2	3	4	5	5	1	.4	.3	.5	2.0
1988	.3	1	1	3	2	3	5	6	2	.7	1	1	2.2
1989	1	1	1	5	4	4	6	5	2	1	1	1	2.7
1990	1	2	2	6	5	7	6	6	2	--	--	--	--



**Figure 25.** Cadmium loads in filtered-water samples and in suspended-sediment samples for selected time periods at the Yakima River at Umtanum, Yakima River Basin, Washington, 1989 (the term “filtered water” represents the portion of a water sample passing through a nominal 0.45 micrometer pore-size filter; non-irrigation season occurs October through March; snowmelt season occurs April through May; irrigation season occurs June through September).

during the irrigation season and the non-irrigation season.

Cadmium concentrations in fish livers and Asiatic clams generally were less than 0.5 µg/g (table 15). Maximum concentrations in mountain whitefish (1.4 µg/g), largescale sucker (0.43 µg/g), and carp (2.5 µg/g) were measured in the lower Yakima Valley in the Yakima River below Toppenish Creek at RM 79.6 near Granger and in the Yakima River at Kiona. In 1989, cadmium concentrations were below the limit of determination in most fish samples; however, detectable concentrations were measured in several samples in the lower Yakima Valley portion of the main stem. For example, the cadmium concentration in mountain whitefish was 1.1 µg/g in the Yakima River at Kiona, and the concentration in largescale sucker was 0.6 µg/g in Sulphur Creek Wasteway near Sunnyside and 0.45 µg/g in the Yakima River at Kiona.

In aquatic plants, variations in cadmium concentrations are generally small. When measured, high concentrations do exist in waterways that carry irrigation return flow and (or) urban runoff. Cadmium concentrations as high as 3.3 µg/g were measured in

waterweed in Spring Creek at mouth at Whitstran as well as concentrations as high as 0.9 µg/g in Curlyleaf pondweed in Cherry Creek at Thrall (Fuhrer, Fluter, and others, 1994). Cadmium in plant tissue at the latter site and the detection of cadmium in filtered-water samples in the Yakima River at Umtanum (located just downstream from Cherry Creek) indicates that a source(s) of cadmium exists in the Kittitas Valley.

Cadmium concentrations in benthic insects do not exceed 0.25 µg/g (table 15). Several areas of cadmium enrichment, however, are evident in the mid-Yakima Valley and in the Kittitas Valley at the Cle Elum site (fig. 26). Cadmium concentrations as high as 0.25 µg/g were found in *Hydropsyche* spp. in Rattlesnake Creek above Little Rattlesnake Creek near Nile and in Rattlesnake Creek above North Fork Rattlesnake Creek near Nile. In relation to other sites in the Yakima River Basin, cadmium concentrations also were high in six other insect taxa in Rattlesnake Creek. Additionally, of the two locations on Rattlesnake Creek, concentrations were higher at the more upstream station (Rattlesnake Creek above North Fork Rattlesnake Creek near Nile) in four of the five taxa common to both locations (Fuhrer, Fluter, and others, 1994). Cadmium (1 µg/g) also was detected in the liver of rainbow trout from this site in 1989. These data indicate that an upstream source of cadmium is present in Rattlesnake Creek.


The median concentration of cadmium (0.2 µg/g) in *Hydropsyche* spp. at the Cle Elum site, although not as large as cadmium concentrations in caddisflies in Rattlesnake Creek, is twice that determined from the 23 sites sampled in 1990. The high concentration in caddisflies at Cle Elum also coincides with concentrations of cadmium in filtered water that were among the highest in the Yakima River Basin.

In 1989, rainbow trout sampled from the Waptus River at mouth near Roslyn had a cadmium concentration of 1.1 µg/g, but rainbow trout sampled in the Jungle Creek near mouth near Cle Elum had a cadmium concentration of only 0.4 µg/g. In the American River, the cadmium concentration in a predaceous stonefly (*Doroneuria* spp.) was 3 to 16 times higher than concentrations in this taxon from any other location in the Yakima River Basin. High concentrations of cadmium in the stonefly also coincide with the maximum concentration of cadmium in streambed sediment (0.8 µg/g) for the biological sampling sites. However,

cadmium enrichment in streambed sediment and stoneflies did not correspond to cadmium enrichment in the liver of rainbow trout or whole body sculpin—both were below the limit of determination (0.2 µg/g).

Cadmium concentrations in most biological samples from the Yakima River Basin indicate natural background concentrations and were similar to concentrations reported by Lynch and others (1988) and Cain and others (1992) for uncontaminated streams (table 40, at back of report).

Cadmium, found in the liver of rainbow and brook trout sampled from the Yakima River Basin in 1990, was near the mean concentration (0.18 µg/g) found in the livers of cutthroat trout from an uncontaminated stream (Moore and others, 1991). Concentrations of cadmium (1.0 µg/g) in rainbow trout from Rattlesnake Creek and the Waptus River at mouth near Roslyn were elevated relative to other sites in the Yakima River Basin and coincided with higher concentrations of cadmium in suspended-sediment and filtered-water samples, but these concentrations were not exceedingly high when compared to concentrations observed in trout and other freshwater fish in California's Toxic Substances Monitoring Program (Rasmussen, 1992; McClenegham and others, 1981). Cadmium found in the livers of other species of fish in the Yakima River Basin rarely exceeded 0.5 µg/g. For comparison, mean cadmium concentrations in northern pike and white sucker from a "control" lake were 0.4 and 0.9 µg/g, respectively (McFarlane and Franzin, 1980). Mean concentrations of cadmium, ranging from 0.12 to 0.92 µg/g (calculated from original wet-weight data; Barak and Mason, 1990), were found in liver from five species of fish: dace (*Leueiscus leueiscus*), chub (*Leueiscus cephalus*), tench (*Tinca tinca*), perch (*Perca fluviatilis*), and pike (*Esox lucius*). The species of fish were collected from two rivers in England. Cadmium in the livers of mountain whitefish and carp (1.43 and 2.55 µg/g, respectively) collected in the Yakima River at Kiona were slightly higher than concentrations noted in Barak and Mason (1990), possibly indicating an increase in bioavailable cadmium at the Kiona site. Some of these concentrations exceed and others are near the 85th percentile concentrations reported in California's Toxic Substances Monitoring Program (Rasmussen, 1992). Cadmium concentrations in Asiatic clams in the Yakima River Basin were indicative of an uncontaminated system. In the San Joaquin River in California, for example, the concentrations of cadmium found in Asiatic clams (*Corbicula fluminea*)



**Figure 26.** Distribution of cadmium concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are graphically represented for caddisflies; sample species: caddisflies [Trichoptera: Hydropsychidae *Hydropsyche* spp.]).

from an area receiving minimal inputs of cadmium were  $<1 \mu\text{g/g}$  (table 39, at back of report; Leland and Scudder, 1990; Luoma and others, 1990).

## Chromium and Nickel

Concentrations of chromium in streambed sediment of the Yakima River Basin range from 21 to  $212 \mu\text{g/g}$  (table 12) and slightly exceed the 8.5 to  $200 \mu\text{g/g}$  range of concentration which encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). The distribution of chromium concentrations at the biological sites generally exceeded those determined from analysis of fine-fraction streambed sediment in other river basins of the United States (table 34, at back of report).

Concentrations of nickel in streambed sediment of the Yakima River Basin ranged from 9.0 to  $260 \mu\text{g/g}$  (table 12); several sites had concentrations in excess of the 3.4 to  $66 \mu\text{g/g}$  range of concentration which encompasses 95 percent of Western United

States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). Chromium concentrations as high as 212 and  $210 \mu\text{g/g}$  were found in the Teanaway River below Forks near Cle Elum (site 5) and the Yakima River at Cle Elum (site 6), respectively. Similarly, nickel concentrations as high as 262 and  $150 \mu\text{g/g}$  were found at these same sites, respectively.


Geologic sources of chromium exist in the Wenatchee Mountains where the pre-Tertiary metamorphic and intrusive rocks geologic unit contains the mineral chromite (Fuhrer, McKenzie, and others, 1994; Tabor and others, 1982; Gualtieri and Simmons, 1989; and Simmons and others, 1983). Fuhrer, McKenzie, and others (1994) reported that concentrations of chromium affected by the mineral chromite, in streambed sediment of the pre-Tertiary rocks geologic unit, ranged from 1,200 to  $1,700 \mu\text{g/g}$ . Another natural source of chromium and nickel in the northern part of the Yakima River Basin is a 1.5-mile segment of the Cle Elum River between Big Boulder Creek and Camp Creek (upstream of Cle Elum Lake and in the pre-Tertiary rock geologic unit) which contains outcrops of nickel-chromium bearing ferruginous laterite

(a highly weathered, iron-rich subsoil) [Lucas, 1975]. The enrichment of chromium and nickel at the Yakima River at Cle Elum probably can be attributed to the presence of chromite and ferruginous laterite. Chromium and nickel enrichment along the geologic contact between the pre-Tertiary rocks geologic unit and the nonmarine sedimentary rocks geologic unit (Lucas, 1975) probably is responsible for the enrichment at the Teanaway River below Forks near Cle Elum. Enrichment of chromium and nickel in streambed sediment of the pre-Tertiary rocks and the nonmarine sedimentary rocks geologic units has been described in detail by Fuhrer, McKenzie, and others (1994).


In addition to chromium and nickel enrichment in some of the northern tributaries of the Yakima River Basin, the upper reach of the main stem has similar, but somewhat attenuated enrichment. The attenuation or concentration decrease along the main stem is attributed to sediment dilution—a decrease in element concentration by the mixing of enriched streambed sediment from (in this instance) a geologic source with less enriched sediment from another geologic source. For example, figure 27 shows enrichment of chromium in streambed sediment of the Cle Elum

River (620  $\mu\text{g/g}$ ) and in streambed sediment of the North Fork of the Teanaway River (212  $\mu\text{g/g}$ ); in addition, chromium was enriched along the main stem at the Yakima River at Cle Elum (210  $\mu\text{g/g}$ ) and again downstream at the Yakima River at Evergreen Farm (160  $\mu\text{g/g}$ ). The concentration (64  $\mu\text{g/g}$ ) of chromium at the Yakima River at Umtanum decreases sharply in comparison to upstream main-stem concentrations. The decrease in chromium concentration was attributed to dilution caused by sediment entering the main stem from Wilson Creek, located a short distance upstream of the Yakima River at Umtanum. Sediment entering the main stem from Wilson Creek was formed from chromium depleted Quaternary deposits and loess and Quaternary flood deposits and contained only 84 and 54  $\mu\text{g/g}$  of chromium, respectively (Fuhrer, McKenzie, and others, 1994). The similar behavior of nickel and chromium was expected because of the relation between nickel and chromium in streambed sediment that originates from the pre-Tertiary metamorphic and intrusive rocks geologic unit.

Concentrations of chromium in suspended sediment at the seven fixed sites ranged from 28 to 160  $\mu\text{g/g}$  (table 13 and fig. 28). The largest temporal varia-



**Figure 27.** Chromium concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987.



#### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.


- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value
- × 1.5 to 3 times the interquartile range from the 25-percentile value

**Figure 28.** Distribution of chromium concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).

tion for suspended chromium was measured in the Yakima River at Umtanum—the interquartile range for chromium was 37 µg/g (table 35, at back of report). In contrast, the smallest temporal variation for suspended chromium was found in Sulphur Creek Wasteway near Sunnyside, where the interquartile range was more than 6 times smaller than Umtanum. Interquartile ranges also were small for sites in the

lower Yakima Valley. The large difference between the distribution of chromium concentrations for the Yakima River at Cle Elum and the Naches River near North Yakima is attributed to geology. The Cle Elum Subbasin has geologic sources of chromium, as high as 1,700 µg/g in streambed sediment, whereas the Naches River Basin does not contain a significant geologic source of chromium (Fuhrer, McKenzie, and others, 1994).

The large temporal variation in chromium concentrations for the Yakima River at Umtanum was indicative of a change in the source of chromium, as well as sediment dilution from chromium-poor sediment in agricultural lands in the Kittitas Valley. Large concentrations of suspended chromium were measured during the non-irrigation season as well as during the early and mid-irrigation season at the Umtanum site. However, the concentrations of suspended chromium decreased sharply in the late irrigation season, during September and October (fig. 29), which coincides with the curtailment of reservoir releases upstream of the Umtanum site, including those from Cle Elum Lake. The lowering of streamflow in the Kittitas Valley is a water-management strategy designed to enhance salmon productivity by forcing spawning salmon to make redds in portions of the streambed that will be covered by water following the irrigation season. Numerous streambed-sediment samples collected in the Kittitas Valley indicate that the Cle Elum River drainage was a probable source of chromium to the main stem; furthermore, suspended-chromium samples from the Yakima River at Cle Elum (although sparse and collected only during the non-irrigation season) were similar in concentration to those found at the Umtanum site and indicated that the Cle Elum River drainage is a probable source of chromium to the main stem. Conversely, streambed samples collected in agricultural land-use areas of the Kittitas Valley indicate that streambed sediment entering the main stem from the Kittitas Valley (especially during the irrigation season) could decrease the suspended-chromium concentrations in the main stem (Fuhrer, McKenzie, and others, 1994). Decreases in the Cle Elum River drainage's contribution of chromium to the main stem, as well as concurrent increases in the proportion of chromium-depleted sediment from agricultural areas in Kittitas Valley, are possible causes for sharp declines in suspended-chromium concentrations.




**Figure 29.** Chromium concentrations in suspended sediment at the Yakima River at Umtanum and the Yakima River at Cle Elum, Yakima River Basin, Washington, 1987–90 (shaded area represents the period of time when streamflow is reduced from reservoirs upstream from the Yakima River at Cle Elum).

Effects from the curtailment of reservoir releases and the diluting effect of incoming sediment, also are measurable in the Yakima River above Ahtanum Creek at Union Gap. Although not as pronounced, concentrations of suspended chromium at Union Gap generally decrease during September and October each year, when reservoir releases from the upper Yakima Valley are curtailed. These decreases probably result from reductions of suspended sediment, enriched in chromium, from the Cle Elum River drainage during September and October, as well as the diluting effect of incoming suspended sediment, depleted in chromium, from tributaries such as Wilson Creek, Selah Creek, Moxee Drain, Wide Hollow Creek, and the Naches River.

As the distance between sites on the main stem and the sources of chromium in the Kittitas Valley increase, concentrations of suspended chromium decrease, especially during rain-on-snow events. Samples from the Yakima River above Ahtanum Creek at Union Gap following rain or snow storms—two in 1989 and one in 1990 (fig. 30)—displayed lower concentrations of chromium compared to monthly samples. Lower chromium concentrations at the Union Gap site were attributed to streambed sediment that was formed in the Columbia River Basalt Group Rocks geologic unit. The median streambed-sediment concentration (40 µg/g; Fuhrer, McKenzie, and others, 1994) in this unit is low, and during rain-on-snow

storms, erosion from this unit probably reduces the suspended-chromium concentrations at the Union Gap site. During one of the storms (May 10, 1989), suspended-chromium concentrations were measured in two small tributaries (Squaw and Burbank Creek) located upstream of the Union Gap site. These tributaries were formed in the Columbia River Basalt Group Rocks geologic unit and, as expected, the suspended-chromium concentrations in these tributaries (40 and 38 µg/g, respectively) are nearly identical to that in streambed sediment from the Columbia River Basalt Group Rocks geologic unit. The low concentrations of chromium in Squaw and Burbank Creek probably are indicative of concentrations in other mid-Yakima Valley tributaries that, when combined, result in lower chromium concentrations during event sampling at the Union Gap site.

During some storms, intrasite variations in suspended-chromium concentrations differed between storm peaks and the periods before and after storms. During a December 5, 1989 rain-on-snow storm (hydrologically affecting all main-stem sites), chromium concentrations in the Yakima River at Umtanum were higher near the peak of the storm than concentrations from monthly samples before and after the storm (fig. 29). Conversely, during the same storm, chromium concentrations in the Yakima River above Ahtanum Creek at Union Gap were lower near the storm's peak than concentrations from monthly sam-



**Figure 30.** Chromium concentrations in suspended sediment at the Yakima River above Ahtanum Creek at Union Gap, Yakima River Basin, Washington, 1987–90 (shaded area represents the period of time when streamflow is reduced from reservoirs upstream from the Yakima River at Cle Elum).

ples preceding and following the storm (fig. 30). Whether chromium concentrations increase or decrease near the peak of the storm probably is related to the magnitude of the chromium source and the proximity of that source to the sampling site. Chromium concentrations measured near the storm's peak at the Umtanum site probably increased because of the close proximity between the site and sources of chromium in the Cle Elum and Teanaway drainages in the Kittitas Valley. During the same storm, chromium concentrations probably decreased near the peak of the storm at the Union Gap site because chromium-rich sediment from the Kittitas Valley mixed with chromium-poor sediment that entered the main stem from the Columbia River Basalt Group Rocks geologic unit in the mid-Yakima Valley.


Chromium concentrations in filtered-water samples ranged from  $<0.5$  to  $1.1 \mu\text{g/L}$  at the seven fixed sites (table 14). Chromium measurements were limited temporally and numerically among fixed sites; 26 measurements were made quarterly, primarily in 1987. The highest chromium concentrations were measured in the Yakima River at Cle Elum and at Kiona ( $1.1$  and  $1.0 \mu\text{g/L}$ , respectively; table 37, at back of report). The median chromium concentration among fixed sites ( $<0.5 \mu\text{g/L}$ ) coincides with the

background concentration of chromium in inland water and is low in comparison to the median chromium concentration measured from the National Stream Quality Accounting Network of the USGS (table 36, at back of report).

Median concentrations of nickel in suspended sediment ranged from  $17$  to  $89 \mu\text{g/g}$  at the fixed sites (table 13) and the lowest and highest median values, respectively, were in the Naches River near North Yakima and Yakima River at Umtanum (fig. 31; and table 35, at back of report). Similar to chromium, the largest temporal variation in suspended nickel was measured in the Yakima River at Umtanum, where the interquartile range for suspended nickel was  $31 \mu\text{g/g}$ . Concentrations of nickel also were variable in the Yakima River at Union Gap, where the interquartile range was  $18 \mu\text{g/g}$ . The concentrations and temporal variability for suspended nickel were reduced greatly in the lower Yakima Valley.

The concentrations of suspended nickel in the Yakima River at Umtanum and the Yakima River at Cle Elum indicate geologic sources of nickel. Nickel- and chromium-enriched deposits exist for a distance of approximately 20 miles between the geologic contacts of the pre-Tertiary rocks and the nonmarine sedimentary rocks geologic unit (Lucas, 1975). Additionally,





#### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.

- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value
- × 1.5 to 3 times the interquartile range from the 25-percentile value

**Figure 31.** Distribution of nickel concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).


streambed sediment in the headwaters of the Cle Elum and Teanaway Rivers, respectively, contained nickel concentrations as high as 1,800 and 1,900 µg/g, respectively (Fuhrer, McKenzie, and others, 1994). Although sources of nickel exist in the Cle Elum and Teanaway River drainages, higher concentrations of nickel were measured in the main stem at the Umtanum site than at the Cle Elum site, because

higher concentrations of suspended nickel reached the main stem from the unregulated Teanaway River than from the Cle Elum River. The Teanaway River flows into the main stem (RM 176.1) downstream of the Yakima River at Cle Elum (RM 183.1) and upstream of the Yakima River at Umtanum (RM 140.4). Wilson Creek also flows into the main stem (RM 147), upstream of the Yakima River at Umtanum, but is an unlikely source of suspended nickel, because the median concentration of nickel in nine streambed-sediment samples from the alluvial sediment of the Kittitas Valley was 21 µg/g (Fuhrer, McKenzie, and others, 1994). This concentration is only one third the median concentration of nickel found in streambed sediment in the headwaters of the Teanaway and Cle Elum Rivers.

The potential contribution of suspended sediment from the Teanaway River to suspended-nickel concentrations in the Yakima River at Umtanum was evident during storms. During a January 29–31, 1989 storm, the daily mean streamflow increased from 505 to 1,290 ft<sup>3</sup>/s at the Yakima River at Cle Elum, from 202 to 931 ft<sup>3</sup>/s at the Teanaway River below Forks near Cle Elum, and from 891 to 2,610 ft<sup>3</sup>/s for the Yakima River at Umtanum. Near the peak of the storm on January 31, 1989, the Teanaway River accounted for 70 percent of the observed increase in streamflow between the Yakima River at Cle Elum and the Yakima River at Umtanum. The concentrations of suspended nickel near the peak of the storm increased from 66 µg/g at the Cle Elum site to 120 µg/g at the Umtanum site. During storms, future samplings, that include the Teanaway River, in addition to the Yakima River at Cle Elum, and the Yakima River at Umtanum are needed to measure the concentration and load of suspended nickel in the Teanaway River.

During storms, suspended-nickel concentrations are related inversely to the distance between the geologic source and the sampling site. The Yakima River at Cle Elum is affected by nickel and chromium in serpentinized peridotite, an ultramafic rock (Fuhrer, McKenzie, and others, 1994). During a December 5, 1989 rain-on-snow storm, the nickel concentration nearly doubled at the Cle Elum site near the peak of the storm (fig. 32) but decreased at the Union Gap site.


Except for storms, the temporal variability for concentrations of nickel at the Union Gap site was similar to the Umtanum site. Nickel concentrations between sites differ; however, at the Umtanum site, nickel concentrations consistently exceeded those at



**Figure 32.** Nickel concentrations in suspended sediment at the Yakima River above Ahtanum Creek at Union Gap and at the Yakima River at Umtanum, Yakima River Basin, Washington, 1987–90 ("Union Gap" represents Yakima River above Ahtanum Creek at Union Gap and "Umtanum" represents Yakima River at Umtanum).

the Union Gap site. The lower concentrations at the Union Gap site (RM 107.2) probably resulted from the mixing of nickel-depleted sediment, from tributaries like the Naches River (RM 116.3) and waterways that carry irrigation return flow such as Moxee Drain (RM 107.6), with nickel-enriched sediment from the Kittitas Valley. Concentrations of suspended nickel abruptly decreased in September and October 1987-89, after the curtailment of reservoir releases in the Kittitas Valley (fig. 32). Similar decreases, measured for suspended chromium, were expected because of existing chromium and nickel in ultramafic rock in the Kittitas Valley (Fuhrer, McKenzie, and others, 1994). Suspended-nickel concentrations at the Umtanum site also increase statistically ( $p \leq 0.001$ ) in proportion to the concentration of suspended chromium (fig. 33). This is consistent with the streambed-sediment chemistry (Fuhrer, McKenzie, and others, 1994) and geologic sources (Tabor and others, 1982; Gualtieri and Simmons, 1989) in the Kittitas Valley.

Chromium concentrations in fish and Asiatic clams typically were less than 2  $\mu\text{g/g}$  (table 15) and varied little among sites in the lower Yakima Valley and in tributaries in the mid-Yakima Valley (Fuhrer, Fluter, and others, 1994). Chromium concentrations in insects generally were lowest in Rattlesnake Creek and in Umtanum Creek near the mouth at Umtanum (table 23). Chromium enrichment, however, was measured in aquatic insects in the North Fork



#### EXPLANATION

Number of observations = 29  
 $Y = 0.74X + 38$   
 where  
 $Y$  = Chromium  
 $X$  = Nickel  
 Correlation coefficient = 0.82  
 Standard error of estimate  
 of mean value of  $Y = 15$

**Figure 33.** Chromium and nickel concentrations in suspended sediment at the Yakima River at Umtanum, Yakima River Basin, Washington, 1987–90.

**Table 23.** Comparison of low and high chromium concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91

[For filtered water and suspended sediment, the low- and high-concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For stream bed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an “H” in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an “L” in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an “\*” in the table. The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale-sucker livers and caddisflies; sample species: largescale sucker (*Catostomus macrocheilus*), caddisfly (*Hydropsyche* spp.), stonefly (*Hesperoperla* sp.), Asiatic clam (*Veneroida*: Corbiculidae *Corbicula fluminea*), and curlyleaf pondweed (*Potamogeton crispus*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Yakima River above Altanum Creek at Union Gap, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987–90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]

Site reference number	Site name	Filtered water	Sediment		Aquatic biota						
			Streambed	Suspended	Largescale- sucker liver	Insects			Asiatic clam	Curlyleaf pondweed	
						Caddisfly	Stonefly				
1	Waputis River at mouth near Roslyn	--	*	--	--	--	--	--	--	--	--
3	Jungle Creek near mouth near Cle Elum	--	H	--	--	--	--	--	--	--	--
4	North Fork Teanaway River below bridge at Dickey Creek Campground	--	--	--	--	--	H	--	--	--	--
5	Teanaway River below Forks near Cle Elum	--	H	--	--	--	--	--	--	--	--
6	Yakima River at Cle Elum	H	H	H	--	*	*	*	--	--	--
7	Naneum Creek below High Creek near Ellensburg	--	*	--	--	*	*	*	--	--	--
8	Taneum Creek at Taneum Meadow near Thorp	--	H	--	--	--	--	*	--	--	--
10	Little Naches River at mouth near Cliffdell	--	L	--	--	*	--	--	--	--	--
12	South Fork Manastash Creek near Ellensburg	--	H	--	--	--	*	*	--	--	--
13	American River at Hells Crossing near Nile	--	L	--	--	--	--	--	--	--	--
14	Cherry Creek above Wipple Wasteway at Thrall	--	L	--	--	*	--	--	--	--	--
16	Cherry Creek at Thrall	--	--	--	--	H	--	--	--	--	H
19	Yakima River at Umtanum	L	*	H	--	H	H	H	--	--	H
20	Umtanum Creek near mouth at Umtanum	--	L	--	--	L	--	--	--	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	--	L	--	--	L	L	L	--	--	--
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	--	*	--	--	L	L	L	--	--	--

**Table 23.** Comparison of low and high chromium concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Filtered water	Sediment		Aquatic biota				
			Streambed	Suspended	Largescale- sucker liver	Insects			Curlyleaf pondweed
						Caddisfly	Stonefly	Asiatic clam	
26	Naches River near North Yakima	*	H	L	*	L	--	--	--
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	--	L	--	--	L	--	--	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	--	*	--	--	*	--	--	*
30	Moxee Drain at Thorp Road near Union Gap	--	*	--	--	--	--	--	--
31	Ahtanum Creek at Union Gap	--	*	--	--	*	--	--	--
32	Yakima River above Ahtanum Creek at Union Gap	L	--	*	--	--	--	--	--
33	Yakima River at Parker	--	*	--	H	*	--	--	*
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	--	L	--	--	*	*	--	--
40	Granger Drain at mouth near Granger	--	*	--	--	H	--	--	--
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger	--	*	--	*	*	--	*	--
43	Toppenish Creek at Indian Church Road near Granger	--	*	--	--	--	--	--	--
47	Satus Creek at gage at Satus	--	*	--	L	H	--	--	L
48	Yakima River at river mile 72 above Satus Creek near Sunnyside	--	*	--	--	--	--	L	L
50	Yakima River at Kiona	H	*	*	*	*	--	*	*
52	Sulphur Creek Wasteway near Sunnyside	*	H	L	--	H	--	--	-
53	Satus Creek below Dry Creek near Toppenish	--	*	--	--	L	--	--	--
54	Spring Creek at mouth at Whitstran	--	*	--	--	H	--	*	--
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	*	*	*	*	*	--	H	*
57	Satus Creek above Wilson-Charley Canyon near Toppenish	--	*	--	--	*	*	--	--

Teanaway River (site 4). Depending on insect species, concentrations of chromium in the North Fork Teanaway River were 4 to 52 times higher (concentrations ranged from 2.2 to 33.4  $\mu\text{g/g}$ ) than the minimum concentrations measured in the Yakima River Basin. Additionally, among the fish collected in 1989, the highest chromium concentration (2.0  $\mu\text{g/g}$ ) was in fish collected from the Teanaway River below Forks near Cle Elum. Fish were not collected at the North Fork Teanaway (site 4).

Concentrations of chromium in most biological samples from the Yakima River Basin were within ranges observed in uncontaminated basins. However, relatively high concentrations were detected in benthic insects in the North Fork Teanaway River. The maximum concentration of chromium observed in benthic insects (33  $\mu\text{g/g}$  in a stonefly [*Perlinoidea* sp.]) was measured at the North Fork Teanaway River and exceeded concentrations in uncontaminated systems reported in other studies (table 40, at back of report).

Concentrations of chromium in Asiatic clams were low because of the large distance between geologic sources of chromium in the Kittitas Valley and the sites sampled for Asiatic clams in the lower Yakima Valley. Chromium concentrations in Asiatic clams from the Yakima River were generally <2  $\mu\text{g/g}$  and were similar to concentrations reported for uncontaminated areas of the San Joaquin River in California (Luoma and others, 1990; table 39, at back of report). By comparison, chromium concentrations of 4 to 15  $\mu\text{g/g}$  were detected in Asiatic clams in areas of the San Francisco Bay estuary that are known to have inputs of chromium from industrial sources (Luoma and others, 1990). The chromium concentration in rainbow trout liver, collected in 1989 from the Teanaway River below Forks near Cle Elum, also exceeds that for fish sampled in California's Toxic Substances Monitoring Program (Rasmussen, 1992).

The maximum concentration of nickel that was detected in rainbow trout liver did not exceed 0.36  $\mu\text{g/g}$  (Fuhrer, Fluter, and others, 1994). Nickel concentrations in Asiatic clams generally are higher than in fish, and concentrations are similar among sites (table 24). Nickel concentrations in fish samples also were similar among sites.

The maximum nickel concentration in benthic insects was found in the North Fork Teanaway River below bridge at Dickey Creek Campground (site 4). Nickel concentrations from insect taxa collected at site 4 were 43 to 102 times greater (concentrations ranged

from 6.4 to 76  $\mu\text{g/g}$ ) than insect taxa collected from other sites (Fuhrer, Fluter, and others, 1994). The caddisflies (*Hydropsyche* spp.) were not abundant enough at site 4 to sample. Additionally, levels of nickel enrichment for insects at site 4 were greater than any other element examined. Fish were not sampled from site 4 in 1990, and the limit of determination for rainbow trout liver collected in 1989 was too high to determine if trout livers were similarly enriched at site 4 (Fuhrer, Fluter, and others, 1994).

The presence of nickel in the Teanaway Subbasin also was evident in benthic insects in the main stem of the Yakima River. The caddisflies (*Hydropsyche* spp.) collected at the Umtanum site represented the maximum concentration (5.5  $\mu\text{g/g}$ ) for the Yakima River Basin and also coincide with the highest concentrations of suspended nickel in the basin. Collectively, high concentrations of nickel in caddisflies and suspended sediment underscores the importance of the Teanaway Subbasin as a source of nickel to the main stem. Nickel concentrations in caddisflies generally remained high in the main stem of the mid-Yakima and lower Yakima Valley, instead of decreasing with distance from the Teanaway Subbasin (fig. 34). This enrichment may result from secondary contributions of nickel from tributaries in the mid-Yakima and lower Yakima Valley. Insects sampled at the mouths of some tributaries (including Cherry Creek, Ahtanum Creek, Spring Creek, Sulphur Creek, and Granger Drain) were enriched moderately with nickel when compared to tributaries at higher elevations in the western part of the Yakima River Basin. For example, the nickel concentration (3.3  $\mu\text{g/g}$ ) in caddisflies (*Hydropsyche* spp.) collected near the mouth of Ahtanum Creek, which receives irrigation return flow and some urban runoff, was more than 4 times higher than at the upstream site (South Fork Ahtanum Creek above Conrad Ranch near Tampico) which does not receive irrigation return flow and is removed from urban effects. The presence of nickel in these lower Yakima Valley tributaries may indicate why nickel concentrations did not decrease down the main stem, as might be expected, if the Kittitas Valley is the sole source of nickel.

Few data are available on nickel in benthic insects; however, concentrations of nickel in several taxa in the North Fork Teanaway River greatly exceeded concentrations reported by Lynch and others (1988) for combined insect samples collected

**Table 24. Comparison of low and high nickel concentrations in sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987-91**


[For filtered water and suspended sediment, the low and high concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For streambed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an "H" in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an "L" in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an "\*" in the table. The term "filtered water" is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale-sucker livers and caddisflies; sample species: largescale sucker (*Catostomus macrocheilus*), caddisfly (*Hydropsyche* spp.), stonefly (*Hesperoperla* sp.), Asiatic clam (*Veneroida*: Corbiculidae *Corbicula fluminea*) and curlyleaf pondweed (*Potamogeton crispus*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Yakima River above Altanum Creek at Union Gap, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987-90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]

Site reference number	Site name	Sediment		Largescale-sucker liver	Aquatic biota		
		Streambed	Suspended		Caddisfly	Stonefly	Curlyleaf pondweed
1	Waptus River at mouth near Roslyn	*	--	--	--	--	--
3	Jungle Creek near mouth near Cle Elum	H	--	--	--	--	--
4	North Fork Teanaway River below bridge at Dickey Creek Campground	--	--	--	--	H	--
5	Teanaway River below Forks near Cle Elum	H	--	--	--	--	--
6	Yakima River at Cle Elum	H	H	--	*	H	--
7	Naneum Creek below High Creek near Ellensburg	*	--	--	L	*	--
8	Taneum Creek at Taneum Meadow near Thorp	H	--	--	--	*	--
10	Little Naches River at mouth near Cliffdell	L	--	--	L	--	--
12	South Fork Manastash Creek near Ellensburg	H	--	--	-	*	--
13	American River at Hells Crossing near Nile	L	--	--	--	--	--
14	Cherry Creek above Wipple Wasteway at Thrall	*	--	--	*	--	-
16	Cherry Creek at Thrall	--	--	--	*	--	*
19	Yakima River at Umtanum	H	H	--	H	*	H
20	Umtanum Creek near mouth at Umtanum	L	--	--	L	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	L	--	--	L	*	--

**Table 24.** Comparison of low and high nickel concentrations in sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Sediment		Aquatic biota				
		Streambed	Suspended	Largescale- sucker liver	Insects	Curlyleaf pondweed	Stonefly	
					Caddisfly			
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	*	--	--	L	L	--	--
26	Naches River near North Yakima	*	L	L	*	--	--	--
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	L	--	--	*	--	--	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	*	--	--	*	--	--	L
30	Moxee Drain at Thorp Road near Union Gap	*	--	--	--	--	--	--
31	Ahtanum Creek at Union Gap	L	--	--	H	--	--	--
32	Yakima River above Ahtanum Creek at Union Gap	--	*	--	--	--	--	--
33	Yakima River at Parker	*	--	H	*	--	--	*
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	L	--	--	L	*	--	--
40	Granger Drain at mouth near Granger	L	--	--	H	--	--	--
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger	*	--	*	*	--	--	--
43	Toppenish Creek at Indian Church Road near Granger	*	--	--	--	--	--	--
47	Satus Creek at gage at Satus	*	--	L	H	--	--	L
48	Yakima River at river mile 72 above Satus Creek near Sunnyside	*	--	--	--	--	--	*
50	Yakima River at Kiona	*	*	L	*	--	--	H
52	Sulphur Creek Wasteway near Sunnyside	H	L	--	H	--	--	--
53	Satus Creek below Dry Creek near Toppenish	*	--	--	*	--	--	--
54	Spring Creek at mouth at Whitstran	*	--	--	H	--	--	--
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	*	*	L	*	--	--	*
57	Satus Creek above Wilson-Charley Canyon near Toppenish	*	-	--	L	*	--	--





**Figure 34.** Nickel concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987–90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are graphically represented for caddisflies; sample species: caddisflies [Trichoptera: Hydropsychidae *Hydropsyche* spp.]).

upstream of a molybdenum mine (table 40, at back of report).


Nickel concentrations in Asiatic clams (*Corbicula* sp.) in the Yakima River were consistent with concentrations reported by Leland and Scudder (1990) for the San Joaquin River in California (table 39, at back of report). The consistently low concentrations of nickel in the Yakima River and the San Joaquin River may be indicative of natural concentrations for Asiatic clams. Nickel in fish liver sampled from the Yakima River Basin was within the 85th percentile concentration for fish sampled in California's Toxic Substances Monitoring Program (Rasmussen, 1992).

## Cobalt

Cobalt concentrations in fish and Asiatic clams were generally less than  $1 \mu\text{g/g}$ , and concentrations of cobalt in benthic insects were higher and more variable in the Yakima River Basin (table 15). The highest concentrations in benthic insects were in the

South Fork of Ahtanum Creek (site 34) and in Ahtanum Creek at Union Gap (site 31). At these sites, concentrations in caddisflies (*Hydropsyche* spp.) were 9.1 and  $6.2 \mu\text{g/g}$ , respectively, and approximately 6 to 9 times higher than the lowest cobalt concentration in the Yakima River Basin (fig. 35). Comparative data for fish are not available for these sites 31 and 34, but cobalt in streambed sediment at site 34 was the maximum concentration among biological sampling sites. Cobalt concentrations at site 34 in benthic insects were as much as 3 to 6 times higher than mean concentrations reported for uncontaminated streams (Elwood and others, 1976; Smock, 1983; table 40, at back of report), indicating some benthic insects in the Yakima River Basin have been enriched from geologic sources of cobalt.

Maximum concentrations of cobalt also were observed in other taxa (for example, caddisfly [*Arctopsyche* sp.], and stoneflies [*Doroneuria* sp., *Pteronarcys* sp., and *Hesperoperla* sp.]) at either the South Fork of Ahtanum Creek or Ahtanum Creek (Fuhrer, Fluter, and others, 1994). High concentrations of cobalt also were measured in caddisflies (*Hydro-*



**Figure 35.** Cobalt concentrations in caddisflies and streambed sediment, Yakima River Basin, Washington, 1987-90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are graphically represented for caddisflies; sample species: caddisflies [Trichoptera: Hydropsychidae *Hydropsyche* spp.]).

*psyche* spp.) in Satus Creek below Dry Creek, Granger Drain at mouth near Granger, and Sulphur Creek Wasteway near Sunnyside. In the main stem of the Yakima River, however, concentrations in insects, fish and Asiatic clams (*Corbicula* sp.) were consistent and relatively low among sites. Inputs of cobalt from some of the enriched tributaries in the basin are apparently diluted by other tributaries carrying lower cobalt concentrations.

## Copper

Concentrations of copper in Yakima River Basin streambed sediment ranged from 17 to 96  $\mu\text{g/g}$  (table 12) and slightly exceed the 4.9 to 90  $\mu\text{g/g}$  range of concentration which encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). Concentrations of copper as high as 93  $\mu\text{g/g}$  were found in streambed sediment of the Naches River near North Yakima (site 26) and Wide Hollow Creek at West Valley Middle School near Ahtanum (site 27). The 93  $\mu\text{g/g}$  of copper

measured near West Valley Middle School was more than twice the concentration in Wide Hollow Creek (near the mouth) at the old Sewage Treatment Plant and probably is the result of light urbanization and industrialization. The median concentration of copper at the biological sampling sites (20  $\mu\text{g/g}$ ) was as much as two times that found in fine-fraction sediment from other river basins in the United States, but the concentration was similar to that found in fine-fraction sediment from randomly sampled lower-order streams in the Yakima River Basin (table 34, at back of report).

Another source of suspended copper in the Yakima River Basin was from the use of highly insoluble, copper-bearing fungicides such as copper hydroxide and copper oxychloride (Dennis Johnson, Washington State University, Prosser Experimental Station, oral commun., 1992). These chemicals are registered for a broad array of uses, including applications on alfalfa, grapes, and peaches (Poplyk, 1989). Although copper has a number of agricultural uses in the basin and is known to readily associate with clay and organic matter in streambed sediment, copper concentrations in streambed sediment of the main stem—as previously mentioned—are within the


expected 95-percent range of copper concentrations found in soils of the Western United States. However, there does appear to be a geologic source of copper within the Naches River drainage (fig. 36). In addition to enrichment at the Naches River site, other sites including American River at Hells Crossing near Nile (site 13), Little Naches River at mouth near Cliffdell (site 10), Rattlesnake Creek above North Fork Rattlesnake Creek near Nile (site 22), and Rattlesnake Creek above Little Rattlesnake Creek near Nile (site 21)—all of which flow to the Naches River—have copper concentrations as high as 79, 48, 43, and 39  $\mu\text{g/g}$ , respectively. Geologic sources of copper exist in the basin; for example, copper enrichment as high as 150  $\mu\text{g/g}$  was reported in streambed sediment of a tributary to Rattlesnake Creek (Fuhrer, McKenzie, and others, 1994). Areas with reported copper enrichment also contained anomalous concentrations of arsenic and zinc; these areas of copper enrichment have been described by Simmons and others (1983) as andesitic tuff and breccia of the Ohanapecosh Formation.

Median concentrations of copper in suspended sediment ranged from 40 to 60  $\mu\text{g/g}$  at the seven fixed


sites and maximum concentrations were as high as 680  $\mu\text{g/g}$  in the Yakima River at Kiona (table 13). The largest variations of copper concentrations were found in Sulphur Creek Wasteway near Sunnyside—the interquartile range for suspended copper was 32  $\mu\text{g/g}$ .

Concentrations of suspended copper in Sulphur Creek Wasteway during the non-irrigation season generally were lower than concentrations measured during the irrigation season (fig. 37).

Concentrations of suspended copper were significantly correlated ( $\rho = 0.0001$ ) to the percentage of suspended sediment finer than 62  $\mu\text{m}$  in diameter and also were significantly correlated negatively ( $\rho = 0.0001$ ) to the concentration of suspended sediment. Thus, high concentrations of suspended copper, associated with an increasing percentage of small-grained-sized suspended sediment, were typical of conditions during the non-irrigation season. Conversely, the lower concentrations of suspended copper, associated with coarse-grained-sized suspended sediment and high concentrations of suspended sediment in Sulphur Creek Wasteway were typical of the irrigation season. As expected, the concentrations of suspended copper



**Figure 36.** Copper concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987.




**Figure 37.** Copper concentrations in suspended sediment at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90 (shaded area represents the irrigation season).

during the irrigation season are similar to the median concentrations of copper (21 µg/g) measured in streambed sediment of the Sunnyside Subbasin by Fuhrer, McKenzie, and others (1994).

In most cases, the concentrations of suspended copper in Sulphur Creek Wasteway were enhanced by increasing quantities of fine-grained-sized suspended sediment. This is typical of sediment dilution (the mixing of copper-rich, fine-grained-size suspended sediment with large quantities of copper-poor, coarse-grained-size suspended sediment) rather than changes in the source of copper between the irrigation and the non-irrigation season. However, two suspended-copper concentrations were in excess of concentrations expected based on relations to particle size and were considered anomalous (fig. 38). These anomalies may be typical of extreme temporal variability that is a characteristic of drainages receiving trace elements from multiple sources, including agricultural runoff, urban runoff, and sewage-treatment plant effluent.

A concentration of 190 µg/g of suspended copper was measured on June 21, 1988 during the irrigation season at Sulphur Creek Wasteway near



**Figure 38.** Copper concentrations in suspended sediment and the percentage of suspended sediment finer than 62 micrometers in diameter at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1987–90.

Sunnyside. This copper concentration was higher than expected based on the relation between copper and fine-grained sized suspended sediment. The concentration of suspended lead measured in the same sample had a near maximum (for Sulphur Creek Wasteway) of 41 µg/g—similar enrichment existed for cadmium. High concentrations of copper, lead, and cadmium may be indicative of sampling during a period of effluent discharge from the Sunnyside Sewage Treatment Plant or urban runoff from the city of Sunnyside. Although no data exist to characterize the trace-element composition of Sunnyside Sewage Treatment Plant effluent, data available for the city of Yakima's Sewage Treatment Plant indicate that unfiltered-water samples of copper in treated effluent ranged from 8 to 30 µg/L (J. Schnebly, City of Yakima Wastewater Treatment Plant, written commun., 1990). Assuming the latter concentration also, at times, exists in Sunnyside Sewage Treatment Plant effluent and using the suspended-sediment concentration for the June 21 sampling at the Sulphur Creek site, effluent from only the sewage treatment plant could account for more than one-half the 190 µg/g of suspended copper measured at the Sulphur Creek site. Statistically significant correlations exist between the concentration of suspended copper in Sulphur Creek Wasteway and arsenic ( $p \leq 0.001$ ), cadmium ( $p \leq 0.001$ ), lead ( $p \leq 0.001$ ), and silver ( $p \leq 0.001$ ). These elements

commonly are associated with sewage-treatment-plant effluent; additionally, these trace elements are attributed to corrosion of pipes in urban-water-supply networks (Forstner and Wittman, 1979, p. 44).

Suspended-copper loads in Sulphur Creek Wasteway near Sunnyside, unlike suspended-copper concentrations, are larger during the irrigation season than during the non-irrigation season. In 1989 for example, the irrigation-season load of copper at the Sulphur Creek site was 3 times higher than the non-irrigation season load (table 25). Suspended-copper loads also were high at the Sulphur Creek site during the snowmelt season; however, a proportion of the snowmelt load results from the spillage of excess water from Roza Canal to Sulphur Creek Wasteway, rather than from only agricultural runoff. The quantity of water spilled is ungaged; consequently, the associated load cannot be accurately determined.

Annual loads of suspended copper were variable among main-stem sites and, without exception, suspended-copper loads were higher in 1989 than in 1988 (table 25), because annual-average streamflow for the Yakima River at Kiona in 1989 was 20 percent higher than in 1988. Between the Yakima River at Umtanum and the Yakima River above Ahtanum Creek at Union Gap, annual loads of suspended copper increased more than 2 times. Loads decreased between Union Gap and Grandview, probably as a result of irrigation-season diversions to Wapato and Sunnyside Canals. Loads increased from Grandview to Kiona and again, the largest increase was measured in 1989.

The suspended-copper load doubled between the Umtanum site and the Union Gap site principally from contributions of copper over a 9.1-mile reach that extends from below the Naches River to the Yakima River above Ahtanum Creek at Union Gap as well as from the Naches River (the largest tributary in the Yakima River Basin). Over this 9.1-mile reach, sources of copper exist in Wide Hollow and the Moxee Subbasins. Copper-bearing fungicides are used in both subbasins (J.F. Rinella, unpub. data, 1996); additionally, urban runoff and effluent from the city of Yakima's sewage-treatment plant are potential sources of copper over this reach. In 1989, the Naches River accounted for about one-fourth of the annual load of copper measured at the Union Gap site; the 9.1-mile reach<sup>8</sup>, however, accounted for nearly one-half the annual load measured at the Union Gap site. During the irrigation season and the non-irrigation season, the

loads over the 9.1-mile reach were similar (11 pounds per day and 12 pounds per day, respectively).

For the Naches River near North Yakima, the Yakima River at Euclid Bridge at RM 55 near Grandview, and the Yakima River at Kiona, a majority of the annual-suspended-copper load was transported during the snowmelt season (table 25). In 1989, more than one-half the annual copper load for the Naches, Grandview, and Kiona site was transported during the snowmelt season. In the Yakima River at Umtanum, however, flow augmentation (by storage reservoirs in the Kittitas Valley) probably affects the irrigation-season load. In 1989, more than one-half the suspended-copper load at the Umtanum site was transported during the irrigation season.

Copper was determined in filtered-water samples by atomic-absorption spectrometry with graphite furnace and ranged in concentration from <0.5 to 20 µg/L at the fixed sites (table 14). Copper concentrations, determined by induction-coupled argon plasma at the 45 synoptic sites in July and November 1987, were all below the limit of determination (10 µg/L). The highest concentrations of copper, typically, were measured in the Kittitas Valley. The distribution of copper concentrations for the Yakima River at Umtanum for example, exceeded those of the other fixed sites (fig. 39; and table 37, at back of report). Additionally, 25 percent of the copper concentrations at the Umtanum site exceeded background concentrations in inland waters (table 36, at back of report). Although other fixed sites are in close proximity to urban sources, none of these sites exhibited percentile values of copper similar or as high as the Umtanum site. For example, although the Yakima River above Ahtanum Creek at Union Gap is located downstream of the city of Yakima, the 75th percentile value for copper at this site (0.9 µg/L) is smaller than the median copper value for the Umtanum site (1.3 µg/L).

The two highest copper concentrations, 20 µg/L on October 12, 1989 and 7.4 µg/L on November 15, 1989, were measured in the Yakima River at Umtanum—these copper concentrations exceeded the acute and chronic criterion, respectively, derived for

---


<sup>8</sup>The load for the reach was determined by subtracting the sum of the loads at the Naches and Umtanum sites from the load at the Union Gap site; a correction was applied to the Umtanum-site load to account for streamflow diverted around the Union Gap site (by way of Roza Canal) during the irrigation season.

**Table 25. Estimated copper loads in suspended sediment at selected-fixed sites, Yakima River Basin, Washington, 1987–90**

[Loads reported as pounds per day; load values are based on calibration data collected from March 1987 to March 1990; --, indicates insufficient data; Jan, January; Feb, February; Mar, March; Apr, April; Aug, August; Sept, September; Oct, October; Nov, November; Dec, December; bold lines represent the irrigation season, and lightly shaded cells represent the snowmelt portion of the irrigation season; non-irrigation season, October through March]

Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Daily mean
<b>Yakima River at Umtanum</b>													
1987	2	3	12	12	14	23	28	27	5	1	1	2	10.8
1988	1	4	5	15	9	15	29	34	12	2	4	5	11.2
1989	5	4	6	28	25	24	35	30	10	5	5	5	15.2
1990	7	9	11	34	26	42	33	36	9	--	--	--	--
<b>Naches River near North Yakima</b>													
1987	--	--	5	16	65	6	1	.3	5	2	.2	.8	--
1988	.4	.9	2	17	21	12	2	1	10	4	1	1	6.0
1989	1	.5	2	27	28	19	1	.8	9	2	.7	2	7.8
1990	9	4	5	--	--	--	--	--	--	--	--	--	--
<b>Yakima River above Ahtanum Creek at Union Gap</b>													
1987	10	15	43	42	70	40	36	33	22	12	5	11	28.2
1988	8	15	15	49	40	40	39	41	30	15	17	18	27.2
1989	19	16	25	82	70	54	41	36	28	15	17	20	35.2
1990	28	28	34	85	64	94	44	46	33	--	--	--	--
<b>Sulphur Creek Wasteway near Sunnyside</b>													
1987	--	--	5	14	15	12	.9	5	3	1	.7	.8	--
1988	1	2	7	12	14	14	7	5	4	2	.8	.8	6.0
1989	1	2	5	18	21	14	9	7	5	2	.8	.8	7.2
1990	1	1	6	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Euclid Bridge at river mile 55 near Grandview</b>													
1987	--	--	60	36	97	12	9	7	6	8	4	11	--
1988	6	11	10	46	23	18	8	10	11	13	19	11	15.6
1989	11	8	31	163	100	27	13	13	12	12	21	18	34.3
1990	32	28	32	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Kiona</b>													
1987	3	7	85	31	137	12	10	8	5	7	3	10	26.5
1988	3	7	6	59	30	24	9	11	13	10	15	8	16.2
1989	6	4	23	226	130	25	12	14	12	14	18	15	41.6
1990	28	16	20	135	72	396	14	62	23	--	--	--	--





**Figure 39.** Distribution of copper concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90 (“Cle Elum” represents Yakima River at Cle Elum; “Umtanum” represents Yakima River at Umtanum; “Naches” represents Naches River near North Yakima; “Union Gap” represents Yakima River above Ahtanum Creek at Union Gap; “Sulphur Creek” represents Sulphur Creek Wasteway near Sunnyside; “Grandview” represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and “Kiona” represents Yakima River at Kiona).

the protection of freshwater aquatic life (table 7). Additionally, these two concentrations of copper were measured after the curtailment of streamflow, which usually occurs during mid-September, from reservoirs in the Kittitas Valley. Mean monthly streamflows at the Umtanum site for October and November 1989 were about 1,000 ft<sup>3</sup>/s and represented about 25 percent of mean monthly streamflows measured during the 1989-irrigation season. The anomalous 7.4 µg/L of copper at the Umtanum site coincided with a

copper concentration of 14 µg/L measured the previous day (November 14, 1989) in the Yakima River at Cle Elum. The copper concentration (14 µg/L) at the Cle Elum site was the maximum value for the period 1987–89. During storms, the copper-bearing pre-Tertiary metamorphic and intrusive rocks geologic unit in the Cle Elum Subbasin (Fuhrer, McKenzie, and others, 1994) may be a potential source of dissolved copper to the main stem. The Cle Elum and the Umtanum sites were sampled in a 10-day interval in which streamflow increased more than 3 times. The anomalies at both sites were probably the result of an early winter storm that affected the Kittitas Valley.

The anomalous 20 µg/L of copper at the Umtanum site did not coincide with a storm event and probably resulted from a local anthropogenic source. The upstream concentration of copper the previous day (October 11, 1989) at the Cle Elum site was low (0.6 µg/L) and was not indicative of a geologic source which may have affected the Umtanum site. High concentrations of copper during the post-irrigation season probably resulted from smaller streamflow in the main stem and, consequently, less dilution water for copper entering the main stem from anthropogenic sources. The Umtanum site is located 6.3 miles downstream from Wilson Creek, which carries irrigation return flow and urban runoff from the Kittitas Valley to the main stem. During 1989, only 5 pounds per day of copper was applied to crops in the Kittitas Irrigation District (Rinella and others, 1992). This application rate is small when compared to the approximately 440 pounds per day applied to crops in the mid- and lower Yakima Valley. Urban sources of copper, measurable at Umtanum, may include the city of Ellensburg’s street runoff that flows into Wilson Creek, but probably does not include significant quantities of effluent from Ellensburg’s Sewage Treatment Plant. The copper in Ellensburg’s Sewage Treatment Plant effluent is small (3 µg/L; Reif, 1989) and streamflow from the plant accounts for less than one-half a percent of the streamflow at the Umtanum site during October and November 1989. Concentrations of copper and lead, in streambed sediment affected by urban runoff, were as large as 65 µg/g and 130 µg/g in Wilson Creek (Fuhrer, McKenzie, and others, 1994). In an oxygen-depleted environment, streambed sediment becomes anaerobic and dissolved copper may increase greatly in the sediment-pore water (Forstner and Whittman, 1979). Oftentimes, the concentration of copper in pore




water of sediment is greater than that predicted based on copper's solubility product with sulfide complexes; such behavior is attributed to the formation of humic-copper complexes (Forstner and Wittmann, 1979, p. 252–253; Moore, 1991, p. 109–139). In July 1987, dissolved oxygen in stream water at the Wilson Creek site was depleted to 72 percent of saturation. Releases of pore water from Wilson Creek streambed sediment may be a potential source of copper to the main stem during periods of low streamflow.

The distribution of copper concentrations at the Cle Elum site was similar to that for the Umtanum site. Higher concentrations generally were associated with winter-storms—storm concentrations of copper at Cle Elum were as large as 14  $\mu\text{g/L}$  (fig. 40). Additionally, copper concentrations in four of the five sampled storms exceeded acute and chronic criteria established for the protection of freshwater aquatic life (table 7). Although flow augmentation from reservoirs in the Kittitas Valley produces streamflow that varies from approximately 1,000  $\text{ft}^3/\text{s}$  to 3,500  $\text{ft}^3/\text{s}$ ; associated copper concentrations only varied slightly


and generally were low from April through September.

The spatial coverage for dissolved copper in the Kittitas Valley was inadequate to separate geologic sources of copper from anthropogenic sources. Storm sampling, however, suggests that urban runoff also may be a potential source of copper at the Cle Elum site. This site was sampled on January 9 and 10, 1990 during a storm; samples were collected on the rising limb and near the peak of the storm hydrograph, and concentrations of copper were measured as 5.3  $\mu\text{g/L}$  and 3.0  $\mu\text{g/L}$ , respectively (fig. 41). A similar pattern was measured during the same storm for suspended-copper concentrations—concentrations on the rising limb and near the peak of the storm were 67 and 47  $\mu\text{g/g}$ , respectively. The Cle Elum site, located 0.15 miles downstream from Crystal Creek, receives urban runoff from the towns of Roslyn and Cle Elum, sewage-treatment-plant effluent from the town of Roslyn, and runoff from numerous open-pit coal mines east of Roslyn. During the storm, Crystal Creek created a turbidity plume which, during the rising limb of the storm, obscured the shallow river bottom in the main stem at the Cle Elum site. Streambed sediment in the Crystal Creek drainage probably is not a geologic source of copper, because the average concentration of 33  $\mu\text{g/g}$  of copper in streambed sediment (Fuhrer, McKenzie, and others, 1994) was low in comparison to storm concentrations of suspended copper at the Cle Elum site. Considering the close proximity of the Cle Elum site to the inflow of Crystal Creek and the apparent absence of a large geological source of copper in the Crystal Creek drainage, urban runoff may be a source of copper measurable in filtered-water samples at the Cle Elum site.

Annual loads of dissolved copper generally were consistent among main-stem sites. The largest difference in loads of dissolved copper was measured between the Yakima River at Cle Elum and the Yakima River at Umtanum, where loads increased by about 2 times (table 26). Increases between these sites were not confined to any particular hydrologic season and were uniformly distributed during the snowmelt, irrigation, and non-irrigation seasons. Within 3.5 miles downstream of the Union Gap site, more than 80 percent of the copper load is diverted into the Wapato and Sunnyside Canals during the irrigation season. Consequently, the irrigation-season load at the Grandview and Kiona sites is affected by irrigation return flow downstream of Parker.



**Figure 40.** Relation between copper concentrations in filtered-water samples and streamflow for selected time periods at the Yakima River at Cle Elum, Yakima River Basin, Washington, 1987–90 (irrigation season occurs June through September; snowmelt season occurs April through May; non-irrigation season occurs October through March).



**Figure 41.** Copper concentrations in filtered-water samples for the Yakima River at Cle Elum and the Yakima River at Umtanum, Yakima River Basin, Washington, 1989–90.

Copper-based pesticides are applied on an annual basis in the lower Yakima Valley. During 1989, an estimated average of 460 pounds per day of copper in pesticide formulations was applied to the Yakima River Basin—major crops that received pesticide applications were hops, cherries, and pears (J.F. Rinella, unpub. data, 1996). Although 330 pounds per day (75 percent of the total copper in pesticide formulations) was applied to the lower Yakima Valley, no appreciable increase in copper load or copper concentration was noted during the irrigation season at the Grandview and Kiona sites. For example, during the 4-month irrigation season in 1989, the copper load at


the Grandview site was only 1,281 pounds, which accounted for only 25 percent of the annual load. Similarly, no appreciable increase in suspended-copper load was noted earlier during the irrigation season at either of these sites. In fact, the total load of copper (in filtered water and suspended sediment) in the Yakima River at Kiona in 1989 represented only 13 percent of that applied to the basin in pesticide formulations.

The increase in copper concentration attributable to pesticide application was estimated for the agricultural soils of the Moxee Subbasin. The estimate was made to determine if pesticide applications were large enough to be measurable in streambed sediment of Moxee Drain. Based on an application rate for Moxee Subbasin of 88 pounds of copper per day and an agricultural land-use area of 40 mi<sup>2</sup> (J.F. Rinella, unpub. data, 1996), soil concentrations were calculated for pesticide-application periods of 10, 20, and 30 years. To determine the volume of soils potentially affected by pesticide applications, a till depth of 1 foot was assumed. Furthermore, as a conservative estimate, 100 percent of the copper in the pesticide formulation was assumed to be sorbed to soil. The concentrations of copper in agricultural soils of the Moxee Subbasin after 10, 20, and 30 years of pesticide application would be expected to be 1.7, 3.5, and 5.2 µg/g, respectively. By the time soils are eroded from agricultural fields and mixed with streambed sediment that contains copper at background concentrations, the effects attributable to pesticide applications probably are not measurable in streambed sediment. Similar findings were true for other agriculturally affected subbasins such as Ahtanum and Wide Hollow. These results indicate that copper in pesticide formulations is not likely to accumulate in significant quantities in streambed sediment nor will pesticide formulations represent a significant source or flux of copper leaving the Yakima River Basin.

The Cle Elum, Umtanum, and Union Gap sites have irrigation-season copper loads that account for a large proportion of their respective annual loads and that are large compared to snowmelt-season and non-irrigation-season loads (fig. 42). In 1989 for example, the irrigation season accounted for about 59 percent of the annual copper load and about 56 percent of the annual streamflow at the Cle Elum site. The similarity between the percentages of load and streamflow was expected because copper concentrations only varied slightly with streamflow (with the exception of storms) and, as stated above, generally were low from

**Table 26.** Estimated copper loads in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90  
 [The term “filtered water” is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; loads reported as pounds per day; load values are based on calibration data collected from March 1987 to March 1990; --, indicates insufficient data; Jan, January; Feb, February; Mar, March; Apr, April; Aug, August; Sept, September; Oct, October; Nov, November; Dec, December; bold lines represent the irrigation season, and lightly shaded cells represent the snowmelt portion of the irrigation season; non-irrigation season, October through March]

Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Daily mean
<b>Yakima River at Cle Elum</b>													
1987	--	--	5	4	4	13	22	29	4	1	0.8	2	--
1988	1	3	3	4	1	5	21	33	9	2	5	7	7.8
1989	8	4	1	8	11	13	28	29	8	2	9	9	10.8
1990	12	11	5	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Umtanum</b>													
1987	5	6	16	12	12	20	28	36	10	3	3	6	13.1
1988	3	7	7	14	8	13	29	46	21	7	12	13	15.0
1989	13	9	8	28	21	21	36	41	19	12	15	15	19.8
1990	17	17	14	33	22	36	34	48	13	--	--	--	--
<b>Naches River near North Yakima</b>													
1987	--	--	4	7	17	5	2	1	4	2	1	1	--
1988	1	2	2	8	9	7	2	2	6	3	2	2	3.8
1989	2	1	3	10	11	8	2	2	6	2	1	3	4.2
1990	5	4	4	--	--	--	--	--	--	--	--	--	--
<b>Yakima River above Ahtanum Creek at Union Gap</b>													
1987	7	9	20	20	29	18	16	14	10	6	4	6	13.3
1988	5	9	10	22	19	18	17	16	12	8	8	9	12.8
1989	10	10	14	32	29	23	18	15	12	8	8	10	15.8
1990	13	14	17	34	27	34	19	18	13	--	--	--	--
<b>Sulphur Creek Wasteway near Sunnyside</b>													
1987	--	--	1	2	2	2	2	1	1	1	.4	.4	--
1988	.4	.4	1	1	2	2	1	1	2	1	.5	.4	1.0
1989	.4	.4	1	2	2	2	2	2	2	1	.5	.4	1.3
1990	.5	.4	1	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Euclid Bridge at river mile 55 near Grandview</b>													
1987	--	--	14	15	17	9	8	8	8	9	8	11	--
1988	11	13	12	15	12	10	8	8	9	10	13	12	11.1
1989	13	12	16	23	18	12	9	9	10	10	14	14	13.3
1990	16	16	17	--	--	--	--	--	--	--	--	--	--
<b>Yakima River at Kiona</b>													
1987	14	18	33	20	24	10	8	8	8	10	10	15	14.8
1988	13	17	14	22	15	12	8	9	11	12	16	16	13.8
1989	17	15	22	37	24	13	9	10	10	13	17	19	17.2
1990	23	22	22	33	21	32	10	13	12	--	--	--	--




**Figure 42.** Percentages of the annual load of copper in filtered-water samples for selected time periods at selected fixed sites, Yakima River Basin, Washington, 1989 (non-irrigation season occurs October through March; snowmelt season occurs April through May; irrigation season occurs June through September).

April through September (irrigation and snowmelt seasons; fig. 40). Conversely, the Grandview and Kiona sites had irrigation-season copper loads that accounted for smaller proportions of their respective annual loads and that, furthermore, were small in comparison to snowmelt-season and non-irrigation season loads (fig. 42). At the Grandview site in 1989, the irrigation season accounted for about 25 percent of the annual copper load and about 22 percent of the annual streamflow; again, as with the Cle Elum site, similarity existed between the percentages of load and streamflow. Because copper concentrations varied only slightly with streamflow, variations in streamflow greatly affected seasonal copper loads. Therefore, copper loads generally are controlled by flow regulation; loads in the main stem increase in the Kittitas Valley as a result of reservoir releases during the irrigation season and decrease in the lower Yakima Valley as a result of streamflow diversions to irrigation canals.

Copper can be transported in dissolved and suspended forms. The suspended form, however, is the major transport mechanism for copper at the Union Gap, Sulphur Creek, Naches, Grandview, and Kiona sites (fig. 43). The dominance of the suspended load is most notable during the snowmelt season and least notable during the non-irrigation season. The largest difference between dissolved and suspended forms of copper for different hydrologic seasons was at Kiona. The snowmelt-suspended load (average monthly load for April and May 1989) of 178 pounds of copper per day was nearly 6 times larger than the respective dissolved copper load.

Copper concentrations in aquatic biota generally were less than 35  $\mu\text{g/g}$  except for rainbow trout liver, carp liver, and some species of stonefly (table 40, at back of report). Differences in copper concentrations among different taxa might reflect species-specific differences in metal accumulation, a common phenomenon. For example, trout tend to accumulate more copper when compared to other fish (Rasmussen, 1992). Among sites, copper concentrations varied less than 2 times in most taxa. For example, in the caddisfly (*Hydropsyche* spp.), concentrations ranged from 10 to 15  $\mu\text{g/g}$  at 20 of the 24 sites where caddisflies were collected in 1990 (fig. 44). The results indicate that biologically available copper is distributed uniformly over much of the Yakima River Basin.

Two sites, however, consistently exhibited high copper concentrations in biological samples. Wide Hollow Creek at the old Sewage Treatment Plant at



**Figure 43.** Copper loads in filtered-water samples and in suspended-sediment samples for selected time periods at fixed sites, Yakima River Basin, Washington, 1989 (the term "filtered water" represents the portion of a water sample passing through a nominal 0.45 micrometer pore-size filter; non-irrigation season occurs October through March; snowmelt season occurs April through May; irrigation season occurs June through September; "Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).



Union Gap had maximum copper concentrations in caddisflies (*Hydropsyche* spp.) of 48 µg/g in 1989 and 21 µg/g in 1990; 480 µg/g in 1990 in rainbow trout liver (trout were not collected at this site in 1989); and curlyleaf pondweed, an aquatic plant. The copper concentration in trout in Wide Hollow Creek was more than an order of magnitude greater than in rainbow trout of the same size in Taneum Creek at Taneum Meadow near Thorp (a site unaffected by agricultural or urban activity). Wide Hollow Creek drains agricultural land and also receives urban runoff from the city of Yakima. Relatively high concentrations of copper (290 µg/g in 1990 and 130 µg/g in 1989) also were measured in trout livers collected from the Yakima River at Umtanum, a site which receives urban runoff from the city of Ellensburg and has episodically high copper concentrations in filtered-water samples, as previously mentioned. The greater variation in copper concentration in rainbow trout livers may indicate differences in the capacity of individual trout to accumulate copper, as well as site-related differences in copper concentrations among samples (Vidal, 1978).


Concentrations of copper in curlyleaf pondweed ranged from 9.2 to 22 µg/g (table 15). Concentrations of copper in pondweed in Cherry Creek at Thrall (22 µg/g) and in Wide Hollow Creek at the old Sewage Treatment Plant at Union Gap (21 µg/g) were among the highest concentrations in the Yakima River Basin (Fuhrer, Fluter, and others, 1994). The water weed (*Elodea* sp.) in Spring Creek at mouth at Whitstran had the highest copper concentration among all aquatic plants. The Spring Creek site carries agricultural return flow to the main stem.

Enrichment of copper in Wide Hollow Creek is measurable in all media sampled. In addition to enrichment in aquatic biota, copper concentrations also were high in filtered water and streambed sediment (suspended sediment was not measured) [table 27]. The concentrations of copper in filtered water from Wide Hollow Creek near West Valley High School (site 27) and in Wide Hollow Creek at the old Sewage Treatment Plant (site 29) were, respectively, 5.0 µg/L and 1.4 µg/L and exceeded the 75th percentile value (1.3 µg/L) for copper determinations at fixed sites 1987–90. Concentrations of copper in streambed sediment at these sites were, respectively, 93 µg/g and 42 µg/g, and equaled or exceeded the 75th percentile for streambed sediment collected from biological sampling sites in the Yakima River Basin. Additionally, concentrations of copper in all media sampled were


larger at the West Valley High School site than at the mouth of Wide Hollow Creek (Old Sewage Treatment Plant site). Water and streambed sediment near the mouth of Wide Hollow Creek may have been affected by the mixing of copper-poor sediment from the Yakima River with copper-rich sediment from Wide Hollow Subbasin.

The patterns of copper enrichment in water and fish in the Yakima River at Umtanum (site 19) are similar to those measured in Wide Hollow Creek; however, the relative concentrations are not as high. The copper concentration in rainbow trout liver, for example, is 290 µg/g in the Yakima River at Umtanum and 480 µg/g in Wide Hollow at old Sewage Treatment Plant at Union Gap. Both concentrations, however, were larger than the median concentration (100 µg/g) for rainbow trout liver. The Umtanum site is located 6.6 miles downstream from Wilson Creek, which is affected by anthropogenic activities in the Kittitas, Wilson-Naneum, and Swauk Subbasins. The city of Ellensburg is located in the Wilson-Naneum Subbasin. Streambed-sediment concentrations of copper, although high (65 µg/g) in the city of Ellensburg's urban, stormwater drain site, were attenuated near the mouth of Wilson Creek. Sediment affected by urban runoff was mixed, near the mouth of Wilson Creek, with copper-poor sediment in agricultural return flow; furthermore, after entering the main stem, sediment concentrations of copper were again attenuated downstream. Comparisons between copper found in rainbow trout collected from the Yakima River at Umtanum and from the Umtanum Creek near mouth at Umtanum (site 20) indicate the presence of an upstream source of copper. Samples collected in the fall of 1989 and 1990 indicated that copper concentrations in liver tissue of rainbow trout from the Yakima River at Umtanum were approximately 2.5 times higher than concentrations from Umtanum Creek near the mouth at Umtanum. Umtanum Creek is located immediately downstream from the Yakima River at Umtanum; Umtanum Creek's drainage is isolated from copper anomalies (excluding atmospheric deposition) in the Kittitas Valley. High concentrations of copper in rainbow trout may result from dissolved forms of copper in the Yakima River at Umtanum. Dissolved forms of copper are adsorbed readily through the gills of fish—rainbow trout in particular. Accordingly, guidelines for protecting fish in low hardness waters are in units of tenths of a microgram per liter (Moore, 1991, p. 119). The highest copper


### MOUNTAIN WHITEFISH LIVERS




### LARGESCALE SUCKER LIVERS



### RAINBOW TROUT LIVERS




### CADDISFLIES



**Figure 44.** Distribution of copper concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, caddisflies, and streambed sediment, Yakima River Basin, Washington, 1987–90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are graphically represented for mountain whitefish, largescale sucker, rainbow trout, and caddisflies; sample species: mountain whitefish [*Prosopium williamsoni*], largescale sucker [*Catostomus macrocheilus*], rainbow trout [*Oncorhynchus mykiss*], and caddisflies [Trichoptera: Hydropsychidae *Hydropsyche* spp.]).



# STREAMBED SEDIMENT



**Figure 44.** Distribution of copper concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, caddisflies, and streambed sediment, Yakima River Basin, Washington, 1987–90—Continued

concentrations (20 and 7.4 µg/L) in filtered-water samples were measured at Umtanum in the fall, a period characterized by low streamflows after the irrigation season. During this period, the Yakima River at Umtanum is susceptible to episodic increases in concentrations of copper in filtered water. Among sites, other factors, such as differences in fish length may affect the distribution of copper in rainbow trout, however.

Overall, copper concentrations in most biological samples from the Yakima River Basin were comparatively low and appear indicative of background concentrations. The enrichment of copper in Wide Hollow Creek and the Yakima River at Umtanum is modest when compared to other river basins. For example, the maximum concentrations in benthic insects in the Yakima River Basin in 1990 generally were lower than the mean concentrations reported for aquatic insects from uncontaminated streams and rivers (table 40, at back of report) [Cain and others, 1992; Gower and Darlington, 1990; Lynch and others, 1988; Miller and others, 1992]. The variation in copper concentration shown in table 40 indicates differences in concentrations among sites and differences in copper

accumulation among different species. More direct comparisons can be made with caddisflies (*Hydropsyche* spp.). Cain and others (1992) reported that the mean and standard deviation for copper concentrations in caddisflies (*Hydropsyche* spp.) from reference streams in Montana were  $18 \pm 0.4$  µg/g. The Montana reference mean is comparable to concentrations of copper in caddisflies (*Hydropsyche* spp.) at all sites in the Yakima River Basin except for the 1989 insect sample (48 µg/g) from Wide Hollow Creek. Concentrations of copper in Wide Hollow Creek, although high relative to caddisflies (*Hydropsyche* spp.) in the Yakima River Basin were similar to those found in aquatic insects in the least contaminated reaches of the Clark Fork River in Montana (Axtmann and others, 1991; Cain and others, 1992).

Concentrations of copper in fish liver in the Yakima River Basin also were not particularly high. All fish-liver samples in the basin had copper concentration that were lower than the 85th percentile concentrations in salmonid and non-salmonid fish liver sampled in California's Toxic Substances Monitoring Program (Rasmussen, 1992). In California's program, copper concentrations in salmonid and non-salmonids, respectively, were 680 and 52 µg/g (calculated from original wet-weight data; Rasmussen, 1992). The concentrations of copper in rainbow trout in Wide Hollow Creek (480 µg/g) and the Yakima River at Umtanum (290 µg/g) were intermediate compared to the 55 to 941 µg/g in the liver of rainbow trout from California rivers (calculated from original wet-weight data or percent-moisture data when available) [McCleneghan and others, 1980]. The highest concentration reported by McCleneghan in rainbow trout liver (941 µg/g) is from a reach of the Sacramento River reportedly affected by acid mine wastes. Moore and others (1991) reported a much lower concentration (15 µg/g) in cutthroat trout livers from an uncontaminated stream, a concentration comparable to that found in the only sample of cutthroat trout collected in the Yakima River Basin (7.2 µg/g in the South Fork of Ahtanum Creek above Conrad Ranch near Tampico).

No evidence of copper contamination was found in Asiatic clams in the Yakima River (table 39, at back of report). In other studies of uncontaminated areas, copper concentrations in Asiatic clams ranged from 20 to 40 µg/g (Luoma and others, 1990; Leland and Scudler, 1990). For comparison, copper concentrations in

**Table 27. Comparison of low and high copper concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91**  
 [For filtered water and suspended sediment, the low and high concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For streambed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an "H" in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an "L" in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an "\*\*" in the table. The term "filtered water" is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale sucker live and caddisflies; sample species: largescale sucker (*Catostomus macrocheilus*), rainbow trout (*Oncorhynchus mykiss*), caddisfly (*Hydropsyche* spp.), stonefly (*Hesperoperla* sp.), Asiatic clam (Veneroida: Corbiculidae *Corbicula fluminea*), and curlyleaf pondweed (*Potamogeton crispus*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Yakima River above Athanum Creek at Union Gap, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987–90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data; NF, North Fork; SF, South Fork; Cr, Creek; RM, river mile]

Site reference number	Site name	Filtered water	Sediment		Aquatic biota						
			Streambed	Suspended	Fish livers			Insects			
					Largescale sucker	Mountain whitefish	Rainbow trout	Caddisfly	Stonefly	Asiatic clam	Curlyleaf pondweed
1	Wapatus River at mouth near Roslyn	--	L	--	--	--	*	--	--	--	--
3	Jungle Creek near mouth near Cle Elum	--	*	--	--	--	L	--	--	--	--
4	NF Teanaway River below bridge at Dickey Creek Campground	--	--	--	--	--	--	--	--	H	--
5	Teanaway River below Forks near Cle Elum	--	*	--	--	--	--	*	--	--	--
6	Yakima River at Cle Elum	L	H	H	--	*	--	*	H	--	--
7	Naneum Creek below High Creek near Ellensburg	--	L	--	--	--	--	H	*	--	--
8	Taneum Creek at Taneum Meadow near Thorp	--	H	--	--	--	L	--	*	--	--
10	Little Naches River at mouth near Cliffdell	--	H	--	--	--	--	L	--	--	--
12	SF Manastash Cr near Ellensburg	--	H	--	--	--	*	--	*	--	--
13	American River at Hells Crossing near Nile	--	H	--	--	--	--	--	--	--	--

**Table 27.** Comparison of low and high copper concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Filtered water	Sediment		Aquatic biota							
			Streambed	Suspended	Fish livers			Insects		Asiatic clam	Curlyleaf pondweed	
					Largescale sucker	Mountain whitefish	Rainbow trout	Caddisfly	Stonefly			
14	Cherry Creek above Wipple Wasteway at Thrall	--	--	--	--	*	--	--	*	--	--	--
16	Cherry Creek at Thrall	--	L	--	--	--	--	--	*	--	--	H
19	Yakima River at Umtanum	H	*	L	--	L	H	--	L	*	--	L
20	Umtanum Creek near mouth at Umtanum	--	*	--	--	--	*	--	H	--	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	--	*	--	--	--	--	--	*	*	--	--
22	Rattlesnake Creek above NF Rattlesnake Creek near Nile	--	H	--	--	--	*	--	*	*	--	--
26	Naches River near North Yakima	L	H	*	--	*	--	--	L	--	--	-
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	--	H	--	--	--	--	--	H	--	--	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	--	H	--	--	--	--	H	H	--	--	H
30	Moxee Drain at Thorp Road near Union Gap	--	L	--	--	--	--	--	--	--	--	--
31	Ahtanum Creek at Union Gap	--	L	--	--	--	--	--	L	--	--	--
32	Yakima River above Ahtanum Creek at Union Gap	*	--	H	--	--	--	--	--	--	--	--
33	Yakima River at Parker	--	*	--	--	L	L	--	*	--	--	L
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	--	*	--	--	--	--	--	*	L	--	--

**Table 27.** Comparison of low and high copper concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Filtered water	Sediment		Aquatic biota							
					Fish livers			Insects			Asiatic clam	Curlyleaf pondweed
					Largescale sucker	Mountain whitefish	Rainbow trout	Caddisfly	Stonefly			
40	Granger Drain at mouth near Granger	--	L	--	--	--	--	H	--	--	--	
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	--	*	--	*	H	--	*	--	*	--	
43	Toppenish Creek at Indian Church Road near Granger	--	*	--	--	--	--	--	-	--	--	
47	Satus Creek at gage at Satus	--	*	--	*	--	--	H	--	--	*	
48	Yakima River at RM 72 above Satus Creek near Sunnyside	--	*	--	--	--	--	--	--	*	*	
50	Yakima River at Kiona	*	*	L	H	H	--	H	--	L	H	
52	Sulphur Creek Wasteway near Sunnyside	H	*	*	--	--	--	*	--	--	--	
53	Satus Creek below Dry Creek near Toppenish	--	*	--	--	--	--	*	--	--	--	
54	Spring Creek at mouth at Whitstran	--	L	--	--	--	--	*	--	*	--	
56	Yakima River at Euclid Bridge at RM 55 near Grandview	*	*	L	*	--	--	*	--	H	*	
57	Satus Creek above Wilson- Charley Canyon near Toppenish	--	L	--	--	--	*	L	L	--	--	

clams from contaminated areas were as high as 200 µg/g (Luoma and others, 1990).

## Mercury

Concentrations of mercury in streambed sediment of the Yakima River Basin ranged from <0.2 to 0.6 µg/g (table 12). Approximately 10 percent of the sites had mercury concentrations that exceeded the baseline value (0.3 µg/g) for Yakima River streambed sediment reported by Fuhrer, McKenzie, and others (1994). Similarly, about 10 percent of the sites sampled had mercury concentrations that exceeded the 0.008 to 0.25 µg/g range of concentration which encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). The median concentration of mercury at the biological sampling sites (0.1 µg/g) also exceeded those determined from analysis of fine-fraction streambed sediment in other river basins of the United States (table 34, at back of report). Concentrations of mercury as high as 0.56 µg/g, 0.40 µg/g, and 0.30 µg/g were found in streambed sediment of Jungle Creek near mouth near Cle Elum (site 3), Taneum Creek at Taneum Meadow near Thorp (site 8), and the Naches River near North Yakima (site 26), respectively.

Mercury enrichment at Jungle Creek near mouth near Cle Elum, a tributary to the North Fork of the Teanaway River, probably was from geologic sources. Streambed sediment in lower-order streams of the north, middle, and west forks of the Teanaway River are enriched similarly (Ryder and others, 1992) and indicate base-metal enrichment in sediment derived from the nonmarine sedimentary rocks geologic unit, which includes the Swauk and Roslyn Formations, and the Miocene and older volcanic rocks geologic unit, which includes the Teanaway Basalt (Fuhrer, McKenzie, and others, 1994).

Similar to Jungle Creek, mercury in sediment at Taneum Creek near Taneum Meadow probably originates from geologic sources. Upstream, concentrations of mercury in the fine-grained-sized sediment in lower-order streams of the Taneum Creek drainage were as high as 0.24 µg/g (Ryder and others, 1992). Mercury in the Naches River near North Yakima probably resulted from geologic sources present in sediment formed from the Miocene and older volcanic rocks, Tertiary granitic and

intermediate intrusive rocks, and marine sedimentary rocks geologic units (Fuhrer, McKenzie, and others, 1994). The concentration of mercury in the Naches River near North Yakima probably comes from lower-order streams of the Naches River that are affected by geologic sources of mercury (Fuhrer, McKenzie, and others, 1994).

Based on three independent samplings, streambed sediment in Wide Hollow Creek contained an average of 0.26 µg/g mercury. Mercury in Wide Hollow Creek at the old Sewage Treatment Plant at Union Gap (site 29) is probably from an anthropogenic source. Wide Hollow Creek drains urbanized and industrialized lowlands in the vicinity of the city of Yakima. Wide Hollow Creek lacks geologic sources of mercury that are known to exist in streambed sediment derived from geologic sources, because its drainage is confined to the mercury-poor Quaternary deposits and loess geologic unit, (Fuhrer, McKenzie, and others, 1994).

Mercury concentrations in most fish samples were less than 0.5 µg/g (table 15). Higher concentrations of mercury were observed in mountain whitefish. For a given species, higher concentrations of mercury were measured in the Yakima River and in the Naches River near North Yakima (site 26) than in tributaries (table 28). For example, concentrations of mercury in mountain whitefish ranged from 0.7 to 1.3 µg/g at sites in the main stem and in the Naches River, compared to only 0.4 µg/g at Cherry Creek above Wipple Wasteway at Thrall (fig. 45).

Similar to mountain whitefish, concentrations of mercury in the livers of largescale suckers were higher at sites in the Yakima River (0.3 to 0.47 µg/g) and at the Naches River near North Yakima (0.14 µg/g, site 26) than, for example, from the Satus Creek at gage at Satus (0.05 µg/g, site 47). However, between species, concentrations of mercury were smaller in livers of largescale suckers than in livers of mountain whitefish (table 15 and fig. 45). Concentrations of mercury in largescale sucker also were lower statistically ( $p \leq 0.0001$ ) than in mountain whitefish.

Mercury concentrations, during the 1990 sampling of rainbow trout liver from the Yakima River at Umtanum (0.27 µg/g), were similar to those in Wide Hollow Creek at the old Sewage Treatment Plant at Union Gap, in Taneum Creek at Taneum Meadow near Union Gap, and in Umtanum Creek near mouth at Umtanum (0.33 to 0.38 µg/g) [fig. 45]. Lower concentrations of mercury were found in trout from

**Table 28. Comparison of low and high mercury concentrations in streambed sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91**

[For filtered water and suspended sediment, the low and high concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For stream bed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an "H" in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an "L" in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an "\*" in the table. Only 1990 data are summarized for largescale sucker live; sample species: largescale sucker (*Catostomus macrocheilus*), mountain whitefish (*Prosopium williamsoni*), rainbow trout (*Oncorhynchus mykiss*), sculpin (*Cottus* spp.), Asiatic clam (*Veneroida: Corbiculidae Corbicula fluminea*), and curlyleaf pondweed (*Potamogeton crispus*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987–90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]


Site reference number	Site name	Streambed sediment	Aquatic biota					
			Fish				Asiatic clam	Curlyleaf pondweed
			Largescale sucker	Mountain whitefish	Rainbow trout	Sculpin		
1	Wapatus River at mouth near Roslyn	*	--	--	*	--	--	--
3	Jungle Creek near mouth near Cle Elum	H	--	--	H	--	--	--
5	Teanaway River below Forks near Cle Elum	H	--	--	L	--	--	--
6	Yakima River at Cle Elum	*	--	*	--	L	--	--
7	Naneum Creek below High Creek near Ellensburg	*	--	--	--	H	--	--
8	Taneum Creek at Taneum Meadow near Thorp	H	--	--	H	H	--	--
10	Little Naches River at mouth near Cliffdeil	*	--	--	--	--	--	--
12	South Fork Manastash Creek near Ellensburg	*	--	--	L	*	--	--
13	American River at Hells Crossing near Nile	*	--	--	--	*	--	--
16	Cherry Creek at Thrall	*	--	L	--	--	--	*
19	Yakima River at Umtanum	*	--	L	*	--	--	L
20	Umtanum Creek near mouth at Umtanum	L	--	--	*	H	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	H	--	--	L	--	--	--
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	*	--	--	--	*	--	--
26	Naches River near North Yakima	H	*	*	--	--	--	--

**Table 28.** Comparison of low and high mercury concentrations in streambed sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987-91—Continued


Site reference number	Site name	Streambed sediment	Aquatic biota					
			Fish					Curlyleaf pondweed
			Largescale sucker	Mountain whitefish	Rainbow trout	Sculpin	Asiatic clam	
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	*	--	--	--	--	--	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	H	--	--	H	--	--	H
30	Moxee Drain at Thorp Road near Union Gap	L	--	--	--	--	--	--
31	Ahtanum Creek at Union Gap	*	--	--	--	*	--	--
33	Yakima River at Parker	H	H	H	--	--	--	*
34	South Fork Ahtanum Creek above Conrad Ranch near Tappico	*	--	--	--	*	--	--
40	Granger Drain at mouth near Granger	L	--	--	--	--	--	--
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger	*	*	H	--	--	H	--
43	Toppenish Creek at Indian Church Road near Granger	*	--	--	--	--	--	--
47	Satus Creek at gage at Satus	*	L	--	--	L	--	L
48	Yakima River at river mile 72 above Satus Creek near Sunnyside	*	--	--	--	--	*	H
50	Yakima River at Kiona	*	*	*	--	--	L	*
52	Sulphur Creek Wasteway near Sunnyside	L	--	--	--	--	--	--
53	Satus Creek below Dry Creek near Toppenish	L	--	--	--	L	--	--
54	Spring Creek at mouth at Whitstran	L	--	--	--	--	H	--
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	*	*	--	--	--	H	L
57	Satus Creek above Wilson-Charley Canyon near Toppenish	*	--	--	*	L	--	--




### MOUNTAIN WHITEFISH LIVERS




### LARGESCALE SUCKER LIVERS



### RAINBOW TROUT LIVERS




### WHOLE SCULPINS



**Figure 45.** Distribution of mercury concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, whole sculpins, and streambed sediment, Yakima River Basin, Washington, 1987–90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are graphically represented for mountain whitefish, largescale sucker, rainbow trout, and sculpins; sample species: mountain whitefish [*Prosopium williamsoni*], largescale sucker [*Catostomus macrocheilus*], rainbow trout [*Oncorhynchus mykiss*], and sculpin [*Cottus* spp.]).

# STREAMBED SEDIMENT




**Figure 45.** Distribution of mercury concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, whole sculpins, and streambed sediment, Yakima River Basin, Washington, 1987–90—Continued

the South Fork of Manastash Creek (0.16  $\mu\text{g/g}$ ). Mercury concentrations in whole sculpin from lower-order tributaries ranged from 0.09 to 0.31  $\mu\text{g/g}$ . The highest concentrations of mercury existed in Naneum Creek below High Creek near Ellensburg (site 7), in Umtanum Creek near mouth at Umtanum (site 20), and in Taneum Creek at Tanum Meadow near Thorp (site 8); the latter sites also coincided with higher mercury concentrations in rainbow trout.

The distribution of mercury concentrations in samples collected in 1989 generally was consistent with that in 1990. The highest concentrations of mercury are in the liver of mountain whitefish in the Yakima River at Cle Elum (0.91  $\mu\text{g/g}$ ). Lower concentrations of mercury existed downstream in the Yakima River at Umtanum (0.6  $\mu\text{g/g}$ ) and in the Yakima River at Kiona (0.58  $\mu\text{g/g}$ ). Mercury concentrations in largescale sucker and rainbow trout were comparatively low (<0.4  $\mu\text{g/g}$ ). The highest concentrations of mercury in rainbow trout were found in the Kittitas Valley in Jungle Creek near mouth near Cle Elum (0.32  $\mu\text{g/g}$ ) and in the Wapatus River at mouth near Roslyn (0.27  $\mu\text{g/g}$ )—these sites were not resampled in

1990. The mercury concentrations in rainbow trout liver at the Jungle Creek site, however, corresponded to the concentration maximum (0.56  $\mu\text{g/g}$ ) for mercury in streambed sediment at the biological sampling sites. Caddisflies analyzed for mercury in 1989 had concentrations of <0.2  $\mu\text{g/g}$ . Additionally, concentrations of mercury among all sites only varied by a factor of 3 (0.05 to 0.16  $\mu\text{g/g}$ ) with the highest concentrations at Rattlesnake Creek (0.16  $\mu\text{g/g}$ ) and Toppenish Creek (0.10  $\mu\text{g/g}$ ). A slight downstream-concentration gradient was apparent in samples from the main stem. Concentrations of mercury were highest at the Cle Elum and Umtanum sites (0.090 and 0.080  $\mu\text{g/g}$ , respectively), and lower in the Yakima River at Parker, the Yakima River at Euclid Bridge at RM 55 near Grandview, and at Kiona (0.066 to 0.070  $\mu\text{g/g}$ ).

The lower concentrations of mercury, in mountain whitefish from Cherry Creek (0.4  $\mu\text{g/g}$ ) and in largescale suckers from Satus Creek (0.05  $\mu\text{g/g}$ ), may result when fish are physically smaller (fig. 46). Inter-




**Figure 46.** Relation between fish length and mercury concentrations of mountain whitefish livers and largescale sucker livers, Yakima River Basin, Washington, 1990 (sample species: mountain whitefish [*Prosopium williamsoni*] and largescale sucker [*Catostomus macrocheilus*]).

site variations in size, however, do not account for all the concentration variability. Variations in mercury concentrations among larger fish from other sites were relatively high; additionally, no significant correlation among all sites existed between fish length and mercury concentration. Thus, site differences were not simply a function of fish size (age). For mountain

whitefish, the variations in mercury content among sites were perhaps best explained by considering the quantity of mercury and total-organic carbon in streambed sediment. Mercury in streambed sediment has an affinity for sorption to organic material (Forstner and Wittmann, 1979, p. 222). The quantity of mercury relative to organic material has been shown to affect the bioavailability of mercury to Asiatic clams (Langston, 1982). Mercury concentrations in streambed sediment in the Yakima River Basin were normalized to the concentrations of total-organic carbon (ratio of mercury in micrograms per gram to total-organic carbon in micrograms per gram) and compared to mercury concentrations in fish liver. The normalized value for the mouth of the Naches River near North Yakima is 0.75 and was exceeded only at the Parker site (0.84) where a mercury concentration of 1.3  $\mu\text{g/g}$  (the maximum concentration for biological sites) was measured in the liver of mountain whitefish. The ratio of mercury in mountain whitefish liver to total-organic carbon is significantly correlated ( $p < 0.05$ ) to mercury accumulation in the liver of mountain whitefish in the Yakima River Basin (fig. 47). The same relation, however, does not exist for mercury in rainbow trout liver (fig. 47), possible because of differences in lipid content between species and from differences in age between species. Mountain whitefish from main-stem sites generally were larger physically (and presumably older) than the smaller (presumably younger) rainbow trout from tributary sites (Fuhrer, Fluter, and others, 1994).

Mercury concentrations in whole sculpin from tributaries to the Yakima River were below 0.31  $\mu\text{g/g}$  which also is below the mean mercury concentration for whole fish collected for the National Contaminant Biomonitoring Program (0.4  $\mu\text{g/g}$ , calculated from original wet-weight data) in 1984 (table 38, at back of report; Schmitt and Brumbaugh, 1990). Mercury concentrations determined as part of the National Contaminant Biomonitoring Program are generally from near the mouths of large rivers and are more likely to be affected by human activities than are headwater tributaries in many of the streams in the Yakima River Basin. In general, concentrations of mercury greater than 0.5  $\mu\text{g/g}$  appear to be indicative of contamination (Saiki and May, 1988).

Mercury concentrations in fish liver have been reported in several studies. Concentrations in uncontaminated tissue appear to be less than 0.5  $\mu\text{g/g}$ . For example, the livers of four species of fish, including



**Figure 47.** Mercury in mountain whitefish livers and rainbow trout livers compared to mercury in streambed sediment normalized to total organic carbon, Yakima River Basin, Washington, 1989–90 (sample species: mountain whitefish [*Prosopium williamsoni*] and rainbow trout [*Oncorhynchus mykiss*]).

whitefish (*Coregonus lavaretus*) from an uncontaminated lake in Finland, had mercury concentrations that ranged from 0.03 to 0.48  $\mu\text{g/g}$  (Lodenius and others, 1982). Mean mercury concentrations in five species from two rivers ranged from 0.16 to 0.64  $\mu\text{g/g}$  (calculated from original wet-weight data in Barak and Mason, 1990). Mean mercury concentrations in bream (*Abramis brama*) and in pike perch (*Stizostedion lucioperca*) from Lake Balaton in western Hungary (determined to be mildly contaminated) were 0.3  $\mu\text{g/g}$  and 0.35  $\mu\text{g/g}$ , respectively (Salanki and others, 1982). In contrast, Koli and others (1977) reported that mercury

concentrations in the livers of contaminated pike and mudfish were 0.94 µg/g and 0.83 µg/g, respectively, whereas concentrations of mercury in muscle tissue were higher than 0.5 µg/g. Compared with these studies, mercury concentrations (0.7 to 1.3 µg/g) in livers of mountain whitefish in Yakima River Basin were moderately enriched. Concentrations of mercury, in mountain whitefish, collected from the mouth of the Naches River, also were enriched (1.0 µg/g); however, mountain whitefish may have moved freely between the Yakima and Naches Rivers. Concentrations of mercury in other species of fish collected with mountain whitefish (for example, largescale sucker and carp) were relatively low (typically less than 0.5 µg/g). However, distribution patterns of mercury in largescale sucker were consistent with those of mountain whitefish; both species indicated enrichment in the Yakima River. Fish collected in the Yakima River at Bridge Avenue at Granger for the National Contaminant Biomonitoring Program, for the period 1978–79, also showed evidence of mercury enrichment (Lowe and others, 1985).

In contrast to fish, Asiatic clam (*Corbicula fluminea*) collected in the Yakima River did not show mercury enrichment. Concentrations of mercury varied between 0.10 and 0.17 µg/g, among stations in the Yakima River, and were similar to concentrations observed in uncontaminated systems (Johns and others, 1988; Leland and Scudder, 1990) [table 39, at back of report].

Mercury concentrations in aquatic plants in the Yakima River Basin were relatively uniform. In curlyleaf pondweed for example, concentrations of mercury ranged from 0.03 to 0.06 µg/g among sites (Fuhrer, Fluter, and others, 1994). The uptake of mercury by aquatic plants has been described as extremely rapid and efficient (Moore, 1991). In another study, moss (*Fontinalis squamosa*) in a rural river that received acid-mine drainage contained mercury in concentrations ranging from 0.12 to 0.48 µg/g (Moore, 1991, p. 200). The absence of appreciable mercury concentrations in aquatic plants sampled at sites in the Yakima River Basin indicates that concentrations of mercury in dissolved and solid forms are relatively low.

## Selenium

Concentrations of selenium in Yakima River streambed sediment ranged from <0.4 to 1.0 µg/g (table 12). Approximately 25 percent of the sites had


selenium concentrations that exceeded the baseline value (0.7 µg/g) for Yakima River streambed sediment as reported by Fuhrer, McKenzie, and others (1994). None of the sites had selenium concentrations that exceeded the 0.04 to 1.4 µg/g range of concentration which encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). Concentrations of selenium as high as 1.0, 0.9, 0.9, 0.8, and 0.7 µg/g were found in streambed sediment of Rattlesnake Creek above North Fork Rattlesnake Creek near Nile (site 22), in Naneum Creek below High Creek near Ellensburg (site 7), in Taneum Creek at Taneum Meadow near Thorp (site 8), in Waptus River at mouth near Roslyn (site 1), and in American River at Hells Crossing near Nile (site 13), respectively.

Selenium in Rattlesnake Creek above North Fork Rattlesnake Creek near Nile (site 22) is probably associated with streambed sediment formed from the marine sedimentary rocks geologic unit. Streambed sediment was sampled by Fuhrer, McKenzie, and others (1994) from three lower order tributaries of Rattlesnake Creek, located upstream of site 22. Although selenium was not measured, large concentrations of arsenic and copper measured at these sites were indicative of selenium enrichment—selenium is known to have strong geochemical associations with arsenic and copper (Levinson, 1980; Rankama and Sahama, 1950). The presence of selenium at the remaining sites probably is due to geological sources; drainages are forested and selenium inputs from human activities are expected to be negligible.


In the Yakima River Basin, several higher order tributaries including Rattlesnake Creek, the American River, Ahtanum Creek, and Wide Hollow Creek, had high selenium concentrations in fish (fig. 48). Selenium concentrations also were high at some sites in the lower Yakima River Basin, particularly in the main stem.

The selenium concentrations in the whole sculpin, collected from 12 sites in the Yakima River Basin in 1990, ranged more than an order of magnitude (table 15). The lowest concentrations of selenium were found at two sites in the upper portion of Satus Creek (above Wilson-Charlie Canyon near Toppenish [0.5 µg/g] and below Dry Creek near Toppenish [0.2 µg/g]). The highest concentrations of selenium were found in Rattlesnake Creek above North Fork Rattlesnake Creek near Nile (5.5 µg/g, site


### MOUNTAIN WHITEFISH LIVERS




### LARGESCALE SUCKER LIVERS



### RAINBOW TROUT LIVERS




### WHOLE SCULPINS



**Figure 48.** Distribution of selenium concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, whole sculpins, and streambed sediment, Yakima River Basin, Washington, 1987–90 (element concentrations are reported in units of micrograms per gram [ $\mu\text{g/g}$ ], dry weight; symbol sizes are proportional to element concentrations; the largest and smallest symbols, respectively, represent the high and low concentration end members; only 1990 data are summarized for mountain whitefish, largescale sucker, rainbow trout, and sculpins; sample species: mountain whitefish [*Prosopium williamsoni*], largescale sucker [*Catostomus macrocheilus*], rainbow trout [*Oncorhynchus mykiss*], and sculpin [*Cottus* spp.]).

## STREAMBED SEDIMENT




**Figure 48.** Distribution of selenium concentrations in mountain whitefish livers, largescale sucker livers, rainbow trout livers, whole sculpins, and streambed sediment, Yakima River Basin, Washington, 1987–90—Continued

22) and in Taneum Creek near Taneum Meadow (5.4  $\mu\text{g/g}$ , site 8). The concentrations of selenium in whole sculpin in Rattlesnake and Taneum Creeks were approximately 3 times higher than the median for the basin. Selenium in sculpin from the Rattlesnake and Taneum Creek sites also coincide with high concentrations of selenium in streambed sediment. Concentrations of selenium in streambed sediment from these two sites are approximately 2.5 times higher than the median for the basin. Concentrations of selenium in sculpin in the American River were intermediate (3.2  $\mu\text{g/g}$ ); however, concentrations of selenium in sculpin at most of the other sites ranged from 1 to 3  $\mu\text{g/g}$ .

Other whole sculpin data indicated that selenium concentrations are higher at sites located near the mouths of creeks that carry irrigation return flow than at sites located above agricultural activity. For example, 2.6  $\mu\text{g/g}$  of selenium was measured in whole sculpin from Ahtanum Creek at Union Gap site. This site is located near the mouth of Ahtanum Creek and receives irrigation return flow from the Ahtanum Subbasin. Upstream of agricultural activity on Ahtanum Creek, concentrations of selenium in sculpin

were only 1.0  $\mu\text{g/g}$  (fig. 49). Similar concentration gradients also existed in the Satus Creek drainage; Satus Creek at gage is located near the mouth and receives irrigation return flow (fig. 49).




**Figure 49.** Selenium concentrations in whole sculpins at sites in Satus Creek and Ahtanum Creek, Yakima River Basin, Washington, 1990 (samples species: sculpin [*Cottus* spp.]; "Wilson-Charley" represents Satus Creek above Wilson-Charley Canyon near Toppenish; "Dry Creek" represents Satus Creek below Dry Creek near Toppenish; "at gage" represents Satus Creek at gage at Satus; "South Fork" represents South Fork Ahtanum Creek near Tampico; "Union Gap" represents Ahtanum Creek at Union Gap).

Selenium concentrations in the livers of mountain whitefish ranged from 4.2 to 15  $\mu\text{g/g}$  (table 15 and fig. 48). In 1990, concentrations of selenium in mountain whitefish liver from Cherry Creek at Thrall and other Kittitas and mid-Yakima Valley sites ranged from 4.5 to 8.2  $\mu\text{g/g}$ ; higher concentrations, however, were found in the lower Yakima Valley. A similar pattern of selenium concentrations was observed in livers of largescale suckers, although, concentrations of selenium were lower than in livers of mountain whitefish (table 15). Concentrations of selenium in largescale suckers from the Naches River near North Yakima and Satus Creek at Gage at Satus were approximately 2  $\mu\text{g/g}$ . Concentrations of selenium in the Yakima River ranged from 2.9 to 4.8  $\mu\text{g/g}$ . The highest selenium concentration observed in liver tissue, 31  $\mu\text{g/g}$ , was found in one sample of rainbow-trout liver collected from Wide Hollow Creek at the old Sewage Treatment Plant at Union Gap. The concentration of selenium in this sample was about 4 to 8

times higher than the concentration in rainbow trout from other sites (fig. 48).

At the American River at Hells Crossing near Nile (site 13), selenium concentrations in sculpin seem to decrease with fish length (fig. 50). At other sites, the relation between fish size and selenium concentration was not evident, because the average length of fish in composite samples was relatively uniform (for example, see Rattlesnake Creek in fig. 50). Size-related effects are of secondary importance when comparing



**Figure 50.** Relation between fish length and selenium concentrations of whole sculpins from the American River at Hells Crossing near Nile and Rattlesnake Creek above North Fork Rattlesnake Creek near Nile, Yakima River Basin, Washington, 1990 (sample species: sculpin [*Cottus* spp.]; "American River" represents American River at Hells Crossing near Nile and "Rattlesnake Creek" represents Rattlesnake Creek above North Fork Rattlesnake Creek near Nile).

selenium concentrations in fish from Rattlesnake Creek to other sites. Selenium concentrations were high in Rattlesnake Creek regardless of whether site comparisons were made with mean-site concentrations or concentrations in fish of similar size. In addition among all sites, significant correlation did not exist between fish length and selenium concentration. In other studies, Saiki (1987) reported no correlation between selenium concentrations in whole fish and fish size. Similarly, selenium in Northern Pike muscle also was unrelated to size (Speyer, 1980). In fish collected from Lake Erie, however, a significant positive correlation was observed between size and selenium concentration in yellow perch at one station, but not in

this species from other sites or in other species of fish (Adams and Johnson, 1977).

Comparison of selenium concentrations in whole sculpin from the Yakima River Basin to existing data for whole fish indicates that the American River, Rattlesnake Creek, and Taneum Creek sites in the Yakima River Basin are moderately enriched in selenium. At these sites, selenium concentrations in whole sculpin exceeded 3 µg/g; for comparison, the mean and 85th percentile-selenium concentrations in whole fish collected for the National Contaminant Biomonitoring Program were 1.7 and 2.9 µg/g, respectively, as calculated from original wet-weight data (Schmitt and Brumbaugh, 1990) [table 38, at back of report]. Selenium concentrations in juvenile striped bass, collected from the San Francisco Bay Estuary which receives low to moderate inputs of selenium, were 1 to 2 µg/g (Saiki and Palawski, 1990). In contrast, selenium concentrations in striped bass, from sites in the San Joaquin River system which receive inputs of selenium in irrigation return flow, ranged from 2.7 to 6.9 µg/g (Saiki and Palawski, 1990). Also in the San Joaquin River in California, mean selenium concentrations in bluegill and common carp were lowest (0.647 to 1.41 and 0.988 to 1.50 µg/g, respectively) in tributaries and main-stem sites that do not receive irrigation return flow, and, conversely, were highest (1.36 to 2.9 and 1.67 to 4.30 µg/g, respectively) in tributaries and main-stem sites affected by irrigation return flow (Saiki and May, 1988). In uncontaminated lakes in the Atchafalaya River Basin in Louisiana, selenium concentrations in whole bodies of seven fish species ranged from about 0.4 to 2.0 µg/g (calculated from original wet-weight data) [Winger and Andreasen, 1985].

Comparison of selenium concentrations in livers of whitefish and largescale sucker from the lower Yakima Valley of the Yakima River Basin to existing data (Wiener and others, 1984; Rassmussen, 1992; and Sorensen and Bauer, 1984) indicates that selenium concentrations in the basin are representative of background concentrations that generally ranged from 2 to 3 µg/g. At sites in the lower Yakima Valley, selenium concentrations in whitefish and largescale sucker livers ranged from 1.9 to 15 µg/g; for comparison, the 85th-percentile selenium concentration in whole fish collected in California's Toxic Substances Monitoring Program was 13.8 µg/g (calculated from original wet-weight data) [Rassmussen, 1992]. Selenium concentrations in common carp from sites in the upper Mis-



Mississippi River ranged from 2.2 to 5.3  $\mu\text{g/g}$  (Wiener and others, 1984). The mean selenium concentration in hepatopancreas (liver plus pancreas) of sunfish from a reference lake was 7.2  $\mu\text{g/g}$ , and in contaminated lakes, selenium concentrations in hepatopancreas of sunfish ranged from 33.6 to 89.2  $\mu\text{g/g}$ . The concentration of selenium measured in rainbow trout liver in 1990 from Wide Hollow Creek at the Old Sewage Treatment Plant at Union Gap (31  $\mu\text{g/g}$ ) clearly exceeded concentrations measured in fish liver from uncontaminated rivers. However, the selenium concentration (0.40  $\mu\text{g/g}$ ) in curlyleaf pondweed from Wide Hollow Creek was not high relative to other pondweed samples in the Yakima River Basin—selenium was not measured in streambed sediment and aquatic insects (table 29). Before concluding that the Wide Hollow drainage (which receives irrigation return flow) is affected by selenium, the high selenium concentration in rainbow trout liver should be confirmed; additionally, selenium should be measured in streambed sediment and filtered water to collectively determine the presence and (or) degree of selenium enrichment. Of the largescale suckers collected in 1989, Sulphur Creek Wasteway (which receives large quantities of irrigation return flow) had the maximum concentration of selenium in liver (4.8  $\mu\text{g/g}$ ) [fig. 48]. Although few filtered-water samples collected in the basin were analyzed for selenium, two of the three samples collected from the Sulphur Creek site had detectable selenium concentrations (1 and 2  $\mu\text{g/L}$ ). Again, as in the Wide Hollow Creek drainage, some evidence of selenium was present in aquatic biota; however, additional data are needed to determine to what degree selenium is present in Sulphur Creek Wasteway. Although not unusually high compared to other studies, the concentration of selenium (11  $\mu\text{g/g}$ ) in the liver of rainbow trout, collected from Rattlesnake Creek in 1989, was the maximum concentration measured in fish liver for 1989. This concentration of selenium also coincided with the relatively high selenium concentration in whole sculpin and streambed sediment (described earlier) in Rattlesnake Creek—selenium was not measured in aquatic plants, insects, and suspended sediment from Rattlesnake Creek. Collectively, streambed sediment and aquatic biota data corroborated the moderate selenium enrichment in Rattlesnake Creek that probably originates from natural geologic sources present in the marine sedimentary rocks geologic unit (Fuhrer, McKenzie, and others, 1994). High concentrations of selenium have been

measured in fish chronically exposed to high concentrations of selenium from natural sources (Kaiser and others, 1979).

Selenium concentrations ranged from 2 to 3  $\mu\text{g/g}$  in Asiatic clams in the Yakima River Basin and were indicative of background concentrations. For example, selenium concentrations ranged from 1 to 3  $\mu\text{g/g}$  in uncontaminated reaches of the San Joaquin and Sacramento Rivers in California (Johns and others, 1988; Leland and Scudder, 1990) [table 39, at back of report]. In reaches of the San Joaquin River that are affected by irrigation return flow, selenium concentrations ranged from 2.5 to 5.13  $\mu\text{g/g}$  (Leland and Scudder, 1990).

## Silver

Median concentrations of silver in suspended sediment ranged from 0.2 to 0.5  $\mu\text{g/g}$  at the fixed sites. The lowest and highest median values of silver were found in the Yakima River at Umtanum and the Yakima River above Ahtanum Creek at Union Gap (table 13 and fig. 51; table 35, at back of report). With the exception of the Union Gap site and Sulphur Creek Wasteway near Sunnyside, interquartile ranges for silver concentrations at most sites were small and indicated little or no temporal variation. The variation in the concentration of suspended silver at these latter two sites may result from their close proximity to point sources.

The Union Gap site is located less than 4 miles downstream from the discharge point for the city of Yakima's Sewage Treatment Plant. In 1989, the city of Yakima's Sewage Treatment Plant had a monthly average discharge of 12.8 Mgal/d (million gallons per day) or about 20  $\text{ft}^3/\text{s}$  (Chris Waurvich, city of Yakima Wastewater Treatment Plant, written commun., 1990). The Sulphur Creek site is located immediately downstream from the discharge point for the city of Sunnyside's Sewage Treatment Plant. Over the period 1987–89, the Sunnyside plant had a monthly average discharge of 1.3 Mgal/d or about 2  $\text{ft}^3/\text{s}$  (Washington State Department of Ecology, 1990). Metallic wastes (dissolved and suspended), passing through municipal waste-treatment plants, may be a source of trace elements (including silver) to streambed sediment (Forstner and Wittmann, 1979).

The concentrations of suspended silver at both the Union Gap site and the Sulphur Creek site were


**Table 29.** Comparison of low and high selenium concentrations in streambed sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91

[For filtered water and suspended sediment, the low and high concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For stream bed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an "H" in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an "L" in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an "\*" in the table. The term "filtered water" is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale sucker and caddisflies; sample species: largescale sucker (*Catostomus macrocheilus*), mountain whitefish (*Prosopium williamsoni*), rainbow trout (*Oncorhynchus mykiss*), sculpin (*Cottus* spp.), Asiatic clam (*Veneroidae*: *Corbiculidae* *Corbicula fluminea*), and curlyleaf pondweed (*Potamogeton crispus*). Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987–90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]

Site reference number	Site name	Streambed sediment	Aquatic biota					
			Fish				Asiatic clam	Curlyleaf pondweed
			Largescale sucker	Mountain whitefish	Rainbow trout	Sculpin		
1	Waptus River at mouth near Roslyn	H	--	--	H	--	--	--
3	Jungle Creek near mouth near Cle Elum	L	--	--	--	--	--	--
5	Teanaway River below Forks near Cle Elum	*	--	--	*	-	--	--
6	Yakima River at Cle Elum	--	--	*	--	*	--	--
7	Naneum Creek below High Creek near Ellensburg	H	--	--	--	L	--	--
8	Taneum Creek at Taneum Meadow near Thorp	H	--	--	*	H	--	--
10	Little Naches River at mouth near Cliffdell	*	--	--	--	--	--	--
12	South Fork Manastash Creek near Ellensburg	L	--	--	L	*	--	--
13	American River at Hells Crossing near Nile	H	--	--	--	H	-	--
14	Cherry Creek above Wipple Wasteway at Thrall	--	--	*	--	--	--	--
16	Cherry Creek at Thrall	--	--	--	--	--	--	L
19	Yakima River at Umtanum	*	--	L	*	--	--	L
20	Umtanum Creek near mouth at Umtanum	L	--	--	*	*	--	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	H	--	--	--	--	--	--

**Table 29.** Comparison of low and high selenium concentrations in streambed sediment and aquatic biota for selected sites, Yakima River Basin, Washington, 1987-91—Continued

Site reference number	Site name	Streambed sediment	Aquatic biota					
			Fish				Asiatic clam	Curlyleaf pondweed
			Largescale sucker	Mountain whitefish	Rainbow trout	Sculpin		
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	H	--	--	*	H	--	--
26	Naches River near North Yakima	--	*	*	--	--	--	--
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	*	--	--	--	--	--	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	--	--	--	H	--	--	H
30	Moxee Drain at Thorp Road near Union Gap	L	--	--	--	-	--	--
31	Ahtanum Creek at Union Gap	L	--	--	--	*	--	--
33	Yakima River at Parker	--	*	L	--	--	--	*
34	South Fork Ahtanum Creek above Conrad Ranch near Tampico	L	--	--	--	*	--	--
40	Granger Drain at mouth near Granger	L	--	--	--	--	--	--
42	Yakima River below Toppenish Creek at river mile 79.6 near Granger	L	H	H	--	--	H	--
43	Toppenish Creek at Indian Church Road near Granger	*	--	--	--	--	--	--
47	Satus Creek at gage at Satus	--	L	--	--	*	--	*
48	Yakima River at river mile 72 above Satus Creek near Sunnyside	*	--	--	--	--	*	L
50	Yakima River at Kiona	--	*	H	--	--	L	H
52	Sulphur Creek Wasteway near Sunnyside	L	--	--	--	--	--	--
53	Satus Creek below Dry Creek near Toppenish	L	--	--	--	L	--	--
54	Spring Creek at mouth at Whitstran	--	--	--	--	--	*	--
56	Yakima River at Euclid Bridge at river mile 55 near Grandview	--	*	--	--	--	L	*
57	Satus Creek above Wilson-Charley Canyon near Toppenish	L	--	--	L	L	--	--





#### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.

- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value
- × 1.5 to 3 times the interquartile range from the 25-percentile value


**Figure 51.** Distribution of silver concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 (“Cle Elum” represents Yakima River at Cle Elum; “Umtanum” represents Yakima River at Umtanum; “Naches” represents Naches River near North Yakima; “Union Gap” represents Yakima River above Ahtanum Creek at Union Gap; “Sulphur Creek” represents Sulphur Creek Wasteway near Sunnyside; “Grandview” represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and “Kiona” represents Yakima River at Kiona).

high during the non-irrigation season and low during the irrigation season (fig. 52). Geologic sources are unlikely to affect the distribution of suspended-silver concentrations at the Union Gap site, because of low suspended-silver concentrations in the Kittitas Valley and in the Naches Subbasin. Similarly, streambed sediment, originating from Quaternary deposits and loess geologic unit in the Sunnyside Subbasin, are an



**Figure 52.** Silver concentrations in suspended sediment at the Yakima River above Ahtanum Creek at Union Gap and at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1989 (“Union Gap” represents Yakima River above Ahtanum Creek at Union Gap and “Sulphur Creek” represents Sulphur Creek Wasteway near Sunnyside).

unlikely source of suspended silver for Sulphur Creek. Instead, the distribution of silver at both sites probably results from the concentration and particle size of the suspended sediment mixing with effluent from point sources. During the irrigation season, for example, the concentration of suspended sediment at the Sulphur Creek site is substantially higher than during the non-irrigation season and, conversely, the quantity of fine-grained-sized, suspended sediment during the irrigation season is smaller than during the non-irrigation season (fig. 53). These factors are consistent



**Figure 53.** Suspended-sediment concentrations and the percentage of suspended sediment finer than 62 micrometers in diameter at Sulphur Creek Wasteway near Sunnyside, Yakima River Basin, Washington, 1989 (shaded area represents the irrigation season).

with the concept of sediment dilution—the mixing of fine-grained-sized, trace-element enriched, suspended sediment with coarse-grained-sized, trace-element depleted, suspended sediment. At the Sulphur Creek site, the irrigation-season decrease in suspended-silver concentration probably resulted from the diluting effect of coarse-grained-sized, agricultural soils eroded to Sulphur Creek Wasteway. Conversely, during the non-irrigation season, the fine-grained-sized sediment was probably enriched in silver from point-source effluent because of the absence of recently eroded agricultural soils.


For the Sulphur Creek Wasteway drainage, comparisons between the concentrations of suspended silver during the irrigation season and silver in streambed sediment were not possible because the limits of determination were lower in suspended sediment (0.1 µg/g) than in streambed sediment (2 µg/g).

## Zinc

Concentrations of zinc in streambed sediment in the Yakima River Basin ranged from 77 to 210 µg/g

(table 12) and slightly exceeded the 17 to 180 µg/g range of concentration which encompasses 95 percent of Western United States soils (R.C. Severson, U.S. Geological Survey, written commun., 1987, based on data in Shacklette and Boerngen, 1984). The median concentrations of zinc at the biological sites (100 µg/g) also exceeded those determined from analysis of fine-fraction streambed sediment in other river basins of the United States (table 34, at back of report). Concentrations of zinc increased slightly down the main stem and in contrast were notably enriched in several tributaries (fig. 54). Concentrations of zinc as high as 210, 200, and 174 µg/g were found in streambed sediment of the American River at Hells Crossing near Nile (site 13), in Wide Hollow Creek at the Old Sewage Treatment Plant at Union Gap (site 29), and in Wide Hollow Creek at West Valley Middle School near Ahtanum (site 27). The enrichment of zinc, measured in the American River, is probably of a geologic nature. Simmons and others (1983, p. 49) noted concentrations of zinc as high as 500 µg/g in streambed sediment of Mesatchee Creek. This upstream tributary to the American River was described as a zone of contact between andesitic rocks of the Ohanapecosh Formation and a granitic pluton. Conversely, the enrichment of zinc in Wide Hollow Creek drainage is not appreciable from a geologic source, rather it is probably the result of agricultural practices and urbanization. Geologic sources of zinc in Wide Hollow Creek Subbasin are small and confined to the Quaternary deposits and the loess geologic unit that contain a median zinc concentration of 88 µg/g (Fuhrer, McKenzie, and others, 1994).

The enrichment of zinc in Wide Hollow Creek at the old Sewage Treatment Plant—located near the mouth of Wide Hollow Creek—was expected because Wide Hollow Creek drains the urban area of Union Gap and because the median concentration of zinc in urban drains of the Yakima River Basin is 210 µg/g (Fuhrer, McKenzie, and others, 1994). However, the 174 µg/g of zinc at the upstream site, Wide Hollow Creek at West Valley High School, was not anticipated because the site is further removed from the higher density, urban effects of the Union Gap area. Most of the Wide Hollow Creek drainage above the West Valley High School site is used to support orchards. Agricultural practices associated with orchard crops may represent a nonpoint source of zinc to streambed sediment. According to a census of pesticide use in the Yakima River Basin for 1989, the zinc based pesticide




**Figure 54.** Zinc concentrations in streambed sediment of the main stem and tributaries, Yakima River Basin, Washington, 1987 ("American River" represents American River at Hells Crossing near Nile; "Wide Hollow Creek at old STP" represents Wide Hollow Creek at old sewage treatment plant at Union Gap; "Wide Hollow Creek" represents Wide Hollow Creek at West Valley Middle School near Ahtanum; "Jungle Creek" represents Jungle Creek near mouth near Cle Elum; and "Satus Creek" represents Satus Creek above Wilson-Charley Canyon near Toppenish).

Ziram (zinc dimethyldithiocarbamate) was applied to orchards in quantities that ranged from 4.7 to 8 pounds of zinc per acre per year (J.F. Rinella, unpub. data, 1996). Based on these applications, the quantity of elemental zinc ranged from 0.8 to 1.3 pounds per acre per year. In the mid-Yakima Valley alone, as much as 3,000 pounds of zinc was applied to apple, pear, and peach orchards in 1989. Soils of several former apple orchards in the Ahtanum Subbasin—the subbasin containing the Wide Hollow Creek drainage—contained concentrations of zinc as high as 150 µg/g, and streambed sediment within the same drainage contained concentrations of zinc as high as 180 µg/g (Fuhrer, McKenzie, and others, 1994). It was noted that the concentration of zinc was only 73 µg/g in a nearby agricultural plot used to grow peas. This concentration of zinc is understandably low (Ziram is not used on peas) compared to the former apple-orchard sites and agrees closely with the median concentration of zinc in the geologic units that form the soils and streambed sediments of the Ahtanum Creek Subbasin (Fuhrer, McKenzie, and others, 1994).

Median concentrations of zinc in suspended sediment ranged from 120 to 190 µg/g at the fixed

sites, and the lowest and highest median values, respectively, were found in the Naches River near North Yakima and the Yakima River at Cle Elum (table 13, fig. 55).

Sources of suspended zinc in the Yakima River at Cle Elum probably originate from the pre-Tertiary metamorphic and intrusive rocks and Miocene and older volcanic rocks geologic units in the northern part of the Kittitas Valley (Fuhrer, McKenzie, and others, 1994). Additionally, sources of zinc at the Cle Elum site probably are identical to geologic sources of antimony, arsenic, chromium, and nickel in the Kittitas Valley. Streambed-sediment samples, collected from lower-order streams in the pre-Tertiary rocks geologic unit of the Cle Elum River drainage and in the drainage upstream of Little Kachess Lake (both in the Kittitas Valley), contained zinc concentrations as high as 180 and 160 µg/g, respectively (Fuhrer, McKenzie, and others, 1994). Neighboring sites in the same drainage and geologic unit had zinc concentrations that were lower than 71 µg/g (Fuhrer, McKenzie, and others, 1994). The large interquartile range for suspended-sediment concentrations of zinc at the Cle Elum site probably results from the large spatial vari-



#### EXPLANATION

Interquartile range equals the value of the 75th percentile minus the value of the 25th percentile.

- More than 3 times the interquartile range from the 75-percentile value
- × 1.5 to 3 times the interquartile range from the 75-percentile value
- Less than 1.5 times the interquartile range from the 75-percentile value
- 75-percentile value
- Median value
- 25-percentile value
- Less than 1.5 times the interquartile range from the 25-percentile value

**Figure 55.** Distribution of zinc concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90 ("Cle Elum" represents Yakima River at Cle Elum; "Umtanum" represents Yakima River at Umtanum; "Naches" represents Naches River near North Yakima; "Union Gap" represents Yakima River above Ahtanum Creek at Union Gap; "Sulphur Creek" represents Sulphur Creek Wasteway near Sunnyside; "Grandview" represents Yakima River at Euclid Bridge at river mile 55 near Grandview; and "Kiona" represents Yakima River at Kiona).

ability of zinc concentrations in streambed sediment in the pre-Tertiary metamorphic and intrusive rocks geologic unit of the Kittitas Valley (Fuhrer, McKenzie, and others, 1994).

Although geologic sources of zinc exist in the Kittitas Valley, fluctuations in suspended-zinc concentrations during storms at the Yakima River at Umtanum indicate that these sources are relatively small in comparison to sources for chromium and nickel. Unlike the increase in concentration for chro-

mium and nickel at the Umtanum site near the peak of a December 5, 1989 rain-on-snow storm, the concentration of suspended zinc (160 µg/g) measured near the storm's peak was lower than that measured after the storm (230 µg/g) on December 13, 1989 (fig. 56). The pattern was the same during an April 6, 1989 storm at the Umtanum site; the zinc concentration measured near the storm's peak was lower than that measured after the storm (180 µg/g) on April 12, 1989. The zinc concentration measured after this storm was similar to that measured the preceding and following month. During the December 5, 1989 storm at Umtanum, similar patterns of decreasing element concentrations for cadmium, copper, and lead were measured near the peak of the storm.

With the exception of zinc concentrations in Asiatic clams in Spring Creek (452 µg/g) and in carp in the Yakima River at Kiona (634 µg/g), zinc concentrations in biota exhibited little variation among sites (table 15 and table 30). Such observations are common and may indicate metabolic regulation of zinc, an essential element (Bryan 1984; O'Grady and Abdullah, 1985; Luoma and others, 1990; Krantzberg and Stokes, 1989; Nuggeoda and Rainbow, 1989; Timmermans and others, 1992).

Zinc concentrations in most taxa from the Yakima River Basin appear typical of a zinc-poor system (table 39, at back of report) [Moore and others, 1991; Salanki and others, 1982; Luoma and others, 1990]. However, in Spring Creek, the mean concentration in Asiatic clams (452 µg/g) was more than twice that observed either at other sites sampled in this study or in the San Francisco Bay/Delta (Luoma and others, 1990). Zinc concentrations also were high in carp (634 µg/g) from the Yakima River at Kiona. Although in another study, carp seem to accumulate zinc to higher concentrations than other fish (Lowe and others, 1985), the concentrations measured in carp at Kiona are more than five times the 85th percentile concentration for all freshwater fish samples in California's Toxic Substances Monitoring Program (Rasmussen, 1992). Thus, some sites in the lower Yakima Valley seem to be enriched in zinc.

The concentrations of zinc ranged from 50 to 187 µg/g in curlyleaf pondweed and from 44 to 239 µg/g in the waterweed (table 15). The maximum concentrations in both species of aquatic plants were found in Wide Hollow Creek at the old Sewage Treatment Plant at Union Gap. High concentrations of zinc (201 µg/g) also were found in Waterweed in Spring



**Table 30. Comparison of low and high zinc concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987-91**

[For filtered water and suspended sediment, the low and high concentration assignments are based on a percentile distribution of the 50-percentile values (median) for each fixed site; For stream bed sediment and aquatic biota, the low and high concentration assignments are based on a percentile distribution of the mean concentrations for each fixed site. High concentrations (denoted with an "H" in the table) represent that portion of the distribution which is greater than or equal to the 75th-percentile value. Low concentrations (denoted with an "L" in the table) represent that portion of the distribution which is less than or equal to the 25th-percentile value. Concentrations greater than 25th, but less than 75th-percentile value are denoted with an "\*" in the table. The term "filtered water" is an operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45 micrometer filter; only 1990 data are summarized for largescale sucker livers and caddisflies; sample species: largescale sucker (*Catostomus macrocheilus*), mountain whitefish (*Prosopium williamsoni*), rainbow trout (*Oncorhynchus mykiss*), caddisfly (*Hydropsyche* spp.), stonefly (*Hesperoperla* sp.), Asiatic clam (*Veneroida*: Corbiculidae *Corbicula fluminea*), and curlyleaf pondweed (*Potamogeton crispus*); NF, North Fork; SF, South Fork; Cr = Creek; RM = river mile. Data statistically summarized for fixed sites are from monthly and selected hydrologic-event samplings from the Yakima River at Cle Elum, Yakima River at Umtanum, Naches River near North Yakima, Ahtanum Creek at Union Gap, Sulphur Creek Wasteway near Sunnyside, Yakima River at Euclid Bridge at river mile 55 near Grandview, and Yakima River at Kiona for the period 1987-90; to avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; --, no data]


Site reference number	Site name	Filtered water	Sediment		Aquatic biota				
					Fish livers			Insects	
			Streambed	Suspended	Largescale sucker	Mountain whitefish	Rainbow trout	Caddisfly	Stonefly
1	Waptus River at mouth near Roslyn	--	L	--	--	--	H	--	--
3	Jungle Creek near mouth near Cle Elum	--	H	--	--	--	*	--	--
4	NF Teanaway River below bridge at Dickey Creek Campground	--	--	--	--	--	--	--	L
5	Teanaway River below Forks near Cle Elum	--	*	--	--	--	H	--	--
6	Yakima River at Cle Elum	H	L	H	--	*	--	L	*
7	Naneum Creek below High Creek near Ellensburg	--	*	--	--	--	--	H	H
8	Taneum Creek at Taneum Meadow near Thorp	--	H	--	--	--	L	--	L
10	Little Naches River at mouth near Cliffdell	--	*	--	--	--	--	H	--
12	SF Manastash Cr near Ellensburg	--	H	--	--	--	L	--	H
13	American River at Hells Crossing near Nile	--	H	--	--	--	--	--	--

**Table 30.** Comparison of low and high zinc concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987-91—Continued

Site reference number	Site name	Filtered water	Sediment		Aquatic biota				
					Fish livers			Insects	
			Streambed	Suspended	Largescale sucker	Mountain whitefish	Rainbow trout	Caddisfly	Stonely
14	Cherry Creek above Wipple Wasteway at Thrall	--	L	--	--	H	--	*	--
16	Cherry Creek at Thrall	--	--	--	--	--	--	*	--
19	Yakima River at Umtanum	*	L	*	--	H	*	L	--
20	Umtanum Creek near mouth at Umtanum	--	*	--	--	--	*	*	--
21	Rattlesnake Creek above Little Rattlesnake near Nile	--	H	--	--	--	--	*	--
22	Rattlesnake Creek above North Fork Rattlesnake Creek near Nile	--	H	--	--	--	*	L	--
26	Naches River near North Yakima	L	H	L	H	L	--	*	--
27	Wide Hollow Creek at West Valley Middle School near Ahtanum	--	H	--	--	--	--	*	--
29	Wide Hollow Creek at old sewage treatment plant at Union Gap	--	H	--	--	--	*	H	--
30	Moxee Drain at Thorp Road near Union Gap	--	L	--	--	--	--	--	--
31	Ahtanum Creek at Union Gap	--	*	--	--	--	--	*	--
32	Yakima River above Ahtanum Creek at Union Gap	*	--	H	--	--	--	--	--
33	Yakima River at Parker	--	*	--	*	L	--	*	--

**Table 30.** Comparison of low and high zinc concentrations in water, sediment, and aquatic biota for selected sites, Yakima River Basin, Washington, 1987–91—Continued

Site reference number	Site name	Filtered water	Sediment		Aquatic biota				
					Fish livers			Insects	
			Streambed	Suspended	Largescale sucker	Mountain whitefish	Rainbow trout	Caddisfly	Stonely
34	South Fork Altanum Creek above Conrad Ranch near Tampico	--	H	--	--	--	--	H	*
40	Granger Drain at mouth near Granger	--	L	--	--	--	--	*	--
42	Yakima River below Toppenish Creek at RM 79.6 near Granger	--	*	--	*	H	--	L	--
43	Toppenish Creek at Indian Church Road near Granger	--	*	--	--	--	--	--	--
47	Satus Creek at gage at Satus	--	*	--	L	--	--	*	--
48	Yakima River at RM 72 above Satus Creek near Sunnyside	--	H	--	--	--	--	--	--
50	Yakima River at Kiona	L	*	H	*	--	--	L	--
52	Sulphur Creek Wasteway near Sunnyside	*	*	*	--	--	--	*	--
53	Satus Creek below Dry Creek near Toppenish	--	L	--	--	--	--	H	--
54	Spring Creek at mouth at Whitstran	--	L	--	--	--	--	*	--
56	Yakima River at Euclid Bridge at RM 55 near Grandview	H	*	L	*	*	--	L	--
57	Satus Creek above Wilson-Charley Canyon near Toppenish	--	H	--	--	--	*	H	*



**Figure 56.** Zinc concentrations in suspended sediment and suspended-sediment concentrations at the Yakima River at Umtanum, Yakima River Basin, Washington, 1989.

Creek at mouth at Whitstran and corresponded with similar enrichment in Asiatic clams, as previously mentioned. High concentrations of zinc in aquatic plants at the Wide Hollow Creek site coincided with the high concentrations of zinc measured in streambed sediment, as previously mentioned, and collectively indicated that Wide Hollow Creek is affected by anthropogenic sources of zinc.

## TEMPORAL VARIATIONS IN ELEMENT CONCENTRATIONS FOR AQUATIC BIOTA SAMPLED AT SITES IN COMMON IN 1989 AND 1990

The aquatic insect order *Trichoptera* (caddisfly) was sampled in 1989 and 1990. Among biological samples, this taxon had the greatest number of sites in common for 1989 and 1990 and, therefore, provided the best basinwide assessment of intrasite differences in element concentrations. Element concentrations in the *Trichoptera* were consistently higher in 1989 than in 1990 (table 31), possibly because of (1) differences between laboratory processing methods, (2) differences in bioaccumulation among *Trichoptera* species included in samples collected in 1989 and 1990, and (3) differences in bioaccumulation (exposure) between 1989 and 1990.

In 1989 and in 1990–91, respectively, the USFWS and the USGS laboratories prepared and analyzed samples from the Yakima River Basin (Fuhrer, Flutter, and others, 1994). One aspect of laboratory processing, which may have contributed to intrasite variability, is the difference in the degree that benthic insects were taxonomically sorted between 1989 and 1990. In 1990, benthic-insect samples were sorted to genus or species prior to chemical analysis. This level of taxonomic separation was not performed on samples collected in 1989; instead, samples were sorted to the order *Trichoptera* (caddisfly). In other studies, differences in the taxonomic composition of samples have been shown to affect comparisons of element concentrations between years (Cain and others, 1992) and may be a factor contributing to intrasite variability in the Yakima River Basin.

Asiatic-clam samples from the Yakima River indicated little change in elemental exposure between 1989 and 1990. For example, element concentrations in Asiatic clams from the Yakima River at Euclid Bridge at RM 55 near Grandview and the Yakima River at Kiona were similar between years (table 32). For most elements, the concentrations in the livers of rainbow trout, mountain whitefish, and largescale suckers also were similar between years (table 33). However, differences for cadmium and mercury in largescale suckers were measured between years. The

**Table 31.** Comparison of selected element concentrations in caddisflies sampled at sites in common in 1989 and 1990, Yakima River Basin, Washington

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, the mean element concentration of each site was statistically summarized; only detectable concentrations were statistically summarized; concentrations are reported in units of micrograms per gram, dry weight; identifications of caddisflies collected in 1989 were verified only to order (Trichoptera); samples for 1990 are Trichoptera: Hydropsychidae *Hydropsyche* spp.]

Element	Number of sites	1989			1990		
		Minimum	Median	Maximum	Minimum	Median	Maximum
Chromium	13	2.0	6.6	11	0.66	2.9	4.0
Copper	14	14	20	48	9.8	14	21
Iron	14	2,060	5,700	13,400	1,020	2,490	5,030
Lead	4	5.0	6.8	10	.81	2.3	5.6
Nickel	12	2.0	7.2	24	.93	3.2	5.7
Zinc	14	83	118	260	67	95	148

**Table 32.** Comparison of selected element concentrations in Asiatic clams sampled at sites in common in 1989 and 1990, Yakima River Basin, Washington

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, the mean element concentration of each site was statistically summarized; concentrations are reported in units of micrograms per gram, dry weight; Grandview, Yakima River at Euclid Bridge at river mile 55 near Grandview (site 56); Kiona, Yakima River at Kiona (site 50); sample species: Asiatic clam (Veneroida: Corbiculidae *Corbicula fluminea*)

Element	Grandview		Kiona	
	1989	1990	1989	1990
Arsenic	5.8	4.6	3.9	4.2
Cadmium	.30	.34	.20	.28
Chromium	2.0	2.0	2.5	1.2
Copper	33	35	23	27
Lead	<4.0	.33	<4.0	.18
Mercury	.15	.16	.10	.10
Nickel	<2.0	.93	2.0	1.2
Selenium	2.3	2.0	1.8	2.6
Zinc	93	99	106	98

minimum concentrations for cadmium and mercury were 13 times and 4 times higher, respectively, in 1989 than in 1990. These differences in concentrations may be related to differences in the sizes of largescale suckers collected between 1989 and 1990. The minimum concentrations of cadmium and mercury were mea-

sured in fish from Satus Creek in 1990. The mean length of these fish was 206 cm. The fish collected at the Satus Creek site in 1989, however, were larger (mean length 413 cm) and also had higher cadmium concentrations (0.4 µg/g). Little difference in either cadmium or mercury concentrations were noted between years at the other sites in the basin where similar sized fish could be compared. Copper concentrations in rainbow trout in the Yakima River at the mouth of Umtanum Creek differed 2 to 3 times between 1989 and 1990 (table 33). These differences did not seem to be related to size—between 1989 and 1990, the average length of fish at the Yakima River at Umtanum varied from only 248 to 293 cm. Analyses of these samples indicate copper exposures in this reach of the river increased between 1989 and 1990, although differences between years also could reflect interlaboratory variability.

## SUMMARY AND CONCLUSIONS

The Yakima River Basin is one of four surface-water pilot project areas selected to test and refine concepts for implementing a National Water Quality Assessment program. As part of the pilot project, major and trace elements were determined from several media that included streambed sediment, suspended sediment, water (filtered-water samples and unfiltered-water samples), and aquatic biota. Spatially, the most extensively sampled medium was streambed sediment; 448 sites were sampled in 1987 to determine

**Table 33.** Comparison of selected element concentrations in fish livers sampled at sites in common in 1989 and 1990, Yakima River Basin, Washington

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, the mean-element concentration of each site was statistically summarized; concentrations are reported in units of micrograms per gram, dry weight; Parker, Yakima River at Parker (site 33); Satus Creek, Satus Creek at gage at Satus (site 47); Kiona, Yakima River at Kiona (site 50); Cle Elum, Yakima River at Cle Elum (site 6); Umtanum Creek, Umtanum Creek near mouth at Umtanum (site 20); Yakima River, Yakima River at Umtanum (site 19); --, not determined; sample species: largescale sucker (*Catostomus macrocheilus*), mountain whitefish (*Prosopium williamsoni*), and rainbow trout (*Oncorhynchus mykiss*)]

Element	Largescale sucker						Mountain whitefish				Rainbow trout			
	Parker		Satus Creek		Kiona		Cle Elum		Kiona		Umtanum Creek		Yakima River	
	1989	1990	1989	1990	1989	1990	1989	1990	1989	1990	1989	1990	1989	1990
Arsenic	<0.20	0.20	<0.20	0.30	0.27	0.30	<0.20	0.10	<0.20	0.20	<0.2	<0.50	0.20	<0.50
Cadmium	<.20	.32	.40	.03	.47	.43	<.20	.13	1.1	1.4	<.30	.23	<.20	.10
Chromium	<1.0	.78	<1.0	.50	<1.0	.59	<1.0	1.0	<1.0	.64	<1.0	.84	<1.0	.75
Copper	11	23	27	26	21	32	6.1	6.0	9.6	7.7	59	100	130	290
Lead	<4.0	.15	<4.0	.17	<4.0	.17	.91	.72	.58	.76	--	--	--	--
Mercury	.21	.47	.27	.05	.28	.30	<2.0	.18	<2.0	<.06	.25	.38	.22	.27
Selenium	2.6	2.9	3.3	1.9	3.7	4.5	<4.0	.22	<4.0	.31	--	--	--	--
Zinc	86	100	120	60	84	85	9.5	8.2	11	15	101	81	109	80

the occurrence and spatial distribution of potentially toxic major and trace elements (Fuhrer, McKenzie, and others, 1994). Thirty-two of the sites that had been sampled for streambed sediment in 1987 were sampled for aquatic biota in 1989–91. Seven of these sites, termed fixed-sampling sites, were sampled for suspended sediment and other water media (monthly and during several hydrologic events) for the period 1987–90. Trace elements were measured in filtered-water samples at least once at 44 sites; most of these sites were sampled over a period of 1 to 2 weeks (synoptic samplings) in July and (or) November 1987. Trace elements generally were measured quarterly in unfiltered-water samples during 1987 at fixed-location sampling sites.

This report describes the occurrence and distribution of potentially toxic trace elements in sediment, water, and aquatic biota. More specifically, this report provides information about:

1. Variations in element concentrations and loads attributable to different hydrologic conditions, including the irrigation season, non-irrigation season, snowmelt season, and storms;
2. Suitability of surface water for the protection of aquatic life and human health based on State and Federal water-quality criteria, guidelines, and regulations;

3. Major natural and human factors in the Yakima River Basin that affect observed water-quality conditions; and
4. Patterns of element enrichment which are common among sediment, water, and aquatic biota media.

**Comparison of trace-element concentrations in water and fish muscle to water-quality guidelines**—Trace-element concentrations in filtered- and unfiltered-water samples were screened against U.S. Environmental Protection Agency (EPA) ambient water-quality criteria for the protection of aquatic life and human health, drinking-water regulations, and drinking-water human-health advisories. Although all EPA ambient water-quality criteria are nonenforceable guidelines, the guidelines were used to screen ambient water-quality data in the Yakima River Basin in order to identify element concentrations that may require study by State and local health agencies. Concentrations of cadmium, chromium, copper, iron, lead, mercury, silver, and zinc in filtered and (or) unfiltered water exceeded the screening value (based on EPA's ambient water-quality criteria for the protection of aquatic organisms) at two or more sites in the Yakima River Basin. Zinc concentrations in filtered water exceeded acute and chronic criteria for aquatic life at several sites, including those receiving irrigation return flow and those located in mountainous areas.

Copper exceedances occur during winter-storm runoff periods, and coincide with seasonal historical patterns of copper exceedances attributed, in part, to the use, past and present, of copper sulfate (a herbicide).

The EPA ambient water-quality criteria for the protection of human health are designed to indicate exposure of humans to a contaminant because of (1) consumption of water and aquatic organisms or (2) consumption of aquatic organisms only. Concentrations of arsenic (a carcinogen) exceeded the human-health screening value for consumption of aquatic organisms and water in 47 percent of the filtered-water samples and exceeded the screening value for consumption of only aquatic organisms in 30 percent of the samples. Exceedances of arsenic were measured predominantly in the lower Yakima Valley. Concentrations of mercury (a noncarcinogen) in filtered-water samples exceeded the human-health screening values for consumption of aquatic organisms and water and consumption of only aquatic organisms in 4 percent of the samples.

Trace-element concentrations, determined from filtered- and unfiltered-water samples, were screened by making comparisons with EPA drinking-water regulations and EPA advisories for human health. Because filtered and unfiltered stream-water samples represent untreated water, element concentrations which exceeded screening values (based on drinking-water regulations) do not indicate that human health is directly at risk. Concentrations of iron in unfiltered-water samples did not meet the screening value in 94 percent of the samples. In filtered-water samples, iron concentrations, however, did meet the screening value; therefore, the exceedances in unfiltered water probably resulted from iron associated with sediment that probably would be removed in a water-treatment process.

Unlike the ambient water-quality criteria for human health, drinking-water health advisories are based only on the consumption of domestic water. In the present study, however, ambient stream water was used to screen for health effects. Concentrations of arsenic (a carcinogen) in filtered-water samples exceeded the screening value in 31 percent of the samples. The largest number of exceedances of arsenic was found in the lower Yakima Valley. Concentrations of mercury (a noncarcinogen) in filtered-water samples rarely exceeded the screening value.


Fish muscle, analyzed for mercury for various fish taxa, was collected from four sites in the Yakima

River Basin in 1991. The median mercury concentration in fish muscle from each site and for each fish species was screened against mercury concentrations in fish that were of potential public-health concern. Muscle samples collected from rainbow trout and mountain whitefish from the four sites contained mercury concentrations that were below the screening value for standard adults (consumers of an average of about one 6-ounce filet per month). Screening values also were calculated for children, recreational fishermen, and subsistence fishermen. The concentration of mercury in fish muscle, consumed by children who ate an average of one 6-ounce filet per month, exceeded the screening value for children for all species of fish sampled and at all sites sampled. Similarly, screening values for recreational fishermen (consumers of about five 6-ounce filets per month) and subsistence fishermen (consumers of about 25 6-ounce filets per month) are exceeded for all species of fish sampled and at all sites sampled.

***Distribution of elements in sediment, water, and aquatic biota***—Most element enrichment results from natural-geologic sources in the forested landscapes of the Kittitas and mid-Yakima Valley, primarily in the Cle Elum, Upper Naches, Teanaway and Tieton Subbasins (see table below and figure on p. 154). These areas might be classified as “pristine”

Element	Dominant Source		
	Geology	Agriculture	Urban and light Industry
Arsenic	Kittitas Valley	Mid-Yakima Valley and lower Yakima Valley	
Chromium	Kittitas Valley		
Copper	Mid-Yakima Valley	Mid-Yakima Valley	
Lead		Mid-Yakima Valley and lower Yakima Valley	
Mercury	Kittitas Valley and Mid-Yakima Valley		
Nickel	Kittitas Valley		
Selenium	Mid-Yakima Valley		
Zinc	Mid-Yakima Valley	Mid-Yakima Valley	





Distinctive-hydrologic reaches of the Yakima River, Washington.

by the casual observer yet they are geologic sources of antimony, arsenic, chromium, copper, mercury, nickel, selenium, and zinc. As an example, arsenic, chromium, and nickel concentrations in streambed sediment (as high as 45, 212, and 260  $\mu\text{g/g}$ , respectively) were nearly 4 to as much as 13 times higher than their respective median concentrations in streambed sediment of agricultural land-use areas in the lower Yakima Valley. As a result of geologic sources, several of these elements, including arsenic, chromium, and nickel, leave chemical signatures that were measurable in streambed sediment and suspended sediment of higher-order streams, including the main stem. In addition to streambed sediment and suspended sediment, some of the geologically derived elements, including chromium, nickel, and selenium, were measurable in aquatic biota of higher-order streams. Only the Teanaway Subbasin was a qualified source of chro-

mium and nickel to sediment and aquatic biota. In the Teanaway Subbasin chromium and nickel concentrations in benthic insects were, respectively, 4 to 52 times and 43 to 102 times higher than concentrations in insect taxa collected from other sites in the Yakima River Basin. Enrichment of chromium and nickel from the Teanaway Subbasin also affected sediment and biota, although to a lesser extent, in the Yakima River at Umtanum. The median concentration of nickel in suspended sediment at the Umtanum site was more than twice that in the Yakima River at Kiona.

Streambed sediment, formed from the Quaternary deposits and loess geologic unit, comprises a majority of the agricultural farmland in the Yakima River Basin and helps control the downstream affects of geologically derived trace elements. Concentrations of several elements, including arsenic, chromium, and nickel, result from

geologic sources and were attenuated by mixing with element-poor streambed sediment from agricultural lands. For example, concentrations of arsenic in the Yakima River at Cle Elum were high as a result of geologic sources in the Cle Elum Subbasin; however, farther downstream at the Yakima River at Umtanum, a site that receives sediment from agricultural land-use areas, arsenic concentrations were 16 times lower than those found in close proximity to geologic sources. Lower arsenic concentrations at Umtanum result from streambed sediment that enters the main stem from agricultural land located in the Quaternary deposits and loess geologic unit in the Kittitas Valley.

In some parts of the Yakima River Basin, human activities (such as farming) can decrease some trace-element concentrations, in other parts they can increase element concentrations. Trace-element distributions of cadmium, copper, lead, mercury, selenium, and zinc increase in areas affected by human activities. Concentrations of the aforementioned elements frequently were highest in the Wide Hollow Subbasin, which drains urbanized and lightly industrialized lowland, as well as agricultural land in the upper reaches of Wide Hollow Subbasin. Concentrations of lead in streambed sediment of Wide Hollow Creek, for example, were more than twice that expected in streambed sediment from geologic sources in Wide Hollow Subbasin and also exceeded the 5.2 to 55  $\mu\text{g/g}$  range of concentrations which encompasses 95 percent of Western United States soils. In addition to urban runoff, past applications of lead arsenate in apple orchards, including applications in the upper reaches of Wide Hollow Subbasin, may be a source of lead. Benthic insects seemed to be sensitive to lead in Wide Hollow Creek. Lead concentrations in caddisflies (*Hydropsyche* spp.) in Wide Hollow Creek not only represented the basin maximum, but also were 15 times higher than concentrations in caddisflies from Umtanum Creek—a reference site not affected directly by human activities. Although lead concentrations were high in caddisflies in Wide Hollow Creek, caddisflies were only slightly enriched compared to insects in other systems; lead concentrations were not high in fish. Among liver samples from the basin's different fish taxa, the concentration of selenium in rainbow trout from Wide Hollow Creek (31  $\mu\text{g/g}$ ) was 4 to 8 times higher than concentrations of selenium in rainbow trout from other sites in the basin. In Wide Hollow Creek, cadmium concentrations in *Hydropsyche californica*, a caddisfly species, increased 3 times from the upstream site (0.07  $\mu\text{g/g}$ ) to the downstream site. This increase probably

results from larger proportions of urban area at the downstream site.

In streambed sediment of the Kittitas Valley, human activities have been shown, in some instances, to affect arsenic concentrations that result from geologic sources. In filtered water, suspended sediment, and aquatic biota of the mid- and lower Yakima Valley, however, human activities can enhance arsenic concentrations in addition to arsenic loads. As with lead, sources of arsenic in the mid-Yakima and lower Yakima Valley included lead-arsenate pesticides that have been applied historically to apple orchards. Consequently, agricultural drains seemed to be good indicators of past arsenic use. Suspended-arsenic concentrations in Sulphur Creek Wasteway, an agricultural drain, were the highest in the basin; these arsenic concentrations ranged from 4.9 to 20  $\mu\text{g/g}$ . The annual suspended-arsenic loads during 1987–90 between the Kittitas Valley and the mid-Yakima Valley increased as much as 3 times. During the irrigation season, in particular, about 2.2 pounds of suspended arsenic per day entered the mid-Yakima Valley over a 9.4-mile reach that receives irrigation return flow from Moxee Subbasin and Wide Hollow Subbasin. This arsenic load represents about one-half the irrigation-season load in the Yakima River above Ahtanum Creek at Union Gap. Irrigation drains probably are large contributors over this reach—Moxee Drain is estimated to contribute nearly 1 pound of suspended arsenic per day. During the irrigation season in the lower Yakima Valley, the June contributions of suspended arsenic (2 pounds per day) from Sulphur Creek Wasteway typically account for most of the suspended-arsenic load in the Yakima River at Euclid Bridge at RM 55 near Grandview—located 6 miles downstream from Sulphur Creek Wasteway.

Concentrations of arsenic in filtered-water samples in Sulphur Creek Wasteway and in the main stem of the lower Yakima Valley were high (exceeded the 90th percentile (3  $\mu\text{g/L}$ ) for the Yakima River Basin) in comparison to fixed sites (<1  $\mu\text{g/L}$ ) in the Kittitas Valley. These high concentrations were found in waters affected primarily by agricultural return flow. In addition to higher concentrations of arsenic in filtered-water samples from agriculturally affected portions of the basin, the load of arsenic in agricultural drains probably represents a large proportion of the arsenic load passing the Yakima River at Kiona, the terminus of the basin. Sulphur Creek Wasteway, for

example, has an annual streamflow representing only about 8 percent of the annual streamflow at the Kiona fixed site, yet accounts for nearly 20 percent of the filtered-arsenic load at Kiona. Comparisons, between loads determined from filtered-water samples (an operational approximation of dissolved load) and loads determined from arsenic in suspended sediment, showed that a majority of the arsenic load in the basin is in dissolved form. For example, the annual dissolved-arsenic loads in the lower Yakima Valley at Sulphur Creek Wasteway, Grandview, and Kiona sites were from 4 to 9 times higher than their respective suspended loads.

Arsenic also was present in aquatic biota in the Yakima River Basin. In curlyleaf pondweed, an aquatic plant, concentrations ranged from 0.48 to 1.5  $\mu\text{g/g}$  and were 3 times higher in the main stem of the lower Yakima Valley than in the main stem of the Kittitas Valley. Concentrations of arsenic in caddisflies, collected from agricultural drains in 1989 in the lower Yakima Valley, were as large as 5.4  $\mu\text{g/g}$  and exceeded the 85th percentile concentration for the basin. Asiatic clams were collected only from the lower Yakima Valley, and arsenic concentrations (3.6 to 4.6  $\mu\text{g/g}$ ) varied little among sites. Compared to other studies, however, arsenic concentrations in Asiatic clams of the lower Yakima Valley were as much as an order of magnitude greater than in clams in uncontaminated or minimally contaminated environments.

Fish taxa provided the most comprehensive spatial coverage for arsenic, mercury, and selenium; however, no single fish taxon was distributed widely across the Yakima River Basin. The aquatic insect taxon *Hydropsyche* spp. (caddisfly) provides the most comprehensive spatial coverage of any single insect taxon. Concentrations of several trace elements, including cadmium, mercury, and selenium, found in various taxa were higher in the main stem of the lower Yakima Valley than in the Kittitas and mid-Yakima Valley. In mountain whitefish livers from the lower Yakima Valley, the concentration of mercury in the Yakima River at Parker (1.3  $\mu\text{g/g}$ ) and of selenium in the Yakima River at Kiona (15  $\mu\text{g/g}$ ) was nearly twice that measured in mountain whitefish in the Kittitas Valley at the Cle Elum site. Similar patterns also were observed for largescale suckers. Compared to other studies of mercury in liver tissue of pike and mud fish, mercury concentrations in some mountain whitefish in the lower Yakima Valley were indicative of moderate enrichment. The concentration of selenium in sculpin

(5.5  $\mu\text{g/g}$ ) in the northeastern Cascade Range was about 10 times higher than in Satus Creek, a lower Yakima Valley tributary. Concentrations of selenium in sculpin in the Yakima River Basin (mean and 85th percentile values) also exceeded those for sculpin in the National Contaminant Biomonitoring Program. Because no single fish species is pervasive throughout the Yakima River Basin, mountain whitefish were good indicators of selenium enrichment in the lower Yakima Valley (where they reside) and sculpin are good indicators of selenium enrichment at sites in close proximity to geologic sources in the northeastern Cascade Range.

**Implications for water-resource monitoring and regulation**—Trace elements that should be considered as part of future water-resource monitoring in the Yakima River Basin include arsenic, cadmium, chromium, copper, lead, nickel, mercury, selenium, and zinc. Some of these trace elements in streams (for example, chromium) are primarily from geologic sources, some (for example, lead) are primarily from anthropogenic sources, and others (for example, arsenic, copper, and zinc) are from both sources. Thus, because sources of trace elements may vary, monitoring-design strategies may need to vary among subbasin(s).

The presence of anomalous concentrations of arsenic, lead, and zinc in streambed sediment of agricultural lands and soils in the mid-Yakima and lower Yakima Valley indicates that agricultural practices are a source of arsenic, lead, and zinc to streambed sediment—particularly, agricultural practices that tend to facilitate soil loss or erosion. Additionally, the presence of arsenic in filtered-water samples from Sulphur Creek Wasteway near Sunnyside (the only agricultural drain sampled) indicates that other agricultural drains may be sources of arsenic to the lower Yakima Valley. Similarly, the presence of arsenic in Sulphur Creek Wasteway, especially the presence of higher arsenic concentrations during the non-irrigation season, indicates that shallow ground water in areas of intense irrigation, although not measured, also may be affected by arsenic.

The screening of ambient water-quality data in the Yakima River Basin was done in order to assist State and local health agencies in the identification of element concentrations that may require further study. Concentrations of arsenic in ambient stream water in

the lower Yakima Valley frequently exceeded the screening values based on EPA's ambient stream-water-quality criteria and EPA human-health advisories for drinking water. These exceedances primarily are associated with waterways affected by irrigation return flow and may be indicative of shallow ground-water quality in some agricultural areas or subbasins.

Concentrations of mercury in fish muscle also frequently exceeded screening values based on EPA's guidance document for assessing chemical contaminant data for use in fish advisories. Depending on the quantity of fish consumed by individuals, the concentration of mercury in fish muscle exceeded the screening value for standard adults, children, recreational fishermen, and subsistence fishermen. Relative to future monitoring activities, the measurement of mercury in streambed sediment was not a sensitive indicator of mercury uptake by biota in the lower Yakima Valley, rather fish-liver and fish-muscle samples were the preferred sample media.

## SELECTED REFERENCES

- Adams, W.J., and Johnson, H.E., 1977, Survey of the selenium content in the aquatic biota of western Lake Erie: *Journal of Great Lakes Research*, v. 3, p. 10–14.
- Arbogast, B.F., ed., 1990, Quality assurance manual for the U.S. Geological Survey Branch of Geochemistry: U. S. Geological Survey Open-File Report 90–668, 184 p.
- Axtmann, E.V., Cain, D.J., and Luoma, S.N., 1991, Distribution of trace metals in fine-grained bed sediments and benthic insects in the Clark Fork River, Montana, p. 1-8; *in* Proceedings of the Clark Fork Symposium, 1989: Missoula, Montana, Montana Academy of Sciences; Butte Montana College of Mineral Sciences and Technology.
- Barak, N.A.E., and Mason, C. F., 1990, Mercury, cadmium and lead concentrations in five species of freshwater fish from eastern England: *The Science of the Total Environment*, v. 92, p. 257–263.
- Bonneville Power Administration, July 1985, Issue Alert, Yakima Basin passage improvement: Report 1A-4-18, 8 p.
- \_\_\_\_\_, August 1988, Issue alert, Yakima and Klickitat salmon and steelhead: Report BP-954, 8 p.
- Bryan, G.W., 1984, Pollution due to heavy metals and their compounds, *in* Kinne, O., ed., *Marine Ecology*: New York, John Wiley and Sons, v. 3, p. 1289–1431.
- Buchanan, T.J., and Somers, W.P., 1969, Discharge measurements at gaging stations: U.S. Geological Survey Techniques of Water-Resources Investigations, book 3, chapter A8, 65 p.
- Cain, D.J., Luoma, S.N., Carter, J.L., and Fend, S.V., 1992, Aquatic insects as bioindicators of trace element contamination in cobble-bottom rivers and streams: *Canadian Journal of Fisheries and Aquatic Science*, v. 49, no. 10, p. 2141–2154.
- Church, S.E., Mosier, E.L., Frisken, J.G., Motooka, J.M., Gruzensky, S.L., Ficklin, W.H., McCollaum, A.D., Willson, W.R., and McDanal, S.K., 1983, Geochemical and statistical analysis of analytical results for stream sediments, rocks, and waters collected from the Goat Rocks Wilderness and adjacent roadless areas, Lewis and Yakima Counties, Washington: U.S. Geological Survey Open-File Report 83–0784, 272 p.
- Cohn, T.A., Caulder, D.L., Gilroy, E.J., Zynjuk, LD., and Summers, R.M., 1992, The validity of a simple statistical model for estimating fluvial constituent loads: An empirical study involving nutrient loads entering Chesapeake Bay: *Water Resources Research*, v. 28, no. 9, p. 2353–2363.
- Colman, J. A., and Sanzolone, R. F., 1991, Surface-water-quality assessment of the Upper Illinois River Basin in Illinois, Indiana, and Wisconsin: Geochemical data for fine-fraction streambed sediment from high- and low-order streams, 1987: U.S. Geological Survey Open-File Report 90–571, 108 p.
- Columbia Basin Inter-Agency Committee, 1964, River-mile index, Yakima River: Columbia River Basin Hydrology Subcommittee Report 15, 39 p.
- Crawford, J.K., and Luoma, S.N., 1993, Guidelines for studies of contaminants in biological tissues for the National Water-Quality Assessment program: U.S. Geological Survey Open-File Report 92–0492, 69 p.
- Edwards, T.K., and Glysson, G.D., 1988, Field methods for measurement of fluvial sediment: U.S. Geological Survey Open-File Report 86–531, 118 p.
- Eisler, Ronald, 1985, Cadmium hazards to fish, wildlife, and invertebrates: A synoptic review: U.S. Fish and Wildlife Service—Contaminant Hazard Reviews, no. 2, 46 p.
- \_\_\_\_\_, 1988, Arsenic hazards to fish, wildlife, and invertebrates: A synoptic review: U.S. Fish and Wildlife Service—Contaminant Hazard Reviews, no. 12, 92 p.
- Elder, J.F., 1988, Metal biogeochemistry in surface-water systems: A review of principles and concepts: U.S. Geological Survey Circular 1013, 43 p.

- Elder, J.F., and Matraw, H.C., Jr., 1984, Accumulation of trace elements, pesticides, and polychlorinated biphenyls in sediments and the clam *Corbicula manilensis* of the Apalachicola River, Florida: *Archive Environmental Contamination and Toxicology*, v. 13, p. 453–469.
- Elwood, J.W., Hildebrand, S.G., and Beauchamp, J.J., 1976, Contribution of gut contents to the concentration and body burden of elements in *Tipula* spp. from a spring-fed stream: *Journal of the Fisheries Research Board of Canada*, v. 33, p. 1930–1938.
- Embrey, S.S., 1992, Surface-water-quality assessment of the Yakima River Basin, Washington: Areal distribution of fecal-indicator bacteria, July 1988: U.S. Geological Survey Water-Resources Investigations Report 91–4073, 34 p.
- Federal Register, 1992, v. 57, no. 246, p. 60887.
- Finley, K.A., 1985, Observations of bluegills fed selenium-contaminated *Hexagenia* nymphs collected from Belews Lake, North Carolina: *Bulletin of Environmental Contamination and Toxicology*, v. 35, p. 816–825.
- Fishman, M.J., and Friedman, L.C., eds., 1989, Methods for determination of inorganic substances in water and fluvial sediments (3d ed.): U. S. Geological Survey Techniques of Water-Resources Investigations, Book 5, Chapter A1, 545 p.
- Food and Drug Administration, 1985, Action levels for poisonous or deleterious substances in human food and animal feed: Washington, D.C., Center for Food Safety and Applied Nutrition, Industry Programs Branch, 13 p.
- Forstner, U., and Wittmann, G.T.W., 1979, Metal pollution in the aquatic environment: New York, Springer-Verlag, 486 p.
- Fuhrer, G.J., 1989, Quality of bottom material and elutriates in the Lower Willamette River, Portland Harbor, Oregon, U.S. Geological Survey Water-Resources Investigations Report 89–4005, 30 p.
- Fuhrer, G.J.; McKenzie, S.W.; Rinella, J.F.; Sanzalone, R.F.; and Skach, K.A., 1994, Surface-water-quality assessment of the Yakima River Basin in Washington: Analysis of major and minor elements in fine-grained streambed sediment, 1987, with sections on Geology by Marshall W. Gannett.: U.S. Geological Survey Open-File Report 93–30, 131 p. (pending publication as U.S. Geological Survey Water Supply Paper 2354-C)
- Fuhrer, G.J., Fluter, S.L., McKenzie, S.W., Rinella, J.F., Crawford, J. K., Cain, D.J., Hornberger, M.I., Bridges, J.L., and Skach, K.A., 1994, Surface-water-quality assessment of the Yakima River Basin in Washington: Major- and minor-element data for water, sediment, and aquatic biota, 1987–91: U.S. Geological Survey Open-File Report 94–308, 223 p.
- Gower, A.M., and Darlington, S.T., 1990, Relationships between copper concentrations in larvae of *Plectrocnemia conspersa* (Curtis) (Trichoptera) and in mine drainage streams: *Environmental Pollution*, v. 65, p. 155–168.
- Gualtieri, J.L., and Simmons, G.C., 1989, Geochemical exploration of the Alpine Lakes study area and additions, Washington, in U.S. Geological Survey and U.S. Bureau of Mines, Mineral resources of the alpine lakes study area and additions, Chelan, King, and Kittitas Counties, Washington: U.S. Geological Survey Bulletin 1542, p. 53–84 and p. 234–317.
- Hem, J.D., 1989, Study and interpretation of the chemical characteristics of natural water: U.S. Geological Survey Water-Supply Paper 2254, 263 p.
- Hirsch, R.M., Alley, W.M., and Wilber, W.G., 1988, Concepts for a National Water-Quality Assessment program: U.S. Geological Survey Circular 1021, 42 p.
- Hopkins, B.S., Clark, D.K., Schlender, M., and Stinson, M., 1985, Basic water monitoring program: Fish tissue and sediment sampling for 1984: Olympia, Washington State Department of Ecology, 43 p.
- Horowitz, A.J., 1991, A primer on sediment-trace element chemistry (2d ed.): Chelsea, Michigan, Lewis Publishers, 136 p.
- Horton, R. E., 1945, Erosional development of streams and their drainage basins; hydrophysical approach to quantitative morphology: *Geological Society of America Bulletin*, v. 56, p. 275–370.
- Jenne, E.A., and Luoma, S.N., 1977, Forms of trace elements in soils, sediments, and associated waters: An overview of their determination and biological availability, in *Biological implications of metals in the environment: Energy Research and Development Administration Series 42, CONF-750929*, p. 110–143.
- Johns, C., and Luoma, S.N., 1990, Arsenic in benthic bivalves of San Francisco Bay and the Sacramento/San Joaquin River Delta: *The Science of the Total Environment*, v. 97/98, p. 673–684.
- Johns, C., Luoma, S.N., and Elrod, V., 1988, Selenium accumulation in benthic bivalves and fine sediments of San Francisco Bay, the Sacramento-San Joaquin Delta, and selected tributaries: *Estuarine, Coastal and Shelf Science*, v. 27, p. 381–396.
- Johnson, A., Norton, D., and Yake, W., 1986, Occurrence and significance of DDT compounds and other contaminants in fish, water, and sediment for the Yakima River Basin: Olympia, Washington Department of Ecology, 89 p.
- Kaiser, I.I., Young, P.A., and Johnson, J.D., 1979, Chronic exposure of trout to waters with naturally high selenium levels: Effects on transfer RNA methylation: *Journal of Fisheries Resources Board of Canada*, v. 36, p. 689–694.

- Kendra, W., 1988, Quality of water, sediment, and biota in Wide Hollow Creek, Washington: Olympia, Washington, Washington State Department of Ecology, 39 p.
- Koli, A.K., Williams, W.R., McClary, E.B., Wright, E.L., and Burrell, T.M., 1977, Mercury levels in freshwater fish of the state of South Carolina: Bulletin of Environmental Contamination and Toxicology, v. 17, p. 82–89.
- Kopp, J.F., and Kroner, R.C., 1968, Trace metals in water in the United States, Oct. 1, 1962–Sept. 30, 1967: U.S. Department of the Interior, Federal Water Pollution Control Administration, 48 p.
- Krantzberg, G., and Stokes, P.M., 1989, Metal regulation, tolerance, and body burdens in the larvae of the genus *Chironomus*: Canadian Journal of Fisheries and Aquatic Sciences, v. 46, p. 389–398.
- Langston, W.J., 1982, The distribution of mercury in British estuarine sediments and its availability to deposit-feeding bivalves: Journal of the Marine Biological Association of the United Kingdom, Plymouth, v. 62, p. 667–684.
- Leahy, P.P., Rosenshein, J.S., and Knopman, D.S., 1990, Implementation plan for the National Water-Quality Assessment program: U.S. Geological Survey Open-File Report 90–174, 10 p.
- Leland, H.V., and Scudder, B.C., 1990, Trace elements in *Corbicula fluminea* from the San Joaquin River, California: The Science of the Total Environment, v. 97/98, p. 641–672.
- Levinson, A.A., 1980, Introduction to exploration geochemistry (2d ed.): Wilmette, Illinois, Applied Publishing Ltd., 924 p.
- Lodenius, M., Seppanen, A., and Herranen, M., 1982, Accumulation of mercury in fish and man from reservoirs in northern Finland: Water, Air, and Soil Pollution, v. 19, p. 237–246.
- Lowe, T.P., May, T.W., Brumbaugh, W.G., and Kane, D.A., 1985, National contaminant biomonitoring program: concentrations of seven elements in freshwater fish, 1978–1981: Archives of Environmental Contamination and Toxicology, v. 14, p. 363–388.
- Lucas, J.M., 1975, The availability of nickel, chromium, and silver in Washington: Washington Department of Natural Resources, Division of Geology and Earth Resources, Open-File Report 75–14, 140 p.
- Luoma, S.N., 1983, Bioavailability of trace metals to aquatic organisms—a review: Science of the Total Environment, v. 28, p. 1–22.
- \_\_\_\_\_, 1986, Cycling of lead into food webs in aquatic environments: Report for the Commission on Lead in the Environment: Ottawa, Ontario, Royal Society of Canada, p. 146–161.
- \_\_\_\_\_, 1989, Can we determine the biological availability of sediment-bound trace elements?: Hydrobiologia, v. 176/177, p. 379–396.
- Luoma, S.N.; Dagovitz, R.; and Axtmann, E., 1990, Temporally intensive study of trace metals in sediments and bivalves from a large river-estuarine system—Suisun Bay/Delta in San Francisco Bay: The Science of the Total Environment, v. 97/98, p. 685–712.
- Lynch, T.R., Popp, C.J. and Jacobi, G.Z., 1988, Aquatic insects as environmental monitors of trace metal contamination: Red River, New Mexico: Water, Air, and Soil Pollution, v. 42, p. 19–31.
- McCleneghan, K., Meinz, T.M., Crane, D., Setu, W., and Lew, T., 1981, Toxic substances monitoring program 1980: State of California: State Water Resources Control Board Water Quality Monitoring Report No. 81–8TS, 54 p., plus appendixes I–VI.
- McFarlane, G.A., and Franzin, W.G., 1980, An examination of Cd, Cu, and Hg concentrations in livers of northern pike, *Esox lucius*, and white sucker, *Catostomus commersoni*, from five lakes near a base metal smelter at Flin Flon, Manitoba: Canadian Journal of Fisheries and Aquatic Science, v. 37, p. 1573–1578.
- McKenzie, S.W., and Rinella, J.F., 1987, Surface-water-quality assessment of the Yakima River Basin, Washington: Project description: U.S. Geological Survey Open-File Report 87–238, 35 p.
- Miller, P.A., Minkittrick, K.R., and Dixon, D. G., 1992, Relationship between concentrations of copper and zinc in water, sediment, benthic invertebrates, and tissues of white sucker (*Catostomus commersoni*) at metal-contaminated sites: Canadian Journal of Fisheries and Aquatic Science, v. 49, p. 978–984.
- Moen, W.S., 1969, Analysis of stream sediment samples in Washington for copper, molybdenum, lead, and zinc: Washington Department of Natural Resources, Division of Geology and Earth Resources, Open-File 69–2, 38 p.
- Moore, J.W., 1991, Inorganic contaminants of surface water: Research and monitoring priorities: New York, Springer-Verlag, 334 p.
- Moore, J.N., Luoma, S.N., and Peters, D., 1991, Downstream effects of mine effluent on an intermontane riparian system: Canadian Journal of Fisheries and Aquatic Sciences, v. 48, p. 222–232.
- National Academy of Sciences, 1972, Water quality criteria 1972—A report of the committee on water quality criteria, environmental studies board: EPA-R3-73-033-March 1973, 594 p.
- Nowell, L.H., and Resek, E.A., 1994, Summary of national standards and guidelines for pesticides in water, bed sediment, and aquatic organisms and their application to water-quality assessments: U.S. Geological Survey Open-file Report 94–44, 115 p.

- Nugegoda, D., and Rainbow, P.S., 1989. Effects of salinity changes on zinc uptake and regulation by the decapod crustaceans *Palaemonetes varians*: Marine Ecology Progress Series, v. 51, p. 57–75.
- Ogle, R.S., Maier, K.J., Kiffney, P., Williams, M.J., Brasher, A., Melton, L.A., and Knight, A.W., 1988, Bioaccumulation of selenium in aquatic ecosystems: Lake and Reservoir Management, v. 4, p. 165–173.
- O'Grady, K.T., and Abdullah, M.I., 1985, Mobility and residence of zinc in brown trout *Salmo trutta*: Results of environmentally induced change through transfer: Environmental Pollution (series A), v. 38, p. 109–127.
- Parker, R.L., 1967, Data of geochemistry—Chapter D: Composition of the earth's crust: U.S. Geological Survey Professional Paper 440–D, 19 p.
- Persaud, D., Jaagumagi, R., and Hayton, A., 1993, Guidelines for the protection and management of aquatic sediment quality in Ontario: Ontario Ministry of the Environment, Water Resources Branch, 24 p.
- Peryea, F.J., 1989, Leaching of lead and arsenic in soils contaminated with lead arsenate pesticide residues: Washington State Water Research Center Project No. A-158-WASH, 59 p.
- Phillips, D.J., 1980, Quantitative aquatic biological indicators: Barking, Essex, England, Applied Science Publishers Ltd., 488 p.
- Poplyk, J., ed., 1989, Farm chemicals handbook: Willoughby, Ohio, Meister Publishing Company, v. 75, 676 p.
- Rankama, K., and Sahama, T.G., 1950, Geochemistry: Chicago, Illinois, University of Chicago Press, 911 p.
- Rasmussen, D., 1992., Toxic substances monitoring program 1990 data report: State Water Resources Control Board, California Environmental Protection Agency, 92–1WQ.
- Reif, D., 1989, City of Ellensburg wastewater treatment plant class II inspection, August 1988: Washington State Department of Ecology, Compliance Monitoring Section, Report no. 18-39-04.
- Rinella, J.F., McKenzie, S.W., and Fuhrer, G.J., 1992, Surface-water-quality assessment of the Yakima River Basin, Washington: Analysis of available water-quality data through 1985 water year: U.S. Geological Survey Open-File Report 91–453, 244 p. (pending publication as U.S. Geological Survey Water Supply Paper 2354-B)
- Ryder, J.L., Sanzolone, R.F., Fuhrer, G.J., and Mosier, E.L., 1992, Surface-water-quality assessment of the Yakima River Basin in Washington: Chemical analyses of major, minor, and trace elements in fine-grained streambed sediment: U.S. Geological Survey Open-File Report 92–520, 60 p.
- Ryder, J. L., Sanzolone, R.F., and Porter, S.P., 1993, Surface-water quality assessment of the Kentucky River Basin in Kentucky—Chemical analyses of major, minor, and trace elements in fine-grained streambed sediment: U.S. Geological Survey Open-File Report 93–326–A, 89 p.
- Saiki, M.K., 1987, Relation of length and sex to selenium concentrations in mosquito fish: Environmental Pollution, v. 47, p. 171–186.
- Saiki, M.K., and May, T.W., 1988, Trace element residues in bluegills and common carp from the lower San Joaquin River, California, and its tributaries: The Science of the Total Environment, v. 74, p. 199–217.
- Saiki, M.K., and Palawski, D.U., 1990, Selenium and other elements in juvenile striped bass from the San Joaquin Valley and San Francisco Bay Estuary, California: Archives of Environmental Contamination and Toxicology, v. 19, p. 717–730.
- Salanki, J. V., Balogh, K., and Berta, E., 1982, Heavy metals in animals of Lake Balaton: Water Resources, v. 16, p. 1147–1152.
- Sanzolone, R.F., and Ryder, J.L., 1989, Quality assessment program and results for the NAWQA surface water pilot studies: U.S. Geological Survey Open-File Report 89–658, 22 p.
- Schmitt, C.J., and Brumbaugh, W.G., 1990, National contaminant biomonitoring program: Concentrations of arsenic, cadmium, copper, lead, mercury, selenium, and zinc in United States freshwater fish, 1976–1984: Archives of Environmental Contamination and Toxicology, v. 19, p. 731–747.
- Shacklette, H.T., and Boerngen, J.G., 1984, Element concentrations in soils and other surficial materials of the conterminous United States: U.S. Geological Survey Professional Paper 1134–A, 18 p.
- Simmons, G.C., Van Noy, R.M., and Zilka, N.T., 1983, Mineral resources of the Cougar Lakes—Mount Aix study area, Yakima and Lewis Counties, Washington, Studies related to Wilderness areas: U.S. Geological Survey Bulletin 1504, 81 p., 3 pl.
- Sittig, Marshall, 1981, Handbook of toxic and hazardous chemicals: Park Ridge, New Jersey, Noyes Publications, 729 p.
- Smith, G.A., Bjornstad, B.N., and Fecht, K.R., 1989, Neogene terrestrial sedimentation on and adjacent to the Columbia Plateau; Washington, Oregon, and Idaho: Geological Society of America Special Paper 239, p. 187–198.
- Smock, L.A., 1983, The influence of feeding habits on whole-body metal concentrations in aquatic insects: Freshwater Biology, v. 13, p. 301–321.



- Sorensen, E.M.B. and Bauer, T.L., 1984, A correlation between selenium accumulation in sunfish and changes in condition factor and organ weight: *Environmental Pollution, Series A*, v. 34, p. 357–366.
- Speyer, M.R., 1980, Mercury and selenium concentrations in fish, sediment, and water of two northwestern Quebec lakes: *Bulletin of Environmental Contamination and Toxicology*, v. 24, p. 427–432.
- Tabor, R.W., Waitt, R.B., Frizzell, V.A., Swanson, G.R., Byerly, G.R., and Bentley, R.D., 1982, Geologic map of the Wenatchee 1:100,000 quadrangle, Washington: U.S. Geological Survey Miscellaneous Investigations Series, Map I-1311, 26 p., 1 pl.
- Tanner, D.Q., Sanzolone, R.F., and Zelt, R.B., 1990, Surface water-quality assessment of the Lower Kansas River Basin, Kansas and Nebraska—Concentrations of major metals and trace elements in streambed sediments, 1987: U.S. Geological Survey Open-File Report 90-581, 73 p.
- Timmermans, K.R., Spijkerman, E., Tonkes, M., and Govers, H., 1992, Cadmium and zinc uptake by two species of aquatic invertebrate predators from dietary and aqueous sources: *Canadian Journal of Fisheries and Aquatic Sciences*, v. 45, p. 655–662.
- U.S. Bureau of Reclamation and U.S. Soil Conservation Service, 1974, Inventory of Yakima River Basin, Washington, diversions and return flows, 1973–74: Yakima, Washington, 145 p., 61 pl.
- U.S. Department of Agriculture, 1986, Washington fruit survey, 1986: Olympia, Washington, Washington Agricultural Statistics Service, 73 p.
- U.S. Environmental Protection Agency, 1980a, Ambient water quality criteria for mercury: Washington, D.C., EPA-440/5-80-058.
- \_\_\_\_\_, 1980b, Water quality criteria documents—Availability: *Federal Register*, v. 45, no. 231.
- \_\_\_\_\_, 1986, Quality criteria for water: Washington, D.C., EPA-440/5-86-001.
- \_\_\_\_\_, 1987, Mercury health advisory: Office of Drinking Water, March 1987, 12 p.
- \_\_\_\_\_, 1989, Risk assessment guidance for superfund, Volume I—Human health evaluation Manual (Part A): EPA/540/1-89/002.
- \_\_\_\_\_, 1991, Is your drinking water safe?: Office of Drinking Water, EPA 570/9-91-005, 24 p.
- \_\_\_\_\_, 1992a, Interim guidance on interpretation and implementation of aquatic life criteria for metals: Washington, D.C., Office of Science and Technology, May 1992, 24 p.
- \_\_\_\_\_, 1992b, Drinking water regulations and health advisories: Washington, D.C., Office of Water, December 1992, 11 p.
- \_\_\_\_\_, 1992c, Bulletin Board SQCCALC.ZIP: Office of Science and Technology, 301/589-0205. (Accessed December 1992)
- \_\_\_\_\_, 1992d, Fish sampling and analysis: A guidance document for issuing fish advisories: Office of Science and Technology, Fish Contamination Section, August 1992, draft.
- \_\_\_\_\_, 1993, Drinking water criteria document on arsenic: Human Risk Assessment Branch, Health and Ecological Criteria Division, EPA Contract No. 68-C8-0033, draft.
- U.S. Geological Survey, 1986, Land use and land cover digital data from 1:250,000- and 1:100,000-scale maps: Data user's guide 4, 36 p.
- U.S. Geological Survey and U.S. Bureau of Mines, 1989, Mineral resources of the Alpine Lakes study area and additions, Chelan, King, and Kittitas Counties, Washington: U.S. Geological Survey Bulletin 1542, 317 p., 2 pl.
- Vidal, I.L., 1978, Copper in the livers of trout caught below a sewage discharge: *New Zealand Journal of Marine and Freshwater Research*, v. 12, p. 217–219.
- Waitt, R.B., Jr., 1985, Case for periodic, colossal jokulhlaups from Pleistocene Lake Missoula: *Geological Society of America Bulletin*, v. 96, p. 1271–1286.
- Walsh, D.F., Berger, B.L., and Bean, J.R., 1977, Residues in fish, wildlife, and estuaries—Mercury, arsenic, lead, cadmium, and selenium residues in fish, 1971–1973: *National pesticide monitoring program: Pesticides Monitoring Journal*, v. 11, no. 1, p. 534.
- Walsh, T.J., Korosec, M.A., Phillips, W.M., Logan, R.L., and Schasse, H.W., 1987, Geologic map of Washington, southwest quadrant: Washington State Department of Natural Resources, Division of Geology and Earth Resources, GM-24, 28 p., 1 pl.
- Washington State Administration Code, 1992, Water quality standards for surface waters of the state of Washington: Olympia, Washington State Administration Code, 14 p.
- Washington State Department of Ecology, 1990: City of Sunnyside, National Pollutant Discharge Elimination System Waste Discharge Permit No. WA-002099-1, 5 p.
- Wiener, J.G., Jackson, G.A., May, T.W., and Cole B.P., 1984, Longitudinal distribution of trace elements (As, Cd, Cr, Hg, Pb, and Se) in fishes and sediments in the upper Mississippi River, in Wiener, J.G., Anderson, R.V., McConville, D.R., eds., *Contaminants in the upper mississippi River*: Stoneham, Massachusetts, Butterworth Publishing, p. 139-170.
- Winger, P.V., and Andreasen, J.K., 1985, Contaminant residues in fish and sediment from lakes in the Atchafalaya River Basin (Louisiana): *Archives of Environmental Contamination and Toxicology*, v. 14, p. 579–586.



---

---

## SUPPLEMENTAL DATA TABLES

---

---



**Table 34.** Comparison of selected-element concentrations in streambed-sediment samples of the Yakima River Basin to streambed-sediment samples in the Lower Kansas River Basin, Upper Illinois River Basin, and the Kentucky River Basin

[Concentrations are in micrograms per gram, dry weight, except for iron which is in percent; sample collection, processing, and analytical protocols are identical among river basins (Sanzolone and Ryder, 1989); statistical summaries are based on sampling sites selected randomly using a Universal-Transverse-Mercator, square-grid-sampling procedure; biological-sampling sites in the Yakima River Basin, however, were not selected randomly; lower-order streams are defined as first- or second-order tributaries, with the smallest unbranched-mapped (1:24,000 map scale) tributaries classified as first-order tributaries and streams receiving only first-order tributaries classified as second-order tributaries (Horton, 1945); --, no data; <, less than]

Element	Lower Kansas <sup>1</sup>				Upper Illinois <sup>2</sup>				Kentucky <sup>3</sup>				Yakima			
					Lower-order stream sites				Biological sites <sup>4</sup>				Lower-order stream sites <sup>5</sup>			
	Percentiles				Percentiles				Percentiles				Percentiles			
	10th	50th	90th		10th	50th	90th		10th	50th	90th		10th	50th	90th	
Antimony	0.5	0.7	1.0		0.4	0.7	1.1		0.3	0.5	1.0		0.2	0.4	0.7	
Arsenic	4.5	6.4	11		5.4	9.3	21		4.7	6.7	13		1.5	3.7	11.9	
Beryllium	1	2	2		<1	2	2		1	2	3		1	1	2	
Cadmium	<2	<2	<2		<2	<2	<2		<2	<2	<2		<2	<2	<2	
Chromium	38	45	57		41	56	74		42	62	86		31	62	167	
Cobalt	8	11	21		9	13	20		13	20	32		16	20	30	
Copper	12	15	19		17	23	35		12	20	32		21	30	70	
Iron	1.8	2.2	2.8		2.2	2.9	4.1		2.5	3.5	4.7		4.3	5.1	6.6	
Lead	17	20	29		19	27	53		19	27	42		11	14	32	
Manganese	330	550	1,400		520	680	1,400		680	1,300	3,400		759	1,015	1,470	
Mercury	<.02	<.02	.04		.02	.04	.12		<.02	.04	.06		<.02	.1	.3	
Nickel	14	18	29		17	26	35		18	30	49		17	27	82	
Selenium	--	--	--		.4	.7	1.3		--	--	--		<.4	.4	.9	
Silver	<2	<2	<2		<2	<2	<2		<2	<2	<2		<2	<2	<2	
Zinc	44	56	79		67	100	240		56	91	150		84	100	168	

<sup>1</sup>Tanner and others, 1990.

<sup>2</sup>Colman and Sanzolone, 1991.

<sup>3</sup>Ryder and others, 1993.

<sup>4</sup>Fuhrer, Fluter, and others, 1994.

<sup>5</sup>Ryder and others, 1992.

**Table 35. Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90**

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; concentrations of major elements are reported as percentages, except suspended organic carbon which is reported as milligrams per liter; concentrations of trace elements are reported as micrograms per gram, dry weight; element names shown in bold print are U.S. Environmental Protection Agency (1992c) Priority Pollutants; Cle Elum, Yakima River at Cle Elum; Umtanum, Yakima River at Umtanum; Naches, Naches River near North Yakima; Union Gap, Yakima River above Ahtanum Creek at Union Gap; Sulphur Creek, Sulphur Creek Wasteway near Sunnyside; Grandview, Yakima River at Euclid Bridge at river mile 55 near Grandview; Kiona, Yakima River at Kiona; <, less than]

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
MAJOR ELEMENTS										
Aluminum										
6	Cle Elum	11	7.5	7.6	8.3	8.5	9.0	9.1	9.2	9.2
19	Umtanum	32	5.8	5.8	6.4	6.9	7.4	7.8	8.7	9.7
26	Naches	20	5.2	5.4	6.4	7.5	7.8	8.0	8.6	8.7
32	Union Gap	35	4.6	6.1	6.5	7.0	7.3	7.6	7.7	8.0
50	Kiona	44	6.3	6.4	6.5	6.8	7.0	7.3	7.4	7.4
52	Sulphur Creek	40	5.6	6.3	6.6	7.0	7.2	7.4	7.6	7.6
56	Grandview	37	6.3	6.5	6.7	6.9	7.0	7.2	7.4	7.5
Calcium										
6	Cle Elum	11	1.2	1.3	1.6	1.8	1.9	2.0	2.0	2.0
19	Umtanum	32	1.5	1.6	1.7	1.9	2.0	2.0	2.4	2.9
26	Naches	20	1.4	1.8	1.9	2.3	2.5	2.8	2.9	2.9
32	Union Gap	35	1.3	1.7	1.9	2.0	2.2	2.5	2.5	2.6
50	Kiona	44	1.9	2.0	2.1	2.2	2.2	2.3	2.3	2.4
52	Sulphur Creek	40	2.3	2.5	2.7	2.9	3.0	3.1	3.5	3.5
56	Grandview	37	2.1	2.2	2.2	2.2	2.3	2.4	2.5	2.5
Carbon, suspended organic										
6	Cle Elum	15	<.1	<.1	.1	.3	.5	1.1	1.9	1.9
19	Umtanum	31	.1	.2	.3	.4	.6	1.2	1.8	1.9
26	Naches	22	.2	.2	.3	.4	1.2	2.7	4.1	4.3
32	Union Gap	32	<.1	.1	.3	.6	1.0	3.1	4.8	4.9
50	Kiona	40	.1	.2	.4	.6	1.3	2.4	2.7	2.8
52	Sulphur Creek	36	.2	.6	.7	1.3	1.9	3.0	3.6	3.9
56	Grandview	34	.4	.4	.6	.7	1.0	2.1	3.1	3.6
Iron										
6	Cle Elum	11	4.0	4.1	5.1	5.4	5.8	5.9	5.9	5.9
19	Umtanum	32	4.3	4.5	4.7	4.9	5.4	5.7	5.9	6.3
26	Naches	20	3.8	4.2	4.8	5.0	5.3	5.6	5.8	5.8
32	Union Gap	35	3.9	4.5	4.8	5.0	5.2	5.4	5.6	6.4
50	Kiona	44	4.9	5.1	5.2	5.4	5.6	5.9	6.2	8.1
52	Sulphur Creek	40	4.6	4.7	4.8	5.2	5.4	5.6	5.7	5.9
56	Grandview	37	4.9	5.1	5.3	5.5	5.7	6.0	6.4	7.5

**Table 35.** Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987-90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Magnesium										
6	Cle Elum	11	1.1	1.2	1.7	1.8	1.9	1.9	1.9	1.9
19	Umtanum	32	1.1	1.2	1.3	1.5	1.6	1.9	2.2	2.4
26	Naches	20	.8	.9	.9	1.0	1.2	1.3	1.5	1.5
32	Union Gap	34	1.0	1.1	1.2	1.2	1.3	1.4	1.4	1.4
50	Kiona	44	1.2	1.2	1.3	1.3	1.4	1.4	1.4	1.4
52	Sulphur Creek	40	1.3	1.3	1.4	1.4	1.5	1.5	1.5	1.6
56	Grandview	37	1.2	1.3	1.3	1.4	1.4	1.5	1.5	1.5
Phosphorus										
6	Cle Elum	11	.14	.14	.14	.15	.15	.16	.16	.16
19	Umtanum	32	.11	.11	.13	.15	.16	.19	.20	.21
26	Naches	20	.10	.11	.12	.13	.15	.20	.22	.22
32	Union Gap	35	.12	.13	.14	.15	.19	.22	.32	.34
50	Kiona	44	.14	.15	.16	.18	.20	.22	.23	1.2
52	Sulphur Creek	39	.11	.12	.13	.16	.22	.27	.29	.33
56	Grandview	37	.14	.15	.16	.17	.18	.20	.20	.21
Potassium										
6	Cle Elum	11	.90	.92	1.1	1.1	1.1	1.2	1.2	1.2
19	Umtanum	32	.70	.89	.90	.98	1.0	1.1	1.3	1.4
26	Naches	20	.70	.73	.82	.97	1.1	1.3	1.5	1.6
32	Union Gap	35	.73	.81	.90	1.0	1.2	1.3	1.3	1.4
50	Kiona	44	1.1	1.1	1.2	1.3	1.3	1.4	1.4	1.5
52	Sulphur Creek	40	1.3	1.4	1.5	1.6	1.6	1.7	1.7	2.4
56	Grandview	37	1.1	1.2	1.2	1.3	1.4	1.4	1.5	1.5
Sodium										
6	Cle Elum	11	1.0	1.0	1.2	1.3	1.4	1.5	1.6	1.6
19	Umtanum	32	.8	1.0	1.1	.13	1.4	1.6	2.0	2.4
26	Naches	20	.9	.9	1.1	1.3	1.5	1.7	1.8	1.8
32	Union Gap	35	.8	1.0	1.1	1.3	1.5	1.7	1.7	1.8
50	Kiona	44	1.0	1.1	1.2	1.2	1.3	1.4	1.5	1.6
52	Sulphur Creek	40	.9	1.0	1.2	1.4	1.6	1.7	1.8	1.8
56	Grandview	37	1.0	1.1	1.2	1.3	1.4	1.4	1.6	1.6



**Table 35.** Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Titanium										
6	Cle Elum	11	0.50	0.52	0.60	0.66	0.71	0.76	0.76	0.76
19	Umtanum	32	.45	.48	.51	.54	.58	.61	.71	.83
26	Naches	20	.41	.46	.48	.55	.60	.64	.65	.65
32	Union Gap	35	.37	.48	.51	.54	.58	.63	.67	.67
50	Kiona	44	.57	.58	.59	.61	.63	.68	.71	.74
52	Sulphur Creek	40	.54	.61	.62	.64	.65	.68	.71	.71
56	Grandview	37	.58	.60	.61	.63	.68	.73	.76	.79
TRACE ELEMENTS										
Antimony										
6	Cle Elum	11	.6	.6	.7	.8	2.3	3.0	3.1	3.1
19	Umtanum	32	.3	.4	.5	.6	.7	.8	.8	.9
26	Naches	20	.5	.5	.5	.6	.6	1.0	1.2	1.2
32	Union Gap	35	.4	.5	.5	.5	.6	.7	.9	1.1
50	Kiona	44	.4	.5	.5	.6	.6	.8	.9	1.2
52	Sulphur Creek	40	.5	.6	.6	.7	.8	.8	.9	.9
56	Grandview	37	.3	.4	.5	.6	.6	.7	.8	1.2
Arsenic										
6	Cle Elum	11	6.3	6.3	6.6	7.0	7.9	8.7	8.8	8.8
19	Umtanum	32	2.8	3.9	4.4	5.1	6.0	6.3	7.1	7.4
26	Naches	20	4.4	4.7	5.4	6.3	8.5	12	13	13
32	Union Gap	35	4.2	4.4	5.0	5.3	6.4	7.9	10	11
50	Kiona	44	5.1	5.4	6.0	7.0	7.9	9.6	10	11
52	Sulphur Creek	40	4.9	5.9	6.8	10	14	16	19	20
56	Grandview	37	5.0	5.2	6.0	7.1	8.7	9.3	9.9	11
Beryllium										
6	Cle Elum	11	<2	<2	<2	<2	<2	<2	2	2
19	Umtanum	32	<2	<2	<2	<2	<2	<2	2	2
26	Naches	20	<2	<2	<2	<2	<2	<2	<2	<2
32	Union Gap	35	<2	<2	<2	<2	<2	<2	<2	2
50	Kiona	44	<2	<2	<2	<2	<2	<2	<2	2
52	Sulphur Creek	40	<2	<2	<2	2	2	2	2	3
56	Grandview	37	<2	<2	<2	<2	<2	2	2	2

**Table 35.** Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Cadmium										
6	Cle Elum	11	0.3	0.3	0.6	0.8	2.6	8.0	9.3	9.3
19	Umtanum	32	<.1	2	3	.5	.8	1.7	3	33
26	Naches	20	.1	.1	4	.7	1.4	2.7	4.4	4.5
32	Union Gap	35	.1	2	3	.4	.6	1.0	1.3	1.5
50	Kiona	44	<.1	2	3	.4	.7	2.1	4.6	15
52	Sulphur Creek	40	<.1	2	2	.4	.5	8	1.5	1.7
56	Grandview	37	<.1	2	4	.6	.8	1.3	1.7	2.6
Chromium										
6	Cle Elum	10	110	110	110	110	120	130	140	140
19	Umtanum	29	59	73	81	100	120	150	160	160
26	Naches	19	28	30	35	42	48	65	83	83
32	Union Gap	32	43	53	58	70	79	88	90	90
50	Kiona	38	53	55	56	58	62	85	93	110
52	SulphurCreek	33	41	46	50	53	56	61	72	89
56	Grandview	31	54	56	58	61	66	94	110	110
Cobalt										
6	Cle Elum	11	20	20	23	25	26	28	28	28
19	Umtanum	32	16	19	20	22	25	26	28	28
26	Naches	20	13	15	17	18	20	22	23	23
32	Union Gap	35	16	18	18	19	21	22	23	25
50	Kiona	44	18	18	20	20	22	23	25	31
52	Sulphur Creek	40	17	18	19	21	22	23	24	25
56	Grandview	37	19	19	20	21	22	23	24	28
Copper										
6	Cle Elum	11	47	47	54	60	67	70	70	70
19	Umtanum	32	31	34	39	43	54	61	67	73
26	Naches	20	26	28	40	46	54	75	78	78
32	Union Gap	35	33	40	42	49	65	93	120	150
50	Kiona	44	32	38	39	43	53	120	440	680
52	Sulphur Creek	40	21	27	31	44	63	110	80	210
56	Grandview	37	29	32	37	40	44	52	56	74

**Table 35.** Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987-90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Lead										
6	Cle Elum	11	12	12	12	17	24	110	120	120
19	Umtanum	32	6	10	12	15	20	26	160	410
26	Naches	20	10	11	14	18	23	26	45	46
32	Union Gap	35	11	12	15	18	25	29	31	33
50	Kiona	44	10	15	17	20	22	24	25	28
52	Sulphur Creek	40	13	17	20	24	26	28	41	43
56	Grandview	37	11	14	16	19	24	26	29	31
Manganese										
6	Cle Elum	11	1,400	1,400	1,500	1,600	2,000	2,500	2,600	2,600
19	Umtanum	32	1,100	1,200	1,400	1,700	2,200	2,600	3,200	3,700
26	Naches	20	1,200	1,300	1,300	1500	2,100	3,300	3,600	3,600
32	Union Gap	35	1,100	1,200	1,300	1,600	2,300	2,900	3,000	3,100
50	Kiona	44	1,200	1,600	1,700	2,600	3,100	3,900	4,100	4,600
52	Sulphur Creek	40	910	950	1,000	1,500	2,800	4,100	5,200	5,400
56	Grandview	37	1,200	1,400	2,000	2,900	3,300	4000	4,500	6,300
Molybdenum										
6	Cle Elum	11	<.1	2	.7	.7	9	1.2	1.3	1.3
19	Umtanum	32	<.1	2	5	.6	.6	.7	.8	.9
26	Naches	20	<.1	.6	.7	.8	1.0	1.1	1.2	1.2
32	Union Gap	35	<.1	3	.6	.7	.8	1.2	1.6	3.0
50	Kiona	44	<.1	<1	.6	.6	.7	.8	.8	1.0
52	Sulphur Creek	40	<.1	<1	.6	.7	1.3	1.7	1.7	2.0
56	Grandview	37	<.1	<1	<1	.6	.8	.8	1.0	1.8
Nickel										
6	Cle Elum	10	62	62	66	72	76	120	130	130
19	Umtanum	29	43	61	69	89	100	140	160	170
26	Naches	19	12	14	16	17	21	30	41	41
32	Union Gap	32	26	31	38	46	56	62	66	71
50	Kiona	38	28	30	33	36	39	44	49	52
52	Sulphur Creek	33	18	23	24	27	30	33	33	34
56	Grandview	31	28	29	34	37	38	41	42	43

**Table 35.** Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Silver										
6	Cle Elum	11	0.2	0.2	0.3	0.4	0.5	0.8	0.9	0.9
19	Umtanum	32	.2	.2	.2	.2	.3	.7	3.8	7.7
26	Naches	20	<.1	.1	.2	.2	.3	.4	.6	.6
32	Union Gap	35	.2	.3	.3	.5	1.0	1.4	1.7	2.0
50	Kiona	44	.2	.3	.4	.4	.5	.6	.8	.8
52	Sulphur Creek	40	.2	.2	.3	.4	.8	1.4	2.4	2.5
56	Grandview	37	.2	.2	.4	.4	.5	.6	.6	.8
Thallium										
6	Cle Elum	11	.3	.3	.3	.3	.4	.5	.5	.5
19	Umtanum	32	.2	.2	.2	.3	.3	.3	.4	.4
26	Naches	20	.2	.2	.2	.3	.4	.4	.5	.5
32	Union Gap	35	.1	.2	.3	.3	.4	.4	.4	.4
50	Kiona	44	.2	.3	.3	.4	.4	.4	.5	.5
52	Sulphur Creek	40	.3	.4	.4	.4	.5	.5	.6	.6
56	Grandview	37	.3	.3	.3	.4	.4	.5	.5	.6
Vanadium										
6	Cle Elum	11	120	120	140	160	170	170	170	170
19	Umtanum	32	110	120	120	130	140	150	160	170
26	Naches	20	110	110	120	130	140	160	170	170
32	Union Gap	35	100	120	120	130	140	140	150	150
50	Kiona	44	130	130	140	140	140	150	160	190
52	Sulphur Creek	40	140	140	140	150	160	170	170	170
56	Grandview	37	130	140	140	150	150	160	180	190
Zinc										
6	Cle Elum	10	140	140	160	190	230	490	520	520
19	Umtanum	29	95	98	130	140	180	250	380	380
26	Naches	19	91	110	120	120	140	170	180	180
32	Union Gap	32	110	120	140	150	170	220	250	300
50	Kiona	38	100	120	130	150	170	180	260	350
52	Sulphur Creek	33	88	97	110	140	180	200	210	220
56	Grandview	31	110	110	120	130	150	180	190	200

**Table 35.** Distribution of major- and trace-element concentrations in suspended sediment at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
OTHER MEASUREMENTS										
Discharge, in cubic feet per second										
6	Cle Elum	17	442	626	1,100	1,440	2,550	3,610	3,610	3,610
19	Umtanum	35	454	907	1,230	1,910	3,450	3,850	4,290	5,600
26	Naches	23	285	453	680	1,470	2,500	3,830	4,720	4,910
32	Union Gap	36	678	1,320	1,870	2,980	3,580	7,280	8,920	8,940
50	Kiona	44	1,040	1,200	1,480	2,325	3,300	5,095	6,360	7,170
52	Sulphur Creek	40	57	66	73	205	255	348	384	422
56	Grandview	37	1,100	1,230	1,500	2,270	2,940	4,380	4,800	4,940
Surface area, in square meters per gram										
6	Cle Elum	16	6.40	7.04	8.72	11.1	13.2	19.8	26.5	26.5
19	Umtanum	34	4.32	9.90	13.7	17.2	20.9	32.7	34.0	35.1
26	Naches	23	5.82	10.1	12.1	18.9	24.3	31.2	32.3	32.4
32	Union Gap	35	8.14	13.0	15.3	17.6	22.8	27.5	32.5	48.5
50	Kiona	44	13.6	15.3	17.7	21.6	27.0	36.0	40.7	63.7
52	Sulphur Creek	40	5.09	6.84	9.58	14.3	19.8	23.4	25.4	32.1
56	Grandview	37	1.11	13.8	17.4	23.5	28.7	34.2	39.3	57.0
Streambed sediment finer than 62 micrometers, in percent										
6	Cle Elum	17	45	47	71	91	95	99	100	100
19	Umtanum	34	34	62	76	84	89	95	96	96
26	Naches	23	56	63	74	87	92	96	98	98
32	Union Gap	34	27	64	82	90	93	96	96	98
50	Kiona	39	60	86	90	93	95	97	97	99
52	Sulphur Creek	40	39	60	70	83	92	94	96	96
56	Grandview	37	19	68	90	94	95	97	98	99
Suspended sediment, in milligrams per liter										
6	Cle Elum	17	2	2	3	6	20	94	130	130
19	Umtanum	34	2	3	6	17	28	88	110	110
26	Naches	23	2	3	5	13	27	78	120	130
32	Union Gap	36	4	7	12	18	43	140	300	1100
50	Kiona	42	4	8	15	25	47	120	150	190
52	Sulphur Creek	40	7	13	32	80	142	220	370	620
56	Grandview	37	7	10	16	25	40	130	220	350

**Table 36. Comparison of selected element concentrations in filtered-water samples from surface waters of the Yakima River Basin to surface waters in the United States**

[Concentrations are in micrograms per liter; NASQAN, National Stream Quality Accounting Network based on data from 300 sites; --, no data; <, less than]

Element	Background concentrations, inland water <sup>1</sup>	North American streams <sup>2</sup>	NASQAN 1974–81 <sup>3</sup>			Yakima River Basin (1953–85 water years) <sup>4</sup>					Yakima River Basin (1987–90 water years) <sup>5</sup>				
			Percentiles			Number of samples	Percentiles				Number of samples	Percentiles			
			25th	50th	75th		25th	50th	75th	90th		25th	50th	75th	90th
Aluminum	<30	--	--	---	--	127	10	20	50	90	48	<10	10	20	40
Antimony	.1	--	--	---	--	24	0	0	0	0	22	<1	<1	<1	<1
Arsenic	2	<10	<1	1	3	218	<5	<5	5	5	119	<1	<1	2	3
Beryllium	.01	<.3	--	--	--	69	<.5	<.5	<1	<1	58	<.5	<.5	<.5	<.5
Cadmium	.07	1	<2	<2	<2	133	<1	<1	<1	2	279	<2	<2	<2	.3
Chromium	.5	5.8	9	10	10	387	0	0	10	10	26	<.5	<.5	.6	1.0
Cobalt	.05	<1	---	---	--	132	<3	<3	<3	<3	58	<3	<3	<3	<3
Copper	1.8	--	--	--	--	491	2	3	10	20	280	.6	.9	1.3	1.9
Iron	<30	--	36	63	157	180	30	48	100	200	58	18	28	40	56
Lead	.2	--	3	4	6	372	<2	4	10	18	279	<.5	<.5	<.5	<.5
Manga-	<5	--	11	24	51	165	<10	10	20	30	58	4	10	16	33
Mercury	.01	( <sup>6</sup> )	.2	.2	.3	167	<.1	<.1	.2	.5	283	<.1	<.1	<.1	<.1
Nickel	.3	10	--	--	--	139	1	1	2	4	<10	<10	<10	<10	<10
Selenium	.1	.2	<1	<1	1	133	0	0	<1	<1	44	<1	<1	<1	<1
Silver	.3	.3	--	--	--	121	<1	<1	1	1	58	<1	<1	<1	1
Zinc	10	20	12	15	21	465	<3	11	20	50	58	<3	4	8	18

<sup>1</sup>Minimally contaminated inland waters (Forstner and Wittmann, 1979).

<sup>2</sup>Hem, 1989

<sup>3</sup>Percentiles are calculated from site-mean concentrations (Smith and others, 1987).

<sup>4</sup>Rinella and others, 1992

<sup>5</sup>Fuhrer, Fluter, and others, 1994

<sup>6</sup>Mercury concentrations rarely exceed a few tenths of a microgram per liter (Hem, 1989).

**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90

[To avoid statistical bias that may be associated with constituents analyzed in duplicate or triplicate at a site, only one element concentration per visit was statistically summarized; all concentrations listed below are in micrograms per liter except bromide, organic carbon, and hardness, which are given in milligrams per liter; element names shown in bold print are U.S. Environmental Protection Agency (1992c) Priority Pollutants; for cadmium, copper, and lead, only samples analyzed by atomic absorption spectroscopy with graphite furnace were statistically summarized below; <, less than; Cle Elum, Yakima River at Cle Elum; Umtanum, Yakima River at Umtanum; Naches, Naches River near North Yakima; Union Gap, Yakima River above Ahtanum Creek at Union Gap; Sulphur Creek, Sulphur Creek Wasteway near Sunnyside; Grandview, Yakima River at Euclid Bridge at river mile 55 near Grandview; Kiona, Yakima River at Kiona]

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
MAJOR ELEMENTS										
Aluminum										
6	Cle Elum	1	10	10	10	10	10	10	10	10
19	Umtanum	2	20	20	20	120	210	210	210	210
26	Naches	3	20	20	20	20	20	20	20	20
32	Union Gap	17	<10	<10	<10	20	30	60	60	60
50	Kiona	15	<10	<10	<10	<10	20	40	50	50
52	Sulphur Creek	6	<10	<10	<10	<10	<10	20	20	20
56	Grandview	4	<10	<10	<10	<10	20	20	20	20
Carbon, Organic										
6	Cle Elum	38	.80	.90	1.2	1.3	1.7	2.2	2.6	2.8
19	Umtanum	40	.40	1.2	1.6	2.0	2.4	3.3	3.7	4.5
26	Naches	38	.60	1.0	1.2	1.4	2.2	3.1	3.4	4.3
32	Union Gap	38	.80	1.3	1.6	2.0	2.7	3.2	4.1	5.8
50	Kiona	41	1.1	1.3	1.9	2.3	2.8	3.6	6.5	7.5
52	Sulphur Creek	43	1.5	1.8	2.2	2.7	3.6	4.7	5.5	8.0
56	Grandview	37	1.0	1.2	1.6	2.0	2.6	4.4	5.7	6.4
Hardness										
6	Cle Elum	41	20	21	22	26	28	32	35	36
19	Umtanum	40	33	35	41	46	57	65	68	70
26	Naches	39	18	21	28	32	37	46	48	57
32	Union Gap	43	28	33	37	44	55	66	79	87
50	Kiona	43	43	53	72	96	110	120	130	140
52	Sulphur Creek	46	70	86	98	130	260	260	270	270
56	Grandview	40	49	60	69	89	100	110	110	120



**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Iron										
6	Cle Elum	5	8	8	9	13	36	44	44	44
19	Umtanum	6	19	19	20	23	100	250	250	250
26	Naches	3	13	13	13	18	35	35	35	35
32	Union Gap	17	5	20	28	36	54	67	67	67
50	Kiona	16	8	12	16	24	36	48	53	53
52	Sulphur Creek	7	16	16	18	22	30	33	33	33
56	Grandview	4	20	20	21	27	64	75	75	75
TRACE ELEMENTS										
Antimony										
6	Cle Elum	4	<1	<1	<1	<1	<1	<1	<1	<1
19	Umtanum	3	<1	<1	<1	<1	1	1	1	1
26	Naches	2	<1	<1	<1	<1	<1	<1	<1	<1
32	Union Gap	4	<1	<1	<1	<1	<1	1	1	1
50	Kiona	4	<1	<1	<1	<1	<1	<1	<1	<1
52	Sulphur Creek	3	<1	<1	<1	<1	<1	<1	<1	<1
56	Grandview	2	<1	<1	<1	<1	<1	<1	<1	<1
Arsenic										
6	Cle Elum	16	<1	<1	<1	<1	<1	<1	<1	<1
19	Umtanum	11	<1	<1	<1	<1	<1	<1	<1	<1
26	Naches	15	<1	<1	<1	<1	<1	<1	1	1
32	Union Gap	23	<1	<1	<1	<1	<1	<1	1	1
50	Kiona	25	<1	<1	1	1	2	3	4	4
52	Sulphur Creek	15	2	2	2	3	7	8	9	9
56	Grandview	14	<1	<1	<1	1	2	2	3	3
Barium										
6	Cle Elum	5	2	2	2	3	4	4	4	4
19	Umtanum	6	7	7	8	10	14	16	16	16
26	Naches	3	<2	<2	<2	4	6	6	6	6
32	Union Gap	17	5	6	8	9	13	27	53	53
50	Kiona	16	7	9	17	20	26	32	40	40
52	Sulphur Creek	7	29	29	36	39	73	79	79	79
56	Grandview	4	11	11	14	24	26	27	27	27

**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Beryllium										
6	Cle Elum	5	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
19	Umtanum	6	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
26	Naches	3	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
32	Union Gap	17	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
50	Kiona	16	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
52	Sulphur Creek	7	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
56	Grandview	4	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
Boron										
6	Cle Elum	1	<10	<10	<10	<10	<10	<10	<10	<10
19	Umtanum	2	10	10	10	20	20	20	20	20
26	Naches	3	10	10	10	10	20	20	20	20
32	Union Gap	5	<10	<10	<10	<10	20	20	20	20
50	Kiona	3	20	20	20	20	30	30	30	30
52	Sulphur Creek	6	20	20	20	20	40	40	40	40
56	Grandview	3	10	10	10	20	20	20	20	20
Bromide										
6	Cle Elum	3	<.01	<.01	<.01	<.01	<.01	<.01	<.01	<.01
19	Umtanum	3	<.01	<.01	<.01	<.01	<.01	<.01	<.01	<.01
26	Naches	2	<.01	<.01	<.01	<.01	<.01	<.01	<.01	<.01
32	Union Gap	3	<.01	<.01	<.01	<.01	<.01	<.01	<.01	<.01
50	Kiona	3	<.01	<.01	<.01	<.01	.02	.02	.02	.02
52	Sulphur Creek	3	<.01	<.01	<.01	.02	.08	.08	.08	.08
56	Grandview	2	<.01	<.01	<.01	.01	.02	.02	.02	.02
Cadmium										
6	Cle Elum	42	<.2	<.2	<.2	<.2	.2	.4	.7	1.5
19	Umtanum	40	<.2	<.2	<.2	<.2	.3	1.0	1.2	2.1
26	Naches	38	<.2	<.2	<.2	<.2	<.2	.3	.4	2.2
32	Union Gap	40	<.2	<.2	<.2	<.2	<.2	.2	.2	.5
50	Kiona	38	<.2	<.2	<.2	<.2	<.2	.2	.2	.6
52	Sulphur Creek	43	<.2	<.2	<.2	<.2	<.2	.3	.6	.6
56	Grandview	38	<.2	<.2	<.2	<.2	<.2	.2	.4	1.0

**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Chromium										
6	Cle Elum	3	<.5	<.5	<.5	<.5	1.1	1.1	1.1	1.1
19	Umtanum	4	<.5	<.5	<.5	<.5	.5	.6	.6	.6
26	Naches	2	<.5	<.5	<.5	<.5	.7	.7	.7	.7
32	Union Gap	4	<.5	<.5	<.5	<.5	.5	.6	.6	.6
50	Kiona	6	<.5	<.5	<.5	<.5	1.0	1.0	1.0	1.0
52	Sulphur Creek	4	<.5	<.5	<.5	<.5	.7	.8	.8	.8
56	Grandview	3	<.5	<.5	<.5	<.5	.6	.6	.6	.6
Cobalt										
6	Cle Elum	5	<3	<3	<3	<3	<3	<3	<3	<3
19	Umtanum	6	<3	<3	<3	<3	<3	<3	<3	<3
26	Naches	3	<3	<3	<3	<3	<3	<3	<3	<3
32	Union Gap	17	<3	<3	<3	<3	<3	<3	<3	<3
50	Kiona	16	<3	<3	<3	<3	<3	<3	<3	<3
52	Sulphur Creek	7	<3	<3	<3	<3	<3	<3	<3	<3
56	Grandview	4	<3	<3	<3	<3	<3	<3	<3	<3
Copper										
6	Cle Elum	42	<.5	<.5	<.5	.8	1.4	4.6	7.2	14
19	Umtanum	39	<.5	.5	.8	1.3	1.9	4.1	7.0	20
26	Naches	38	<.5	<.5	<.5	.6	.8	1.0	1.4	1.5
32	Union Gap	41	<.5	<.5	.6	.9	1.2	1.9	3.1	3.6
50	Kiona	40	<.5	<.5	.7	1.0	1.3	2.8	5.1	5.5
52	Sulphur Creek	42	<.5	.7	.9	1.1	1.3	1.7	2.3	2.5
56	Grandview	38	<.5	<.5	.7	.9	1.3	1.5	1.8	2.9
Lead										
6	Cle Elum	42	<.5	<.5	<.5	<.5	<.5	<.5	<.5	<.5
19	Umtanum	39	<.5	<.5	<.5	<.5	<.5	.6	.9	1.8
26	Naches	37	<.5	<.5	<.5	<.5	<.5	.6	.7	.7
32	Union Gap	41	<.5	<.5	<.5	<.5	<.5	<.5	<.5	.8
50	Kiona	39	<.5	<.5	<.5	<.5	<.5	.6	.7	.8
52	Sulphur Creek	43	<.5	<.5	<.5	<.5	<.5	<.5	1.0	1.2
56	Grandview	38	<.5	<.5	<.5	<.5	<.5	.6	1.1	1.9

**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Lithium										
6	Cle Elum	5	<4	<4	<4	<4	4	5	5	5
19	Umtanum	6	<4	<4	<4	<4	6	16	16	16
26	Naches	3	<4	<4	<4	<4	<4	<4	<4	<4
32	Union Gap	17	<4	<4	<4	<4	<4	<4	<4	<4
50	Kiona	16	<4	<4	<4	<4	<4	4	5	5
52	Sulphur Creek	7	<4	<4	<4	<4	8	9	9	9
56	Grandview	4	<4	<4	<4	<4	<4	<4	<4	<4
Manganese										
6	Cle Elum	5	2	2	2	2	5	7	7	7
19	Umtanum	6	4	4	4	6	17	21	21	21
26	Naches	3	2	2	2	3	4	4	4	4
32	Union Gap	17	6	7	8	10	14	21	31	31
50	Kiona	16	<1	2	2	10	12	24	34	34
52	Sulphur Creek	7	13	13	13	18	86	110	110	110
56	Grandview	4	10	10	16	36	49	53	53	53
Mercury										
6	Cle Elum	42	<.1	<.1	<.1	<.1	<.1	<.1	.2	.6
19	Umtanum	40	<.1	<.1	<.1	<.1	<.1	<.1	.2	.6
26	Naches	38	<.1	<.1	<.1	<.1	<.1	<.1	<.1	.2
32	Union Gap	42	<.1	<.1	<.1	<.1	<.1	<.1	.2	.2
50	Kiona	43	<.1	<.1	<.1	<.1	<.1	<.1	.2	.3
52	Sulphur Creek	41	<.1	<.1	<.1	<.1	<.1	<.1	.1	.3
56	Grandview	37	<.1	<.1	<.1	<.1	<.1	<.1	<.1	.1
Molybdenum										
6	Cle Elum	5	<10	<10	<10	<10	<10	<10	<10	<10
19	Umtanum	6	<10	<10	<10	<10	<10	<10	<10	<10
26	Naches	3	<10	<10	<10	<10	<10	<10	<10	<10
32	Union Gap	17	<10	<10	<10	<10	<10	<10	<10	<10
50	Kiona	16	<10	<10	<10	<10	<10	<10	<10	<10
52	Sulphur Creek	7	<10	<10	<10	<10	<10	<10	<10	<10
56	Grandview	4	<10	<10	<10	<10	<10	<10	<10	<10

**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Nickel										
6	Cle Elum	5	<10	<10	<10	<10	<10	<10	<10	<10
19	Umtanum	6	<10	<10	<10	<10	<10	<10	<10	<10
26	Naches	3	<10	<10	<10	<10	<10	<10	<10	<10
32	Union Gap	17	<10	<10	<10	<10	<10	<10	<10	<10
50	Kiona	16	<10	<10	<10	<10	<10	<10	<10	<10
52	Sulphur Creek	7	<10	<10	<10	<10	<10	<10	<10	<10
56	Grandview	4	<10	<10	<10	<10	<10	<10	<10	<10
Selenium										
6	Cle Elum	4	<1	<1	<1	<1	<1	<1	<1	<1
19	Umtanum	3	<1	<1	<1	<1	<1	<1	<1	<1
26	Naches	2	<1	<1	<1	<1	<1	<1	<1	<1
32	Union Gap	15	<1	<1	<1	<1	<1	<1	<1	<1
50	Kiona	15	<1	<1	<1	<1	<1	<1	1	1
52	Sulphur Creek	3	<1	<1	<1	1	2	2	2	2
56	Grandview	2	<1	<1	<1	<1	<1	<1	<1	<1
Silver										
6	Cle Elum	5	<1	<1	<1	<1	<1	<1	<1	<1
19	Umtanum	6	<1	<1	<1	<1	1	1	1	1
26	Naches	3	<1	<1	<1	2	2	2	2	2
32	Union Gap	17	<1	<1	<1	<1	<1	1	1	1
50	Kiona	16	<1	<1	<1	<1	<1	<1	1	1
52	Sulphur Creek	7	<1	<1	<1	<1	1	1	1	1
56	Grandview	4	<1	<1	<1	<1	2	2	2	2
Strontium										
6	Cle Elum	5	23	23	26	30	36	41	41	41
19	Umtanum	6	42	42	45	60	70	74	74	74
26	Naches	3	21	21	21	53	65	65	65	65
32	Union Gap	17	31	34	44	50	65	92	96	96
50	Kiona	16	49	65	84	110	140	150	150	150
52	Sulphur Creek	7	110	110	140	150	330	330	330	330
56	Grandview	4	71	71	86	130	140	140	140	140

**Table 37.** Distribution of major- and trace-element concentrations in filtered-water samples at fixed sites, Yakima River Basin, Washington, 1987–90—Continued

Site reference number	Site name	Number of samples	Minimum value	Value at indicated percentile						Maximum value
				10	25	50	75	90	95	
Vanadium										
6	Cle Elum	5	<6	<6	<6	<6	<6	<6	<6	<6
19	Umtanum	6	<6	<6	<6	<6	<6	<6	<6	<6
26	Naches	3	<6	<6	<6	<6	<6	<6	<6	<6
32	Union Gap	17	<6	<6	<6	<6	<6	<6	<6	<6
50	Kiona	16	<6	<6	<6	<6	10	10	10	10
52	Sulphur Creek	7	10	10	10	10	20	20	20	20
56	Grandview	4	<6	<6	<6	10	10	10	10	10
Zinc										
6	Cle Elum	5	4	4	6	10	18	19	19	19
19	Umtanum	6	<3	<3	<3	4	9	12	12	12
26	Naches	3	<3	<3	<3	<3	18	18	18	18
32	Union Gap	17	<3	<3	<3	5	8	32	40	40
50	Kiona	16	<3	<3	<3	<3	7	11	12	12
52	Sulphur Creek	7	<3	<3	<3	4	5	9	9	9
56	Grandview	4	<3	<3	<3	10	26	29	29	29

**Table 38.** Comparison of arsenic, mercury, and selenium concentrations in freshwater fish collected in 1984 for the National Contaminant Biomonitoring Program to concentrations in whole-body sculpin collected from the Yakima River Basin, Washington, 1990

[NCBP, National Contaminant Biomonitoring Program (Schmitt and Brumbaugh, 1990); concentrations are reported as micrograms per gram, dry weight; concentrations were converted to dry weight assuming a moisture content of 75 percent; statistics for the Yakima River Basin are based on site means in 1990, not individual sample concentrations; arsenic concentrations below the limit of determination were statistically summarized as one-half their value; sample species: sculpin (*Cottus* spp.)]

Element	Yakima River Basin				NCBP		
	Minimum	Mean	85th percentile	Maximum	Mean	85th percentile	Maximum
Arsenic	<0.10	0.16	0.31	0.45	0.56	4.3	6.0
Mercury	.09	20	.19	.31	.40	.68	1.4
Selenium	.20	2.1	5.3	5.4	1.7	2.9	9.2



**Table 39.** Comparison of selected trace-element concentrations in *Corbicula* species collected from uncontaminated or minimally contaminated aquatic environments in other basins to Asiatic clams collected from the Yakima River Basin, Washington, 1990

[Statistics for the Yakima River Basin are based on site means for 1990 data, not individual sample concentrations; concentrations are reported in micrograms per gram, dry weight; reference basins are: (1) Apalachicola River [Elder and Matraw, 1984], (2) San Francisco Bay and Sacramento/San Joaquin River Delta [Johns and Luoma, 1990], (3) San Francisco Bay, Sacramento-San Joaquin Delta, and selected tributaries [Johns and others, 1988], (4) San Joaquin River [Leland and others, 1988], (5) San Joaquin River [Leland and Scudder, 1990], (6) Suisun Bay/Delta in San Francisco Bay [Luoma and others, 1990], (7) selected river basins in the State of California [McCleneghan and others, 1981]; data cited from McCleneghan and others (1981) were converted from fresh-weight concentrations using the reported values for percent moisture; Yakima River Basin sample species: Asiatic clam (Veneroidae: Corbiculidae *Corbicula fluminea*)]

Element	Yakima River Basin			Other Basins	
	Minimum	Median	Maximum	Site mean	Reference basin
Arsenic	3.6	4.2	4.6	<0.1–0.36	1
				6.9–7.2	2
				5.3–7.7	4
				0.2–1.4	6
Cadmium	.23	.31	.39	<0.1–0.25	1
				0.24–0.48	4
				<1	5
Chromium	1.2	1.3	2.1	<0.1–2.0	1
				<2	5
				0.8	6
Copper	25	29	35	1.0–13	1
				16–66	4
				20–40	5
				48	3, 6
Lead	.18	.31	.40	<0.1–0.2	1
				1.6	5
Mercury	.10	.16	.17	0.086–0.184	3
				0.11–0.13	4
Nickel	.88	1.1	1.3	0.75–1.35	4
Selenium	2.0	2.6	3.0	2.8–3.1	3
				1.26–2.02	4
Zinc	98	107	452	2.1–26	1
				120–177	5
				167	6

**Table 40.** Comparison of selected trace-element concentrations in benthic insects collected from uncontaminated or minimally contaminated aquatic environments in other basins to benthic insects collected from the Yakima River Basin, Washington, 1990

[Statistics for the Yakima River Basin are based on site means for 1990 data, not individual-sample concentrations; concentrations are reported in micrograms per gram, dry weight; N, number of taxa measured at all sites; reference basins are: (1) cobble-bottom rivers and streams [Cain and others, 1992] (2) spring-fed stream [Elwood and others, 1976], (3) mine-drainage streams [Gower and Darlington, 1990], (4) Red River, New Mexico [Lynch and others, 1988], (5) metal-contaminated sites [Miller and others, 1992], (6) Uncontaminated streams [Smock, 1983]]

Element	Yakima River Basin				Other Basins	
	N	Minimum	Median	Maximum	Site mean	Reference basin
Cadmium	87	0.11	0.12	0.58	0.1–0.2	1
					1.9	4
Chromium	92	.31	1.7	33	14.2	2
					4.9	4
					7.4	6
Cobalt	94	.20	1.0	9.1	2.9	2
					6.2	6
Copper	94	5.6	16	39	11–34	1
					29	3
					43	4
					57–99	5
Lead	85	.06	.73	24	0.1–1.8	1
					0.5	4
Nickel	90	.12	1.2	77	7.1	4
Zinc	94	67	138	450	11–34	1
					29	3
					43	4
					57–99	5



---

---

## APPENDIX

---

---



## APPENDIX

Equations for calculating estimated loads for arsenic, cadmium, and copper at fixed sites in the Yakima River Basin, Washington, 1987–90

[Equations derived using model in Cohn & others (1992); ln, natural logarithm; L, load, in grams per day; Q, flow, in cubic feet per second; T, time, in decimal years; sin, sine; cos, cosine]

Element	Sample medium	Equation
<b>Yakima River at Cle Elum</b>		
Cadmium	Filtered water	$\ln(L) = 0.9544\ln(Q) - 0.8720$
Copper	Filtered water	$\ln(L) = 1.7504\ln(Q) - 0.1496\sin(2\pi T) + 0.9285\cos(2\pi T) - 4.547$
<b>Yakima River at Umtanum</b>		
Arsenic	Suspended sediment	$\ln(L) = 1.8952\ln(Q) + 0.2113\sin(2\pi T) + 0.3696\cos(2\pi T) - 8.252$
Cadmium	Suspended sediment	$\ln(L) = 1.5532\ln(Q) + 0.9503\sin(2\pi T) + 0.2455\cos(2\pi T) - 7.558$
Cadmium	Filtered water	$\ln(L) = 1.3087\ln(Q) - 3.532$
Copper	Suspended sediment	$\ln(L) = 1.5635\ln(Q) - 3.554$
Copper	Filtered water	$\ln(L) = 1.5360\ln(Q) - 0.3383\sin(2\pi T) + 0.5223\cos(2\pi T) - 2.866$
<b>Naches River near North Yakima</b>		
Arsenic	Suspended sediment	$\ln(L) = 2.3185\ln(Q) + 0.0774\sin(2\pi T) + 0.6033\cos(2\pi T) - 10.916$
Cadmium	Suspended sediment	$\ln(L) = 2.1101\ln(Q) + 0.7819\sin(2\pi T) + 0.7520\cos(2\pi T) - 11.819$
Copper	Suspended sediment	$\ln(L) = 2.1312\ln(Q) - 7.709$
Copper	Filtered water	$\ln(L) = 1.1817\ln(Q) - 0.8732$
<b>Yakima River above Ahtanum Creek at Union Gap</b>		
Arsenic	Suspended sediment	$\ln(L) = 1.4355\ln(Q) - 11.73$
Cadmium	Suspended sediment	$\ln(L) = 0.9314\ln(Q) + 0.6929\sin(2\pi T) + 0.2841\cos(2\pi T) - 2.923$
Cadmium	Filtered water	$\ln(L) = 1.6426\ln(Q) - 7.203$
Copper	Suspended sediment	$\ln(L) = 1.3215\ln(Q) - 1.346$
Copper	Filtered water	$\ln(L) = 0.9334\ln(Q) + 0.1263\sin(2\pi T) - 0.0300\cos(2\pi T) + 1.276$
<b>Yakima River at Euclid Bridge at river mile 55 near Grandview</b>		
Arsenic	Suspended sediment	$\ln(L) = 2.099\ln(Q) - 0.3149\sin(2\pi T) - 0.6459\cos(2\pi T) - 9.219$
Arsenic	Filtered water	$\ln(L) = 0.0884\ln(Q) + 8.258$
Cadmium	Suspended sediment	$\ln(L) = 1.4354\ln(Q) + 0.3762\sin(2\pi T) - 0.3698\cos(2\pi T) - 6.577$
Copper	Suspended sediment	$\ln(L) = 2.4962\ln(Q) - 0.2799\sin(2\pi T) - 0.6514\cos(2\pi T) - 10.474$
Copper	Filtered water	$\ln(L) = 0.7494\ln(Q) + 2.737$
<b>Yakima River at Kiona</b>		
Arsenic	Filtered water	$\ln(L) = 0.5026\ln(Q) + 5.039$
Cadmium	Suspended sediment	$\ln(L) = 2.4629\ln(Q) + 0.1347\sin(2\pi T) - 0.7258\cos(2\pi T) - 14.994$
Copper	Suspended sediment	$\ln(L) = 2.9597\ln(Q) - 0.5866\sin(2\pi T) - 1.1695\cos(2\pi T) - 14.160$
<b>Sulphur Creek Wasteway near Sunnyside</b>		
Arsenic	Suspended sediment	$\ln(L) = 1.4907\ln(Q) + 0.5104\sin(2\pi T) - 0.1757\cos(2\pi T) - 2.004$
Arsenic	Filtered water	$\ln(L) = 0.1476\ln(Q) + 6.5755$
Cadmium	Suspended sediment	$\ln(L) = 1.0724\ln(Q) + 0.6953\sin(2\pi T) - 0.4716\cos(2\pi T) - 3.166$
Copper	Suspended sediment	$\ln(L) = 1.0648\ln(Q) + 0.6039\sin(2\pi T) - 0.5476\cos(2\pi T) + 1.751$
Copper	Filtered water	$\ln(L) = 0.9674\ln(Q) + 1.118$



## GLOSSARY OF SELECTED TERMS

- alluvium.** A deposit of clay, silt, sand, gravel, mud, or similar unconsolidated detrital material formed by flowing water.
- amphibolite.** A crystalloblastic (metamorphic) rock consisting mainly of amphibole and plagioclase with little or no quartz.
- anadromous.** Migrating from the sea up a river to spawn.
- andesite.** A dark-colored, fine-grained volcanic rock composed essentially of plagioclase feldspar, resembling trachyte in appearance; the extrusive equivalent of diorite.
- anthropogenic.** Of or pertaining, or resulting from to human activities.
- arsenopyrite.** A tin-white or silver-white to steel-gray orthorhombic mineral:  $\text{FeAsS}$ ; it constitutes the principal ore of arsenic.
- asphalt.** A dark brown to black viscous liquid or low-melting solid bitumen formed in oil-bearing rocks by the evaporation of the volatiles.
- basalt.** The dark, dense igneous rock of a lava flow or minor intrusion, composed essentially of labradorite and pyroxene and often displaying a columnar structure.
- batholith.** A large body of igneous rock, bounded by irregular, cross-cutting surfaces or fault planes, and believed to have crystallized at a considerable depth below the earth's surface.
- benthic.** Pertaining to the aggregate of organisms living on or at the bottom of a body of water.
- bioavailability.** The accumulation of trace elements in the tissues of an organism from both food and water sources.
- breccia.** A coarse-grained, clastic rock composed of large, angular, and broken rock fragments that are cemented together in a fine-grained matrix and that can be of any composition, origin, or mode of accumulation; the consolidated equivalent of rubble.
- Cretaceous.** The final period of the Mesozoic era (after the Jurassic and before the Tertiary period of the Cenozoic era), thought to have covered the span of time between 136 and 65 million years ago; also, the corresponding system of rocks.
- deleterious.** Injurious to health.
- diabase.** An intrusive rock whose main components are labradorite and pyroxene and which is characterized by ophitic texture.
- elutriation.** A process in which streambed sediment is mixed with stream water, and after a settling period, the liquid portion is removed, filtered, and chemically analyzed.
- Eocene.** An epoch of the lower Tertiary period, after the Paleocene and before the Oligocene; also, the corresponding worldwide series of rocks.
- eutrophication.** The process by which waters become more eutrophic, especially the artificial or natural enrichment of a lake by an influx of nutrients required for the growth of aquatic plants, such as algae, that are vital for fish and animal life.
- first-order tributary.** Smallest unbranched mapped (1:24,000 map scale) tributaries.
- filtered water.** An operational definition referring to the chemical analysis of that portion of a water-suspended sediment sample that passes through a nominal 0.45-micrometer ( $\mu\text{m}$ ) filter.
- fluvial.** Of or pertaining to a river or rivers.
- gabbro.** A group of dark-colored, basic intrusive igneous rocks composed principally of ferromagnesian minerals and calcium-rich plagioclase feldspar; the approximate intrusive equivalent of basalt.
- galena (lead sulfate).** A bluish-gray to lead-gray mineral:  $\text{PbS}$ ; almost always contains silver; the most important ore of lead.
- gneiss.** A metamorphic rock, generally made up of bands that differ in color and composition, some bands being rich in feldspar and quartz, others rich in hornblende or mica. Varieties are distinguished by texture, characteristic minerals, or general composition and/or origins.
- granitic rock.** A term loosely applied to any light-colored, coarse-grained plutonic rock containing quartz as an essential component, along with feldspar and mafic minerals.
- higher-order stream.** Third-order or larger tributary.
- hydrograph.** A graph showing stage, flow, velocity, or other characteristics of water with respect to time.
- intrusive rock.** An igneous rock mass that has been forced by magmatic activity, while molten or plastic, into fissures or other openings or between layers of other rocks.
- irrigation season.** An operational period defined as June through September for the Yakima River Basin.
- Jurassic.** The second period of the Mesozoic era (after the Triassic and before the Cretaceous), thought to have covered the span of time between 195–190 and 136 million years ago; also, the corresponding system of rocks.
- Kittitas Valley.** The part of the Yakima River Basin that extends 67.5 river miles from the foot of Keechelus Dam to just upstream of Wilson Creek (fig. 3).
- loess.** A widespread, homogeneous, commonly non-stratified, porous, friable, unconsolidated but slightly coherent, usually highly calcareous, fine-grained, blanket deposit of marl or loam, consisting predominantly of silt with subordinate grain sizes ranging from clay to fine sand.
- lower limit of determination (LLD).** An operational limit determined as three times the standard deviation of the blank sample added to the average of the blank.
- lower limit of quantitation (LLQ).** An operational limit determined as ten times the standard deviation of the blank sample added to the average of the blank; a value that has a high probability of being the actual concentration.
- lower-order stream.** First- or second-order tributary.
- lower Yakima Valley.** The part of the Yakima River Basin that extends 107.5 river miles from the city of Union Gap to the mouth of the Yakima River (fig. 3).



**metamorphic rock.** Any rock derived from pre-existing rocks by mineralogical, chemical, and structural changes in response to marked changes in temperature, pressure, shearing stress, and chemical environment.

**mid-Yakima Valley.** That part of the Yakima River Basin that extends a distance of 39.5 river miles from Wilson Creek to the city of Union Gap (fig. 3).

**Miocene.** An epoch of the upper Tertiary period, after the Oligocene and before the Pliocene; also, the corresponding worldwide series or rocks.

**non-irrigation season.** An operational period defined as October through March for the Yakima River Basin.

**peridotite.** A general term for a coarse-grained plutonic rock composed chiefly of olivine with or without other mafic minerals and containing little or no feldspars.

**periphyton.** Micro-organisms that coat rocks, plants, and other surfaces on the bottom of lakes and streams.

**phyllite.** A metamorphic rock in which clay minerals have recrystallized into microscopic micas, giving the rock a silky sheen.

**Pleistocene.** An epoch of the Quaternary period, after the Pliocene of the Tertiary and before the Holocene; also, the corresponding worldwide series of rocks.

**predaceous.** Predatory, rapacious.

**pyrite (iron sulfide).** A common, pale-bronze or brass-yellow, isometric mineral:  $\text{FeS}_2$ ; an important ore of sulfur.

**pyroclastic.** Pertaining to clastic rock material formed by volcanic explosion or aerial expulsion from a volcanic vent; also, pertaining to rock texture of explosive origin.

**redd.** The spawning area or nest of trout or salmon.

**reference dose (RfD).** An estimate of the daily exposure of a non-carcinogen to the human population that is likely to be without appreciable risk of deleterious health effects during a lifetime of 70 years.

**schist.** A metamorphic rock characterized by coarse-grained minerals oriented approximately parallel.

**second-order tributary.** Stream receiving only first-order tributaries.

**sedimentary rock.** A rock resulting from the consolidation of loose sediment that has accumulated in layers.

**serpentinite.** A rock consisting almost wholly of serpentine-group minerals derived from the alteration of previously existing ferromagnesian silicate minerals.

**shale.** A fine-grained sedimentary rock that has a pronounced splitting capability.

**siltstone.** An indurated or somewhat indurated silt having the texture and composition, but lacking the fine lamination or fissility, of shale.

**snowmelt season.** An operational period defined as April through May for the Yakima River Basin.

**sphalerite (zinc-iron sulfide).** A brown or black, sometimes yellow or white, isometric mineral:  $(\text{Zn}, \text{Fe})\text{S}$ ; a widely distributed ore of zinc.

**Tertiary.** The first period of the Cenozoic era (after the Cretaceous of the Mesozoic era and before the Quaternary), thought to have covered the span of time between 65 and three to two million years ago.

**third-order tributary.** Stream receiving only first- and second-order tributaries.

**tuff.** A rock formed from fine-grained pyroclastic particles (ash and dust).

**ultramafic rock.** An igneous rock composed chiefly of mafic minerals.

**unfiltered water.** An operational definition referring to the chemical analysis of a water sample that has not been filtered or centrifuged, nor in any way altered from the original matrix.

**volcanic rock.** A generally finely crystalline or glassy igneous rock resulting from volcanic action at or near the Earth's surface, either ejected explosively or extruded as lava.

**volcaniclastic.** Pertaining to a clastic rock containing volcanic material in whatever proportion, and without regard to its origin or environment.