

U.S. DEPARTMENT OF THE INTERIOR

U.S. GEOLOGICAL SURVEY

GEOCHEMICAL DATA OF DRILL CORE SAMPLES OF CARBONATITES AND ASSOCIATED
IGNEOUS ROCKS, MAGNET COVE COMPLEX, ARKANSAS

by

Marta J.K. Flohr¹ and J. Michael Howard²

Open-File Report 95-621

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards (or with the North American Stratigraphic Code). Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

¹National Center, MS 959
Reston, Virginia 22092

²Arkansas Geological Commission
Vardelle Parham Geology Center
3815 West Roosevelt Road
Little Rock, Arkansas 72204

CONTENTS

PAGE

Introduction	1
Analytical Methods	1
Rock Types	2
Acknowledgments	3
References Cited	3

TABLE

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas .	5
---	---

Introduction

Drill core M5 sampled carbonatites and alkaline igneous rocks of the Magnet Cove Complex, Hot Spring County, Arkansas. Representative samples of these rocks were obtained as part of a study of the Magnet Cove Complex and other alkaline igneous intrusions in the Southern Midcontinent of the United States. Recent detailed studies of the Magnet Cove Complex and its associated Ti-V-Nb mineralization were presented by Erickson and Blade (1963), Flohr and Ross (1989, 1990), Willis (1992), and Flohr (1994); earlier studies are cited by these authors. The Magnet Cove Complex is one of numerous igneous intrusions in the region, some of which crop out and others of which are not exposed at the surface, but which have been sampled by deep drilling and (or) identified by gravity and magnetic surveys (e.g., McKeown, 1982; Hendricks, 1988; see Fig. 1 of Flohr and Howard, 1994, for a sketch map of the region that shows the locations of these intrusions).

Drill core M5 was obtained by Molycorp, Inc. as part of their mineral exploration program. The drilling location was in the northwest corner of Sec. 19, TS3, R17W. The total depth of the core was 750 feet. The core is curated by the Arkansas Geological Commission in the Norman F. Williams Well Sample Depository, Little Rock, Arkansas. This report contains major-, minor-, and trace-element geochemical data of 33 rocks from the core. Short descriptions of the rock types sampled are included. Drill core samples other than those for which geochemical data were obtained were also collected, but these samples are not discussed here.

Analytical Methods

Samples were prepared for geochemical analysis using standard procedures (Arbogast, 1990) by USGS personnel. Outer surfaces of samples were removed before the samples were ground to ensure that no contamination by the coring equipment was present.

All analytical work was performed by USGS personnel. The following methods were used for the analyses reported in Table 1. Where a given element was determined by more than one method, all values are reported without attempting to evaluate the superiority of one method over another method. Si, Ti, Al, total Fe, Mn, Mg, Ca, Na, K, and P were analyzed using wavelength-dispersive X-ray fluorescence spectrometry (WDXRF; Taggart and others, 1987) by J. S. Mee and D. F. Siems and are reported as weight percent of the oxides. Ferrous iron, reported as FeO, was determined by H. Smith who used the colorimetric titration method of Peck (1964). Ferric iron, reported as Fe₂O₃, was calculated from the total iron as obtained by WDXRF and the measured ferrous iron content. F⁻ and Cl⁻ were determined by ion-selective electrode potentiometry using either the method of Kirschenbaum (1988) or of Bodkin (1977) for F and that of Aruscavage and Campbell (1983) for Cl (analysts: C.J. Skeen and N.H. Elsheimer). Total S was analyzed by first combusting the sample in a sulfur analyzer and then measuring the evolved sulfur dioxide by an infrared (IR) detector (Kirschenbaum, 1983; analysts: C.J. Skeen and N.H. Elsheimer). CO₂ was measured by first digesting the sample with HClO₄, a process during which CO₂ is evolved and carried into a coulometric cell. The CO₂ was then converted into a strong acid by ethanolamine and was titrated coulometrically (Engleman and others, 1985; analysts: C. Papp and H. Smith). H₂O⁺, or nonessential water, was determined by weighing the sample before and after drying it for one hour at 100°C (Shapiro, 1975; analyst: H. Smith). Total water was determined by H. Smith who used the method of

Jackson and others (1987): the sample was mixed with a flux, heated to 950°C, and the evolved water was determined coulometrically by Karl-Fischer titration. H_2O^+ , or bound water, is the difference between the total water and H_2O^- . Loss on ignition (LOI) was determined as part of the WDXRF analytical work-up and was determined by weighing the sample before and after heating at 925°C for 45 minutes. A large number of elements, including the rare-earth elements (REE), were determined by instrumental neutron activation analysis (INAA; Baedeker and McKown, 1987) by P. A. Baedeker and J.N. Grossman. Additional important trace elements were analyzed using inductively coupled plasma-atomic emission spectrometry (ICP-AES; Lichte and others, 1987) by M. W. Doughten or using energy-dispersive X-ray fluorescence spectrometry (EDXRF; Johnson, 1984; Johnson and King, 1987) by J. Kent.

Polished thin sections of all samples, except the altered clay-rich rocks, were obtained. Examination of these thin sections and limited X-ray powder diffraction (XRD) analysis provide the basis for the brief mineralogical descriptions presented in the next section.

Rock Types

Lithologies sampled by drill core M5 include a variety of carbonatites and altered rocks. Altered rocks include (1) rocks that contain abundant clay and may be the product of relatively late-stage hydrothermal alteration (these rocks are termed altered clay-rich rocks in Table 1) and (2) altered silicate rocks that are the products of intense metasomatism and reaction between silicate rocks and carbonatite. The following brief descriptions present only the gross characteristics of the rocks. Some of the carbonatites are complex and contain clasts of partly altered silicate rock or earlier-formed carbonatite. Within each of the following groups discussed below, the analyses (Table 1) are ordered from low to high Si content to emphasize the ranges of compositions present in each of these three broad and compositionally diverse groups.

Carbonatites. The dominant lithologies in the M5 drill core are coarse-to-medium-grained calcite carbonatites. These calcite carbonatites are readily distinguished in hand sample and in thin section on the basis of the abundance of accessory silicate, oxide, phosphate, and sulfide minerals. Intervals with concentrations of one or more of the accessory minerals are interspersed with zones of almost pure coarse-grained calcite (barren) carbonatite. The accessory mineral-bearing and barren carbonatite intervals range from a few feet to tens of feet in thickness. The most common accessory minerals are monticellite, apatite, magnetite, perovskite, and pyrite. Barren calcite, monticellite-rich, and apatite-rich carbonatites are the most common. Vuggy REE-mineralized coarse-grained carbonatite (e.g., sample M5-47, with bastnaesite and tentatively identified synchesite) forms a thin zone near the top of the core. Coarse-grained pyrite is most abundant in calcite carbonatite where the carbonatite is in contact with fine-grained mafic silicate rock. Phlogopitized coarse-grained calcite carbonatite (sample M5-549.8) contains masses of phlogopite with subordinate aegirine.

A distinct medium-grained calcite carbonatite contains accessory phlogopite, magnetite, apatite, and sulfide. In some occurrences, particularly in the 706 ft to 727 ft interval (e.g., samples M5-717, M5-721), this carbonatite is faintly laminated, with millimeter-wide bands rich in mica and magnetite. Medium-grained carbonatite is less common at shallower depths, where coarser-grained carbonatites are more common.

Ankeritic carbonatites, which contain calcite and ankerite, are not common in the M5 drill core. Accessory minerals include hematite, apatite, pyrite, quartz, and

arfvedsonite. A Ti-mineralized zone (represented by sample M5-387) contains, in addition to calcite and ankerite, rutile, feldspar, pyrite, mica, and a 14Å clay mineral, which is represented by several weak XRD peaks that are consistent with smectite group minerals.

Altered silicate rocks. Aphanitic mafic rock occurs at the bottom of the core at about 729 ft to 750 ft and at several shallower intervals. Pyroxene grains replaced by biotite and oxide minerals, fine-grained recrystallized groundmasses, and local development of abundant pyrite characterize these rocks. Other altered silicate rocks are more syenitic and contain abundant pyroxene and zoned red garnet. Mafic and felsic rocks that are in contact with carbonatite are the most altered.

Altered clay-rich rocks. Altered clay-rich rocks occur mainly in the interval from about 619 ft to 662 ft, although altered rocks are present at other depths. Smectite and subordinate kaolinite are the clay minerals present. Other minerals in these rocks include calcite, ankerite, siderite, orthoclase, biotite, and sulfide. Enrichments of Ti and REE are also found (e.g., samples M5-658.5 and M5-658.5, respectively).

Acknowledgements

The assistance of Mr. Jack D. Stephenson and Mr. Benjamin F. Clardy, recently retired, (Arkansas Geological Commission) during the first author's visit to Little Rock is gratefully acknowledged.

References Cited

- Arbogast, B.F., ed., 1990, Quality assurance manual for the Branch of Geochemistry: U.S. Geological Survey Open-File Report 90-668, 184 p.
- Aruscavage, P.J. and Campbell, E.Y., 1983, An ion selective electrode method for the determination of chlorine in geological materials: *Talanta*, v. 30, p. 745-749.
- Baedecker, P.A. and McKown, D.M., 1987, Instrumental neutron activation analysis of geochemical materials, *in* Baedecker, P.A., ed., *Methods of geochemical analysis*, U.S. Geological Survey Bulletin 1770, p. H1-H14.
- Bodkin, J.B., 1977, Determination of fluorine in silicates by use of an ion-selective electrode following fusion with lithium metaborate: *Analyst*, v. 102, 409-413.
- Engleman, E.E., Jackson, L.L., and Norton, D.R., 1985, Determination of carbonate carbon in geological materials by coulometric titration: *Chemical Geology*, v. 53, p. 125-128.
- Erickson, R.L., and Blade, L.V., 1963, Geochemistry and petrology of the alkalic igneous complex at Magnet Cove, Arkansas: U.S. Geological Survey Professional Paper 425, 95 p.
- Flohr, M.J.K., 1994, Titanium, vanadium, and niobium mineralization and alkali metasomatism from the Magnet Cove Complex, Arkansas: *Economic Geology*, v. 89, p. 105-130.
- Flohr, M.J.K., and Ross, M., 1989, Alkaline igneous rocks of Magnet Cove Arkansas: Metasomatized ijolite xenoliths from Diamond Jo quarry: *American Mineralogist*, v. 74, p. 113-131.
- Flohr, M.J.K., and Ross, M., 1990, Alkaline igneous rocks of Magnet Cove Arkansas: Mineralogy and geochemistry of syenites: *Lithos*, v. 26, p. 67-98.
- Flohr, M.J.K., and Howard, J. M., 1994, Geochemical data of drill core samples of carbonatites and associated igneous rocks, Benton, Arkansas: U.S. Geological Survey Open-File Report 94-450, 21 p.
- Hendricks, J.D., 1988, Bouguer gravity of Arkansas: U.S. Geological Survey Professional

- Paper 1474, 31 p.
- Jackson, L.L., Brown, F.W., and Neil, S.T., 1987, Major and minor elements requiring individual determination, classical whole rock analysis, and rapid rock analysis, *in* Baedecker, P.A., ed., Methods of geochemical analysis, U.S. Geological Survey Bulletin 1770, p. G1-G23.
- Johnson, R.G., 1984, Trace element analysis of silicates by means of energy-dispersive X-ray spectrometry: X-Ray Spectrometry, v. 13, p. 64-68.
- Johnson, R.G. and King, B.-S.L., 1987, Energy-dispersive X-ray spectrometry, *in* Baedecker, P.A., ed., Methods of geochemical analysis, U.S. Geological Survey Bulletin 1770, p. F1-F5.
- Kirschenbaum, H., 1983, The classical chemical analysis of rocks - the old and the new: U.S. Geological Survey Bulletin 1547, 55 p.
- Kirschenbaum, H., 1988, The determination of fluoride in silicate rocks by ion selective electrode: an update: U.S. Geological Survey Open-File Report, 88-588, 5 p.
- Lichte, F.E., Golightly, D.W., and Lamothe, P.J., 1987, Inductively coupled plasma-atomic emission spectrometry, *in* Baedecker, P.A., ed., Methods of geochemical analysis, U.S. Geological Survey Bulletin 1770, p. B1-B10.
- McKeown, F.A., 1982, Overview and discussion, *in* McKeown, F.A. and Pakiser, L.C., eds., Investigations of the New Madrid, Missouri, earthquake region, U. S. Geological Survey Professional Paper 1236, p. 1-14.
- Peck, L.C., 1964, Systematic analysis of silicates: U.S. Geological Survey Bulletin 1170, 89 p.
- Shapiro, L., 1975, Rapid analysis of silicate, carbonate, and phosphate rocks - revised edition: U.S. Geological Survey Bulletin 1401, 76 p.
- Taggart, J.E., Lindsey, J.R., Scott, B.A., Vivit, D.V., Bartell, A.J., and Stewart, K.C., 1987, Analysis of geological materials by wavelength dispersive X-ray fluorescence spectrometry, *in* Baedecker, P.A., ed., Methods of geochemical analysis, U.S. Geological Survey Bulletin 1770, p. E1-E19.
- Willis, M.A., 1992, Calculation of thermodynamic properties of aqueous titanium species and constraints on the mobility of Ti in aqueous solution in geologic environments: Unpublished M.A. thesis, Washington University, St. Louis, 323 p.

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas

[Sample - number following core number (M5) is depth in feet; Lab No. - USGS laboratory sample number; pct - percent; WDXRF - wavelength-dispersive X-ray fluorescence spectrometry; col. titra. - colorimetric titration; SIE - selective ion electrode; comb./IR - combustion/IR spectroscopy; coul. titra. - coulometric titration; diff. - by difference (methods for obtaining H_2O^+ and H_2O^- are given in the text); calc. - calculated; INAA - instrumental neutron activation analysis; LOI - loss on ignition; 925°C - LOI determined after heating sample to 925°C; EDXRF - energy-dispersive X-ray fluorescence; ICP-AES - inductively coupled plasma-atomic emission spectrometry; nd - not determined due to interference; ppm - parts per million; ppb - parts per billion; -- - not analyzed; ank - ankerite; cal - calcite; cbt - carbonatite; phlg - phlogopitized; alt sil rock - altered silicate rock; alt clay-rich - altered clay-rich rock]

			Sample	M5-288	M5-591.5	M5-722	M5-717	M5-721	M5-115.5	M5-47
			Lab No.	W257414	W257421	W257425	W257423	W257424	W257410	W257409
			Rock type	cal cbt	cal-ank cbt	cal cbt	cal cbt	cal cbt	cal cbt	cal cbt
G	SiO ₂ pct	WDXRF 0.46	1.02	1.39	1.88	2.37	2.96	3.51
	TiO ₂ pct	WDXRF <0.2	<0.2	0.12	0.11	0.05	<0.2	<0.2
	Al ₂ O ₃ pct	WDXRF <0.10	0.25	0.34	0.44	0.67	0.77	1.25
	Fe ₂ O ₃ pct	calc. 0.02	0.64	0.79	1.39	0.64	1.12	0.82
	FeO pct	col. titra	0.05	0.40	0.40	0.99	1.44	0.10	0.36
	MnO pct	WDXRF 0.06	0.26	0.25	0.11	0.07	0.87	0.35
	MgO pct	WDXRF 0.13	0.49	0.18	1.18	1.3	0.62	0.3
	CaO pct	WDXRF 54.6	52.8	52.8	51.4	50.6	50.8	49.3
	Na ₂ O pct	WDXRF <0.15	<0.15	<0.05	<0.15	<0.15	<0.15	<0.15
	K ₂ O pct	WDXRF 0.02	0.07	0.05	0.12	0.51	0.30	0.02
	P ₂ O ₅ pct	WDXRF 1.73	2.58	0.11	1.84	0.82	2.52	<0.05
	F ⁻ pct	SIE 0.14	0.25	0.04	0.12	0.12	0.41	0.28
	Cl ⁻ pct	SIE 0.008	0.014	0.010	0.002	0.001	0.004	0.024
	Total S	.. pct	comb./IR 0.01	0.46	0.46	0.73	0.77	<0.01	0.20
	CO ₂ pct	coul. titra.	.. 42.0	39.6	42.9	38.9	39.9	38.7	40.4
	H ₂ O ⁺ pct	diff. 0.02	0.04	0.03	0.13	0.22	0.41	0.65
	H ₂ O ⁻ pct	diff. 0.03	0.09	0.09	0.09	0.08	0.29	0.68
	-F≡oxy	.. pct	calc. 0.06	0.11	0.02	0.05	0.05	0.17	0.12
	-Cl≡oxy	. pct	calc. 0.002	0.003	0.002	0	0	0.001	0.005
	-S≡oxy	. pct	calc. 0.005	0.23	0.23	0.365	0.385	0	0.10
	Sum pct	calc. 99.3	99.3	100.2	99.8	100.0	100.0	98.4
LOI pct	925°C 41.2	39.1	41.1	37.2	38.0	38.6	40.9	
Na pct	INAA 0.034	0.060	0.123	0.024	0.025	0.046	<3	
K pct	INAA <0.2	<0.2	<0.8	<0.2	0.40	0.36	--	
Ca pct	INAA 38.1	37.4	39.4	33.7	37.2	35.8	38.9	
Fe pct	INAA 0.043	0.771	0.925	1.73	1.62	0.927	0.936	
V ppm	ICP 62	35	43	76	38	78	21	
Li ppm	ICP 10	7.0	<5	6.8	<5	169	29	
Ba ppm	EDXRF	... 450	355	192	750	550	510	134	

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

		Sample	M5-288	M5-591.5	M5-722	M5-717	M5-721	M5-115.5	M5-47
9	Cu ppm .. EDXRF ...	<10	16	10	144	24	<10	17
	Ni ppm .. EDXRF ...	<10	<10	<10	<10	<10	<10	10
	Zn ppm .. EDXRF ...	<10	<10	<10	32	13	<10	12
	Sc ppm .. INAA	0.19	2.34	11.13	0.52	0.29	4.79	6.63
	Cr ppm .. INAA	<0.6	<1	3.8	<1	<0.7	4.3	4.6
	Co ppm .. INAA	0.20	2.66	0.68	18.63	9.65	2.11	2.78
	Ni ppm .. INAA	<6	<12	<20	<10	<11	<14	30
	Zn ppm .. INAA	<2	<4	<5	34.4	16.7	8.5	9.4
	As ppm .. INAA	3.66	7.2	19.6	9.7	2.00	5.3	<26
	Rb ppm .. INAA	1.9	<2	<3	5.1	25.7	15.6	<8
	Sr ppm .. INAA	5642	5503	2401	5875	7229	2134	2108
	Zr ppm .. INAA	<14	<5	<60	31	<23	<50	<190
	Mo ppm .. INAA	<2	<4	8	<0.3	<1	5	<15
	Sb ppm .. INAA	0.171	0.096	1.72	0.219	<0.05	0.237	1.33
	Cs ppm .. INAA	<0.04	0.137	<0.1	0.186	0.443	0.172	1.90
	Ba ppm .. INAA	487	392	197	704	577	545	147
	La ppm .. INAA	184	133	358	113	54.0	137	4521
	Ce ppm .. INAA	174	157.7	549	104.6	48.8	211	5425
	Nd ppm .. INAA	40.1	59.9	201	22.1	11.6	102	1463
	Sm ppm .. INAA	8.06	20.1	36.7	3.68	1.68	33.2	206
	Eu ppm .. INAA	2.66	7.21	9.21	1.19	0.481	11.24	48.3
	Tb ppm .. INAA	1.035	4.11	2.38	0.392	0.150	5.53	1137
	Ho ppm .. INAA	1.17	4.59	<3	<1	<0.4	6.2	<90
	Yb ppm .. INAA	2.29	7.62	11.0	1.05	0.41	10.8	11.2
	Lu ppm .. INAA	0.26	0.91	1.55	0.13	0.06	1.29	1.37
	Hf ppm .. INAA	<0.03	<0.1	<0.09	0.238	<0.08	0.116	0.38
	Ta ppm .. INAA	<0.03	0.102	0.155	0.160	0.058	0.069	<0.1
	Th ppm .. INAA	1.08	55.4	73.2	1.37	0.713	39.4	387
	U ppm .. INAA	0.82	3.62	0.42	1.73	0.60	1.74	6.7
	Au ppb ... INAA	<5	<5	<8	<5	3.4	<9	<40
		Lab No.	W258893	W258900	W258904	W258902	W258903	W258889	W258888
	Y ppm .. ICP	23	120	45	11	2.6	140	97
	Sr ppm .. ICP	4700	6100	2000	5800	6400	2100	1700
	Zr ppm .. ICP	8.7	12	13	39	13	16	36
	Ba ppm .. ICP	450	520	210	870	640	670	130
	Mo ppm .. ICP	<1	<1	3.7	3.8	<1	3.3	2.2
	Nb ppm .. ICP	<1	10	37	83	6.8	3.1	4.7

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

			Sample	M5-387	M5-374	M5-394	M5-436.5	M5-207	M5-125.5	M5-689
			Lab No.	W257416	W257418	W257417	W257419	W257412	W257411	W257422
			Rock type	cal-ank cbt	cal cbt	cal-ank cbt	cal cbt	cal cbt	cal-ank cbt	cal cbt
SiO ₂	pct	WDXRF	4.04	4.74	6.15	7.61	8.89	9.12	9.16
TiO ₂	pct	WDXRF	2.60	0.19	0.16	0.74	1.06	0.04	0.1
Al ₂ O ₃	...	pct	WDXRF	0.83	0.65	0.44	1.45	2.22	0.51	0.47
Fe ₂ O ₃	...	pct	calc.	0.24	3.93	3.31	0.82	0.14	0.42	4.60
FeO	pct	col. titra	3.11	1.20	0.95	1.49	1.27	1.32	1.60
MnO	pct	WDXRF	0.78	0.31	0.26	0.51	1.46	0.48	0.54
MgO	pct	WDXRF	6.36	3.11	1.43	3.78	5.83	4.23	6.05
CaO	pct	WDXRF	39.4	47.8	47.0	42.7	38.3	42.8	44.9
Na ₂ O	...	pct	WDXRF	<0.15	<0.15	<0.15	0.17	<0.15	<0.15	<0.15
K ₂ O	pct	WDXRF	0.08	0.10	0.12	1.48	1.94	0.40	0.08
P ₂ O ₅	...	pct	WDXRF	<0.05	8.67	9.80	9.13	0.53	0.59	9.58
F ⁻	pct	SIE	0.13	0.47	0.52	1.36	0.98	0.61	0.57
Cl ⁻	pct	SIE	0.010	0.006	0.005	0.010	0.018	0.005	0.008
Total S	..	pct	comb./IR	0.14	0.11	0.14	0.11	0.18	0.01	0.14
CO ₂	pct	coul. titra.	39.9	26.1	27.2	27.2	32.8	37.2	21.0
H ₂ O ⁺	...	pct	diff.	0.26	0.29	0.26	0.18	0.30	0.19	0.29
H ₂ O ⁻	pct	diff.	0.32	0.14	0.09	0.09	0.08	0.18	0.10
-F≡oxy	..	pct	calc.	0.05	0.20	0.22	0.57	0.41	0.26	0.24
-Cl≡oxy	..	pct	calc.	0.002	0.001	0.001	0.002	0.004	0.001	0.002
-S≡oxy	..	pct	calc.	0.07	0.055	0.07	0.055	0.09	0.005	0.07
Sum	pct	calc.	98.3	98.1	98.1	99.5	96.5	98.4	99.5
LOI	pct	925°C	40.0	26.9	27.7	27.5	33.9	38.6	20.6
Na	pct	INAA	0.098	0.077	0.100	0.158	0.121	0.041	0.056
K	pct	INAA	<2	<2	<2	1.7	<0.6	0.33	<2
Ca	pct	INAA	28.8	34.6	35.4	32.6	26.8	32.0	33.0
Fe	pct	INAA	2.66	3.81	3.20	1.87	1.206	1.400	4.83
V	ppm	ICP	192	497	571	248	95	197	730
Li	ppm	ICP	37	19	73	438	340	364	46
Ba	ppm	EDXRF	198	500	415	520	>5000	335	380
Cu	ppm	EDXRF	<10	15	16	11	<10	<10	<10
Ni	ppm	EDXRF	<10	<10	10	<10	<10	<10	<10
Zn	ppm	EDXRF	29	60	53	63	59	22	66
Sc	ppm	INAA	12.81	2.12	2.21	1.16	1.29	3.94	2.73
Cr	ppm	INAA	10.9	<1	<1	<2	2.4	1.2	<1
Co	ppm	INAA	6.67	9.89	3.99	6.79	2.95	5.43	13.68

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

		Sample	M5-387	M5-374	M5-394	M5-436.5	M5-207	M5-125.5	M5-689
8	Ni ppm .. INAA <22	<17	<18	<20	<12	<7	<16
	Zn ppm .. INAA 26.6	78.7	63.4	56.4	64.2	24.9	93.7
	As ppm .. INAA <2	22.1	25.7	5.0	<3	2.46	49.1
	Rb ppm .. INAA 5.6	5.3	5.8	63.6	76.5	25.1	6.6
	Sr ppm .. INAA 2247	4557	4265	4262	3179	4323	4041
	Zr ppm .. INAA <60	--	--	<4	<60	<30	<20
	Mo ppm .. INAA 506	--	<12	<5	<12	<0.5	<0.9
	Sb ppm .. INAA 6.12	0.17	0.68	0.500	0.887	0.414	0.24
	Cs ppm .. INAA 0.26	0.24	0.40	0.68	1.05	0.192	0.266
	Ba ppm .. INAA 181	488	428	554	7430	352	372
	La ppm .. INAA 975	665	754	671	1435	78.2	664
	Ce ppm .. INAA 1352	721	805	776	1845	78.6	651
	Nd ppm .. INAA 354	167	186	226	432	20.5	138
	Sm ppm .. INAA 52.1	29.5	33.4	53.4	44.9	4.26	23.5
	Eu ppm .. INAA 12.79	9.85	11.18	17.8	9.34	1.5	8.15
	Tb ppm .. INAA 3.33	3.63	4.25	7.72	2.27	0.753	3.02
	Ho ppm .. INAA 3.6	4.6	6.7	9.3	<10	1.20	4.0
	Yb ppm .. INAA 7.69	7.13	9.41	13.8	9.40	4.37	8.34
	Lu ppm .. INAA 1.05	0.86	1.18	1.52	1.28	0.61	1.08
	Hf ppm .. INAA 0.22	<0.3	0.60	0.50	<0.07	0.538	0.160
	Ta ppm .. INAA 12.52	0.142	0.115	1.52	2.34	0.078	<0.07
	Th ppm .. INAA 97.1	0.44	0.66	8.01	12.46	2.03	0.59
	U ppm .. INAA 2.36	6.22	5.8	5.82	<1	1.76	4.5
	Au ppb ... INAA <14	<11	<12	<9	<17	<4	<9
		Lab No.	W258895	W258897	W258896	W258898	W258891	W258890	W258901
	Y ppm .. ICP 39	90	130	190	50	36	94
	Sr ppm .. ICP 1900	4000	4300	3700	2600	5200	3700
	Zr ppm .. ICP 17	30	38	20	7	24	20
	Ba ppm .. ICP 200	510	460	560	8000	480	390
	Mo ppm .. ICP 608	<1	<1	<1	<1	<1	<1
	Nb ppm .. ICP 1170	199	149	211	218	17	20

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

		Sample	M5-251	M5-365	M5-549.8	M5-704.5B	M5-547.5	M5-750	M5-740.5
		Lab No.	W257413	W257415	W257420	W257450	W257444	W257452	W257451
		Rock type	cal cbt	cal cbt	cal cbt	alt sil rock	alt sil rock	alt sil rock	alt sil rock
SiO ₂ pct	WDXRF	12.7	15.7	19.3	35.0	38.2	41.5	42.8
TiO ₂ pct	WDXRF	0.04	0.16	0.65	0.09	1.03	3.12	2.86
Al ₂ O ₃ pct	WDXRF	0.13	0.42	5.25	13.8	14.3	18.3	19.4
Fe ₂ O ₃ pct	calc.	0.94	2.87	2.97	1.69	3.71	1.28	2.52
FeO pct	col. titra	1.24	1.26	1.19	0.48	0.27	7.72	6.08
MnO pct	WDXRF	0.42	0.67	0.27	0.32	0.19	0.28	0.29
MgO pct	WDXRF	7.57	10.7	7.74	1.96	2.54	3.69	2.49
CaO pct	WDXRF	46.6	32.7	29.5	19.0	20.4	8.41	8.98
Na ₂ O pct	WDXRF	<0.15	1.76	1.23	5.94	4.35	3.54	4.10
K ₂ O pct	WDXRF	0.02	0.53	2.92	2.57	1.05	4.45	4.97
P ₂ O ₅ pct	WDXRF	0.34	1.47	7.73	0.06	0.29	0.72	0.48
F ⁻ pct	SIE	0.10	0.54	1.89	0.27	0.31	0.38	0.42
Cl ⁻ pct	SIE	0.007	0.002	0.003	0.007	0.007	0.006	0.038
Total S pct	comb./IR	0.07	0.01	0.19	0.31	0.39	2.93	0.52
CO ₂ pct	coul. titra.	28.8	30.3	15.7	13.5	6.59	1.02	0.72
H ₂ O ⁺ pct	diff.	0.42	0.82	0.96	3.99	5.18	2.76	2.21
H ₂ O ⁻ pct	diff.	0.12	0.29	0.40	0.26	0.75	0.25	0.20
-F≡oxy pct	calc.	0.04	0.23	0.80	0.11	0.13	0.16	0.18
-Cl≡oxy pct	calc.	0.002	0	0.001	0.002	0.002	0.001	0.009
-S≡oxy pct	calc.	0.035	0.005	0.095	0.16	0.20	1.47	0.26
Sum pct	calc.	99.6	100.4	98.8	99.0	99.2	98.7	98.6
LOI pct	925°C	29.0	30.3	16.7	16.9	11.3	2.76	2.07
Na pct	INAA	0.024	1.40	0.953	4.66	3.44	2.71	3.20
K pct	INAA	<0.1	0.65	2.41	3.1	1.02	4.2	4.5
Ca pct	INAA	32.2	22.6	21.3	13.5	14.8	6.6	7.53
Fe pct	INAA	1.64	3.21	3.19	1.6	2.94	7.21	6.76
V ppm	ICP	25	174	986	97	212	418	336
Li ppm	ICP	9.6	61	354	79	24	33	18
Ba ppm	EDXRF	350	260	>5000	790	520	2600	2800
Cu ppm	EDXRF	12	10	12	26	32	65	36
Ni ppm	EDXRF	<10	<10	<10	<10	<10	<10	10
Zn ppm	EDXRF	34	52	80	28	164	148	130
Sc ppm	INAA	1.504	3.28	18.41	0.287	2.18	6.86	5.11
Cr ppm	INAA	1	0.9	4	<2	2.12	3.3	6.4
Co ppm	INAA	11.35	18.09	6.57	3.78	8.29	20	19.5
Ni ppm	INAA	<10	<14	<21	<8	<15	25	<26

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

		Sample	M5-251	M5-365	M5-549.8	M5-704.5B	M5-547.5	M5-750	M5-740.5
Zn ppm ..	INAA	37.6	66.3	75.8	34.2	162	146	135
As ppm ..	INAA	2.08	14	16.1	3.3	8.9	12.4	9.6
Rb ppm ..	INAA	<2	7.3	153	60.8	32.9	168	179
Sr ppm ..	INAA	4026	2532	3395	2002	2390	2391	2830
Zr ppm ..	INAA	<60	90	<197	263	246	349	384
Mo ppm ..	INAA	<2	<1	<7	6.9	12.1	7.5	<9
Sb ppm ..	INAA	0.138	0.64	0.28	0.282	0.76	3.38	3.03
Cs ppm ..	INAA	0.13	0.65	3.22	12.37	1.18	3.95	7.82
Ba ppm ..	INAA	348	284	9787	784	520	2616	2826
La ppm ..	INAA	66.8	167	256	35.3	105.2	147	136
Ce ppm ..	INAA	65.9	188	298	40.2	131.3	264	232
Nd ppm ..	INAA	16.6	42.3	81	8.9	34.1	94	78.6
Sm ppm ..	INAA	3.94	8.53	15.9	1.87	6.31	16.4	14
Eu ppm ..	INAA	1.52	2.86	5.03	0.643	1.96	4.31	3.77
Tb ppm ..	INAA	0.929	1.31	2.39	0.383	0.873	1.53	1.41
Ho ppm ..	INAA	1.79	1.99	3.3	--	--	--	--
Yb ppm ..	INAA	7.29	6.79	6.49	2.38	3.52	3.37	3.4
Lu ppm ..	INAA	1.05	1.07	0.72	0.321	0.467	0.425	0.421
Hf ppm ..	INAA	<0.05	6.15	4.52	3.59	4.66	6.68	6.08
Ta ppm ..	INAA	0.039	0.108	3.01	1.50	6.16	18.2	17.9
Th ppm ..	INAA	<0.1	0.369	13.75	2.84	12.02	21.2	18.7
U ppm ..	INAA	0.44	1.41	20.3	3.97	7.8	5.62	5.93
Au ppb ...	INAA	<3	<7	<8	15.4	7.9	<7	22.9
		Lab No.	W258892	W258894	W25899	W258929	W258923	W258931	W258930
Y ppm ..	ICP	45	43	85	21	35	47	30
Sr ppm ..	ICP	3900	2200	3100	1900	2100	2900	2300
Zr ppm ..	ICP	11	99	130	240	320	510	330
Ba ppm ..	ICP	370	320	9900	840	460	3800	5700
Mo ppm ..	ICP	<1	<1	<1	5.8	12	5.5	3.5
Nb ppm ..	ICP	46	121	227	191	220	290	317

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

			Sample	M5-567	M5-616.5	M5-626	M5-583	M5-704.5A	M5-658.5	M5-533
			Lab No.	W257445	W257447	W257448	W257446	W257449	W257480	W257477
			Rock type	alt sil rock	alt sil rock	alt sil rock	alt sil rock	alt sil rock	alt clay-rich	alt clay-rich
11	SiO ₂ pct ...	WDXRF 46.0	46.6	48.0	48.1	49.9	34.3	39.2
	TiO ₂ pct ...	WDXRF 2.65	0.96	2.17	1.58	0.22	3.85	0.75
	Al ₂ O ₃ pct ...	WDXRF 18.5	17.9	17.7	17.9	20.9	7.51	14.9
	Fe ₂ O ₃ pct ...	calc. 1.88	5.57	2.91	3.93	2.25	5.9	2.1
	FeO pct ...	col. titra 6.61	1.45	4.40	1.86	0.82	3.5	1.9
	MnO pct ...	WDXRF 0.24	0.39	0.21	0.19	0.23	0.37	0.31
	MgO pct ...	WDXRF 2.74	0.99	2.02	1.40	0.60	4.05	3.65
	CaO pct ...	WDXRF 7.29	7.65	7.47	6.04	4.72	9.79	12.5
	Na ₂ O pct ...	WDXRF 3.56	4.65	6.00	7.49	6.13	0.76	2.07
	K ₂ O pct ...	WDXRF 5.66	6.63	4.23	6.39	6.70	0.07	2.07
	P ₂ O ₅ pct ...	WDXRF 0.64	0.28	0.49	0.35	0.14	<0.05	0.32
	F ⁻ pct ...	SIE 0.42	0.13	0.25	0.14	0.05	1.1	0.48
	Cl ⁻ pct ...	SIE 0.016	0.042	0.016	0.268	0.007	0.011	0.012
	Total S pct ...	comb./IR 0.93	0.21	0.76	0.60	0.11	2.5	0.28
	CO ₂ pct ...	coul. titra. 0.47	1.27	0.62	1.11	1.6	14.7	12.7
	H ₂ O ⁺ pct ...	diff. 1.88	2.85	1.90	1.30	3.94	3.2	3.3
	H ₂ O ⁻ pct ...	diff. 0.27	0.80	0.20	0.17	0.43	2.9	2.4
	-F≡oxy pct ...	calc. 0.18	0.05	0.11	0.06	0.02	0.46	0.20
	-Cl≡oxy pct ...	calc. 0.004	0.009	0.004	0.061	0.002	0.002	0.003
	-S≡oxy pct ...	calc. 0.47	0.11	0.38	0.30	0.06	1.25	0.14
	Sum pct ...	calc. 99.1	98.2	98.9	98.4	98.7	92.8 ¹	98.6
11	LOI pct ...	925°C 1.42	4.26	1.95	1.80	5.55	17.1	18.1
	Na pct ...	INAA 2.72	3.64	4.74	5.95	4.85	<19	1.66
	K pct ...	INAA 4.9	6.0	3.7	5.8	6.2	<100	2.01
	Ca pct ...	INAA 6.4	5.2	6.2	4.57	3.58	<15	9.0
	Fe pct ...	INAA 6.74	5.29	5.66	4.38	2.33	6.16	3.13
	V ppm ..	ICP 260	398	263	186	142	290	346
	Li ppm ..	ICP 25	19	6.4	7.6	144	113	163
	Ba ppm ..	EDXRF	.. 1350	4100	1350	1850	1900	nd	395
	Cu ppm ..	EDXRF 36	<10	38	21	<10	12	63
	Ni ppm ..	EDXRF 14	<10	<10	16	<10	26	<10
	Zn ppm ..	EDXRF 116	205	108	95	124	170	94
	Sc ppm ..	INAA 5.29	0.744	5.03	3.05	0.072	4.13	8.2
	Cr ppm ..	INAA 4.0	1.5	9.3	2.8	<1	<11	6.1
	Co ppm ..	INAA 19.2	5.34	17.8	11.38	0.918	20.7	6.71
	Ni ppm ..	INAA 20	<19	<24	<14	<8	<140	<23

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

		Sample	M5-567	M5-616.5	M5-626	M5-583	M5-704.5A	M5-658.5	M5-533
Zn ppm ..	INAA	113	209	110	97.4	121	265	74.1
As ppm ..	INAA	12.8	8.9	6.2	4.5	4.4	<120	4.89
Rb ppm ..	INAA	199	207	127	183	134	<40	58
Sr ppm ..	INAA	1567	2823	1499	1474	1378	2020	1670
Zr ppm ..	INAA	348	349	333	323	294	<600	99
Mo ppm ..	INAA	8.1	10.1	3.1	3.4	5.8	<70	4.5
Sb ppm ..	INAA	2.18	2.07	1.12	0.92	0.5	1.76	1.38
Cs ppm ..	INAA	2.61	3.67	0.98	5.47	4.61	4.1	4.42
Ba ppm ..	INAA	1345	4073	1375	1859	1932	<190	421
La ppm ..	INAA	128	175	110	101.2	137	17800	67.2
Ce ppm ..	INAA	231	268	189	166	151	14350	91.1
Nd ppm ..	INAA	82.6	54	67	54	20.8	2920	32.7
Sm ppm ..	INAA	14.09	5.9	11.2	8.73	1.69	511	8.67
Eu ppm ..	INAA	3.58	1.44	2.8	2.28	0.368	138.2	2.63
Tb ppm ..	INAA	1.32	0.498	1.015	0.847	0.109	29.6	1.13
Yb ppm ..	INAA	2.95	1.78	2.56	2.33	0.44	5.9	2.91
Lu ppm ..	INAA	0.378	0.231	0.337	0.308	0.063	0.80	0.395
Hf ppm ..	INAA	6.58	3.99	6.68	6.36	3.65	0.84	1.60
Ta ppm ..	INAA	16.3	4.23	10.15	8.27	1.42	11.8	3.65
Th ppm ..	INAA	20.2	23.8	20.5	21.3	19.2	839	8.36
U ppm ..	INAA	5.27	19.23	6.16	7.53	5.01	8.2	1.78
Au ppb ...	INAA	59	16.4	<8	<7	<5	<140	4.7
		Lab No.	W258924	W258926	W258927	W258925	W258928	W258959	W258956
Y ppm ..	ICP	41	29	34	24	3.3	180	26
Sr ppm ..	ICP	1900	4100	1800	1400	1100	1900	1500
Zr ppm ..	ICP	520	660	510	430	320	77	90
Ba ppm ..	ICP	2100	7100	2100	2400	1800	140	450
Mo ppm ..	ICP	9.4	9.5	1.5	4.3	5.6	8.3	4.6
Nb ppm ..	ICP	271	267	179	157	101	952	177

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

			Sample	M5-657.5	M5-728	M5-656.5	M5-619	M5-662
			Lab No.	W257479	W257483	W257478	W257481	W257482
			Rock type	alt clay-rich	alt clay-rich	alt clay-rich	alt clay-rich	alt clay-rich
13	SiO ₂ pct	WDXRF 40.0	41.6	43.8	44.0	45.4
	TiO ₂ pct	WDXRF 3.16	1.03	3.24	1.73	3.61
	Al ₂ O ₃ pct	WDXRF 15.0	19.5	17.2	19.8	19.5
	Fe ₂ O ₃ pct	calc. 5.9	4.4	5.0	0.85	2.8
	FeO pct	col. titra 4.8	3.2	2.3	0.88	1.2
	MnO pct	WDXRF 0.46	0.2	0.18	0.14	0.09
	MgO pct	WDXRF 3.81	4.14	1.75	1.21	1.64
	CaO pct	WDXRF 5.97	6.36	5.19	8.13	4.07
	Na ₂ O pct	WDXRF 1.11	1.93	1.12	1.21	1.41
	K ₂ O pct	WDXRF 1.57	2.60	4.80	4.33	2.17
	P ₂ O ₅ pct	WDXRF 0.49	0.3	0.78	0.52	0.75
	F ⁻ pct	SIE 1.3	0.69	0.49	0.38	0.35
	Cl ⁻ pct	SIE 0.019	0.014	0.005	0.005	0.005
	Total S	.. pct	comb./IR 0.82	0.25	2.0	0.18	1.6
	CO ₂ pct	coul. titra. 9.4	5.3	5.0	6.1	2.8
	H ₂ O ⁺ pct	diff. 3.7	4.9	3.2	4.2	4.6
	H ₂ O ⁻ pct	diff. 5.0	3.9	3.0	3.3	5.0
	-F≡oxy	.. pct	calc. 0.55	0.29	0.21	0.16	0.15
	-Cl≡oxy	.. pct	calc. 0.004	0.003	0.001	0.001	0.001
	-S≡oxy	.. pct	calc. 0.41	0.13	1.0	0.09	0.8
	Sum pct	calc. 101.6	99.9	97.8	96.7	96
LOI pct	925°C 17.6	14.4	11.3	14.1	13.8	
Na pct	INAA 0.879	1.52	0.884	0.941	1.10	
K pct	INAA 1.62	2.33	4.41	4.02	2.08	
Ca pct	INAA 3.91	4.57	4.40	6.22	2.82	
Fe pct	INAA 5.92	5.0	5.28	2.23	3.65	
V ppm	ICP 323	542	395	493	440	
Li ppm	ICP 57	174	127	154	45	
Ba ppm	EDXRF 770	1100	2100	2300	1400	
Cu ppm	EDXRF 32	23	49	32	30	
Ni ppm	EDXRF 10	<10	<10	<10	<10	
Zn ppm	EDXRF 44	46	27	260	530	
Sc ppm	INAA 5.76	2.78	7.51	2.54	6.09	
Cr ppm	INAA 7.1	2.1	9.3	<1	5	
Co ppm	INAA 28.9	6.22	31.1	6.97	28.7	
Ni ppm	INAA <60	<21	<28	<18	<27	

Table 1. Geochemical data of M5 drill core samples, Magnet Cove Complex, Arkansas - Continued

		Sample	M5-657.5	M5-728	M5-656.5	M5-619	M5-662
Zn ppm ..	INAA	37.7	45.2	31.1	219	572
As ppm ..	INAA	7.9	2.55	20.9	9.4	11.7
Rb ppm ..	INAA	36.6	108	125	91	37.2
Sr ppm ..	INAA	1240	870	2250	1970	1340
Zr ppm ..	INAA	210	345	300	429	440
Mo ppm ..	INAA	14.3	3.1	11.4	5.2	12.3
Sb ppm ..	INAA	2.16	0.72	2.78	1.3	4.66
Cs ppm ..	INAA	1.33	13.1	3.48	2.19	2.07
Ba ppm ..	INAA	810	1120	2160	2330	1470
La ppm ..	INAA	109	49.9	164	83.3	122
Ce ppm ..	INAA	145	55	258	97.4	182
Nd ppm ..	INAA	46.9	13.6	85.9	28	62.1
Sm ppm ..	INAA	10.67	2.81	14.33	5.72	10.2
Eu ppm ..	INAA	3.44	0.95	3.37	1.77	2.46
Tb ppm ..	INAA	1.7	0.531	1.14	0.788	0.752
Yb ppm ..	INAA	2.81	3.00	2.99	2.02	1.42
Lu ppm ..	INAA	0.375	0.417	0.407	0.276	0.196
Hf ppm ..	INAA	4.34	4.73	6.45	5.19	8.25
Ta ppm ..	INAA	14.66	3.44	15.22	5.58	17.7
Th ppm ..	INAA	22.1	4.54	18.8	9.34	22.8
U ppm ..	INAA	9.64	2.99	3.8	4.43	6.54
Au ppb ...	INAA	<4	<3	<5	11.5	12.5
		Lab No.	W258958	W258962	W258957	W258960	W258961
Y ppm ..	ICP	31	21	19	22	11
Sr ppm ..	ICP	1200	950	1900	2100	1100
Zr ppm ..	ICP	240	300	350	650	430
Ba ppm ..	ICP	900	1400	2300	3100	1500
Mo ppm ..	ICP	16	3.6	13	5.9	13
Nb ppm ..	ICP	411	192	251	273	251

¹ low sum attributed to high concentrations of REE and other elements that are commonly considered trace elements in many igneous rocks