

U.S. Department of the Interior

U.S. Geological Survey

Digital Release of Stream-Sediment, Heavy-Mineral-Concentrate,  
and Other Geochemical Data Collected in the  
Circle 1° x 3° Quadrangle, Alaska

By

S.L. George

Open-File Report 95-626

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature. Any use of trade names is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey.

U.S. Geological Survey, DFC, Box 25046, MS 973, Denver, CO 80225

1995

#### DISCLAIMER

This diskette publication was prepared by an agency of the United States Government. Neither the United States Government nor any agency thereof nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed in this report or represents that its use would not infringe privately owned rights. Reference therein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. Any views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Although all data and software published on this diskette have been used by the U.S. Geological Survey, no warranty, expressed or implied, is made by the U.S. Geological Survey as to the accuracy of the data and related materials and/or the functioning of the software. The sample description and geologic codes reported in the data files reflect the subjective judgement of the individual sample submitters. Therefore the codes were not necessarily reported consistently and some records are more completely coded than others. The data in these files conform with data found in the NGDB as of August 1995. Further studies and information may add or correct data records for this quadrangle dataset in the NGDB.

## INTRODUCTION

The historic geochemical data presented here were compiled from the Rock Analysis Storage System (RASS) of the U.S. Geological Survey National Geochemical Database (NGDB). The RASS database consists of multi-element chemical and spectrographic analyses for approximately 700,000 geochemical samples collected from the mid-1960's to the late 1980's.

Reconnaissance geochemical sampling and analysis was conducted in the Circle quadrangle in the 1980's as part of the Alaska Mineral Resource Assessment Program (AMRAP). Although collected primarily for mineral resource studies these geochemical data may be useful for environmental or other mineral resource studies in the region. The purpose of this report is to release these data in a more modern, easy-to-use format. While compiling the data for this report, sample coding and geochemical data were inspected and gross errors were corrected. Most of the chemical data included in this report were previously published in hard-copy format, generally without the sample coding information provided here. Previous data reports for the Circle quadrangle include Bailey and others (1987), Cathrall and others (1988; 1991), O'Leary and others (1986), Sutley and others (1987), and Tripp and others (1986).

The Circle quadrangle is bounded by latitude 65° N to 66° N and longitude 144° W to 147° W. The analytical results for 1,630 stream-sediment, 1,140 heavy-mineral-concentrate, and 33 organic samples are given in this report. The data files included on this diskette are separated by sample media type. All data files are in dBase III .dbf format. The first three letters of the filename refer to the quadrangle. Letters following the underscore refer to sample media: CONC, heavy-mineral concentrates; SED, stream sediments; and ORG, organic samples.

## METHODS OF STUDY

### Sample Media

The chemical composition of stream-sediment samples reflects the overall chemistry of rocks contained within the drainage basins. Such information is useful in identifying those basins which contain concentrations of elements that may be related to mineral deposits. Moss-trap sediments represent eroded material in the active stream channel. These samples are similar to stream-sediment samples and were collected in some areas containing thick loess cover (Sutley and others, 1987).

Heavy-mineral concentrates from stream sediment are selectively enriched in certain minerals, including many that may be ore-related. This concentration process permits detection of some elements that are not easily or reliably detected in bulk stream sediment.

Organic sample media generally refer to vegetation samples collected from specimens growing in the flood plains of a stream or river. These samples were usually collected at or near a corresponding stream-sediment or heavy-mineral-concentrate sample locality. Organic sample media for the Circle quadrangle include willow leaves.

### Sample Collection

Stream-sediment samples consist of active alluvium collected from first-order (unbranched) and second-order (below the junction of two first-order) streams as shown on USGS topographic maps (scale = 1:63,360). Each sample was composited from several localities within a 100-ft radius of the sample site. Heavy-mineral concentrates were collected in conjunction with stream sediment sampling by panning sediment at the sample site. No details are available for the soil and organic samples collected.

### Sample Preparation

The stream-sediment samples were air dried, then sieved through an 80-mesh (0.17 mm) stainless-steel sieve. The minus-80-mesh fraction was saved for analysis. Moss-trap-sediment samples were screened and the organic material separated by flotation. After oven drying, the sediment was sieved using 30-mesh (0.50 mm) and 80-mesh (0.17 mm) stainless-steel sieves. Both the minus-30-mesh to plus-80-mesh and the minus-80-mesh fractions were saved for analysis. Sieve size fraction information is recorded in the data tables.

Some stream-sediment samples were prepared with an oxalic-acid leaching technique. The secondary iron and manganese oxides coating stream-sediment grains are scavenging agents that concentrate elements leached from bedrock and colluvium and migrating as free ions in solution. The oxide components are extracted from the minus-80-mesh fraction using a weak, hot oxalic-acid solution (Alminas and Mosier, 1976).

The heavy-mineral concentrates generally were sieved to minus-30 or minus-40 mesh. The sample was further separated with the heavy liquid bromoform into two fractions: a light-mineral fraction (specific gravity 2.86 or less) and a heavy-mineral fraction (specific gravity greater than 2.86). Following heavy-liquid separation, magnetite and other strongly magnetic minerals were removed from the heavy-mineral fraction by use of a hand magnet and a Frantz isodynamic magnetic separator set at 0.2 ampere and saved for analysis. The remaining fraction was again sent through the Frantz separator at a setting of 0.6 amperes and the non-magnetic fraction was retained for analysis.

The willow leaves initially were air dried in cloth bags. The leaves were then pulverized in a blender and ashed in a muffle furnace at a peak temperature of 500° C. The ash was saved for analysis.

### Sample Analysis

The stream-sediment, heavy-mineral-concentrate, and both moss-trap-sediment samples were analyzed using a semiquantitative, direct-current arc emission spectrographic method (Grimes and Marranzino, 1968). Spectrographic results were determined by visually comparing spectra derived from the sample against spectra obtained from laboratory reference standards. Standard concentrations are geometrically spaced over any given order of magnitude of concentration such that values reported for each sample are reported in the geometric sequence 10, 15, 20, 30, 50, 70, 100 etc. The precision of the Grimes and Marranzino (1968) method is plus or minus one reporting interval at 83 percent, or two intervals at 96 percent confidence (Motooka and Grimes, 1976). The elements analyzed and their nominal limits of determination are listed in table 1.

Selected stream-sediment samples were analyzed for gold, copper, and lead by atomic-absorption methods (Ward and others, 1969). Ultraviolet fluorometry (modification of Centanni and others, 1956) was used to determine uranium in selected stream-sediment samples. Mercury was determined by a mercury-vapor detector developed by Vaughn and McCarthy (1964). The lower limits of determination for these elements in parts per million (ppm) are: gold, 0.05; copper, 5; lead, 5; uranium, 0.05; and mercury, 0.02.

Selected stream-sediment samples and the minus-80 fraction of the moss-trap-sediment samples were analyzed for arsenic, bismuth, cadmium, antimony, and zinc by atomic-absorption spectrometry (O'Leary and Viets, 1986). The lower limits of determination for these elements in parts per million are: arsenic, 5 or 10; bismuth, 1; cadmium, 0.1; antimony, 2; and zinc, 5.

The heavy-mineral concentrates and oxide residues were analyzed by the Grimes and Marranzino (1968) emission spectrographic procedure as described above, with the following modification: to eliminate the spectral interferences caused by high concentrations of iron, 5 mg of prepared sample was used instead of 10 mg, thus raising the lower limit of determination by two steps (Table 1).

Selected heavy-mineral-concentrate samples were analyzed for the six platinum group elements (PGE) by fire-assay/ICP mass spectrographic analysis. The lower limits of determination for these elements in parts per million are: platinum, 0.0005; palladium, 0.0008; rhodium, 0.0005; ruthenium, 0.0005, osmium, 0.001; and iridium, 0.0005.

Selected heavy-mineral-concentrate samples were analyzed for silver, bismuth, cadmium, copper, cobalt, nickel, lead, and zinc by atomic absorption (modification of Nakagawa, 1975). Gold, indium, and thallium contents were determined by atomic absorption (modification of Hubert and Lakin, 1973). The lower limits of determination for these elements in parts per million are: silver, 0.2; bismuth, 5; cadmium, 0.2; copper, 1; cobalt, 1; nickel, 1; lead, 5; zinc, 1; gold, 0.2; indium, 0.2; thallium, 0.2. Selected magnetic concentrates were measured to determine equivalent uranium. The radiation of each magnetic concentrate was reported as percent equivalent uranium and the lower limit of determination is 0.003 percent (Pan and others, 1980).

The ashed willow leaves were analyzed by a semiquantitative, direct-current arc emission spectrographic method developed by Mosier (1972) for the analysis of plant ash.

#### DESCRIPTION OF DATA

Sample description, geologic, and analytical data are presented in each of the sample media files. Sample site locations are given as latitude and longitude both in decimal degree and degree-minute-second formats in the tables. The following list summarizes table structure and sample description column headings. For table structures, "A8" refers to an alphanumeric format eight characters wide, while "N" indicates a numeric column format. Sample description code explanations are listed in Appendix A.

Table Structure	Column Identifier	Description
A6	Jobnum	assigned laboratory job number
A7	Labnum	assigned sample laboratory number
A8	Fieldnum	sample field identification number
A8	Date_sub	date sample submitted to laboratory
A20	Submitter	submitter name
A20	Subm2	secondary submitter name
A9	Lat_dms	latitude in degree-minute-second
A11	Lon_dms	longitude in degree-minute-second
N	D_lat	latitude in decimal degrees
N	D_lon	longitude in decimal degrees
A2	LL_precis	latitude/longitude precision
A1	St	type of sample media
A1	Mc	method sample collected
A1	Sc	sample source
A1	Rt	rock type
A1	Ga	geologic age
A2	Ms	mesh/sieve code
A2	Sd	additional sample description information
A2	Sd3	additional sample description information
A2	Sd4	additional sample description information
A2	Sd5	additional sample description information
A2	Sp1	sample lab preparation information
A2	Sp2	sample lab preparation information
A2	Sp3	sample lab preparation information
A2	Cm	lab/submitter comments

Chemical data follows the above sample description information in the data files. The chemical data are accurate to two significant digits. Trailing zeros are nonsignificant. Columns in which the element headings

contain the prefix "S" represent emission-spectrographic data. The prefix "Aa" indicates atomic absorption analyses; "Inst" indicates instrumental method (fluorometric or mercury-vapor); the prefix "As" indicates fire-assay/ICP-MS; and "Samp\_wt" indicates sample weight in grams for fire assay analyses for PGE. The suffix "p" indicates a partial digestion; "t" indicates a total digestion; and "sw" indicates sample weight in grams. The column heading "Equiv\_u" indicates percent equivalent uranium. The results for all elements are reported in parts per million (ppm) except for iron, magnesium, calcium, titanium, sodium, and phosphorus, which are given in percent (pct).

Definitions of the qualifier codes used in the tables are as follows: B, sample not analyzed for this element; N, not detected at the specified level of detection; L, detected, but below the specified limit of determination; G, greater than the specified upper limit of determination; and H, values not determined due to interference.

#### REFERENCES CITED

- Alminas, H.V., and Mosier, E.L., 1976, Oxalic acid leaching of rock, soil, and stream-sediment samples as an anomaly-accentuation technique: U.S. Geological Survey Open-File Report 76-275.
- Bailey, E.A., Lee, G.K., and Light, T.D., 1987, Semiquantitative emission spectrographic analyses of stream-sediment samples collected in the Livengood and western one-third of the Circle 1 x 3 degree quadrangles, Alaska: U.S. Geological Survey Open-File Report 87-264, 74 p.
- Cathrall, J.B., Carlson, R.R., Antweiler, J.C., and Mosier, E.L., 1991, Platinum group elements in native gold, alluvium concentrates, and mineralized rock concentrates for some gold mining districts of Alaska: U.S. Geological Survey Open-File Report 91-348, 36 p.
- Cathrall, J.B., Tripp, R.B., McDanal, S.K., Mosier, E.L., and VanTrump, G., 1988, Analytical results, geochemical signatures, mineralogical data, and sample locality map of placer gold and heavy-mineral-concentrates from the Circle mining district, Circle quadrangle, Alaska: U.S. Geological Survey Open-File Report 88-676, 48 p.
- Centanni, F.A., Ross, A.M., and DeSesa, M.A., 1956, Fluorometric determination of uranium: Analytical Chemistry, v. 28, p. 1651.
- Grimes, D.J., and Marranzino, A.P., 1968, Direct-current arc and alternating current spark emission spectrographic field methods for the semiquantitative analysis of geologic materials: U.S. Geological Survey Circular 591, 6 p.
- Mosier, E.L., 1972, A method for semiquantitative spectrographic analysis of plant ash for use in biochemical and environmental studies: Applied Spectroscopy, v. 26, no. 6, p. 636-641.
- Motooka, J.M., and Grimes, D.J., 1976, Analytical precision of one-sixth order semiquantitative spectrographic analysis: U.S. Geological Survey Circular 738, 25 p.
- O'Leary, R.M., and Viets, J.G., 1986, Determination of arsenic, antimony, bismuth, cadmium, copper, lead, molybdenum, silver, and zinc in geologic materials by atomic absorption spectrophotometry using a hydrochloric acid-hydrogen peroxide digestion: Atomic Spectroscopy, v. 7, p. 4-8.
- O'Leary, R.M., Hoffman, J.D., Risoli, D.A., and Tripp, R.B., 1986, Analytical results and sample locality map of stream-sediment and heavy-mineral-concentrate samples from the Circle quadrangle, Alaska: U.S. Geological Survey Open-File Report 86-204, 123 p.
- Pan, K.L., Overstreet, W.C., Robinson, K., Hubert, A.E., and Crenshaw, G.L., 1980, Equivalent uranium and selected minor elements in magnetic concentrates from the Candle quadrangle, Solomon quadrangle, and elsewhere in Alaska: U.S. Geological Survey Professional Paper 1135, 115 p.
- Sutley, S.J., O'Leary, R.M., Lee, G.K., and Light, T.D., 1987, Analytical results and sample locality map of stream-sediment, moss-trap-sediment, and heavy-mineral-concentrate samples from the White Mountains Recreation Area, Livengood and Circle quadrangles, east-central Alaska: U.S. Geological Survey Open-File Report 87-285, 110 p.

Tripp, R.B., Crim, W.D., Hoffman, J.D., O'Leary, R.M., and Risoli, D.A., 1986, Mineralogical and geochemical maps showing the distribution of selected minerals and elements found in the minus-80-mesh stream-sediment and related minus-30-mesh heavy-mineral-concentrate samples from the Circle quadrangle, Alaska: U.S. Geological Survey Open-File Report 83-170 F,G,H, 7 p.

Ward, F.N., Nakagawa, H.M., Harms, T.F., and Van Sickle, G.H., 1969, Atomic-absorption methods of analysis useful in geochemical exploration: U.S. Geological Survey Bulletin 1289, 45 p.

Table 1.--Limits of determination for Emission Spectrographic Analysis  
Numbers in () were limits before 1988

Elements	Sediments		Concentrates	
	Lower limit	Upper limit	Lower limit	Upper limit
Percent				
Iron (Fe)	.05	20	.1	50
Magnesium (Mg)	.02	10	.05	20
Calcium (Ca)	.05	20	.1	50
Sodium (Na)	.2	5	.5	10
Titanium (Ti)	.002	1	.005	2
Phosphorus (P)	.2	5	.5	10
Parts per million				
Silver (Ag)	.5	5,000	1.0	10,000
Arsenic (As)	200	10,000	500	20,000
Gold (Au)	10	500	20	1,000
Boron (B)	10	2,000	20	5,000
Barium (Ba)	20	5,000	50	10,000
Beryllium (Be)	1	1,000	2	2,000
Bismuth (Bi)	10	1,000	20	2,000
Cadmium (Cd)	20	500	50	1,000
Cobalt (Co)	(5) 10	2,000	(10) 20	5,000
Chromium (Cr)	(5) 10	5,000	(10) 20	10,000
Copper (Cu)	5	20,000	10	50,000
Gallium (Ga)	5	500	10	1,000
Germanium (Ge)	10	100	20	200
Indium (In)	2	--	--	--
Lanthanum (La)	(20) 50	1,000	(50) 100	2,000
Lithium (Li)	200	--	--	--
Manganese (Mn)	10	5,000	20	10,000
Molybdenum (Mo)	5	2,000	10	5,000
Niobium (Nb)	(10) 20	2,000	(20) 50	5,000
Nickel (Ni)	5	5,000	10	10,000
Lead (Pb)	10	20,000	20	50,000
Palladium (Pd)	--	--	5	1,000
Platinum (Pt)	--	--	20	1,000
Antimony (Sb)	100	10,000	200	20,000
Scandium (Sc)	5	100	10	200
Tin (Sn)	10	1,000	20	2,000
Strontium (Sr)	100	5,000	200	10,000
Thorium (Th)	100	2,000	200	5,000
Thallium (Tl)	2	--	--	--
Vanadium (V)	10	10,000	20	20,000
Tungsten (W)	(50) 20	10,000	(100) 50	20,000
Yttrium (Y)	10	2,000	20	5,000
Zinc (Zn)	200	10,000	500	20,000
Zirconium (Zr)	10	1,000	20	2,000

APPENDIX A  
Explanation of sample description codes

Sample Type (St)

A rock  
B unconsolidated sediment  
C organic material  
D soil  
E water  
F other  
G gas

Method collected (Mc)

A single (grab)  
B composite  
C channel  
D other

Sample source (Sc)

A outcrop  
B mine  
C dump or prospect pit  
D float  
E drill hole, well  
F marine  
G other  
H stream  
I spring  
J lake  
K aquaduct, canal, irr. ditch  
L atmosphere

Rock type (Rt)

A unidentified rock  
B sedimentary rock  
C metamorphic rock  
D igneous rock  
E unconsolidated sediment  
F conglomerate  
G sandstone  
H siltstone  
I claystone  
J shale  
K limestone or dolomite  
L carbonate  
M gneiss  
N schist  
O quartzite  
P marble  
Q skarn  
R phyllite or slate  
S felsic igneous  
T intermediate igneous  
U mafic igneous  
V ultramafic igneous  
W feldspathoidal  
X chert or jasperoid  
Y other

#### Geologic age of sample (Ga)

A Precambrian undifferentiated  
B Early Precambrian  
C Middle Precambrian  
D Late Precambrian  
E Paleozoic undifferentiated  
F Cambrian  
G Ordovician  
H Silurian  
I Devonian  
J Mississippian  
K Pennsylvanian  
L Permian  
M Mesozoic undifferentiated  
N Triassic  
P Jurassic  
Q Cretaceous  
R Tertiary undifferentiated  
S Paleocene  
T Eocene  
U Oligocene  
V Miocene  
W Pliocene  
X Quaternary undifferentiated  
Y Pleistocene  
Z Holocene

#### Mesh/sieve fraction (Ms)

A unknown, assumed to be -80 mesh  
B identified as -80 mesh  
C identified as -100 mesh  
D identified as -120 mesh  
E identified as -150 mesh  
F identified as -200 mesh  
G identified as -60 mesh  
H identified as -40 mesh  
I identified as -35 mesh  
J identified as -30 mesh  
K identified as -24 mesh  
L identified as -20 mesh  
M identified as -30+80 mesh

#### Sample description (Sd)

AL alluvium  
AS ash  
CL clay  
CV colluvium  
C pan or artificial concentrate  
C1 concentrate, high magnetic fraction  
C2 concentrate, moderately magnetic fraction  
C3 concentrate, low or non-magnetic fraction  
GV gravel  
GT grit  
HS heavy sand  
LO loess  
MD mud  
OZ ooze  
SN sand  
SD stream sediment  
SI silt  
TI till

#### Sample description (Sd3)

AN animal parts  
CS combined splits of  
heavy-mineral-concentrate  
DM detrital magnetites  
MS splits of magnetites  
NS non-magnetic splits from  
heavy-mineral-concentrate  
OA oxalic acid leachate  
VG vegetation

#### Sample description (Sd5)

MI mill tailings  
MT moose pellets  
WL willow leaves

#### Sample description (Sd4)

GD glacial debris  
MT moss-trap-sediment sample  
PT peat material  
SP spruce

#### Sample preparation (Sp1)

AD ashed  
BR bromoform  
GR ground

#### Sample preparation (Sp2)

FR Frantz isodynamic separator  
PV pulverized

Sample preparation (Sp3)

FS fire assay PGE  
HG hand ground  
HM separated by hand magnet  
RT split into red tops

Lab/submitter comments (Cm)

HG high organic content  
RS rock/soil survey  
VG visible gold

Lat/lon precision (LL\_precis)

A apparently accurate to nearest  
second  
B apparently accurate to nearest  
minute  
C apparently accurate to nearest  
degree

Information for monthly listing of USGS publications:

OF-95-626. Digital release of stream-sediment, heavy-mineral-concentrate, and other geochemical data collected in the Circle 1 x 3 degree quadrangle, Alaska, by S.L. George. 1995. One 3-1/2 inch diskette.

Requirements: IBM PC or compatible computer; 3-1/2 inch, 1.44 Mb floppy disk drive; PC or MS DOS; software capable of importing dBase III .DBF files.