



**INTRODUCTION**

This report, prepared in cooperation with the Fort Bend Subsidence District, presents maps for the Chicot and Evangeline aquifers in Fort Bend County and adjacent areas showing the approximate water-level altitudes in wells in 1990 (figs. 1, 2). The most recent previously published water-level-altitude maps (and water-level-change maps) for the two aquifers are by Coplin and others (1997). The earliest water-level-altitude maps (and water-level-change maps) for the Chicot aquifer are by Wesselman (1972). The first maps of water-level altitudes (and water-level changes) for the Chicot and Evangeline aquifers are by Locke (1990).

**GEOHYDROLOGY**

The Chicot aquifer consists of sediments of Holocene and Pleistocene age, and the underlying Evangeline aquifer consists of sediments of Pliocene and Miocene age. The sediments of the Chicot aquifer are sand and clay. In most of Fort Bend County, the Evangeline aquifer is composed of sand. The basis for differentiation of the Chicot and Evangeline aquifers in Fort Bend County is differences in stratigraphic position, lithology, and hydraulic conductivity (Wesselman, 1972, p. 9, p. 16). A weak hydraulic connection between land surface and the Chicot aquifer and between the Chicot and Evangeline aquifers allows vertical movement of water into and between the aquifers; the aquifer system thus is characterized as "leaky" (Gabrysch and Coplin, 1990, p. 2).

**WATER-LEVEL MEASUREMENTS**

Water-level measurements used to prepare this report were obtained by steel tape, airline, electronic sensor, and from reports by well operators. Most wells are pumped once daily, but some are pumped more frequently. Multiple measurements were made when wells were not being pumped. However, antecedent pumping conditions were not always known. Water-level measurements were made in January and February, the months when water levels usually are highest. The wells selected for water-level measurements had comparable depths and screened intervals. Some water-level measurements from wells outside the study area are obtained to increase water-level-contour control.

**REFERENCES CITED**

- Coplin, L.S., Kasmarek, M.C., Santos, H.X., and Noble, J.E., 1997, Water-level altitudes 1997 and water-level changes 1990–97 and 1996–97 in the Chicot and Evangeline aquifers, Fort Bend County and adjacent areas, Texas: U.S. Geological Survey Open-File Report 97-180, 6 sheets.  
Gabrysch, R.K., and Coplin, L.S., 1990, Land-surface subsidence resulting from ground-water withdrawals in the Houston-Galveston region, Texas, through 1987: Harris-Galveston Coastal Subsidence District Report of Investigations 90-01, 53 p.  
Locke, G.L., 1990, Ground-water withdrawals, water-level changes, land-surface subsidence, and ground-water quality in Fort Bend County, Texas, 1969–87: U.S. Geological Survey Water-Resources Investigations Report 90-4012, 155 p.  
Wesselman, J.B., 1972, Ground-water resources of Fort Bend County, Texas: Texas Water Development Board Report 155, 185 p.

**VERTICAL DATUM**

**Sea level:** In this report, "sea level" refers to the National Geodetic Vertical Datum of 1929 (NGVD of 1929)—a geodetic datum derived from a general adjustment of the first-order level nets of the United States and Canada, formerly called Sea Level Datum of 1929.

**EXPLANATION**

- -50 --- Water-level contour—Shows altitude at which water level would have stood in tightly cased well. Contour interval 50 feet. Datum is sea level  
----- Boundary of study area  
o Data point—Well in which water-level measurement was made. One point can represent more than one well

**Figure 1.** Map showing approximate water-level altitudes in the Chicot aquifer, Fort Bend County and adjacent areas Texas, January–February 1990.

**WATER-LEVEL ALTITUDES IN WELLS COMPLETED IN THE CHICOT AND EVANGELINE AQUIFERS,  
FORT BEND COUNTY AND ADJACENT AREAS, TEXAS, JANUARY–FEBRUARY 1990**

By  
Mark C. Kasmarek  
1997