



Carbon Storage and Late Holocene Chronostratigraphy of a Mississippi River Deltaic Marsh, St. Bernard Parish, Louisiana

First Report: Mississippi Basin Carbon Project Process Studies

Edited by Helaine Walsh Markewich

U.S. GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards (or with the North American Stratigraphic Code). Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Contributing Authors (listed alphabetically):

Louis D. Britsch

U.S. Army Corps of Engineers, New Orleans District, New Orleans Louisiana 70160-0267

Gary R. Buell

U.S. Geological Survey, 3039 Amwiler Rd., Atlanta, Georgia 30360-2824

Douglas L. Dillon

U.S. Army Corps of Engineers, New Orleans District, New Orleans, Louisiana 70160-0267

Carmen M. Fraticelli

Center for Excellence in Palynology (CENEX), Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803

Terry L. Fries

U.S. Geological Survey, 345 Middlefield Rd., MS 962, Menlo Park, California 94025

Helaine Walsh Markewich

U.S. Geological Survey, 3039 Amwiler Rd., Atlanta, Georgia 30360-2824

John P. McGeehin

U.S. Geological Survey, MS 955, National Center, Reston, Virginia 20192

Jodi B. Pracht

Center for Excellence in Palynology (CENEX), Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803

John A. Robbins

Great Lakes Environmental Research Laboratory, National Oceanic and Atmospheric Administration, 2205 Commonwealth Blvd., Ann Arbor,

Bruce M. Samuel

Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803

John H. Wrenn

Center for Excellence in Palynology (CENEX), Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803

CONTENTS - Hyperlinks in CONTENTS are internal. External links to specific figures and tables are within the document.

ACKNOWLEDGEMENTS

INTRODUCTION

H.W. Markewich

GEOLOGIC AND HYDROGEOLOGIC SETTING

L.D. Britsch, D.L. Dillon, and H.W. Markewich

FIELD METHODS

L.D. Britsch and D.L. Dillon

¹⁴C AGE DETERMINATIONS

J.P. McGeehin

¹³⁷Cs ANALYSIS

T.L. Fries and J.H. Robbins

BIOCHRONOSTRATIGRAPHY

J.H. Wrenn, J.B. Pracht, and C.M. Fraticelli, and B.M. Samuel

COMPOSITIONAL ANALYSIS

T.L. Fries, G.R. Buell, and H.W. Markewich

DATA INTERPRETATION AND SUMMARY

H.W. Markewich, G.R. Buell, T.L. Fries, and J.P. McGeehin

REFERENCES

FIGURES

FIGURE 1. Location of study area in relation to the Mississippi River basin

FIGURE 2. Physiographic provinces of southern Louisiana

FIGURE 3. Major Holocene deltaic lobes of the Mississippi River

FIGURE 4A. Depositional environments in the vicinity of the St. Bernard core locality.

FIGURE 4B. Map showing St. Bernard core locality in relation to New Orleans, Chalmette, and the Mississippi River Gulf Outlet Canal.

FIGURE 5. Photograph of vibracore set up.

FIGURE 6. ¹⁴C bomb curve for the core SB1a.

FIGURE 7. ¹³⁷Cs curve for the core SB1b.

FIGURE 8. Pollen diagram for core SB1c.

FIGURE 9. Diagnostic pollen and other microfossils for core SB1c -photographic plate.

FIGURE 10A. 1845 map of the Mississippi River delta from New Orleans to the Gulf of Mexico.

FIGURE 10B. Blow up (zoom) of the 1845 map showing locations of plantations and railroads near the St. Bernard core locality.

FIGURE 11. Phytolith and other siliceous microfossil diagram for core SB1c, St. Bernard Parish, LA.

FIGURE 12. Diagnostic phytoliths and siliceous microfossils, core SB1c -photographic plate.

FIGURE 13. Concentration diagram of charcoal and carbonaceous spherules for core SB1c.

FIGURE 14. Carbonaceous spherules present in core SB1c - photographic plate.

FIGURE 15. Composite diagram of diagnostic microfossils for core SB1c.

FIGURE 16. Carbon, nitrogen, and bulk density profiles for cores SB1a and SB1c.

FIGURE 17. Elemental chemistry profiles for cores SB1a and SB1c.

FIGURE 18. Carbon storage for core SB1a and SB1c.

FIGURE 19A, B. Decomposition rates

TABLES

TABLE 1. Data for ^{14}C age determinations for peat samples from core SB1c.

TABLE 2. Data for ^{14}C age determinations for peat samples from core SB1a.

TABLE 3. ^{137}Cs data for bomb-curve analysis of core SB1b.

TABLE 4. St. Bernard Parish, Louisiana cores - events and potential stratigraphic markers.

TABLE 5. Sample depths and analyses for microscopic markers in core SB1c.

TABLE 6. Discussion of pollen zones depicted in figure 8.

TABLE 7. Selected entries of the 1860 census of the large slave holders of Louisiana.

TABLE 8. Phytolith and other siliceous microfossil data for core SB1c.

TABLE 9. Carbonaceous spherule data for core SB1c.

TABLE 10A. Physical and chemical data for samples from core SB1a.

TABLE 10B. Major-element and trace-element data for samples from core SB1a.

TABLE 10C. Trace-element data for samples from core SB1a.

TABLE 11A. Physical and chemical data for samples from core SB1c.

TABLE 11B. Major-element and trace-element data for samples from core SB1c.

TABLE 11C. Trace-element data for samples from core SB1c.

TABLE 12. Mercury concentrations for selected samples from core SB1c.

TABLE 13. Organic carbon cumulative mass and age-interval accumulation rates for cores SB1a and SB1c.

ACKNOWLEDGEMENTS

We would like to thank Caryl Wipperfurth and Brian Bossak (U.S. Geological Survey, Atlanta, GA) for producing most of the figures in this report.

INTRODUCTION

Helaine Walsh Markewich

Today, the causes, results, and time scale(s) of climate change, past and potential, are the focus of much research, news coverage, and pundit

speculation. Many of the US government scientific agencies have some funds earmarked for research into past and (or) future climate change (National Science and Technology Council, 1997). The Mississippi Basin Carbon Project (MBCP) is part of the U.S. Geological Survey (USGS) effort in global change research. The project is motivated by the need to increase our understanding of the role of terrestrial carbon in the global carbon cycle, particularly in the temperate latitudes of North America. The global land area between 30° and 60°N is thought to be a large sink for atmospheric CO₂ (IPCC, 1996). The identity of this sink is unknown, but is in part the soil and sediment that makes up the upper several meters of the Earth's surface. The MBCP focuses on the Mississippi River basin, the third largest river system in the world (fig. 1), that drains an area of 3.3 x 10⁶ km² (1.27 x 10⁶ mi²). The Mississippi River basin includes more than 40 percent of the land surface, and is the home of more than one-third of the population, of the conterminous United States. Because climate, vegetation, and land use vary greatly within the Mississippi River basin, the primary terrestrial sinks for carbon need to be identified and quantified for representative parts of the basin.

The primary goal of the MBCP is to quantify the interactive effects of land-use, erosion, sedimentation, and soil development on carbon storage and nutrient cycles within the Mississippi River basin. The project includes spatial analysis of a wide variety of geographic data, estimation of whole-basin and sub-basin carbon and sediment budgets, development and implementation of terrestrial carbon-cycle models, and site-specific field studies of relevant processes. Areas can be studied and compared, and estimates can be made for whole-basin carbon storage and flux.

Site specific studies are directed at estimating rates of carbon accumulation, decomposition, erosion, transport, and deposition; and particularly at assessing the sensitivity of these rates to climatic, hydrologic, topographic and land-use gradients. To date, process studies are focused on that part of the Mississippi River basin between 82° and 100°W longitude. Study areas include open- and closed-system lakes, primarily in the northern part of the Mississippi River basin; farmland in Iowa, Ohio, and Mississippi; and marshes and swamps in the Mississippi River deltaic plain (figs. 1 and 2).

One of the most important aspects of the project is *time*, and the ability to determine the rate(s) of carbon accumulation per unit of time. Therefore, attention had to be given to the techniques available for looking at small intervals of time. With this goal in mind, the first field study initiated by the MBCP was an investigation into the rates of carbon accumulation in the Mississippi River deltaic plain. The goal is to determine the accumulation rates on a decadal time scale for the last several hundred years. This ongoing effort is collaborative with the U.S. Army Corps of Engineers (COE), New Orleans District, Louisiana; the Center for Excellence in Palynology (CENEX), Geology Department Louisiana State University, Baton Rouge; and the Great Lakes Environmental Research Laboratory, National Ocean and Atmospheric Administration, Ann Arbor, Michigan.

Field sites were chosen to include most of the major geomorphic and (or) ecological environments characteristic of the Mississippi River deltaic plain. Since the fresh, brackish, and salt marshes comprise the most prominent deltaic environment, approximately 1,500 km² (Sasser and others, 1995), the first year's efforts were concentrated in three brackish- to fresh-water marshes located in the western, central, and eastern parts of the deltaic plain (fig. 1). Chronostratigraphic reconstruction of the pre-human development part of the geologic record relied on ¹⁴C isotopic, compositional, and palynomorphologic analyses. Construction of a chronostratigraphy for the post-European development part of the geologic record involved most of the same analytical techniques. However, ¹³⁷Cs and ¹⁴C bomb spike analyses were used to establish "dated" horizons for the last 30-45 years, and diagnostic microstratigraphic markers were used to identify horizons representing the time period from 1700 to 1900, too recent a time interval for standard ¹⁴C isotopic analysis. Carbon accumulation rates and inventory estimates were calculated using data from this study, data from published sources, and data from unpublished sources that were made available to this investigation.

FIGURE 1. The Mississippi River drains an area of 3.3 x 10⁶ km² and forms a well-defined bird's foot delta as it enters the Gulf of Mexico. Mississippi Basin Carbon Project core localities in the Mississippi River deltaic plain are indicated by different colored circles: purple, marsh; dark brown, swamp; green, floodplain and (or) back levee. Core locations are named for the parish in which they are located; parish names are shown. The St. Bernard core locality is in the easternmost part of the study area. Triangles indicate localities for which surface and near-surface organic carbon data are available from the National Soil Survey Laboratory in Lincoln, Nebraska (NSSL, 1994).

This report is the first of the Mississippi Basin Carbon Project process studies. It presents findings from the easternmost core locality in the Mississippi River deltaic plain, a marsh on a western border of Lake Borgne in St. Bernard Parish, southeastern Louisiana. It includes: (a) the geologic setting and Holocene geologic history of the easternmost part of the Mississippi River deltaic plain, (b) a brief description of investigative methods used for chronostratigraphic analysis, and (c) results, including the rate(s) of organic carbon accumulation and the amount of organic carbon represented by the organic matter. This report's companion, Markewich (*in press*), contains detailed descriptions of the principal field and laboratory methods used for investigations of the MBCP in the Mississippi River deltaic plain. Data from this study, along with other available data on surface and near-surface organic matter will be used to estimate the carbon inventory and flux, wholly and in part, for the Mississippi River basin.

REFERENCES

[Return to CONTENTS](#)

GEOLOGIC AND HYDROGEOLOGIC SETTING

L.D. Britsch, D.L. Dillon, and H.W. Markewich

Geomorphology

The St. Bernard core locality is within the St. Bernard subdelta of the Mississippi River deltaic plain. The deltaic plain covers an area of approximately 29,785 km² (11,500 mi.²) (fig. 2), encompassing the active and abandoned deltas of the Mississippi River. The deltaic plain is a distinct physiographic unit, bounded on the west by the coastal marshlands of the Chenier Plain, on the east and south by the Gulf of Mexico, and on the north by the

Mississippi River Alluvial Valley and older gulfward-dipping Pleistocene deposits (Kolb and Van Lopik, 1958) (fig. 2). The most prominent physiographic feature of the deltaic plain is the vast expanse of fresh, brackish, and salt marshes. Natural levee ridges along active and abandoned distributaries, inland swamp, tidal channels, shallow lakes and bays, and sandy barrier islands and beaches comprise the majority of the remaining surface features.

FIGURE 2. Physiographic provinces of southern Louisiana. Numerous gulfward-dipping Pleistocene deposits underlie the untitled area north of the Deltaic and Chenier plains.

The modern alluvial valley and deltaic plain of southern Louisiana have been formed by progradation of the present and former Mississippi River courses and deltas. Each time the Mississippi River has built a major delta lobe seaward, it has subsequently abandoned the lobe in favor of a shorter, more direct route to the sea. These river course (meander belt) changes and accompanying shifts in centers of deposition have resulted in the present distribution of deltaic sediments along the southeast Louisiana coast. Marine transgression, caused in part by compaction and subsidence of deltaic sediments, begins soon after a delta lobe is abandoned. However, for much of the Holocene, the net result between the advancing deltas and the encroaching sea has been an overall increase in the size of the deltaic plain (Kolb and Van Lopik, 1966). During the last 50 years, coastal land loss has accelerated as human activities (particularly sediment trapping by the Mississippi River lock and dam system), combined with relative subsidence and erosion, have resulted in a net rise in sea level and loss of land.

Geologic History of the Mississippi River Deltaic Plain

The geologic history of the Lower Mississippi Valley and the Mississippi River deltaic plain has been determined from more than 30,000 borings and hundreds of radiocarbon age determinations. Some of the important contributions to the understanding of the history of the alluvial valley and the deltaic plain have been made by Fisk (1944, 1952, 1955), Fisk and McFarlan (1955), McFarlan (1961), Kolb and Van Lopik (1966), Frazier (1967), and Saucier (numerous papers, see Saucier (1994) for complete listing). Despite differences in calculations and interpretations of radiocarbon ages, information gained from these data indicates that over the past 8 k yr the Mississippi River has changed its course several times, forming a complex setting in which to observe the various aspects of fluvial and coastal sedimentation. (see Coleman and Roberts (1989) for a description of the Mississippi River alluvial valley and deltaic plain.)

During the last glacial advance, the Late Wisconsinan (about 26 to 12 ka; Martinson and others, 1987), continental ice accumulation caused sea level to be lowered some 90 m (295 ft) below its present level (Dillon and Oldale, 1978). As a result, the Louisiana shoreline was as far as 160 km (100 mi.) south of its present position (Kolb and Van Lopik, 1958). Lowered sea level led to the entrenchment of gulfward-flowing streams and their tributaries into the newly exposed deposits of the last interglacial Pleistocene Prairie Formation (Fisk, 1938). Entrenchment of the ancestral Mississippi River into the late Pleistocene Prairie Formation formed an alluvial valley with branching tributary valleys approximately 16- to 40-km (10- to 25-mi.) wide which trended southeast across the deltaic plain. (The Prairie Formation is not shown on fig. 3, but forms the surface unit updip of the Holocene alluvial/deltaic sediments)

Sea level was at its lowest position about 18 ka (Martinson and others, 1987) and subsequently began to rise as a result of glacial melting and regional subsidence of the coast (Kolb and Van Lopik, 1966; Nummedal, 1983). Streams alluviated the entrenched valley with coarse sediments in order to adjust to the rise in base level. As sea level continued to rise, deposition of coarse sediments was forced farther up the alluvial valley. Closer to the Gulf of Mexico, shallow marine sediments were deposited over coarse basal fluvial sediments as the shoreline transgressed northward. As sea level continued to rise, both the quantity and grain size of detritus supplied to the streams decreased, leaving only fine sands, silts, and clays for deltaic deposition (Kolb and Van Lopik, 1966).

Between 8 and 4 ka a stillstand of sea level occurred at approximately its present level (Nummedal, 1983). The Mississippi River began building a series of lobate deltas in a gulfward direction as a result of a stationary sea level, displacing Gulf of Mexico waters that had extended up the Mississippi River alluvial valley to the latitude of Baton Rouge, Louisiana (Kolb and Van Lopik, 1966). The Mississippi River and its associated deltas shifted several times during this gulfward growth of land, resulting in a deltaic plain composed of an active and several inactive deltaic complexes extending some 288 km (180 mi.) across southeast Louisiana. From oldest to youngest, the deltaic complexes are the Maringouin, Teche, St. Bernard, Lafourche, and the Plaquemine-modern (fig. 3). The relative ages of these complexes are well established, but the absolute ages are less accurate. Delta ages have been derived from radiocarbon analysis as well as archeological evidence (McIntire, 1958; McFarlan, 1961; Frazier, 1967; Törnqvist and others, 1996).

FIGURE 3. Major Holocene deltaic lobes of the Mississippi River (modified after Frazier, 1967): Maringouin, 6-8 ka; Teche, 5.8-3.5 ka; St. Bernard, 3.5-2 ka; LaFourche, 2-1 ka; Plaquemine-Modern, 1 ka to the present. Red dot marks St. Bernard Parish core locality SB1.

The earliest delta lobe, the Maringouin, probably prograded sometime between 6 and 8 ka, during a short stillstand when sea level was approximately 12 to 18 m (40 to 60 ft) lower than its present level (Frazier, 1967). Frazier (1967) mapped the Maringouin as the most extensive delta lobe in the Mississippi River deltaic plain. Erosion, subsidence, and burial by subsequent deltaic deposition have made it difficult to reconstruct the exact limits, location, and upvalley extent of the Maringouin delta complex. About 5.8 ka, the initial progradation of the Teche delta began in the western part of the Mississippi River deltaic plain (Frazier, 1967). Gradually, the major locus of Teche deposition shifted eastward depositing sediments in a southeastward direction. The Teche-Mississippi River System was actively depositing sediments in this area until approximately 3.5 ka when the primary flow of the Mississippi River shifted far to the east and continued building the St. Bernard delta. The Mississippi River continued to build the St. Bernard delta until sometime between 2 ka (Frazier, 1967) and 1.4 ka (Törnqvist and others, 1996), when flow was diverted westward and the Lafourche delta began to prograde seaward. Abandonment of the Lafourche course for the Plaquemine-Modern delta lobe occurred around 0.5 ka (Frazier, 1967) or as early as 1.3 ka as suggested by Törnqvist and others (1996).

Geologic History of the St. Bernard Marsh Locality

The St. Bernard core locality (N. Lat. 29°58'53", W. Long. 89°55'27") is in a brackish marsh on the southwestern edge of Lake Borgne. Sediments at this locality are part of the St. Bernard delta complex (fig. 4), which began receiving Mississippi River deltaic sediments in the middle to late Holocene

(about 4.7 ka, Frazier (1967); about 3.5 ka, Törnqvist and others (1996)). Based on data from borings, Dunbar, and others (1994) suggest that at this time the area was characterized by open marine water approximately 18 m (60 ft) deep. Homogeneous prodelta clays were the first deltaic deposits to enter the study area. As the Mississippi River and its distributaries prograded, interdistributary bay deposits began to be deposited. Interdistributary bay deposits include those sediments deposited in low areas between active distributary channels, usually under brackish water conditions. Sediment charged water during flood stage overflows the natural levees of the distributary channels, depositing the coarsest sediment (silt) near the channel on the natural levee flank. The finer sediment (silty clay and clay) is transported away from the distributary channel and settles out of suspension as interdistributary deposits. The 15 cm of interdistributary deposits at the base of the St. Bernard long core (designated SB1c) contains numerous fragments of highly degraded plant material. The early Holocene radiocarbon (^{14}C) age of $8,420 \pm 60$ yr BP (table 1) for charcoal in the sediment supports an interpretation as a fluvial deposit containing organic matter of various ages, but probably does not reflect the actual age of the deposit. Deposition of interdistributary clay and silt continued until the height of the deposit reached sea level, at which time the sediments were populated by marsh vegetation. Radiocarbon age determinations for the core SB1c confirm that marshes began to form in this part of the deltaic plain about $3,180 \pm 50$ yr BP (table 1) and have continued to exist relatively undisturbed to the present. Typical depositional environments characteristic of deltaic progradation are found adjacent to the study site. Natural levee, point bar, swamp, and abandoned distributaries located in the vicinity of the study site are shown on fig. 4A.

FIGURE 4A. Deltaic environments in the vicinity of the St. Bernard core locality; these are characteristic of progradation in this part of the Mississippi River deltaic plain. Vegetation associated with the environments of deposition: brackish marsh, predominantly *Spartina patens* (wiregrass) with minor amounts of *Scirpus olneyi* (three-cornered grass); natural levee and point bar, predominantly live oaks and water oaks, with minor amounts of pecan, hackberry, and bald cypress; swamp, predominantly bald cypress trees with minor amounts of tupelo gum, live oaks, water oaks, elms, hackberry, and sweet gum trees.

FIGURE 4B. Map showing St. Bernard core locality in relation to New Orleans, Chalmette, and the Mississippi River Gulf Outlet Canal (MR-GO canal). Fresh-to-brackish water is dark blue; Gulf of Mexico water is light blue. Names of water bodies are blue; parish names are brown.

Active deltaic deposition shifted away from the study site about 2 ka (Frazier, 1967), or about 3 ka as indicated by ^{14}C data for this study (see table 2). However, the area around the St. Bernard core locality continued to receive periodic deposition related to crevassing on the Mississippi River until the early 1900s when levees began to be built along the Mississippi River for flood protection. Marsh accretion due to organic and mineral matter accumulation is the dominant process presently active in the study area

Stratigraphy of the St. Bernard Marsh Locality

The uppermost 4 m of marsh sediment at the St. Bernard core locality was sampled using vibracore techniques (see section on Field Methods). The core SB1c was described in the COE, New Orleans District, soil test laboratory. Both measured and compaction corrected (in parentheses) depths are given in the description. Many of the samples analyzed for this study were from this core and are assumed to be representative of the marsh environment on the western edge of Lake Borgne.

St. Bernard core SB1c

0-55.0 cm (0-77.6 cm) peat; fibrous, plant fragments more identifiable toward surface; less compact toward surface; no apparent wood fragments; few very-small seeds; plant material appears to be reed and grass fragments; 0-7 cm, 10YR 2.5/1-5Y 3/1, 7-26 cm, 2.5Y 2.5/0, 26-55 cm, 2.5YR 2.5/0

55.0-69.0 (77.6-97.3 cm) peat; fibrous, more compact than overlying interval, compaction uniform throughout; no obvious wood fragments, numerous reed and grass; uniform 2.5R 2.5/0

69.0-82.5 cm (97.3-116.3 cm) peat; fibrous, same as above except color change to 5YR 2/1.5

82.5-86.5 cm (116.3-122.0 cm) clay; upper and lower contacts are irregular and abrupt; 5Y 3/1

86.5-121.0 cm (122.0-170.6 cm) peat; compact, fibrous, particularly odorous, peat; finer textured than in 0-82.5 cm, fewer plant fragments than in 0-82.5 cm; 7.5YR 2.5/1; abrupt to clear lower contact

121-140.5 cm (170.6-198.1 cm) peat; 121-124.7 cm very finely disseminated organics, smooth with few to no plant fragments, 7.5YR 2.5/0; 124.7-140.5 cm increasing number of fragments or fibers downward with identifiable changes (abrupt to clear contacts) at 134.5 cm and with the underlying organic clay.

140.5-162.5 cm (198.1-229.1 cm) organic clay; organic matter content appears to decrease downward to 162.5 cm; 140.5-146.0 cm, 7.5YR 2.5/0; 146-158.5 cm 10YR 3-2.5/1; 158.5-162.5 cm, 7.5YR 2.5/0; very finely disseminated organics throughout, very few fibers, clear lower contact

162.5-218.0 cm (229.1-307.4 cm) peat; fibrous, clay content increases upward, 7.5YR 2.5/0; 190.25-191.0 cm clay lens, 7.5YR 3/0; abrupt lower contact.

218.0-233.5 cm (307.4-329.2 cm) organic clay; indistinct layers, more organic and fibrous in basal 3.0 cm with some fibers throughout, bioturbated throughout; 7.5YR 3/0; clear to abrupt lower contact.

233.5-277.0 cm (329.2-390.6 cm) peat; very finely disseminated organics with only few discernible plant fragments, large 4 cm-wide x 6 cm-deep burrows at top; 0.5-0.75-cm thick organic clay at 249.0 cm; bioturbation and clay content increases downward in basal 20.0 cm; abrupt stepped lower contact (irregular); 7.5YR 3/0

277.0-282.0⁺ cm (390.6-397.6⁺ cm) gray clay; N4; small black organic blobs with plant material throughout.

REFERENCES

Return to CONTENTS

FIELD METHODS

L.D. Britsch and D.L. Dillon

Two or more cores were taken at each study site. Sampling technique(s) were determined by conditions at the site from the numerous techniques commonly used for sampling the saturated peats of the Mississippi River deltaic plain. The main considerations at each site were the degree of saturation of the substrate and the depth of sample. In the marshes, all access was by boat, and the water table was at or above the marsh surface.

At the St. Bernard locality, one long core, about 4 m in length, and two short cores, each about 1 meter in length, were extracted. All cores were taken using a vibracore sampler (fig. 5). This technique allows retrieval of continuous, undisturbed samples from unconsolidated saturated sediments. The vibracore works on the principle of liquefaction in fine-grained sediments by displacing sediment to allow passage of the core barrel (Smith, 1984). The effectiveness of the vibracore in relation to penetration and recovery is directly related to the engineering properties of the material being sampled. The vibracore works best in saturated clays, silty clays, silts, and fine sands, but is very inefficient in firm clays and medium to coarse sands. The vibracore system is portable allowing retrieval of relatively long (up to 10 m) undisturbed samples in marsh settings.

FIGURE 5. Photograph of vibracore equipment used to drill the marsh site in St. Bernard Parish, southeastern LA.

The vibracore equipment consists of a 5 horsepower gasoline engine connected to a hydraulic pump and a 15.24 m (50 ft) flexible hydraulic hose attached to a hydraulic vibrator head. For the short core the vibrator head was attached to a 7.62 cm (3 in) inside diameter, 1.27 m (5 ft) length of aluminum irrigation pipe by a quick release clamp. The bottom of the sample tube was cut on a 45 degree angle and sharpened to aid the penetration during sampling. To obtain the sample the pipe was vibrated into the marsh at a low frequency. After the pipe had reached its maximum penetration, the pipe was cut with a hacksaw approximately 0.3 m (1.0 ft) above ground surface. Compaction was then measured as the difference between the marsh surface and the top of the sample inside the pipe. A 7.62 cm (3 in) diameter packer was placed in the end of the pipe and tightened to create a vacuum preventing sample loss. Extracting the core was accomplished by digging a shallow trench adjacent to the sample tube, tilting the sample tube into the trench, and lifting the sample from the substrate. After retrieval, the sample was measured, the sample tube was wrapped in aluminum foil, secured with duct tape on each end, and labeled to indicate top and bottom and location. The same procedure was followed to obtain the longer core except that a 5 m (19.7 ft) section of irrigation pipe was used to obtain a greater depth of penetration. Sample tubes were transported to the laboratory in the horizontal position to minimize movement of material in the tubes.

REFERENCES

Return to CONTENTS

¹⁴C AGE DETERMINATIONS

J.P. McGeehin

Two cores from the St. Bernard parish core locality were analyzed for ¹⁴C resulting in 40 radiocarbon ages. Both cores were taken using vibracore techniques. Dating of the organic carbon sequestered in the cores provides chronostratigraphic constraints that can be used to estimate variations in rates of carbon accumulation for marsh environments of the Mississippi River deltaic plain. The longer of the two cores (SB1c) is approximately 4 m (compaction corrected) and composed of peat to 3.91 m depth. The lowermost 11 cm of core is a blue-gray interdistributary clay. Samples were selected for ¹⁴C dating based on visually observed stratigraphic breaks. The shorter core (SB1a) is about 1 m long, composed entirely of peat, and was sampled to analyze the peat for the ¹⁴C bomb spike. The "bomb spike" is a measure of elevated ¹⁴C activity as a function of the sharp rise in atmospheric ¹⁴C levels resulting from above-ground nuclear testing in the 1950s and 60s. The atmospheric ¹⁴C bomb peak occurred in the northern hemisphere between 1963 and 1964, although elevated levels of atmospheric ¹⁴C can be traced from the early 1950s to the present day (Levin and others, 1985; Burchuladze and others, 1989). A similar peak exists for ¹³⁷Cs, and it was hoped that the peaks for the two isotopes would match and provide a useful marker with which to measure very recent carbon accumulation rates.

Fifteen ¹⁴C analyses were obtained for samples taken from core SB1c. Most of these were run on bulk peat. Two samples were run on grass pieces with corresponding bulk peats at the same depth intervals. One sample was run on degraded peat fragments found in the basal interdistributary silt and clay. All samples were chemically pretreated to remove the inorganic carbon and loosely adsorbed humic acid fractions. The dried samples were then

processed to graphite according to the method described in Markewich (*in press*). The graphite was pressed into targets and sent to Lawrence Livermore Laboratory's Center for Mass Spectrometry (CAMS) for ^{14}C analysis. Splits of the chemically-pretreated samples were analyzed for $\delta^{13}\text{C}$ using stable isotope mass spectrometers at the USGS in Reston, Virginia, Denver, Colorado, and Menlo Park, California. The ^{14}C ages were adjusted accordingly to account for naturally occurring fractionation effects.

Data for the SB1c core, including ^{14}C ages, the corresponding fraction modern values, $\delta^{13}\text{C}$ values, and corrected versus uncorrected depths are shown in table 1. Analytical ages for the bulk peat samples range from approximately 200 to 3,200 yr BP. The sampling errors and the effects of environmental factors on radiocarbon age determinations, and the need to calibrate radiocarbon ages to available tree-ring data, are discussed in the last section of this report *Data Interpretation and Summary*.

TABLE 1. Data for ^{14}C age determinations for peat samples from core SB1c, St. Bernard Parish, LA.

Twenty-five ^{14}C analyses were obtained for core SB1a. The core was sampled at 2.54-cm (1 in) intervals for the entire one meter depth. The bulk of the bomb curve was determined to be above approximately 30 cm, so all the samples collected were not run. Because identifiable grass blades and (or) roots as well as more decomposed peat were found at some depths, for each depth interval, samples of both were selected for analysis. At the 25.9 cm interval, grass seeds were analyzed, in addition to the grass and peat. Possibly, when all the age data (isotope ages and microscopic stratigraphic markers) for the Mississippi River deltaic plain are available, some trend will be identifiable. Then, we may be able to ascertain which peat component is a most reliable age indicator.

TABLE 2. Data for ^{14}C age determinations for peat samples from core SB1a, St. Bernard Parish, LA.

All samples for core SB1a were pretreated and reduced to graphite. Sample results are shown in table 2. Note that the ^{14}C activity is expressed as D^{14}C in addition to the ^{14}C age in years Before Present (yr BP). This is because the ^{14}C activity in the samples affected by the bomb pulse is higher than the activity of the modern standard, resulting in a $>\text{MODERN}$ age. The $\text{Delta}^{14}\text{C}$ results are expressed as parts per thousand (per mil) values above the activity of the modern standard corrected to the current year from the "zero" year for ^{14}C of 1950. The $\text{Delta}^{14}\text{C}$ data for core SB1a have been plotted in figure 6. The 1963-64 atmospheric bomb carbon peak can be seen clearly at approximately 15 cm. This results in an accumulation rate of approximately 0.5 cm yr^{-1} from 1963-64 to 1996 when the core was taken.

Note that all ^{14}C ages are reported as standard radiocarbon years, in years before present (yr BP), where BP represents 1950, the zero year for radiocarbon dating. The data have not been calibrated for variations in atmospheric ^{14}C level over time as measured in tree rings. The calibration corrections, which can be made using one of several computer software programs, will be made for the data as efforts proceed to accurately estimate the rates and amounts of organic carbon accumulation in the Mississippi River deltaic plain.

FIGURE 6. ^{14}C bomb curve for core SB1a, St. Bernard Parish, LA. Data corrected for compaction and ^{13}C .

REFERENCES

[Return to CONTENTS](#)

^{137}Cs ANALYSIS

T.L. Fries and J.H. Robbins

Ritchie and McHenry (1984) give an overview of the extensive use of ^{137}Cs for measuring soil erosion and sediment accumulation in reservoirs and lakes. The time dependence of ^{137}Cs fallout ($t_{1/2}=30.2 \text{ y}$) resulting from above-ground testing of nuclear weapons in the 1950s and 60s makes it an excellent tracer for determining rates of sediment accumulation. Commonly a distinct ^{137}Cs peak can be detected by analyzing closely (1-2 cm) spaced core samples. This peak is associated with the maximum rate of ^{137}Cs deposition during 1963 and 64. Rarely, a minor peak can be identified. In the United States this is commonly a secondary peak associated with testing in 1959 (Callendar and Robbins, 1993).

In the Mississippi River deltaic marshes, where there has been only in-situ sediment (peat) production, measurement of ^{137}Cs was carried out to develop an estimate of a mean linear rate of peat accumulation over the last 35 to 40 years. Hand-pushed cores are preferable for ^{137}Cs isotopic analysis, however, at the St. Bernard core locality, we could not get the sample material to stay in the sampling tube when the tube was pushed by hand. Therefore, ^{137}Cs measurements for the St. Bernard core locality are from core SB1b, one of two short (approximately 1m-long) vibracores. The vibracore technique produced intact short cores with less than ten percent compaction. By using a meter long sampling tube to obtain a short vibracore, instead of sampling the top of a longer vibracore, we minimized (or eliminated) the potential problem of contamination of shallow material with water that has been in contact with material from greater depths.

The whole frozen core (SB1b) was cut into 2-cm sections on a band saw. Each saw cut consumed 1 mm of sample length and each 2-cm measurement was made from the edge of the previous cut. As each section was cut the sample and the resulting aluminum ring were transferred to a pre-weighed plastic bag already marked with identification information. Air was removed from the bag, the bag was sealed, and the side of the bag corresponding to the top of the sample was marked. The sample/ring/bag combination was then weighed and the weight recorded.

Upon completion of ^{137}Cs measurements the bags were split open and the entire sample/ring/bag combination was allowed to air dry. When the samples reached constant weight the aluminum rings were removed, cleaned, dried and weighed. A dry sample weight was then calculated for use in ^{137}Cs calculations. Dry sample material was then archived.

Samples from the SB1b core were analyzed by placing the sample/ring/bag assemblage directly on a gamma counter. Each 2-cm section was counted for up to one day and data were recorded for both ^{137}Cs and ^{40}K . ^{137}Cs (661.6 KeV) and ^{40}K (1460.7 KeV) were determined by gamma spectroscopic analyses using a high resolution lithium-drifted planar (GeLi) detector coupled to a multichannel analyzer. Calibration was achieved by doping comparable materials with precisely known amounts of NIST-traceable standard solutions of radiocesium and weights of KCl. Detection limits for ^{137}Cs and ^{40}K were 2x counting error in decays per minute per gram (dpm g $^{-1}$). These detection levels account for most of the fractional counting errors of 1.5 -5 percent for the peat (table 3).

A plot of ^{137}Cs count rates (decays) versus depth (fig. 7) shows a characteristic shape with a peak normally associated with the years of maximum fallout, 1963-1964. Because a large portion of the shallow material from this core was discarded in the field and another section was lost during preparation it was not possible to accurately model and date deposition throughout the core. A date of 1963.5 can probably be assigned to the maximum ^{137}Cs count rate. Maximum ^{137}Cs activity was found at a compaction corrected center-of-section (midpoint) depth of 16.0 cm. Since the ^{137}Cs peak could be located anywhere in the sample interval, and the compaction corrected length of this interval is 2.2 cm, the uncertainty in the depth estimate is approximately 14 percent (about ± 1.1 cm).

The ^{137}Cs curve (fig. 7) for core SB1b is a reasonable match for the ^{14}C bomb spike curve (fig. 6) for core SB1a. Based on the ^{137}Cs data the average linear accumulation rate at the SB1b site since 1963-64 is estimated to be 0.50 \pm 0.04 cm yr $^{-1}$. This is the same rate indicated by the ^{14}C bomb spike data.

FIGURE 7. ^{137}Cs curve for core SB1b, St. Bernard Parish, LA.

TABLE 3. ^{137}Cs data for bomb-curve analysis of core SB1b, St. Bernard Parish, LA.

REFERENCES

[Return to CONTENTS](#)

BIOCHRONOSTRATIGRAPHY

J.H. Wrenn, J.B. Pracht, C.M. Fraticelli, and B.M. Samuel

Potentially Useful Microscopic Stratigraphic Markers

The biochronostratigraphic portion of the USGS Mississippi River Basin Carbon Project seeks to use stratigraphic markers, in conjunction with the historical record, to provide relative and (or) absolute ages for core samples from marsh, swamp, lake, and floodplain environments. For cores from St. Bernard Parish, Louisiana (figs. 1, 4A-B), the ultimate purpose is to provide chronological control for sediments deposited between calendar year 1750 and 1950 (only the year, not A.D., is used herein). Potentially useful markers include: pollen grains, plant phytoliths, charcoal fragments, carbonaceous spherules, fungal spores, and lithologic storm indicators. The targeted era of the MBCP includes the colonial period (1717-1803) and the Americanization period (1803-1900) of Louisiana. The area around the St. Bernard core locality includes New Orleans, which was settled early and extensively. The historical documents of this era provide a detailed history with which to correlate sedimentary deposits. Historical syntheses, maps, personal journals, memoirs, and census data provide information on past patterns of land use, regional geography, and farming of the region. These records document several events which have produced chronologically significant markers. The events and their potential stratigraphic markers are discussed below and outlined in table 4 in the order of their occurrences.

TABLE 4. St. Bernard Parish, Louisiana cores - events and potential stratigraphic markers.

Heliophyte pollen. --The earliest dated event that would produce strategic stratigraphic markers for the colonial period is the settlement of New Orleans in 1717 by French colonists. As the native forest was felled for building material and to make room for agricultural fields, the thinning and/or removal of the canopy should be reflected in the sediment record as an increase in the pollen of sun-loving plants, heliophytes, especially members of the Compositae family (Whitehead and Sheehan, 1985).

Cultigen pollen and phytoliths. --Commonly in the historical sediment record, the increase in Compositae pollen is followed by the appearance of non-indigenous agricultural crop pollen. In the St. Bernard area, sugarcane (*Saccharum officinarum*), rice (*Oryza sativa*), cotton (*Gossypium hirsutum*), potatoes (*Solanum tuberosum*), sweet potatoes (*Ipomoea batatas*), and peas and beans (Leguminosae family) were introduced and cultivated. Corn (*Zea mays*) is indigenous to the Americas; however, European colonists adopted corn and expanded its cultivation.

Phytoliths, in conjunction with pollen, can be used to achieve higher taxonomic resolution within the Gramineae fossils. Sugar cane and corn were the major cash crops cultivated in the St. Bernard area (Menn, 1964). Pollen of sugar cane and corn are not readily distinguishable. They are both members of the Gramineae family, a family whose taxa produce similar pollen. However, Piperno (1988) suggests that some taxa of this family may be differentiated based on phytolith morphology.

Sporomiella spores. -- Domesticated livestock came with European settlement and cultivation. Davis (1987) identified the fungus *Sporomiella* as a palynological marker for determining the arrival of these animals. *Sporomiella* is commonly associated with the dung of herbivores. Although *Sporomiella* spores have been found in some prehistoric sediments, their abundance increases following the introduction of domesticated grazing animals. The first census of New Orleans, taken in November 1721, records 36 cattle, nine horses, and no pigs or sheep (DuFour, 1968), therefore, *Sporomiella* spores may be present or shown an increase in the sediments of that period.

Charcoal layers. -- Charcoal layers can be associated with historical events, and, in association with other stratigraphic markers, may be chronologically significant. In 1788, and again in 1794, when virtually all structures were wood, New Orleans was heavily damaged by fire. The 1788 fire destroyed about 75 percent of the city, (de Rojas, 1937). The fire of 1794, although not as destructive as the 1788 fire (Arthur, 1937), destroyed one third of all residences (Fortier, 1904). As a result of fire, particulate charcoal would have blanketed the surrounding landscape, and runoff would move charcoal from the burned city into small waterways that feed the estuaries, where it would be incorporated into the sediment record.

Carbonaceous spherules. -- Carbonaceous spherules are indicators of urbanization and industrialization. The spherules are produced when resinous wood, oil, or coal are incompletely burnt and the volatile components vaporized, leaving a microscopic spheroidal skeleton of elemental carbon (McCrone and Delly, 1973). Following their release to the atmosphere via chimneys and smokestacks, carbonaceous spherules become incorporated into the sedimentary record.

Decrease in Taxodium distichum (bald cypress) pollen. -- The early settlers of Louisiana quickly learned that *Taxodium distichum* (bald cypress) wood resists decay, making it ideal for construction and for use in shipping containers. Although the technological infrastructure required to log Louisiana's extensive cypress swamps developed slowly, by 1890 there was industrial-scale logging operations. By 1925, the cypress swamps were so heavily denuded that commercial logging was no longer cost efficient (Norgress, 1947). The result was also an abrupt decline of *Taxodium distichum* in the sediment record.

Hurricane layers. -- Several hurricane landfalls in Louisiana have been documented in the historic records of the study area. Category 4 or 5 strength hurricanes on the Saffir-Simpson scale leave distinctive sedimentary strata (Liu and Fearn, 1993). Although the Saffir-Simpson scale was not developed until 1977, eyewitness descriptions indicate that some of the early hurricanes approached or achieved these intensities. For example, the hurricane of September 11, 1722, destroyed every home in the fledgling village of New Orleans (DuFour, 1968). In addition to stratigraphic signatures, hurricane surges can deposit open marine microfossils in coastal zones. Both the strata and the microfossils can be diagnostic stratigraphic markers.

Sample Processing

To examine the St. Bernard sediment record for the historic events, each of the potentially diagnostic microscopic stratigraphic markers had to be isolated. The isolation of these markers required four different processing techniques. Details of these processing procedures are given in Markewich (in press). The samples processed for each analysis are listed in table 5.

Pollen grains and fungal spores were isolated from twenty-two, 0.9 ml core samples according to the methods of Faegri and Iversen (1964). Sediment samples were processed to recover dispersed phytoliths by oxidizing organic matter and dissolving carbonate using procedures developed in the CENEX laboratory at LSU. Reference phytoliths were prepared from plant samples by heating in a furnace to reduce the organic matter to ash according to standard processing procedures. Charcoal and opaque carbonaceous spherule analyses (OCS) were conducted on the same sample preparations and slides. These were prepared by acid digestion to remove mineral matter and oxidation to remove all organic matter.

TABLE 5. Sample depths and analyses for microscopic markers in core SB1c, St. Bernard Parish, LA.

Pollen

Data. -- Absolute pollen counts of selected taxa observed in the SB1c core are presented in figure 8. In the interest of clarity, rare and chronologically/environmentally non-diagnostic taxa were omitted from the diagram. Seven zones were defined in the core and are described in table 6, from zone 1 (oldest) at the base of the core to zone 7 (youngest) at the top of the core.

FIGURE 8. Pollen diagram for core SB1c, St. Bernard Parish, Louisiana.

FIGURE 9. Photographic plate of diagnostic pollen and other microfossils for core SB1c, St. Bernard Parish, LA. A. TCT (Taxodiaceae/Cupressaceae/Thuja); B. Chen-Am (Chenopodiaceae-Amaranthaceae); C. *Typha latifolia*; D. Gramineae; E. Cyperaceae; F. Compositae (high spine); G. *Vigna luteola*; H. foram test lining.

TABLE 6. Discussion of pollen zones depicted in Figure 8.

Discussion of the Pollen-Defined 1700 to 1900 Sediments. -- Based on their pollen assemblages, Zones 5 and 6 are considered to represent the time intervals 1350 to 1890, and 1890 into the mid-twentieth century, respectively (uncalibrated conversion from ^{14}C ages to calendar years). These zones correspond to the radiometrically undatable zone targeted by this research project. Although some of the predicted stratigraphic markers were observed in core SB1c, others were not. In addition, unanticipated markers made substantial contributions to the interpretation of Zones 5 and 6. Four of the strategic markers are discussed in this section.

1. As predicted, a rise in Compositae pollen served to delineate the pre-historic from the historic sediments.

----1a. The appearance of *Vigna luteola* in this zone, unexpectedly, confirmed this interpretation. *Vigna luteola* is a legume which originated in central Africa and was long cultivated throughout Africa, south Asia, and the Mediterranean. It was introduced into the New World via the Spanish West Indies and diffused from there. As early as 1714, *Vigna luteola* was being cultivated in North Carolina (McKee, 1948, p. 725).

Because of its Old World origin, the appearance of *Vigna luteola* provides a maximum age of contact (1492), however it is more probable that the legume was introduced into the study area after settlement began in 1717.

FIGURE 10A. Photograph of 1845 map showing the historical geography of Mississippi River from just east of New Orleans to the bird's foot delta (La Tourrette, 1845). The St. Bernard core locality is indicated by the white circle.

FIGURE 10B. Photograph of 1845 map showing: (1) position of St. Bernard Parish core locality (white circle) on Conseil Plantation lot, and (2) the railroad used for transportation of agriculture and wood products. This is a close up of part of map in figure 10A (La Tourrette, 1845).

Core SB1c was taken from land which was once part of the Conseil Plantation of the Villeré family in a region historically rich in agricultural products (fig. 10). General Jacques Philippe Villeré, the founder of Conseil, was Louisiana's second governor under the American regime and the first native born governor of the state (Villere, 1981). Due to General Villeré's prominent social status, many records exist concerning his public and private life. In 1828, Karl Bernard, the Duke of Saxe-Weimar-Eisenach, visited and described the Villeré property.

"Back of the elegant mansion-house stand the negro cabins, like a camp, and behind the sugar-cane fields, which extend to the marshy cypress woods about a mile back, called the cypress swamp... The whole [of the living quarters] is surrounded by cane fields, of which some were brought in, and others all cut down. A field of this description must rest fallow for five years and be manured, before being set out again in plants. For manure, a large species of bean is sown which is left to rot in the field, and serves the purpose very well." (Bernard, 1828)

TABLE 7. Selected Entries of the 1860 Census of the Large Slave holders of Louisiana (Menn, 1964).

An 1860 census of the large slave holders of Louisiana records the products of the plantations in the area of St. Bernard core locality (Menn, 1964). The census included the Conseil Plantation, which, in 1860, was under the management of Edmon Villeré, General Villeré's grandson (table 7). The data provide an index of types of pollen to be expected in the core.

No sugar cane pollen was indentified in sediment from the St. Bernard cores. Sugarcane (*Saccharum officinarum*) is a member of the Gramineae family. Its pollen is very similar to that of other grasses. If sugar cane pollen grains were present in the core, they were probably grouped in the general Gramineae category. The fact that sugar cane rarely flowers in Louisiana (it is replanted from cuttings) suggests that sugar cane pollen would not be common in the marsh sediment

A search of the historical literature of the study area revealed that *Vigna luteola* can probably be considered a "pollen proxy" for sugar cane. In Bernard's journal entry of his visit to the Conseil Plantation, he describes General Villeré's practice of restoring nitrogen to his sugar cane fields with "a large species of bean" which was sown and left to rot in the fields. Literature on southern agriculture of this period indicates that this bean was likely *Vigna luteola*, a cowpea (Gray, 1941, p. 824; Sitterson, 1953, p.126).

As discussed above, *Vigna luteola* makes its appearance in Zone 5, and although never abundant, its presence reflects the cultivation of sugarcane in the area. Because *Vigna luteola* is insect pollinated (Mackie, 1946), and its pollen is not produced in large quantities, it is unlikely that its pollen would have been deposited at the core site without the Villeré's practice of letting the peas rot in place, a practise not common in Louisiana until the 1840s (Sitterson, 1953). The association between *Vigna luteola* and *Saccharum officinarum* provides a new and distinctive marker for the cultivation of sugarcane in Louisiana, the southeastern United States, and possibly the Caribbean Islands.

-----1b. The census of 1860 records Indian Corn (*Zea mays*) as Conseil's second largest crop, yet there is no pollen evidence of this cultigen in the St. Bernard core. Although *Zea mays* produces 2,500 pollen grains per anther, the grains are large and not very buoyant in the air (Lewis and others, 1983). As a result, *Zea mays* pollen typically settles out within 0.5 km from the parent plant (Purseglove, 1972). To increase the probability of finding the rare, and typically large, pollen grain of cultigens in pollen preparations, a sieving and large fraction scanning (LFS) method was developed by Gish (1994). Although we conducted LFS, no *Zea mays* pollen was recovered. The 0.5 km travel distance of *Zea mays* pollen is much less than the 2 -3 km distance between the Villeré corn fields and the core locality and probably explains why no *Zea mays* pollen was found in the St. Bernardcore.

Given the placement of the St. Bernard core on the plantation, it is an enigma why only one cultigen pollen was recovered from the core sediment. We hypothesize that the taphonomy of the expected cultigen pollen (i.e. pollen production, dispersal,deposition, and fossilization) did not allow widespread pollen dispersal.

2. Despite the region's 1860 domestic herbivore population (Table 6), *Sporomiella* spores were not positively identified in core SB1c. The absence is probably due to location of the core locality in the deltaic marsh, at the back of the Villeré property, beyond the cypress swamp where the plantation's domestic herbivores were not likely to roam.
3. A decline in *Taxodium distichum* (bald cypress) pollen was predicted based on (1) the intensive logging of this species between 1890 and 1925, and (2) census records and accounts from the Villeré plantation suggesting that clearing for row crop cultivation was common in the early 1800's in the area around the St. Bernard core locality. A decline is seen in the TCT (Taxodiaceae/Cupressaceae/Thuja) category in Zone 6. Although the pollen of the TCT category are difficult to distinguish from one another, the decline of TCT pollen in this area is being attributed to the historically documented decline of *Taxodium distichum* due to logging, most likely in the mid1800's.
4. No hurricane sediment layers were noted in the SB1c core.

Phytoliths

Introduction. --Phytoliths are deposits of opaline silica (SiO₂) or calcium oxalate secreted in or between plant cells, usually as a normal part of growth (Piperno, 1989; Jones and Bryant, 1992). Phytoliths of either composition may exhibit distinctive morphologies (Piperno, 1989). This study is concerned only with siliceous phytoliths (hereafter referred to as "phytoliths").

Phytolith taxonomy is complicated. The leaves, stem and roots of many plants may produce quite different looking phytoliths. In addition, similar looking phytoliths may be produced by different plant species, genera or even families. This contrasts markedly with pollen or spores, a single morphology of which is typically characteristic of a particular plant genus, or even species, and carries the name of that plant (at least with regards to extant plants).

Another complicating factor is that relatively few plants have been processed to determine what types of phytoliths they actually produce. This is especially true of Old World domesticated plants (Piperno, 1988). It is important to have a phytolith reference collection of plants in a study area with which to compare dispersed phytoliths recovered from sediments.

For the above reasons, phytoliths are often designated by descriptive morphologic terms (for example, circles, wagon wheels, bi-spinose rods, and so on) rather than by binomial names of the Linnean classification system. A morphologic designation is used until a correlation between a phytolith morphologic type and a particular plant taxa is made, permitting the name of that plant to be assigned to that phytolith morphologic type. Most forms reported here are referred to by informal morphologic designations.

FIGURE 11. Phytoliths and other siliceous microfossils diagram for core SB1c, St. Bernard Parish, LA.

No phytoliths of cultigens (for example, corn, sugar cane, cow peas, and so on) have been identified with certainty in the St. Bernard core sediment. Despite the negative results, the analytical results are discussed because of the potential in using phytoliths as strategic stratigraphic markers.

Results. --Some of the dumbbell or rod shaped phytoliths attributed to grasses may well have been produced by sugar cane. Abundant and diverse phytoliths were recovered from all samples processed. (See " Processing for Strategic Microscopic Stratigraphic Markers " in Markewich [*in press*] for the processing procedures used.) Commonly, a non-terrestrial siliceous component consisting of diatoms, sponge spicules and rare silicoflagellates were present. All elements were included during counting to characterize the entire siliceous residue from each sample. In addition, inertinite (charcoal) was counted along with the siliceous microfossils as a possible indicator of fire. Absolute counts of phytolith types observed in core SB1c are presented in figure 11 and table 8.

Most phytolith types recovered during this study have not been attributed to a specific plant taxa. Specimens of modern marsh plants are being processed to build a comparative phytolith collection with which to make such attributions (See discussion of the "CENEX Phytolith Reference Collection" in (Markewich, 1997). The most abundant phytoliths recovered are the various types of rods and irregular forms, though in some samples sedge (Cyperaceae) and saddle phytoliths are numerous. Dumbbell, biserial and uniserial spinose rods, non-spinose rods and saddle shaped phytoliths have been attributed to grasses based on studies in other areas and environments (Mulholland and Rapp, 1992; Twiss, 1992; Twiss and others, 1969). Marsh grasses probably produced many of the forms recovered from the St. Bernard core samples, though sugar cane, itself a grass, also produces dumbbell phytoliths (fig. 12) similar to those of other grasses. It is not possible at this time to determine if any of the dumbbell phytoliths in the core samples were produced by sugar cane or if all were produced by marsh grasses.

FIGURE 12. Diagnostic phytoliths and other siliceous microfossils present in core SB1c, St. Bernard Parish, LA. A. Dumbbell, produced by various grasses; B. Biserial spinose rod; C. Non-spinose rod; D. In situ dumbbell phytoliths in a modern sugarcane fragment; E. Cyperaceae (type); F. Cyperaceae (type), brownish coloration does not show but is probably due to burning from fire; G. centric and pennate diatoms.

TABLE 8. Phytoliths and other siliceous microfossil data for core SB1c, St. Bernard Parish, LA.

Phytolith morphologies that can be attributed to particular plant groups include:

GRASS:

dumbbell, cross, saddle, circle, biserial spinose rod, uniserial spinose rod, non-spinose rod (Mulholland and Rapp, 1992; Twiss, 1992). Non-spinose rods also are produced by some cereals, such as wheat (Kaplan, Smith, and Sneddon, 1992). The core locality is in an area that has been a marsh on the margin of a cypress swamp at least since European settlement; therefore, wheat has probably never been grown in the area.

SEDGE (Cyperaceae):

polygonal plates with a verrucate surface and satellite bumps, or rounded and smooth hat-shaped forms. (Some samples contained brown or grey phytoliths that indicate the sedges were burned.)

PALM:

spheres (some)

CRYSTOPHYTES:

cysts

UNKNOWN:

spheres (some), irregular shapes, wagon wheels

Charcoal

The concentration of charcoal in sediments can be used in conjunction with other microstratigraphic marker data to refine the chronostratigraphy

of the cores, particularly in reconstructing the history of natural fires, for example, (Terasmae and Weeks, 1979) or man-made fires (Iversen, 1941; Tolonen, 1978). The latter include those ignited for land clearing, the sacking of cities, industrial activity, and by accident. (See Tolonen (1986) for an overview of charcoal analysis.) Potential contributors of particulate charcoal in the St. Bernard cores include: natural marsh fires; burning of agricultural fields; clearing of the cypress forests; wood burning sugar mills, river boats and trains; major fires in New Orleans; military action during the War of 1812 and the Civil War; and redeposition of old carbon (inertinite). Data for core SB1c are in table 9 and figure 13.

FIGURE 13. Concentration diagram of charcoal and carbon spherules from SB1c, St. Bernard Parish, LA.

Organic Carbonaceous Spherules

Introduction. --One of the particulate emissions produced by the high temperature combustion of oil, coal or resinous woods is organic carbonaceous spherules (OCS), commonly composed of elemental carbon (Rose, 1990). OCS can be transported considerable distances, depending on their aerodynamic characteristics and local weather conditions at the time of burning. Because they are inert, they are preserved and accumulate in sediments (Renberg and Wik, 1985). Although OCS appear similar when viewed with a transmitted light microscope (fig. 14A), OCS produced by oil (fig. 14B), and those produced by wood or coal (fig. 14C), generally can be distinguished from one another using scanning electron microscope (Griffin and Goldberg, 1979).

OCS have been used to document the geographic and stratigraphic distribution in sediments of atmospheric pollutants generated by burning fossil fuels (Griffin and Goldberg, 1981; Renberg and Wik, 1985; Rose, Harlock, and others, 1995). They also have been used stratigraphically, in conjunction with palynology and ^{210}Pb radiometric dating, to provide relative ages of sediments, such as intertidal sediments in eastern Long Island Sound (Clark and Patterson III, 1984). OCS and historic records are used in this study to mark the advent of coal and oil burning, and to refine age estimates of the deposits.

The numerous sugar cane plantations in the St. Bernard area in the 1800's used railroad lines as well as wagons for transportation of goods. The core locality is within 7 km of both the old line of the Mexican Gulf Railroad (built in 1853) and the Mississippi River (figs. 10A, 10B), both were sources of OCS. Similarly, the steam powered sugar-processing mills on the plantations, and the various river and ocean-going steamboats that plied the Mississippi River were all generating OCS. The type of OCS produced depended on the fuel being used - wood, coal, and (or) oil; the type produced changed through time.

Coal OCS. -- The first coal powered steamboat to arrive in New Orleans, the *New Orleans*, docked in 1814 (Hunter, 1949), however, before the 1850's steamboats on the Lower Mississippi rarely burned coal, and then it only on the down river leg. Wood was burned on the return, up river voyage because it was cheaper and readily available (Hunter, 1949). Wood was the major steamboat fuel until well after the Civil War.

The first coal barges reached the New Orleans area in 1829; their contents consigned to the Labranch Plantation (Fossier, 1957, p. 25) to power the plantation sugarmill. This event provides the maximum age for coal-generated OCS in the area.

Coal became widely used in the lower valley by the 1880's for sugar mills, steamboats and steam locomotives. It was the dominant fuel used in all ocean going steamships, and merchant or naval vessels, that visited New Orleans until the beginning of the 20th century (Milton, 1953). Even by the beginning of the Second World War, 60 percent of all oceangoing steamships were still coal powered (Milton, 1953).

Oil OCS (also called liquid fuel, crude petroleum, or petroleum residue). --Oil was little used, alone or mixed with crushed coal, on merchant or warships before 1900 (Beig, 1906). It was well into the first decade of the 20th century before navies and merchant marines began switching to oil. Laminated OCS (fig. 9B), produced by the incomplete combustion of fuel oil (Griffin and Goldberg, 1979) provide a horizon marker for the first decade of the 20th century.

TABLE 9. Carbonaceous spherule data for core SB1c, St. Bernard Parish, LA.

Discussion of OCS data. --The lowest verified occurrence of OCS is at 85 cm. These are wood or coal OCS that probably mark the advent of high temperature wood or coal burning in the area. The maximum age for OCS at 85 cm, if produced by burning coal, could be 1814 when the first steamboat arrived in New Orleans; it was coal burning (Hunter, 1949). However, significant steamboat traffic did not develop in the Lower Mississippi Valley until the 1820's, and these were wood burners until the 1850's. If the OCS at 85 cm were produced by the burning of coal, the maximum age would more likely coincide with the arrival of the first coal shipment to New Orleans in 1829 (see *Coal OCS* above). The appearance of oil OCS at 56 cm suggests a maximum age of 1900-1910 (Milton, 1953; Gardiner and Lambert, 1992).

If the first appearance of coal OCS was around 1830, and a linear rate of accumulation is assumed, then the rate of organic matter accumulation in the marsh from 1830 to 1900 was about 0.4 cm yr^{-1} . If a linear rate is assumed, then from 1900 to 1996 OCS data suggest a rate of about 0.6 cm yr^{-1} . These rates bracket the 0.5 cm yr^{-1} rate suggested by ^{14}C (core SB1a) and ^{137}Cs (core SB1b) data (tables 2 and 3) for the period from 1964 to 1996.

FIGURE 14. Carbonaceous spherules present in core SB1c, St. Bernard Parish, LA.

Summary/Conclusions for Microscopic Stratigraphic Markers

Microscopic stratigraphic markers were used to subdivide the segment of core SB1c that covers a time interval no older than 1717 to about 1900, from 85 to 55 cm depth (fig. 13). Diagnostic stratigraphic markers were also used to achieve chronological resolution within this sediment segment.

FIGURE 15. Composite diagram of diagnostic microfossils from core SB1c, St. Bernard Parish, LA.

1. Carbon isotope ages and the presence of diagnostic microscopic stratigraphic markers suggest that the founding of New Orleans in 1717

occurred sometime during deposition of sediments between 106 and 85 cm depth. A ^{14}C age of 550 ± 50 yr BP at 106 cm (about 1350-1450 in uncalibrated conversion from ^{14}C ages) (table 1) provides a lower age limit. The age estimate at 85 cm is interpreted from the pollen and OCS data which indicate that this level dates to about 1830 (tables 5 and 8).

The relative age of about 1830 at 85 cm was determined by three stratigraphic markers. First, the appearance of *Vigna luteola* at 85 cm provides a maximum age equal to that of settlement, 1717. The probable age of sediment at 85 cm is about 1830. This is based on records of the Conseil Plantation which describe the use of *Vigna luteola* for fertilizing the plantation fields date from the mid 1820s. Second, the estimated number of OCS cm^{-3} at 85 cm is too few to plot on figure 13, however the first appearance of coal OCS at 85 cm suggests an 1829 maximum age for this level, coincident with coal being burned in area sugar mills. And, third, the beginning of the increase in Compositae pollen at 85 cm reflects land clearing activity and is consistent with the *Vigna* and coal spherule record.

2. The peak of wood/coal OCS at 71 cm provides the second chronological horizon within the targeted sediments and is attributed to wood and coal combustion in mills, steamboats, and steam locomotives during the mid 1800's, probably between 1860 and 1880. The drop in TCT, probably *Taxodium distichum* (bald cypress), at this level could indicate a later date of 1890, the period of large-scale regional cypress logging. The charred Cyperaceae phytoliths and abundant charcoal at this level suggest that this is a fire horizon, probably associated with cypress logging. Since land clearing in the area of the St. Bernard core began in the late 1700's and continued through the 1800's, the drop in TCT pollen probably documents accelerated local cypress logging in the mid 1800's rather than the 1890 regional increase in logging activity.
3. The uppermost identified horizon within the targeted sediments is at 56 cm. It is identified by the appearance of oil produced OCS, which came into common use around 1900. This is the transition period from steam to internal combustion engines.
4. The rise in aquatic taxa above 56 cm indicate a greater water to land ratio in the upper sediments of the core. This aquatic characteristic is in contrast to sediments below 56 cm which have more terrestrial components. This shift is likely due to the removal of the cypress trees, the roots of which acted as sediment traps, and to the construction of the MR-GO and other canals through the marshes.
5. Finally, an important diagnostic marker in the St. Bernard SB1c core is *Vigna luteola*. It is important not only as a stratigraphic marker, but also as a proxy for sugar cane cultivation. This is particularly relevant given the low flowering frequency of sugarcane in Louisiana. Using *Vigna luteola* as a proxy for sugar cane may prove to be an important new palynological tool for dating historical deposits in Louisiana, the southeastern United States, and perhaps the Caribbean area.

REFERENCES

[Return to CONTENTS](#)

COMPOSITIONAL ANALYSIS

T. L. Fries, G.R. Buell, and H.W. Markewich

Chemical and Isotopic Analysis and Carbon Fractionation

We analyzed the chemical composition of the peat in order to determine (1) the relative content of organic carbon (OC) and inorganic constituents, and (2) track the relations between relative percents of organic and inorganic constituents to climate and (or) human activity. *Chemical and Isotopic analysis.* -- Samples from each of the three St. Bernard cores were analyzed for bulk density (BD) (tables 3, 10, and 11). Cores SB1a and SB1c were analyzed for (a) total carbon and inorganic carbon by a UIC CO_2 coulometer, equipped with a furnace for the determination of total carbon and an acidification module for the determination of carbonate carbon; (b) OC by difference; (c) total N, ^{15}N , total C, and ^{13}C by a VG Optima elemental analyzer/isotope ratio mass spectrometer system; and (d) inorganic elemental analysis (tables 10B and 11B) by extracting approximately 100 mg of air dried sample with concentrated nitric acid. This procedure provides a near total extraction for many elements including Cu, Pb, Zn, Mn and Fe (Luoma and Bryan, 1981; Graney and others, 1995). Elemental concentrations were determined using a Perkin-Elmer 6000 inductively coupled plasma mass spectrometer (ICP-MS).

Carbon fractionation. -- The total OC did not exceed 40 percent for any sample; values were commonly between 25 and 35 percent (tables 10A, 11A). Carbon fractionation procedures were designed to provide data that might allow differences in organic matter composition, and associated inorganic components, to be detected and related to transport or in-situ production. As previously mentioned, detailed procedures are described in Markewich (*in press*).

The limited amount of sample material available, and the high percentage of the total sample in the low density fraction, made it impossible to carry the procedure beyond the initial density fractionation for most samples. For the available sample size only those samples with less than 10 percent total carbon, dry weight basis, contained sufficient high density material for further fractionation. With the exception of the sample of interdistributary clay from the base of the SB1c core, the high density material from these low carbon samples contained less than 10 percent of the total carbon in the sample. The low-carbon samples were from sections of the core that had been visually described as clay. The low carbon content and small sample size made further fractionation impossible.

Summary and Interpretation of Compositional Data

Data for the SB1a core show shifts in organic C (OC), total N (N), ^{15}N , bulk density (BD) and carbon storage (CS) at two depth ranges (table 10A).

OC and N decrease; BD increases. These shifts are also accompanied by shifts in a number of inorganic constituents including increases in aluminum (Al), iron (Fe), lithium (Li), lanthanum (La), cerium (Ce), thorium (Th) (table 10B), and calculated percent mineral matter (table 10A).

Some samples from the uppermost meter of core SB1c were analyzed for OC, N, and BD (fig. 16). Although values for the two cores are similar at comparable depths, the fine-scale variability present in the uppermost meter of peat is not adequately defined by the limited data from SB1c.

FIGURE 16. Organic carbon, nitrogen, and bulk density profiles for cores SB1a and SB1c, St. Bernard Parish, LA.

FIGURE 17. Elemental chemistry profiles for cores SB1a and SB1c, St. Bernard Parish, LA: (A) aluminum, (B) iron, (C) manganese; (D) lithium, (E) sodium, (F) potassium; (G) calcium, (H) magnesium, (I) strontium; (J) arsenic, (K) cadmium, (L) lead; (M) copper, (N) nickel, (O) zinc; (P) vanadium, (Q) chromium, (R) cobalt, (S) lanthanum, (T) cerium, and (U) thorium.

Chemistry. --The first shift in constituents occurs from 18-12 cm. Many elements show peak concentrations at about 15 cm. ^{137}Cs data indicate that sediment at 16 cm in core SB1b (fig. 6, table 3) was the surface in 1963-64. Assuming an accumulation rate of about 0.5 cm yr^{-1} , the 18-12 cm interval in core SB1a is assigned an age range of 8 to 10 years, representing most the 1960s. This interval corresponds in time to the construction of the Mississippi River-Gulf Outlet (MR-GO) Canal (fig. 1), 1958-1968 (US Army COE, 1975). Higher concentrations of metals, such as nickel (Ni), copper (Cu), chromium (Cr), vanadium (V), and cobalt (Co) (figs. 17E, F) are coincident with the ^{137}Cs and ^{14}C peaks (figs. 6, 7). However, they are probably not related. The higher (albeit still low) trace metal concentrations are more likely related to a decrease in OC accumulation due to dry climatic conditions (Cook and others, 1997) or possibly to local emissions from industrial sources and (or) to construction of the MR-GO Canal. The higher concentrations of crustal elements such as Al, Fe, manganese (Mn), Ce, La, and Th in this 18-12 cm depth interval support a natural rather than an anthropogenic source, suggesting that the increase in inorganic components is due to a decrease in peat production during a dry climatic interval. It is also possible that the increase in inorganics is due to a large-scale Mississippi River flood or to a local disturbance, such as dredging for the MR-GO canal. Disturbance due to hurricane activity could also result in an increase in crustal and trace elements. We propose that the increase in inorganics (major and trace elements) is relative and that the increase in inorganics results in an increase in BD and corresponds to an area drought in the early 1960's (Cooke and others, 1997).

Below 18 cm, the Ni and Cu levels drop off rapidly; similarly high levels are not seen in any other part of the core.

The second shift in core SB1a occurs between 42-29 cm with increases in the concentrations of almost all the inorganic constituents, including Al, and a 1.7X increase in BD (fig. 17). Although we ran whole-peat ^{14}C ages, which suggest that this section of core is 200-400 years old, we cannot calibrate the ages, so are not using them for age assignments to this part of the SB1a core material. OCS data for core SB1c indicate that peat in the depth interval of 50-56 cm was deposited around the turn of the century, probably between 1900 and 1910 (table 9). If we assume a linear rate of organic matter accumulation (table 12) between 1905 and 1952, then the peak in concentrations of inorganic constituents and BD values occurred in 1936-37. As with the shift in values from 18-12 cm, there is no identifiable clay and (or) silt lens associated with the shift between 42 and 29 cm. We speculate that the increase in BD values results from the increase in inorganic constituents. We suggest that the increases are climatically related and correspond to (1) the 1925 drought (Cooke and others, 1997), or (2) 1930's regional drying in the south-central U.S. The absence of an increase in Ca supports number 1, but neither hypothesis is proved by the available data.

Trends in the inorganic constituents show that the two intervals with noticeable shifts in data trends, 18-12 and 42-29 cm in core SB1a, occur during a period of generally increasing concentrations of metals such as Cr, Co, V, lead (Pb), and Cu beginning at a depth of about 60 cm. This gradual increase in metal concentrations is probably associated with the continued increase in the urbanization and industrialization of the New Orleans area. Zinc has a slightly different pattern than other inorganic constituents. Zinc (Zn) concentrations increase sharply at about 80 cm depth. If the age estimates are correct, then Zn concentrations increase during the 1800's, remain relatively high through the early 1900's, and then decline markedly until 1960 when they again increase. We hypothesize that the increase in Zn concentrations in the 1800's may be associated with the presence of a U.S. mint in New Orleans at that time (Collins, 1971), but when more data become available we may see that Zn is a proxy for human activity.

At corrected depths greater than 80 cm there appears to be an increasing trend in the concentrations of Al, Fe, Li, potassium (K), Co, La, Ce, Th, Pb, Zn, as well as in calculated percent mineral matter and BD. This trend continues to the bottom of the core SB1a at 100 cm. If looked at stratigraphically, then inorganics and BD *increase* upward toward an overlying clay layers and *decrease* upward from the clay layer. The description of SB1c core in the *Geologic and Hydrogeologic Setting* section of this report support the analytical data.

TABLE 10A. Physical and chemical data for samples from core SB1a, St. Bernard Parish, LA

TABLE 10B. Major-element and trace-element data for samples from core SB1a, St. Bernard Parish, LA

TABLE 10C. Trace-element data for samples from core SB1a, St. Bernard Parish, LA

TABLE 11A. Physical and chemical data for samples from core SB1c, St. Bernard Parish, LA.

TABLE 11B. Major-element and trace-element data for samples from core SB1c, St. Bernard Parish, LA.

TABLE 11C. Trace-element data for samples from core SB1c, St. Bernard Parish, LA.

Mercury (Hg) was run for five samples from core SB1c (table 13) by methods described in Elrick and Horowitz (1986). There is an apparent increase in concentration toward the surface, similar to trends documented as early as the 1970s, from Canada (Thomas, 1972) and England (Aston and others, 1973). However, because all Hg values were at or below background values for most terrestrial sediment (commonly from 0.1-2 ppm; Förstner and

Wittman, 1979), no other samples were analyzed.

TABLE 12. Mercury values for selected samples from core SB1c, St. Bernard Parish, LA.

Bulk density (BD). -- In general, BD data for SB1a (table 10a) and SB1b (table 3) are highly correlated, $r^2=0.9$. SB1a BD values range from 0.07-0.16 g cm⁻³; SB1b BD values range from 0.07-0.18 g cm⁻³. BD values for 10 and 64 cm depths in core SB1a are 0.11 and 0.09 g cm⁻³, respectively. Values for the same depths, 10 and 65 cm, in core SB1c are equivalent, 0.10 and 0.09 g cm⁻³.

There are clear shifts upward in BD values at the same depth increments, 12-18 and 36-42 cm, as seen in the inorganic constituents, suggesting that the increase in BD may result from the increase in percent inorganics (fig. 16).

REFERENCES

[Return to CONTENTS](#)

DATA INTERPRETATION AND SUMMARY

H.W. Markewich, G.R. Buell, T.L. Fries, and J.P. McGeehin

Data for cores taken in a brackish marsh in St. Bernard Parish, near Chalmette, Louisiana can be used to indicate the rate of organic carbon (OC) accumulation for about the last 3 k yr. Compositional data allow estimates of the rate of decay as well as the rate of accumulation. Trends in the elemental chemistry of the peat deposits and in the microstratigraphic markers allow age/depth relations to be determined on a decadal time scale for the last 200 years. The combination of age and chemical data allows estimates of carbon storage for the cores. This section summarizes the data and shows how the data were used to determine sedimentation rates and carbon storage.

The successful application of various analytical techniques for determination of sediment age fulfilled a major goal of our investigation in the Mississippi River deltaic plain. The techniques and the results of their application are also summarized in this section.

Age Data

One of the problems inherent with using different methods of sample collection, particularly in peat, is accounting for the degree of compaction. Even when using one method in uniform parent material, just one change, such as the diameter of the sample tube, results in a major change in the degree of compaction. Using the same method with exactly the same equipment in what appears to be exactly the same type of parent material does not necessarily ensure the same degree of compaction. For instance, at the St. Bernard core locality, the 1-m cores SB1a and SB1b were less than 7 m apart and taken using the same vibracore techniques and identical 3-in (7.62 cm) diameter sample tubes (aluminum irrigation pipe). The measured compaction for SB1a was 7.3 percent; for SB1b it was 9.7 percent. Core length also affects compaction, as demonstrated by the measured compaction of 29 percent in 4-m long core SB1c located in close proximity to cores SB1a and SB1b. Thus, even with correcting for compaction, assigning an age to a given depth in one core to the same depth in another core is problematic without verifying data. In the case of the St. Bernard cores, C and Cs isotopic analyses were used for age determinations for the uppermost 25 cm of sediment. The comparison of first-appearance and peak-occurrence of microscopic stratigraphic markers with historical records was used for age determinations for the 85-50 cm depth interval (from about 1830 to 1910). The age of sediment below 1 m depth was determined by radiocarbon (¹⁴C) analysis.

Several summary statements can be made concerning the ages of specific depth intervals in the surface 4 m of sediment at the St. Bernard core locality.

- Cs and C isotopic data for cores SB1b and SB1a, respectively, indicate that the sediment at 15-16 cm was land surface in 1963-64; the sediment from 25 -28 cm was land surface in the early 1950s, and is here considered to be background for above testing and is assigned to the year 1952.
- An increase in microforam linings in the uppermost 11 cm of sediment in core SB1c suggests that this interval of sediment post-dates cutting of the Mississippi River - Gulf Outlet canal (MR-GO canal) , 1958-1968.
- The appearance of oil-produced OCS at 56 cm indicates that sediment at this depth was the land surface no earlier than 1900, and marks the transition from steam to internal combustion engines.
- The peak of wood/coal OCS at 71 cm is attributed to the combustion of wood and coal in mills, steamboats and steam locomotives during the mid-1800's, probably between 1860 and 1880. The drop in TCT, probably *Taxodium distichum* (bald cypress), at this level could indicate a later date of 1890, the period of regional cypress logging. However, large-scale logging of cypress in and around New Orleans, including this part of St. Bernard Parish, occurred earlier than the region-wide clearing. So the decrease in TCT in the St. Bernard core samples reflects clearing activities in the mid 1800's.
- Assigned ages for the 56-200 cm depth interval are probably the most uncertain.

First, there is the sampling error inherent with any compressed core material (discussed in the first paragraph of this section). Within the

uppermost 1 m of core the sampling error probably puts a \pm of at least a decade on all assigned ages. Pollen, specifically *Vigna luteola*, and OCS data indicate that the probable age for peat at 85 cm is about 1830. However, if the increase in charcoal concentration at 71 cm is related to the 1788 New Orleans fire, and not to the logging of *Taxodium* (bald cypress), then the assigned age of 1830, based on microscopic stratigraphic markers, could be too young by 40 to 60 years. This is possible since it is known that *Vigna luteola* was introduced into the New World by the late 1700's. One argument against this older age assignment is the fact that the Conseil Plantation, which includes the St. Bernard core locality, was not established until 1790, and *Vigna luteola* was just not reported in the New Orleans area until 1828.

If the 85 cm depth is assumed to represent the 1830 surface horizon, and a 0.5 cm yr^{-1} rate of peat accumulation is applied backwards in time, then the peat at 106 cm was the land surface in 1790. The reported ^{14}C age for sediment at 106 cm is $550 \pm 50 \text{ yr BP}$ (table 1). We suggest that neither the 1790 nor the ^{14}C age are definite. Even if a 40 to 60 year correction is applied to account for sampling error, the ages determined by matching diagnostic stratigraphic markers with the historical record are at least 300 years younger than the 550 ± 50 radiocarbon age. As with the other ^{14}C ages the $550 \pm 50 \text{ yr BP}$ age need to be tree-ring calibrated and corrected for reservoir effects.

One source of error for all the radiocarbon ages is the use of just one standard deviation for an error bar. A more realistic error should probably be two standard deviations (table 1). This would make the 1140 yr BP age at 166.4 cm and the 1000 yr BP age at 194.6 cm indistinguishable. The same would be true for the 1920 yr BP age at 235.5 cm and the 1980 yr BP age at 265.1 cm.

Also, all ^{14}C ages should be, but have not yet been, tree-ring calibrated. For example, the 550 yr BP age, including all error bars, when calibrated would most probably be equivalent to 1421-1327.

Another problem in the ^{14}C age determinations may be related to allochthonous material transported with the interdistributary silt and clay deposits and incorporated into the adjacent peat. There are two lenses of interdistributary silt and clay in the surface 2 m, at 82 and 158 cm. The material in these lenses could be the same age, but probably is older than the surrounding autochthonous peat.

The effects of environmental factors on ^{14}C ages also need to be considered. One example is the effect of marine water (Stuiver and Braziunas, 1993). Peats in the St. Bernard cores are saturated with brackish water. The reservoir effect of the water on the age of the peat needs to be determined.

Another source of error may be the effect of old carbon on plants growing in a substrate of similar but decomposed material such as discussed by Kilian and others (1995). We are presently involved in specific sampling and analyses of peat in the region to see if the problems with age determinations in this part of the sediment record can be resolved.

- Pollen and ^{14}C data indicate that by 1 ka marsh vegetation was fully dominant in this part of the St. Bernard delta; the Mississippi River had by this time migrated westward to form the LaFourche delta (fig. 3).
- ^{14}C ages and palynomorphic data from this study, and ^{14}C ages from previous studies (Frazier, 1967; Törnqvist and others, 1996), indicate that floodplain and marsh vegetation began forming about 3 ka in this part of the St. Bernard delta.
- The $8420 \pm 60 \text{ yr BP}$ age on organic material in the interdistributary clay (at 396 cm in core SB1c) suggest a 5 k yr hiatus between the clay and the overlying peat. We hypothesize that the $8420 \pm 60 \text{ yr BP}$ age does not represent the real age of the clay, but only the age of the allochthonous (transported) material included in the clay. More data are needed to determine the true age of the clay.

Carbon Inventory - Accumulation and Storage

Data for incremental carbon storage for core SB1a shows a decreasing trend from 100 to 50 cm (fig. 18). The trend may possibly be related to the apparent increase in mineral matter content (table 10A) over the same period. Carbon storage in this depth interval also shows dramatic short-term shifts at those depths where shifts in bulk density are observed, 42-29 and 18-12 cm. The change in the pattern of incremental storage above and below 85 cm may be an artifact of not having good age control for the sediment record from 200-85 cm. Cumulative carbon storage is relatively unaffected by any of the perturbations in incremental carbon storage (fig. 18).

TABLE 13. Organic carbon cumulative mass and age-interval accumulation rates for cores SB1a and SB1c.

FIGURE 18. Carbon storage: (A) incremental; (B) cumulative for cores SB1a and SB1c, St. Bernard Parish, LA.

An evaluation of carbon storage as a function of time (table 13, fig. 9), based on ^{14}C ages for core SB1c and ^{14}C bomb spike determinations for core SB1a and ^{137}Cs analysis of core SB1b, suggest that (1) the OC accumulation rate is reasonably constant through time; or, if the near surface data (0-85 cm) is viewed separately, then (2) the OC accumulation rates for about the last 200 years may be an order of magnitude greater than long term rates. Neither possibility can be proven or excluded using the data from this study. The higher rates in the upper 85 cm of core may be an artifact of our ability to differentiate time intervals. The 170 years, represented by the uppermost 85 cm of sediment, has an average OC accumulation rate of $0.20 \text{ kg m}^{-2} \text{ yr}^{-1}$. While this seems high when compared to the long-term OC accumulation rate of $0.04 \text{ kg m}^{-2} \text{ yr}^{-1}$, the 170 year interval represents only 1/8 to 1/4 the time represented by the sediment intervals between ^{14}C "dated" samples below 85 cm. It is entirely possible that periods of low or no deposition included in these longer time intervals are just not present in the shorter recent record. It may also be possible, that the ^{14}C ages, which have not been corrected for environmental variables, are significantly older than the actual ages. This possibility is reflected by the difference in grass and peat ^{14}C activities shown in figure 6. The most probable explanation is that there is an inherited OC fraction in the whole peat sample. As mentioned above, we are investigating the various sources of error in the analytical age determinations.

FIGURE 19. Plots of carbon mass/area versus time (years before present) for the SB1c core, St. Bernard Parish, LA. (A) arithmetic plot showing apparent difference in recent (first four data points) and long term rates, (B) semi-log plot showing an exponential relation, and

(C) log/log plot showing, that once the corrected age data become available, a decay constant can be calculated for the decomposition rates of marsh organic matter.

The wide variation in apparent OC accumulation rates for the last 200 years and for the period from 1,000-3,000 yr BP may also be explained by decomposition of organic matter over time, shown for the marsh environment as the difference in ^{14}C values for grass and peat in figure 6. Studies of organic matter breakdown in peat bogs have shown an order of magnitude difference in rates of loss of organic matter observed for young peats compared with the rates for older peats. Changes in decomposition processes with depth (aerobic to anaerobic) have been proposed as the major cause of this difference (Clymo, 1965, 1984). By fitting an exponential decay function to the time versus OC accumulation data, turnover times for organic matter can be estimated (fig. 19A, B, C). If OC accumulation is thought of as production, then the amount OC at a given time (t^1) is equal to the amount at t^0 plus the amount accumulating yr^{-1} minus the amount decomposed yr^{-1} . This method assumes a constant rate of loss as well as a constant rate of production and is limited by the accuracy of the dating methods. At present, the age data are not definitive enough for us to determine if the change in accumulation rates above and below 85 cm depth is real. The probability of material at a given depth being a certain age will be calculated as data from other marsh cores become available. Decay (decomposition) functions will be determined at that time. We anticipate that for cores SB1a and SB1c the long-term turnover (decay or decomposition) time will be between 2,200 and 5,000 years and that short-term turnover time will be between 80 and 250 years, similar to those described in Clymo (1984).

A few summary statements can be made about OC accumulation rates and OC storage from analysis of the St. Bernard Parish, Louisiana, core SB1a and SB1c.

- Incremental OC accumulation values range from 0.15-0.37 $\text{kg m}^{-2} \text{cm}^{-1}$ for core SB1a and 0.05-0.44 $\text{kg m}^{-2} \text{cm}^{-1}$ for core SB1c.
- Cumulative storage for core SB1a is 27.8 kg m^{-2} . Cumulative storage for core SB1c is 132 kg m^{-2} .
- Variations in incremental storage do not seem to affect the near linear trend of cumulative storage.
- There appears to be a relatively constant primary production of organic material, despite annual variability. The OC accumulation rate decreases exponentially with age and is governed by a constant decay rate from the time of burial when all decomposition is anaerobic.
- OC inventory for the surface 1 m compares with values shown by Bliss and Waltman (1995) for this area. Their map, based on the STATSGO data base (USDA, 1991), places this part of the Mississippi River deltaic plain in two categories, from 24-41 kg m^{-2} and from 41-69 kg m^{-2} . An average depth of 1 m is assumed.
- The values shown on soil carbon map (Bliss and Waltman, 1995) are comparable to the 41.5 kg m^{-2} cumulative storage value for 127 cm depth in core SB1c. The rate of OC accumulation shown for core SB1a in table 13 corresponds well to rates determined by Smith and others (1983) for brackish and freshwater marshes in southeastern Louisiana.

REFERENCES

This page is <URL:http://geochange.er.usgs.gov/pub/carbon/OFR_98-36/>

Maintained by Peter Schweitzer

Last updated 9-Feb-1998

Carbon Storage and Late Holocene Chronostratigraphy of a Mississippi River Deltaic Marsh, St. Bernard Parish, Louisiana

First Report: Mississippi Basin Carbon Project Process Studies

Edited by Helaine Walsh Markewich

U.S. GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

REFERENCES

Introduction

IPCC (Intergovernmental Panel on Climate Change), 1996, *Climate Change 1995 -- The Science of Climate Change*: Cambridge, Cambridge University Press, 572 p.

Markewich, H.W., in press, Detailed descriptions for sampling, sample preparation and analysis of cores from St. Bernard Parish, Louisiana: U.S. Geological Survey Open-File Report

National Science and Technology Council, Committee on Environment and Natural Resources, Subcommittee on Global Change Research, 1997, *Our Changing Planet - the FY 1998 U.S. Global Change Research Program*, 118 p.

NSSL (National Soil Survey Laboratory), 1994, National soil characterization data compact disc, May 1994, Natural Resource Conservation Service (formerly the Soil Conservation Service).

Sasser, et al., 1995, Floating marshes in the Barataria and Terrebonne Basins, Louisiana Coastal Ecology Institute, Baton Rouge, LSU-CEI-94-03.

Geologic and Hydrogeologic Setting

Coleman, J.M. and Roberts, H.H., 1989, Mississippi River depositional system - Baton Rouge to New Orleans, Louisiana, in *Coastal and Marine Geology of the United States, Field Trip Guidebook T170*, American Geophysical Union, p. 1-35.

Dillon, W.D., and Oldale, R.N., 1978, Late Quaternary sea-level curve: Reinterpretation based on glaciotectionic influence: *Geology*, v. 6, p 56-60.

Dunbar, J. B., Blaes, M. R., Dueitt, S. E., May, J. R., and Stroud, K. W., 1994, Geological investigation of the Mississippi River deltaic plain: Technical Report No. GL-84-15, US Army Engineer Waterways Experiment Station, Vicksburg Mississippi.

Fisk, H.N. 1938, *Geology of Grant and LaSalle Parishes*: Louisiana Department of Conservation Geology Bulletin, v. 10, 246 p.

Fisk, H.N. 1944, *Geological investigation of the Alluvial Valley of the Lower Mississippi River*: US Army Corps of Engineers, Mississippi River Commission, Vicksburg, Mississippi.

- Fisk, H.N., 1952, Geological investigation of the Atchafalaya Basin and problems of Mississippi River diversion, Vol. I and II: US Army Corps of Engineers, Mississippi River Commission, Vicksburg, Mississippi.
- Fisk, H.N., 1955, Sand facies of Recent Mississippi Delta deposits: 4th World Petroleum Congress Proceedings, Rome, Italy, Section I, p. 377-398.
- Fisk, H.N., and McFarlan, E., 1955, Late Quaternary deposits of the Mississippi River, in, *Crust of the Earth: Special Paper No. 62*. Geological Society of America, Boulder, Colorado., p. 279-302.
- Frazier, D.E., 1967, Recent deltaic deposits of the Mississippi River: Their development and chronology: *Transactions Gulf Coast Association Geological Society*, v. 17, p. 287-315.
- Kolb, C.R., and Van Lopik, J.R., 1958, Geology of the Mississippi River deltaic plain, southeastern Louisiana: Technical Report 3-483, US Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
- Kolb, C.R. and Van Lopik, 1966, Depositional environments of the Mississippi River deltaic plain, southeastern Louisiana: *in* Shirley, M. L. and Ragsdale, J. A., eds., *Deltas*, Houston Geological Society, Houston, Texas, p.17-62.
- Martinson, D.G., Psias, N.G., Hays, J.D., Imbrie, J., Moore, T.C., Jr., and Shackleton, N.J., 1987, Age dating and orbital theory of the ice ages: Development of a high-resolution 0 - 300,000 - year chronostratigraphy: *Quaternary Research*, v. 27, p. 1-29.
- McFarlan, E., Jr., 1961, Radiocarbon dating of late Quaternary deposits, South Louisiana: *Geological Society of America Bulletin*, v. 72, p. 129-158.
- McIntire, W. G. 1958, Prehistoric Indiana settlements of the changing Mississippi River Delta: Louisiana State University, Coastal Studies Series No. 1, Baton Rouge, Louisiana.
- Nummedal, D. 1983, Rates and Frequencies of Sea-Level Changes: A review with an approach to predict future sea levels in Louisiana: *Gulf Coast Association Geological Society Transactions*, v. 33, p 361-366.
- Saucier, R.T., 1994, *Geomorphology and Quaternary history of the Lower Mississippi River Valley*: U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi, December 1994, volumes I and II.
- Törnqvist, T. E., Kidder, T. R., Autin, W.J., van der Borg, K., de Jong, A.F. M., Klerks, C.J., Snijders, E.M.A., Storms, J.E.A., van Dam, R.L., Wiemann, M.C., 1996, A revised chronology for Mississippi River subdeltas: *Science*, v. 273, p. 1693-1696.

Field Methods

- Smith, D.G., 1984, Vibracoring fluvial and deltaic sediments: Tips on improving penetration and recovery: *Journal of Sedimentary Petroleum*, v. 54, no. 2, p. 660-663.

¹⁴C Age Determinations

- Burchuladze, A. A., Chudy, M., Eristavi, I. V., Pagava, S. V., Povinec, P., Sivo, A, and Togonidze, G. I., Anthropogenic ¹⁴C variations In atmospheric CO₂ and wines, 1989, *Radiocarbon*, v. 31, no. 3, p. 771-776
- Kilian, M.R. van der Plicht, J. and van Geel, B., 1995, Dating raised bogs: new aspects of AMS ¹⁴C wiggle matching, a reservoir effect and climatic change: *Quaternary Science Reviews*, v. 14, p. 959-966.
- Levin, I., Kromer, B., Schoch-Fischer, H., Bruns, M., Munnich, M., Berdau, D., Vogel, J., Munnich, K. O., 1985, *Radiocarbon*, v., 27, no. 1, p. 1-19.
- Markewich, H.W., in press, Detailed descriptions for sampling, sample preparation and analysis of cores from St. Bernard Parish, Louisiana: U.S. Geological Survey Open-File Report
- Stuiver, M. and Braziunas, T. F., 1993, Modeling atmospheric ¹⁴C influences and ¹⁴C ages of marine samples to 10,000 B.C.: *Radiocarbon*, v. 35, no. 1, p. 137-189.
- Stuiver, M. and Reimer, P. J., 1993, Extended ¹⁴C data base and revised Calib 3.0 ¹⁴C Age Calibration Program: *Radiocarbon*, v. 35, no. 1, p. 215-230.

¹³⁷Cs Analysis

Callendar, E., and Robbins, J.R., 1993, Transport and accumulation of radionuclides and stable elements in a Missouri River reservoir. *Water Resources Research*, v. 29, p. 1787-1804.

Ritchie, J.C., and McHenry, J.R., 1984, Application of radioactive ^{137}Cs for measuring soil erosion and sediment accumulation rates and patterns: a review: *Journal of Environmental Quality*, v. 19, p. 215-233.

Biochronostratigraphy

U.S. Army Corps of Engineers (COE), 1975, Draft environmental statement: Composite draft environmental statement for operation and maintenance work on three navigation projects in the Lake Borgne vicinity, Louisiana (The Mississippi River Gulf Outlet, Bayous La Loutre, St. Malo, and Yscloskey Bayou Dupre). U.S. Army Engineer District, New Orleans, Louisiana, April, 1975.

Arthur, S.C., 1937, *Old New Orleans*: New Orleans, Harmanson, 246 p.

Barss, M.S., and Williams, G.L., 1973, Palynology and nanofossil processing techniques: *Geological Survey of Canada* 73-26, 25 p.

Beig, F.C., 1906, *Naval Boilers, A text-book for the instruction of midshipmen at the U.S. Naval Academy*: Annapolis, The United States Naval Institute, 372 p.

Bernhard, K.D.O.S.-W.-E., 1828, *Travels Through North America During the years 1825 and 1826*: Philadelphia, Carey, Lea, and Carey, 238 p..

Clark, J.S., and Patterson, W.A., III, 1984, Pollen, Pb-210, and opaque spherules: an integrated approach to dating and sedimentation in the intertidal environment: *Journal of Sedimentary Petrology*, v. 54, no. 4, p. 1251-1265.

Clark, R.L., 1982, Point count estimation of charcoal and pollen preparations and thin sections of sediments: *Pollen et Spores*, v. 24, no. 3-4, p. 523-535.

Coleman, J.M. and Roberts, H.H., 1989, Mississippi River depositional system - Baton Rouge to New Orleans, Louisiana, in *Coastal and Marine Geology of the United States, Field Trip Guidebook T170*, American Geophysical Union, p. 1-35.

Davis, O.K., 1987, Spores of the dung fungus *Sporormiella*: Increased Abundance in Historic Sediments and before Pleistocene Megafaunal Extinction: *Quaternary Research*, v. 28, p. 290-294.

de Rojas, L., 1937, The Great Fire of 1788 in New Orleans: Reprinted from *The Louisiana Historical Quarterly*, New Orleans, v. 20, no. 3, 14 p.

DuFour, C., 1968, The people of New Orleans, in Carter, H., ed., *The Past as Prelude: New Orleans 1718-1968*, Tulane University, New Orleans, LA., 383 p..

Fortier, A., 1904, *A History of Louisiana*: New York, Goupil and Co., 4 volumes.

Fossier, A.A., 1957, *New Orleans: The Glamour Period, 1800-1840*: New Orleans, Pelican Publishing, 520 p.

Frazier, D.E., 1967, Recent deltaic deposits of the Mississippi River: Their development and chronology.: *Transactions of the Gulf Coast Association of Geological Societies*, v. 17, p. 287-315.

Gardiner, R., and Lambert, A., eds., 1992, *Steam, steel and shellfire: The steam warship 1815-1905*: Conway Maritime Press, Ltd., London, England, 192 p.

Gish, J., 1994, Large fraction pollen scanning and its application in archaeology, in Davis, O.K., ed., *Aspects of archaeological palynology: methodology and application*: American Association of Stratigraphic Palynologists Contributions Series, no. 29, American Association of Stratigraphic Palynologists Foundation, Dallas, Texas, p. 92-100.

Gray, L.C., 1941, *History of agriculture in the Southern United States to 1860*: New York, Peter Smith, v. 2, 1086 p.

Griffin, J.J., and Goldberg, E.D., 1979, Morphologies and origin of elemental carbon in the environment: *Science*, v. 206, no. 2, p. 563-565.

Griffin, J.J., and Goldberg, E.D., 1981, Sphericity as a characteristic of solids from fossil fuel burning in a Lake Michigan sediment: *Geochim et Cosmochim, Acta*, v. 45, p. 763-769.

Hunter, L., 1949, *Steamboats on the western rivers*: Cambridge, Harvard University Press, 684 p.

Iversen, J., 1941, Land occupation in Denmark's Stone Age: *Danmarks Geologiske Undersogelse*, v. 2, no. 66, p. 68 p.

Jones, J.G., and Bryant, V.M., Jr., 1992, Phytolith taxonomy in selected species of Texas cacti, in Rapp, G., Jr., and Mulholland, S.C., eds., *Phytolith*

systematics: New York, Plenum, p. 215-238.

Kaplan, L., Smith, M.B., and Sneddon, L.A., 1992, Cereal grain phytoliths of Southwest Asia and Europe, in Rapp, G., Jr., and Mulholland, S.C., eds., *Phytolith systematics*: New York, Plenum, p. 149-174.

La Tourrette, J., 1845, La Tourrette's reference map of the state of Louisiana, scale 1:380,160.

Lewis, W., Vinay, P., and Zenger, V., 1983, *Airborne and allergenic pollen of North America*: Baltimore, The Johns Hopkins University Press, 254 p.

Liu, K.B., and Fearn, M.L., 1993, Lake-sediment record of late Holocene hurricane activities from coastal Alabama: *Geology*, v. 21, p. 793-796.

Mackie, W.W., 1946, Blackeyed beans in California: *California Agricultural Experimental Station Bulletin*, 56 p.

Markewich, H.W., in press, Detailed descriptions for sampling, sample preparation and analysis of cores from St. Bernard Parish, Louisiana: U.S. Geological Survey Open-File Report

McCrone, W.C., and Delly, J.G., 1973, *The Particle Atlas*: Ann Arbor, Ann Arbor Science Publishers, v. 6.

McKee, R., 1948, *The legumes of many uses, grass: The yearbook of agriculture 1948*: Washington, D.C., U.S. Government Printing Office, 892 p.

Menn, J.K., 1964, *The large slaveholders of Louisiana - 1860*: New Orleans, Pelican Publishing Company, 432 p.

Milton, J.L., 1953, *Marine Steam Boilers*: London, George Newnes, 256 p.

Mulholland, S., and Rapp, G., Jr., 1992, A morphological classification of grass silica-bodies, in Rapp, G., Jr., and Mulholland, S.C., eds., *Phytolith systematics*: New York, Plenum, p. 65-89.

Norgress, R.E., 1947, The history of the cypress lumber industry in Louisiana: *The Louisiana Historical Quarterly*, v. 30, no. 3, 83 p.

Piperno, D., 1988, *Phytolith analysis*: Academic Press, San Diego, 280 p.

Piperno, D.R., 1989, The occurrence of phytoliths in the reproductive structures of selected tropical angiosperms and their significance in tropical paleoecology, paleoethnobotany and systematics: *Review of Paleobotany and Palynology*, v. 61, p. 147-173.

Purseglove, J.W., 1972, *Tropical Crops: Monocotyledons*: London, Longman, 719 p.

Rapp, G., Jr., and Mulholland, S.C., eds., 1992, *Phytolith systematics*: New York, Plenum, 350 p.

Renberg, I., and Wik, M., 1985, Carbonaceous particles in lake sediments-pollutants from fossil fuel combustion: *Ambio*, v. 14, no. 3, p. 161-163.

Rose, N.L., 1990, A method for the extraction of carbonaceous particles from lake sediments: *Journal of Paleolimnology*, v. 3, p. 45-53.

Rose, N.L., Harlock, S., Appleby, P.G., and Battarbee, R.W., 1995, Dating of recent lake sediments in the United Kingdom and Ireland using spheroidal carbonaceous particle (SCP) concentration profiles: *The Holocene*, v. 5, p. 328-335.

Sitterson, J.C., 1953, *Sugar Country*, University of Kentucky Press, 414 p.

Terasmae, J., and Weeks, N.C., 1979, Natural fires as an index of paleoclimate: *Canadian Field-Naturalist*, v. 93, p. 116-125.

Tolonen, K., 1986, Charred Particle Analysis, in Berglund, B.E., ed., *Handbook of Holocene Palaeoecology and Palaeohydrology*: Chichester, John Wiley and Sons, p. 485-496.

Tolonen, M., 1978, Paleoecology of annually laminated sediments in Lake Ahvenainen, South Finland. I: Pollen and charcoal analyses and their relation to human impact: *Annals of Botanica Fennici*, v. 15, p. 155-208.

Twiss, P.C., 1992, Predicted world distribution of C3 And C4 grass phytoliths, in Rapp, G., Jr., and Mulholland, S.C., eds., *Phytolith systematics*: New York, Plenum, p. 113-128.

Twiss, P.C., Suess, E., and Smith, R.C., 1969, Morphological classification of grass phytoliths: *Proceedings of the Soil Science of America*, v. 33, p. 109-115.

Villere, S.L., 1981, Jacques Philippe Villere, first native-born governor of Louisiana 1816-1820: New Orleans, The Historic New Orleans Collection, 135 p.

Whitehead, D.R., and Sheehan, M.C., 1985, Holocene vegetational changes in the Tombigbee River valley, Eastern Mississippi: *The American Midland Naturalist*, v. 113, no. 1, p. 122-137.

Compositional Analysis

Aston, S., Bruty, D., Chester, R., and Padgham, R., 1973, Mercury in lake sediments: a possible indicator of technological growth: *Nature* (London), v. 241, p. 450-451.

Collins, Charles J., 1971, A history of the United States Mint, New Orleans, Louisiana: Baton Rouge, 47 p.

Cook, E., Meko, D., Stahle, D., and Cleaveland, M., 1997, Reconstruction of past drought across the coterminous United States from a network of climatically sensitive tree-ring data: National Oceanic and Atmospheric Administration web site accessed December 4, 1997, on the World Wide Web at URL <http://www.ngdc.noaa.gov/paleo/usclient.html>.

Elrick, K.A., and Horowitz, A.J., 1986, Analysis of rocks and sediments for mercury, by wet digestion and flameless cold vapor atomic absorption: U.S. Geological Survey Open-File Report 86-596, 12 p.

Graney, J.R., et al., 1995, Isotopic record of lead pollution in lake sediments from the northeastern United States: *Geochimica et Cosmochimica Acta*, v. 59, no. 9, p. 1715-1720.

Förstner, U., and Wittman, G.T.W., 1979, Metal pollution in the aquatic environment: Springer-Verlag, New York, 486 p.

Luoma, S.N., and Bryan, G.W., 1981, A statistical assessment of the form of trace metals in oxidized estuarine sediments employing chemical extractants: *The Science of the Total Environment*, v. 17, p. 165-196.

Markewich, H.W., in press, Detailed descriptions for sampling, sample preparation and analysis of cores from St. Bernard Parish, Louisiana: U.S. Geological Survey Open-File Report

Thomas, R., 1972, The distribution of mercury in the sediment of Lake Ontario: *Canadian Journal of Earth Science*, v. 9, p. 636-651.

Data Interpretation and Summary

Bliss, N.B. and Waltman, S., 1995, Soil organic carbon kg/m² total profile: unpublished map, 1:5,000,000.

Clymo, R.S., 1965, Experiments on breakdown of Sphagnum in two bogs, *Journal of Ecology*, v. 53, p. 747-757.

Clymo, R.S., 1984, The limits to peat bog growth, *Philosophical Transactions of the Royal Society of London*, v. 303, p. 605-654.

Frazier, D.E., 1967, Recent deltaic deposits of the Mississippi River: Their development and chronology: *Transactions Gulf Coast Association Geological Society*, v. 17, p. 287-315.

Kilian, M.R. van der Plicht, J. and van Geel, B., 1995, Dating raised bogs: new aspects of AMS ¹⁴C wiggle matching, a reservoir effect and climatic change: *Quaternary Science Reviews*, v. 14, p. 959-966.

Smith, C.J., DeLaune, R.D., and Patrick, W. H., Jr., 1983, CO₂ emissions and carbon accumulation in coastal wetlands: *Estuarine, Coastal, and Shelf Science*, v. 17, p. 21-29.

Stuiver, M. and Braziunas, T F, 1993, Modeling atmospheric ¹⁴C influences and ¹⁴C ages of marine samples to 10,000 B.C., *Radiocarbon*, v. 35, no. 1, p. 137-189.

Törnqvist, T. E., Kidder, T. R., Autin, W.J., van der Borg, K., de Jong, A.F. M., Klerks, C.J., Snijders, E.M.A., Storms, J.E.A., van Dam, R.L., Wiemann, M.C., 1996, A revised chronology for Mississippi River subdeltas: *Science*, v. 273, p. 1693-1696.

USDA (U.S. Department of Agriculture), 1991, State soil geographic data base (STATSGO), data users guide: Washington, D.C., Natural Resources Conservation Service (formerly Soil Conservation Service) Miscellaneous Publication 1492, 88 p.

This page is <URL:http://geochange.er.usgs.gov/pub/carbon/OFR_98-36/DeltaOFA-refs.html>

Maintained by Peter Schweitzer

Last updated 9-Feb-1998

TABLE 1. Data for ^{14}C age determinations for peat samples from core SB1c, St. Bernard Parish, Louisiana.								
St. Bernard core SB1c (location: 29°58'53" N, 89°55'27" W; sampled: 03/20/1996).								
Percent compaction (applied linearly): 29.1.								
Corrected depth sample midpoint (cm) ¹	Original depth sample midpoint (cm)	Lab I.D. (WW- nnn) ²	Material	Fraction modern ³	Age (yr BP ⁴ ± one standard deviation)		$\delta^{13}\text{C}$ (parts per thousand)	$\delta^{13}\text{C}$ lab ⁵
43.7	31	685	peat	0.9704	240	60	-20.0	R
69.1	49	746	peat	0.9732	220	60	-20.5	D
84.6	60	747	peat	0.9595	330	60	-21.9	D
84.6	60	748	grass	0.9734	220	50	-15.9	D
105.8	75	748	peat	0.9334	550	50	-20.0	M
126.9	90	750	peat	0.8976	870	60	-20.1	D
166.4	118	751	peat	0.8672	1140	60	-20.1	D
194.6	138	752	peat	0.8825	1000	60	-20.2	D
235.5	167	753	peat	0.7875	1920	60	-17.6	D
265.1	188	754	peat	0.7813	1980	50	-21.5	D
304.6	216	755	peat	0.7459	2360	60	-26.6	D
345.4	245	756	peat	0.7171	2670	50	-27.8	D
345.4	245	757	grass	0.7212	2630	50	-25.8	D
383.5	272	698	peat	0.6728	3180	50	-25.9	D
396.2	281	802	clay	0.3507	8420	60	-24.9	D
¹ Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.41 to correct for compaction.								
² ^{14}C data are archived in a world-wide database indexed by U.S. Geological Survey laboratory I.D. numbers "WW- nnn" where "WW" indicates AMS ages.								
³ Ratio of the ^{14}C activity of the sample to the activity of the modern standard, which is used in various calculations related to radiocarbon ages.								
⁴ ^{14}C age determinations not calibrated to tree-ring ages and not corrected for ^{14}C enrichment by seawater.								
⁵ R=Reston, VA; D=Denver, CO; M=Menlo Park, CA.								
Source: U.S. Geological Survey Open-File Report 98-36. Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana (H.W. Markewich, ed.).								

Last updated on 2/5/98
By Gary R. Buell

TABLE 2. Data for ^{14}C age determinations for peat samples from core SB1a, St. Bernard Parish, Louisiana.

St. Bernard core SB1a (location: 29°58'53" N, 89°55'27" W; sampled: 03/18/1996).

Percent compaction (applied linearly): 6.8.

Cor- rected depth sample mid- point (cm) ¹	Original depth sample mid- point (in) ²	Lab I.D. (WW- nnn) ³	Material	Fraction modern ⁴	Age (yr BP ⁵ ± one sigma)		$\Delta^{14}\text{C}$ (parts per thousand ± one standard deviation)		$\delta^{13}\text{C}$ (parts per thou- sand)	$\delta^{13}\text{C}$ lab ⁶
6.8	2.7	796	peat	1.1304	>MODERN		124.1	9.1	-18.5	D
9.5	3.5	797	peat	1.2081	>MODERN		201.4	8.2	-17.9	D
12.3	4.5	798	peat	1.2698	>MODERN		262.7	7.0	-17.5	D
15.0	5.5	763	peat	1.5288	>MODERN		520.3	8.4	-17.2	D
17.7	6.5	764	peat	1.2950	>MODERN		287.8	7.1	-26.8	D
20.4	7.5	765	peat	1.0319	>MODERN		26.2	6.7	-17.3	D
23.2	8.5	766	peat	0.9889	90	60	-16.5	6.5	-17.2	D
25.9	9.5	767	peat	0.9699	250	60	-35.5	6.3	-24.1	D
28.5	10.5	768	peat	0.9546	370	50	-50.7	5.3	-22.2	D
31.3	11.5	769	peat	0.9515	400	50	-53.7	5.3	-21.2	D
34.0	12.5	770	peat	0.9488	420	60	-56.4	5.9	-20.6	D
36.7	13.5	771	peat	0.9497	410	60	-55.6	6.2	-22.4	D
6.8	2.5	799	grass	1.1203	>MODERN		114.1	7.1	-16.5	D
9.5	3.5	800	grass	1.1892	>MODERN		182.6	7.6	-14.5	D
12.3	4.5	801	grass	1.2398	>MODERN		232.9	8.0	-16.1	D
15.0	5.5	772	grass	1.5530	>MODERN		544.4	10.0	-15.9	D
17.7	6.5	773	grass	1.5518	>MODERN		543.2	9.9	-17.0	D
20.4	7.5	774	grass	1.4642	>MODERN		456.0	9.4	-16.7	D
23.2	8.5	775	grass	1.1208	>MODERN		114.6	6.3	-15.4	D
25.9	9.5	776	grass	0.9726	220	60	-32.8	6.4	-17.3	D
28.5	10.5	777	grass	0.9729	220	50	-32.5	5.5	-14.9	D
31.3	11.5	778	grass	0.9673	270	50	-38.1	5.5	-14.0	D
34.0	12.5	779	grass	0.9661	280	50	-39.3	5.5	-15.0	D
36.7	13.5	780	grass	0.9725	220	50	-32.9	5.4	-14.3	D
25.9	9.5	781	seeds	0.9693	250	60	-36.0	6.1	-27.8	D

¹Corrected depth equals original depth (midpoint of sample interval) converted to centimeters (2.54 cm in⁻¹) and multiplied by 1.0743 to correct for compaction.²Top 2 inches (5.08 cm) discarded in the field.³ ^{14}C data are archived in a world-wide database indexed by U.S. Geological Survey laboratory I.D. numbers "WW-nnn" where "WW" indicates AMS ages.⁴Ratio of the ^{14}C activity of the sample to the activity of the modern standard, which is used in various calculations related to radiocarbon ages.⁵ ^{14}C age determinations not calibrated to tree-ring ages and not corrected for ^{14}C enrichment by seawater.⁶R=Reston, VA; D=Denver, CO; M=Menlo Park, CA.Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

TABLE 3. ^{137}Cs data for bomb-curve analysis of core SB1b, St. Bernard Parish, Louisiana.

St. Bernard core SB1b (location: 29°58'53' N, 89°55'27' W; sampled 03/18/1996).

Percent compaction (applied linearly): 9.7. dpm, disintegrations per minute.

Sample I.D.	Corrected depth sample midpoint ¹ (cm)	Net weight, air-dried (g)	Bulk density, air-dried (g cm ⁻³)	^{137}Cs activity (dpm \pm one standard deviation)		^{40}K activity (dpm \pm one standard deviation)	
SB1b-1 ²	6.7	--	--	--	--	--	--
SB1b-2	9.1	8.00	0.08	0.128	0.020	0.311	0.017
SB1b-3	11.4	7.40	0.07	0.145	0.017	0.268	0.014
SB1b-4	13.7	13.90	0.14	1.561	0.039	0.418	0.021
SB1b-5	16.0	13.29	0.13	2.814	0.050	0.430	0.021
SB1b-6	18.4	11.53	0.12	2.137	0.082	0.332	0.036
SB1b-7	20.7	11.67	0.12	0.759	0.037	0.298	0.023
SB1b-8	23.0	11.12	0.11	0.446	0.028	0.311	0.020
SB1b-9	25.3	13.06	0.13	0.302	0.024	0.446	0.020
SB1b-10	27.7	13.16	0.13	0.149	0.024	0.463	0.024
SB1b-11	30.0	12.20	0.12	0.052	0.014	0.334	0.013
SB1b-12	32.3	10.98	0.11	0.072	0.022	0.294	0.019
SB1b-13	34.7	11.50	0.11	0.039	0.020	0.260	0.018
SB1b-19	48.6	11.59	0.12	-0.026	0.018	0.304	0.017
SB1b-25	62.6	9.61	0.10	0.013	0.019	0.206	0.018

¹Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.1074 to correct for compaction.²Top 2 inches (5.08 cm) discarded in the field. No analysis of sample at corrected depth of 6.7 cm.Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).Last Updated on 2/5/98
By Gary R. Buell

TABLE 4. St. Bernard Parish, Louisiana cores - events and potential stratigraphic markers.		
Year	Historic event	Predictive stratigraphic marker
1893	Hurricane	Hurricane sediment layer and marine fossils
1890	Initiation of industrial-scale bald cypress (<i>Taxodium distichum</i>) cutting	Decline in <i>Taxodium distichum</i>
1860's	Initiation of coal-fueled steamboats and trains	Increase in coal-based carbonaceous spherules
1850's	Initiation of regional railroad network	Increase in wood-based carbonaceous spherules
1831	Hurricane	Hurricane sediment layer and marine fossils
1829	Coal barges begin supplying sugar mills	First appearance of coal-based carbonaceous spherules
1814	Beginning of steamboat activity on the Mississippi River	Appearance of wood-based carbonaceous spherules
1812	Hurricane	Hurricane sediment layer and marine fossils
1794	Second fire of New Orleans	Abundance of charcoal
1790	Major production of sugarcane begins	Increase in <i>Saccharum officinarum</i> phytoliths and pollen
1788	First fire of New Orleans	Abundance of charcoal
1751	Introduction of sugarcane (<i>Saccharum officinarum</i>)	Appearance of <i>Saccharum officinarum</i> phytoliths and pollen
1731	Peas and beans exported to the West Indies	Increase of pea and bean pollen and phytoliths
1722	Two hurricanes	Hurricane sediment layers and marine fossils
1721	New Orleans census records cattle and horses	Dramatic increase in <i>Sporomella</i> spores
1720	Introduction of rice (<i>Oryza sativa</i>) in region	Appearance of <i>Oryza sativa</i> phytoliths and pollen
1717	New Orleans settled by the French	Increase in Compositae pollen
Source: U.S. Geological Survey Open-File Report 98-36. Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana (H.W. Markewich, ed.).		

Last Updated on 2/5/98
By Gary R. Buell

TABLE 5. Sample depths and analyses for microscopic markers in core SB1c, St. Bernard Parish, Louisiana.

St. Bernard core SB1c (location: 29°58'53" N, 89°55'27" W; sampled 03/20/1996).

Percent compaction (applied linearly): 29.1.

Corrected depth sample midpoint (cm) ¹	Original depth sample midpoint (cm)	Palynology	Charcoal/ carbon spherules	Phytoliths/ diatoms
4.2	3	X		
11.3	8	X		
14.1	10		X	X
18.3	13	X		
28.2	20		X	X
32.4	23	X		
42.3	30		X	X
43.7	31	X		
56.4	40		X	X
69.1	49	X		
70.5	50		X	X
84.6	60	X	X	X
98.7	70		X	X
105.8	75	X		
112.8	80		X ²	
126.9	90	X	X ²	
166.4	118	X		
194.6	138	X		

¹Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.41 to correct for compaction.²Samples to be processed at a later date.Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).Last Updated on 2/5/98
By Gary R. Buell

TABLE 6. Discussion of pollen zones depicted in figure 8.

Zone 7 0 - 11 cm 1996 to < 1963	<ul style="list-style-type: none"> • Arboreal taxa stable except for <i>Quercus</i> which increases. • Compositae (high and low-spine) and Gramineae increase. • Coiled microforams increase dramatically. 	Modern zone, defined by an increase in microforam linings (fig. 9H) possibly resulting from the rise in salinity levels after construction of the Mississippi River/ Gulf Outlet Canal in the late 1950's.
Zone 6 11 - 77 cm < 1963 to > 1830	<ul style="list-style-type: none"> • Arboreal taxa decline, especially TCT. • Non-arboreal taxa remain unchanged except Cheno-Am which fluctuates. • Compositae (low-spine) rises dramatically, then begins declining. 	Defined by the dramatic decrease in TCT and the large increase in low-spine Compositae. The TCT decline records the historic decline in <i>Taxodium distichum</i> (bald cypress) resulting from intensive logging between 1890 and 1925 and the slightly earlier logging associated with the expansion of New Orleans. Continued increase of low-spine Compositae is the result of logging and/or expanding settlement.
Zone 5 77 - 106 cm > 1830 to 550 ± 50 yr BP (1350 - 1450; uncalibrated)	<ul style="list-style-type: none"> • Arboreal taxa remain the same except <i>Quercus</i> which declines. • Cheno-Am and Compositae (high-spine) remain low. • Compositae (low-spine) begins increase. • Cyperaceae, Gramineae, <i>Sagittaria</i>, and <i>Typha</i> decline. • <i>Vigna luteola</i> appears for the first time. 	Most noticeable is the increase in low-spine Compositae (fig. 9F) and the appearance of <i>Vigna luteola</i> (fig. 9G) (cow pea) at 85 cm, an Old World domesticate that probably made its first appearance after colonization of New Orleans in 1717. The increase in the sun-loving, low-spine Compositae supports the hypothesis that Zone 5 reflects the period of colonization and clearing of the New Orleans area.
Zone 4 106 - 214 cm 550 ± 50 (1450 - 1350; uncalibrated) to >1000 ± 60 yr BP	<ul style="list-style-type: none"> • <i>Pinus</i>, <i>Quercus</i>, <i>Salix</i>, and TCT increase. • Cheno-Am and Compositae remain low. • Cyperaceae and Gramineae are abundant. • <i>Sagittaria</i> appears for the first time. • <i>Typha</i> reaches a peak near the top of this zone. 	The return of wetland taxa such as <i>Salix</i> , Cyperaceae (fig. 9E) and <i>Typha</i> indicates an end to the drier conditions of Zone 3. Although Zone 4 is characterized by a return to wetter conditions, the abundance of Cyperaceae and Gramineae, and the appearance of <i>Sagittaria</i> , make this wetland qualitatively different from the wetland of Zone 2. Zone 4 has a stronger marsh character and probably records the beginning of the disintegration of the St. Bernard deltaic plain following its maximum development.
Zone 3 214 - 305 cm >1000 ± 60 to 2360 ± 60 yr BP	<ul style="list-style-type: none"> • <i>Juglans</i> and Piperaceae make their only appearance. • <i>Pinus</i>, <i>Quercus</i>, and Compositae (low-spine) are common. • <i>Salix</i> and TCT decline. • Cheno-Am, Compositae (high-spine), and <i>Typha</i> decline. • Cyperaceae and Gramineae increase at the initiation of the zone and subsequently disappear and decline, respectively. 	The presence of <i>Juglans</i> (and possibly Piperaceae) indicates a shift to drier, more terrestrial, conditions than those reflected in Zone 2. This conclusion is supported by the decline of <i>Salix</i> , Cheno-Am, and <i>Typha</i> . Following the delta model (Coleman and Roberts, 1989), Zone 3 is composed of the vegetation of the mature delta plain. That is, as the delta prograded further seaward of the core site, the site became more terrestrial in character.
Zone 2 305 - 384 cm 2360 ± 60 to 3180 ± 50 yr BP	<ul style="list-style-type: none"> • <i>Pinus</i> and <i>Quercus</i> common. • <i>Salix</i> reaches a peak. • TCT (Taxodiaceae/Cupressaceae/Thuja) generally increases. • Cheno-Am (Chenopodiaceae-Amaranthaceae) fluctuates but maintains a strong presence throughout. • Compositae (high-spine), Gramineae and <i>Typha</i> increase in the middle of the zone. • Compositae (low-spine) and Cyperaceae are present but rare. 	The presence of <i>Quercus</i> , <i>Salix</i> and TCT (fig. 9A) reflects the proximity of floodplain forests. Cheno-Am (fig. 9B) and <i>Typha</i> (fig. 9C) are typical pond or stream-side taxa in fresh to brackish water marshes. The presence of Compositae (high-spine) and Gramineae (fig. 9D) is consistent with a marsh environment, perhaps along the delta margin. Zone 2 is synchronous with the developmental phase of the St. Bernard delta.
Zone 1 384 - 396 cm 3180 ± 50 to 8420 ± 60 yr BP	<ul style="list-style-type: none"> • Pyrite abundant in sample. • Pollen not abundant. 	¹⁴ C age is old compared to previous work in the area (Frazier, 1967). The date, on decomposed plant fragments in the interdistributary clay, probably reflects older, redeposited, organic material.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

TABLE 7. Selected entries of the 1860 census of the large slaveholders of Louisiana (Menn, 1964).

Crops and livestock	Conseil Plantation (of St. Bernard Parish)	St. Bernard Parish	Orleans Parish
Cane sugar (1,000-pound hogheads)	250	3,884	1,750
Molasses (gallons)	16,000	218,850	109,000
Ginned cotton (400-pound bales)	0	0	400
Indian corn (bushels)	3,000	68,300	22,500
Irish potatoes (bushels)	1,500	4,150	1,950
Sweet potatoes (bushels)	1,200	4,280	2,600
Peas and beans (bushels)	0	639	2,430
Horses	4	63	44
Asses and mules	36	579	196
Milk cows	12	119	82
Oxen	15	124	63
Sheep	30	268	284
Swine	0	72	387

Source: U.S. Geological Survey Open-File 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

Last Updated on 2/5/98
By Gary R. Buell

TABLE 8. Phytolith and other siliceous microfossil data for core SB1c, St. Bernard Parish, Louisiana.

St. Bernard core SB1c (location: 29°58'53" N, 89°55'27" W; sampled: 03/20/1996).

Percent compaction (applied linearly): 29.1.

Microfossil type	Number counted (14-cm depth) ¹	Per-cent	Number counted (28-cm depth) ¹	Per-cent	Number counted (42-cm depth) ¹	Per-cent	Number counted (56-cm depth) ¹	Per-cent	Number counted (71-cm depth) ¹	Per-cent
Biseriate spinose rods	8	4	10	5	9	4	35	16	55	25
Diatoms - centric (nonterrestrial component)	26	12	3	1	24	12	2	1	0 ²	0
Diatoms - pennate (nonterrestrial component)	44	21	55	27	62	31	13	6	0	0
Circle	31	15	34	17	18	9	6	3	5	2
Cross	0	0	0	0	0	0	0	0	1	0
Cyperaceae - type 1	2	1	1	0	8	4	8	4	0	0
Cyperaceae - type 2	0	0	2	1	0	0	4	2	61	28
Cyst	0	0	0	0	0	0	0	0	1	0
Dumbbell	3	1	6	3	3	1	5	2	2	1
Hair	0	0	0	0	1	0	0	0	1	0
Intertine (charcoal)	20	9	10	5	1	0	17	8	4	2
Irregular forms	52	25	51	25	47	23	69	31	52	24
Non-spinose rods	10	5	9	4	11	5	26	12	19	9
Saddle	6	3	18	9	13	6	13	6	10	5
Silicoflagellate (<i>Distephanus</i>)	1	0	0	0	0	0	0	0	0	0
Sphere	2	1	0	0	1	0	3	1	0	0
Sponge spicule (nonterrestrial component)	2	1	2	1	3	1	5	2	6	3
Uniseriate spinose rods	3	1	1	0	0	0	3	1	2	1
Wagon wheel	2	1	0	0	0	0	15	7	0	0
Total count	212	100	202	100	201	100	224	100	219	100

¹Compaction-corrected depth at sample midpoint.²Rare, broken fragments.Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).Last Updated on 2/5/98
By Gary R. Buell

TABLE 9. Carbonaceous spherule data for core SB1c, St. Bernard Parish, Louisiana.

St. Bernard core SB1c (location: 29°58'53" N, 89°55'27" W; sampled: 03/20/1996).

Percent compaction (applied linearly): 29.1.

Sample depth (cm) ¹	Total number of carbon spherules	Markers counted/ Total markers added during processing	Estimated number of carbon spherules/ sediment volume (1.2 cm ³) ²	Spherule type
14	135	360 / 12,542	4703	oil, coal, or wood
28	18	230 / 12,542	982	oil, coal, or wood
42	107	505 / 12,542	2657	oil
56	27	602 / 12,542	563	oil, coal, or wood
71	8	407 / 12,542	247	--
85	2	594 / 12,542	42	coal or wood

¹Compaction-corrected depth at sample midpoint.²Estimated number of carbon spherules per cubic centimeter of sediment (formula of Clarke and Patterson [1984]).Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).Last Updated on 2/5/98
By Gary R. Buell

TABLE 10A. Physical and chemical data for samples from core SB1a, St. Bernard Parish, Louisiana.

St. Bernard core SB1a (location: 29°58'53" N, 89°55'27" W; sampled 03/18/1996). Percent compaction (applied linearly): 6.8. OC, organic carbon. N, nitrogen. Ppt, parts per thousand. SU, pH standard units.

Sample I.D.	Material	Corrected depth (cm) ¹	Total OC (%) ²	Mineral matter (%) ³	$\delta^{13}\text{C}$ (ppt)	Total N (%)	$\delta^{15}\text{N}$ (ppt)	C:N ratio	H ₂ O, air-dried (%)	H ₂ O, 65°C (%)	Bulk density, air-dried (g/cm ³) ⁴	Bulk density, 65°C (g/cm ³) ⁴	pH (SU)	Weight fraction >2 mm (%)
--	fibrous peat ⁵	2.7	--	--	--	--	--	--	--	--	--	--	--	--
SB1a-1	same	6.8	26.2	47.6	-15.8	1.5	2.46	17.0	86.7	87.6	0.109	0.108	6	28.6
SB1a-2	same	9.6	26.4	47.3	-15.0	1.3	2.91	20.0	86.4	87.3	0.107	0.106	7	11.9
SB1a-3	same	12.3	14.8	70.3	-14.6	0.9	3.29	16.2	86.6	87.7	0.102	0.100	7	2.4
SB1a-4	same	15.1	15.0	70.1	-19.8	0.9	4.03	17.7	80.8	81.7	0.162	0.160	6	3.3
SB1a-5	same	17.8	26.0	48.1	-21.5	1.6	4.71	16.7	86.7	87.9	0.101	0.100	6	2.0
SB1a-6	same	20.5	27.8	44.4	-18.3	1.7	2.12	16.5	86.5	87.7	0.097	0.096	6	10.6
SB1a-7	same	23.3	28.8	42.3	-19.2	1.8	2.14	16.3	88.7	89.6	0.083	0.082	7	2.7
SB1a-8	same	26.0	27.3	45.3	-23.0	2.0	1.97	13.7	88.5	89.4	0.091	0.090	7	2.1
SB1a-9	same	28.8	25.6	48.8	-21.7	1.7	1.86	15.1	86.3	87.3	0.106	0.105	7	0.7
SB1a-10	same	31.5	23.7	52.6	-21.0	1.4	1.60	17.3	84.6	85.7	0.127	0.125	8	3.5
SB1a-11	same	34.2	12.3	75.4	-22.8	0.8	2.64	15.6	81.8	83.0	0.153	0.151	8	0.0
SB1a-12	same	37.0	17.3	65.4	-23.1	1.2	3.12	14.2	84.2	85.3	0.130	0.129	8	0.0
SB1a-13	same	39.5	23.1	53.8	-21.1	1.4	1.14	16.7	87.2	88.2	0.109	0.108	8	0.3
SB1a-14	same	42.2	29.8	40.4	-20.2	1.7	1.01	17.1	88.7	89.6	0.090	0.090	8	0.6
SB1a-15	same	45.0	30.0	40.1	-20.5	1.8	0.78	16.3	89.5	90.4	0.078	0.077	7	1.2
SB1a-16	same	47.7	29.7	40.7	-20.3	1.8	0.70	16.4	89.7	90.5	0.080	0.079	7	0.5
SB1a-17	same	50.4	31.2	37.6	-20.2	1.8	0.30	17.7	90.6	91.1	0.075	0.074	8	3.7
SB1a-18	same	53.1	31.7	36.6	-21.6	1.9	0.58	16.8	90.3	91.1	0.081	0.080	7	2.3
SB1a-19	same	55.8	32.8	34.5	-21.3	1.9	0.45	17.3	90.3	91.1	0.084	0.083	8	0.3
SB1a-20	same	58.6	33.4	33.2	-21.6	2.0	0.39	17.0	89.4	90.3	0.086	0.085	8	0.3
SB1a-21	same	61.3	35.4	29.2	-19.5	2.1	-0.18	16.8	89.4	90.4	0.091	0.090	8	0.2
SB1a-22	same	64.0	38.1	23.8	-22.0	2.1	-0.92	18.0	90.7	91.6	0.087	0.086	8	0.2
SB1a-23	same	66.7	39.2	21.5	-23.0	2.2	-0.56	18.0	90.9	91.7	0.094	0.093	8	0.0
SB1a-24	same ⁶	69.3	36.8	26.5	-23.2	2.2	0.06	16.6	90.5	91.4	0.084	0.083	7	0.7
SB1a-25	same ⁶	72.2	39.2	21.5	-24.5	2.0	-0.01	20.0	89.8	90.9	0.090	0.089	7	5.1
SB1a-26	same ⁶	74.9	36.0	28.0	-23.2	1.8	-0.79	19.8	89.7	90.8	0.083	0.082	7	1.0
SB1a-27	same ⁶	77.6	32.8	34.4	-21.4	1.7	-0.06	19.6	86.9	88.2	0.111	0.109	7	0.9
SB1a-28	same ⁶	80.4	26.2	47.6	-21.2	1.4	-1.23	19.3	87.6	88.7	0.102	0.100	7	6.1
SB1a-29	same ⁶	83.1	33.4	33.2	-19.5	1.7	-1.04	19.8	87.8	88.9	0.099	0.098	7	0.7
SB1a-30	same ⁶	85.8	33.6	32.8	-20.1	1.8	-1.04	19.1	88.4	89.4	0.091	0.090	7	1.4
SB1a-31	same ⁶	88.5	31.9	36.2	-21.5	2.0	-0.84	15.9	87.6	88.8	0.100	0.099	6	4.2
SB1a-32	clay	91.3	31.7	36.6	-22.0	1.6	-1.18	19.4	85.4	86.7	0.118	0.116	6	0.0
SB1a-33	peat ⁷	94.0	25.7	48.6	-22.9	1.5	-0.74	17.0	84.2	85.6	0.131	0.129	7	0.0
SB1a-34	peat ⁷	96.7	23.6	52.8	-22.7	1.4	-0.56	16.4	84.0	85.4	0.150	0.147	7	0.0

¹Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.0743 to correct for compaction.

²Total organic carbon = total carbon - inorganic carbon. Inorganic carbon less than 0.01% for all samples.

³Mineral matter (percent) = 100 - (2 x organic carbon).

⁴Bulk-density values reported to three decimal places for visual comparison of air-dried and oven-dried (65°C) values.

⁵Top 2 inches (5.08 cm) discarded in the field. No sample for corrected depth at 2.7 cm.

⁶More compact (dense) than overlying interval.

⁷Peat below clay lense is denser and finer than peat above clay lense.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

TABLE 10B. Major-element and trace-element data for samples from core SB1a, St. Bernard Parish, Louisiana.

St. Bernard core SB1a (location: 29°58'53" N., 89°55'27" W; sampled 03/18/1996). Percent compaction (applied linearly): 6.8. Concentrations of Al, Fe, Na, K, Mg, and Ca reported in percent. All other concentrations reported in micrograms per gram (µg/g, equivalent to parts per million [ppm]).

Sample I.D.	Cor- rected depth (cm) ¹	Al	Fe	Mn	Li	Na	K	Mg	Ca	Sr	Ba	La	P	As
-- ²	2.7	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1a-1	6.8	0.98	1.05	97	9	1.52	0.33	0.60	0.41	24	42	11	436	4
SB1a-2	9.6	0.96	1.05	85	9	1.91	0.36	0.62	0.34	22	43	12	300	4
SB1a-3	12.3	1.03	1.31	107	10	1.78	0.38	0.60	0.32	20	43	13	233	3
SB1a-4	15.1	0.65	1.11	84	6	1.31	0.26	0.49	0.33	21	37	12	190	2
SB1a-5	17.8	0.74	0.90	161	6	1.87	0.29	0.72	0.57	37	52	8	168	3
SB1a-6	20.5	0.73	0.65	59	6	1.94	0.27	0.68	0.55	31	40	7	215	2
SB1a-7	23.3	0.62	0.63	59	6	2.36	0.27	0.75	0.58	36	38	7	246	2
SB1a-8	26.0	0.59	0.56	49	5	2.50	0.25	0.75	0.66	38	44	7	264	3
SB1a-9	28.8	0.66	0.65	47	6	1.96	0.25	0.67	0.55	34	43	7	246	2
SB1a-10	31.5	0.80	0.70	52	8	1.84	0.32	0.63	0.45	27	45	9	227	3
SB1a-11	34.2	1.00	1.61	126	11	1.16	0.33	0.55	0.31	21	47	14	148	3
SB1a-12	37.0	0.92	1.40	130	9	1.26	0.31	0.55	0.36	23	45	12	248	4
SB1a-13	39.5	0.85	1.14	101	8	1.74	0.30	0.65	0.50	30	47	10	220	3
SB1a-14	42.2	0.58	0.82	79	5	1.82	0.24	0.62	0.50	34	41	7	276	3
SB1a-15	45.0	0.56	0.66	62	5	1.94	0.22	0.65	0.54	33	38	6	245	3
SB1a-16	47.7	0.63	0.82	73	5	1.90	0.22	0.65	0.54	35	39	6	255	3
SB1a-17	50.4	0.51	0.78	69	5	1.94	0.20	0.67	0.54	34	38	5	227	3
SB1a-18	53.1	0.47	0.66	66	3	2.02	0.20	0.67	0.64	36	40	5	298	4
SB1a-19	55.8	0.47	0.88	67	4	2.13	0.19	0.66	0.59	34	43	5	236	3
SB1a-20	58.6	0.41	0.66	66	3	1.84	0.17	0.61	0.58	36	44	5	244	3
SB1a-21	61.3	0.25	0.28	49	2	2.18	0.13	0.65	0.60	35	37	3	166	4
SB1a-22	64.0	0.21	0.23	51	1	2.29	0.13	0.70	0.65	37	38	2	183	3
SB1a-23	66.7	0.36	0.18	68	1	1.78	0.13	0.66	0.71	41	44	3	205	3
SB1a-24	69.3	0.48	0.20	86	1	1.93	0.12	0.70	0.72	41	43	3	152	3
SB1a-25	72.2	0.23	0.37	51	1	1.6	0.10	0.61	0.64	36	39	2	137	2
SB1a-26	74.9	0.29	0.32	53	1	1.73	0.13	0.63	0.71	38	42	3	166	4
SB1a-27	77.6	0.39	0.24	55	3	1.4	0.16	0.60	0.60	35	47	4	152	3
SB1a-28	80.4	0.54	0.34	50	4	1.35	0.19	0.54	0.50	29	50	6	134	3
SB1a-29	83.1	0.46	0.32	53	4	1.36	0.18	0.53	0.48	30	53	5	115	2
SB1a-30	85.8	0.46	0.30	55	4	1.51	0.18	0.58	0.54	31	52	5	145	3
SB1a-31	88.5	0.47	0.30	63	3	1.56	0.16	0.60	0.61	35	60	6	164	3
SB1a-32	91.3	0.61	0.43	65	5	1.26	0.20	0.53	0.52	31	63	7	116	3
SB1a-33	94.0	0.69	0.63	68	6	1.14	0.22	0.50	0.46	28	65	8	143	2
SB1a-34	96.7	0.94	0.60	77	8	1.11	0.25	0.54	0.46	26	68	8	160	3

¹Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.0473 to correct for compaction.

²Top 2 inches (5.08 cm) discarded in the field. No sample for corrected depth at 2.7 cm.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

Last Updated on 2/5/98
By Gary R. Buell

TABLE 10C. Trace-element data for samples from core SB1a, St. Bernard Parish, Louisiana.

St. Bernard core SB1a (location: 29°58'53" N., 89°55'27" W; sampled 03/18/1996). Percent compaction (applied linearly): 6.8. Element concentrations reported in micrograms per gram (µg/g, equivalent to parts per million [ppm]). Silver (Ag) concentrations were <1 µg/g for all samples, so the data column is not shown.

Sample I.D.	Cor- rected depth (cm) ¹	Ti	V	Cr	Co	Ni	Cu	Zn	Y	Mo	Pb	Cd	Sn	Ce	Th	U
-- ²	2.7	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1a-1	6.8	42	50	24	5	63	50	269	2	5	44	4	5	25	7	24
SB1a-2	9.6	37	55	23	4	79	42	80	2	5	39	1	3	25	9	26
SB1a-3	12.3	33	45	24	8	244	75	56	3	5	32	1	4	30	11	22
SB1a-4	15.1	46	36	17	9	554	120	<1	2	3	26	1	3	29	7	10
SB1a-5	17.8	30	49	18	4	158	46	13	2	3	36	1	7	18	6	9
SB1a-6	20.5	44	43	19	2	56	38	79	2	4	34	1	4	16	5	7
SB1a-7	23.3	38	898	17	2	45	33	48	2	2	44	1	4	16	4	8
SB1a-8	26.0	42	47	13	2	39	21	189	2	1	31	1	9	15	2	5
SB1a-9	28.8	51	696	17	3	38	19	77	2	1	20	1	8	18	3	5
SB1a-10	31.5	48	44	19	2	48	25	120	2	2	22	1	12	22	7	4
SB1a-11	34.2	39	39	23	6	20	16	149	3	1	29	1	2	33	12	5
SB1a-12	37.0	43	40	20	5	19	19	346	3	1	32	1	5	27	9	5
SB1a-13	39.5	42	41	17	5	56	31	257	3	2	27	1	13	22	7	5
SB1a-14	42.2	50	38	10	4	37	22	365	2	2	32	1	8	16	5	4
SB1a-15	45.0	47	32	15	3	38	22	302	2	3	21	1	12	13	4	4
SB1a-16	47.7	54	34	15	3	36	21	166	2	1	25	1	25	13	4	5
SB1a-17	50.4	53	29	11	3	47	24	193	1	1	22	1	22	13	3	4
SB1a-18	53.1	61	29	13	3	42	17	165	1	2	22	1	15	12	3	4
SB1a-19	55.8	68	28	10	2	44	19	192	1	1	15	1	88	11	3	3
SB1a-20	58.6	60	26	12	2	28	15	340	1	2	13	<1	18	11	3	3
SB1a-21	61.3	43	20	5	1	13	6	248	1	1	6	1	5	6	1	2
SB1a-22	64.0	46	20	5	1	17	6	461	1	1	14	1	4	5	1	2
SB1a-23	66.7	45	19	9	1	16	6	149	1	1	9	1	5	7	2	2
SB1a-24	69.3	43	17	6	1	33	10	370	1	2	10	<1	3	7	2	2
SB1a-25	72.2	59	16	5	1	24	5	309	1	1	9	1	31	5	2	2
SB1a-26	74.9	50	19	6	1	13	4	<1	1	1	<1	<1	30	7	2	2
SB1a-27	77.6	54	21	7	1	18	7	<1	1	1	<1	<1	8	10	3	2
SB1a-28	80.4	55	25	10	1	43	13	<1	1	1	<1	<1	9	12	5	2
SB1a-29	83.1	66	20	7	1	23	13	<1	1	2	1	<1	7	11	4	2
SB1a-30	85.8	48	20	7	1	21	10	<1	1	1	2	<1	7	11	4	2
SB1a-31	88.5	43	18	7	1	20	11	17	1	1	6	<1	5	13	4	3
SB1a-32	91.3	49	24	12	2	27	9	3	2	1	7	<1	2	16	5	3
SB1a-33	94.0	36	26	15	2	36	14	193	2	1	9	1	5	19	7	3
SB1a-34	96.7	47	30	16	2	30	11	271	2	3	11	<1	12	19	7	4

¹Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.0473 to correct for compaction.

²Top 2 inches (5.08 cm) discarded in the field. No sample for corrected depth at 2.7 cm.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

Last Updated on 2/5/98
By Gary R. Buell

TABLE 11A. Physical and chemical data for samples from core SB1c, St. Bernard Parish, Louisiana.

St. Bernard core SB1c (location: 29°58'53" N., 89°55'27" W.; sampled 03/20/1996). Percent compaction (applied linearly): 29.1. OC, organic carbon. N, nitrogen. Ppt, parts per thousand.

Sample I.D. ¹	Material	Sampled interval (cm)	Uncorrected depth (cm)	Corrected depth (cm) ²	Total OC (%) ³	Mineral matter (%) ⁴	δ ¹³ C (ppt)	Total N (%)	δ ¹⁵ N (ppt)	C:N ratio	H ₂ O, air-dried (%)	Bulk density, air-dried (g/cm ³)
SB1c-10	fibrous peat	0-55	10	14.1	--	--	--	--	--	--	89	0.1
SB1c-21.5	same	0-55	21.5	30.3	31.9	36.2	-15.8	1.9	0.57	17	--	--
SB1c-40	same	0-55	40	56.4	34	32	-19	1.9	0.34	18	--	--
SB1c-45	same	0-55	45	63.5	--	--	--	--	--	--	89.7	0.09
SB1c(65)	fibrous peat, more compact	55-69	65	91.7	32.7	34.7	-20	1.8	-1.53	18.3	--	--
SB1c(79)	fibrous peat as above, color change	69-82.5	79	111.4	20.9	58.1	-20.5	1.1	-1.65	19	--	--
SB1c(83)	clay	82.5-86.5	83	117	10.9	78.2	-23.5	0.6	-0.71	17.9	--	--
SB1c-84	clay	82.5-86.5	84	118.4	--	--	--	--	--	--	63.4	0.28
SB1c(97)	fibrous peat	86.5-121	97	136.8	21.2	57.7	-19.4	0.9	-1.45	22.8	--	--
SB1c-108	same	86.5-121	108	152.3	--	--	--	--	--	--	80.8	0.14
SB1c(110)	same	86.5-121	110	155.1	26.1	47.8	-19.9	1.3	-1.29	20.9	--	--
SB1c(128)	peat changing downward	121-140.5	128	180.5	21.6	56.8	-20.2	1	-0.88	20.8	--	--
SB1c(145)	organic clay	140.5-162.5	145	204.5	10.3	79.5	-16	0.4	-0.89	29.3	--	--
SB1c(145)D	same	140.5-162.5	145	204.5	10.2	79.5	--	--	--	--	--	--
SB1c-158	same	140.5-162.5	158	222.8	--	--	--	--	--	--	71.4	0.23
SB1c-158D	same	140.5-162.5	158	222.8	--	--	--	--	--	--	--	0.23
SB1c(177)	fibrous peat	162.5-218	177	249.6	38.2	23.6	-19.7	1.6	-1.03	23.4	--	--
SB1c-194	same	162.5-218	194	273.5	--	--	--	--	--	--	85.7	0.09
SB1c(202)	same	162.5-218	202	284.8	37.7	24.7	-26	2.3	-0.33	16.7	--	--
SB1c-227.5	organic clay	218-233.5	227.5	320.8	--	--	--	--	--	--	62.4	0.29
SB1c(229)	same	218-233.5	229	322.9	12.6	74.8	-26.1	0.7	-0.43	17.5	--	--
SB1c(250)	peat	233.5-277	250	352.5	37.3	25.5	-26.8	2.1	-0.58	17.8	--	--
SB1c-251	same	233.5-277	251	353.9	--	--	--	--	--	--	81.2	0.12
SB1c(265)	same	233.5-277	265	373.7	32.7	34.5	-26.5	2	-0.09	16.4	--	--
SB1c-280	gray clay	277-285	280	394.8	--	--	--	--	--	--	--	--
SB1c(284)	same	277-285	284	400.4	1.8	96.3	-28.2	< 0.05	5.05	--	--	--

¹Chemical analyses were not done on samples with depths in parentheses. "D", duplicate sample.

²Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.41 to correct for compaction.

³Total organic carbon = total carbon - inorganic carbon. Inorganic carbon less than 0.01% for all samples.

⁴Mineral matter (percent) = 100 - (2 x organic carbon).

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

TABLE 11B. Major-element and trace-element data for samples from core SB1c, St. Bernard Parish, Louisiana.

St. Bernard core SB1c (location: 29°58'53' N., 89°55'27' W.; sampled 03/20/1996). Percent compaction (applied linearly): 29.1. Concentrations of Al, Fe, Na, K, Mg, and Ca reported in percent. All other concentrations reported in micrograms per gram (µg/g, equivalent to parts per million [ppm]).

Sample I.D. ¹	Cor- rected depth (cm) ²	Al	Fe	Mn	Li	Na	K	Mg	Ca	Sr	Ba	La	P	As
SB1c-10	14.1	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-21.5	30.3	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-40	56.4	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-45	63.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(65)	91.7	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(79)	111.4	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(83)	117.0	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-84	118.4	1.41	1.14	80	16	0.74	0.40	0.81	0.30	69	97	14	129	2
SB1c(97)	136.8	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-108	152.3	0.63	0.58	86	8	1.18	0.23	0.73	0.54	105	132	11	226	2
SB1c(110)	155.1	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(128)	180.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(145)	204.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(145)D	204.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-158	222.8	0.83	0.80	73	8	0.79	0.27	0.57	0.28	53	124	16	110	2
SB1c-158D	222.8	0.79	0.88	80	10	0.74	0.32	0.58	0.30	58	129	17	143	3
SB1c(177)	249.6	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-194	273.5	0.29	0.31	130	2	1.99	0.16	0.84	0.79	104	173	4	233	4
SB1c(202)	284.8	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-227.5	320.8	1.22	1.09	116	13	0.55	0.43	0.74	0.28	54	177	16	78	2
SB1c(229)	322.9	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(250)	352.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-251	353.9	0.44	1.74	139	4	1.41	0.13	0.68	0.67	91	166	7	208	10
SB1c(265)	373.7	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-280	394.8	1.09	1.46	143	12	0.38	0.38	0.66	0.25	47	178	19	184	4
SB1c(284)	400.4	--	--	--	--	--	--	--	--	--	--	--	--	--

¹Chemical analyses were not done on samples with depths in parentheses. "D", duplicate sample.

²Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.41 to correct for compaction.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

Last Updated on 2/5/98
By Gary R. Buell

TABLE 11C. Trace-element data for samples from core SB1c, St. Bernard Parish, Louisiana.

St. Bernard core SB1c (location: 29°58'53' N., 89°55'27' W.; sampled 03/20/1996). Percent compaction (applied linearly): 29.1. Element concentrations reported in micrograms per gram (µg/g, equivalent to parts per million [ppm]). "<", concentration less than detection limit. Silver (Ag), Ti, and Sn concentrations were <1 µg/g for all samples, so the data columns are not shown.

Sample I.D. ¹	Cor- rected depth (cm) ²	V	Cr	Co	Ni	Cu	Zn	Y	Mo	Pb	Cd	Ce	Th	U
SB1c-10	14.1	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-21.5	30.3	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-40	56.4	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-45	63.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(65)	91.7	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(79)	111.4	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(83)	117.0	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-84	118.4	21	9	4	35	18	35	9	43	7	1	28	20	11
SB1c(97)	136.8	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-108	152.3	15	<1	2	7	9	<1	8	3	<1	<1	21	11	5
SB1c(110)	155.1	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(128)	180.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(145)	204.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(145)D	204.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-158	222.8	16	5	3	8	16	<1	10	9	8	<1	33	18	7
SB1c-158D	222.8	22	10	3	8	15	<1	11	4	9	<1	33	18	7
SB1c(177)	249.6	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-194	273.5	12	<1	1	8	6	<1	3	6	<1	<1	8	4	4
SB1c(202)	284.8	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-227.5	320.8	23	15	4	20	20	<1	10	6	16	59	32	21	8
SB1c(229)	322.9	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c(250)	352.5	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-251	353.9	16	<1	5	21	14	<1	5	8	<1	<1	13	8	10
SB1c(265)	373.7	--	--	--	--	--	--	--	--	--	--	--	--	--
SB1c-280	394.8	24	9	8	24	22	<1	14	<1	<1	<1	37	22	4
SB1c(284)	400.4	--	--	--	--	--	--	--	--	--	--	--	--	--

¹Chemical analyses were not done on samples with depths in parentheses. "D", duplicate sample.

²Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.41 to correct for compaction.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** (H.W. Markewich, ed.).

Last Updated on 2/5/98
By Gary R. Buell

TABLE 12. Mercury concentrations for selected samples from core SB1c, St. Bernard Parish, Louisiana.		
St. Bernard core SB1c (location: 29°58'53' N, 89°55'27' W; sampled: 03/20/1996).		
Percent compaction (applied linearly): 29.1.		
Analysis for mercury by USGS Sediment Partitioning Research Laboratory, Atlanta, Georgia. Ppm, parts per million (equivalent to micrograms per gram).		
Sample I.D.	Corrected depth sample midpoint (cm) ¹	Mercury concentration (ppm)
SB1c-40	56.4	0.07
SB1c-145	204.5	0.04
SB1c-152	214.3	0.05
SB1c-224	315.8	0.05
SB1c-259	365.2	0.04
¹ Corrected depth equals original depth (midpoint of sample interval) multiplied by 1.41 to correct for compaction.		
Source: U.S. Geological Survey Open-File Report 98-36. Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana (H.W. Markewich, ed.).		

Last Updated on 2/5/98
By Gary R. Buell

TABLE 13. Organic carbon cumulative mass and age-interval accumulation rates for cores Sb1a and SB1c, St. Bernard Parish, Louisiana.

St. Bernard cores SB1a and SB1c (location: 29°58"53' N, 89°55"27' W; SB1a sampled 03/18/1996, SB1c sampled 03/20/1996).

Corrected depth intervals by age range (cm) ¹	Assignments by age interval ²	Interval period (yr)	Organic carbon cumulative mass (kg m ⁻²) ³		Organic carbon accumulation rate (kg m ⁻² yr ⁻¹)	
			SB1a	SB1c	SB1a	SB1c ⁴
0-15	1964-96	32	3.85	4.99	0.12	0.16
15-28	1952-64	12	7.16	9.20	0.28	0.35
28-52	1905-52	47	13.04	16.71	0.13	0.16
52-85	1830-1905	75	23.40	27.53	0.14	0.14
85-127	870 yr BP-1830	704	--	41.50	--	0.02-0.02
127-166	1140-870 yr BP	270	--	53.76	--	0.03-0.08
166-236	1920-1140	780	--	75.73	--	0.02-0.03
236-265	1980-1920	60	--	85.25	--	0.06-xxx ⁵
265-305	2360-1980	380	--	98.24	--	0.03-0.05
305-345	2670-2360	310	--	113.85	--	0.04-0.08
345-384	3180-2670	510	--	129.37	--	0.03-0.04

¹Depths corrected for compaction, 6.8% (core SB1a) and 29.1% (core SB1c). Depths for 1964 (15 cm) and 1952 (28 cm) were taken from ¹⁴C data (fig. 6) for core SB1a and used both for cores SB1a and SB1c.

²Ages determined from radiocarbon analysis and diagnostic stratigraphic markers. Greatest uncertainty in ages between 85 and 195 cm (see section *Data Interpretation and Summary*).

³Organic carbon (OC) cumulative mass = Σ (OC x bulk density x 10000) / 1000.

⁴Minimum and maximum accumulation rates are reported for intervals in SB1c below 85 cm based on the errors in ¹⁴C age determinations (table 1).

⁵A maximum accumulation rate cannot be calculated for this interval because the ages for the interval boundaries "cross over". The assigned ages of 1980 yr BP and 1920 yr BP may not be statistically different.

Source: U.S. Geological Survey Open-File Report 98-36. **Carbon storage and late Holocene chronostratigraphy of a Mississippi River deltaic marsh, St. Bernard Parish, Louisiana** H.W. Markewich, ed.).

Last Updated on 2/5/98
By Gary R. Buell


US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

[illegible]

CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 2. Physiographic provinces of southern Louisiana. Numerous gulfward-dipping Pleistocene deposits underlie the untitled area north of the Deltaic and Chenier plains.



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 3. Major Holocene deltaic lobes of the Mississippi River (modified after Frazier, 1967): Maringouin, 6-8 ka; Teche, 5.8-3.5 ka; St. Bernard, 3.5-2 ka; LaFourche, 2-1 ka; Plaquemine-Modern, 1 ka to the present. Red dot marks St. Bernard Parish core locality SB1 (See figure 4B.)



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 4A. Deltaic environments in the vicinity of the St. Bernard core locality; these are characteristic of progradation in this part of the Mississippi River deltaic plain. Vegetation associated with the environments of deposition: brackish marsh, predominantly *Spartina patens* (wiregrass) with minor amounts of *Scirpus olneyi* (three-cornered grass); natural levee and point bar, predominantly live oaks and water oaks, with minor amounts of pecan, hackberry, and bald cypress; swamp, predominantly bald cypress trees with minor amounts of tupelo gum, live oaks, water oaks, elms, hackberry, and sweet gum trees.



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 4B. Map showing St. Bernard core locality in relation to New Orleans, Chalmette, and the Mississippi River Gulf Outlet Canal (MR-GO canal). Fresh to brackish water is dark blue; Gulf of Mexico water is light blue. Names of water bodies are blue; parish names are brown.



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 5. Photograph of vibracore equipment used to drill the marsh site in St. Bernard Parish, southeastern LA.



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 6. ^{14}C bomb curve for core SB1a, St. Bernard Parish, LA. Data corrected for compaction and ^{13}C .



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 7. ^{137}Cs curve for core SB1b, St. Bernard Parish, LA.




CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 8. Pollen diagram for core SB1c, St. Bernard Parish, LA.


**Pollen Diagram
core SB1c, St. Bernard Parish, SE LA**



**CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER
DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA**

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 9. Photographic plate of diagnostic pollen and other microfossils for core SB1c, St. Bernard Parish, LA. A. TCT (Taxodiaceae/ Cupressaceae/Thuja); B. Cheno-Am (Chenopodiaceae-Amaranthaceae); C. *Typha latifolia*; D. Graminea; E. Cyperaceae; F. Compositae (high spine); G. *Vigna luteola*; H. foram test lining.



**CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER
DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA**

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 10A. Photo of 1845 map showing the historical geography of the Mississippi River from just east of New Orleans to the bird's foot delta (La Tourrette, 1845). The St. Bernard core locality is indicated by the white circle.



**CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER
DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA**


US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 10B. Photo of 1845 map showing: (1) the position of St. Bernard Parish core locality (white circle) on Conseil Plantation lot, and (2) the railroad (black line that parallels the Mississippi River on the north side and then runs eastward through plantation land) used for transportation of agriculture and wood products. This is a close up of part of map in figure 10A (La Tourrette, 1845).



US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


Phytolith - Siliceous Microfossil Diagram
core SB1c, St. Bernard Parish, SE LA



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 12. Diagnostic phytoliths and other siliceous microfossils present in core SB1c, St. Bernard Parish, LA. A. Dumbbell, produced by various grasses; B. Biserial spinose rod; C. Non-spinose rod; D. In situ dumbbell phytoliths in a modern sugarcane fragment; E. Cyperaceae (type); F. Cyperaceae (type), brownish does not show but is probably due to burning from fire; G. centric and pennate diatoms.



**CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER
DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA**

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 13. Concentration diagram of charcoal and carbon spherules, core SB1c, St. Bernard Parish, Southeast Louisiana



**CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER
DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA**

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 14. Carbonaceous spherules present in core SB1c, St. Bernard Parish, LA.



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 15. Composite diagram of diagnostic microfossils from core SB1c, St. Bernard Parish, LA.



CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 16. Organic carbon (A), nitrogen (B), and bulk density (C) profiles for cores SB1a and SB1c, St. Bernard Parish, LA.







CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA


US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36.


FIGURE 17. Elemental chemistry profiles for cores SB1a and SB1c, St. Bernard Parish, LA: (A) aluminum, (B) iron, (C) manganese; (D) lithium, (E) sodium, (F) potassium; (G) calcium, (H) magnesium, (I) strontium; (J) arsenic, (K) cadmium, (L) lead; (M) copper, (N) nickel, (O) zinc; (P) vanadium, (Q) chromium, (R) cobalt, (S) lanthanum, (T) cerium, and (U) thorium.












CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36


FIGURE 18. Carbon storage: (A) incremental; (B) cumulative for cores SB1a and SB1c, St. Bernard Parish, LA.




CARBON STORAGE AND LATE HOLOCENE CHRONOSTRATIGRAPHY OF A MISSISSIPPI RIVER DELTAIC MARSH, ST. BERNARD PARISH, LOUISIANA

US GEOLOGICAL SURVEY OPEN-FILE REPORT 98-36

FIGURE 19. Plots of carbon mass/area versus time (years before present) for the SB1c core, St. Bernard Parish, Louisiana. (A) arithmetic plot showing apparent difference in recent (first four data points) and long-term rates (B) semi-log plot showing an exponential relation, and (C) log/log plot showing, that once the corrected age data become available, a decay constant can be calculated for the decomposition rates of marsh organic matter.



B



C

