

U.S. DEPARTMENT OF THE INTERIOR

U.S. GEOLOGICAL SURVEY

**Geochemical data for stream-sediment, heavy-mineral-concentrate, and rock samples
collected from the Fortyseven Creek gold-arsenic-antimony-tungsten prospect,
southwestern Alaska**

By

John E. Gray, Gregory K. Lee, Richard M. O'Leary, and Peter M. Theodorakos

Open-File Report 99-584

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards. Any use of trade names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

U.S. Geological Survey, P.O. Box 25046, MS 973, Federal Center, Denver, CO 80225-0046

1999

U.S. DEPARTMENT OF THE INTERIOR
Bruce Babbitt, *Secretary*
U.S. GEOLOGICAL SURVEY
Charles G. Groat, *Director*

For additional information

write to:

Chief Scientist, Central Region
Mineral Resources Program
U.S. Geological Survey
Box 25046, Federal Center
Denver, CO 80225-0046

Copies of this report can

be purchased from:

Books and Open-File Reports
U.S. Geological Survey
Federal Center
Box 25046
Denver, CO 80225-0046

TABLE OF CONTENTS

Introduction.....	4
Site Description and History.....	4
Geology.....	4
Methods of Study.....	5
Sample Media.....	5
Sample Preparation.....	5
Analytical Methods.....	5
Semiquantitative Spectrography.....	5
Inductively Coupled Plasma-Atomic Emission Spectrometry.....	6
Atomic Absorption Spectrophotometry.....	6
Visible Spectrophotometry.....	6
Selective Ion Analysis.....	6
X-ray Fluorescence.....	7
Coulometric Titration.....	7
Digital Data.....	7
Description of the Data Tables.....	7
Acknowledgments.....	7
References.....	7

FIGURES

Figure 1. Location of the Fortyseven Creek prospect.....	10
Figure 2. Location of samples collected in the Fortyseven Creek area	11

TABLES

Table 1. Limits of determination for the spectrographic analysis of stream-sediment samples.....	12
Table 2. Limits of determination for the analysis of stream-sediment and rock samples by several analytical techniques.....	13
Table 3. Geochemical data for stream-sediment samples in the Fortyseven Creek study.....	14
Table 4. Geochemical data for heavy-mineral-concentrate samples in the Fortyseven Creek study	17
Table 5. Geochemical data for rock samples in the Fortyseven Creek study.....	19

INTRODUCTION

In the summer of 1991, we conducted a reconnaissance geochemical survey around the Fortyseven Creek Au-As-Sb-W prospect that is located in the southwestern part of the Sleetmute quadrangle. At that time, this project was a small part of a more comprehensive Alaska Mineral Resource Assessment Program (AMRAP) study of the Sleetmute quadrangle. AMRAP studies were conducted by the U.S. Geological Survey (USGS) to fulfill requirements of the Alaska National Interests Lands Conservation Act (Public Law 96-487, 1980) to survey certain federal lands to determine their mineral potential. Although AMRAP is no longer in operation, this study represents a small topical study that was conducted during the Sleetmute quadrangle AMRAP study. The objective of the Fortyseven Creek work was to characterize the geochemistry of samples collected downstream from the Fortyseven Creek prospect, as well as mineralized and altered rock samples collected from the prospect. In this report, we describe the samples collected in 1991, the methods used for the analysis of the samples, and the geochemical data for these samples. The data in this report are also available in digital form on computer diskette in Gray and others (1999). An interpretation of these data appears in Gray and others (1998).

SITE DESCRIPTION AND HISTORY

The Fortyseven Creek Au-As-Sb-W prospect is located about 95 km southeast of Aniak, on the ridge crest at the headwaters of Fortyseven Creek, a tributary of the Holitna River (fig. 1). This lode and an associated gold placer mine on Fortyseven Creek about 3 km downstream from the prospect were discovered in 1947 by Russell Schaeffer. The lode was first described by Cady and others (1955) as a quartz-scheelite-gold-bearing shear zone cutting highly silicified graywacke and shale of the Kuskokwim Group. Additional exploration and geologic mapping of the Fortyseven Creek area was conducted by Hawley (1989), who described quartz veins cutting silicified sedimentary rock hornfels of the Kuskokwim Group; granite porphyry intrusions also cut the sedimentary rocks in this area. Hawley (1989) also found minor amounts of arsenopyrite, scheelite, stibnite, gold, pyrite, wolframite, jamesonite, and argentite in the veins, with locally abundant sericite (hydrothermal muscovite); these veins were traceable laterally for greater than 1,000 m. Some mineralized samples contain greater than 34 ppm Au (1 oz/t) (Hawley, 1989). The vein has not been mined. However, between 1948-1954, about 28 kg of gold (891 oz) and about 1,900 kg (5000 lbs) of scheelite, containing 67-77 percent WO_3 , were recovered by Schaeffer from the placer mine on Fortyseven Creek downstream from the lode (Hawley, 1989).

Several companies conducted exploration of the Fortyseven Creek area in the 1970's and 1980's including Homestake, Amax, American Copper and Nickel, and Anaconda (Hawley, 1989). More recently, from 1991-1994, Nevada Star Resources conducted exploration of the area that included minor prospect trenching, an extensive soil geochemical survey, and exploration drilling (Maynard, 1995). Nevada Star collected over 200 soil samples along the ridges at the headwaters of Fortyseven Creek and found highly elevated concentrations of Au (as much as 1.3 ppm), As (as much as 8,230 ppm), Bi (as much as 100 ppm), Cu (as much as 322 ppm), Sb (as much as 106 ppm), and W (as much as 510 ppm) in these soils. Nevada Star used soil anomalies of Au, As, and Bi to delineate several target areas for three exploration drill holes. Drilling was difficult due to circulation loss and hole caving problems. Consequently, only one drill hole penetrated significant mineralized rock about 44-105 m (145-345 ft) below the surface that contained as much as 1.2 ppm Au (Maynard, 1995).

The Fortyseven Creek prospect has geologic and geochemical characteristics similar to other gold deposits in southwestern Alaska such as the Au-As-W Golden Horn mine near Flat, and the Au-As-Sb Donlin Creek deposit (fig. 1). There is presently considerable exploration for such gold deposits in southwestern Alaska, but these deposits have not been well studied.

GEOLOGY

Rocks in the Fortyseven Creek area consist mostly of interbedded graywacke and siltstone of the Cretaceous Kuskokwim Group that are cut by small granite porphyry intrusions of probable Late Cretaceous or early Tertiary age (Cady and others, 1955; Hawley, 1989). The Kuskokwim Group is a sequence of flysch representing turbidite fan, foreslope, shallow-marine, and shelf facies deposited into an elongate, northeast trending, fault-controlled Cretaceous basin (Decker and Hoare, 1982; Bundtzen and Gilbert, 1983). Fossil ages for the Kuskokwim Group range from Albian to Campanian (Cady and others, 1955; Hoare and Coonrad, 1959; Box and Murphy, 1987; Box and Elder, 1992; Miller and Bundtzen, 1994). Granite porphyry intrusions in the Fortyseven Creek area are similar to those found elsewhere in southwestern Alaska. These intrusions are generally peraluminous in composition, with Al_2O_3 exceeding $Na_2O + K_2O + CaO$, and locally contain igneous garnet

(Bundtzen and Swanson, 1984; Moll-Stalcup, 1994). The granite porphyry near Fortyseven Creek has not been dated, but similar rocks in southwestern Alaska are Late Cretaceous and early Tertiary (about 72 to 61 Ma), based on K-Ar determinations (Reifenstuhl and others, 1984; Robinson and Decker, 1986; Decker and others, 1986; 1995; Miller and Bundtzen, 1994; Bundtzen and Miller, 1997). Gray and others (1998) obtained a $^{40}\text{Ar}/^{39}\text{Ar}$ plateau age of 67.1 ± 0.1 Ma for hydrothermal sericite separated from a sample of mineralized quartz vein from the Fortyseven Creek prospect. This age represents the best estimate of the timing of mineralization at Fortyseven Creek and is coeval with Late Cretaceous-early Tertiary granite porphyry intrusions found throughout southwestern Alaska. Late Cretaceous and early Tertiary magmatism is interpreted to be related to a broad subduction arc (Moll-Stalcup, 1994; Szumigala, 1993). Granite porphyry intrusions are important in southwestern Alaska because they show a close spatial and temporal association with Au-As-Sb-W and Hg-Sb veins (Cady and others, 1955; Bundtzen and Miller, 1997; Gray and others, 1997).

METHODS OF STUDY

Sample Media

In this study, stream-sediment and heavy-mineral-concentrate samples were collected to provide information about the rocks eroded from the drainage basin upstream from each sample site (fig. 2). Analyses of stream-sediment samples are representative of the geochemistry of rocks contained within the drainage basins. Heavy-mineral-concentrate samples provide information about higher density minerals that are eroded from rocks in the drainage basins. The heavy-mineral-concentrate collection procedure selectively concentrates the minerals with high specific gravity, many of which may be related to mineral deposits. Collection of heavy-mineral-concentrate samples facilitates concentration of some elements that are not easily detected in stream-sediment samples. Altered and mineralized rock samples were collected from the Fortyseven Creek prospect to characterize the geochemistry of this type of mineral deposit.

The stream-sediment material collected consisted of alluvium (bed sediment) in the active stream channel. Where possible, samples were composited by collecting sediment from several localities in the active channel. The stream sediment was screened to minus-10 mesh (2 mm) and collected in a stainless steel gold pan. About 2 kg of sediment was taken from the pan and saved as the stream-sediment sample.

Heavy-mineral-concentrate samples were collected from the same active alluvium (bed sediment) as the stream-sediment samples. At each sample site, the gold pan was filled with stream sediment screened to minus-10 mesh and panned until most of the less dense minerals, organic materials, and clays were removed. This resulting panned-concentrate sample was later processed in the laboratory as described below to obtain a heavy-mineral-concentrate sample.

Sample Preparation

In the laboratory, the stream-sediment samples were air dried, sieved to minus-80 mesh (0.18 mm), pulverized to less than 100 mesh (0.15 mm), and chemically analyzed by the methods described in the following section. The heavy-mineral-concentrate samples were sieved to minus-35 mesh (0.50 mm), and then separated using bromoform (specific gravity 2.85) to remove any remaining lighter minerals, primarily quartz and feldspar. The resultant heavy-mineral-concentrate samples were then separated into magnetic, paramagnetic, and nonmagnetic fractions using a modified Frantz Isodynamic Separator. The most magnetic material was removed at a setting of 0.25 ampere and contained mostly magnetite. The paramagnetic fraction was removed at 1.75 ampere and consisted largely of ferromagnesian silicates and iron oxides. The nonmagnetic fraction of the heavy-mineral-concentrate samples contained sulfide minerals, gold, and some nonmagnetic oxides and silicates; this was the only fraction chemically analyzed. Rock samples were crushed, ground, pulverized to less than 100 mesh, and analyzed by the analytical methods described below.

ANALYTICAL METHODS

Semiquantitative Spectrography (SQS)

The rock, minus-80-mesh stream-sediment, and minus-35-mesh nonmagnetic-heavy-mineral-concentrate samples were analyzed by semiquantitative, direct-current arc emission spectrography (SQS) using a technique adapted from Grimes and Marranzino (1968). Spectrographic results were determined by visually comparing spectra derived from the sample against spectra obtained from laboratory reference standards made from pure

oxides and carbonates. Standard concentrations are geometrically spaced over any given order of magnitude as follows: 100, 50, 20, 10, etc. Samples whose concentrations were estimated to fall between those values were assigned values of 70, 30, 15, etc. The precision of this analytical technique is approximately \pm one reporting interval at the 83 percent confidence level and \pm two reporting intervals at the 96 percent confidence level (Motooka and Grimes, 1976). Values determined for the major elements Fe, Mg, Ca, Na, Ti, and P are given in weight percent; all other values are in parts per million (micrograms/gram). In addition, Pd and Pt were determined in the heavy-mineral-concentrate samples by emission spectrography. The elements determined by emission spectrography and their limits of determination are listed in table 1. Data for stream-sediment samples determined by this technique are listed in table 3; the heavy-mineral-concentrate data are listed in table 4, and the rock data are listed in table 5.

Inductively Coupled Plasma-Atomic Emission Spectroscopy (ICP-AES)

Concentrations of Ag, As, Au, Bi, Cd, Cu, Mo, Pb, Sb, and Zn were determined in the rock and stream-sediment samples by inductively coupled plasma-atomic emission spectroscopy (ICP-AES) using the procedure developed by Motooka (1996). The sediments were decomposed with concentrated hydrochloric acid and hydrogen peroxide in a hot-water bath. The metals were extracted in diisobutyl ketone (DIBK) in the presence of ascorbic acid and potassium iodide. The DIBK phase was then aspirated directly into the plasma and element concentrations were determined simultaneously with a multichannel ICP instrument.

Atomic Absorption Spectrophotometry (AAS)

Concentrations of Au, Te, and Tl in the rock and stream-sediment samples were determined by an atomic absorption spectrophotometry (AAS) technique adapted from Hubert and Chao (1985). The samples were digested using a series of hydrogen peroxide, hydrofluoric acid, aqua-regia, and hydrobromic acid-bromine solutions. Gold, Te, and Tl were separated and concentrated by extraction into methyl isobutyl ketone and determined by flame atomic absorption spectrophotometry. Concentrations for Au in the range of 0.002 to 0.050 ppm were determined by graphite furnace atomic absorption spectrophotometry on samples that were shown to be less than 0.050 ppm by the flame atomic absorption spectrophotometry technique. The graphite furnace atomic absorption spectrophotometry technique for Au was adapted from Meier (1980).

Mercury was measured in the rock and stream-sediment samples using a modified version of the cold-vapor atomic AAS technique (Kennedy and Crock, 1987). The samples were decomposed with nitric acid and sodium dichromate. Mercury (II) was reduced to elemental mercury with hydroxylamine hydrochloride/sodium chloride and stannous chloride in a continuous flow system releasing the vapor into a quartz cell of an atomic absorption spectrophotometer where the concentration of mercury was determined.

The rock and stream-sediment samples were analyzed for Se using continuous-flow hydride generation AAS (Sanzolone and Chao, 1987). In this method, the stream-sediment samples were digested using nitric, perchloric, and hydrofluoric acids; hydrochloric acid was added to reduce Se (VI) to Se (IV), which is necessary for the accurate determination of Se using hydride generation AAS. A mixture of hydrochloric acid, sodium borohydride, and sodium hydroxide was added to produce selenium hydride, which is then stripped and transported with inert gas to the atomizer of the atomic absorption spectrophotometer where Se concentration was determined.

Visible Spectrophotometry (VS)

Tungsten was determined in the rocks and stream-sediment samples by a visible spectrophotometric (VS) method by decomposing the samples with nitric, hydrofluoric, and hydrochloric acids (Welsch, 1983). Stannous chloride and dithiol solution was added to reduce the W, forming the blue tungsten-dithiol complex, which was then extracted into heptane. The color intensity of the tungsten-dithiol complex is proportional to the concentration of W in the sample. Tungsten concentrations were determined using a visible absorption spectrophotometer.

Selective Ion Electrode Analysis (SIE)

Fluoride concentrations were determined in the rock and stream-sediment samples by selective ion electrode (SIE) analysis using a method modified from Bodkin (1977). Samples were fused with lithium metaborate, and then dissolved in nitric acid. A complexing buffer was added, and concentrations were determined using a fluoride-selective ion electrode.

X-Ray Fluorescence (XRF)

The rock samples were also analyzed for major element oxides by a wavelength dispersive X-ray fluorescence spectroscopy (XRF) method (Taggart and others, 1990). In this procedure the samples were fused with lithium tetraborate and formed into a glass disk. The disks were irradiated with X-rays, and X-ray photons emitted by the elements present in the sample were counted and concentrations determined using calibrated standards. The weight of the sample loss on ignition (LOI) at 925°C was determined gravimetrically. The elements determined by this technique and the concentration range limitations are shown in table 2.

Coulometric Titration (CT)

Concentrations of CO₂ and H₂O were measured in the rock samples. The amount of CO₂ was determined by a coulometric titration (CT) technique following sample digestion with perchloric acid (Jackson and others, 1987). Total water was determined by coulometry using a titration method after fusion with a PbO-PbCrO₄-CaCO₃ flux at 900°C (Norton and Papp, 1990). The amount of moisture (H₂O) was determined by weight loss after heating the samples at 110°C. The amount of H₂O⁺ was determined by the calculation: [H₂O_{Total} - H₂O = H₂O⁺]. Lower limits of determination for CO₂ and H₂O are shown in table 2.

DIGITAL DATA

The geochemical data in this report are also part of the USGS National Geochemical Database in Denver that contains both descriptive geological information and analytical data. This report may also be obtained on the USGS web site at: greenwood.cr.usgs.gov/pub/open-file-reports/ofr-99-0584/.

DESCRIPTION OF DATA TABLES

Table 3 contains the geochemical data for the stream-sediment samples collected during this study; table 4 contain the data for the heavy-mineral-concentrate samples, and table 5 contain the data for the rock samples. Sample site locations are given in latitude and longitude and decimal degrees in the tables, and these sample sites are shown on figure 2. The analytical method for each element shown in tables is abbreviated as a suffix in the column headings. The designation "SQS" indicates semiquantitative emission spectrography, "ICP" indicates inductively coupled plasma-atomic emission spectroscopy, "AAS" indicates atomic absorption spectrophotometry, "VS" indicates visible spectrophotometry, "SIE" indicates selective ion electrode analysis, "XRF" indicates X-ray fluorescence spectroscopy, and "CT" indicates coulometric titration. Discrepancies in analyses for certain elements duplicated by different analytical methods, such as values determined for Au in stream-sediment samples, may be attributable to the particulate nature of minerals that contain Au (i.e., nugget effect), different sample aliquots used, and different extraction procedures. Due to the larger sample aliquot analyzed, the atomic absorption spectrophotometry analysis of Au generally provides more representative results than emission spectrography. For example, a 10-gram sample aliquot is used for the atomic absorption spectrophotometry analysis, whereas a 10-milligram sample aliquot is used in the emission spectrography technique. There is no data for heavy-mineral-concentrate sample 91FS02C because there was insufficient material for geochemical analysis following preparation in the laboratory.

Acknowledgments--John Bullock, Phil Hageman, Jerry Motooka, Clara Papp, Dave Siems, and Eric Welsch (USGS) provided chemical analyses of samples. Carter Borden assisted with field work and sample collection.

REFERENCES

- Bodkin, J.B., 1977, Determination of fluorine in silicates by use of an ion-selective electrode following fusion with lithium metaborate: *The Analyst*, v. 102, no. 1215, p. 409-413.
- Box, S.E., and Elder, W.P., 1992, Depositional and biostratigraphic framework of the Upper Cretaceous Kuskokwim Group, southwest Alaska, *in* Bradley, D.C., and Ford, A.B., eds., *Geologic studies in Alaska by the U.S. Geological Survey, 1990: U.S. Geological Survey Bulletin 1999*, p. 8-16.

- Box, S.E., and Murphy, J.M., 1987, Late Mesozoic structural and stratigraphic framework, eastern Bethel quadrangle, southwest Alaska, *in* Hamilton, T.D., and Galloway, J.P., eds., *Geologic studies in Alaska by the U.S. Geological Survey during 1986: U.S. Geological Survey Circular 998*, p. 78-82.
- Bundtzen, T.K., and Gilbert, W.G., 1983, Outline of geology and mineral resources of upper Kuskokwim region, Alaska: *Journal of the Alaska Geological Society*, v. 3, p. 101-119.
- Bundtzen, T.K., and Miller, M.L., 1997, Precious metals associated with Late Cretaceous-early Tertiary igneous rocks of southwestern Alaska, *in* Goldfarb, R.J., and Miller, L.D., eds., *Mineral Deposits of Alaska: Economic Geology Monograph 9*, 242-286.
- Bundtzen, T.K., and Swanson, S.E., 1984, Geology and petrology of igneous rocks in the Innoko River area, western Alaska: *Geological Society of America Abstracts with Programs*, v. 16, no. 5, p. 273.
- Cady, W.M., Wallace, R.E., Hoare, J.M., and Webber, E.J., 1955, The central Kuskokwim region, Alaska: U.S. Geological Survey Professional Paper 268, 132 p.
- Decker, John, Bergman, S.C., Blodgett, R.B., Box, S.E., Bundtzen, T.K., Clough, J.G., Coonrad, W.L., Gilbert, W.G., Miller, M.L., Murphy, J.M., Robinson, M.S., and Wallace, W.K., 1994, Geology of southwestern Alaska, *in* Plafker, George, and Berg, H.C., eds., *The Geology of Alaska: Geological Society of America, Boulder, Colorado, The Geology of North America*, v. G-1, p. 285-310.
- Decker, John, and Hoare, J.M., 1982, Sedimentology of the Cretaceous Kuskokwim Group, southwest Alaska, *in* Coonrad, W.L., ed., *U.S. Geological Survey in Alaska, Accomplishments during 1980: U.S. Geological Survey Circular 844*, p. 81-83.
- Decker, John, Reifensuhl, R.R., Robinson, M.S., and Waythomas, C.F., 1986, Geologic map of the Sleetmute A-5, A-6, B-5, and B-6 quadrangles, Alaska: Alaska Division of Geological and Geophysical Surveys Professional Report 93, 22 p., 1 sheet, scale 1:250,000.
- Decker, John, Reifensuhl, R.R., Robinson, M.S., Waythomas, C.F., and Clough, J.G., 1995, Geology of the Sleetmute A-5, A-6, B-5, and B-6 quadrangles, southwestern Alaska: Alaska Division of Geological and Geophysical Surveys Professional Report 99, 16 p.
- Decker, John, Robinson, M.S., Murphy, J.M., Reifensuhl, R.R., and Albanese, M.D., 1984, Geologic map of the Sleetmute A-6 quadrangle: Alaska Division of Geological and Geophysical Surveys Report of Investigations 84-8, scale 1:40,000.
- Gray, J.E., Lee, G.K., O'Leary, R.M., and Theodorakos, P.M., 1999, Digital geochemical data for stream sediment, heavy-mineral-concentrate, and rock samples collected from the Fortyseven Creek gold-arsenic-antimony-tungsten prospect, southwestern Alaska: U.S. Geological Survey Open-File Report 99-584, 1.2 MB Diskette.
- Gray, J.E., Gent, C.A., Snee, L.W., and Wilson, F.H., 1997, Epithermal mercury-antimony and gold-bearing vein deposits of southwestern Alaska, *in* Goldfarb, R.J., and Miller, L.D., eds., *Mineral Deposits of Alaska: Economic Geology Monograph 9*, p. 287-305.
- Gray, J.E., Gent, C.A., Snee, L.W., and Theodorakos, P.M., 1998, Age, isotopic, and geochemical studies of the Fortyseven Creek Au-As-Sb-W prospect and vicinity, southwestern Alaska, *in* Gray, J.E., and Riehle, J.R., eds., *Geologic Studies by the U.S. Geological Survey, 1996: U.S. Geological Survey Professional Paper 1595*, p. 17-29.
- Grimes, D.J., and Marranzino, A.P., 1968, Direct-current arc and alternating-current spark emission spectrographic field methods for the semiquantitative analysis of geological materials: U.S. Geological Survey Circular 591, 6 p.
- Hawley, C.C., 1989, The Forty Seven Creek prospect, southwest Alaska, summary of data and recommendations: unpublished report prepared for Holitna Basin Mining and Exploration, Inc., 7 p.
- Hoare, J.M., and Coonrad, W.L., 1959, Geology of the Bethel quadrangle, Alaska: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-285, scale 1:250,000.
- Hubert, A.E., and Chao, T.T., 1985, Determination of gold, indium, tellurium and thallium in the same sample digestion of geological materials by atomic-absorption spectroscopy and two-step solvent extraction: *Talanta*, v. 32, p. 383-387.
- Jackson, L.L., Brown, F.W., and Neil, S.T., 1987, Major and minor elements requiring individual determination, classical whole rock analysis, and rapid rock analysis, *in* Baedeker, P.A., ed., *Geochemical Methods of Analysis: U.S. Geological Survey Bulletin 1770*.
- Kennedy, K.R., and Crock, J.G., 1987, Determination of mercury in geological materials by continuous flow, cold-vapor, atomic-absorption spectrophotometry: *Analytical Letters*, v. 20, p. 899-908.
- Maynard, G.B., 1995, 47 Creek Property Southwestern Alaska, Results of the Phase Three Exploration Program:

- unpublished report prepared for Nevada Star Resource Corp., Seattle, Washington.
- Meier, A.L., 1980, Flameless atomic-absorption determination of gold in geological materials: *Journal of Geochemical Exploration*, v. 13, p. 77-85.
- Miller, M.L., and Bundtzen, T.K., 1994, Generalized geologic map of the Iditarod quadrangle, Alaska, showing potassium-argon, major-oxide, trace-element, fossil, paleocurrent, and archaeological sample localities: U.S. Geological Survey Miscellaneous Field Studies Map MF-2219-A, 1 plate, scale 1:250,000.
- Moll-Stalcup, E.J., 1994, Latest Cretaceous and Cenozoic magmatism in mainland Alaska, *in* Plafker, George, and Berg, H.C., eds., *The Geology of Alaska*: Geological Society of America, Boulder, Colorado, *The Geology of North America*, v. G-1, p. 589-619.
- Motooka, J.M., 1996, Organometallic halide extraction for 10 elements by inductively coupled plasma-atomic emission spectrophotometry, *in* Arbogast, B.F., ed., *Analytical methods manual for the Mineral Resources Surveys Program*, U.S. Geological Survey: U.S. Geological Survey Open-File Report 96-525, p. 102-108.
- Motooka, J. M., and Grimes, D. J., 1976, Analytical precision of one-sixth order semiquantitative spectrographic analyses: U.S. Geological Survey Circular 738, 25 p.
- Norton, D.R., and Papp, C.S.E., 1990, Determination of moisture and total water in silicate rocks, *in* Arbogast, B.F., ed., *Quality Assurance Manual for the Branch of Geochemistry*, U.S. Geological Survey: U.S. Geological Survey Open-file Report 90-668, 184 p.
- Reifensuhl, R.R., Robinson, M.S., Smith, T.E., Albanese, M.D., and Allegro, G.A., 1984, Geologic map of the Sleetmute B-6 quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Report of Investigations 84-12, scale 1:40,000.
- Robinson, M.S., and Decker, John, 1986, Preliminary age dates and analytical data for selected igneous rocks from the Sleetmute, Russian Mission, Taylor Mountains, and Bethel quadrangles, southwestern Alaska: Alaska Division of Geological and Geophysical Surveys Public-Data File 86-99, 9 p.
- Sanzolone, R.F., and Chao, T.T., 1987, Determination of selenium in thirty-two geochemical reference materials by continuous-flow hydride generation atomic absorption spectrophotometry: *Geostandards Newsletter*, v. 11, p. 81-85.
- Szumigala, D.J., 1993, Gold mineralization related to Cretaceous-Tertiary magmatism in the Kuskokwim Mountains of west-central and southwestern Alaska: Ph.D. dissertation, University of California, Los Angeles, 301 p.
- Taggart, J.E., Jr., Bartel, Ardith, and Siems, D.F., 1990, High precision major element analysis of rocks and minerals by wavelength dispersive X-ray fluorescence spectroscopy, *in* Arbogast, B.F., ed., *Quality Assurance Manual for the Branch of Geochemistry*: U.S. Geological Survey: U.S. Geological Survey Open-file Report 90-668, 184 p.
- Welsch, E.P., 1983, A rapid geochemical spectrophotometric determination of tungsten with dithiol: *Talanta*, v. 30, p. 876-878.




Figure 1. Map showing location of the Fortyseven Creek prospect in southwestern Alaska. Geology generalized from Cady and others (1955), Hoare and Coonrad (1959), Decker and others (1984), Decker and others (1994), Decker and others (1995), Miller and Bundtzen (1994), and Bundtzen and Miller (1997).




Figure 2. Location of samples collected in the Fortyseven Creek study. Circles indicate stream-sediment and heavy-mineral-concentrate sites and triangles indicate rock sample sites.

Table 1. Limits of determination for the spectrographic analysis of stream-sediment samples, based on a 10-mg sample.

[The spectrographic limits of determination for heavy-mineral-concentrate samples are based on a 5-mg sample weight, and are therefore two reporting intervals higher than the limits given for stream sediments]

Elements	Lower determination limit	Upper determination limit
	Percent	
Iron (Fe)	0.05	20
Magnesium (Mg)	0.02	10
Calcium (Ca)	0.05	20
Sodium (Na)	0.2	5
Titanium (Ti)	0.002	1
Phosphorous (P)	0.2	10
	Parts per million	
Silver (Ag)	0.5	5,000
Arsenic (As)	200	10,000
Gold (Au)	10	500
Boron (B)	10	2,000
Barium (Ba)	20	5,000
Beryllium (Be)	1	1,000
Bismuth (Bi)	10	1,000
Cadmium (Cd)	20	500
Cobalt (Co)	10	2,000
Chromium (Cr)	10	5,000
Copper (Cu)	5	20,000
Gallium (Ga)	5	100
Germanium (Ge)	10	100
Lanthanum (La)	50	1,000
Manganese (Mn)	10	5,000
Molybdenum (Mo)	5	2,000
Niobium (Nb)	20	2,000
Nickel (Ni)	5	5,000
Lead (Pb)	10	20,000
Antimony (Sb)	100	10,000
Scandium (Sc)	5	100
Tin (Sn)	10	1,000
Strontium (Sr)	100	5,000
Thorium (Th)	100	2,000
Vanadium (V)	10	10,000
Tungsten (W)	20	10,000
Yttrium (Y)	10	2,000
Zinc (Zn)	200	10,000
Zirconium (Zr)	10	1,000
Palladium (Pd)*	5	1,000
Platinum (Pt)*	20	1,000

*Determined in heavy-mineral-concentrate samples only.

Table 2. Limits of determination for the analysis of stream-sediment and rock samples by several analytical techniques.

[ICP-AES, inductively coupled plasma-atomic emission spectroscopy; AA, atomic absorption spectrophotometry; VS, visible spectrophotometry; SIE, selective ion electrode analysis; CT, coulometric titration, and XRF, X-Ray fluorescence spectroscopy;].

Element	Analytical Method	Lower limit	Upper limit
Silver (Ag)	ICP-AES	0.045 ppm	1,500 ppm
Arsenic (As)	ICP-AES	0.6 ppm	3,000 ppm
Gold (Au)	ICP-AES	0.15 ppm	2,400 ppm
Bismuth (Bi)	ICP-AES	0.6 ppm	1,500 ppm
Cadmium (Cd)	ICP-AES	0.03 ppm	500 ppm
Copper (Cu)	ICP-AES	0.03 ppm	1,200 ppm
Molybdenum (Mo)	ICP-AES	0.09 ppm	1,500 ppm
Lead (Pb)	ICP-AES	0.6 ppm	12,000 ppm
Antimony (Sb)	ICP-AES	0.6 ppm	800 ppm
Zinc (Zn)	ICP-AES	0.03 ppm	500 ppm
Tellurium (Te)	AAS	0.050 ppm	
Thallium (Tl)	AAS	0.050 ppm	
Gold (Au)	AAS	0.002 ppm	
Mercury (Hg)	AAS	0.02 ppm	
Selenium (Se)	AAS	0.1 ppm	
Tungsten (W)	VS	1.0 ppm	
Fluoride (F)	SIE	200 ppm	
Tungsten (W)	ICP-MS	1.0 ppm	
Carbon dioxide (CO ₂)	CT	0.01 %	
Water (H ₂ O)	CT	0.1 %	
Calcium (CaO)	XRF	0.02 %	
Iron (Fe _(T) O ₃)	XRF	0.04 %	
Magnesium (MgO)	XRF	0.1 %	
Sodium (Na ₂ O)	XRF	0.15 %	
Phosphorous (P ₂ O ₅)	XRF	0.05 %	
Titanium (TiO ₂)	XRF	0.02 %	
Manganese (MnO)	XRF	0.01 %	
Silica (SiO ₂)	XRF	0.1 %	
Alumina (Al ₂ O ₃)	XRF	0.1 %	
Potassium (K ₂ O)	XRF	0.02 %	

Table 3. Geochemical data for stream-sediment samples collected from Fortyseven Creek.

Lab no.	Field no.	Latitude			Longitude			Latitude dec. deg.	Longitude dec. deg.
		deg.	min.	sec.	deg.	min.	sec.		
US03 D387843	91FS08S	61	3	10	-158	10	45	61.05280	-158.17920
US03 D387845	91FS03S	61	2	37	-158	8	47	61.04360	-158.14639
US03 D387846	91FS06S	61	3	4	-158	10	20	61.05110	-158.17220
US03 D387847	91FS01S	61	2	9	-158	6	38	61.03580	-158.11060
US03 D387849	91FS04S	61	2	48	-158	9	15	61.04670	-158.15421
US03 D387850	91FS00S	61	1	49	-158	5	22	61.03030	-158.08940
US03 D387851	91FS07S	61	3	17	-158	10	33	61.05470	-158.17580
US03 D387852	91FS02S	61	2	28	-158	7	58	61.04110	-158.13280
US03 D387853	91FS05S	61	2	57	-158	9	47	61.04920	-158.16310

Lab no.	Field no.	Ag	As	Au	Bi	Cd	Cu	Mo	Pb
		ICP-ppm							
US03 D387843	91FS08S	0.28	330	1.10	8.7	0.45	46	2.4	15
US03 D387845	91FS03S	0.15	140	<.10	<.67	0.20	33	1.3	11
US03 D387846	91FS06S	0.33	240	0.10	2.1	0.45	49	2.5	15
US03 D387847	91FS01S	0.10	110	<.10	<.67	0.15	11	0.62	7.6
US03 D387849	91FS04S	0.22	170	<.10	<.67	0.29	37	1.7	12
US03 D387850	91FS00S	0.12	8.3	<.10	<.67	0.08	9.6	0.19	8.8
US03 D387851	91FS07S	0.19	140	<.10	0.82	0.39	44	1.6	14
US03 D387852	91FS02S	0.21	150	<.10	<.67	0.25	27	1.6	12
US03 D387853	91FS05S	0.15	50	<.10	<.67	0.26	37	1.1	14

Lab no.	Field no.	Sb	Zn	Au	Se	Te	Tl	Hg	W
		ICP-ppm	ICP-ppm	AAS-ppm	AAS-ppm	AAS-ppm	AAS-ppm	AAS-ppm	VS-ppm
US03 D387843	91FS08S	29	150	1.000	0.7	0.25	0.3	0.8	800
US03 D387845	91FS03S	7.8	130	<0.002	0.4	0.05	0.2	1.5	110
US03 D387846	91FS06S	5.2	180	0.300	0.6	<0.05	0.3	0.7	350
US03 D387847	91FS01S	1.9	81	0.008	0.4	<0.05	0.4	0.6	11
US03 D387849	91FS04S	4.1	140	0.004	0.4	0.05	0.3	1.0	100
US03 D387850	91FS00S	0.9	73	0.100	0.2	<0.05	0.3	0.3	12
US03 D387851	91FS07S	2.7	170	0.004	0.6	0.05	0.3	0.8	25
US03 D387852	91FS02S	3.4	110	0.004	0.5	0.05	0.3	0.8	26
US03 D387853	91FS05S	3.3	130	0.008	0.5	0.10	0.3	2.0	50

Table 3. Geochemical data for stream-sediment samples collected from Fortyseven Creek.

Lab no.	Field no.	F SIE-ppm	CaO SQS-%	FeTO3 SQS-%	MgO SQS-%	MnO SQS-%	Na2O SQS-%	P2O5 SQS-%	TiO2 SQS-%
US03 D387843	91FS08S	700	0.10	6.6	1.2	0.19	1.2	<0.5	0.5
US03 D387845	91FS03S	300	<0.07	2.6	1.2	0.04	1.8	<0.5	0.3
US03 D387846	91FS06S	300	0.07	6.6	0.8	0.26	1.2	<0.5	0.5
US03 D387847	91FS01S	300	0.07	6.6	0.8	0.04	2.4	<0.5	0.3
US03 D387849	91FS04S	300	0.07	4.0	1.2	0.13	1.8	<0.5	0.5
US03 D387850	91FS00S	300	0.07	2.6	0.8	0.01	1.8	<0.5	0.3
US03 D387851	91FS07S	300	0.10	6.6	1.2	0.13	2.4	<0.5	0.8
US03 D387852	91FS02S	300	0.07	6.6	0.8	0.19	1.2	<0.5	0.5
US03 D387853	91FS05S	300	0.10	6.6	1.2	0.06	1.8	<0.5	0.5

Lab no.	Field no.	Ag SQS-ppm	As SQS-ppm	Au SQS-ppm	B SQS-ppm	Ba SQS-ppm	Be SQS-ppm	Bi SQS-ppm	Cd SQS-ppm
US03 D387843	91FS08S	<0.5	<200	<10	50	1000	1	<10	<20
US03 D387845	91FS03S	<0.5	<200	<10	15	500	<1	<10	<20
US03 D387846	91FS06S	<0.5	<200	<10	50	700	<1	<10	<20
US03 D387847	91FS01S	<0.5	<200	<10	15	500	<1	<10	<20
US03 D387849	91FS04S	<0.5	<200	<10	30	700	<1	<10	<20
US03 D387850	91FS00S	<0.5	<200	<10	20	500	<1	<10	<20
US03 D387851	91FS07S	<0.5	<200	<10	30	1000	<1	<10	<20
US03 D387852	91FS02S	<0.5	<200	<10	30	1000	<1	<10	<20
US03 D387853	91FS05S	<0.5	<200	<10	20	700	<1	<10	<20

Lab no.	Field no.	Co SQS-ppm	Cr SQS-ppm	Cu SQS-ppm	Ga SQS-ppm	Ge SQS-ppm	La SQS-ppm	Mo SQS-ppm	Nb SQS-ppm
US03 D387843	91FS08S	30	200	50	30	<10	<50	<5	<20
US03 D387845	91FS03S	10	150	30	30	<10	<50	<5	<20
US03 D387846	91FS06S	20	200	70	20	<10	<50	<5	<20
US03 D387847	91FS01S	10	100	10	20	<10	<50	<5	<20
US03 D387849	91FS04S	20	200	70	50	<10	<50	<5	<20
US03 D387850	91FS00S	<10	100	15	30	<10	<50	<5	<20
US03 D387851	91FS07S	30	150	70	30	<10	<50	<5	<20
US03 D387852	91FS02S	20	100	30	15	<10	<50	<5	<20
US03 D387853	91FS05S	20	200	50	30	<10	<50	<5	<20

Table 3. Geochemical data for stream-sediment samples collected from Fortyeven Creek.

Lab no.	Field no.	Ni SQS-ppm	Pb SQS-ppm	Sb SQS-ppm	Sc SQS-ppm	Sn SQS-ppm	Sr SQS-ppm	Th SQS-ppm	V SQS-ppm
US03 D387843	91FS08S	70	10	<100	7	50	<100	<100	150
US03 D387845	91FS03S	30	<10	<100	5	<10	<100	<100	70
US03 D387846	91FS06S	70	10	<100	7	<10	<100	<100	100
US03 D387847	91FS01S	30	<10	<100	<5	<10	<100	<100	70
US03 D387849	91FS04S	70	15	<100	7	10	<100	<100	150
US03 D387850	91FS00S	30	<10	<100	5	<10	<100	<100	100
US03 D387851	91FS07S	100	10	<100	7	30	<100	<100	150
US03 D387852	91FS02S	70	<10	<100	5	<10	<100	<100	200
US03 D387853	91FS05S	70	10	<100	7	<10	<100	<100	200

Lab no.	Field no.	W SQS-ppm	Y SQS-ppm	Zn SQS-ppm	Zr SQS-ppm	Description
US03 D387843	91FS08S	1000	10	<200	70	minus-80-mesh stream sediment
US03 D387845	91FS03S	50	<10	<200	50	minus-80-mesh stream sediment
US03 D387846	91FS06S	200	10	<200	70	minus-80-mesh stream sediment
US03 D387847	91FS01S	<20	<10	<200	30	minus-80-mesh stream sediment
US03 D387849	91FS04S	50	10	<200	100	minus-80-mesh stream sediment
US03 D387850	91FS00S	<20	<10	<200	150	minus-80-mesh stream sediment
US03 D387851	91FS07S	<20	10	<200	70	minus-80-mesh stream sediment
US03 D387852	91FS02S	<20	15	<200	70	minus-80-mesh stream sediment
US03 D387853	91FS05S	30	10	<200	70	minus-80-mesh stream sediment

Table 4. Geochemical data for nonmagnetic heavy-mineral-concentrate samples collected from Fortytseven Creek.

Lab no.	Field no.	Latitude			Longitude			Latitude	Longitude
		deg.	min.	sec.	deg.	min.	deg.	deg.	
US66 D388830	91FS08C	61	3	10	-158	10	45	61.05280 -158.17920	
US66 D388832	91FS03C	61	2	37	-158	8	47	61.04360 -158.14639	
US66 D388833	91FS06C	61	3	4	-158	10	20	61.05110 -158.17220	
US66 D388834	91FS01C	61	2	9	-158	6	38	61.03580 -158.11060	
US66 D388836	91FS04C	61	2	48	-158	9	15	61.04670 -158.15421	
US66 D388837	91FS00C	61	1	49	-158	5	22	61.03030 -158.08940	
US66 D388838	91FS07C	61	3	17	-158	10	33	61.05470 -158.17580	
US66 D388840	91FS05C	61	2	57	-158	9	47	61.04920 -158.16310	

Lab no.	Field no.	CaO SQS-%	FeTO3 SQS-%	MgO SQS-%	MnO SQS-%	Na2O SQS-%	P2O5 SQS-%	TiO2 SQS-%	Ag SQS-ppm
US66 D388830	91FS08C	7.0	<0.1	<0.1	0.004	<0.6	<1.1	0.3	7
US66 D388832	91FS03C	2.8	0.7	0.5	0.013	<0.6	3.4	>3.3	100
US66 D388833	91FS06C	9.8	0.4	0.2	0.019	<0.6	1.6	2.5	300
US66 D388834	91FS01C	1.4	0.9	1.2	0.026	<0.6	4.6	>3.3	70
US66 D388836	91FS04C	7.0	0.4	0.2	0.065	<0.6	4.6	>3.3	10
US66 D388837	91FS00C	0.7	0.3	0.2	0.003	<0.6	2.3	>3.3	30
US66 D388838	91FS07C	4.2	1.3	0.3	0.065	<0.6	3.4	>3.3	300
US66 D388840	91FS05C	9.8	0.3	0.2	0.019	<0.6	6.9	>3.3	150

Lab no.	Field no.	As SQS-ppm	Au SQS-ppm	B SQS-ppm	Ba SQS-ppm	Be SQS-ppm	Bi SQS-ppm	Cd SQS-ppm	Co SQS-ppm
US66 D388830	91FS08C	<500	<20	20	10000	<2	2000	<50	<20
US66 D388832	91FS03C	<500	500	1000	3000	5	70	<50	<20
US66 D388833	91FS06C	<500	1000	100	10000	<2	150	<50	<20
US66 D388834	91FS01C	<500	300	1000	200	<2	<20	<50	<20
US66 D388836	91FS04C	<500	100	70	5000	<2	200	<50	<20
US66 D388837	91FS00C	<500	500	150	200	<2	<20	<50	<20
US66 D388838	91FS07C	<500	>1000	300	>10000	<2	2000	<50	<20
US66 D388840	91FS05C	<500	300	70	5000	<2	<20	<50	<20

Table 4. Geochemical data for nonmagnetic heavy-mineral-concentrate samples collected from Fortyseven Creek.

Lab no.	Field no.	Cr SQS-ppm	Cu SQS-ppm	Ga SQS-ppm	Ge SQS-ppm	La SQS-ppm	Mo SQS-ppm	Nb SQS-ppm	Ni SQS-ppm
US66 D388830	91FS08C	<20	<10	<10	<20	<100	<10	<50	<10
US66 D388832	91FS03C	500	<10	<10	<20	150	<10	<50	<10
US66 D388833	91FS06C	20	<10	<10	<20	100	<10	<50	<10
US66 D388834	91FS01C	1000	<10	<10	<20	150	<10	<50	<10
US66 D388836	91FS04C	30	<10	<10	<20	<100	<10	<50	<10
US66 D388837	91FS00C	100	<10	<10	<20	<100	<10	<50	<10
US66 D388838	91FS07C	100	10	<10	<20	100	<10	<50	<10
US66 D388840	91FS05C	20	<10	<10	<20	<100	<10	<50	<10

Lab no.	Field no.	Pb SQS-ppm	Sb SQS-ppm	Sc SQS-ppm	Sn SQS-ppm	Sr SQS-ppm	Th SQS-ppm	V SQS-ppm	W SQS-ppm
US66 D388830	91FS08C	20	<200	<10	>2000	200	<200	30	>20000
US66 D388832	91FS03C	<20	1000	10	>2000	300	<200	150	10000
US66 D388833	91FS06C	30	<200	<10	>2000	300	<200	100	>20000
US66 D388834	91FS01C	<20	<200	10	>2000	<200	<200	150	2000
US66 D388836	91FS04C	<20	1000	<10	>2000	200	<200	100	20000
US66 D388837	91FS00C	<20	1000	<10	>2000	<200	<200	100	2000
US66 D388838	91FS07C	20	<200	<10	>2000	500	<200	150	10000
US66 D388840	91FS05C	<20	1000	<10	>2000	700	<200	100	>20000

Lab no.	Field no.	Y SQS-ppm	Zn SQS-ppm	Zr SQS-ppm	Pd SQS-ppm	Pt SQS-ppm	Description
US66 D388830	91FS08C	30	<500	50	<100	<100	heavy-mineral-concentrate
US66 D388832	91FS03C	150	<500	>2000	<100	<100	heavy-mineral-concentrate
US66 D388833	91FS06C	150	<500	>2000	<100	<100	heavy-mineral-concentrate
US66 D388834	91FS01C	70	<500	>2000	<100	<100	heavy-mineral-concentrate
US66 D388836	91FS04C	100	<500	>2000	<100	<100	heavy-mineral-concentrate
US66 D388837	91FS00C	150	<500	>2000	<100	<100	heavy-mineral-concentrate
US66 D388838	91FS07C	150	<500	>2000	<100	<100	heavy-mineral-concentrate
US66 D388840	91FS05C	70	<500	<20	<100	<100	heavy-mineral-concentrate

Table 5. Geochemical data for rock samples collected near Fortyseven Creek.

Lab no.	Field no.	Latitude			Longitude			Latitude dec. deg.	Longitude dec. deg.
		deg.	min.	sec.	deg.	min.	sec.		
US06 D387931	47EST	61	3	56	-158	10	32	61.06560	-158.17560
US06 D387932	47BSTDA	61	3	58	-158	10	49	61.06610	-158.18030
US06 D387934	47BST	61	3	58	-158	10	49	61.06610	-158.18030
US06 D387936	47DGW	61	3	56	-158	10	37	61.06560	-158.17700
US06 D387937	47CST	61	3	56	-158	10	44	61.06560	-158.17889
US06 D387940	47FST	61	3	57	-158	10	24	61.06580	-158.17329
US06 D387941	47FGW	61	3	57	-158	10	24	61.06580	-158.17329
US06 D387943	47XGW	61	3	22	-158	11	45	61.05610	-158.19580
US06 D387944	47GST	61	3	57	-158	10	17	61.06580	-158.17140
US06 D387946	47BGW	61	3	58	-158	10	49	61.06610	-158.18030
US06 D387948	47EGW	61	3	56	-158	10	32	61.06560	-158.17560
US06 D387950	47GGW	61	3	57	-158	10	17	61.06580	-158.17140
US06 D387951	47DST	61	3	56	-158	10	37	61.06560	-158.17700

Lab no.	Field no.	CaO	FeTO3	MgO	Na2O	P2O5	TiO2	MnO	SiO2
		XRF-%							
US06 D387931	47EST	0.46	10.6	3.00	1.03	0.42	0.72	0.27	60.7
US06 D387932	47BSTDA	0.32	7.07	1.85	0.89	0.28	0.93	0.19	63.1
US06 D387934	47BST	0.32	6.50	1.59	1.14	0.25	0.73	0.17	69.0
US06 D387936	47DGW	0.22	5.27	1.46	1.72	0.19	0.63	0.16	78.2
US06 D387937	47CST	0.32	6.67	2.28	1.03	0.30	0.87	0.08	63.1
US06 D387940	47FST	0.29	6.19	2.36	0.81	0.31	1.01	0.07	58.2
US06 D387941	47FGW	0.21	3.42	1.03	1.44	0.19	0.42	0.03	81.7
US06 D387943	47XGW	0.16	6.64	2.13	0.90	0.28	0.97	0.07	61.6
US06 D387944	47GST	0.32	6.10	2.05	1.10	0.32	0.87	0.04	63.0
US06 D387946	47BGW	0.62	3.69	0.64	1.68	0.20	0.40	0.16	81.2
US06 D387948	47EGW	0.26	3.88	1.17	1.60	0.22	0.43	0.03	80.4
US06 D387950	47GGW	0.19	4.63	1.36	1.32	0.20	0.60	0.06	76.1
US06 D387951	47DST	0.38	6.44	2.27	1.09	0.30	0.86	0.09	63.0

Lab no.	Field no.	Al2O3	K2O	Ag	As	Au	Bi	Cd	Cu
		XRF-%	XRF-%	ICP-ppm	ICP-ppm	ICP-ppm	ICP-ppm	ICP-ppm	ICP-ppm
US06 D387931	47EST	14.7	2.16	0.092	3.2	<0.10	<0.67	0.35	54
US06 D387932	47BSTDA	16.9	3.53	<0.067	72	<0.10	<0.67	0.24	53
US06 D387934	47BST	13.4	2.54	0.220	82	<0.10	<0.67	0.22	89
US06 D387936	47DGW	8.67	0.85	<0.067	9.4	<0.10	<0.67	0.26	33
US06 D387937	47CST	16.5	3.33	<0.067	6.7	<0.10	<0.67	0.06	28
US06 D387940	47FST	18.8	4.12	<0.067	2.6	<0.10	<0.67	0.14	50
US06 D387941	47FGW	7.78	1.05	0.140	8.5	<0.10	<0.67	0.09	23
US06 D387943	47XGW	17.8	3.74	<0.067	200	<0.10	<0.67	0.10	52
US06 D387944	47GST	16.6	3.27	0.078	5.7	<0.10	<0.67	0.13	55
US06 D387946	47BGW	7.57	1.17	0.098	40	<0.10	<0.67	0.14	28
US06 D387948	47EGW	8.26	1.06	0.097	9.5	<0.10	<0.67	0.09	22
US06 D387950	47GGW	10.0	1.61	0.085	11	<0.10	<0.67	0.11	36
US06 D387951	47DST	16.3	3.40	0.083	8.7	<0.10	<0.67	0.17	41

Table 5. Geochemical data for rock samples collected near Fortyseven Creek.

Lab no.	Field no.	Mo ICP-ppm	Pb ICP-ppm	Sb ICP-ppm	Zn ICP-ppm	Au AAS-ppm	Se AAS-ppm	Te AAS-ppm	Tl AAS-ppm
US06 D387931	47EST	1.0	14	1.1	190	<0.002	0.80	<0.05	0.15
US06 D387932	47BSTDA	0.88	6.4	5.0	180	0.033	0.60	<0.05	0.30
US06 D387934	47BST	0.37	12	4.7	160	0.003	0.90	<0.05	0.30
US06 D387936	47DGW	0.92	11	0.92	100	0.002	0.70	<0.05	0.05
US06 D387937	47CST	2.6	7.0	<0.067	110	<0.002	0.90	<0.05	0.15
US06 D387940	47FST	1.0	9.7	0.90	160	0.004	0.40	<0.05	0.30
US06 D387941	47FGW	1.1	9.7	0.70	89	<0.002	0.60	<0.05	0.05
US06 D387943	47XGW	1.0	5.1	6.7	130	0.002	0.70	<0.05	0.30
US06 D387944	47GST	2.1	15	1.3	170	<0.002	0.90	0.05	0.30
US06 D387946	47BGW	1.2	9.9	4.9	120	0.002	0.70	<0.05	0.30
US06 D387948	47EGW	1.3	9.7	0.78	92	<0.002	1.0	<0.05	0.10
US06 D387950	47GGW	0.79	13	0.79	120	0.004	0.60	<0.05	0.20
US06 D387951	47DST	1.1	11	1.4	170	0.004	1.2	<0.05	0.25

Lab no.	Field no.	Hg AAS-ppm	W S-ppm	F SIE-ppm	CO2 CT-%	H2O CT-%	H2O+ CT-%	H2O- CT-%	CaO SQS-%
US06 D387931	47EST	0.09	2	500	<0.01	5.2	4.7	0.6	0.07
US06 D387932	47BSTDA	0.03	3	500	<0.01	4.1	3.7	0.4	0.07
US06 D387934	47BST	0.05	2	500	<0.01	3.3	3.0	0.3	<0.07
US06 D387936	47DGW	0.09	2	400	<0.01	2.5	2.3	0.2	<0.07
US06 D387937	47CST	0.02	<1	400	<0.01	4.5	4.0	0.4	<0.07
US06 D387940	47FST	0.14	1	300	<0.01	5.6	4.9	0.7	<0.07
US06 D387941	47FGW	0.06	<1	<200	<0.01	1.9	1.8	0.1	<0.07
US06 D387943	47XGW	0.04	10	300	<0.01	4.3	3.9	0.4	<0.07
US06 D387944	47GST	0.13	<1	400	<0.01	4.6	4.3	0.3	0.07
US06 D387946	47BGW	0.03	3	300	0.66	1.5	1.4	0.1	0.07
US06 D387948	47EGW	0.06	<1	300	<0.01	2.1	2.0	0.1	<0.07
US06 D387950	47GGW	0.12	<1	<200	<0.01	3.0	2.8	0.2	<0.07
US06 D387951	47DST	0.05	3	400	<0.01	4.5	4.0	0.5	<0.07

Lab no.	Field no.	FeTO3 SQS-%	MgO SQS-%	MnO SQS-%	Na2O SQS-%	P2O5 SQS-%	TiO2 SQS-%	Ag SQS-ppm	As SQS-ppm
US06 D387931	47EST	13	2.5	0.13	1.8	<0.5	0.5	<0.5	<200
US06 D387932	47BSTDA	9.3	1.7	0.13	2.4	<0.5	0.8	<0.5	<200
US06 D387934	47BST	9.3	1.2	0.09	2.4	<0.5	0.5	<0.5	<200
US06 D387936	47DGW	6.6	0.5	0.06	1.2	<0.5	0.3	<0.5	<200
US06 D387937	47CST	6.6	1.7	0.02	1.8	<0.5	0.5	<0.5	<200
US06 D387940	47FST	6.6	1.7	0.01	1.8	<0.5	0.5	<0.5	<200
US06 D387941	47FGW	2.6	0.3	0.003	1.2	<0.5	0.2	<0.5	<200
US06 D387943	47XGW	6.6	1.7	0.02	1.8	<0.5	0.3	<0.5	<200
US06 D387944	47GST	4.0	1.7	0.009	2.4	<0.5	0.5	<0.5	<200
US06 D387946	47BGW	2.6	0.3	0.06	2.4	<0.5	0.3	<0.5	<200
US06 D387948	47EGW	2.6	0.5	0.004	1.8	<0.5	0.3	<0.5	<200
US06 D387950	47GGW	4.0	0.8	0.01	1.8	<0.5	0.3	<0.5	<200
US06 D387951	47DST	6.6	2.5	0.04	2.4	<0.5	0.5	<0.5	<200

Table 5. Geochemical data for rock samples collected near Fortyseven Creek.

Lab no.	Field no.	Au SQS-ppm	B SQS-ppm	Ba SQS-ppm	Be SQS-ppm	Bi SQS-ppm	Cd SQS-ppm	Co SQS-ppm	Cr SQS-ppm
US06 D387931	47EST	<10	15	700	<1	<10	<20	20	150
US06 D387932	47BSTDA	<10	100	2000	<1	<10	<20	50	300
US06 D387934	47BST	<10	70	1000	<1	<10	<20	20	200
US06 D387936	47DGW	<10	<10	300	<1	<10	<20	15	150
US06 D387937	47CST	<10	20	1500	<1	<10	<20	<10	200
US06 D387940	47FST	<10	30	2000	<1	<10	<20	15	300
US06 D387941	47FGW	<10	<10	200	<1	<10	<20	<10	50
US06 D387943	47XGW	<10	200	1000	<1	<10	<20	<10	300
US06 D387944	47GST	<10	30	2000	<1	<10	<20	15	200
US06 D387946	47BGW	<10	15	200	<1	<10	<20	10	50
US06 D387948	47EGW	<10	<10	300	<1	<10	<20	<10	50
US06 D387950	47GGW	<10	15	500	<1	<10	<20	15	70
US06 D387951	47DST	<10	20	1500	<1	<10	<20	15	300

Lab no.	Field no.	Cu SQS-ppm	Ga SQS-ppm	Ge SQS-ppm	La SQS-ppm	Mo SQS-ppm	Nb SQS-ppm	Ni SQS-ppm	Pb SQS-ppm
US06 D387931	47EST	50	30	<10	<50	<5	<20	70	10
US06 D387932	47BSTDA	50	50	<10	<50	<5	<20	70	15
US06 D387934	47BST	100	50	<10	<50	<5	<20	70	15
US06 D387936	47DGW	30	<5	<10	<50	<5	<20	50	<10
US06 D387937	47CST	30	50	<10	<50	<5	<20	30	<10
US06 D387940	47FST	50	50	<10	<50	<5	<20	50	<10
US06 D387941	47FGW	15	<5	<10	<50	<5	<20	15	<10
US06 D387943	47XGW	50	70	<10	<50	<5	<20	15	15
US06 D387944	47GST	70	70	<10	<50	<5	<20	70	15
US06 D387946	47BGW	20	10	<10	<50	<5	<20	30	<10
US06 D387948	47EGW	20	5	<10	<50	<5	<20	30	<10
US06 D387950	47GGW	50	20	<10	<50	<5	<20	50	<10
US06 D387951	47DST	50	50	<10	<50	5	<20	50	10

Lab no.	Field no.	Sb SQS-ppm	Sc SQS-ppm	Sn SQS-ppm	Sr SQS-ppm	Th SQS-ppm	V SQS-ppm	W SQS-ppm	Y SQS-ppm
US06 D387931	47EST	<100	5	<10	<100	<100	150	<20	10
US06 D387932	47BSTDA	<100	10	<10	<100	<100	200	<20	15
US06 D387934	47BST	<100	5	<10	<100	<100	150	<20	10
US06 D387936	47DGW	<100	<5	<10	<100	<100	70	<20	<10
US06 D387937	47CST	<100	7	<10	<100	<100	150	<20	<10
US06 D387940	47FST	<100	10	<10	<100	<100	200	<20	10
US06 D387941	47FGW	<100	<5	<10	<100	<100	50	<20	<10
US06 D387943	47XGW	<100	10	<10	<100	<100	200	<20	<10
US06 D387944	47GST	<100	10	<10	<100	<100	200	<20	15
US06 D387946	47BGW	<100	<5	<10	<100	<100	50	<20	<10
US06 D387948	47EGW	<100	<5	<10	<100	<100	70	<20	<10
US06 D387950	47GGW	<100	<5	<10	<100	<100	100	<20	<10
US06 D387951	47DST	<100	7	<10	<100	<100	150	<20	<10

Table 5. Geochemical data for rock samples collected near Fortyseven Creek.

Lab no.	Field no.	Zn SQS-ppm	Zr SQS-ppm	description
US06 D387931	47EST	<200	50	Kuskokwim Group siltstone
US06 D387932	47BSTDA	<200	100	silicified and oxidized Kuskokwim Group siltstone
US06 D387934	47BST	<200	100	silicified and oxidized Kuskokwim Group siltstone
US06 D387936	47DGW	<200	70	silicified with quartz veins in Kuskokwim Group graywacke
US06 D387937	47CST	<200	70	silicified Kuskokwim Group siltstone
US06 D387940	47FST	<200	70	weakly silicified Kuskokwim Group siltstone
US06 D387941	47FGW	<200	70	silicified with quartz veinlets in Kuskokwim Group graywacke
US06 D387943	47XGW	<200	70	silicified and oxidized Kuskokwim Group graywacke
US06 D387944	47GST	<200	100	Kuskokwim Group siltstone with small quartz vein
US06 D387946	47BGW	<200	70	silidified and oxidized Kuskokwim Group graywacke
US06 D387948	47EGW	<200	50	Kuskokwim Group graywacke
US06 D387950	47GGW	<200	50	weakly silicified Kuskokwim Group graywacke
US06 D387951	47DST	<200	30	silicified Kuskokwim Group siltstone