

Mercator projection
Geodetic Reference System 1980; North American Datum 1983
Longitude of central meridian 70° 19' W; latitude of true scale 41° 39' N
False easting 0 m; false northing 0 m
This map is not intended for navigational purposes.

SCALE 1:25 000
ONE CENTIMETER ON THE MAP REPRESENTS 250 METERS ON THE SEA FLOOR
CONTOUR INTERVAL 5 METERS
DATUM MEAN LOWER LOW WATER

DISCUSSION

Introduction - The Stellwagen Bank National Marine Sanctuary Mapping Project is a cooperative effort of the U.S. Geological Survey and the National Oceanic and Atmospheric Administration, with support from the University of New Brunswick and the Canadian Hydrographic Survey. The multibeam echosounder survey was conducted on four cruises over a two-year period from the fall of 1994 to the fall of 1996. This map shows one of a series of 16 quadrangles (see location map) in which sea floor depth information is depicted in sun-illuminated (or shaded relief) view at a scale of 1:25,000, with topographic contours overlaid in blue. The image shown here uses a sun elevation angle of 45 degrees above the horizon from an azimuth of 350 degrees and a vertical exaggeration of four times. In effect, topographic relief is enhanced by having the sun illuminate the sea floor from a position 10 degrees west of north so that shadows are cast on the southern flanks of seabed features. Some features in the images are artifacts of data collection. They are especially noticeable where the seabed is smooth and include small highs and lows and unnatural-looking features and patterns that are oriented parallel or perpendicular to survey tracklines. For a depiction of the topographic contours alone, and for an explanation of survey and topographic data processing methods, see the companion map by Valentine and others (1997). Topographic contour maps for all 16 quadrangles of the map series are available on a CD-ROM in EPS, PS, Arc export, and PDF file formats (Valentine and others, 1998). Blank areas represent places where no data exist.

Regional seabed features - The major topographic features depicted in the map series were formed by glacial processes. In broad terms, these features are interpreted here to represent a geologic history that developed in several stages. Ice containing rock debris moved across the region, sculpting its surface and depositing sediment to form the large basins, banks, ridges, and valleys. Many other features observed here represent the later stages of deglaciation. They are the result of processes at work when much of the area was covered by stationary retreating ice, and when at the same time small valley glaciers and ice falls were active in and near areas of high topographic relief. The sea invaded the region formerly occupied by ice, and seabed features were partly eroded and some new sedimentary deposits were formed. Today, the sea floor mainly is modified by strong southward-flowing bottom currents caused by storm winds from the northeast. These currents erode sediments from the shallow banks and transport them into the basins. With time, the banks affected by these currents become coarser, as sand and mud are removed but gravel remains; and the western flanks of the banks, and adjacent basins, are built up by deposits of mud and sand.

Quadrangle 15 features - The regional slope of the sea floor in Quadrangle 15 increases from approximately 70 m water depth in the southwest to 130 m in the east and northeast. The shallow and deep parts of the seabed are very different in character. The southwestern part of the quadrangle is characterized by low hills and shallow valleys (and one deep valley) that are related topographically to the well-defined banks and deep valleys that occur in the adjacent Quadrangle 14 to the west (Valentine and others, 1999). The low hills are covered with gravel, including boulder piles and ridges. The gravel is covered in places by a thin veneer of sand that is more extensive on the hilltops. Some of the linear boulder ridges (42 33.9'N, 70 11.4'W; 42 35.2'N, 70 09.2'W; 42 35.4'N, 70 12.0'W) resemble eskers (sand and gravel deposited by running water in channels within stationary glacial ice). Other ridges are located along the upper edges of shallow glaciated valleys (42 33.9'N, 70 11.9'W; 42 35.65'N, 70 13.15'W) and are interpreted to be lateral moraines (linear deposits of rock debris formed at the edges of moving ice). The valleys are floored by sand. Lateral moraines also are present lower on the valley walls (42 36.9'N, 70 12.5'W and 42 34.35'N, 70 11.3'W). Broad depressions in the tops of some hills possibly indicate the former locations of large masses of melting ice (42 34.8'N, 70 09.0'W). The sea floor in the remainder of the quadrangle exhibits a southeast-trending grain of small hills and basins that probably reflect the movement of a regional ice sheet. The hills are low and rounded and typically have 5 to 10 m of relief. They are covered with gravel patches of boulders and cobbles are common that is partly covered with a thin veneer of sand. Between the hills, the seabed is chiefly sand with patches of gravel with scattered cobbles and boulders. A series of elongate, shallow, southeast-trending basins is present in the central and southeastern part of the quadrangle. Some of the basins reach water depths of 120 to more than 150 m, and their floors are 15 to 20 m below the surrounding seabed. The floors of the deepest basins are relatively smooth and are covered with fine-grained sand and mud. The most striking topographic features in Quadrangle 15 are many long, narrow grooves that dominate

all but the southwestern part of the map area. They typically are 50 to 100 m wide, up to 5 to 10 m deep, and up to several kilometers long. These grooves are interpreted to be marks made by the jagged bottoms of icebergs that gouged the seabed by grounding here during the late stages of the last glaciation. The iceberg grooves are most common in the deeper eastern part of the quadrangle, and they disappear in the 75 to 80 m depth interval in the shallow southwestern part of the quadrangle. The grooves appear to have a dominant northeast-southwest orientation. They become less distinct toward the southwest where they appear to be partly filled by sandy sediment transported from shallow areas in the western part of this quadrangle and the eastern part of Quadrangle 14.

REFERENCES CITED

- Valentine, P.C., Baker, J.L., Unger, T.S., and Roworth, E.T., 1997, Sea floor topography of Quadrangle 15 in the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Open-File Report 97-728, scale 1:25,000, 1 sheet.
- Valentine, P.C., Baker, J.L., Unger, T.S., and Polloni, C., 1998, Sea floor topographic map and perspective view imagery of Quadrangles 1-18, Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Open-File Report 98-138, 1 CD-ROM.
- Valentine, P.C., Unger, T.S., and Baker, J.L., 1999, Sun-illuminated sea floor topography of Quadrangle 14 in the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Geologic Investigations Series Map I-2714, scale 1:25,000, 1 sheet.


SUN-ILLUMINATED SEA FLOOR TOPOGRAPHY OF QUADRANGLE 15 IN THE STELLWAGEN BANK NATIONAL MARINE SANCTUARY OFF BOSTON, MASSACHUSETTS

By

Page C. Valentine, Jessica L. Baker, and Tanya S. Unger

1999

AUTHOR AFFILIATIONS
U.S. Geological Survey, Woods Hole Field Center
384 Woods Hole Road, Woods Hole, MA 02543



Location map - Shows mapped quadrangle outlined. Stellwagen Bank National Marine Sanctuary (SBNMS) boundary shown as dashed line. Bathymetric contours in meters.