

CONTENTS

Abstract	1
Introduction	1
Use of Vapor Diffusion Samplers for Monitoring of Volatile Organic Compounds	2
Volatile Organic Compounds Detected in Vapor Diffusion Samples	4
References	8

PLATE (in back pocket)

1. Map showing location of study area, data collection sites, and total concentrations of volatile organic compounds from vapor diffusion samplers, and graphs of trichloroethylene and tetrachloroethylene concentrations at collection sites, Centredale Manor Site, North Providence, Rhode Island, September 1999

FIGURES

1. Graph showing gage height of Woonasquatucket River, U.S. Route 44 bridge, North Providence, Rhode Island, September 1999..... 3

TABLES

1. Minimum reporting limits for target volatile organic compounds, Centredale Manor, North Providence, Rhode Island, September 1999..... 4
2. Concentrations of volatile organic compounds in vapor diffusion samplers at Centredale Manor, North Providence, Rhode Island, September 1999..... 5

CONVERSION FACTORS, VERTICAL DATUM, AND ABBREVIATIONS

CONVERSION FACTORS

Multiply	By	To obtain
acre	0.4047	hectare
inches (in.)	2.54	centimeters
feet (ft)	0.3048	meters
miles (mi)	1.609	kilometers

Temperature in degrees Celsius (°C) can be converted to degrees Fahrenheit (°F) as follows:
 $^{\circ}\text{F} = 1.8^{\circ}\text{C} + 32$

VERTICAL DATUM

Sea Level: In this report, “sea level” refers to the National Geodetic Vertical Datum of 1929 (NGVD of 1929)—a geodetic datum derived from a general adjustment of the first-order level nets of the United States and Canada, formerly called Sea Level Datum of 1929.

ABBREVIATIONS

mL	milliliter
ppb v	parts per billion by volume
µg/L	micrograms per liter
mm	millimeter