

WATER-LEVEL CHANGES

HYDROLOGIC CONDITIONS IN WEST-CENTRAL FLORIDA

The Floridan aquifer system consists of the Upper and Lower Floridan aquifers separated by a middle confining unit. The middle confining unit and the Lower Floridan aquifer in west-central Florida generally contain highly mineralized water. The water-bearing units containing fresh water are herein referred to as the Upper Floridan aquifer. The Upper Floridan aquifer is a highly productive aquifer and supplies more than 10 times the amount of water pumped from either the surficial aquifer system or the intermediate aquifer system in most of the study area (Duerr and others, 1988).

This map report shows the potentiometric surface of the Upper Floridan aquifer measured in September 2003. The potentiometric surface is an imaginary surface connecting points of equal altitude to which water will rise in tightly-cased wells that tap a confined aquifer system. (Lohman, 1979). This map represents water-level conditions near the end of the wet season, when ground-water levels usually are at an annual high and withdrawals for agricultural use typically are low. The cumulative average rainfall for west-central Florida (from October 2002 through September 2003) was 13.15 inches above the historical cumulative average of 53.15 inches (Southwest Florida Water Management District, September 2003). Historical cumulative averages are calculated from regional rainfall summary reports (1915 to most recent complete calendar year) and are updated monthly by the Southwest Florida Water Management District.

This report, prepared by the U.S. Geological Survey in cooperation with the Southwest Florida Water Management District, is part of a semi-annual series of Upper Floridan aquifer potentiometric-surface map reports for west-central Florida. Potentiometric-surface maps have been prepared for January 1964, May 1969, May 1971, May 1973, May 1974, and for each May and September since 1975. Water-level data are collected in May and September each year to show the approximate annual low and high water-level conditions, respectively. Most of the water-level data for this map were collected by the U.S. Geological Survey during the period September 16-20, 2003. Supplemental water-level data were collected by other agencies and companies. A corresponding potentiometric-surface map was prepared for areas east and north of the Southwest Florida Water Management District boundary by the U.S. Geological Survey office in Altamonte Springs, Florida (Kinman, S.L., 2004). Most water-level measurements were made during a 5-day period, therefore, measurements do not represent a "snapshot" of conditions at a specific time, nor do they necessarily coincide with the seasonal high water-level condition.

Water levels in about 80 percent of the wells measured in September 2003 were higher than the September 2002 water levels (Blanchard and others, 2003). Data from 424 wells indicate the September 2003 levels ranged from about 4 feet below to about 20 feet above the September 2002 water levels (fig. 1). The largest water-level declines occurred in northwestern Highlands County, southern Sumter and southwestern Pasco County. The largest water-level rise was in north-central Pasco and southern Hernando Counties (fig. 1).

Water levels in about 99 percent of the wells measured in September 2003 were higher than in May 2003 (Blanchard and others, 2004). Data from 421 wells indicate the September 2003 water levels ranged from about 0 to 27 feet above the May 2003 water levels. The largest water-level decline was in west-central Pinellas County and the largest increase in water levels was in south-central Hillsborough County.

ACKNOWLEDGMENTS

The authors thank the Southwest Florida Water Management District, Tampa Bay Water and CFI Industries for their assistance with the preparation and production of this report.

SELECTED REFERENCES

Duerr, A.D., Hunn, J.D., Lewelling, B.R., and Trommer, J.T., 1988, Geology and 1985 water withdrawals of the aquifer systems in southwest Florida, with emphasis on the intermediate aquifer system: U.S. Geological Survey Water-Resources Investigations Report 1987-4259, 115 p.

Blanchard, R.A., Knochenmus, L.A., Seidenfeld, A.V., and McCulloch, D.S., 2003, Potentiometric surface of the Upper Floridan aquifer, west-central Florida, September 2002: U.S. Geological Survey Open-File Report 2003-223, 1 sheet.

Blanchard, R.A., Seidenfeld, A.V., and McCulloch, D.S., 2004, Potentiometric surface of the Upper Floridan aquifer, west-central Florida, May 2003: U.S. Geological Survey Open-File Report 2004-1079, 1 sheet.

Kinman, S.L., and Knowles Jr., L., 2004, Potentiometric surface of the Upper Floridan aquifer in the St. Johns River Water Management District and vicinity, Florida, September 2003: U.S. Geological Survey Open-File Report 2004-1288, 1 sheet.

Lohman, S.W., 1979, Ground-water hydraulics: U.S. Geological Survey Professional Paper 708, 79 p.

Southwest Florida Water Management District, 2002, Hydrologic conditions for the month of September 2003, 79 p.

EXPLANATION

Areas where potentiometric surface changed, in feet

- 10 and greater
- 5 to 9.99
- 0 to 4.99
- Less than 0 to -5


(Positive values indicate an increase in water-level altitudes)

Figure 1. Change in potentiometric surface of the Upper Floridan aquifer from September 2002 to September 2003 in west-central Florida.

EXPLANATION

- MUNICIPAL WELL FIELD
- POTENTIOMETRIC CONTOUR - - Shows altitude at which water would have stood in tightly cased wells. Contour interval is 10 feet. National Geodetic Vertical Datum (NGVD) of 1929. Hechures indicate depressions. Dashed where approximately located.
- BOUNDARY OF SOUTHWEST FLORIDA WATER MANAGEMENT DISTRICT
- OBSERVATION WELL - - Number is altitude of water level in feet above or below NGVD of 1929.
- OBSERVATION WELL - - Number is altitude of water level in feet above or below NGVD of 1929. Italic number indicates water levels corrected for salinity.
- SPRING
- CITY OR TOWN

NOTE: The potentiometric contours are generalized to synoptically portray the head in a dynamic hydrologic system, taking due account of the variations in hydrogeologic conditions, such as differing depths of wells, nonsimultaneous measurements of water levels, variable effects of pumping, and changing climatic influence. The potentiometric contours may not conform exactly with the individual measurements of water level.



POTENTIOMETRIC SURFACE OF THE UPPER FLORIDAN AQUIFER, WEST-CENTRAL FLORIDA, SEPTEMBER 2003
By R.A. Blanchard, A.V. Seidenfeld, and D.S. McCulloch