

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryna glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina biora</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormosinella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
0-2	12	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	2	-	26	-	
4-6	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	25	-	
8-10	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	10	-	
12-14	14	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	6	-	8	-	
16-18	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	26	-	
20-22	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	44	-	
24-26	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	8	-	26	-	
28-30	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	16	-	
32-34	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	6	-	
36-38	2	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	-	10	-	
40-42	1	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	2	-	10	-	
44-46	3	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	1	-	2	-	
48-50	2	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	2	-	5	-	
52-54	3	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	2	-	15	-	
56-58	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	39	-	
60-62	3	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	3	-	16	-	
64-66	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	9	-	
68-70	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	26	-	
76-78	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	20	-	
80-80	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	9	-	
84-86	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	34	-	
88-90	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	18	-	
92-94	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	-	
96-98	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	32	-	
100-02	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	27	-	
104-06	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	41	-	
112-14	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	30	-	
116-18	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	25	-	
120-22	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	33	-	
128-30	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	16	-	
132-34	2	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	2	-	34	-	
136-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	29	-	
140-42	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	-	
144-46	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	46	-	
148-50	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	28	-	
152-54	7	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	17	-	
156-58	5	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	2	-	42	-	
160-62	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	17	-	
164-66	1	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	7	-	17	-	
168-70	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	32	-	
172-74	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	62	-	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina bitor</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormonsinella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
176-78	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	22	-	
180-82	7	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	-	50	-	
184-86	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	66	-	
188-90	11	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	54	-	
192-94	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	40	-	
196-98	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	32	-	
200-02	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	38	-	
204-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	35	-	
208-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	22	-	
212-14	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	84	-	
216-18	2	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	2	-	56	-	
220-22	3	-	-	-	-	-	-	-	-	-	-	-	-	1	1	-	11	34	-	1	-	15	1	
224-26	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	20	14	-	-	-	17	-	
228-30	2	-	-	-	-	-	-	-	-	-	-	-	-	2	1	-	-	-	-	1	-	41	-	
232-34	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	26	-	
236-38	7	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	25	-	
240-42	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	46	-	
244-46	1	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	1	-	18	-	
248-50	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	55	-	
252-54	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	10	10	-	-	-	21	-	
256-58	2	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	1	2	-	1	-	16	-	
260-62	4	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	2	1	-	-	-	46	-	
264-66	14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	1	-	27	-	
268-70	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	12	15	-	-	-	27	-	
272-74	8	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	-	-	37	-	
276-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	80	-	
280-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	21	8	-	-	-	50	-	
284-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	20	-	
288-90	3	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	4	-	-	-	-	26	-	
292-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	72	-	
296-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	53	-	
300-02	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	-	55	-	
304-06	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	59	-	
308-10	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	52	-	
312-14	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	61	-	
316-18	2	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	63	-	
320-22	3	-	-	-	-	-	-	-	-	-	-	-	-	1	1	-	-	-	-	-	-	70	-	
324-26	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	50	-	
328-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	47	-	
332-34	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	-	-	28	-	
336-38	4	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	2	-	30	-	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina biora</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormonsinella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
340-42	2																			1		32		
348-50														3								71		
352-54	3													3						6		10		
356-58							1										21	13		1		42		
360-62														2			27	10						
364-66	1																38	13		1		16		
368-70																								
372-74																	1				2		85	
376-78	2																	1					56	
380-82																	6	9					26	
384-86	1																21	13					6	
388-90																							12	
392-94							1										2	2					66	
396-98	2																						63	
400-02	2						1										8	8			2		11	
404-06	3													1							2		64	
408-10	1													1							2		36	
412-14	1																						61	
416-18																					1		67	
420-22																							49	
424-26																							17	
428-30	1																						42	
432-34	1													2										
436-38	1													2									14	
440-42																							62	
444-46	1													1									62	
448-50																							75	
452-54																							50	
456-58																							59	
460-62	1																						74	
464-66																					5		82	
468-70	2																						74	
472-74	14													4							8		20	
476-78																					1		79	
480-82	3																				1		77	
484-86														1									52	
488-90																	1						91	
492-94														2									81	
496-98	1													2									78	
500-02																					1		70	
504-06															1								76	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)																							
508-10	1	<i>Adercotryma glomeratum</i>																					
512-14	2	<i>Alabaminella wedellensis</i>																					
516-18		<i>Angulogerina earlandi</i>																					
520-22		<i>Angulogerina pauperata</i>																					
524-26		<i>Astronion echolsi</i>																					
528-30		<i>Astronion antarctica</i>																					
532-34		<i>Bolivina psuedopunctata</i>																					
536-38		<i>Cassidulina neocarinata</i>																					
540-42	1	<i>Cassidulinoides prava</i>																					
544-46	1	<i>Cibicides lobatulus</i>																					
548-50		<i>Cibicides refulgans</i>																					
552-54	1	<i>Cibicides spp.</i>																					
556-58		<i>Conotrocammina alternans</i>																					
560-62		<i>Cribrostomoides arenacea</i>												2									
564-66		<i>Cribrostomoides sphaeriloculus</i>																					
568-70		<i>Epistominella exigua</i>																					
572-74		<i>Globocassidulina bitor</i>																					
576-78		<i>Globocassidulina subglobosa</i>																					
580-82	1	<i>Haplophragmoides wiesneri</i>																					
584-86		<i>Hormoninella ovicula</i>																					
588-90		<i>Labrospira jeffersi</i>																					
592-94	1	<i>Melonis affinis</i>																					
596-98		<i>Miliammina arenacea</i>																					
600-02		<i>*Neogloboquadrina pachyderma</i>																					
604-06																							
608-10																							
612-14																							
616-18																							
620-22	3																						
624-26	2																						
628-30																							
632-34																							
636-38																							
640-42																							
644-46																							
648-50																							
652-54																							
656-58																							
660-62																							
664-66																							
668-70																							

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina biora</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
672-74																								
676-78	1														1						1		68	
680-82																							77	
684-86																							87	
688-90																							78	
688-90														1							1		73	
692-94																							94	
700-02																							69	
704-06																							67	
708-10																	1						48	
712-14																							5	
716-18																							11	
720-22																							69	
724-26														1									59	
728-30														1									89	
732-34	1																						73	
736-38																	11						79	
740-42																							79	
744-46																							78	
748-50																							74	
752-54																	4						64	
756-58																							84	
760-62																					1		67	
764-66																							65	
768-70																							51	
772-74																	1						87	
776-78																							69	
780-82																							81	
784-86																							72	
788-90																							94	
792-94																							66	
796-98																							91	
800-02																							51	
804-06														1									68	
808-10			1														8						43	
812-14																1	3	3					22	
816-18																							94	
820-22																	5						72	
824-26																	25	9					31	
828-30																							68	
832-34																							89	
836-38																							95	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotyma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina pseudopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribr stomoides arenacea</i>	<i>Cribr stomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina biora</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
840-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	67	-	
844-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
848-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	86	-	
852-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
856-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
860-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	-	
864-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	85	-	
868-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	65	-	
872-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	89	-	
876-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	78	-	
880-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	-	
884-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
888-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	75	-	
892-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	29	-	
896-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
900-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
904-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	98	-	
908-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
912-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
916-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	-	
920-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
924-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	67	-	
928-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
932-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	93	-	
936-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
940-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	-	
944-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
948-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	-	
952-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	93	-	
956-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	-	
960-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	-	
964-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
968-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	82	-	
972-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	88	-	
976-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	102	-	
980-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	-	
984-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
988-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
992-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	-	
996-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	1	-	-	-	67	-	
1000-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	76	-	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina bitor</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
1004-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	55	-	
1008-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	93	-	
1012-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6-	12	-	-	-	24	-	
1016-18	-	-	-	-	-	-	-	-	-	-	1	1	-	-	-	-	39	4	-	-	-	51	-	
1020-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	59	-	
1024-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	84	-	
1028-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	85	-	
1032-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	-	
1036-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	83	-	
1040-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	58	-	
1044-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1-	7	-	-	-	76	-	
1048-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	-	
1052-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	96	-	
1056-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
1060-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	92	-	
1064-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	98	-	
1068-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	96	-	
1072-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	95	-	
1076-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	91	-	
1080-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1084-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	-	
1088-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1092-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	-	
1096-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
1100-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	-	
1104-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	92	-	
1108-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	89	-	
1112-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
1116-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	18	6	-	-	-	73	-	
1120-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
1124-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	-	
1128-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	90	-	
1132-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	83	-	
1136-38	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	90	-	
1140-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	23	8	-	-	-	61	-	
1144-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	-	
1148-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	81	-	
1152-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	3	-	-	-	76	-	
1156-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	3	-	-	-	76	-	
1160-62	-	-	-	-	-	-	-	-	-	1	1	-	-	-	-	-	4	-	-	-	-	61	-	
1164-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	-	-	30	-	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina pseudopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina bitor</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
1168-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	1	-	-	-	49	-	
1172-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	21	-	-	-	71	-	
1176-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	89	-	
1180-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1184-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1188-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
1192-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	-	
1196-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	-	
1200-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
1204-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1208-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	93	-	
1212-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
1216-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
1220-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	-	
1224-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	85	-	
1228-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
1232-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	95	-	
1236-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	101	-	
1240-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	-	
1244-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	-	
1248-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1252-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	-	
1256-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	40	-	
1260-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	-	
1264-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	93	-	
1268-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	-	
1272-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	-	
1276-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	-	
1280-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	54	-	
1284-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1288-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	88	-	
1292-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	-	
1296-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	-	
1300-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	98	-	
1304-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	88	-	
1308-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1312-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	-	
1316-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	-	
1320-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	-	
1324-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	-	
1328-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	-	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina bitoria</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
1332-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	
1336-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1340-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	
1344-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	
1348-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	90	
1352-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	
1356-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	
1360-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	79	
1364-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	
1368-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	61	
1372-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	78	
1376-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	95	
1380-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	79	
1384-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	56	
1388-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	91	
1392-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	61	
1396-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	71	
1400-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	93	
1404-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	
1408-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	
1412-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	79	
1416-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	80	
1420-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	63	
1424-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	71	
1428-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	
1432-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	25	2	-	-	-	-	66	
1436-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	35	4	-	-	-	-	48	
1440-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3-	3	-	-	-	-	52	
1444-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	
1448-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	
1452-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	81	
1456-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	86	
1460-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	95	
1464-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	82	
1468-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	
1472-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	
1476-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	
1480-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	
1484-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	85	
1488-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	
1492-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	88	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotyma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina biora</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
1496-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	38	
1500-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	71	
1504-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	
1508-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	
1512-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	
1516-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	22	
1520-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	
1524-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1528-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	
1532-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	42	
1536-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1540-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	
1544-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	76	
1548-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	92	
1552-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	
1556-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1560-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	55	
1564-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	
1568-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	
1572-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1576-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	90	
1580-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	99	
1584-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1588-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	95	
1592-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	53	
1596-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	94	
1600-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	
1604-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	-	-	-	-	62	
1608-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	12	-	-	-	-	-	25	
1612-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11	-	-	-	-	-	29	
1616-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	6	-	-	-	-	15	
1620-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	2	-	-	-	-	13	
1624-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	4	-	-	-	-	8	
1628-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	-	-	-	-	10	
1632-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	19	4	-	-	-	-	11	
1636-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	96	
1640-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	12	12	-	-	-	-	4	
1644-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	58	
1648-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	8	15	-	-	-	-	5	
1652-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1-	8	-	-	-	-	45	
1656-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	13	11	-	-	-	-	22	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina psuedopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina biora</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormoninella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
1660-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	9	7	-	-	-	7	1	
1664-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	12	-	-	-	6	1	
1668-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	9	9	-	-	-	11	-	
1672-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	11	16	-	-	-	16	-	
1676-78	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	4	9	13	-	-	-	24	-	
1680-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	4	-	-	-	23	-	
1684-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	4	-	-	-	22	-	
1688-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	2	6	-	-	-	12	-	
1692-94	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1	2	11	-	-	-	5	-	
1696-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	5	5	-	-	-	12	-	
1700-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	3	13	-	-	-	11	-	
1704-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	77	1	
1708-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	7	-	-	-	34	-	
1712-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	23	6	-	-	-	32	-	
1716-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	4	14	-	-	-	3	1	
1720-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	-	-	11	2	
1724-26	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	2	6	2-	-	-	-	7	-	
1728-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	14	-	-	-	25	-	
1732-34	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	19	-	-	-	15	-	
1736-38	-	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	12	-	-	-	42	3	
1740-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	20	-	-	-	33	-	
1744-46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7	10	-	-	-	22	-	
1748-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	12	-	-	-	27	-	
1752-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	3	-	-	-	51	-	
1756-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1	2	-	-	-	50	-	
1760-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	5	-	-	-	46	-	
1764-766	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	5	-	-	-	54	-	
1768-770	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-	-	-	49	-	
1772-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	4	-	-	-	32	-	
1776-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	9	-	-	-	52	-	
1780-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	3	-	-	-	58	-	
1784-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	70	-	
1788-90	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	4	3	-	-	-	52	1	
1792-94	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	65	-	
1796-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	84	-	
1800-02	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	80	1	
1804-06	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	-	66	1	
1808-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	79	1	
1812-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	87	1	
1816-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	84	-	
1820-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	81	-	

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	<i>Adercotryma glomeratum</i>	<i>Alabaminella wedellensis</i>	<i>Angulogerina earlandi</i>	<i>Angulogerina pauperata</i>	<i>Astrononion echolsi</i>	<i>Astrononion antarctica</i>	<i>Bolivina pseudopunctata</i>	<i>Cassidulina neocarinata</i>	<i>Cassidulinoides prava</i>	<i>Cibicides lobatulus</i>	<i>Cibicides refulgans</i>	<i>Cibicides spp.</i>	<i>Conotrocammina alternans</i>	<i>Cribrostomoides arenacea</i>	<i>Cribrostomoides sphaeriloculus</i>	<i>Epistominella exigua</i>	<i>Globocassidulina bioira</i>	<i>Globocassidulina subglobosa</i>	<i>Haplophragmoides wiesneri</i>	<i>Hormosinella ovicula</i>	<i>Labrospira jeffersi</i>	<i>Melonis affinis</i>	<i>Miliammina arenacea</i>	<i>*Neogloboquadrina pachyderma</i>
1824-26	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1	86	-
1828-30	-	-	-	-	1	-	-	1	-	-	-	-	-	-	-	-	-	2	-	-	-	-	70	-
1832-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	72	-
1836-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	7	-	-	-	-	41	-
1840-42	-	-	-	-	3	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	44	-
1844-46	-	-	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	51	2
1848-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	-	-	-	53	4
1852-54	-	-	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	65	1
1856-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	45	-
1860-62	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	80	1
1864-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	91	-
1868-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	86	-
1872-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	84	-
1876-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1880-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	47	-
1884-86	-	-	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	85	-
1888-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	2	-	-	-	-	39	2
1892-94	-	2	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	-	-	39	-
1896-98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	53	2
1900-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	43	-	
1904-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11	-	-	-	-	-	26	-
1908-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	6	-	-	-	-	-	33	-
1912-14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	-	-	-	-	1	70	-
1916-18	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	1	-	-	-	-	-	-	85	-
1920-22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	-	-	-	71	-
1924-26	-	-	-	-	2	1	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	58	-
1928-30	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1	3	-	-	-	-	-	80	-
1932-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	4	-	-	-	-	-	70	-
1936-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	73	-
1940-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	-	-	-	-	-	54	-
1944-46	-	-	-	-	1	1	-	1	-	-	-	-	-	-	-	2	4	-	-	-	-	-	24	-
1948-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	2	-	-	-	-	-	13	-
1952-54	-	-	18	2	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	56	-
1956-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	74	-
1960-62	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	91	-
1964-66	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	4	-	-	-	-	-	5	-
1968-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	44	1
1972-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11	-	-	-	-	-	8	-
1976-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	-	-	-	-	-	25	-
1980-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	15	-
1984-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	20	-

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuilina minuta</i>	Total	Percent Agglutinated	Percent Calcareous
0-2	-	-	8	2	-	-	1	-	1	-	-	52	-	-	-	6	111	76.58	0.00
4-6	-	-	10	-	-	-	-	-	2	-	-	56	-	-	-	4	101	75.25	0.00
8-10	-	-	6	2	-	-	1	-	1	-	-	62	-	-	4	2	99	89.90	0.00
12-14	-	-	5	1	-	-	-	-	2	-	-	57	-	-	3	2	99	91.92	0.00
16-18	-	-	3	-	-	-	-	-	1	-	-	60	-	-	1	2	99	73.74	0.00
20-22	-	-	11	1	-	-	-	-	-	-	-	35	-	-	2	5	100	56.00	0.00
24-26	-	-	3	1	-	-	-	-	-	-	-	57	-	-	-	4	100	74.00	0.00
28-30	-	-	10	-	-	-	-	-	1	-	-	66	-	1	-	-	103	84.47	0.00
32-34	-	-	10	1	-	-	-	-	1	-	-	71	-	-	1	2	102	94.12	0.00
36-38	-	-	9	-	-	-	-	-	4	-	1	67	-	-	2	3	100	89.00	1.00
40-42	-	-	11	-	-	1	-	-	3	-	-	57	-	-	6	9	101	90.10	0.00
44-46	-	-	14	-	-	-	-	-	3	-	-	67	-	-	3	8	102	98.04	0.00
48-50	-	-	20	-	-	-	-	-	3	-	-	52	-	-	3	10	99	94.95	0.00
52-54	-	-	17	1	-	-	-	-	7	-	-	50	-	-	2	-	100	85.00	0.00
56-58	-	-	10	-	-	-	-	-	1	-	-	41	-	-	2	4	99	60.61	0.00
60-62	-	-	21	1	-	-	-	-	3	-	-	46	-	-	1	5	102	84.31	0.00
64-66	-	-	5	-	-	1	-	-	2	-	-	64	-	-	1	6	98	90.82	0.00
68-70	-	-	5	1	-	-	-	-	2	-	-	60	-	-	-	-	104	75.00	0.00
76-78	-	-	5	-	-	1	-	-	-	-	-	67	-	-	-	2	100	80.00	0.00
80-80	-	-	17	-	-	-	1	-	1	-	-	62	-	-	-	3	98	90.82	0.00
84-86	-	-	5	-	-	-	-	-	1	-	-	53	-	-	2	2	103	66.99	0.00
88-90	-	-	8	-	-	-	-	-	2	-	-	64	-	-	2	2	98	81.63	0.00
92-94	-	-	9	1	-	-	-	-	-	-	-	55	-	-	2	8	104	74.04	0.00
96-98	-	-	12	-	-	-	-	-	2	-	-	52	-	-	1	4	104	69.23	0.00
100-02	-	-	2	-	-	-	-	-	2	-	-	60	-	-	4	3	100	73.00	0.00
104-06	-	-	4	1	-	-	-	-	-	-	-	50	-	-	1	3	104	60.58	0.00
112-14	-	-	6	-	-	-	-	-	2	-	-	59	-	-	1	7	106	71.70	0.00
116-18	-	-	1	-	-	-	-	-	1	-	-	64	-	-	2	3	102	75.49	0.00
120-22	-	-	3	-	-	-	-	-	-	-	-	58	-	-	-	3	100	67.00	0.00
128-30	-	-	6	-	-	-	-	-	-	-	-	69	-	-	5	1	102	84.31	0.00
132-34	-	-	3	-	-	-	-	-	1	-	-	61	-	-	-	-	104	67.31	0.00
136-38	-	-	3	1	-	-	-	-	-	-	-	68	-	-	-	2	104	72.12	0.00
140-42	-	-	1	-	-	-	-	-	3	-	-	69	-	-	2	-	101	75.25	0.00
144-46	-	-	7	1	-	-	-	-	1	-	-	37	-	-	2	3	103	55.34	0.00
148-50	-	-	5	-	-	-	-	-	-	-	-	58	-	-	4	4	103	72.82	0.00
152-54	-	-	11	-	-	-	-	-	-	-	-	63	-	-	1	3	103	83.50	0.00
156-58	-	-	14	-	-	-	-	-	2	-	-	23	-	-	4	3	98	57.14	0.00
160-62	-	-	16	3	-	-	-	-	4	-	-	45	-	-	-	4	100	83.00	0.00
164-66	-	-	13	-	-	-	-	-	-	-	-	54	-	-	2	4	99	82.83	0.00
168-70	-	-	8	-	-	-	-	-	1	-	-	55	-	-	-	3	105	69.52	0.00
172-74	-	-	9	1	-	-	-	-	-	-	-	21	-	-	3	1	100	38.00	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuilina minuta</i>	Total	Percent Agglutinated	Percent Calcareous
176-78	-	-	31	1	-	-	-	-	2	-	-	37	-	-	2	-	103	78.64	0.00
180-82	-	-	10	1	-	-	-	-	1	-	-	29	-	-	3	-	103	51.46	0.00
184-86	-	-	7	-	-	-	-	-	-	-	-	16	-	-	-	3	102	35.29	0.00
188-90	-	-	8	-	-	-	1	-	-	-	-	17	-	-	3	5	101	46.53	0.00
192-94	-	-	3	-	-	-	1	-	3	-	-	44	-	-	1	6	102	60.78	0.00
196-98	-	-	9	3	-	-	-	-	2	-	-	46	-	-	1	5	104	69.23	0.00
200-02	-	-	6	-	-	-	-	-	1	-	-	51	-	-	1	2	100	62.00	0.00
204-06	-	-	2	-	-	-	-	-	-	-	-	65	-	-	3	1	107	67.29	0.00
208-10	-	-	5	-	-	-	-	-	3	-	6	65	-	-	1	1	104	73.08	5.77
212-14	-	-	-	-	-	-	-	-	1	-	-	12	-	-	-	2	104	19.23	0.00
216-18	-	-	10	2	-	-	1	-	1	-	-	16	-	-	1	3	97	42.27	0.00
220-22	-	-	7	2	-	-	1	-	2	-	1	14	-	-	6	1	100	39.00	46.00
224-26	-	-	1	-	-	-	-	-	3	-	3	5	-	-	7	1	72	25.00	51.39
228-30	-	-	6	1	-	-	-	-	-	-	1	35	-	-	4	-	94	55.32	1.06
232-34	-	-	7	1	-	-	-	-	8	-	-	16	-	-	8	-	70	62.86	0.00
236-38	-	-	2	2	-	-	-	-	3	-	-	47	-	-	4	8	99	74.75	0.00
240-42	-	-	4	-	-	-	-	-	2	-	-	35	-	-	3	9	101	54.46	0.00
244-46	-	-	5	-	-	-	-	-	-	-	-	67	-	-	4	3	101	82.18	0.00
248-50	-	-	2	1	-	-	-	-	5	-	-	36	-	-	-	3	103	46.60	0.00
252-54	-	15	1	1	-	-	-	-	-	25	15	-	-	-	-	2	103	21.36	58.25
256-58	-	1	3	1	-	-	-	-	1	-	11	51	-	-	3	3	99	68.69	15.15
260-62	-	-	5	1	-	-	8	-	3	-	-	20	1	-	5	-	98	50.00	3.06
264-66	-	1	13	5	-	-	-	-	-	18	5	-	-	11	1	1	98	51.02	21.43
268-70	-	2	10	1	-	-	-	-	2	-	9	5	-	-	7	2	94	30.85	40.43
272-74	-	-	4	1	-	-	-	-	-	-	-	31	-	-	7	8	99	62.63	0.00
276-78	-	-	-	-	-	-	-	-	-	-	-	14	-	-	3	3	100	20.00	0.00
280-82	-	8	1	-	-	-	-	-	-	2	7	-	-	-	-	-	97	8.25	40.21
284-86	-	-	5	-	-	-	-	-	2	-	-	10	-	-	6	11	54	62.96	0.00
288-90	-	-	8	1	-	-	-	-	2	-	-	34	1	-	5	7	92	67.39	4.35
292-94	-	-	3	-	-	-	-	-	-	-	-	19	-	-	-	2	96	25.00	0.00
296-98	-	-	6	1	-	-	-	-	1	-	-	3-	-	-	2	7	100	47.00	0.00
300-02	-	-	1	1	-	-	-	-	-	-	-	37	-	-	-	4	100	45.00	0.00
304-06	-	-	3	-	-	-	-	-	2	-	-	32	-	-	-	2	100	41.00	0.00
308-10	-	-	11	-	-	-	-	-	-	-	-	31	-	-	2	3	100	48.00	0.00
312-14	-	-	19	-	-	-	-	-	-	-	-	1-	-	-	3	3	103	40.78	0.00
316-18	-	-	10	2	-	-	-	-	-	-	-	22	-	-	-	2	102	38.24	0.00
320-22	-	-	8	1	-	-	-	-	-	-	-	15	-	-	1	4	104	32.69	0.00
324-26	-	-	5	1	-	-	-	-	-	-	-	36	-	-	-	5	103	51.46	0.00
328-30	-	-	7	1	-	-	-	-	2	-	-	42	-	-	-	3	103	54.37	0.00
332-34	-	-	13	-	-	-	-	-	6	-	-	51	-	-	1	1	102	72.55	0.00
336-38	-	-	14	1	-	-	-	-	4	-	-	31	-	-	-	4	93	67.74	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuilina minuta</i>	Total	Percent Agglutinated	Percent Calcareous
340-42	-	-	11	-	-	-	-	-	7	-	-	41	-	-	-	3	97	67.01	0.00
348-50	-	-	5	-	-	-	-	-	1	-	-	21	-	-	1	1	103	31.07	0.00
352-54	-	-	19	-	-	-	-	-	4	-	-	49	-	-	-	4	98	89.80	0.00
356-58	-	3	2	1	-	-	-	-	-	-	12	8	-	-	-	-	104	11.54	48.08
360-62	-	-	8	-	-	1	-	-	-	-	-	13	-	-	3	2	66	43.94	56.06
364-66	1	5	5	-	-	-	-	-	-	-	11	10	-	-	-	-	101	16.83	67.33
368-70	-	-	10	1	-	-	-	-	-	-	-	27	-	-	7	3	48	100.00	0.00
372-74	-	-	6	-	-	-	-	-	3	-	-	10	-	-	-	-	107	19.63	0.93
376-78	-	-	4	-	-	-	1	-	-	-	4	29	-	-	2	1	100	39.00	5.00
380-82	-	-	10	-	-	1	-	-	1	-	4	34	-	-	-	2	93	51.61	20.43
384-86	1	8	1	-	-	-	-	-	1	-	41	5	-	-	2	3	103	12.62	81.55
388-90	-	-	-	-	-	-	-	-	-	-	-	1	-	-	2	-	15	20.00	0.00
392-94	-	-	7	-	-	-	1	-	1	-	-	10	-	-	4	5	99	28.28	5.05
396-98	-	-	6	-	-	-	-	-	1	-	-	24	-	-	5	3	104	39.42	0.00
400-02	-	10	14	1	-	-	-	-	1	-	20	17	-	-	1	6	102	43.14	46.08
404-06	-	-	2	-	-	-	-	-	-	-	-	27	-	-	1	-	100	36.00	0.00
408-10	-	-	6	-	-	-	-	-	-	-	-	41	-	-	6	6	99	63.64	0.00
412-14	-	-	10	-	-	-	-	-	1	-	-	28	-	-	1	1	103	40.78	0.00
416-18	-	-	5	-	-	-	-	-	2	-	-	15	-	-	5	5	100	33.00	0.00
420-22	-	-	11	-	-	-	-	-	4	-	-	24	-	-	2	8	98	50.00	0.00
424-26	-	-	5	1	-	-	-	-	1	-	-	9	-	-	6	6	45	62.22	0.00
428-30	-	-	3	-	-	-	1	-	4	-	-	37	-	-	3	8	99	57.58	0.00
432-34	-	-	2	-	-	-	-	-	-	-	-	11	-	-	6	3	25	100.00	0.00
436-38	-	-	4	-	-	-	-	-	-	-	-	27	-	-	5	3	56	75.00	0.00
440-42	-	-	3	-	-	-	-	-	1	-	-	27	-	-	2	3	98	36.73	0.00
444-46	-	-	2	-	-	-	-	-	-	-	-	24	-	-	-	8	98	36.73	0.00
448-50	-	-	5	-	-	-	-	-	-	-	-	17	-	-	-	3	100	25.00	0.00
452-54	-	-	-	-	-	-	-	-	-	-	1	3	-	-	2	-	56	8.93	1.79
456-58	-	-	4	1	-	-	-	-	2	-	-	23	-	-	3	6	98	39.80	0.00
460-62	-	-	3	-	-	-	-	-	-	-	-	17	-	-	-	2	97	23.71	0.00
464-66	-	-	8	-	-	1	-	-	1	-	-	2	-	-	1	-	100	18.00	0.00
468-70	-	-	3	3	-	-	-	-	1	-	-	14	-	-	2	-	99	25.25	0.00
472-74	-	-	21	-	-	-	-	-	1	-	-	3	-	-	2	2	102	80.39	0.00
476-78	-	-	2	1	-	-	-	-	1	-	-	17	-	-	-	1	102	22.55	0.00
480-82	-	-	4	-	-	-	-	-	-	-	-	12	-	-	-	1	98	21.43	0.00
484-86	-	-	9	-	-	-	2	-	1	-	-	25	-	-	3	2	95	45.26	0.00
488-90	-	-	4	-	-	-	-	-	-	-	-	4	-	-	2	-	102	9.80	0.98
492-94	-	-	5	-	-	-	-	-	2	-	-	8	-	-	3	1	102	20.59	0.00
496-98	-	-	3	1	-	-	-	-	-	-	-	8	-	-	-	2	95	17.89	0.00
500-02	-	-	5	1	-	-	-	-	2	-	-	18	-	-	2	2	101	30.69	0.00
504-06	-	-	5	-	-	-	-	-	1	-	-	17	-	-	-	2	102	25.49	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuilina minuta</i>	Total	Percent Agglutinated	Percent Calcareous
508-10	-	-	24	1	-	-	1	-	4	-	-	22	-	-	3	2	96	67.71	0.00
512-514	-	-	17	-	-	-	-	-	3	-	-	31	-	-	2	8	99	66.67	1.01
516-18	-	-	13	-	-	-	-	-	-	-	-	2	-	-	-	5	99	20.20	0.00
520-22	-	-	15	-	-	-	-	-	8	-	-	15	-	-	3	6	99	53.54	0.00
524-26	-	-	2	-	-	-	-	-	1	-	-	3	-	-	-	-	101	5.94	0.00
528-30	-	-	7	-	-	-	-	-	1	-	-	4	-	-	-	4	100	17.00	0.00
532-34	-	-	6	-	-	-	-	-	1	-	-	25	-	-	2	1	106	35.85	0.00
536-38	-	-	6	-	-	-	-	-	3	-	-	44	-	-	-	3	97	58.76	0.00
540-42	-	-	22	-	-	-	-	-	1	-	-	4	-	-	2	1	102	30.39	0.00
544-46	-	-	6	-	-	-	-	-	-	-	-	2	-	-	-	4	99	13.13	0.00
548-50	-	-	3	-	-	-	-	-	1	-	-	8	-	-	3	1	99	16.16	0.00
552-54	-	-	2	-	-	-	-	-	2	-	-	15	-	-	1	4	102	28.43	0.98
556-58	-	-	20	-	-	-	-	-	-	-	-	16	-	-	1	-	101	38.61	0.00
560-62	-	-	6	1	-	-	-	-	-	-	-	18	-	-	7	4	97	38.14	0.00
564-66	-	-	5	-	-	-	-	-	1	-	-	6	-	-	8	2	74	37.84	0.00
568-70	-	-	1	-	-	-	-	-	2	-	-	4	-	-	1	-	100	8.00	0.00
572-74	-	-	2	-	-	-	-	-	2	-	-	18	-	-	-	1	100	24.00	0.00
576-78	-	-	2	-	-	-	-	-	-	-	-	35	-	-	-	-	104	35.58	0.00
580-82	-	-	5	-	-	-	-	-	3	-	-	49	-	-	2	8	99	68.69	0.00
584-86	-	-	3	-	-	-	-	-	2	-	-	15	-	-	1	3	77	31.17	0.00
588-90	-	-	15	-	-	-	-	-	2	-	-	12	-	-	1	1	103	33.98	0.00
592-94	-	-	1	1	-	-	-	-	-	-	-	2	-	-	-	3	101	7.92	0.00
596-98	-	-	9	-	-	-	1	-	1	-	-	18	-	-	1	5	96	37.50	0.00
600-02	-	-	31	1	-	-	-	-	4	-	-	11	-	-	4	7	101	60.40	0.00
604-06	-	-	18	-	-	-	-	-	1	-	-	6	-	-	4	3	98	33.67	0.00
608-10	-	-	6	-	-	-	2	-	-	-	-	2	-	-	2	-	101	12.87	0.00
612-14	-	-	14	-	-	-	-	-	-	-	-	4	-	-	4	2	93	25.81	0.00
616-18	-	-	14	-	-	-	-	-	-	-	-	3	-	-	3	2	99	25.25	2.02
620-22	-	-	8	-	-	-	-	-	3	-	-	23	-	-	-	2	98	39.80	0.00
624-26	-	-	24	-	-	-	-	-	9	-	-	22	-	-	11	5	97	77.32	0.00
628-30	-	-	9	-	-	-	-	-	5	-	-	7	-	-	5	3	98	32.65	0.00
632-34	-	-	10	-	-	-	-	-	7	-	-	18	-	-	5	2	94	46.81	0.00
636-38	-	-	11	-	-	-	1	-	6	-	-	16	-	-	2	1	99	37.37	0.00
640-42	-	33	1	-	-	-	-	-	-	-	8	5	-	-	-	1	100	7.00	87.00
644-46	-	-	9	-	-	-	-	-	3	-	-	16	-	-	5	2	97	38.14	0.00
648-50	-	-	11	-	-	-	-	-	4	-	-	33	-	-	1	5	99	54.55	0.00
652-54	-	-	12	-	-	-	-	-	2	-	-	20	-	-	-	8	97	44.33	0.00
656-58	-	-	7	-	-	-	-	-	3	-	-	16	-	-	1	5	102	31.37	0.00
660-62	-	-	10	-	-	-	-	-	1	-	-	15	-	-	2	5	99	35.35	0.00
664-66	-	-	3	-	-	-	-	-	-	-	-	3	-	-	1	2	98	10.20	0.00
668-70	-	-	12	-	-	-	-	-	2	-	-	10	-	-	3	1	100	28.00	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuilina minuta</i>	Total	Percent Agglutinated	Percent Calcareous
672-74	-	-	5	-	-	-	-	-	-	-	-	20	-	-	2	-	96	29.17	0.00
676-78	-	-	7	-	-	-	-	-	-	-	1	16	-	-	7	1	111	29.73	0.90
680-82	-	-	5	-	-	-	-	-	1	-	-	4	-	-	1	1	99	12.12	0.00
684-86	-	-	9	-	-	-	1	-	1	-	-	7	-	-	1	2	99	21.21	0.00
688-90	-	-	13	-	-	-	-	-	-	-	-	5	-	-	4	5	102	28.43	0.00
692-94	-	-	3	-	-	-	-	-	-	-	-	-	-	-	3	-	100	6.00	0.00
700-02	-	-	10	-	-	1	1	-	1	-	-	4	-	-	2	7	95	27.37	0.00
704-06	-	-	21	-	-	-	-	-	-	-	-	-	-	-	4	4	96	30.21	0.00
708-10	-	-	27	-	-	-	-	-	3	-	-	6	-	-	6	4	95	48.42	1.05
712-14	-	-	4	-	-	-	-	-	2	-	-	3	-	-	5	-	19	73.68	0.00
716-18	-	-	4	-	-	-	-	-	-	-	-	16	-	-	9	4	44	75.00	0.00
720-22	-	-	3	-	-	-	-	-	-	-	-	16	-	-	7	5	100	31.00	0.00
724-26	-	-	13	-	-	-	-	-	6	-	-	16	-	-	6	2	103	42.72	0.00
728-30	-	-	6	-	-	-	-	-	1	-	-	1	-	-	1	3	102	12.75	0.00
732-34	-	-	11	-	-	-	-	-	-	-	-	7	-	-	5	2	99	26.26	0.00
736-38	-	-	3	-	-	-	-	-	1	-	-	4	-	-	3	-	101	10.89	10.89
740-42	-	-	9	-	-	-	-	-	2	-	-	8	-	-	-	5	103	23.30	0.00
744-46	-	-	11	-	-	-	-	-	-	-	-	8	-	-	2	1	100	22.00	0.00
748-50	-	-	6	-	-	-	-	-	6	-	-	8	-	-	2	1	97	23.71	0.00
752-54	-	-	8	-	-	-	-	-	2	-	1	21	-	-	-	1	101	31.68	4.95
756-58	-	-	6	-	-	-	-	-	1	-	-	9	-	-	4	-	104	19.23	0.00
760-62	-	-	6	-	-	-	-	-	1	-	-	14	-	-	3	8	100	33.00	0.00
764-66	-	-	18	-	-	1	-	-	4	-	-	11	-	-	1	1	101	35.64	0.00
768-70	-	-	7	-	-	-	-	-	2	-	-	32	-	-	2	3	97	47.42	0.00
772-74	-	-	4	-	-	-	-	-	1	-	-	3	-	-	3	-	99	11.11	1.01
776-78	-	-	13	-	-	-	-	-	4	-	-	11	-	-	-	4	101	31.68	0.00
780-82	-	-	11	-	-	-	-	-	2	-	-	4	-	-	4	-	102	20.59	0.00
784-86	-	-	20	-	-	-	-	-	2	-	-	4	-	-	1	-	99	27.27	0.00
788-90	-	-	4	-	-	-	-	-	-	-	-	1	-	-	2	-	101	6.93	0.00
792-94	-	-	11	-	-	-	-	-	7	-	-	10	-	-	2	2	98	32.65	0.00
796-98	-	-	4	-	-	-	-	-	-	-	-	1	-	-	2	1	99	8.08	0.00
800-02	-	-	26	-	-	-	-	-	6	-	-	8	-	-	3	3	97	47.42	0.00
804-06	-	-	21	-	-	-	-	-	1	-	-	3	-	-	1	2	97	29.90	0.00
808-10	2	27	9	-	-	-	-	-	1	-	2	7	-	-	1	-	101	17.82	39.60
812-14	-	50	1	-	-	-	-	-	-	-	18	-	-	-	1	-	99	2.02	75.76
816-18	-	-	3	-	-	-	-	-	1	-	-	1	-	-	1	1	101	6.93	0.00
820-22	-	-	4	-	-	-	-	-	5	-	-	1	-	-	6	4	97	20.62	5.15
824-26	-	24	2	-	-	-	-	-	-	-	8	-	-	-	-	1	100	3.00	66.00
828-30	-	-	14	-	-	-	-	-	5	-	-	5	-	-	7	4	103	33.98	0.00
832-34	-	-	11	-	-	-	-	-	-	-	-	-	-	-	1	-	101	11.88	0.00
836-38	-	-	6	-	-	-	-	-	2	-	-	-	-	-	-	-	103	7.77	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentatiniiformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuliara minuta</i>	Total	Percent Agglutinated	Percent Calcareous
840-42	-	-	20	-	-	-	-	-	3	-	-	5	-	-	1	-	97	29.90	1.03
844-46	-	-	1	-	-	-	-	-	-	-	-	2	-	-	5	-	102	7.84	0.00
848-50	-	-	7	-	-	-	-	-	-	-	-	4	-	-	2	-	99	13.13	0.00
852-54	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	100	1.00	0.00
856-58	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	-	101	1.98	0.00
860-62	-	-	5	-	-	-	-	-	1	-	-	2	-	-	-	-	99	8.08	0.00
864-66	-	-	8	-	-	-	-	-	1	-	-	5	-	-	1	1	102	15.69	0.98
868-70	-	-	16	-	-	-	-	-	1	-	-	17	-	-	2	5	106	38.68	0.00
872-74	-	-	6	-	-	-	-	-	-	-	-	-	-	-	-	-	95	6.32	0.00
876-78	-	-	17	-	-	-	-	-	-	-	-	2	-	-	2	1	100	22.00	0.00
880-82	-	-	8	-	-	-	-	-	-	-	-	-	-	-	4	-	37	32.43	0.00
884-86	-	-	4	-	-	-	-	-	-	-	-	-	-	-	2	-	100	6.00	0.00
888-90	-	-	3	-	-	-	-	-	-	-	-	-	-	-	2	-	80	6.25	0.00
892-94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	30	3.33	0.00
896-98	-	-	5	-	-	-	-	-	-	-	-	-	-	-	-	-	104	4.81	0.00
900-02	-	-	2	-	-	-	-	-	-	-	-	-	-	-	1	-	98	3.06	0.00
904-06	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	102	0.98	2.94
908-10	-	-	-	-	-	-	-	-	1	-	-	1	-	-	-	-	101	1.98	0.00
912-14	-	-	2	-	-	-	-	-	-	-	-	1	-	-	1	-	98	4.08	0.00
916-18	-	-	2	-	-	-	-	-	-	-	-	-	-	-	1	-	103	2.91	0.00
920-22	-	-	4	-	-	-	-	-	-	-	-	-	-	-	-	2	101	5.94	0.00
924-26	-	-	30	-	-	-	-	-	1	-	-	1	-	-	1	1	101	33.66	0.00
928-30	-	-	3	-	-	-	-	-	1	-	-	-	-	-	2	1	102	6.86	0.00
932-34	-	-	4	-	-	-	-	-	-	-	-	1	-	-	2	-	100	7.00	0.00
936-38	-	-	1	-	-	-	-	-	2	-	-	1	-	-	4	-	100	8.00	0.00
940-42	-	-	5	-	-	-	-	-	1	-	-	1	-	-	-	1	99	8.08	0.00
944-46	-	-	5	-	-	-	-	-	-	-	-	-	-	-	1	-	98	6.12	0.00
948-50	-	-	8	-	-	-	-	-	-	-	-	-	-	-	-	-	104	7.69	0.00
952-54	-	-	5	-	-	-	-	-	1	-	-	-	-	-	1	-	100	7.00	0.00
956-58	-	-	1	-	-	-	-	-	1	-	-	-	-	-	-	-	100	2.00	0.00
960-62	-	-	8	-	-	-	-	-	2	-	-	-	-	-	-	-	101	9.90	0.00
964-66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-
968-70	-	-	15	-	-	-	-	-	4	-	-	-	-	-	-	1	102	19.61	0.00
972-74	-	-	10	-	-	-	-	-	-	-	-	2	-	-	-	-	100	12.00	0.00
976-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	102	0.00	0.00
980-82	-	-	2	-	-	-	-	-	-	-	-	1	-	-	-	-	101	2.97	0.00
984-86	-	-	6	-	-	-	-	-	-	-	-	-	-	-	1	-	99	7.07	0.00
988-90	-	-	11	-	-	-	-	-	-	-	-	-	-	-	1	-	106	11.32	0.00
992-94	-	-	8	-	-	-	-	-	-	-	-	-	-	-	1	-	100	9.00	0.00
996-98	-	-	22	-	-	-	-	-	1	-	-	-	-	-	5	-	100	28.00	5.00
1000-02	-	-	12	-	-	-	-	-	2	-	-	-	-	-	5	-	95	20.00	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentatiniiformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuliara minuta</i>	Total	Percent Agglutinated	Percent Calcareous
1004-06	-	-	2	-	-	-	-	-	-	-	-	1	-	-	5	-	65	12.31	3.08
1008-10	-	1	2	-	-	-	-	-	-	-	-	2	-	-	3	-	102	6.86	1.96
1012-14	-	0	-	-	-	-	-	-	-	-	4	-	-	-	1	-	101	0.99	75.25
1016-18	-	-	6	-	-	-	-	-	-	-	-	-	-	-	2	-	103	7.77	42.72
1020-22	-	-	38	-	-	1	-	-	1	-	-	-	-	-	3	-	102	42.16	0.00
1024-26	-	-	9	-	-	-	-	-	2	-	-	-	-	-	5	-	100	16.00	0.00
1028-30	-	-	10	-	-	-	-	-	-	-	-	-	-	-	3	-	98	13.27	0.00
1032-34	-	-	4	-	-	-	-	-	1	-	-	-	-	-	1	-	102	5.88	0.00
1036-38	-	-	17	-	-	-	-	-	1	-	-	-	-	-	1	-	102	18.63	0.00
1040-42	-	-	20	-	-	-	-	-	-	-	-	1	-	-	8	-	87	33.33	0.00
1044-46	-	-	7	-	-	-	-	-	-	-	-	-	-	-	4	-	104	10.58	16.35
1048-50	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	98	1.02	0.00
1052-54	-	-	4	-	-	-	-	-	-	-	-	-	-	-	-	-	101	3.96	0.99
1056-58	-	-	9	-	-	-	-	-	-	-	-	-	-	-	1	-	102	9.80	0.00
1060-62	-	-	5	-	-	-	-	-	-	-	-	-	-	-	1	-	101	5.94	2.97
1064-66	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	101	1.98	0.99
1068-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	1	102	3.92	1.96
1072-74	-	-	1	-	-	-	-	-	-	-	-	-	-	-	2	-	99	3.03	1.01
1076-78	-	-	3	-	-	-	-	-	-	-	1	-	-	-	-	-	96	3.13	2.08
1080-82	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	-	101	1.98	0.00
1084-86	-	-	1	-	-	-	-	-	-	-	-	-	-	-	2	-	101	2.97	0.00
1088-90	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	100	1.00	0.00
1092-94	-	-	3	-	-	-	-	-	-	-	-	-	-	-	1	-	100	4.00	0.00
1096-98	-	-	1	-	-	-	-	-	1	-	-	-	-	-	2	-	99	4.04	0.00
1100-02	-	-	5	-	-	-	-	-	-	-	-	-	-	-	1	-	102	5.88	0.00
1104-06	-	-	7	-	-	-	-	-	-	-	-	-	-	-	-	-	100	7.00	1.00
1108-10	-	-	7	-	-	-	-	-	1	-	-	-	-	-	1	2	100	11.00	0.00
1112-14	-	-	4	-	-	-	-	-	1	-	-	-	-	-	1	-	98	6.12	0.00
1116-18	-	-	1	-	-	-	-	-	-	-	3	-	-	-	-	-	101	0.99	26.73
1120-22	-	-	8	-	-	-	-	-	-	-	-	-	-	-	2	-	104	9.62	0.00
1124-26	-	-	2	-	-	-	-	-	-	-	-	-	-	-	1	-	103	2.91	0.00
1128-30	-	-	10	-	-	-	-	-	-	-	-	-	-	-	-	-	100	10.00	0.00
1132-34	-	4	9	-	-	-	-	-	-	-	-	1	-	-	3	-	100	13.00	4.00
1136-38	-	-	8	-	-	-	-	-	-	-	-	-	-	-	1	-	100	9.00	1.00
1140-42	-	7	1	-	-	-	-	-	-	-	2	-	-	-	2	-	104	2.88	38.46
1144-46	-	-	9	-	-	-	-	-	-	-	-	-	-	-	-	-	100	9.00	0.00
1148-50	-	-	16	-	-	-	1	-	-	-	-	-	-	-	1	-	99	18.18	0.00
1152-54	1	6	3	-	-	-	1	-	-	-	5	-	-	-	-	-	101	3.96	20.79
1156-58	-	3	8	-	-	-	-	-	-	-	5	-	-	-	1	-	101	8.91	15.84
1160-62	-	9	3	-	-	-	-	-	-	2	14	-	-	-	4	-	98	7.14	30.61
1164-66	-	8	2	-	-	-	-	-	-	-	49	-	-	-	8	-	101	9.90	60.40

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina bifurcata</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuilina minuta</i>	Total	Percent Agglutinated	Percent Calcareous
1168-70	-	4	9	-	-	-	-	-	-	-	33	-	-	-	-	-	98	9.18	40.82
1172-74	-	1	5	-	-	-	-	-	-	-	-	1	-	-	-	-	101	5.94	23.76
1176-78	-	-	7	-	-	-	1	-	-	-	-	1	-	-	1	-	99	10.10	0.00
1180-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	100	1.00	0.00
1184-86	-	-	4	-	-	-	-	-	-	-	-	1	-	-	1	-	105	5.71	0.00
1188-90	-	-	1	-	-	-	-	-	-	-	-	-	-	-	3	-	99	4.04	0.00
1192-94	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	99	2.02	0.00
1196-98	-	-	2	-	-	-	2	-	-	-	-	-	-	-	1	-	101	4.95	0.00
1200-02	-	-	7	-	-	-	-	-	-	-	-	-	-	-	1	1	101	8.91	0.00
1204-06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	100	1.00	0.00
1208-10	-	-	6	-	-	-	-	-	-	-	-	-	-	-	3	-	102	8.82	0.00
1212-14	-	-	6	-	-	-	2	-	-	-	-	-	-	-	-	-	102	7.84	0.00
1216-18	-	-	6	-	-	-	-	-	2	-	-	-	-	-	1	-	103	8.74	0.00
1220-22	-	-	5	-	-	-	-	-	1	-	-	-	-	-	3	-	103	8.74	0.00
1224-26	-	-	10	-	-	-	-	-	3	-	-	-	-	-	-	-	98	13.27	0.00
1228-30	-	-	6	-	-	-	-	-	-	-	-	-	-	-	-	-	101	5.94	0.00
1232-34	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	97	2.06	0.00
1236-38	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	102	0.98	0.00
1240-42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	98	0.00	0.00
1244-46	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	102	0.98	0.00
1248-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	100	1.00	0.00
1252-54	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	-	98	2.04	0.00
1256-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	41	2.44	0.00
1260-62	-	-	2	-	-	-	-	-	-	-	-	1	-	-	-	-	100	3.00	0.00
1264-66	-	-	4	-	-	-	-	-	-	-	-	-	-	-	-	-	97	4.12	0.00
1268-70	-	-	1	-	-	-	-	-	1	-	-	-	-	-	-	-	102	1.96	0.00
1272-74	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	98	1.02	0.00
1276-78	-	-	3	-	-	-	-	-	-	-	-	-	-	-	1	-	99	4.04	0.00
1280-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	54	0.00	0.00
1284-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	100	1.00	0.00
1288-90	-	-	10	-	-	-	-	-	-	-	-	-	-	-	2	-	100	12.00	0.00
1292-94	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	99	2.02	0.00
1296-98	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	-	100	2.00	0.00
1300-02	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	101	1.98	0.99
1304-06	-	-	8	-	-	-	-	-	1	-	-	-	-	-	1	-	99	10.10	1.01
1308-10	-	-	2	-	-	-	-	-	-	-	-	-	-	-	1	-	102	2.94	0.00
1312-14	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	102	1.96	0.00
1316-18	-	-	8	-	-	-	-	-	-	-	-	-	-	-	1	-	101	8.91	0.00
1320-22	-	-	7	-	-	-	-	-	-	-	-	-	-	-	-	-	105	6.67	0.00
1324-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	100	1.00	0.00
1328-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	99	2.02	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentatiniiformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuliara minuta</i>	Total	Percent Agglutinated	Percent Calcareous
1332-34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	100	1.00	0.00
1336-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	101	0.99	0.00
1340-42	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	101	1.98	0.00
1344-46	-	-	7	-	-	-	-	-	1	-	-	-	-	-	-	-	99	8.08	0.00
1348-50	-	2	1	-	-	-	-	-	-	-	7	-	-	-	-	-	102	0.98	10.78
1352-54	-	-	6	-	-	-	-	-	1	-	-	-	-	-	-	-	98	7.14	0.00
1356-58	-	-	27	-	-	-	-	-	-	-	-	-	-	-	-	-	122	22.13	0.00
1360-62	-	-	20	-	-	-	-	-	1	-	-	-	-	-	1	-	101	21.78	0.00
1364-66	-	-	7	-	-	-	-	-	-	-	-	-	-	-	1	-	99	8.08	0.00
1368-70	-	-	31	-	-	-	-	-	1	-	-	-	-	-	-	-	93	34.41	0.00
1372-74	-	-	20	-	-	-	-	-	1	-	-	-	-	-	1	-	100	22.00	0.00
1376-78	-	-	3	-	-	-	-	-	-	-	-	-	-	-	1	-	100	4.00	1.00
1380-82	-	-	19	-	-	-	-	-	-	-	-	-	-	-	-	-	98	19.39	0.00
1384-86	-	-	41	-	-	-	-	-	2	-	-	-	-	-	3	-	102	45.10	0.00
1388-90	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	95	2.11	2.11
1392-94	-	-	35	-	-	-	-	-	1	-	1	-	-	-	-	-	98	36.73	1.02
1396-98	-	-	26	-	-	-	-	-	1	-	-	-	-	-	-	-	99	27.27	1.01
1400-02	-	-	4	-	-	-	-	-	1	-	-	-	-	-	-	-	98	5.10	0.00
1404-06	-	-	7	-	-	-	-	-	-	-	-	-	-	-	1	-	100	8.00	0.00
1408-10	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	97	2.06	0.00
1412-14	-	-	21	-	-	-	-	-	1	-	-	-	-	-	-	-	101	21.78	0.00
1416-18	-	-	7	-	-	-	-	-	-	-	10	-	-	-	-	-	99	7.07	12.12
1420-22	-	-	12	-	-	-	-	-	-	-	26	-	-	-	-	-	102	11.76	26.47
1424-26	-	-	10	-	-	-	-	-	-	-	8	-	-	-	1	-	91	12.09	9.89
1428-30	-	-	4	-	-	-	-	-	-	-	-	-	-	-	1	-	101	4.95	0.00
1432-34	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	95	2.11	28.42
1436-38	1	1	-	-	-	-	-	-	2	-	14	-	-	-	-	-	105	1.90	52.38
1440-42	1	-	2	-	-	-	-	-	-	-	14	-	-	-	-	-	102	1.96	47.06
1444-46	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	97	2.06	0.00
1448-50	-	-	2	-	-	-	-	-	1	-	-	-	-	-	-	-	101	2.97	0.00
1452-54	-	-	21	-	-	-	-	-	1	-	-	-	-	-	-	-	103	21.36	0.00
1456-58	-	-	6	-	-	-	-	-	2	-	-	-	-	-	1	-	97	9.28	2.06
1460-62	-	-	4	-	-	-	-	-	-	-	1	-	-	-	1	-	102	4.90	1.96
1464-66	-	-	18	-	-	-	-	-	-	-	3	-	-	-	-	-	103	20.39	0.00
1468-70	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	-	98	2.04	0.00
1472-74	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	97	1.03	0.00
1476-78	-	-	8	-	-	-	-	-	-	-	-	-	-	-	-	-	100	8.00	0.00
1480-82	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	97	0.00	0.00
1484-86	-	-	6	-	-	-	-	-	-	-	-	-	-	-	4	-	96	10.42	1.04
1488-90	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	99	2.02	0.00
1492-94	-	-	12	-	-	-	-	-	-	-	-	-	-	-	-	-	100	12.00	0.00

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuliara minuta</i>	Total	Percent Agglutinated	Percent Calcareous
1496-98	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	40	5.00	0.00
1500-02	-	-	4	-	-	-	-	-	-	-	-	-	-	-	-	-	75	5.33	0.00
1504-06	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	99	2.02	0.00
1508-10	-	-	3	-	-	-	-	-	-	-	-	-	-	-	-	-	98	3.06	0.00
1512-14	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	100	2.00	0.00
1516-18	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	22	0.00	0.00
1520-22	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	100	2.00	0.00
1524-26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	0.00	0.00
1528-30	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	100	1.00	0.00
1532-34	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	43	2.33	0.00
1536-38	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	0.00	0.00
1540-42	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	100	1.00	0.00
1544-46	-	-	17	-	-	-	-	-	-	-	3	-	-	-	-	-	99	17.17	6.06
1548-50	-	-	6	-	-	-	-	-	1	-	-	-	-	-	1	-	100	8.00	0.00
1552-54	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	100	1.00	0.00
1556-58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	0.00	0.00
1560-62	-	-	6	-	-	-	-	-	2	-	-	-	-	-	1	-	64	14.06	0.00
1564-66	-	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	100	2.00	0.00
1568-70	-	-	1	-	-	-	-	-	-	-	-	-	-	-	1	-	100	2.00	0.00
1572-74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	0.00	0.00
1576-78	-	-	6	-	-	-	-	-	2	-	-	-	-	-	-	-	100	8.00	2.00
1580-82	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	100	1.00	0.00
1584-86	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	0.00	0.00
1588-90	-	-	5	-	-	-	-	-	-	-	-	-	-	-	-	-	100	5.00	0.00
1592-94	-	-	3	-	-	-	-	-	-	-	-	-	-	-	2	-	60	8.33	3.33
1596-98	-	-	4	-	-	-	-	-	2	-	-	-	-	-	-	-	100	6.00	0.00
1600-02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	100	0.00	0.00
1604-06	-	-	15	-	-	-	-	-	-	-	1	-	-	-	-	-	83	18.07	7.23
1608-10	-	15	4	-	-	-	-	-	-	42	-	-	-	-	-	-	101	3.96	71.29
1612-14	-	11	1	-	-	-	-	-	-	45	-	-	-	-	-	-	97	1.03	69.07
1616-18	-	17	3	-	-	-	-	-	-	53	-	-	-	-	-	-	100	3.00	82.00
1620-22	2	11	-	-	-	-	-	-	-	68	-	-	-	-	-	-	98	0.00	86.73
1624-26	-	5	2	-	-	-	-	-	-	77	-	-	-	-	-	-	97	2.06	89.69
1628-30	-	6	6	-	-	-	-	1	-	67	-	-	-	1	-	-	100	7.00	83.00
1632-34	-	12	1	-	-	-	-	1	-	49	-	-	-	-	-	-	98	1.02	87.76
1636-38	-	-	4	-	-	-	-	-	-	-	-	-	-	-	-	-	100	4.00	0.00
1640-42	-	43	-	-	-	-	-	-	-	27	-	-	-	-	-	-	99	0.00	95.96
1644-46	-	-	33	-	-	-	-	-	-	-	-	-	-	-	3	-	97	37.11	3.09
1648-50	1	26	-	-	-	-	-	-	-	44	-	-	-	-	-	-	101	0.00	95.05
1652-54	1	10	10	-	-	-	-	-	-	17	-	-	-	1	-	-	102	10.78	45.10
1656-58	1	11	3	-	-	-	-	-	-	38	-	-	-	-	-	-	99	3.03	74.75

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Psuedobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuliara minuta</i>	Total	Percent Agglutinated	Percent Calcareous
1660-62	1	23	2	-	-	-	-	-	-	-	48	-	-	-	-	-	97	2.06	90.72
1664-66	-	20	4	-	-	-	-	-	-	-	45	-	-	-	-	-	90	4.44	88.89
1668-70	-	41	3	-	-	-	-	-	-	-	23	-	-	-	-	-	97	3.09	85.57
1672-74	-	7	7	-	-	-	-	-	-	-	39	-	-	-	-	-	100	7.00	77.00
1676-78	-	9	8	-	-	-	-	-	-	-	34	-	-	-	-	-	102	7.84	68.63
1680-82	-	4	9	-	-	-	-	-	-	-	54	-	-	-	-	-	98	9.18	67.35
1684-86	-	9	1	-	-	-	-	-	-	1	60	-	-	-	1	-	100	2.00	76.00
1688-90	1	13	1	-	-	-	-	-	-	-	58	-	-	-	1	-	98	2.04	85.71
1692-94	-	57	2	-	-	-	-	-	-	-	18	-	-	-	-	-	97	2.06	92.78
1696-98	-	52	4	-	-	-	-	-	-	-	22	1	-	-	-	-	102	4.90	83.33
1700-02	2	32	6	-	-	-	-	-	-	-	28	-	-	-	-	-	99	6.06	82.83
1704-06	-	1	4	-	-	-	-	-	-	-	15	-	-	-	-	-	100	4.00	19.00
1708-10	-	19	5	-	-	-	-	-	1	-	30	-	-	-	-	-	101	5.94	60.40
1712-14	-	26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	87	0.00	63.22
1716-18	-	57	-	-	-	-	-	-	-	-	-	-	-	-	-	-	70	0.00	95.71
1720-22	1	42	2	-	-	-	-	-	-	-	35	-	-	-	-	-	96	2.08	86.46
1724-26	-	44	-	-	-	-	-	-	-	-	29	-	-	-	-	-	109	0.00	93.58
1728-30	-	26	2	-	-	-	-	-	-	-	31	-	-	-	1	-	101	2.97	72.28
1732-34	-	33	1	-	-	-	-	-	-	-	33	-	-	-	-	-	102	0.98	84.31
1736-38	-	18	4	-	-	-	-	-	-	-	19	-	-	-	-	-	97	4.12	52.58
1740-42	2	16	6	-	-	-	-	-	-	-	16	-	-	-	1	-	96	7.29	58.33
1744-46	-	17	-	-	-	-	-	-	-	-	47	-	-	-	-	-	103	0.00	78.64
1748-50	-	16	1	-	-	-	-	-	-	-	30	-	-	-	-	-	91	1.10	69.23
1752-54	-	23	2	-	-	-	-	-	-	-	20	-	-	-	-	-	101	1.98	47.52
1756-58	-	12	1	-	-	-	-	-	-	-	35	-	-	-	-	-	102	0.98	50.00
1760-62	-	9	-	-	-	-	-	-	-	-	37	-	-	-	-	-	99	0.00	53.54
1764-766	-	2	6	-	-	-	-	-	-	-	30	-	-	-	-	-	100	6.00	40.00
1768-770	-	5	2	-	-	-	-	-	-	-	41	-	-	-	-	-	102	1.96	50.00
1772-74	-	18	-	-	-	-	-	-	-	-	46	-	-	-	-	-	103	0.00	68.93
1776-78	-	23	1	-	-	-	-	-	-	-	18	-	-	-	-	-	105	0.95	49.52
1780-82	-	19	3	-	-	-	-	-	-	-	15	-	-	-	-	-	99	3.03	38.38
1784-86	-	15	-	-	-	-	-	-	-	-	15	-	-	-	-	-	102	0.00	31.37
1788-90	-	25	-	-	-	-	-	-	-	-	17	-	-	-	-	-	102	0.00	49.02
1792-94	-	12	2	-	-	-	-	-	-	-	16	-	1	-	-	-	97	3.09	29.90
1796-98	-	5	-	-	-	-	-	-	-	-	13	-	-	-	-	-	102	0.00	17.65
1800-02	-	2	-	-	-	-	-	-	-	-	14	-	-	-	-	-	97	0.00	17.53
1804-06	-	14	1	-	-	-	-	-	-	-	14	-	-	-	-	-	100	1.00	33.00
1808-10	-	14	-	-	-	-	-	-	-	-	7	-	-	-	-	-	101	0.00	21.78
1812-14	-	5	-	-	-	-	-	-	-	-	5	-	-	-	-	-	99	0.00	12.12
1816-18	-	2	-	-	-	-	-	-	-	-	14	-	-	-	-	-	102	0.00	17.65
1820-22	-	5	-	-	-	-	-	-	-	-	8	-	-	-	-	-	94	0.00	13.83

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	Nodosarids	<i>Nonionella iridea</i>	<i>Portatrochammina</i> spp.	<i>Pseudobolivina antarctica</i>	<i>Pullinella</i> sp.	<i>Rhabdammina abyssorum</i>	<i>Reophax subdentaliniformis</i>	<i>Spiroloculina</i> spp.	<i>Spiroplectommina biformis</i>	<i>Stainforthia davisii</i>	<i>Stainforthia fusiformis</i>	<i>Textularia wiesneri</i>	<i>Trochammina bullata</i>	<i>Trochammina conica</i>	<i>Trochammina intermedia</i>	<i>Verneuliara minuta</i>	Total	Percent Agglutinated	Percent Calcareous
1824-26	-	1	-	-	-	-	-	-	-	-	15	1	-	-	-	-	106	0.94	17.92
1828-30	-	5	-	-	-	-	-	-	-	-	25	-	-	-	-	-	104	0.00	32.69
1832-34	-	5	-	-	-	-	-	-	-	-	20	-	-	-	-	-	98	0.00	26.53
1836-38	-	8	-	-	-	-	-	-	-	-	46	-	-	-	-	-	104	0.00	60.58
1840-42	-	6	-	-	-	-	-	-	-	-	47	-	-	-	-	-	102	0.00	56.86
1844-46	-	12	1	-	-	-	-	2	-	-	28	-	-	-	-	-	98	1.02	46.94
1848-50	-	19	-	-	-	-	-	-	-	-	19	-	-	-	-	-	97	0.00	45.36
1852-54	-	15	-	-	-	-	-	-	-	-	17	-	-	-	1	-	101	0.99	34.65
1856-58	1	11	-	-	-	-	-	-	-	-	41	-	-	-	1	-	102	0.98	54.90
1860-62	-	1	1	-	-	-	-	-	-	-	13	-	-	-	-	-	98	1.02	17.35
1864-66	-	2	1	-	-	-	-	-	-	-	6	-	-	-	1	-	101	1.98	7.92
1868-70	-	5	1	-	-	-	-	-	-	-	8	-	-	-	1	-	102	1.96	13.73
1872-74	-	5	1	-	-	-	-	1	-	-	11	-	-	-	-	-	102	0.98	16.67
1876-78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-
1880-82	-	9	-	-	-	-	-	-	-	-	41	-	-	-	-	-	98	0.00	52.04
1884-86	-	4	-	-	-	-	-	-	-	-	8	-	-	-	-	-	100	0.00	15.00
1888-90	-	32	-	-	-	-	-	-	-	-	26	-	-	-	1	-	101	0.99	60.40
1892-94	-	34	-	-	-	-	-	-	-	-	22	-	-	-	-	-	102	0.00	61.76
1896-98	1	10	-	-	-	-	-	-	-	-	32	-	-	-	-	-	97	0.00	45.36
1900-02	-	13	-	-	-	-	-	-	-	-	42	-	-	-	-	-	100	0.00	57.00
1904-06	-	22	1	-	-	-	-	-	-	-	36	-	-	-	-	-	96	1.04	71.88
1908-10	-	12	-	-	-	-	-	-	-	-	47	-	-	-	-	-	100	0.00	67.00
1912-14	-	3	-	-	-	-	-	-	-	-	21	-	-	-	-	-	101	0.00	30.69
1916-18	-	3	-	-	-	-	-	-	-	2	11	-	-	-	-	-	103	0.00	17.48
1920-22	-	5	-	-	-	-	-	-	-	-	21	-	-	-	-	-	101	0.00	29.70
1924-26	-	3	-	-	-	-	1	-	-	-	34	-	-	-	-	-	102	0.98	42.16
1928-30	-	4	-	-	-	-	1	-	-	-	10	-	-	-	-	-	100	1.00	19.00
1932-34	-	11	1	-	-	-	-	-	-	-	12	-	-	-	2	-	102	2.94	28.43
1936-38	-	10	2	-	1	-	-	-	-	-	7	-	-	-	1	-	96	3.13	20.83
1940-42	-	6	-	-	-	-	-	-	-	-	34	-	-	-	-	-	97	0.00	44.33
1944-46	-	7	-	-	-	-	-	-	-	-	60	-	-	-	1	-	101	0.99	75.25
1948-50	-	3	-	-	-	-	-	-	-	-	78	-	-	-	-	-	97	0.00	86.60
1952-54	-	1	-	-	-	-	-	-	-	2	16	-	-	-	4	-	100	4.00	40.00
1956-58	-	-	-	-	-	-	-	19	-	-	1	-	-	-	8	-	102	7.84	19.61
1960-62	-	-	-	-	-	-	-	-	-	-	5	-	-	-	4	-	101	3.96	5.94
1964-66	-	21	-	-	-	-	-	-	-	-	64	-	-	-	2	-	101	1.98	93.07
1968-70	-	18	-	-	-	-	-	-	-	-	36	-	-	-	1	-	101	0.99	55.45
1972-74	-	44	1	-	-	-	-	-	-	-	26	-	-	-	-	-	90	1.11	90.00
1976-78	-	34	-	-	-	-	-	-	-	-	32	-	-	-	-	-	100	0.00	75.00
1980-82	-	5	-	-	-	-	-	-	-	-	78	-	-	-	1	-	99	1.01	83.84
1984-86	1	10	-	-	-	-	-	-	-	-	68	-	-	-	-	-	99	0.00	79.80

*Planktonic species not used in statistical analysis

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
0-2	0.26	0.95	0.01
4-6	0.23	0.97	0.01
8-10	0.03	0.99	0.03
12-14	0.06	0.98	0.04
16-18	0.22	0.97	0.02
20-22	0.65	0.75	0.07
24-26	0.24	0.96	0.01
28-30	0.05	0.99	0.01
32-34	0.11	0.99	0.03
36-38	0.04	1.00	0.01
40-42	0.02	0.99	0.03
44-46	0.16	0.98	0.04
48-50	0.10	0.96	0.04
52-54	0.10	0.97	0.01
56-58	0.54	0.83	0.05
60-62	0.13	0.95	0.01
64-66	0.05	0.99	0.03
68-70	0.22	0.97	0.01
76-78	0.10	0.99	0.00
80-80	0.05	0.99	0.02
84-86	0.38	0.92	0.03
88-90	0.09	0.99	0.00
92-94	0.26	0.96	0.02
96-98	0.36	0.93	0.03
100-02	0.23	0.96	0.02
104-06	0.49	0.87	0.05
112-14	0.28	0.96	0.02
116-18	0.18	0.97	0.01
120-22	0.33	0.94	0.03
128-30	0.04	0.99	0.01
132-34	0.32	0.94	0.03
136-38	0.22	0.97	0.02
140-42	0.16	0.98	0.01
144-46	0.66	0.75	0.07
148-50	0.26	0.96	0.02
152-54	0.07	0.99	0.01
156-58	0.75	0.62	0.07
160-62	0.16	0.96	0.01
164-66	0.11	0.98	0.01
168-70	0.33	0.94	0.03
172-74	0.87	0.47	0.10

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
176-78	0.44	0.89	0.05
180-82	0.28	0.82	0.00
184-86	0.76	0.64	0.08
188-90	0.91	0.39	0.10
192-94	0.87	0.45	0.09
196-98	0.53	0.84	0.05
200-02	0.41	0.91	0.03
204-06	0.31	0.94	0.03
208-10	0.13	0.98	0.09
212-14	0.95	0.29	0.11
216-18	0.89	0.43	0.10
220-22	0.16	0.28	0.13
224-26	0.33	0.13	0.21
228-30	0.63	0.77	0.09
232-34	0.70	0.62	0.05
236-38	0.29	0.94	0.01
240-42	0.67	0.73	0.07
244-46	0.07	0.99	0.00
248-50	0.73	0.67	0.08
252-54	0.28	0.37	0.75
256-58	0.08	0.97	0.21
260-62	0.82	0.52	0.08
264-66	0.60	0.25	0.51
268-70	0.56	0.19	0.36
272-74	0.62	0.75	0.05
276-78	0.94	0.32	0.11
280-82	0.78	0.21	0.22
284-86	0.70	0.54	0.04
288-90	0.43	0.88	0.04
292-94	0.91	0.40	0.11
296-98	0.77	0.63	0.08
300-02	0.72	0.68	0.08
304-06	0.79	0.61	0.09
308-10	0.75	0.65	0.08
312-14	0.91	0.32	0.10
316-18	0.87	0.48	0.10
320-22	0.92	0.36	0.11
324-26	0.69	0.71	0.07
328-30	0.62	0.78	0.07
332-34	0.31	0.94	0.02
336-38	0.53	0.82	0.04

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
340-42	0.45	0.88	0.04
348-50	0.90	0.43	0.11
352-54	0.01	0.97	0.03
356-58	0.65	0.21	0.38
360-62	0.14	0.28	0.08
364-66	0.14	0.16	0.34
368-70	0.18	0.93	0.05
372-74	0.95	0.27	0.12
376-78	0.79	0.59	0.16
380-82	0.39	0.83	0.16
384-86	0.08	0.04	0.85
388-90	0.95	0.23	0.11
392-94	0.94	0.30	0.11
396-98	0.86	0.50	0.10
400-02	0.07	0.52	0.67
404-06	0.84	0.52	0.10
408-10	0.51	0.85	0.05
412-14	0.82	0.56	0.09
416-18	0.92	0.37	0.10
420-22	0.79	0.58	0.08
424-26	0.72	0.57	0.05
428-30	0.62	0.77	0.06
432-34	0.18	0.86	0.07
436-38	0.28	0.95	0.02
440-42	0.84	0.54	0.10
444-46	0.85	0.50	0.10
448-50	0.92	0.37	0.11
452-54	0.97	0.21	0.14
456-58	0.85	0.50	0.10
460-62	0.92	0.37	0.11
464-66	0.97	0.18	0.11
468-70	0.93	0.33	0.11
472-74	0.32	0.81	0.01
476-78	0.93	0.35	0.11
480-82	0.94	0.30	0.11
484-86	0.81	0.57	0.09
488-90	0.97	0.19	0.12
492-94	0.96	0.25	0.12
496-98	0.96	0.25	0.12
500-02	0.91	0.40	0.11
504-06	0.92	0.37	0.11

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
508-10	0.60	0.65	0.04
512-14	0.54	0.79	0.04
516-18	0.97	0.19	0.11
520-22	0.84	0.46	0.07
524-26	0.97	0.18	0.12
528-30	0.97	0.20	0.12
532-34	0.87	0.49	0.10
536-38	0.53	0.84	0.06
540-42	0.93	0.22	0.11
544-46	0.97	0.18	0.12
548-50	0.96	0.24	0.12
552-54	0.93	0.35	0.11
556-58	0.88	0.40	0.10
560-62	0.89	0.43	0.10
564-66	0.94	0.28	0.10
568-70	0.97	0.19	0.12
572-74	0.92	0.37	0.11
576-78	0.79	0.59	0.10
580-82	0.37	0.92	0.03
584-86	0.90	0.42	0.11
588-90	0.92	0.33	0.11
592-94	0.98	0.17	0.12
596-98	0.89	0.44	0.10
600-02	0.73	0.39	0.06
604-06	0.93	0.26	0.10
608-10	0.98	0.18	0.12
612-14	0.95	0.22	0.11
616-18	0.96	0.20	0.11
620-22	0.85	0.51	0.10
624-26	0.43	0.68	0.00
628-30	0.95	0.26	0.10
632-34	0.85	0.48	0.09
636-38	0.90	0.40	0.10
640-42	0.04	0.02	0.34
644-46	0.90	0.41	0.10
648-50	0.68	0.72	0.07
652-54	0.85	0.50	0.09
656-58	0.92	0.37	0.11
660-62	0.91	0.38	0.10
664-66	0.97	0.19	0.12
668-70	0.94	0.29	0.11

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
672-74	0.90	0.43	0.11
676-78	0.92	0.35	0.12
680-82	0.97	0.20	0.12
684-86	0.96	0.24	0.11
688-90	0.96	0.23	0.11
692-94	0.98	0.15	0.12
700-02	0.96	0.22	0.11
704-06	0.94	0.17	0.10
708-10	0.84	0.28	0.08
712-14	0.50	0.47	0.01
716-18	0.36	0.83	0.01
720-22	0.92	0.37	0.11
724-26	0.89	0.41	0.09
728-30	0.98	0.17	0.12
732-34	0.95	0.25	0.11
736-38	0.96	0.19	0.13
740-42	0.96	0.26	0.11
744-46	0.95	0.26	0.11
748-50	0.96	0.26	0.11
752-54	0.87	0.45	0.13
756-58	0.96	0.26	0.12
760-62	0.92	0.36	0.10
764-66	0.91	0.32	0.10
768-70	0.74	0.66	0.08
772-74	0.97	0.19	0.12
776-78	0.93	0.32	0.11
780-82	0.97	0.21	0.11
784-86	0.94	0.22	0.11
788-90	0.98	0.16	0.12
792-94	0.93	0.30	0.10
796-98	0.98	0.16	0.12
800-02	0.85	0.31	0.08
804-06	0.94	0.21	0.11
808-10	0.75	0.24	0.24
812-14	0.27	0.00	0.52
816-18	0.98	0.16	0.12
820-22	0.97	0.16	0.12
824-26	0.49	0.03	0.38
828-30	0.95	0.23	0.10
832-34	0.98	0.16	0.12
836-38	0.98	0.15	0.12

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
840-42	0.93	0.24	0.11
844-46	0.98	0.17	0.12
848-50	0.97	0.20	0.12
852-54	0.98	0.15	0.12
856-58	0.98	0.15	0.12
860-62	0.98	0.17	0.12
864-66	0.97	0.21	0.12
868-70	0.89	0.41	0.10
872-74	0.98	0.15	0.12
876-78	0.96	0.19	0.11
880-82	0.93	0.17	0.10
884-86	0.98	0.15	0.12
888-90	0.98	0.15	0.12
892-94	0.98	0.15	0.12
896-98	0.98	0.15	0.12
900-02	0.98	0.15	0.12
904-06	0.98	0.15	0.13
908-10	0.98	0.16	0.12
912-14	0.98	0.16	0.12
916-18	0.98	0.15	0.12
920-22	0.98	0.15	0.12
924-26	0.90	0.19	0.10
928-30	0.98	0.15	0.12
932-34	0.98	0.16	0.12
936-38	0.98	0.16	0.12
940-42	0.98	0.16	0.12
944-46	0.98	0.15	0.12
948-50	0.98	0.16	0.12
952-54	0.98	0.15	0.12
956-58	0.98	0.15	0.12
960-62	0.98	0.16	0.12
964-66	-	-	-
968-70	0.97	0.16	0.11
972-74	0.97	0.18	0.12
976-78	0.98	0.15	0.12
980-82	0.98	0.16	0.12
984-86	0.98	0.15	0.12
988-90	0.98	0.16	0.12
992-94	0.98	0.16	0.12
996-98	0.93	0.17	0.11
1000-02	0.97	0.16	0.11

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
1004-06	0.97	0.17	0.12
1008-10	0.98	0.17	0.12
1012-14	0.23	0.02	0.19
1016-18	0.71	0.08	0.16
1020-22	0.83	0.18	0.08
1024-26	0.98	0.16	0.12
1028-30	0.98	0.16	0.12
1032-34	0.98	0.15	0.12
1036-38	0.96	0.16	0.11
1040-42	0.93	0.19	0.10
1044-46	0.94	0.14	0.14
1048-50	0.98	0.15	0.12
1052-54	0.98	0.15	0.12
1056-58	0.98	0.16	0.12
1060-62	0.98	0.15	0.13
1064-66	0.98	0.15	0.12
1068-70	0.98	0.15	0.12
1072-74	0.98	0.15	0.12
1076-78	0.98	0.15	0.13
1080-82	0.98	0.15	0.12
1084-86	0.98	0.15	0.12
1088-90	0.98	0.15	0.12
1092-94	0.98	0.15	0.12
1096-98	0.98	0.15	0.12
1100-02	0.98	0.15	0.12
1104-06	0.98	0.15	0.12
1108-10	0.98	0.16	0.12
1112-14	0.98	0.15	0.12
1116-18	0.91	0.12	0.19
1120-22	0.98	0.16	0.12
1124-26	0.98	0.15	0.12
1128-30	0.98	0.16	0.12
1132-34	0.97	0.17	0.13
1136-38	0.98	0.16	0.12
1140-42	0.84	0.10	0.21
1144-46	0.98	0.16	0.12
1148-50	0.97	0.16	0.11
1152-54	0.95	0.14	0.21
1156-58	0.96	0.15	0.20
1160-62	0.91	0.13	0.36
1164-66	0.38	0.04	0.91

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
1168-70	0.74	0.11	0.66
1172-74	0.91	0.14	0.15
1176-78	0.98	0.17	0.12
1180-82	0.98	0.15	0.12
1184-86	0.98	0.16	0.12
1188-90	0.98	0.15	0.12
1192-94	0.98	0.15	0.12
1196-98	0.98	0.15	0.12
1200-02	0.98	0.16	0.12
1204-06	0.98	0.15	0.12
1208-10	0.98	0.15	0.12
1212-14	0.98	0.15	0.12
1216-18	0.98	0.15	0.12
1220-22	0.98	0.15	0.12
1224-26	0.98	0.16	0.12
1228-30	0.98	0.15	0.12
1232-34	0.98	0.15	0.12
1236-38	0.98	0.15	0.12
1240-42	0.98	0.15	0.12
1244-46	0.98	0.15	0.12
1248-50	0.98	0.15	0.12
1252-54	0.98	0.15	0.12
1256-58	0.98	0.15	0.12
1260-62	0.98	0.16	0.12
1264-66	0.98	0.15	0.12
1268-70	0.98	0.15	0.12
1272-74	0.98	0.15	0.12
1276-78	0.98	0.15	0.12
1280-82	0.98	0.15	0.12
1284-86	0.98	0.15	0.12
1288-90	0.98	0.16	0.12
1292-94	0.98	0.15	0.12
1296-98	0.98	0.15	0.12
1300-02	0.98	0.15	0.12
1304-06	0.98	0.16	0.12
1308-10	0.98	0.15	0.12
1312-14	0.98	0.15	0.12
1316-18	0.98	0.16	0.12
1320-22	0.98	0.15	0.12
1324-26	0.98	0.15	0.12
1328-30	0.98	0.15	0.12

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
1332-34	0.98	0.15	0.12
1336-38	0.98	0.15	0.12
1340-42	0.98	0.15	0.12
1344-46	0.98	0.16	0.12
1348-50	0.97	0.14	0.20
1352-54	0.98	0.15	0.12
1356-58	0.95	0.17	0.11
1360-62	0.96	0.17	0.11
1364-66	0.98	0.16	0.12
1368-70	0.88	0.18	0.10
1372-74	0.96	0.17	0.11
1376-78	0.98	0.15	0.12
1380-82	0.96	0.17	0.11
1384-86	0.80	0.18	0.08
1388-90	0.98	0.15	0.13
1392-94	0.86	0.18	0.11
1396-98	0.93	0.17	0.11
1400-02	0.98	0.15	0.12
1404-06	0.98	0.16	0.12
1408-10	0.98	0.15	0.12
1412-14	0.95	0.17	0.11
1416-18	0.96	0.15	0.24
1420-22	0.85	0.14	0.48
1424-26	0.96	0.16	0.23
1428-30	0.98	0.15	0.12
1432-34	0.89	0.12	0.16
1436-38	0.68	0.07	0.38
1440-42	0.75	0.09	0.38
1444-46	0.98	0.15	0.12
1448-50	0.98	0.15	0.12
1452-54	0.95	0.17	0.11
1456-58	0.98	0.15	0.12
1460-62	0.98	0.15	0.13
1464-66	0.96	0.20	0.11
1468-70	0.98	0.15	0.12
1472-74	0.98	0.15	0.12
1476-78	0.98	0.16	0.12
1480-82	0.98	0.15	0.12
1484-86	0.98	0.15	0.12
1488-90	0.98	0.15	0.12
1492-94	0.97	0.16	0.12

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
1496-98	0.98	0.15	0.12
1500-02	0.98	0.15	0.12
1504-06	0.98	0.15	0.12
1508-10	0.98	0.15	0.12
1512-14	0.98	0.15	0.12
1516-18	0.98	0.15	0.12
1520-22	0.98	0.15	0.12
1524-26	0.98	0.15	0.12
1528-30	0.98	0.15	0.12
1532-34	0.98	0.15	0.12
1536-38	0.98	0.15	0.12
1540-42	0.98	0.15	0.12
1544-46	0.95	0.16	0.16
1548-50	0.98	0.15	0.12
1552-54	0.98	0.15	0.12
1556-58	0.98	0.15	0.12
1560-62	0.98	0.16	0.12
1564-66	0.98	0.15	0.12
1568-70	0.98	0.15	0.12
1572-74	0.98	0.15	0.12
1576-78	0.98	0.15	0.12
1580-82	0.98	0.15	0.12
1584-86	0.98	0.15	0.12
1588-90	0.98	0.15	0.12
1592-94	0.98	0.15	0.12
1596-98	0.98	0.15	0.12
1600-02	0.98	0.15	0.12
1604-06	0.95	0.16	0.14
1608-10	0.33	0.02	0.93
1612-14	0.38	0.03	0.92
1616-18	0.09	0.01	0.99
1620-22	0.03	0.02	0.99
1624-26	0.05	0.02	0.98
1628-30	0.01	0.02	0.99
1632-34	0.02	0.04	0.96
1636-38	0.98	0.15	0.12
1640-42	0.09	0.07	0.67
1644-46	0.86	0.17	0.10
1648-50	0.09	0.06	0.91
1652-54	0.78	0.11	0.49
1656-58	0.28	0.00	0.88

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
1660-62	0.05	0.05	0.96
1664-66	0.06	0.04	0.97
1668-70	0.07	0.04	0.65
1672-74	0.15	0.01	0.88
1676-78	0.35	0.02	0.84
1680-82	0.24	0.03	0.95
1684-86	0.19	0.01	0.98
1688-90	0.03	0.02	1.00
1692-94	0.04	0.05	0.48
1696-98	0.07	0.01	0.57
1700-02	0.07	0.03	0.78
1704-06	0.94	0.14	0.31
1708-10	0.56	0.06	0.77
1712-14	0.56	0.04	0.24
1716-18	0.04	0.06	0.21
1720-22	0.05	0.03	0.78
1724-26	0.04	0.07	0.68
1728-30	0.36	0.01	0.81
1732-34	0.13	0.03	0.82
1736-38	0.74	0.09	0.56
1740-42	0.63	0.06	0.56
1744-46	0.23	0.00	0.95
1748-50	0.45	0.03	0.82
1752-54	0.78	0.10	0.52
1756-58	0.71	0.09	0.69
1760-62	0.66	0.08	0.74
1764-766	0.79	0.11	0.59
1768-770	0.66	0.09	0.74
1772-74	0.40	0.03	0.90
1776-78	0.78	0.09	0.49
1780-82	0.86	0.12	0.41
1784-86	0.91	0.13	0.36
1788-90	0.79	0.09	0.47
1792-94	0.91	0.13	0.38
1796-98	0.95	0.14	0.28
1800-02	0.95	0.14	0.29
1804-06	0.91	0.13	0.36
1808-10	0.94	0.14	0.24
1812-14	0.97	0.14	0.19
1816-18	0.95	0.14	0.29
1820-22	0.96	0.14	0.23

Distribution of Foraminiferal Data for NBP00-03-JPC-38

Core depth (cm)	PC1 <i>M. arenacea</i>	PC2 <i>T. wiesneri</i>	PC3 <i>S. fusiformis</i>
1824-26	0.94	0.15	0.29
1828-30	0.88	0.12	0.46
1832-34	0.91	0.13	0.39
1836-38	0.53	0.06	0.84
1840-42	0.57	0.07	0.81
1844-46	0.78	0.10	0.61
1848-50	0.82	0.10	0.49
1852-54	0.89	0.12	0.39
1856-58	0.62	0.08	0.78
1860-62	0.95	0.14	0.28
1864-66	0.97	0.15	0.19
1868-70	0.96	0.14	0.22
1872-74	0.95	0.14	0.26
1876-78	-	-	-
1880-82	0.65	0.08	0.75
1884-86	0.96	0.14	0.22
1888-90	0.59	0.06	0.65
1892-94	0.60	0.06	0.59
1896-98	0.76	0.10	0.63
1900-02	0.59	0.07	0.79
1904-06	0.37	0.02	0.88
1908-10	0.43	0.04	0.90
1912-14	0.90	0.13	0.41
1916-18	0.96	0.14	0.25
1920-22	0.90	0.13	0.41
1924-26	0.78	0.11	0.61
1928-30	0.95	0.14	0.25
1932-34	0.93	0.13	0.31
1936-38	0.95	0.14	0.24
1940-42	0.76	0.10	0.64
1944-46	0.22	0.01	0.97
1948-50	0.02	0.01	0.98
1952-54	0.87	0.11	0.35
1956-58	0.95	0.13	0.11
1960-62	0.97	0.15	0.18
1964-66	0.09	0.04	0.98
1968-70	0.64	0.08	0.74
1972-74	0.01	0.04	0.68
1976-78	0.33	0.01	0.79
1980-82	0.05	0.01	0.98
1984-86	0.13	0.00	0.98