Effects of Mining on the Environment — A National Overview

S.E. Church, T.S. Schmidt,
C.A. San Juan, and G.S. Plumlee

NRDAR Conference, Phoenix, AZ

April 25, 2007

Two of the most important points of this talk are:

- Knowing the geology of the mine site is critical to understanding the environmental impact of mining at any given site
- Hydrothermal alteration associated with the mineral deposit has a major effect on the premining geochemical background as well as on the postmining environmental impact on the watershed

What is a mineral deposit?

An area within the Earth's crust where geologic processes have concentrated one or more mineral commodities

What is an ore deposit?

A mineral deposit in which the commodities (i.e., metals) are present in sufficient concentration (grade) and amount (tonnage) that they can (ideally) be extracted economically

Metallic mineral deposits

- Concentrations of metallic elements and minerals in the Earth's crust
- Form by:
 - Crystallization of magmas
 - Cooling, boiling, mixing of hydrothermal fluids in the Earth's crust
 - Chemical precipitation of minerals on the sea floor or in sediment
 - Chemical reaction of two or more different fluids
 - Chemical reaction of a hydrothermal fluid with a rock

Mineral deposit models

- Mineral deposits classified by geologic characteristics and geologic environment of formation
- Mineral deposit models developed for more than 40 deposit types
- Components:
 - Descriptive (general geologic characteristics)
 - Genetic (how they form)
 - Grade and tonnage
 - Hydrothermal alteration

Hydrothermal alteration

- Acid-sulfate alteration
- Potassic alteration (ore zone)
- Quartz-sericite-pyrite (QSP)
- Argillic (clay minerals)
- Propylitic

Geoenvironmental mineral deposit models

- Use geologic characteristics of mineral deposits
 - Sulfide mineralogy
 - Hydrothermal alteration
 - Ore-deposit mineralogy
- Time frame of historical mining activity
 - Metal mobility from mine-waste dumps
 - Water chemistry
 - Smelter signatures in soils
 - Sediment geochemistry in surface streams

Take-home message

- A mineral deposit may not be an ore deposit. The term “ore” is an economic descriptor
- Physical processes of formation of the deposit are the primary drivers controlling the impact of the mined site on the environment
- Hydrothermal alteration controls the distribution of pyrite in many mineral deposits

Potential environmental and health impacts of mining

A complex **but predictable** function of:

- Mineral deposit geology
 - Environmental signatures, physical footprint
- The mining and mineral processing methods used
 - Typically mandated by the deposit geology
- Geochemical and biogeochemical processes
 - Bacterial activity can increase reaction rates by a million times
- The effect of climate on rates of weathering

Effective prediction, mitigation, and remediation strategies are also dependent upon characteristics of the surrounding watersheds

- Geology, size, topography, ecology
- Number and footprint of other environmental stressors

What data are available to indicate the extent of land disturbance throughout the Nation?

- USGS database of historical mines provides a representative look at disturbance by mining

Mines, prospects, and occurrences

(USGS MAS/MILS data base)

What data are in the USGS MAS/MILS databases?

- Four types of data that show the extent of land disturbance
 - Mines
 - Active or
 - Inactive
 - These data are current as of the time of data entry
 - Prospects—minimal ground disturbance
 - Occurrences—no ground disturbance

Significant mineral deposits

- Cu, Pb, Zn, Au, Ag (Long et al., 2002)
 - Defined about 1,000 major ore deposits by deposit type
 - Upper 10 % of producers by commodity (Singer, 1995)
- Uranium
- Phosphate
- Iron
- Molybdenum
- Mercury

Significant gold deposits

Porphyry copper and molybdenum deposits

Bingham, Utah

Grasberg, Indonesia

Marcopper, Philippines

Schematic acid-drainage potential of a porphyry copper deposit

Modified from Park and Guilbert, 1986

Near-neutral waters, some metals

Development of pit lakes

- Fracture flow hydrology, pit wall geology, lake limnology, climate are key parameters

Significant porphyry deposits

Porphyry deposits represent the biggest environmental challenge that Federal Land Management Agencies face in terms of acid mine drainage (AMD)

- Many porphyry deposits are still being mined
- Most porphyry deposits occur in arid regions and are on private land
- Most deposits that have been mined out continue to be held by mining companies and are listed as inactive mines
- In the future these will be the major AMD issues FLMA will face

Epithermal vein and Polymetallic vein deposits

Significant vein deposits

Red Mountains, Silverton, Colo.

Footprint of hydrothermal alteration

Animas River watershed study area

<http://amli.usgs.gov>

Figure 8.

Formation of Al sulfate precipitates

Paradise portal, Animas River watershed

- Dilution by fresh ground water triggers Al precipitation

VMS deposits

- Form on the sea floor
- Very pyritic
- Little alteration
- Buried by fresh volcanic rock or by sediment to be preserved
- Very large acid generators

Significant volcanogenic massive sulfide deposits

What controls water chemistry in AMD?

- The amount of acid-generating minerals
 - Pyrite isn't the only one!
- The amount and type of hydrothermal alteration
- The acid-neutralization potential in the watershed
- The amount of water available for dilution of AMD
- Climate

Acid-generating minerals

PYRITE (FeS₂):

- By oxygen:

- By ferric iron:

- Bacteria speed up the key acid-drainage formation reaction (b) by 1 million times
- Dissolved ferric iron is a corrosive oxidant

Acid-generating minerals

Pyrite (FeS_2)

Pyrrhotite (Fe_{1-x}S)

Enargite (Cu_3AsS_4)

Realgar (As_2S_3)

Others

Marcasite (FeS_2)

Arsenopyrite (FeAsS)

Tennantite ($\text{Cu}_{12}\text{As}_4\text{S}_{13}$)

Orpiment (AsS)

If ferric iron is oxidant, above minerals plus:

Chalcopyrite (CuFeS_2)
(ZnS)

Covellite (CuS)

Sphalerite

Chalcocite (Cu_2S)

Acanthite (Ag_2S)

Galena (PbS)

If metal hydroxides (solid or aqueous) form, above minerals plus:

Siderite (FeCO_3)

Rhodochrosite (MnCO_3)

Comparison of water chemistry from “natural” and AMD sites

Differences in rate of some types of reactions that influence metal mobility

Temporal Scale, as approximate half-times of reaction

seconds minutes hours days months years

solute-solute

gas-water

solute-water

solubility

adsorption-desorption

oxidation-reduction

solid-water

mineral
recrystallization

Precipitation of iron minerals

Pyrite weathering

Geological and geochemical controls on mine- and natural-drainage water compositions

Sampling water from the Reynolds adit

Seep, 8/95

- pH 3.1
- Conductivity 8,900 μS
- Cu 825 mg/L
- Fe 2,000 mg/L

Geologic controls on mine-drainage composition

Symbols depict waters from deposits with similar geologic characteristics

Plumlee et al.,
1999, (all
samples
filtered)

Colloidal Fe in the stream

The effect of iron on trace-element concentrations in water

Preferentially removing iron from a system may result in increased dissolved concentrations of other metals

Mixing zone, Snake River drainage, Colo.

Example of changes in precipitation of colloids as a function of pH in a mixing zone downstream of the confluence of an acidic stream (undisturbed by historical mining) and a neutral stream

Acid-neutralizing minerals

- Carbonate minerals and some other minerals (some silicates, volcanic glasses) in mineral deposits, their host rocks, and watershed rocks:
 - Can react with and help consume acid generated by sulfide oxidation
 - Can also generate alkalinity in ground and surface waters, thereby increasing the waters' ability to buffer acid

Limestone

http://www.sirendesigns.com/elora_resources/gallery_1/pages/limestone_cliff_gif.htm

Calcite on
pyrite,
Silesia,
Poland

What geochemical data are available that would show where potentially toxic mine sites might be?

- In the conterminous U.S.
 - 330,000 analyzed NURE sediments
 - 150,000 analyzed USGS sediments
- In Alaska
 - 65,000 analyzed NURE sediments
 - 25,000 analyzed USGS sediments

Abandoned Mines (MAS/MILS)

Hazard quotient Copper – Obs. Conc. / PEC 149 ppm)

Hazard quotient – Cadmium (Obs. Conc. / PEC 4.98 ppm)

Hazard quotient – Zinc (Obs. Conc. / PEC 459 ppm)

<http://mrdata.usgs.gov/website/MRData-US/viewer.htm>

U.S. Geological Survey - MRData.USGS.gov

Active Mines and Plants

Rec	Commodity	Name	Owner/Operator	State	County	Latitude	Longitude	Mine or Plant
1	Sand and Gravel	Huber Sand Pit #2	Huber Sand Co.	Kansas	Finney	37.94778	-100.8448	M/P
2	Sand and Gravel	Pit #2	Smith Sand Co., Inc.	Kansas	Finney	37.98383	-100.9057	M/P
3	Sand and Gravel	Site #3 - Holcomb	Klotz Sand Co., Inc.	Kansas	Finney	37.97828	-100.9971	M/P
4	Sand and Gravel	Kendall #1	Klotz Sand Co., Inc.	Kansas	Finney	37.9434	-100.7715	M/P

Refresh Map

Map Layers

☐ Mineral Resources

Visible Active

- ☐ MRDS, producers
- ☐ MRDS, all
- ☒ Active Mines and Plants
- ☐ Mine claim activity 1976-2004

☐ Geochemistry

Visible Active

- ☐ US Geochemical Survey
- ☐ Igneous Rock, Pluto
- ☐ Sediment, Pluto
- ☐ Sediment, RASS
- ☐ Soil, Pluto
- ☐ Soil, RASS
- ☐ Soil, Shacklette

☒ Geophysics☒ Geology☒ Hydrology☐ Boundaries

Visible Active

Central Colorado Assessment Project

Central Colorado Assessment Project

- Provide an environmental assessment of the central 1/3 of the State of Colorado
- Develop geochemical and biological baseline data
 - Sampled watersheds underlain by single geologic units to provide a geologic signature
 - Sampled watersheds underlain by hydrothermally altered rock
 - Sampled watersheds in which historical mining had occurred

Ephemeroptera

Plecoptera

Trichoptera

Diversity and Abundance

[Trace Metals]

Chronic Criterion Accumulation Ratio (CCAR)

$$\sum \frac{\text{Site-specific [gill-metal]}}{\text{US EPA standard water [gill-metal] @ CCC}}$$

- **New model derived from
Toxic Unit Model**
 - **BLM derived**
(Sprague, 1970; Playle, 2004)

Development of biological baselines

- **Develop a model of bioavailability**
 - **Determine that lithologic classes can differentiate bioavailability**
 - **Validate that model can predict indigenous taxa responses**
 - **Also develop bio-indicator species for assessment purposes**
 - **Infer adverse effect concentrations**

Can lithologic classes differentiate bioavailability?

Can hydrothermal alteration differentiate bioavailability?

Are water-quality criteria protective?

- **Water Quality Criteria**
 - Protect 95% of all taxa
 - Genus-level mean responses
- **30% loss in richness**
- **70% loss in abundance**

Appropriate conservative benchmarks are not applicable for all lithologies

Summary

- **Lithology and hydrothermal alteration control bioavailability of metals**
- **Not all BMI endpoints are appropriate**
- **CCAR is more descriptive than CCU**
- **Are Water Quality Criteria protective endpoint?**
- **Are remedial actions based on conservative benchmark concentrations appropriate?**

Colloidal Fe in the stream

Natural?