

### HIGH RESOLUTION GEOLOGIC MAPPING OF THE INNER CONTINENTAL SHELF: CAPE ANN TO SALISBURY BEACH, MASSACHUSETTS

U.S. Geological Survey Open File Report 2007-1373  
Map Sheet 1: Shaded-relief topography of seafloor (color)

**Introduction**  
A series of five map sheets shows the seafloor topography and geology of the inner continental shelf between Cape Ann and Salisbury Beach, Massachusetts. This map sheet (1) shows seafloor topography in shaded-relief view and colored by water depth with selected bathymetric contours overlain in black. Sheet 2 shows shaded-relief topography in gray scale, sheet 3 shows gray-scale backscatter intensity, sheet 4 shows shaded-relief topography colored by backscatter intensity, and sheet 5 shows seafloor geology. These maps were produced as part of a cooperative effort by the U.S. Geological Survey (USGS) and the Massachusetts Office of Coastal Zone Management (CZM) to systematically map the seafloor geology offshore of Massachusetts. This map sheet is accompanied by a more extensive report available on the Internet and a DVD-ROM that presents a full description of the data-collection, processing, and analysis procedures used to create the maps. The report also includes copies of selected data layers in a Geographic Information System (GIS).

The shaded-relief imagery in this map was created by artificially illuminating the bathymetry by a light source at an azimuth of 335° and from an altitude of 45° above the horizon. Topographic features, such as submarine ridges and hills, are enhanced by strong illumination on northeast-facing slopes and by shadows cast on southeast-facing slopes. The hillshading accentuates small features that are not effectively illustrated by contours alone at this scale. Bathymetric contours are shown at 10-m intervals and, for clarity, have been simplified or omitted in areas of complex topography.

Additional data are included on this map to show the regional topography in areas adjacent to the area of the new survey. To the north and west, seafloor topography in shaded-relief view is shown at a reduced resolution of 90 m/pixel from the data published by the National Geophysical Data Center (NGDC) Coastal Relief Model (Dunn and Metzger, 2007). Hillshaded bathymetry to the east and southeast from Butman and others (2007) is displayed at a resolution of 10 m/pixel. Onshore topography displayed at a resolution of 5 m/pixel is from the Massachusetts Geographic Information System (MassGIS, 2005). Some features in the image are artifacts of data collection or environmental conditions. They include small highs and lows and unnatural-looking patterns oriented parallel or perpendicular to survey tracklines, which were oriented northeast-southwest in the map area.

**Data and Methods**  
Approximately 325 km<sup>2</sup> of the inner shelf in water depths of 2–92 m were mapped on two separate cruises in 2004 and 2005. Several different geophysical systems were used, including multibeam and interferometric sonars (bathymetry), sidescan and multibeam sonars (acoustic backscatter intensity), and chirp sonar reflection profiles (subsurface stratigraphy and structure).

In deeper offshore areas, bathymetric data on this map were collected by a Reson 8101 multibeam echo sounder (MBES) operating at a frequency of 240 kHz (SAC, 2004). The system has a 101-beam transducer that collects depth data in a continuous swath on either side of the vessel. Adjacent swaths were overlapped to achieve 100 percent coverage of the seafloor. Depth data were processed with SABER software developed by Science Applications International Corporation (SAIC). The bathymetric data have a vertical resolution of approximately 0.5 percent of water depth and a grid cell size of 5 m. Tidal offsets, calculated by using a discrete tidal-zoning model and observations from National Oceanic and Atmospheric Administration (NOAA) Tidal Station #8418150 at Portland, Maine (NOA CO-QRS, 2005), were used to reference the soundings to the local mean lower low water datum.

In shallower nearshore areas, bathymetric data were collected by a SEA SwathPlus 2000 series interferometric sonar that operates at a frequency of 234 kHz. The system has two transducers that collect depth data in a continuous swath on either side of the vessel. Adjacent swaths were overlapped to obtain 100 percent coverage of the seafloor. Data were processed and gridded with the CARIS Hydrographic Information Processing System (HIPS) ver. 6.11. Minor gaps were filled by creating a Triangulated Irregular Network (TIN). The bathymetric data have a vertical resolution of approximately 1 percent of water depth and a grid cell size of 5 m. Navigation was provided by a Real-Time Kinematic Global Positioning System (RTK-GPS) from a base station established onshore. Tidal offsets were calculated by using RTK-GPS onshore elevations (GEOID 03, NAVD83) and were applied to soundings during post processing. Soundings were referenced to local mean lower low water using an ellipsoid to chart datum offsets obtained from NOAA Tidal Station #8440452 located at the Parker River National Wildlife Refuge.

Bottom photography and sediment sampling at 87 stations were used to validate interpretations of the remotely-sensed geophysical data. A Seafloor Observation and Sampling System (SEAMOS; Valentine and others, 2000) recorded continuous video and still photographs along drift tracks that typically lasted 5–15 minutes at each station. Sediment samples were collected at locations where the bottom was not covered with boulders or ledge and analyzed for grain size by the methods outlined in Pappas and others (2005).

**Features**  
Maps depicting topography and surficial materials on the inner continental shelf play an important role in representing the geologic history of the region and the processes that have shaped the seafloor. Igneous and metamorphic rocks spanning millions of years of Earth history control the overall geometry of the coast and inner continental shelf (Zan and others, 1983). Erosion-resistant granitic rocks form rugged headlands and submarine shoals that define the northern and southern margins of the study area. Glaciation and relative sea-level change are the most important processes to have acted on the region and have produced a heterogeneous mix of bottom types on the inner continental shelf. Late Pleistocene glaciers reached their maximum extent south of Cape Cod about 21,000 years before present (yr B.P.) and retreated northward as the climate warmed. The retreating ice margin passed the present coast of Cape Ann about 14,500 yr B.P. (Kaye and Barghorn, 1964). The glaciers smoothly eroded the underlying bedrock and largely buried it with boulders, till and muddy glacial marine sediment (Stone and others, 2006). Thick deposits of coarser grained fluvial, deltaic, and littoral sediment accumulated above the glacial deposits and masked the underlying topography beneath a generally smooth, seaward-dipping seafloor. These younger sediment accumulations represent the variety of different environments that existed on the inner shelf after the end of the last Ice Age.

Large excursions in relative sea level, primarily caused by a combination of global climate change and local isostatic effects, have driven profound changes in the position of the coastline and strongly influenced the geologic evolution of the region. Relative sea level fell from a highstand of +33 m at 14,500 yr B.P. to a lowstand of about -50 m at 12,000 yr B.P. (Osone and Piper, 1982; Oldale and others, 1993). An isostatic rebound decreased relative sea level rise at different rates to the present elevation. The ongoing Holocene transgression (i.e., landward migration of the coast) has reworked a wide expanse of the inner shelf, and eroded a prominent unconformity overlain by relatively thin, discontinuous deposits of mobile sand. Large mounds of boulder-strewn clasts represent the remnants of ice-contact deposits such as drumlins and moraines, which have been eroded by marine and terrestrial processes. In water depths greater than about 50 m, below the sea level lowstand, the generally smooth seafloor is dominated by fine-grained marine mud interrupted by a few widely separated outcrops of ledge and boulders. Scoured moats around these deep-water outcrops indicate persistent erosion by bottom currents.

The Merrimack River, which drains a large area of Massachusetts and New Hampshire, enters the Gulf of Maine just north of Cape Ann. Sediment supplied by the river and derived from erosion of coastal and inner-shelf deposits has built the longest chain of sandy barriers in the Gulf of Maine (FitzGerald and others, 1994). The barrier islands and spits are pinned to bedrock or glacial promontories along the coast and stretch approximately 30 km from Ipswich Bay to the New Hampshire border.

**Acknowledgments**  
This series of seafloor maps is dedicated to the late Susan Snow-Cotter (1961–2006), who for many years was the leader in efforts to balance environmental policies with human uses of ocean resources in Massachusetts. She was instrumental in establishing and maintaining the cooperative seafloor-mapping program, which is funded by the Coastal and Marine Geology Program of USGS and the CZM. We wish to thank Anthony Wilbur and Daniel Sampson of CZM for their encouragement and support of marine research. Assistance in the field was provided by Emily Bergstrom, William Danforth, Barry Irwin, and Charles Worley of the USGS. We also thank Jane Denny, William Danforth, and David Foster of USGS for their help in processing the large amounts of acoustic data. The maps benefited from reviews by Jane Denny, S. Jeffrey Williams, Matthew Arsenault, Anthony Wilbur, and Daniel Sampson.

**References Cited**  
Ackerman, S., Butman, B., Barnhardt, W., Danforth, W., and Crocker, J., 2006. High-resolution geologic mapping of the inner continental shelf, Boston Harbor and Approaches, Massachusetts. U.S. Geological Survey Open File Report 2006-1008. Available on DVD and online at <http://pubs.usgs.gov/of/2006/1008/>

Barnhardt, W.A., Andrews, B.D., and Butman, B., 2006. High-resolution geologic mapping of the inner continental shelf, Nahant to Gloucester, Massachusetts. U.S. Geological Survey Open File Report 2005-1293. Available on DVD and online at <http://pubs.usgs.gov/of/2005/1293/>

Butman, Bradford, Valentine, P.C., Middleton, T.J., and Danforth, W.W., 2007. A GIS Library of multibeam data for Massachusetts Bay and the Stellwagen Bank National Marine Sanctuary offshore of Boston, Massachusetts. U.S. Geological Survey Data Series 99, DVD-ROM. Available online at <http://pubs.usgs.gov/ds/99/>

Dunn, D.L., and D. Metzger, 2007. NGDC Coastal Relief Model, accessed June 15, 2007 at <http://www.ngdc.noaa.gov/mgg/coastal/coastal.html>

FitzGerald, D.M., Rosen, P.S., and van Heteren, S., 1994. New England barriers, in: Davis, R.A., ed., *Geology of holocene barrier island systems*. Springer-Verlag, Berlin, Germany, p. 305-394.

Kaye, C.A. and Barghorn, E.S., 1964. Late Quaternary sea-level change and crustal rise at Boston, Massachusetts, with a note on the autocompaction of peat. *Geological Society of America Bulletin*, v. 75, p. 63-80.

Massachusetts Geographic Information System (MassGIS) 2005. Statewide digital elevation model (1:50,000) February 2005. Available online at [http://www.mass.gov/eng/mgis/eng\\_elev3d.html](http://www.mass.gov/eng/mgis/eng_elev3d.html)

National Oceanic and Atmospheric Administration, National Ocean Service, Center for Operational Oceanographic Products and Services (NOA CO-QRS), 2005. accessed October 20, 2005 at <http://oeps.noaa.gov/benchmarks/8418150.html>

Oldale, R.N., Colman, S.M., and Jones, G.A., 1993. Radiocarbon ages from two submerged strandline features in the western Gulf of Maine and a sea-level curve for the northeastern Massachusetts coastal region. *Quaternary Research*, v. 40, p. 38-45.

Pappas, L., Williams, S.J., and Paskovitch, V.F., eds., 2005. USGS East-Coast Sediment Analysis Procedures, Database, and GIS Data. U.S. Geological Survey Open File Report 2005-1001, supercedes 2000-358 DVD ROM. Available online <http://pubs.usgs.gov/of/2005/1001/>


Science Applications International Corporation (SAIC), 2004. Southern Merrimack embayment multibeam survey: Survey Report. Prepared for U.S. Geological Survey by SAIC, Newport, RI.

Stone, B.D. and Piper, J.D., 1982. Topographic control of depauperation of eastern Massachusetts ice lobe and marine incursion, in: B.J. Larson and B.D. Stone, eds. *Late Wisconsinan glaciation of New England: Kendall/Hunt*, Dubuque, Iowa, p. 145-166.

Stone, B.D., Stone, J.R., and DiGiacomo-Cohen, M.L., 2006. Surficial geologic map of the Salem-Newburyport East-Wilmington-Rockport 16 Quadrangle Area in Northeast Massachusetts. U.S. Geological Survey Open File Report 2006-1260-B. Available online at <http://pubs.usgs.gov/of/2006/1260/B/>

Valentine, P.C., Blackwood, D.B., and Parobski, K.F., 2000. Seabed observation and sampling system. U.S. Geological Survey Fact Sheet FS-142-00. Available online at <http://pubs.usgs.gov/fs/fs142-00/fs142-00.pdf>

Zen, Evan, Goldsmith, Richard, Ratcliffe, N.M., Robinson, Peter, and Stanley, R.S., 1983. Bedrock geologic map of Massachusetts. U.S. Geological Survey, Washington D.C., scale 1:250,000, 3 sheets.



### HIGH-RESOLUTION GEOLOGIC MAPPING OF THE INNER CONTINENTAL SHELF: CAPE ANN TO SALISBURY BEACH, MASSACHUSETTS

Sheet 1. Shaded-relief topography of seafloor (color)

By  
Walter A. Barnhardt, Brian D. Andrews, Seth D. Ackerman, Wayne E. Baldwin, and Christopher J. Hein

