

HIGH RESOLUTION GEOLOGIC MAPPING OF THE INNER CONTINENTAL SHELF: CAPE ANN TO SALISBURY BEACH, MASSACHUSETTS

Map Sheet 3. Backscatter intensity of the seafloor (grayscale)

Introduction
A series of five map sheets shows the seafloor topography and geology of the inner continental shelf between Cape Ann and Salisbury Beach, Massachusetts. This map (sheet 3) shows acoustic backscatter intensity in grayscale derived from sidescan-sonar imagery, and with selected topographic contours overlain in blue. Sheet 1 shows shaded-relief topography in color, sheet 2 shows shaded-relief topography in grayscale, sheet 4 shows shaded-relief topography colored by backscatter intensity, and sheet 5 shows seafloor geology. These maps are produced as part of a cooperative effort by the U.S. Geological Survey (USGS) and the Massachusetts Office of Coastal Zone Management (CZM) to systematically map the seafloor geology offshore of Massachusetts. This map sheet is accompanied by a more extensive report available on the Internet and a DVD-ROM that presents a full description of the data-collection, processing, and analysis procedures used to create the maps. The report also includes copies of selected data layers in a Geographic Information System (GIS).

The intensity of acoustic backscatter shown on this map is represented by 256 shades of gray, ranging from lighter shades (high-backscatter values) to darker shades (low-backscatter values). High-backscatter values indicate that the seafloor is generally composed of coarse sand, gravel, cobbles, boulders, and rock. In contrast, moderate to low-backscatter values generally indicate a bottom composed of finer grained mixtures of sand and mud. Relief on the seafloor also influences backscatter intensity. For example, a sloping surface that faces the sound source (leeward) produces higher backscatter values than a surface that slopes away from the source. High-relief rocky areas may exhibit shadows, indicating that no acoustic signal is returned. Accurate interpretation of substrate properties from backscatter data requires validation by sediment sampling, bottom photography, and video.

Additional data are included on this map to show the regional topography in areas adjacent to the area of the new survey. To the north and west, seafloor topography in shaded-relief view is shown at a reduced resolution of 90 m/pixel from the data published by the National Geophysical Data Center (NGDC) Coastal Relief Model (Drons and Metzger, 2007). Hillsloped bathymetry to the east and southeast from Butman and others (2007) is displayed at a resolution of 10 m/pixel. Onshore topography displayed at a resolution of 5 m/pixel is from the Massachusetts Geographic Information System (MassGIS, 2005). Some features in the image are artifacts of data collection and environmental conditions. They include narrow stripes parallel to survey tracks, which were oriented northwest-southeast in the map area, and slight mismatches in grayscale tones between adjacent swaths.

Data and Methods
Approximately 325 km² of the inner shelf in water depths of 2–92 m were mapped on two separate cruises in 2004 and 2005. Several different geophysical systems were used, including multibeam and interferometric swath (bathymetry), sidescan and multibeam sonars (acoustic backscatter intensity), and chirp seismic-reflection profilers (subsurface stratigraphy and structure).

In deeper offshore areas, bathymetric data on this map were collected by a Reson 8101 multibeam echo sounder (MBES) operating at a frequency of 240 kHz (SAIC, 2004). The system has a 101-beam transducer that collects depth data in a continuous swath on either side of the vessel. Adjacent swaths were overlapped to achieve 100 percent coverage of the seafloor.

In shallower nearshore areas, acoustic backscatter data were collected with a Klein 3000 sidescan sonar that operates at dual frequencies of 132 and 445 kHz. The system has two transducers that image a continuous swath on either side of the vessel. Adjacent swaths were overlapped to obtain 100 percent coverage of the seafloor. Sidescan-sonar data were corrected for beam angle and slant-range distortions by using Xsonar/Showimage as described in Danforth (1997), and mosaicked using PCI Geomatics GPC works (PCI Geomatics ver 8.2). The 100-kHz data were used for the final mosaic, which is presented here with a pixel size of 1 m. Navigation for the backscatter data used signal from U.S. Coast Guard Differential Global Positioning System (DGPS).

Bottom photography and sediment sampling at 87 stations were used to validate interpretations of the remotely-sensed geophysical data. A SEABed Observation and Sampling System (SEABOSS; Valentine and others, 2005) recorded continuous video and still photographs along drift tracks that typically lasted 5–15 minutes at each station. Sediment samples were collected at locations where the bottom was not covered with boulders or ledge and analyzed for grain size by the methods outlined in Poppe and others (2006).

Features
Maps depicting topography and surficial materials on the inner continental shelf play an important role in representing the geologic history of the region and the processes that have shaped the seafloor, igneous and metamorphic rocks spanning millions of years of Earth history control the overall geometry of the coast and inner continental shelf (Zen and others 1983). Erosion-resistant granitic rocks form rugged headlands and submarine shoals that define the northern and southern margins of the study area. Glaciation and relative sea-level change are the most important processes to have acted on the region and have produced a heterogeneous mix of bottom types on the inner continental shelf. Late Pleistocene glaciers reached their maximum extent south of Cape Cod about 21,000 years before present (ys B.P.) and retreated northward as the climate warmed. The retreating ice margin passed the present coast of Cape Ann about 14,500 ys B.P. (Kaye and Barghoorn, 1964). The glaciers smoothly eroded the underlying bedrock and largely buried it with fine-grained till and muddy glacial-marine sediment (Stone and others, 2006). Thick deposits of coarser grained fluvial, deltaic, and littoral sediment accumulated above the glacial deposits and masked the underlying topography beneath a generally smooth, seaward-dipping seafloor. These younger sediment accumulations represent the variety of different environments that existed on the inner shelf after the end of the last Ice Age.

Large excursions in relative sea level, primarily caused by a combination of global climate change and local isostatic effects, have driven profound changes in the position of the coastline and strongly influenced the geologic evolution of the region. Relative sea level fell from a highstand of +33 m at 14,500 ys B.P. to a lowstand of about -50 m at 12,000 ys B.P. (Stone and Poppe, 1982; Oldale and others, 1994). As isostatic rebound decreased relative sea level rose at different rates to the present elevation. The ongoing Holocene transgression (i.e. landward migration of the coast) has revealed a wide expanse of the inner shelf, and eroded a prominent unconformity overlain by relatively thin, discontinuous deposits of mobile sand and silt. Boulder-strewn clasts represent the remnants of ice-contact deposits such as drumlins and moraines, which have been eroded by marine and terrestrial processes. In water depths greater than about 50 m, below the sea-level lowstand, the generally smooth seafloor is dominated by fine-grained marine mud interrupted by a few widely separated outcrops of ledge and boulders. Scoured moats around these deep-water outcrops indicate persistent erosion by bottom currents.

The Merrimack River, which drains a large area of Massachusetts and New Hampshire, enters the Gulf of Maine just north of Cape Ann. Sediment supplied by the river and derived from erosion of coastal and inner-shelf deposits has built the longest chain of sand barriers in the Gulf of Maine (FitGerald and others, 1994). The barrier islands and spits are pinned to bedrock or glacial promontories along the coast and stretch approximately 30 km from Ipswich Bay to the New Hampshire border.

Acknowledgments
This series of seafloor maps is dedicated to the late Susan Snow-Cotter (1961–2006), who for many years was the leader in efforts to balance environmental policies with human uses of ocean resources in Massachusetts. She was instrumental in establishing and maintaining the cooperative seafloor-mapping program, which is funded by the Coastal and Marine Geology Program of USGS and the CZM. We wish to thank Anthony Wilbur and Daniel Sampson of CZM for their encouragement and support of marine research. Assistance in the field was provided by Emily Bergeron, William Danforth, Barry Irwin, and Charles Woolsey of the USGS. We also thank Jane Denny, William Danforth, and David Foster of USGS for their help in processing the large amount of acoustic data. The maps benefited from reviews by Jane Denny, S. Jeffress Williams, Matthew Arvanou, Anthony Wilbur, and Daniel Sampson.

References Cited
Ackerman, S., Butman, B., Barnhardt, W., Danforth, W., and Crocker, J., 2006. High-resolution geologic mapping of the inner continental shelf: Boston Harbor and Approaches, Massachusetts. U.S. Geological Survey Open-File Report 2006-1008. Available on DVD and online at <http://pubs.usgs.gov/of/2006/1008/>.

Barnhardt, W.A., Andrews, B.D., and Butman, B., 2006. High-resolution geologic mapping of the inner continental shelf, Nahant to Gloucester, Massachusetts. U.S. Geological Survey Open File Report 2005-1293. Available on DVD and online at <http://pubs.usgs.gov/of/2005/1293/>.

Butman, Bradford, Valentine, P.C., Middleton, T.J., and Danforth, W.W., 2007. A GIS Library of multibeam data for Massachusetts Bay and the Stellwagen Bank National Marine Sanctuary offshore of Boston, Massachusetts. U.S. Geological Survey Data Series 99. DVD-ROM. Available online at <http://pubs.usgs.gov/ds/99/>.

Danforth, W.W., 1997. Xsonar/Showimage: a complete system for rapid sidescan sonar processing and display. U.S. Geological Survey Open-File Report 97-456, 77 p.

Drons, D.L., and D. Metzger. NGDC Coastal Relief Model, accessed June 15, 2007 at <http://www.ngdc.noaa.gov/rggc/coastal/coastal.html>

FitGerald, D.M., Rosen, P.S., and van Heteren, S., 1994. New England barriers. In: Davis, R.A., ed., *Geology of holocene barrier island systems*. Berlin, Germany: Springer-Verlag, pp. 305-394.

Kaye, C.A. and Barghoorn, E.S., 1964. Late Quaternary sea-level change and crustal rise at Boston, Massachusetts, with a note on the auto-compaction of peat. *Geological Society of America Bulletin*, v. 75, p. 63-80. Available online at http://www.mass.gov/mgis/mg_elev5k.htm

Massachusetts Geographic Information System (MassGIS), 2005. Statewide digital elevation model (1:5000), February 2005. Available online at http://www.mass.gov/mgis/mg_elev5k.htm

Oldale, R.N., Colman, S.M., and Jones, G.A., 1993. Radiocarbon ages from two submerged strandline features in the western Gulf of Maine and a sea-level curve for the northeastern Massachusetts coastal region. *Quaternary Research*, v. 40, p. 38-45.

Poppe, L., Williams, S.J., and Paskevich, V.F., eds., 2005. USGS East-Coast Sediment Analysis: Procedures, Database, and GIS Data. U.S. Geological Survey Open-File Report 2005-1001, supercedes 2000-358 DVD ROM. Available online <http://pubs.usgs.gov/of/2005/1001/>



Science Applications International Corporation (SAIC), 2004. Southern Merrimack embayment multibeam survey. Survey Report. Prepared for U.S. Geological Survey by SAIC, Newport, RI.

Stone, B.D. and Peper, J.D., 1982. Topographic control of deglaciation of eastern Massachusetts—ice lobation and marine incursion. In: B.J. Larson and B.D. Stone eds., *Late Wisconsinan glaciation of New England: Kendall/Hunt, Dubuque, Iowa*, p. 145-166.

Stone, B.D., Stone, J.R., and DiGiacomo-Cohen, M.L., 2006. Surficial geologic map of the Salem-Newburyport East-Wilmington-Rowley 16 Quadrangle Area in Northeast Massachusetts. U.S. Geological Survey Open-File Report 2006-1260-B. Available online at <http://pubs.usgs.gov/of/2006/1260/B/>

Valentine, P.C., Blackwood, D.B., and Paroski, K.F., 2000. Seabed observation and sampling system. U.S. Geological Survey Fact Sheet FS-142-00. Available online at <http://pubs.usgs.gov/fs/142-00/142-00.pdf>

Zen, Ean, Goldsmith, Richard, Ratcliffe, N.M., Robinson, Peter, and Stanley, R.S., 1983. Bedrock geologic map of Massachusetts. U.S. Geological Survey, Washington D.C., scale 1:250,000, 3 sheets.



HIGH-RESOLUTION GEOLOGIC MAPPING OF THE INNER CONTINENTAL SHELF: CAPE ANN TO SALISBURY BEACH, MASSACHUSETTS

Sheet 3. Backscatter intensity of seafloor (gray scale).

By
Walter A. Barnhardt, Brian D. Andrews, Seth D. Ackerman, Wayne E. Baldwin, and Christopher J. Hein

