





Prepared in cooperation with the  
NATIONAL PARK SERVICE (NPS) AND  
THE NATIONAL AERONAUTICS AND SPACE ADMINISTRATION (NASA)



OPEN FILE REPORT 2007-1431  
TILE 64 of 116 (BE)

Brock, J.C., Wright, C.W., Nayegandhi, A., Patterson, M., Wilson, I., and Travers, L.J., 2007, USGS-NPS-NASA Bare Earth (BE) Topography-Padre Island National Seashore U. S. Geological Survey Open File Report 2007-1431 (On DVD).



### Project Description

This Lidar-derived topographic map was produced as a collaborative effort between the U.S. Geological Survey (USGS) Coastal and Marine Geology Program, the Northeast Coastal and Barrier Network of the National Park Service (NPS) Inventory and Monitoring Program, the South Florida Caribbean Network of the NPS Inventory and Monitoring Program, and the National Aeronautics and Space Administration (NASA) Wallops Flight Facility. The aim of the partnership that created this product is to develop advanced survey techniques for mapping nearshore bathymetry and habitat, and to provide a baseline for monitoring and detecting change within National Seabees. This product is based on data from an innovative airborne Light Detection and Ranging (LiDAR) instrument under development at the NASA Wallops Flight Facility, the NASA Experimental Advanced Airborne Research Lidar (EAARL).

## Data Description

The laser soundings used to create a map were collected during September and October of 2005, by the NASA EAARL system mounted on a Cessna 310 aircraft. The EAARL uses a "waveform-formatted" green laser capable of mapping submarine and subaerial (land) topography in a single overflight. The EAARL system is typically flown at 300m altitude above AGL, resulting in a 240m swath for each flightline. Data collection occurred with approximately 50% overlap between flightlines, resulting in about one laser sounding per square meter. The data were processed using the software package LIDAR 360 to produce a point cloud. The point cloud was then projected to a geographic coordinate system and ingested into a Geographic Information System (GIS). The data were organized as 2 km by 2 km data tiles in 32-bit floating-point integer GeoTIFF format. Contour line and hillshade layers were generated from the Lidar data tile and incorporated into this map product.

### Further Reading

Brock, J.C., and Sallenger, A., 2001, Airborne topographic Lidar mapping for coastal science and resource management;

Brock, J.C., Wright, C.W., Nayegandhi, A., Clayton, T., Hansen, M., Longenecker, J., Gesch, D., and Crane, M., 2002, Initial results from a test of the NASA EAARL Lidar in the Tampa Bay Region: Transactions of the Gulf Coast Association of Geological Societies, v. 52, p. 89-98.

Wright, C.W. and Brock, J.C., 2002. EAARL: A Lidar for mapping shallow coral reefs and other coastal environments, in the Proceedings of the Seventh International Conference on Remote Sensing for Marine and Coastal Environments, Miami, May 20-22, 2002: Ann Arbor, MI. Veridian International Conferences. 1 computer optical disc.

Nayegandhi, A., Brock, J.C., Wright, C.W., OConnell, M.J., 2006, Evaluating a small footprint, waveform-resolving lidar over coastal vegetation communities, *Photogrammetric Engineering and Remote Sensing*, Vol. 72, No. 12, pp. 1407-1417.

NORTH AMERICAN VERTICAL DATUM OF 1988  
CONTOUR INTERVAL 1 METER

Padre Island National Seashore  
USGS-NPS-NASA EAARL Bare Earth (BE) Lidar Topography  
Map Tile 658000e\_2984000n\_14z

By John C. Brock<sup>1</sup>, C. Wayne Wright<sup>2</sup>, Amar Nayegandhi<sup>3</sup>, Matt Patterson<sup>4</sup>, Iris Wilson<sup>4</sup>, and Laurinda J. Travers<sup>5</sup>

<sup>1</sup> U. S. Geological Survey, FISC, St. Petersburg, FL

<sup>2</sup> NASA Wallops Flight Facility, Wallops Island, VA

<sup>2</sup> ETI Professionals, Contracted to USGS, St. Petersburg, FL

<sup>4</sup> NPS South Florida/Caribbean Network Inventory and Monitoring Program, Miami, FL

<sup>5</sup> Eckerd College, Contracted to USGS, St. Petersburg, FL

