

DESCRIPTION OF MAP UNITS

DEPTH ZONE 2—INTER-TIDAL TO 30 METERS WATER DEPTH

SLOPE CLASS 1—0 TO 5 DEGREES

- Fine to medium-grained smooth sediment—Low backscatter, low rugosity; typically mud to medium-grained sand; often rippled and (or) burrowed
- Mixed smooth sediment and rock—Moderate to very high backscatter, low rugosity; typically coarse-grained sand, gravel, cobbles, and bedrock
- Rock and boulder, rugose—High backscatter, high rugosity; typically boulders and rugose bedrock
- Medium to coarse-grained sediment—Very high backscatter, low rugosity; typically medium to coarse-grained sediment, with varying amounts of shell hash, in scour depressions

SLOPE CLASS 2—5 TO 30 DEGREES

- Fine to medium-grained smooth sediment—Low backscatter, low rugosity; typically mud to medium-grained sand; often rippled and (or) burrowed
- Mixed smooth sediment and rock—Moderate to very high backscatter, low rugosity; typically coarse-grained sand, gravel, cobbles, and bedrock
- Rock and boulder, rugose—High backscatter, high rugosity; typically boulders and rugose bedrock

DEPTH ZONE 3—30 METERS TO 100 METERS WATER DEPTH

SLOPE CLASS 1—0 TO 5 DEGREES


- Fine to medium-grained smooth sediment—Low backscatter, low rugosity; typically mud to medium-grained sand; often rippled and (or) burrowed
- Mixed smooth sediment and rock—Moderate to very high backscatter, low rugosity; typically coarse-grained sand, gravel, cobbles, and bedrock
- Rock and boulder, rugose—High backscatter, high rugosity; typically boulders and rugose bedrock
- Medium to coarse-grained sediment—Very high backscatter, low rugosity; typically medium to coarse-grained sediment, with varying amounts of shell hash, in scour depressions

SLOPE CLASS 2—5 TO 30 DEGREES

- Fine to medium-grained smooth sediment—Low backscatter, low rugosity; typically mud to medium-grained sand; often rippled and (or) burrowed
- Mixed smooth sediment and rock—Moderate to very high backscatter, low rugosity; typically coarse-grained sand, gravel, cobbles, and bedrock
- Rock and boulder, rugose—High backscatter, high rugosity; typically boulders and rugose bedrock

EXPLANATION OF MAP SYMBOLS

- Area of "no data"—Areas near shoreline not mapped owing to insufficient high-resolution seafloor mapping data; areas beyond 3-nautical-mile limit of California's State Waters were not mapped as part of California Seafloor Mapping Program
- 3-nautical-mile limit of California's State Waters
- Bathymetric contour (in meters)—Derived from modified 2-m-resolution bathymetry grid. Contour interval: 10 m



(that is, sediment typically forming a veneer over bedrock, or rock outcrops having little to no relief) makes up 14.9 percent (27.2 km²) of the map area; 5.4 km² is in Depth Zone 2, and 2.1 km² is in Depth Zone 3. Rock and boulder, rugose (rock and boulder outcrops having high surface complexity) makes up 2.9 percent (5.3 km²) of the map area; 1.5 km² is in Depth Zone 2, and 3.8 km² is in Depth Zone 3. Medium to coarse-grained sediment (in scour depressions consisting of material that is coarser than the surrounding seafloor) makes up 6.2 percent (11.3 km²) of the map area; 1.2 km² is in Depth Zone 2, and 1.0 km² is in Depth Zone 3 (table 1).

Table 1. Coverage of classified seafloor, in square kilometers (sq km) and percent of total area, broken into California Marine Life Protection Act Depth Zones 2 and 3.

Seafloor Class	Total		Depth Zone 2 (water depth 0–30 m)		Depth Zone 3 (water depth 30–100 m)	
	percent	sq km	percent of total	sq km	percent of total	sq km
Fine to medium-grained smooth sediment	76.0	138.3	17.2	31.2	58.8	107.1
Mixed smooth sediment and rock	14.9	27.2	2.9	5.4	12.0	21.8
Rock and boulder, rugose	2.9	5.3	0.8	1.5	2.1	3.8
Medium to coarse-grained sediment	6.2	11.3	0.7	1.2	5.5	10.1



DISCUSSION

This seafloor-character map of the Offshore of Point Reyes map area in northern California was produced using video-supervised, maximum-likelihood classification of the bathymetry and backscatter (intensity of return signals from sonar systems) in summary of the video data collected for the purpose of supervising the classification is shown on sheet 6. Rugosity (a GIS-derived characterization of roughness) and backscatter intensity were used as variants in the classification. The interpreted classifications were then draped over shaded-relief bathymetry (see sheet 2). The substrate classes mapped in this area have been divided into the following California Marine Life Protection Act depth zones: Depth Zone 2 (inter-tidal to 30 m), and Depth Zone 3 (30 to 100 m). In addition, the following slope classes are represented on this map (Coastal and Marine Ecological Classification Standard slope zones are shown in parentheses): Slope Class 1, 0° to 5° (flat), and Slope Class 2, 5° to 30° (steep). Depth Zone 1 (essentially, Depth Zones 4 and 5 [greater than 100 m], and Slope Classes 3 to 5, greater than 30° [steeply sloping to overhang]), are not present in this map area.

Fine to medium-grained smooth sediment (sand and mud) makes up 76.0 percent (138.3 km²) of the map area; 31.2 km² is in Depth Zone 2, and 107.1 km² is in Depth Zone 3. Mixed smooth sediment (sand and gravel) and rock




Figure 1. Detailed view of substrate classes mapped offshore of Point Reyes peninsula (see Box A, on map, for location). Depth Zone 3 (30 to 100 m), Slope Class 1 (0°–5°), and Slope Class 2 (5°–30°). Fine to medium-grained smooth sediment is shown in shades of green; mixed smooth sediment and rock is shown in shades of pink; and medium to coarse-grained sediment is shown in shades of yellow. Bathymetric contour (30 m) added for depth reference.




Figure 2. Detailed view of substrate classes mapped offshore of Point Reyes peninsula, west of Point Reyes headland (see Box B, on map, for location). Depth Zone 2 (inter-tidal to 30 m), Depth Zone 3 (30 to 100 m), Slope Class 1 (0°–5°), and Slope Class 2 (5°–30°). Fine to medium-grained smooth sediment is shown in shades of green; mixed smooth sediment and rock is shown in shades of pink; and medium to coarse-grained sediment is shown in shades of yellow. Bathymetric contour (30 m) added for depth reference.




Figure 3. Rugosity (characterization of roughness derived from bathymetry) draped over shaded-relief bathymetry (see sheet 2) for same area as Figure 1 (Box A on map). Rugosity values are displayed in muted "rainbow" color spectrum that ranges from purple (low rugosity) through green (medium rugosity) to red (high rugosity). Interpreted substrate classes from Figure 1 included for comparison.




Figure 4. Rugosity (characterization of roughness derived from bathymetry) draped over shaded-relief bathymetry (see sheet 2) for same area as Figure 2 (Box B on map). Rugosity values are displayed in muted "rainbow" color spectrum that ranges from purple (low rugosity) through green (medium rugosity) to red (high rugosity). Interpreted substrate classes from Figure 2 included for comparison.

Bathymetric contour data from NOAA Coastal Service Center's Digital Coast (available at <http://coast.noaa.gov/digitalcoast/>) and from U.S. Geological Survey's National Ocean Service, available at <http://nauticalcharts.noaa.gov/>. Drifted shaded-relief bathymetry from map sheet 2. This report, California State Waters limit from NOAA Office of Coast Survey, Division of Technical Services, Zone 105.



Seafloor Character, Offshore of Point Reyes Map Area, California
By
Mercedes D. Erdey and Guy R. Cochrane
2015

Seafloor character mapped in 2013. Bathymetric contours by Mercedes D. Erdey, 2013. GIS database and digital cartography by Mercedes D. Erdey and Guy R. Cochrane. Manuscript approved for publication June 8, 2015.

This map was printed on an electronic plate directly from digital files. Dimensional calibration may vary between electronic plates and printed versions. This map is for informational purposes only. It is not intended for navigation. For more information, contact the U.S. Geological Survey, Office of Coastal and Estuarine Science, 1225 National Center, Reston, VA 20192. Telephone: 703/648-5855. Digital file available at <http://dx.doi.org/10.7927/H4TJ-1114>. Suggested Citation: Erdey, M.D., and Cochrane, G.R., 2015, Seafloor character, Offshore of Point Reyes map area, California, sheet 5 of 10, Open-File Report 2015-1114, U.S. Geological Survey, Reston, VA, 10 p. USGS Open-File Report 2015-1114, pamphlet 30. For more information, contact the U.S. Geological Survey, Office of Coastal and Estuarine Science, 1225 National Center, Reston, VA 20192. Telephone: 703/648-5855. Digital file available at <http://dx.doi.org/10.7927/H4TJ-1114>.