

Manuscript approved for publication September 30, 2015

UNCONSOLIDATED CONTINENTAL SHELF SEDIMENTS	
Soft, unconsolidated sediment (fine sand and mud), predominantly rippled	
Soft, unconsolidated sediment (sand), predominantly rippled	
Unconsolidated, dynamic mound of soft sediment (sand)	
Unconsolidated, dynamic mound of soft sediment (sand), in delta zone	
Soft, unconsolidated delta sediment (sand), predominantly rippled	
Soft, mobile sediment window that has unconsolidated and rippled sediment waves, overlying scoured lag pavement of sand and gravel (sorted bedforms)	

Sme_c/u	Mixed habitat of soft, unconsolidated sediment, locally overlying hard, consolidated sedimentary bedrock
<b>HARD SUBSTRATE ON CONTINENTAL SHELF</b>	
Shd_c/d/f	Hard, consolidated sedimentary rock, boulders, or pinnacle
Sh(b)p_c	Hard, consolidated boulders or pinnacles of sedimentary rock

**Contact**

**Area of “no data”**—Areas near shoreline not mapped owing to insufficient high-resolution seafloor mapping data; areas beyond 3-nautical-mile limit of California’s State Waters were not mapped as part of California Seafloor Mapping Program

**3-nautical-mile limit of California’s State Waters**

**Bathymetric contour (in meters)**—Derived from modified 2-m-resolution bathymetry

This map shows "potential" marine benthic habitats in the Offshore of Scott Creek map area. Marine benthic habitats represent a particular type of substrate, geomorphology, seafloor process, or any other attribute that may provide a habitat for a specific species or an assemblage of organisms. Such maps are based largely on seafloor bathymetry data collected from the Offshore of Scott Creek map area: high-resolution bathymetry (sheet 1); shaded-relief imagery (sheet 2); backscatter (sheet 3); seafloor character (sheet 5); and ground-truth information (sheet 6). This map also uses information from the useABED bottom-sampling compilation by Reid and others (2006). The combination of remotely observed data (for example, multibeam bathymetry and backscatter, seismic-reflection profiles) and ground-truth data (for example, sediment samples) translates to higher confidence in the ability to interpret broad areas of the seafloor (fig. 1).

To avoid any possible misunderstanding of the term "habitat," the term "potential habitat" (as defined by Greene and others, 2005) is used herein to describe a set of distinct seafloor conditions that in the future may qualify as an "actual habitat." Once habitat associations of a species are determined, they can be used to create maps that depict actual habitats, which then need to be confirmed by "ground-truth" surveying using in situ observations, video, and (or) photographic documentation.

Marine benthic habitats are classified using the Benthic Marine Potential Habitat Classification Scheme, a mapping-attribute code developed by Greene and others (1999, 2007). In this map series, habitat-classification codes are based on the deepwater habitat-characterization scheme developed by Greene and others (1999), which was created to not only easily distinguish marine benthic habitats but also to facilitate ease of use and queries within GIS and database programs. The code, which is summarized in chapter 6 in the accompanying pamphlet, is derived from several categories of the Benthic Marine Potential Habitat Classification Scheme (Greene and others, 1999, 2007), and it can be subdivided on the basis of the areal scale of the data.

High-resolution, multibeam-sonar data, converted to bathymetric depth grids (seafloor digital elevation models; sheet 1), are essential to development of the habitat map, as is shaded-relief imagery (sheet 2), which allows for visualization of seafloor terrain, and provides a foundation for interpretation of submarine landforms. Areas of seafloor bedrock exposures are identified by their common sharp edges and high relative relief; these may be contiguous outcrops, isolated parts of outcrop protruding through sediment cover (pinnacles or knobs), or isolated bedrock.

Backscatter maps (sheet 3) are also essential for developing potential benthic habitat maps. High backscatter is further indication of "hard" bottom, consistent with interpretation as rock or coarse sediment. In many locations, areas within or around a rocky feature appear to be covered by a thin veneer of sediment, identified on the habitat map as "mixed" induration (in other words, containing both rock and sediment). Broad, generally smooth areas of the seafloor that lack sharp and angular edge characteristics are mapped as "bedrock" and "fine-grained sediment." The latter category includes various sedimentary features such as seasonal oases and depressions, well as depositional features such as dunes, mounds, or sand waves. Low backscatter, indicative of a "soft" bottom, also significantly aids identification and classification of sedimentary habitats.

Delineated in the Offshore of Scott Creek map area are 9 potential marine benthic habitat types, all of which are located on the continental shelf ("Shell" megahabitat). The meso- and macrohabitats include soft unconsolidated sediment (6 habitat types) such as fine sand and mud, sand, and dynamic features such as mobile sand sheets, sediment waves, and rippled sediment depressions; mixed substrate (1 habitat type) such as soft sand and gravels overlying consolidated sedimentary bedrock and gravel pavement; and hard substrate (2 habitat types) such as dolomite and carbonate bedrock, and gravel pavement with bedrock.

Backscatter data (see sheet 3) show that most of the area is underlain by "soft" materials, consistent with the interpretation that unconsolidated sediments dominate habitat in the map area. Sedimentary processes are quite active, especially on the inner shelf just south of Point Año Nuevo, just north of the map area (see fig. 1-1 in pamphlet), and, thus, habitats are highly dynamic, with sediment transport primarily to the southeast. An extensive bedrock exposure of deformed and differentially eroded bedrock is found offshore just south of Point Año Nuevo, and it extends along the nearshore to Davenport, potentially providing good habitat for rockfish (*Sebastes* spp.).


Of the 128.49 km<sup>2</sup> on the continental shelf in the map area, soft, unconsolidated sediment is the dominant habitat type, covering 109.01 km<sup>2</sup> (84.8 percent). Hard rock covers 18.58 km<sup>2</sup> (14.5 percent), whereas 0.91 km<sup>2</sup> (0.7 percent) consists of mixed hard-soft substrate.

Greene, H.G., Bizzarro, J.J., O'Connell, V.M., and Brylinsky, C.K., 2007. Construction of digital Benthic and Marine habitat biotop maps using a coded classification scheme and its application, in: Todal, P.J. and Greene, H.G., eds. Mapping the seafloor for habitat characterization: Geological Association of Canada Special Publication 53, 141-145.

Greene, H.G., Bizzarro, J.J., Tilden, J.E., Lopez, H.L., and Efridy, M.Z., 2005. The benefits and pitfalls of geographic information systems in marine benthic habitat mapping, in: Wright, D.J. and Scholz, A.J., eds. Place matters: Portland, Oregon State University Press, p. 34-46.

Greene, H.G., Naeem, M., Sogard, S.M., and Efridy, M.Z., 2006. W.V. Sullivan, D.E. McKen, J.E. McKen, J.E. and Caillet, G.M., 1999. A classification scheme for deep seafloor habitats: *Oceanologica Acta*, v. 22, p. 663-678.

Reid, J.A., Reid, J.M., Jenkins, C.J., Zimmerman, M., Williams, S.J., and Field, M.E., 2006. *unSEABED* – Pacific Coast (California, Oregon, and Washington) surficial sediment data release: U.S. Geological Survey Data Series 182, available at <http://pubs.usgs.gov/of/2006/182/>



**Figure 1.** Map showing video-observation locations and sample localities for Offshore of Scott Creek main area.